Heterogeneous endocrine cell composition defines human islet functional phenotypes

AUTHORS

Carmella Evans-Molina[1,2,3,4,5,6,7,18*], Yasminye D. Pettway[8,18], Diane C. Saunders[9,18], Seth A. Sharp[10,18], Thomas SR. Bate[9], Han Sun[10], Heather Durai[9], Shaojun Mei[9], Anastasia Coldren[9], Corey Davis[9], Conrad V. Reihsmann[9], Alexander L. Hopkirk[9], Jay Taylor[9], Amber Bradley[9], Radhika Aramandla[9], Greg Poffenberger[9], Adel Eskaros[9], Regina Jenkins[9], Danni Shi[11], Hakmook Kang[11], Varsha Rajesh[10], Swaraj Thaman[10], Fan Feng[9], Jean-Philippe Cartailler[12], Alvin C. Powers[8,9,13], Kristin Abraham[14], Anna L. Gloyn[10,15,16*], Joyce C. Niland[17*], and Marcela Brissova[9,19*] on behalf of the Integrated Islet Distribution Program

AUTHOR AFFILIATIONS AND FOOTNOTES

Departments of [1]Pediatrics, [2]Medicine, [3]Anatomy, Cell Biology, and Physiology, and [4]Biochemistry and Molecular Biology, the [5]Center for Diabetes and Metabolic Diseases, and the [6]Herman B. Wells Center for Pediatric Research, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[7]Richard L. Roudebush VA Medical Center, Indianapolis, IN 46202, USA

[8]Department of Molecular Physiology and Biophysics, Vanderbilt University School of Medicine, Nashville, TN 37232, USA

[9]Division of Diabetes, Endocrinology, and Metabolism, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN 37232, USA

[10]Department of Pediatrics, Stanford School of Medicine, Stanford, CA 94305, USA

[11]Department of Biostatistics, Vanderbilt University Medical Center, Nashville, TN 37232, USA

[12]Vanderbilt Center for Stem Cell Biology, Vanderbilt University, Nashville, TN 37232, USA

[13]VA Tennessee Valley Healthcare System, Nashville, TN 37232, USA

[14]Division of Diabetes, Endocrinology, & Metabolic Diseases, National Institutes of Health, Bethesda, MD 20892, USA

[15]Department of Genetics, Stanford School of Medicine, Stanford, CA, USA

[16]Stanford Diabetes Center, Stanford School of Medicine, Stanford, CA, USA

[17]Department of Diabetes and Cancer Discovery Science, Arthur Riggs Diabetes and Metabolism Research Institute, Beckman Research Institute, City of Hope, Duarte, CA 91010, USA

[18]These authors contributed equally

[19]Lead Contact

*Corresponding Authors (in alphabetical order)

Address correspondence and requests for reprints to: Marcela Brissova, marcela.brissova@vanderbilt.edu; Carmella Evans-Molina, cevansmo@iu.edu; Anna L Gloyn, agloyn@stanford.edu; and Joyce C Niland, jniland@coh.org

SUMMARY

Phenotyping and genotyping initiatives within the Integrated Islet Distribution Program (IIDP), the largest source of human islets for research in the U.S., provide standardized assessment of islet preparations distributed to researchers, enabling the integration of multiple data types. Data from islets of the first 299 organ donors without diabetes, analyzed using this pipeline, highlights substantial heterogeneity in islet cell composition associated with hormone secretory traits, sex, reported race and ethnicity, genetically predicted ancestry, and genetic risk for type 2 diabetes (T2D). While α and β cell composition influenced insulin and glucagon secretory traits, the abundance of δ cells showed the strongest association with insulin secretion and was also associated with the genetic risk score (GRS) for T2D. These findings have important implications for understanding mechanisms underlying diabetes heterogeneity and islet dysfunction and may provide insight into strategies for personalized medicine and β cell replacement therapy.

KEYWORDS

Human islets, alpha cells, beta cells, delta cells, insulin secretion, glucagon secretion, T1D, and T2D genetic risk score

INTRODUCTION

Diabetes mellitus impacts over 11% of Americans and is the eighth leading cause of death in the U.S., with an estimated economic cost of over $413 billion annually[1,2]. Type 1 diabetes (T1D) accounts for 5-10% of all diabetes cases and results from autoimmune-mediated destruction and loss of the insulin-producing β cells, whereas type 2 diabetes (T2D), the most prevalent diabetes form, is characterized by β cell dysfunction and peripheral insulin resistance. While the etiologies of T1D and T2D are largely distinct, both forms require loss and/or dysfunction of pancreatic β cells, have a strong genetic component, and are heterogeneous regarding disease pathophysiology, progression, and response to therapeutic interventions [3–7]. Increasingly, data from clinical cohorts, tissues derived from human organ donors and mouse models highlight a role for impaired function of other islet cell subtypes, including α cells and δ abnormalities in diabetes pathophysiology[8–18]. Murine models have advanced our understanding of disease pathophysiology; however, there are critical differences in the function and architecture of murine and human pancreatic islets necessitating the comprehensive study of human islets to translate basic science research findings[19–25].

The Integrated Islet Distribution Program (IIDP), formerly the Islet Cell Resource[26], has served as the main source of human islets for research within the U.S., with nearly 300 million islets supplied for use in more than 600 unique studies to date (https://iidp.coh.org/). Located at City of Hope (Duarte, CA) and funded by the National Institutes of Diabetes and Digestive and Kidney Diseases (NIDDK) with targeted programmatic funding from Breakthrough T1D (formerly JDRF), the IIDP's mission is to distribute high-quality human islets to the diabetes research community to support basic science and translational research. The IIDP subcontracts with a group of expert islet isolation centers at institutions across the country, who procure donated pancreata from their local Organ Procurement Organizations (OPOs). Rigorous protocols are followed to isolate the islets while maximizing the yield, viability and purity. An automated algorithm developed by the IIDP ensures fair and equitable islet distribution, matching the islet offers to awaiting researchers who have subscribed to receive islets and other non-islet biospecimens from the IIDP[27,28].

In response to the NIH's call to enhance rigor, reproducibility, and transparency in biomedical research, the IIDP initiated the Human Islet Phenotyping Program (HIPP) at Vanderbilt University Medical Center and the Human Islet Genotyping Initiative (HIGI) Program at Stanford University to provide systematic phenotypic and genotypic assessment of IIDP-supported human islet preparations (**Extended Data Figure 1**)[29]. This workflow comprehensively evaluates islet morphology, purity, and viability (**Extended Data Figure 2**). For

the first time, it also assesses islet cell composition and concurrently examines the *in vitro* β and

α cell function[12,30–35]. It also generates GRS for T1D and T2D while predicting genetic ancestry

across a diverse group of organ donors in the U.S. (see **Figure 1A**). This centralized approach

reduces redundancy and provides cost savings to investigators to accelerate the speed of

scientific discoveries made using these islets. Importantly, the IIDP provides this rich integrated

islet data generated by the HIPP and HIGI for researchers to explore via the online IIDP

Research Data Repository (RDR). Here, we present an analysis of the first 299 IIDP-supported

islet preparations from donors without a clinical diagnosis of diabetes analyzed using the HIPP

and HIGI pipelines. Our integrated analysis of this dataset highlights marked heterogeneity in

islet secretory traits and points to a previously underappreciated heterogeneity in islet cell

composition. We found that the heterogeneity concerning three primary endocrine cell types, α,

β, and δ cells, is associated not only with islet hormone secretion traits but also with sex,

reported race and ethnicity, genetically predicted ancestry, and T2D GRS. While α and β cell

composition influenced insulin and glucagon secretory traits, most notably, the abundance of a

relatively small δ cell population showed the strongest association with insulin and glucagon

secretion traits. Delta cell composition was also associated with the T2D GRS. These findings

have important implications for understanding the mechanisms underlying personalized

signatures—whether genetic, functional, or architectural—that predispose individuals to

diabetes. Additionally, they may influence the progression and treatment outcomes of the

disease, as well as inform β cell replacement therapy using human islets for transplantation.

RESULTS

We present an analysis of the first release of 299 human islet preparations from donors

without diabetes, for which all relevant donor information was available (**Figure 1B**). This

dataset was derived from a diverse pool of donors with respect to reported race, ethnicity, age,

and BMI. Islets from male donors were overrepresented in the dataset (62%), which is reflective

of overall trends in available islets and organ donors in the U.S.[36]. The distribution of reported

race or ethnicity differed between male and female donors (p = 0.014), where the dataset

included a higher percentage of female donors that were reported Black or African American

(15% vs. 5%, p = 0.006). A higher percentage of male donors were Hispanic or Latino, though

this difference was not statistically significant). In 90% of the human islet preparations for which

genotyping data was available, the genetic ancestry of the donor was predicted (n = 268;

Figure 1C). Overall, there was a high concordance between reported race or ethnicity and

primary predicted genetic ancestry within the two datasets (κ= 0.867, p < 0.001). Similar to the

larger dataset, there was a lower percentage of male donors with primary African ancestry

(2.5% vs. 16%, p = 0.0002) and a higher proportion of male donors with Admixed American

ancestry (38.5% vs. 22%, p = 0.009).

There were no significant differences in age, BMI, or HbA1c between male and female

donors by Wilson rank sum test. The mean age of male and female donors was 43 (range 15 –

68 years) and 46 years (range 20 – 65 years), and BMI averaged 29 kg/m^2 in both male (range

13.7 – 46.8 kg/m^2) in female donors (range 17.6 – 47.6 kg/m^2). The mean HbA1c of male and

female donors was 5.4% (range 4 – 6.3%) and 5.3% (range 4.3 – 6.4%). Of note, this dataset

includes islets from donors within a normal HbA1c range, defined by the American Diabetes

Association as <5.7% (n = 223, mean ± SD: 5.2 ± 0.3, range 4.0 – 5.6), as well as a subset of

islets from donors with an HbA1c ≥ 5.7% (n = 76, mean ± SD: 5.9 ± 0.2, range 5.7 – 6.4). In the

group of donors where genetic ancestry was predicted, the average age was 44 (range 15 – 68

years) and 46 years (range 20 – 65 years) in male and female donors, respectively. The

average and distribution of BMI and HbA1c were similar to the larger dataset. Together, these

data form a powerful dataset to understand population-based differences in human islet

phenotype.

Islet processing traits associated with islet purity and viability

IIDP-affiliated islet isolation centers identify, process donor pancreata, and distribute

human islet preparations for phenotypic assessment by the HIPP, genotypic assessment by the

HIGI, and scientific investigation by researchers (**Supplemental File 1**). The IIDP RDR includes

data on islet purity and viability, which are measured by islet isolation centers and made

available to researchers at broadcast (i.e., prior to distribution, at the time the islets are offered

to investigators). The post-shipment islet purity and viability is assessed by the HIPP on the day

of arrival. The RDR also includes data related to islet processing, including organ cold ischemia

duration, pre-shipment culture time, and transit time. The current dataset included human islet

preparations from five islet isolation centers: Scharp-Lacy Research Institute (n = 113),

Southern California Islet Cell Resource Center (n = 94), University of Miami Diabetes Research

Institute (n = 30), University of Pennsylvania Islet Transplant Center (n = 21), and the University

of Wisconsin Human Islet Core (n = 35). Pre-shipment culture time averaged 57.5 hours (range

8 – 237 hours), and the average transit time to the HIPP was 25 hours (range 10 – 102 hours).

The mean islet purity measured by the HIPP was 73% (range 15 – 95%), and the mean

dispersed islet cell viability was 76% (range 43 – 93%). Islet transit time and dispersed cell

viability were positively correlated (Spearman r =0.2, unadjusted p-value (p_{unadj}) = 0.001). The

HIPP assessment of islet purity did not show an association with cold ischemia duration, pre-
shipment culture time, or islet transit time.

Islet secretory function is highly heterogeneous amongst donors

A dynamic perifusion system was used to investigate simultaneously β and α cell
hormone secretion in response to both physiologic and pharmacologic stimuli with co-
stimulatory and opposing effects on insulin and glucagon secretion. This included basal glucose
(5.6 mM), high glucose (16.7 mM), high glucose with isobutylmethylxanthine (IBMX), low
glucose (1.7 mM) with adrenaline (Ad), and potassium chloride (KCl) (**Figure 2A-B**). On
average, high glucose led to a 7.9-fold increase in stimulated insulin secretion and inhibited
glucagon secretion by 0.4-fold on average compared to basal glucose. In contrast, low glucose
with adrenaline led to the opposite effect, reducing insulin secretion by approximately 0.2-fold
and increasing glucagon secretion by 5.1-fold compared to basal glucose. Exposure to IBMX, a
phosphodiesterase inhibitor that increases intracellular cAMP levels, potentiated glucose-
stimulated insulin secretion (14-fold) and stimulated glucagon secretion (6.5-fold) compared to
basal glucose level. Additionally, KCl-induced membrane depolarization potently stimulated both
insulin and glucagon secretion by approximately 31- and 11-fold on average, respectively.
To understand how biological variables and islet processing features impacted hormone
secretion across donors, we derived 11 insulin and 9 glucagon secretion traits from the
respective hormone secretion traces (**Figure 2C-D**). For each human islet preparation, we
derived the following secretion traits from each insulin trace: basal insulin secretion, glucose-
stimulated 1st phase secretion (G 16.7 1st area under the curve; AUC), glucose-stimulated 2nd
phase secretion (G 16.7 2nd phase secretion), overall glucose-stimulated insulin secretion (G
16.7 AUC), the glucose stimulation index (SI; G 16.7 SI), cAMP-potentiated insulin secretion (G
16.7 + IBMX 100 AUC), cAMP-potentiated SI (G 16.7 + IBMX 100 SI), response to low glucose
plus adrenaline (G 1.7 + Ad 1 AUC), low glucose plus adrenaline inhibition index (II; G 1.7 + Ad
1 II), KCl-mediated insulin secretion (KCl 20 AUC), and KCl stimulation index (KCl 20 SI). We
also derived the following secretory traits from each glucagon trace: basal secretion, glucose-
inhibited glucagon secretion (G 16.7 AUC), glucose inhibition index (G 16.7 II), cAMP-
potentiated glucagon secretion, cAMP-potentiated SI, low glucose plus adrenaline-induced
secretion, low glucose plus adrenaline SI, KCl-mediated secretion, and the KCl stimulation
index. These data highlight the marked heterogeneity in responses, with the distribution of
values shown for the total dataset and separately by sex and reported race and ethnicity or
primary predicted ancestry in **Extended Data Tables 1-4**. Spearman correlation analyses
revealed a significant correlation between donor age, sex, HbA1c, and BMI with multiple

secretory traits (**Extended Data Figure 3**). Basal insulin secretion was positively correlated with

HbA1c (r = 0.14, p_{unadj} = 0.02) and BMI (r = 0.14, p_{unadj} = 0.02). Age was negatively correlated

with cAMP-potentiated insulin secretion (r = -0.15, p_{unadj} = 0.01) and the glucose stimulation

index (r = -0.12, p_{unadj} = 0.047). The glucose stimulation index also was negatively correlated

with sex (r = -0.12, p_{unadj} = 0.04), BMI (r = -0.14, p_{unadj} =0.02), and HbA1c (r = -0.14, p_{unadj} =

0.02). Additionally, HbA1c was positively correlated with cAMP-potentiated (r = 0.16, p_{unadj} =

0.005) and KCl-mediated glucagon secretion (r = 0.12, p_{unadj} = 0.04).

In multivariable models adjusting for donor age, sex, BMI, HbA1c, reported race or

ethnicity, and islet isolation center, both pre-shipment culture time and islet transit time were

significantly associated with multiple insulin and glucagon secretion traits (**Extended Data**

Figure 4A-B). Given this, we added pre-shipment culture time and islet transit time into future

multivariable regression models as additional covariates unless otherwise stated (8 covariates

total). Of note, islet purity was associated with 3 islet functional traits; however, these data were

unavailable for all observations for inclusion as a covariate in the regression model.

To remove potential confounding effects of other demographic and processing variables,

we next explored associations between donor traits and secretory traits in multivariable

analyses controlling for the other seven covariates. In these analyses, donor BMI was

significantly associated with glucagon secretory traits (**Extended Data Figure 4C-F**).

Specifically, BMI was negatively associated with basal (regression coefficient, b = -0.20, p =

0.005) and KCl-mediated depolarization of glucagon secretion (b = -0.17, p = 0.03) and

positively associated with glucose-mediated glucagon inhibition (b = 0.15, p = 0.03; **Extended**

Data Figure 4D). Donor age, sex, and reported race or ethnicity were associated with multiple

secretory traits prior to, but not after adjusting for multiple comparisons (**Figure 3-4, Extended**

Data Figure 4). Interestingly, we found no differences in insulin or glucagon secretion between

donors with HbA1c < 5.7% (n = 223) and those with HbA1c ≥ 5.7% (n = 76) (**Extended Data**

Figure 4C-D).

For the human islet preparations for which genotyping data were available, we explored

relationships between predicted genetic ancestry and islet function. We found that the

relationships between primary predicted genetic ancestry and secretion were similar to those

observed between reported race or ethnicity and insulin secretion. In addition, genetic ancestry

had a statistically significant effect on basal insulin secretion (**Extended Data Figure 4E,**

Extended Data Figure 5A).

Islet composition strongly influences hormone secretory response

Next, we explored associations between islet function, morphology (i.e., diameter, area,

perimeter), composition, and hormone content (**Figure 5**). Interestingly, in both Spearman

correlation and multivariable analyses, islet composition was highly associated with multiple

secretion traits, especially insulin secretion (**Figure 5A-B, Extended Data Figure 3**). A higher

percentage of β cells was associated with an increase in glucose-stimulated (b = 0.23, p = 2.31

x 10^{-4}), cAMP-potentiated (b = 0.35, p = 1.18 x 10^{-8}), and depolarization-mediated insulin

secretion (b = 0.30, p = 1.16 x 10^{-6}). These secretory traits were negatively associated with the

percentage of islet α and δ cells. Additionally, the percentage of δ cells within the islets was

negatively associated with the glucose stimulation index (b = -0.13, p = 0.02) and inhibition

index of insulin secretion in response to low glucose and adrenaline (b = -0.19, p = 1.76x10^{-3}).

These associations were marked by relatively large effect sizes, as evidenced by the absolute

value of the scaled regression coefficient (0.13 – 0.35).

Regarding the glucagon secretion traits, the percentage of α cells was positively

associated with an increase in basal glucagon secretion (b = 0.18, p = 3.42 x10^{-3}) and total

glucagon secreted in response to low glucose with adrenaline (b = 0.21, p = 3.42 x10^{-3}), while

the percentage of β cells was negatively associated with these two traits (**Figure 5B**).

Additionally, the percentage of α cells was negatively associated with the glucagon stimulation

indices in response to both low glucose with adrenaline (b =-0.17, p =1.58 x10^{-2}) and KCl (b = -

0.18, p = 9.28 x10^{-3}), while the opposite was true for their association with the percentage of δ

cells.

As expected, β and α cell composition were strongly associated with insulin and

glucagon content, respectively (**Figure 5C, Extended Data Figure 3**). However, associations

between hormone content were more striking for glucagon secretion traits than insulin secretion

traits after controlling for the eight covariates and adjusting for multiple comparisons (**Figure**

5A-B). Glucagon content was strongly associated with multiple glucagon secretion traits (**Figure**

5B), including basal glucagon secretion (b = 0.40, p =1.13 x10^{-13}) and total glucagon secreted in

response to all four secretagogues; further, these associations were marked by relatively large

effect sizes (0.21 – 0.42).

In contrast to the impact of islet composition and hormone content, we found no

association between morphological traits (diameter, area, perimeter) of islets purified by hand-

picking that underwent dynamic perifusion and any insulin or glucagon secretion traits (**Figure**

5A-B). Similar results were found when examining the relationship between the morphology-

related traits of the islet preparation as a whole and islet functional traits (**Figure 5A-B**).

Islet composition is influenced by sex, reported race/ethnicity, and predicted genetic ancestry

Given the strong association between islet composition and hormone secretion, we analyzed relationships between islet composition and donor characteristics. On average, analyzed islets were composed of 58% β cells (range 25 – 92%), 34% α cells (range 3 – 68%), and 8% δ cells (range 1 – 19%). Similar to islet hormone secretion, there was significant donor-to-donor heterogeneity in islet endocrine cell composition (**Figure 6A**). Donor sex and reported race or ethnicity had a global effect on the percentage of islet β and α cells in multivariable models incorporating the other seven covariates (**Figure 6B-C**). Specifically, female sex was associated with a higher percentage of α cells and a lower percentage of β cells. Additionally, individuals reported as Asian had a higher percentage of β cells (p = 0.011) and a lower percentage of α cells (p = 0.009). Similar to relationships observed with reported race, donors with predicted East Asian ancestry had a higher percentage of β cells (p = 0.007) and a lower percentage of α cells (p = 0.005) versus those with African, Admixed American, or European ancestry (**Figure 6D**). We observed no significant effect of donor age, BMI, HbA1c, isolation center, pre-shipment culture time or islet transit time on islet composition in multivariable analyses (**Extended Data Figure 6A-B**).

We next investigated associations between islet hormone content, donor demographic, and islet processing traits (**Figure 6E-L, Extended Data Figure 6C-D**). Insulin content was positively associated with donor age (b = 0.14, p = 0.0496) and negatively associated with transit time (b = -0.15, p = 0.03) in multivariable models controlling for the other seven covariates (**Extended Data Figure 6C-D**). In contrast, we found no associations between insulin content and donor sex, reported race or ethnicity, genetic predicted ancestry, BMI, HbA1c, or isolation center in similar multivariable analyses (**Figure 6F-H; Extended Data Figure 6C-D**). Finally, there was a global effect of sex on glucagon content (p = 0.015), but no other associations were noted between glucagon content and other demographic or processing traits in multivariable models (**Figure 5J-L; Extended Data Figure 6C-D**).

Genetic risk for diabetes is associated with islet composition

Both T1D and T2D GWAS signals map to loci associated with islet-enriched genes and their regulatory elements[6,37]. Furthermore, diabetes risk alleles have been associated with *in vivo* measures of islet function in participants without diabetes and linked to worsened islet function in those with T1D[38–40]. Thus, we were interested in determining how genetic risk for diabetes influenced *in vitro* hormone secretion, content, and islet composition in our study. In the current release of human islet preparations where donor genotyping data were available (n

= 268), we generated T1D and T2D GRS, utilizing 67 and 338 single nucleotide variants,

respectively, based on previously published models[7,41]. Using these data, we investigated

whether genetic risk for diabetes predicted any observed differences in islet function,

composition or hormone content in our dataset of human islet preparations from donors without

diabetes. We further investigated the role of "process-specific" partitioned genetic risk scores

previously published from clustering of intermediate traits to delineate heterogeneity in

polygenic risk[42]. We incorporated either the complete or partitioned GRS into multivariable

models that included the following as covariates: donor age, BMI, HbA1c, sex, isolation center,

pre-shipment culture time, islet transit time, and the first five principal components explaining

genetic ancestry (12 covariates total). Interestingly, the T2D GRS was positively associated with

the percentage of islet δ cells (b = 0.17, p = 0.019; **Figure 7**) and, at the same time, δ cells

strongly influenced the islet hormone secretory traits (**Figure 5**), especially those of β cells. We

detected no significant associations between T1D and T2D GRS and either hormone secretion

or content (**Figure 7**). To determine whether donor HbA1c might mediate associations between

the calculated GRS and islet secretory traits, hormone content, or composition, we performed a

similar analysis excluding HbA1c as a covariate (**Extended Data Figure 7**). The significant

association between the T2D GRS and the percentage of islet δ cells remained; however, we

also detected a significant positive association between the T2D GRS and glucagon content

(**Extended Data Figure 7**).

DISCUSSION

Islet physiology is disrupted in both T1D and T2D. The pathways and cell-cell

communications that underlie islet dysfunction in diabetes are poorly understood, yet they may

have important implications in treatment strategies employing both pharmacologic agents and

islet or β cell replacement therapy. Recent emphasis on precision medicine approaches coupled

with increased utilization of single-cell technologies has highlighted an important role for

heterogeneity in diabetes classification, treatment, and molecular phenotypes[43–47]. To

understand how biological variation in humans impacts islet function and diabetes risk, we

generated this research resource from islets distributed through the NIH-funded IIDP. These

islets underwent pre-shipment assessment at one of five islet isolation centers followed by

centralized phenotyping and genotyping through the HIPP and HIGI programs. A strength of our

study is the inclusion of islets from a diverse group of U.S. organ donors with over 40% of

donors having a reported race or ethnicity other than White, non-Hispanic or Latino/a. The IIDP-

HIPP-HIGI workflow has enabled investigation for associations between inherent donor

characteristics, islet composition, genetic risk scores for T1D and T2D, and multiple insulin and
glucagon secretory traits. Our data can be accessed through the IIDP RDR, where researchers
can view and interrogate the dataset used for this analysis and propose their own unique
questions related to human islet biology using the larger RDR dataset. Images presented in this
report are available on the open access Pancreatlas™ platform[48], which supports interactive
exploration of full-resolution islet imaging data.

Our phenotyping workflow includes an assessment of islet purity and viability, insulin and
glucagon secretion measured in a dynamic perifusion system, and a quantitative assessment of
islet composition in a sample of islets distinct from the islet aliquot used for perifusion. Studies
of the human pancreas suggest a greater than four-fold variation in β cell mass amongst
individuals without diabetes[49]. Not surprisingly, we observed a wide range of β cell secretory
responses to glucose, IBMX, and KCl, in line with previous reports [50–52]. While clinical studies
suggest that glucose tolerance declines with age, whether β cell function declines with age
independently of changes in insulin resistance and weight in individuals who maintain normal
glucose tolerance is not clear[53,54]. Similarly, the impact of donor age on insulin secretion *in vitro*
has been mixed in previous reports, with studies finding either no effect or a negative impact of
age on glucose-stimulated insulin secretion[51,55–58]. In addition, many studies report an effect of
BMI on multiple insulin secretory traits, while others have found no correlation[50,51,57,59]. In the
current study, Spearman correlation analyses revealed a significant positive relationship
between BMI and basal insulin secretion and a negative relationship between either BMI or age
and the glucose stimulation index. However, we detected no significant association between any
donor trait and insulin secretion in multivariable regression models adjusting for 7 other donor
and islet processing-related covariates. Our studies differ from these previous studies in that we
included a larger sample size of donors and controlled for multiple potentially confounding
demographic and islet processing features, which we and others have shown to impact *in vitro*
secretory responses[51,57,60].

In contrast to the weak effects of donor demographics on insulin secretion, islet
composition exhibited strong and consistent associations with insulin secretory traits in both
Spearman correlation and multivariable analyses. Consistent with observed heterogeneity in
hormone secretory responses, islet composition was highly heterogeneous amongst islet
preparations, with an approximately 4, 22, and 19-fold variation in the percentages of β, α, and
δ cells, respectively. Beta cell composition or a higher percentage of β cells was associated with
increased glucose-stimulated, cAMP-potentiated, and depolarization-mediated insulin secretion.
While islet isolation severs vascular and neuronal connections, the autonomy of the islet as a

mini-organ is well preserved after isolation, as demonstrated by the persistence of paracrine
interactions and electrical coupling that modulate islet cell-cell communication within islets[61].
Interestingly, although δ cells comprised a relatively low percentage of islet endocrine cell
composition, a higher percentage of δ cells negatively impacted most insulin secretion traits.
This finding is notable given a recent report showing that δ cells regulate the glycemic set point
in healthy mice and that ablation of the δ cell population leads to blood glucose lowering and
increased insulin secretion[62].

Finally, compared to studies that have assessed islet insulin secretion, considerably less
is known about the impact of donor and processing traits on *in vitro* glucagon secretion in islets
from organ donors without diabetes[63]. In living individuals without diabetes, Fræch and
colleagues reported an inverse relationship between fasting glucagon levels and insulin
sensitivity in individuals with normal glucose tolerance[11]. Individuals with reduced insulin
sensitivity also displayed impaired glucagon suppression in response to an oral glucose
challenge, even after adjusting for plasma glucose levels[11]. We found that higher BMI was
associated with lower basal glucagon levels and depolarization-induced glucagon secretion *in*
vitro; however, consistent with the previous study, total glucagon secreted in response to high
glucose was higher, suggesting impaired glucose-mediated glucagon suppression.

The mechanisms that regulate glucagon secretion are not well understood but likely
include a combination of intrinsic and paracrine mechanisms[64,65]. Multiple glucagon secretory
traits were associated with islet composition; however, we found even stronger associations
between glucagon content and total glucagon secreted in response to all secretagogues used in
this study. Interestingly, this same relationship was not observed between insulin content and
insulin secretory responses. One study comparing murine β and α cells noted faster rates of
glucagon granule exocytosis, larger readily releasable pools, and a higher rate of refilling of
these pools after depletion in α cells[66]. Therefore, it is possible that these same mechanisms are
at play in human islets and could explain differences in the association between hormone
content and secretion between β and α cells.

Our observations on the impact of islet composition on secretory traits raise an important
question of whether differences in cell composition are influenced by donor demographics
and/or genetics. Interestingly, we found a global effect of sex and reported race or ethnicity on
the percentage of islet β and α cells. Here, female donors had a higher percentage of α cells,
and donors reported as Asian exhibited a higher percentage of β cells. We note the relatively
small number of donors reported as Asian in the dataset, highlighting a need to confirm this
observation with larger sample sizes. Remarkably, a high T2D GRS was positively associated

with the percentage of islet δ cells, which was consistent with the finding that δ cell composition

negatively impacted most insulin secretion traits, highlighting the inhibitory potential of δ cell-

secreted factors, including somatostatin. Together, these data suggest that, in some individuals,

a genetic predisposition to having more δ cells may impact insulin secretory capacity and

glycemic setpoints, making one more susceptible to the development of T2D. However, the

mechanisms by which genetic variation associated with altered T2D risk influences δ cell

composition are unknown. Recent studies have utilized the abundance of known endocrine

marker proteins to infer associations with gene and protein expression[52,60]; however, this is the

first time to our knowledge that direct quantitative assessment of islet composition has been

compared to hormone secretion.

In summary, this study has identified important relationships between donor

demographics, genetics, islet processing, islet composition, hormone content, function, using

human islet preparations from donors without diabetes. Given their impact on human islet

function, these factors should be considered in interpreting future human islet studies. Further,

they may have implications in strategies for β cell replacement therapy[67]. For example, in 2022

the FDA approved Lantidra[68], the first allogeneic pancreatic islet cellular therapy made from

deceased donor pancreatic cells for the treatment of T1D to ameliorate problematic

hypoglycemia with glucocorticoid-free immunosuppression[69]. Release criteria for islet grafts

have historically included measures of glucose-stimulated insulin secretion. Our data highlight

the impact of cell-cell interactions on islet function and suggest that islet composition may

provide valuable insight into graft function post-transplant. Similarly, we expect that integrating

these findings from 299 donors without diabetes with datasets generated using the same

phenotyping and genotyping pipelines for islets from individuals with T1D and T2D, such as

those generated by the Human Pancreas Analysis Program (HPAP)[70,71] will inform our

understanding of diabetes pathophysiology and lead to the design of new therapies and disease

prevention. Further, whilst we examined factors impacting glucose-stimulated islet responses in

detail, we did not investigate responses to other nutrients, such as fatty acids or amino acids,

which also display significant donor-to-donor heterogeneity, as highlighted by a recent report

from Kolic et al[52]. Finally, while this study is currently the largest of its type, we had multiple

observations that were nominally significant, supporting the need for larger studies and

expanded sample sizes. The diverse IIDP-HIPP-HIGI datasets represent a valuable and

continuously growing resource for the islet biology community that may be used to generate

new scientific hypotheses and guide mechanistic studies of human islets.

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Marcela Brissova, PhD (marcela.brissova@vumc.org). Requests for access to genetic data should be directed to Anna L Gloyn, DPhil (agloyn@stanford.edu).

Materials Availability

This study did not generate new unique reagents.

Data and code availability

Data from the first release of 299 de-identified human islet preparations reported in this paper is available in **Supplemental File 2.** The genetic risk scores and predicted ancestry for each donor and the larger integrated dataset hosted are available for download from the IIDP Research Data Repository. For more information on available datasets and how to access visit https://iidp.coh.org/Resources-Offered/Research-Data-Repository. Genotyping data has been deposited in the European Genome Phenome Archive (EGA) and will be available at publication through a data access agreement.

All original code is publicly available as of the date of publication. Code related to generation of statistical models is available at https://github.com/hakmook/HIPP_project. Code related to prediction of genetic ancestry is available at https://github.com/gloynlab/GeneticAncestry.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

ACKNOWLEDGMENTS

The authors would like to thank the organ donors and their families for their invaluable donations and acknowledge the IIDP collaborators who continue to make this work possible (**Supplemental File 1**). The authors would also like to thank Danielle Gibson (Vanderbilt University Medical Center) for her assistance to the HIPP and Dr. Emily Anderson-Baucum (Indiana University School of Medicine) for her helpful advice and edits. Research reported in this publication was supported by U42 RR017673, UC4 DK098085, U24 DK098085, 1-RSC-2018-561-I-X, 1-RSC-2019-712-S-B, 1-RSC-2020-890-S-B, F30DK134041, T32GM007347,

AUTHOR CONTRIBUTIONS

Conceptualization: M.B., Y.D.P., H.K., D.C.S., S.A.S., A.L.G., J.C.N., and C.E.M.; Data curation:
Y.D.P., D.C.S., S.A.S., H.S., A.L.K., A.B., and F.F.; Formal analysis: H.K., D.S., Y.D.P., S.A.S.,
and D.C.S.; Funding acquisition: C.E.M., K.A., A.L.G., J.C.N., and M.B.; Investigation: H.D.,
S.M., A.C., C.D., C.V.R., J.T., T.SR.B., R.A., G.P., A.E., R.J. V.S., and S.T.; Methodology: S.A.S.,
H.S., D.S., H.K., A.C.P., S.A.S. H.S. A.L.G and M.B.; Project administration: C.E.M., K.A.,
A.L.G., J.C.N., and M.B.; Software: JP.C.; Resources: M.B., C.E.M., A.L.G., J.C.N., JP.C., and
A.C.P.; Supervision: M.B., C.E.M., A.L.G, J.C.N., H.K., A.C.P., and K.A.; Visualization: Y.D.P.
and D.C.S.; Writing – original draft: Y.D.P., C.E.M., and M.B.; Writing – review & editing: all
authors

DECLARATION OF INTERESTS

410 A.L.G's spouse is an employee of Genentech and holds stock options in Roche.

 FIGURE LEGENDS

Figure 1. Integration of phenotyping and genotyping assessment of a diverse set of IIDP-
derived human islet preparations. (**A**) Schematic illustrating the human islet phenotyping and
genotyping pipelines utilized by the HIPP and HIGI. The HIPP performs analysis of islet
morphology, purity, viability, histology, and function; the remaining sample is archived in a
biorepository for additional analyses. The HIGI receives a non-islet tissue sample and isolates
DNA to sequence for genetic ancestry prediction and calculation of genetic risk scores for type 1
and type 2 diabetes. (**B**) Descriptive statistics of donor demographics by sex and reported race
or ethnicity (n = 299). Values are reported as mean ± standard deviation, followed by the range.
(**C**) Descriptive statistics of donor demographics by sex and primary predicted ancestry (n =
268). Values are reported as mean ± standard deviation, followed by the range. **p < 0.01 vs.
male donors; ***p < 0.001 vs. male donors.

Figure 2. Islet secretory function is highly heterogeneous amongst donors. Average (**A**)
insulin and (**B**) glucagon secretion from donor islets normalized to islet equivalents (IEQs). Data
are displayed as mean ± 95% confidence interval (n = 299). (C) Schematic and tables
describing insulin and (**D**) glucagon secretion traits derived from individual islet secretion traces.
Descriptive statistics by sex and reported race/ethnicity are provided in **Extended Data Tables**
1 and 2. G – Glucose (mM); IBMX – isobutylmethylxanthine (µM); A – adrenaline (µM); KCl –
potassium chloride (mM).

Figure 3. Islet secretion traits by sex. Violin plots comparing (**A**) insulin and (**B**) glucagon
secretion traits between female (n = 115) and male (n = 184) donors. The solid line in each
violin plot depicts the median, and the dotted lines represent the 1st and 3rd quartiles. Global P-
value is based on the F-test while controlling for the seven covariates (age, sex, BMI, HbA1c,
islet isolation center, pre-shipment culture time, and islet transit time) and adjusting for multiple
comparisons. Comparisons where the unadjusted p-value (p_{unadj}) was significant are noted in
italics.

Figure 4. Islet secretion traits by reported race or ethnicity. Violin plots comparing (**A**)
insulin and (**B**) glucagon secretion traits by reported race or ethnicity (n = 12 Asian, n = 26 Black
or African American, n = 86 Hispanic or Latino/a, n = 175 White). The solid line in each violin
plot depicts the median, and the dotted lines represent the 1st and 3rd quartiles. Global p-value is
based on the F-test while controlling for the seven covariates (age, sex, BMI, HbA1c, islet
isolation center, pre-shipment culture time, and islet transit time) and adjusting for multiple

comparisons. Comparisons where the unadjusted p-value (p_{unadj}) was significant are noted in

italics.

Figure 5. Islet composition and hormone content are associated with multiple islet

secretory traits. (A-B) Heatmaps depicting regression coefficients of each morphologic

variable for islets that underwent perifusion (n = 186), islet composition variable (n = 299), and

total islet insulin and glucagon content (n = 299) after being incorporated into multivariable

regression models controlling for age, sex, BMI, HbA1c, islet isolation center, reported race or

ethnicity, pre-shipment culture time, and islet transit time for each (**A**) insulin or (**B**) glucagon

secretion trait. (**C**) Heatmaps depicting regression coefficients of each islet composition variable

after being incorporated into multivariable regression models for total insulin or glucagon

content after adjusting for the eight covariates (n = 299). Adjusted p-values are indicated where

*$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$. Comparisons where the unadjusted p-value (p_{unadj}) was

significant are noted in italics.

Figure 6. Islet composition and hormone content are highly variable and influenced by

donor sex and ancestry. (A) Representative images of collagen gel-embedded islets and

associated quantification of islet composition (n = 299). Violin plots comparing islet composition

by (**B**) donor sex, (**C**) reported race or ethnicity, and (**D**) predicted primary genetic ancestry. (**E**)

Islet insulin content normalized to IEQ (n = 299). Violin plots comparing islet insulin content by

(**F**) donor sex, (**G**) reported race or ethnicity, and (**H**) predicted primary genetic ancestry. (**F**) Islet

glucagon content normalized to IEQ (n = 299). Violin plots comparing islet glucagon content by

(**J**) donor sex, (**K**) reported race or ethnicity, and (**L**) predicted primary genetic ancestry. Donor

sex: n = 115 female, n = 184 male. Reported race or ethnicity: n = 12 Asian, n = 26 Black or

African American, n = 86 Hispanic or Latino/a, n = 175 White. Primary predicted genetic

ancestry: n = 86 Admixed American, n = 21 African, n = 11 East Asian, n = 150 European. *$p <$

0.05, **$p < 0.01$. For sex, reported race or ethnicity, and genetic ancestry, the global adjusted p-

value is based on the F-test while controlling for the other seven covariates. INS – insulin; GCG

– glucagon; SST – somatostatin.

Figure 7. Associations between genetic risk scores for type 1 and type 2 diabetes and

islet function, hormone content, and composition. (A-B) Heatmaps depicting coefficients of

the GRS for (**A**) T1D and (**B**) T2D along with partitioned GRS variables after being incorporated

into multivariable regression models for each secretion trait, hormone content, and islet

composition variable (n = 268). Each model included the following covariates: donor age, sex,

BMI, HbA1c, islet isolation center, pre-shipment culture time, islet transit time, and the first 5

principal components explaining primary genetic ancestry. "T1D GRS" encompasses the genetic

risk for T1D (67 total variants). "HLA DR/DQ" represents the sum of T1D risk at HLA-DR/DQ

haplotypes. "HLA Class 1" and "HLA Class 2" represent the sum of T1D risk at HLA class 1 and

2 alleles, respectively, excluding HLA DR/DQ haplotypes. "Non-HLA" represents the sum of T1D

risk from variants outside the HLA region, genome-wide. "T2D GRS" encompasses genetic risk

for T2D (338 total variants). "Beta Cell," "Proinsulin," "Obesity," "Lipodystrophy," and "Liver/Lipid"

represent partitioned T2D GRS clusters encoding the related variants. $*p < 0.05$. Comparisons

where the unadjusted p-value (p_{unadj}) was significant are noted in italics.

EXTENDED DATA FIGURE AND TABLE LEGENDS

Extended Data Figure 1, related to Figure 1. Schematic of multimodal data integration by phenotyping and genotyping pipelines within Integrated Islet Distribution Program (IIDP) and data deposition in Resource Data Repository (RDR).

Extended Data Figure 2, related to Figure 1. Example HIPP procedures. (**A**) Representative images of a human islet preparation stained with dithizone (DTZ) to delineate islet versus non-islet tissue. Using the Count and Measure function in cellSens, images of DTZ-stained islets were utilized to quantify the total islet equivalents (IEQs), purity, and islet morphology. Tissue depicted in red on the image mark-up indicates positively identified islet tissue, while green marks non-islet tissue. (**B**) Representative images of handpicked islets pre- and post-perifusion. The red color on the image markup indicated positively identified islet tissue by the Count and Measure function in cellSens. The scale bar is 500 µm.

Extended Data Table 1, related to Figure 2. Insulin secretion is highly heterogeneous amongst donors. Descriptive statistics of the eleven insulin secretion traits by donor sex and reported race or ethnicity. Data are displayed as mean ± SD, followed by the range.

Extended Data Table 2, related to Figure 2. Glucagon secretion is highly heterogeneous amongst donors. Descriptive statistics of the nine glucagon secretion traits by donor sex and reported race or ethnicity. Data are displayed as mean ± SD, followed by the range.

Extended Data Table 3, related to Figure 2. Distribution of insulin secretion traits is similar in n = 268 subset. Descriptive statistics of the eleven insulin secretion traits by donor sex and genetic ancestry. Data are displayed as mean ± SD, followed by the range.

Extended Data Table 4, related to Figure 2. Distribution of glucagon secretion traits is similar in n = 268 subset. Descriptive statistics of the nine glucagon secretion traits by donor sex and genetic ancestry. Data are displayed as mean ± SD, followed by the range.

Extended Data Figure 3, related to Figure 2. Relationship between donor demographics, islet processing, islet morphology, and functional traits. Heatmap of Spearman r correlation coefficients between the indicated variables in the corresponding labeled row and column. For donor sex, the male was coded as 0 and female as 1. *$p < 0.05$; †$p < 0.01$; ‡$p < 0.001$. All p-values are unadjusted.

Extended Data Figure 4, related to Figures 3 and 4. Donor and islet processing traits correlate with islet function. (**A-B**) Heatmaps depicting regression coefficients of each islet

processing trait after being incorporated into multivariable regression models for each (**A**) insulin

and (**B**) glucagon secretion trait. Each model included the following covariates: age, sex,

reported race or ethnicity, BMI, HbA1c, and islet isolation center. Pre-shipment culture time and

islet transit time were included as additional covariates for all subsequent models, unless noted

otherwise. (**C-D**) Heatmaps depicting regression coefficients of donor age, BMI, and

prediabetes status after being incorporated into multivariable regression models for each (**C**)

insulin and (**D**) glucagon secretion trait while controlling for the other seven final covariates.

Adjusted p-values are indicated where *p < 0.05, **p < 0.01, and **p < 0.001. For associations

that are significant prior to adjusting to multiple comparisons, the unadjusted p-value is noted in

italics. (**E-F**) Tables summarizing the global P-value when comparing differences by donor sex,

reported race or ethnicity, genetic ancestry, and isolation center for each (**E**) insulin and (**F**)

glucagon secretion trait. The global p-value is based on the F-test while controlling for the other

seven covariates. Analyses involving genetic ancestry controlled for all covariates except

reported race or ethnicity. n.s. indicates an insignificant global adjusted p-value and un-adjusted

p-values. For results that were significant prior to adjusting to multiple comparisons, the

unadjusted p-value is noted in italics. n = 299 for all analyses except those involving islet purity

(n = 269), elevated HbA1c (n = 76 elevated HbA1c, n = 223 normal HbA1c), and genetic

ancestry (n = 268).

Extended Data Figure 5, related to Figure 4. Islet secretion traits by genetic ancestry.

Violin plots comparing (**A**) insulin and (**B**) glucagon secretion traits by primary predicted genetic

ancestry (n = 86 Admixed American, n = 21 African, n = 11 East Asian, n =150 European). The

solid line in each violin plot depicts the median, and the dotted lines represent the 1^{st} and 3^{rd}

quartiles. *p < 0.05 after adjusting for multiple comparisons. Comparisons where the unadjusted

p-value (p_{unadj}) was significant are noted in italics. Global p-value is based on the F-test while

controlling for the seven covariates (donor age, sex, BMI, HbA1c, islet isolation center, pre-

shipment culture time, and islet transit time).

Extended Data Figure 6, related to Figure 6. Correlation between donor demographic and

islet processing variables with islet composition and hormone content. (**A**) Heatmaps

depicting regression coefficients of donor and islet processing variables when incorporated into

multivariable regression models with a composition variable (% β, α, or δ cells) as the outcome

variable. For models related to islet processing traits, the following covariates were included:

donor age, sex, reported race or ethnicity, BMI, HbA1C, and islet isolation center. For donor

traits, pre-shipment culture time and islet transit time were also included as covariates. (**B**) Table

summarizing the global p-value when comparing differences by isolation center for each composition variable. The global p-value is based on the F-test while controlling for the other seven covariates. (**C**) Heatmaps depicting regression coefficients of donor and islet processing variables when incorporating into multivariable regression models insulin and glucagon content. For models related to islet processing traits, the following covariates were included: age, sex, reported race or ethnicity, BMI, HbA1C, and islet isolation center. For donor traits, pre-shipment culture time and islet transit time were also included as covariates. (**D**) Table summarizing the global P-value when comparing differences by isolation center for hormone content. The global P-value is based on the F-test while controlling for the other seven covariates. (**E**) Correlation between insulin content with donor age, and (**F**) transit time. n = 299 for all analyses; n.s. indicates insignificant global adjusted p-value and unadjusted p-values.

Extended Data Figure 7, related to Figure 7. Associations between T1D/T2D GRSs and islet function, hormone content, and composition, disregarding HbA1c. (**A-B**) Heatmaps depicting coefficients of the GRS for (**A**) T1D and (**B**) T2D along with partitioned GRS variables after being incorporated into multivariable regression models for each secretion, hormone content, and islet composition variable (n = 268). Each model included the following covariates: donor age, sex, BMI, islet isolation center, pre-shipment culture time, islet transit time, and the first five principal components explaining primary genetic ancestry. *p < 0.05. For associations that are significant prior to adjusting to multiple comparisons, the unadjusted p-value is noted in italics.

METHODS

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Primary human islet isolation

Human islet preparations from cadaveric organ donors (n = 299) were isolated from five affiliated islet isolation centers across the U.S and distributed to the Human Islet Phenotyping Program through the IIDP from 2016-2024. Donor demographic information is detailed in **Figure 1**. De-identified human pancreatic specimens do not qualify as human subjects research under the Vanderbilt or the Stanford Institutional Review Boards. All assays were performed on the day of islet arrival. Islet preparations included in this manuscript (n = 299) were isolated from five IIDP-affiliated isolation centers: Scharp-Lacy (n = 113), Southern California Islet Cell Resource Center (n = 94), University of Miami (n = 30), University of Pennsylvania Islet Transplant Center (n = 27), and the University of Wisconsin Human Islet Core (n = 35).

METHOD DETAILS

All human islet preparations were assessed according to standardized protocols established by the HIPP and HIGI of the IIDP and are publicly available at https://iidp.coh.org/SOPs. All HIPP-related assessments are conducted on the day-of-arrival. Methods are briefly summarized below:

Islet Purity and Morphology

A representative sample of the unpicked human islet preparation was prepared by combining a 320 µL aliquot of islet preparation and 680 µL CMRL medium. The representative sample was stained with 50 µL of 1 mg/mL dithizone/DMSO/PBS (DTZ; Sigma, catalog #D5130) for 1-2 min at room temperature. After adding 1 mL of CMRL medium, brightfield and darkfield images of stained islets were captured at 10X magnification. Islet (stained) and non-islet (unstained) area was determined using the Count and Measure function in cellSens (Olympus). Islet purity was calculated as the following:

$$Islet\ Purity\ (\%) = \frac{Total\ islet\ tissue\ area}{Total\ islet\ tissue\ area + Total\ non-islet\ tissue\ area} \times 100$$

Using the Count and Measure function in cellSens, morphological characteristics of each islet were also determined, including average islet radius, diameter, perimeter, and area. Measurements were performed in triplicate to ensure accuracy.

Islet Viability

Qualitative assessment of islet viability upon arrival to the HIPP was performed using fluorescein diacetate (FDA) and propidium iodide (PI). In brief, 400 µL of well-mixed total islet suspension was added to a cell culture dish, and 10 µL of PI (Sigma Aldrich, catalog # P4170) and 10 µL of FDA (Sigma Aldrich, catalog # F-7378) were added to the islet suspension in succession. The plate was incubated in the dark for 15 mins, and the preparation was imaged immediately using an Olympus SZX12 stereomicroscope system. Multiple fields of view were captured from two replicates per donor to ensure that FDA/PI staining was visualized in 50-100 islets per preparation.

Additionally, a quantitative assessment of islet viability upon arrival to the HIPP was performed using trypan blue dye. Two aliquots of ~100 handpicked islets were prepared in 1.5 mL tubes and washed three times using 2 mM EDTA/PBS via centrifugation at 200 rcf for 1 min at 4 °C. Islets were dispersed into a single cell suspension using Accutase (Innovative Cell Technologies, Inc, catalog # AT-104) and triturated at a slow speed for 10 minutes using an electronic multichannel pipet with tips. The reaction was quenched with CMRL-1066 media

containing 10% FBS, 2 mM L-glutamine, and 1% penicillin/streptomycin, and cells were washed

twice via centrifugation at 500 rcf for 3 mins at 4 °C. The cell pellet was resuspended in 50-75

µL of CMRL-1066 medium by gently flicking the tube. Next, 10 µL of the cell suspension was

transferred to a new 1.5 mL tube and mixed with 10 µL Trypan Blue dye (Invitrogen, catalog #

T10282). The cell chamber slide was loaded with 10 µL of the mixture in each compartment,

and cell counts were acquired using the Countess II cell counter following instrument guidelines.

Cell counts were repeated for the second aliquot of handpicked islets. Reported viability

represents the average percentage of viable cells for duplicates of purified islets based on two

cell counts per aliquot.

Islet Function in a Dynamic Perifusion System

Islet function was assessed via dynamic perifusion. On the day islets were received, half

of the islet shipment (1000-5000 islet equivalents (IEQs)) was plated in a 10-cm non-tissue

culture treated dish and cultured in CMRL-1066 plus 10% FBS media at 37 °C and 5% CO_2 for

2 hours prior to perifusion. Under microscope guidance, 267-300 IEQs were handpicked, islet

size and count were recorded, and high-resolution brightfield and darkfield images were

captured at 10x magnification. All islets were then transferred to a clear Eppendorf tube for

loading into the perifusion chamber, fractions were collected at 1 mL/min, and secretagogues

were changed at predetermined fractions. The following secretagogues were used for these

perifusion studies: 5.6 mM glucose (baseline media; G 5.6), 16.7 mM glucose (G 16.7), 16.7

mM glucose with 100 µM isobutylmethylxanthine (IBMX; G 16.7 + IBMX 100), 1.7 mM glucose

with 1 µM adrenaline (G 1.7 + Ad 1), and 5.6 mM glucose with 20 mM KCl (G 5.6 + KCl 20). All

secretagogues were prepared in a perifusion medium containing 3.2 g $NaHCO_3$, 0.58 g L-

glutamine, 0.11 g sodium pyruvate, 1.11 g HEPES, 1 bottle DMEM, 1 g RIA-grade BSA, and 70

mg ascorbate in 1 L ultra-purified water. The perifusion medium was filtered and de-gassed prior

to preparation of secretagogues.

After retrieving islets from the perifusion chamber and placing them into a 60-mm non-

treated culture dish, the IEQ of the retrieved islets was determined and islets were centrifuged

for 3 min at 200 rcf at room temperature. Islet hormone extracts were prepared by incubating

the retrieved islet pellet in 200 µL of fresh acid ethanol (50 µL 5N HCl + 5.5 mL 95% ethanol) for

24 hours. Samples were then spun at 3000 rcf for 5 min, aliquoted to 2 mL screwcap tubes, and

stored at -80 °C until further processing.

Insulin concentration in perifusates and islet extracts was measured via ELISA (2020 –

present; Human Insulin ELISA Kit, Mercodia, catalog # 10-1113-10) or RIA (Human Insulin-

Specific RIA, Millipore catalog # HI-14K). In the same samples, glucagon was also measured

via ELISA or RIA (2021 – present: Quantikine Glucagon ELISA Kit, R&D Systems catalog #

DGCG0; 2020 – 2021: HTRF Glucagon Detection Kit, Cisbio catalog #62CGLPEH; prior to

2020: Glucagon RIA, Millipore catalog #GL-32K). For each assay, standards and quality

controls were measured in duplicates, and perifusate and islet extract samples were measured

in duplicates (RIA) or single point (ELISA). Secreted insulin and glucagon concentrations were

normalized to IEQs and expressed as ng/100 IEQs/min. Alternatively, secreted insulin and

glucagon concentrations were normalized to their respective total hormone content and

expressed as %content/min.

Islet Composition

To assess islet composition and acinar cell components, an aliquot of the human islet

preparation was immobilized in a collagen I gel, embedded, and sectioned for histologic

analysis. An aliquot containing approximately 500 IEQs of the human islet preparation was

transferred into a 1.5 mL tube and washed three times in 1X PBS via centrifugation at 200 *xg* for

3 min. After removing the supernatant, 150 µL of collagen I working solution (563 µL collagen I

stock, 167 µL ultrapurified H_2O, 200 µL 5X DMEM, 20 µL HEPES, and 50 µL $NaHCO_3$) was

added to the islet pellet and the mixture was transferred into a well in the center of a 96-well

plate. The plate was incubated at 37 °C for 1.5 hours to allow the gel to solidify, then fixed on ice

for 15 min, filling the well with cold 4% paraformaldehyde/PBS. After the brief fixation, the

paraformaldehyde solution was removed and, using a 28 G needle, the gel was loosened from

the sides of the well and transferred with a spatula into a 12-well plate containing cold 4%

paraformaldehyde/PBS. The gel was fixed for an additional 45 min, shaking on ice, before being

washed three times with 1X PBS for 20 min shaking on ice, transferring the gel to a new well

containing fresh 1X PBS for each wash. Following an overnight incubation in 30% sucrose/1X

PBS, the collagen I gel was embedded into cryomolds containing OCT, imaged, and flash frozen

on dry ice. Islet gels were stored at -80 °C until sectioning.

Following sectioning of collagen gels, immunofluorescence staining of islets was

performed. In brief, 8-µm islet cryosections were allowed to thaw at room temperature and air-

dried for 30 mins. Sections were washed with 1X PBS 3 times for 5 mins to remove the OCT,

permeabilized with 0.2% Triton for 15 mins at room temperature, and washed 3 times in 1X

PBS. Sections were blocked with 5% normal donkey serum in 1X PBS at room temperature for

90 mins in a humidified chamber, then incubated overnight at 4 °C in a humidified chamber with

primary antibodies diluted in 0.1% Triton-X-100/1% BSA/1X PBS. Primary antibodies included:

C-peptide (rat, Developmental Studies Hybridoma Bank, GN-ID4, RRID:AB_2631151, 1:100),

Glucagon (mouse, Abcam, ab10988, RRID:AB_297642, 1:250), Glucagon (rabbit, Cell Signaling

Technology, 2760S, RRID:AB_659831, 1:100), Somatostatin (goat, Santa Cruz Biotechnology,
sc-7819, RRID:AB_2302603, 1:500), and HPX1 (mouse, Novus Biologicals, NBP1-18951,
RRID:AB_1625456, 1:100). Sections were washed with 1X PBS 3 times for 10 mins each, then
incubated for 1.5 hours at room temperature in a humidified chamber with secondary antibodies
diluted in 0.1% Triton/1% BSA/1X PBS. Secondary antibodies were purchased from Jackson
ImmunoResearch: Rat IgG-Cy2 (donkey, 712-225-150, RRID:AB_2340673, 1:500), Rat IgG-
Cy5 (donkey, 712-175-150, RRID:AB_2340671, 1:200), Mouse IgG-Cy3 (donkey, 715-165-150,
RRID:AB_2340813, 1:500), Rabbit IgG-Cy5 (donkey, 711-175-152, RRID:AB_2340607, 1:200),
Goat IgG-Cy5 (donkey, 705-175-147, RRID:AB_2340415, 1:200), and Mouse IgG-Cy3 (donkey,
715-165-150, RRID:AB_2340813, 1:500). Sections were counterstained with 1:25,000
DAPI/PBS for 10 mins at room temperature, then washed 3 times for 15 mins each in 1X PBS.
Sections were mounted with SlowFade Gold mounting medium, and islet sections were imaged
using a high-resolution whole slide scanning system (ScanScope FL, Aperio/Leica) connected
to a web-based digital slide repository powered by eSlide Manager and housed in the Vanderbilt
University Medical Center Data Center. A quantitative assessment of islet cell composition and
endocrine/acinar cell compartments was performed using a tissue classifier algorithm (Halo™,
Indica Labs) to analyze 50-100 islets per labeling experiment. Images presented herein are
deposited in the publicly available Pancreatlas™ platform[48], which allows the exploration of full-
resolution islet imaging data in an interactive manner
(https://pancreatlas.org/datasets/853/explore).

Genetic Analysis

DNA on 246/299 donors included in this release was isolated from human islet donor
acinar tissue at Vanderbilt using the Wizard Genomic DNA Purification Kit (Promega, catalog #
A1120). Quantification and quality assessment was performed using the Genomic DNA
ScreenTape assay. DNA was normalized to 50 ng/µL prior to shipment to the HIGI for further
analysis. For 30 samples no acinar tissue was available. For 9 of these, DNA was extracted
from FPPE sections at Stanford using the Zymo QuickDNA FFPE Extraction Kit. For the
remaining 23 donors DNA was isolated from acinar tissue at Stanford using the DNAExtraction
(Qiagen DNeasyBlood&TissueKit, catalog # 69504).
Prior to genotyping all DNA samples were re-quantified using both the Nanodrop and
Qubit platforms. Samples containing less than 500 ng total DNA were speed-vacuumed and
concentrated prior to genotyping. All samples were genotyped using the Illumina Infinium
Omni2.5Exome V1.5 array with ~2.6 million single nucleotide variant (SNV) sites measured.
Genotyping data underwent quality control by filtering of SNVs with excess missing genotypes

(>2%), filtering of SNVs that do not conform to Hardy-Weinberg equilibrium (p<1x10^-6), and
removal of samples with either excess missing genotypes (>3%) or discordant genetic vs
recorded sex. For donor samples with excess missing genotypes, 10 um thick formalin-fixed
paraffin embedded pancreas tissue slides were requested from IIDP. DNA was extracted using
the Zymo QuickDNA FFPE Extraction Kit and quantified by Nanodrop and Qubit. Samples were
then restored using the Infinium HD FFPE Restore Protocol and then re-genotyped. Since FFPE
samples have considerable DNA degradation, if these samples did not pass restoration, thicker
20 um FFPE slides were shipped from IIDP and DNA was extracted and sent for restoration and
genotyping. For samples that had discordant sex, an XY PCR was performed on the original
sample and an additional donor sample from IIDP (10-μm thick formalin-fixed paraffin-
embedded pancreas tissue slides). From the 270 donors passing QC coordinates were then
aligned to the positive strand on genome build GRCh38 and the data imputed against the
TOPMed R3 reference panel. After quality control and imputation, 69.4 million high-quality SNVs
(R2 > 0.3) and 269 samples were available for analysis. Genetic risk scores (GRS) for T1D and
T2D were generated from previously published models using 67 and 338 SNVs respectively[7,41].
T2D partitioned genetic risk scores (pGRS) were generated using 94 SNVs with predetermined
weights from a previous soft-clustering[42]. Genetic ancestry was determined using a random
forest classifier trained on the first 20 principal components of the 1000 Genomes reference
panel (https://github.com/gloynlab/GeneticAncestry).

QUANTIFICATION AND STATISTICAL ANALYSIS

Hormone secretion traits

Using the insulin and glucagon secretion traces normalized to IEQ, eleven insulin
secretion traits and nine glucagon secretion traits were defined and calculated as illustrated in
Figure 2C-D.

Statistical analysis

To determine demographic, islet processing, and morphologic variables associated with
differences in islet secretory function, we first performed Spearman correlation analyses. For
variables with a statistically significant association with islet secretory traits, we utilized
multivariable regression models to examine the relationship among the explanatory variables
and the outcome variable of interest. For each model, we controlled for the following potential
confounders, unless otherwise noted: donor age, sex, BMI, isolation center, HbA1c, reported
race or ethnicity, pre-shipment culture time, and transit time. A statistically significant regression

coefficient for the explanatory variable (p-value < 0.05) denoted a significant association. Given

multiple p-values, we adjusted for the false discovery rate (FDR) at 0.05 to control for multiple

comparisons.

Similarly, to explore potential differences based on prediabetes status or sex, we used

multivariable regression models. In these models, an indicator variable (0 for normal, 1 for pre-

diabetes participants; 0 for male, 1 for female) acted as the explanatory variable. For

prediabetes-related associations, we controlled for all eight of the aforementioned potential

confounders except HbA1c. For sex-related associations, we controlled all potential

confounders except donor sex. The significance of the regression coefficient (b) for the group

indicator (p-value < 0.05) indicated a notable difference in the outcome variable between the

two groups.

To ensure comparability of regression coefficients across models, we normalized the

outcome variables and all continuous explanatory variables, including covariates. This

normalization involved centering the variables by subtracting their mean and scaling them by

their standard deviation. However, this procedure was not applied to categorical covariates in

the model, such as center, self-reported ancestry, and sex. Statistical analyses were all

performed using R Statistical Software version 4.3.0 (R Foundation for Statistical Computing,

Vienna, Austria) and GraphPad Prism version 10.

REFERENCES

1. Centers for Disease Control and Prevention National Diabetes Statistics Report.
https://www.cdc.gov/diabetes/php/data-research/index.html.

2. Parker, E.D., Lin, J., Mahoney, T., Ume, N., Yang, G., Gabbay, R.A., ElSayed, N.A., and
Bannuru, R.R. (2023). Economic Costs of Diabetes in the U.S. in 2022. Diabetes Care *47*, 26–
43. https://doi.org/10.2337/dci23-0085.

3. Mattis, K.K., and Gloyn, A.L. (2020). From Genetic Association to Molecular Mechanisms for
Islet-cell Dysfunction in Type 2 Diabetes. J. Mol. Biol. *432*, 1551–1578.
https://doi.org/10.1016/j.jmb.2019.12.045.

4. Viñuela, A., Varshney, A., Bunt, M. van de, Prasad, R.B., Asplund, O., Bennett, A., Boehnke,
M., Brown, A.A., Erdos, M.R., Fadista, J., et al. (2020). Genetic variant effects on gene
expression in human pancreatic islets and their implications for T2D. Nat Commun *11*, 4912.
https://doi.org/10.1038/s41467-020-18581-8.

5. Rich, S.S., Erlich, H., and Concannon, P. (2018). Genetics of Type 1 Diabetes. In Diabetes in
America. 3rd edition. Bethesda (MD): National Institute of Diabetes and Digestive and Kidney
Diseases (US), C. C. Cowie, S. S. Casagrande, A. Menke, M. A. Cissell, M. S. Eberhardt, J. B.
Meigs, E. W. Gregg, W. C. Knowler, E. Barrett-Connor, D. J. Becker, et al., eds., p. Chapter 12.

6. Krentz, N.A.J., and Gloyn, A.L. (2020). Insights into pancreatic islet cell dysfunction from type
2 diabetes mellitus genetics. Nat. Rev. Endocrinol. *16*, 202–212. https://doi.org/10.1038/s41574-
020-0325-0.

7. Mahajan, A., Spracklen, C.N., Zhang, W., Ng, M.C.Y., Petty, L.E., Kitajima, H., Yu, G.Z.,
Rüeger, S., Speidel, L., Kim, Y.J., et al. (2022). Multi-ancestry genetic study of type 2 diabetes
highlights the power of diverse populations for discovery and translation. Nat. Genet. *54*, 560–
572. https://doi.org/10.1038/s41588-022-01058-3.

8. Unger, R.H., Aguilar-Parada, E., Müller, W.A., and Eisentraut, A.M. (1970). Studies of
pancreatic alpha cell function in normal and diabetic subjects. J Clin Invest *49*, 837–848.
https://doi.org/10.1172/jci106297.

9. Gerich, J.E., Langlois, M., Noacco, C., Karam, J.H., and Forsham, P.H. (1973). Lack of
Glucagon Response to Hypoglycemia in Diabetes: Evidence for an Intrinsic Pancreatic Alpha
Cell Defect. Science *182*, 171–173. https://doi.org/10.1126/science.182.4108.171.

10. Sherr, J., Tsalikian, E., Fox, L., Buckingham, B., Weinzimer, S., Tamborlane, W.V., White,
N.H., Arbelaez, A.M., Kollman, C., Ruedy, K.J., et al. (2014). Evolution of Abnormal Plasma
Glucagon Responses to Mixed-Meal Feedings in Youth With Type 1 Diabetes During the First 2
Years After Diagnosis. Diabetes Care *37*, 1741–1744. https://doi.org/10.2337/dc13-2612.

11. Færch, K., Vistisen, D., Pacini, G., Torekov, S.S., Johansen, N.B., Witte, D.R., Jonsson, A.,
Pedersen, O., Hansen, T., Lauritzen, T., et al. (2016). Insulin Resistance Is Accompanied by
Increased Fasting Glucagon and Delayed Glucagon Suppression in Individuals With Normal
and Impaired Glucose Regulation. Diabetes *65*, 3473–3481. https://doi.org/10.2337/db16-0240.

12. Brissova, M., Haliyur, R., Saunders, D., Shrestha, S., Dai, C., Blodgett, D.M., Bottino, R.,
Campbell-Thompson, M., Aramandla, R., Poffenberger, G., et al. (2018). α Cell Function and
Gene Expression Are Compromised in Type 1 Diabetes. Cell Reports *22*, 2667–2676.
https://doi.org/10.1016/j.celrep.2018.02.032.

13. Camunas-Soler, J., Dai, X.-Q., Hang, Y., Bautista, A., Lyon, J., Suzuki, K., Kim, S.K.,
Quake, S.R., and MacDonald, P.E. (2020). Patch-Seq Links Single-Cell Transcriptomes to
Human Islet Dysfunction in Diabetes. Cell Metab *31*, 1017-1031.e4.
https://doi.org/10.1016/j.cmet.2020.04.005.

14. Omar-Hmeadi, M., Lund, P.-E., Gandasi, N.R., Tengholm, A., and Barg, S. (2020).
Paracrine control of α-cell glucagon exocytosis is compromised in human type-2 diabetes. Nat
Commun *11*, 1896. https://doi.org/10.1038/s41467-020-15717-8.

15. Kellard, J.A., Rorsman, N.J.G., Hill, T.G., Armour, S.L., Bunt, M. van de, Rorsman, P.,
Knudsen, J.G., and Briant, L.J.B. (2020). Reduced somatostatin signalling leads to
hypersecretion of glucagon in mice fed a high-fat diet. Mol. Metab. *40*, 101021.
https://doi.org/10.1016/j.molmet.2020.101021.

16. Vergari, E., Denwood, G., Salehi, A., Zhang, Q., Adam, J., Alrifaiy, A., Asterholm, I.W.,
Benrick, A., Chibalina, M.V., Eliasson, L., et al. (2020). Somatostatin secretion by Na+-
dependent Ca2+-induced Ca2+ release in pancreatic delta cells. Nat. Metab. *2*, 32–40.
https://doi.org/10.1038/s42255-019-0158-0.

17. Dai, X.-Q., Camunas-Soler, J., Briant, L.J.B., Santos, T. dos, Spigelman, A.F., Walker, E.M.,
Drigo, R.A. e, Bautista, A., Jones, R.C., Avrahami, D., et al. (2021). Heterogenous impairment
of α cell function in type 2 diabetes is linked to cell maturation state. Cell Metab *34*, 256-268.e5.
https://doi.org/10.1016/j.cmet.2021.12.021.

18. Hill, T.G., Gao, R., Benrick, A., Kothegala, L., Rorsman, N., Santos, C., Acreman, S., Briant,
L.J., Dou, H., Gandasi, N.R., et al. (2024). Loss of electrical β-cell to δ-cell coupling underlies
impaired hypoglycaemia-induced glucagon secretion in type-1 diabetes. Nat. Metab., 1–12.
https://doi.org/10.1038/s42255-024-01139-z.

19. Brissova, M., Fowler, M.J., Nicholson, W.E., Chu, A., Hirshberg, B., Harlan, D.M., and
Powers, A.C. (2005). Assessment of Human Pancreatic Islet Architecture and Composition by
Laser Scanning Confocal Microscopy. J Histochem Cytochem *53*, 1087–1097.
https://doi.org/10.1369/jhc.5c6684.2005.

20. Cabrera, O., Berman, D.M., Kenyon, N.S., Ricordi, C., Berggren, P.-O., and Caicedo, A.
(2006). The unique cytoarchitecture of human pancreatic islets has implications for islet cell
function. Proc National Acad Sci *103*, 2334–2339. https://doi.org/10.1073/pnas.0510790103.

21. Rodriguez-Diaz, R., Abdulreda, M.H., Formoso, A.L., Gans, I., Ricordi, C., Berggren, P.-O.,
and Caicedo, A. (2011). Innervation Patterns of Autonomic Axons in the Human Endocrine
Pancreas. Cell Metab. *14*, 45–54. https://doi.org/10.1016/j.cmet.2011.05.008.

22. Dai, C., Brissova, M., Hang, Y., Thompson, C., Poffenberger, G., Shostak, A., Chen, Z.,
Stein, R., and Powers, A.C. (2012). Islet-enriched gene expression and glucose-induced insulin
secretion in human and mouse islets. Diabetologia *55*, 707–718.
https://doi.org/10.1007/s00125-011-2369-0.

23. Brissova, M., Shostak, A., Fligner, C.L., Revetta, F.L., Washington, M.K., Powers, A.C., and
Hull, R.L. (2015). Human Islets Have Fewer Blood Vessels than Mouse Islets and the Density of
Islet Vascular Structures Is Increased in Type 2 Diabetes. J. Histochem. Cytochem. *63*, 637–
645. https://doi.org/10.1369/0022155415573324.

24. Alcazar, O., and Buchwald, P. (2019). Concentration-Dependency and Time Profile of
Insulin Secretion: Dynamic Perifusion Studies With Human and Murine Islets. Front. Endocrinol.
10, 680. https://doi.org/10.3389/fendo.2019.00680.

25. Gloyn, A.L., Ibberson, M., Marchetti, P., Powers, A.C., Rorsman, P., Sander, M., and
Solimena, M. (2022). Every islet matters: improving the impact of human islet research. Nat.
Metab. *4*, 970–977. https://doi.org/10.1038/s42255-022-00607-8.

26. Kaddis, J.S., Olack, B.J., Sowinski, J., Cravens, J., Contreras, J.L., and Niland, J.C. (2009).
Human Pancreatic Islets and Diabetes Research. Jama *301*, 1580–1587.
https://doi.org/10.1001/jama.2009.482.

27. Qian, D., Kaddis, J., and Niland, J.C. (2007). A matching algorithm for the distribution of
human pancreatic islets. Comput. Stat. Data Anal. *51*, 5494–5506.
https://doi.org/10.1016/j.csda.2007.02.030.

28. Niland, J.C., Stiller, T., Cravens, J., Sowinski, J., Kaddis, J., and Qian, D. (2010).
Effectiveness of a Web-Based Automated Cell Distribution System. Cell Transplant. *19*, 1133–
1142. https://doi.org/10.3727/096368910x505486.

29. Brissova, M., Niland, J.C., Cravens, J., Olack, B., Sowinski, J., and Evans-Molina, C. (2019).
The Integrated Islet Distribution Program answers the call for improved human islet phenotyping
and reporting of human islet characteristics in research articles. Diabetologia *62*, 1312–1314.
https://doi.org/10.1007/s00125-019-4876-3.

30. Haliyur, R., Tong, X., Sanyoura, M., Shrestha, S., Lindner, J., Saunders, D.C., Aramandla,
R., Poffenberger, G., Redick, S.D., Bottino, R., et al. (2018). Human islets expressing HNF1A
variant have defective β cell transcriptional regulatory networks. J. Clin. Investig. *129*, 246–251.
https://doi.org/10.1172/jci121994.

31. Hart, N.J., Aramandla, R., Poffenberger, G., Fayolle, C., Thames, A.H., Bautista, A.,
Spigelman, A.F., Babon, J.A.B., DeNicola, M.E., Dadi, P.K., et al. (2018). Cystic fibrosis–related
diabetes is caused by islet loss and inflammation. JCI Insight *3*, e98240.
https://doi.org/10.1172/jci.insight.98240.

32. Shrestha, S., Saunders, D.C., Walker, J.T., Camunas-Soler, J., Dai, X.-Q., Haliyur, R.,
Aramandla, R., Poffenberger, G., Prasad, N., Bottino, R., et al. (2021). Combinatorial
transcription factor profiles predict mature and functional human islet α and β cells. Jci Insight *6*,
e151621. https://doi.org/10.1172/jci.insight.151621.

33. Haliyur, R., Walker, J.T., Sanyoura, M., Reihsmann, C.V., Shrestha, S., Aramandla, R.,
Poffenberger, G., Ramirez, A.H., Redick, S.D., Babon, J.A.B., et al. (2021). Integrated Analysis
of the Pancreas and Islets Reveals Unexpected Findings in Human Male With Type 1 Diabetes.
J Endocr Soc *5*, bvab162. https://doi.org/10.1210/jendso/bvab162.

34. Doliba, N.M., Rozo, A.V., Roman, J., Qin, W., Traum, D., Gao, L., Liu, J., Manduchi, E., Liu,
C., Golson, M.L., et al. (2022). α Cell dysfunction in islets from nondiabetic, glutamic acid
decarboxylase autoantibody–positive individuals. J. Clin. Investig. *132*, e156243.
https://doi.org/10.1172/jci156243.

35. Walker, J.T., Saunders, D.C., Rai, V., Chen, H.-H., Orchard, P., Dai, C., Pettway, Y.D.,
Hopkirk, A.L., Reihsmann, C.V., Tao, Y., et al. (2023). Genetic risk converges on regulatory
networks mediating early type 2 diabetes. Nature *624*, 621–629. https://doi.org/10.1038/s41586-
023-06693-2.

36. Organ Procurement and Transplantation Network (OPTN), and Scientific Registry of
Transplant Recipients (SRTR) (2024). OPTN/SRTR 2022 Annual Data Report. U.S. Department
of Health and Human Services, Health Resources and Services Administration; 2024.

37. Fløyel, T., Kaur, S., and Pociot, F. (2015). Genes Affecting β-Cell Function in Type 1
Diabetes. Curr. Diabetes Rep. *15*, 97. https://doi.org/10.1007/s11892-015-0655-9.

38. Dimas, A.S., Lagou, V., Barker, A., Knowles, J.W., Mägi, R., Hivert, M.-F., Benazzo, A.,
Rybin, D., Jackson, A.U., Stringham, H.M., et al. (2014). Impact of Type 2 Diabetes
Susceptibility Variants on Quantitative Glycemic Traits Reveals Mechanistic Heterogeneity.
Diabetes *63*, 2158–2171. https://doi.org/10.2337/db13-0949.

39. Brorsson, C.A., Nielsen, L.B., Andersen, M.L., Kaur, S., Bergholdt, R., Hansen, L.,
Mortensen, H.B., Pociot, F., Størling, J., and Diabetes, H.S.G. on C. (2016). Genetic Risk Score
Modelling for Disease Progression in New-Onset Type 1 Diabetes Patients: Increased Genetic
Load of Islet-Expressed and Cytokine-Regulated Candidate Genes Predicts Poorer Glycemic
Control. J. Diabetes Res. *2016*, 9570424. https://doi.org/10.1155/2016/9570424.

40. Snethlage, C.M.F., Balvers, M., Ferwerda, B., Rampanelli, E., Groen, P. de, Roep, B.O.,
Herrema, H., McDonald, T.J., Raalte, D.H. van, Weedon, M.N., et al. (2024). Associations
between diabetes-related genetic risk scores and residual beta cell function in type 1 diabetes:
the GUTDM1 study. Diabetologia, 1–12. https://doi.org/10.1007/s00125-024-06204-6.

41. Sharp, S.A., Rich, S.S., Wood, A.R., Jones, S.E., Beaumont, R.N., Harrison, J.W.,
Schneider, D.A., Locke, J.M., Tyrrell, J., Weedon, M.N., et al. (2019). Development and
Standardization of an Improved Type 1 Diabetes Genetic Risk Score for Use in Newborn
Screening and Incident Diagnosis. Diabetes Care *42*, 200–207. https://doi.org/10.2337/dc18-
1785.

42. DiCorpo, D., LeClair, J., Cole, J.B., Sarnowski, C., Ahmadizar, F., Bielak, L.F., Blokstra, A.,
Bottinger, E.P., Chaker, L., Chen, Y.-D.I., et al. (2022). Type 2 Diabetes Partitioned Polygenic
Scores Associate With Disease Outcomes in 454,193 Individuals Across 13 Cohorts. Diabetes
Care *45*, 674–683. https://doi.org/10.2337/dc21-1395.

43. Tobias, D.K., Merino, J., Ahmad, A., Aiken, C., Benham, J.L., Bodhini, D., Clark, A.L.,
Colclough, K., Corcoy, R., Cromer, S.J., et al. (2023). Second international consensus report on
gaps and opportunities for the clinical translation of precision diabetes medicine. Nat. Med. *29*,
2438–2457. https://doi.org/10.1038/s41591-023-02502-5.

44. Dorrell, C., Schug, J., Canaday, P.S., Russ, H.A., Tarlow, B.D., Grompe, M.T., Horton, T.,
Hebrok, M., Streeter, P.R., Kaestner, K.H., et al. (2016). Human islets contain four distinct
subtypes of β cells. Nat. Commun. *7*, 11756. https://doi.org/10.1038/ncomms11756.

45. Wigger, L., Barovic, M., Brunner, A.-D., Marzetta, F., Schöniger, E., Mehl, F., Kipke, N.,
Friedland, D., Burdet, F., Kessler, C., et al. (2021). Multi-omics profiling of living human
pancreatic islet donors reveals heterogeneous beta cell trajectories towards type 2 diabetes.
Nat Metabolism *3*, 1017–1031. https://doi.org/10.1038/s42255-021-00420-9.

46. Weng, C., Gu, A., Zhang, S., Lu, L., Ke, L., Gao, P., Liu, X., Wang, Y., Hu, P., Plummer, D.,
et al. (2023). Single cell multiomic analysis reveals diabetes-associated β-cell heterogeneity
driven by HNF1A. Nat. Commun. *14*, 5400. https://doi.org/10.1038/s41467-023-41228-3.

47. Wang, G., Chiou, J., Zeng, C., Miller, M., Matta, I., Han, J.Y., Kadakia, N., Okino, M.-L.,
Beebe, E., Mallick, M., et al. (2023). Integrating genetics with single-cell multiomic
measurements across disease states identifies mechanisms of beta cell dysfunction in type 2
diabetes. Nat. Genet. *55*, 984–994. https://doi.org/10.1038/s41588-023-01397-9.

48. Saunders, D.C., Messmer, J., Kusmartseva, I., Beery, M.L., Yang, M., Atkinson, M.A.,
Powers, A.C., Cartailler, J.-P., and Brissova, M. (2020). Pancreatlas: Applying an Adaptable
Framework to Map the Human Pancreas in Health and Disease. Patterns *1*, 100120.
https://doi.org/10.1016/j.patter.2020.100120.

49. Meier, J.J., Butler, A.E., Saisho, Y., Monchamp, T., Galasso, R., Bhushan, A., Rizza, R.A.,
and Butler, P.C. (2008). β-Cell Replication Is the Primary Mechanism Subserving the Postnatal
Expansion of β-Cell Mass in Humans. Diabetes *57*, 1584–1594. https://doi.org/10.2337/db07-
1369.

50. Kayton, N.S., Poffenberger, G., Henske, J., Dai, C., Thompson, C., Aram, la, R., Shostak,
A., Nicholson, W., Brissova, M., et al. (2015). Human islet preparations distributed for research
exhibit a variety of insulin-secretory profiles. Am J Physiol Endocrinol Metab *308*, E592–E602.
https://doi.org/10.1152/ajpendo.00437.2014.

51. Henquin, J. (2018). Influence of organ donor attributes and preparation characteristics on
the dynamics of insulin secretion in isolated human islets. Physiol. Rep. *6*, e13646.
https://doi.org/10.14814/phy2.13646.

52. Kolic, J., Sun, W.G., Cen, H.H., Ewald, J.D., Rogalski, J.C., Sasaki, S., Sun, H., Rajesh, V.,
Xia, Y.H., Moravcova, R., et al. (2024). Proteomic predictors of individualized nutrient-specific
insulin secretion in health and disease. Cell Metab. *36*, 1619-1633.e5.
https://doi.org/10.1016/j.cmet.2024.06.001.

53. Chang, A.M., Smith, M.J., Galecki, A.T., Bloem, C.J., and Halter, J.B. (2006). Impaired β-
Cell Function in Human Aging: Response to Nicotinic Acid-Induced Insulin Resistance. J. Clin.
Endocrinol. Metab. *91*, 3303–3309. https://doi.org/10.1210/jc.2006-0913.

54. Ferrannini, E., Natali, A., Muscelli, E., Nilsson, P.M., Golay, A., Laakso, M., Beck-Nielsen,
H., Mari, A., and Investigators, R. (2011). Natural history and physiological determinants of
changes in glucose tolerance in a non-diabetic population: the RISC Study. Diabetologia *54*,
1507–1516. https://doi.org/10.1007/s00125-011-2112-x.

55. Ihm, S.-H., Matsumoto, I., Sawada, T., Nakano, M., Zhang, H.J., Ansite, J.D., Sutherland,
D.E.R., and Hering, B.J. (2006). Effect of Donor Age on Function of Isolated Human Islets.
Diabetes *55*, 1361–1368. https://doi.org/10.2337/db05-1333.

56. Niclauss, N., Bosco, D., Morel, P., Demuylder-Mischler, S., Brault, C., Milliat-Guittard, L.,
Colin, C., Parnaud, G., Muller, Y.D., Giovannoni, L., et al. (2011). Influence of Donor Age on
Islet Isolation and Transplantation Outcome. Transplantation *91*, 360–366.
https://doi.org/10.1097/tp.0b013e31820385e6.

57. Lyon, J., Fox, J.E.M., Spigelman, A.F., Kim, R., Smith, N., O'Gorman, D., Kin, T., Shapiro,
A.M.J., Rajotte, R.V., and MacDonald, P.E. (2016). Research-Focused Isolation of Human Islets
From Donors With and Without Diabetes at the Alberta Diabetes Institute IsletCore.
Endocrinology *157*, 560–569. https://doi.org/10.1210/en.2015-1562.

58. Shrestha, S., Erikson, G., Lyon, J., Spigelman, A.F., Bautista, A., Fox, J.E.M., Santos, C.
dos, Shokhirev, M., Cartailler, J.-P., Hetzer, M.W., et al. (2022). Aging compromises human islet
beta cell function and identity by decreasing transcription factor activity and inducing ER stress.
Sci Adv *8*, eabo3932. https://doi.org/10.1126/sciadv.abo3932.

59. D'Aleo, V., Guerra, S.D., Gualtierotti, G., Filipponi, F., Boggi, U., Simone, P.D., Vistoli, F.,
Prato, S.D., Marchetti, P., and Lupi, R. (2010). Functional and Survival Analysis of Isolated

Human Islets. Transplant. Proc. *42*, 2250–2251.
https://doi.org/10.1016/j.transproceed.2010.05.132.

60. Ewald, J.D., Lu, Y., Ellis, C.E., Worton, J., Kolic, J., Sasaki, S., Zhang, D., Santos, T. dos,
Spigelman, A.F., Bautista, A., et al. (2024). HumanIslets.com: Improving accessibility,
integration, and usability of human research islet data. Cell Metab.
https://doi.org/10.1016/j.cmet.2024.09.001.

61. Walker, J.T., Saunders, D.C., Brissova, M., and Powers, A.C. (2021). The Human Islet:
Mini-organ with Mega-impact. Endocr Rev *42*, bnab010-.
https://doi.org/10.1210/endrev/bnab010.

62. Huang, J.L., Pourhosseinzadeh, M.S., Lee, S., Krämer, N., Guillen, J.V., Cinque, N.H.,
Aniceto, P., Momen, A.T., Koike, S., and Huising, M.O. (2024). Paracrine signalling by
pancreatic δ cells determines the glycaemic set point in mice. Nat. Metab. *6*, 61–77.
https://doi.org/10.1038/s42255-023-00944-2.

63. Pettway, Y.D., Saunders, D.C., and Brissova, M. (2023). The human α cell in health and
disease. J Endocrinol. https://doi.org/10.1530/joe-22-0298.

64. Walker, J.N., Ramracheya, R., Zhang, Q., Johnson, P.R.V., Braun, M., and Rorsman, P.
(2011). Regulation of glucagon secretion by glucose: paracrine, intrinsic or both? Diabetes,
Obes. Metab. *13*, 95–105. https://doi.org/10.1111/j.1463-1326.2011.01450.x.

65. Hughes, J.W., Ustione, A., Lavagnino, Z., and Piston, D.W. (2018). Regulation of islet
glucagon secretion: Beyond calcium. Diabetes Obes Metabolism *20*, 127–136.
https://doi.org/10.1111/dom.13381.

66. Barg, S. (2003). Mechanisms of Exocytosis in Insulin-Secreting B-Cells and Glucagon-
Secreting A-Cells. Pharmacol. Toxicol. *92*, 3–13. https://doi.org/10.1034/j.1600-
0773.2003.920102.x.

67. Perez-Frances, M., Bru-Tari, E., Cohrs, C., Abate, M.V., Gurp, L. van, Furuyama, K., Speier,
S., Thorel, F., and Herrera, P.L. (2024). Regulated and adaptive in vivo insulin secretion from
islets only containing β-cells. Nat. Metab. *6*, 1791–1806. https://doi.org/10.1038/s42255-024-
01114-8.

68. RIOS, P., MCGARRIGLE, J.J., LAMONICA, G., LI, Y., COOK, J., JOSHI, I., GHANI, S.,
COOK, D., LOPEZ, D., NASIR, H., et al. (2024). 35-OR: Lantidra, First FDA-Approved Cellular
Therapy to Treat Type 1 Diabetes. Diabetes *73*. https://doi.org/10.2337/db24-35-or.

69. Shapiro, A.M., Lakey, J.R., Ryan, E.A., Korbutt, G.S., Toth, E., Warnock, G.L., Kneteman,
N.M., and Rajotte, R.V. (2000). Islet transplantation in seven patients with type 1 diabetes
mellitus using a glucocorticoid-free immunosuppressive regimen. New Engl J Medicine *343*, 230
238. https://doi.org/10.1056/nejm200007273430401.

70. Kaestner, K.H., Powers, A.C., Naji, A., Consortium, H., and Atkinson, M.A. (2019). NIH
Initiative to Improve Understanding of the Pancreas, Islet, and Autoimmunity in Type 1
Diabetes: The Human Pancreas Analysis Program (HPAP). Diabetes *68*, 1394–1402.
https://doi.org/10.2337/db19-0058.

71. Shapira, S.N., Naji, A., Atkinson, M.A., Powers, A.C., and Kaestner, K.H. (2022).
Understanding islet dysfunction in type 2 diabetes through multidimensional pancreatic
phenotyping: The Human Pancreas Analysis Program. Cell Metab. *34*, 1906–1913.
https://doi.org/10.1016/j.cmet.2022.09.013.

A

B

N = 299

Sex: ● Female ● Male

Reported race or ethnicity:
● Asian
● Black or African American
● Hispanic or Latino/a
● White

Reported race or ethnicity	Female (38%)				Male (62%)			
	% (N)	Age (yr)	BMI (kg/m²)	HbA1C (%)	% (N)	Age (yr)	BMI (kg/m²)	HbA1C (%)
Asian	3% (4)	49 ± 6 (43 – 57)	28.5 ± 4.8 (23.2 – 32.9)	5.5 ± 0.17 (5.3 – 5.6)	4% (8)	49 ± 13 (31 – 66)	26.5 ± 3.0 (21.6 – 29.9)	5.4 ± 0.33 (5.0 – 5.9)
Black or African American	15%** (17)	48 ± 14 (20 – 65)	30.6 ± 7.2 (18.2 – 47.6)	5.5 ± 0.42 (4.5 – 6.2)	5% (9)	47 ± 9 (37 – 62)	30.7 ± 6.9 (25.0 – 46.8)	5.5 ± 0.36 (5.0 – 6.1)
Hispanic or Latino/a	23% (26)	44 ± 12 (24 – 64)	30.0 ± 4.4 (22.9 – 38.7)	5.4 ± 0.42 (4.3 – 6.0)	33% (60)	40 ± 13 (15 – 62)	29.3 ± 4.7 (18.2 – 39.5)	5.5 ± 0.34 (4.5 – 6.1)
White	59% (68)	46 ± 10 (21 – 65)	28.9 ± 6.2 (17.6 – 45.5)	5.3 ± 0.43 (4.4 – 6.4)	58% (107)	45 ± 12 (17 – 68)	28.6 ± 4.8 (13.7 –40.7)	5.3 ± 0.39 (4.0 – 6.3)
Combined	38% (115)	46 ± 11 (20 – 65)	29.4 ± 5.9 (17.6 – 47.6)	5.3 ± 0.42 (4.3 – 6.4)	62% (184)	43 ± 13 (15 – 68)	28.8 ± 4.9 (13.7 – 46.8)	5.4 ± 0.37 (4.0 – 6.3)

Values reported as mean ± SD (range)

C

N = 268

Sex: ● Female ● Male

Primary predicted ancestry:
● East Asian
● African
● Admixed American
● European

Primary predicted ancestry	Female (40%)				Male (60%)			
	% (N)	Age (yr)	BMI (kg/m²)	HbA1C (%)	% (N)	Age (yr)	BMI (kg/m²)	HbA1C (%)
East Asian	5% (5)	48.2 ± 4 (43 – 57)	29.0 ± 4.3 (23.2 – 32.9)	5.4 ± 0.2 (5.2 – 5.6)	4% (6)	51.3 ± 15 (31 – 66)	25.9 ± 3.3 (21.6 – 29.9)	5.5 ± 0.3 (5.1 – 5.9)
African	16%*** (17)	48.5 ± 14 (20 – 65)	30.6 ± 7.2 (18.2 – 47.6)	5.5 ± 0.4 (4.5 – 6.2)	2.5% (4)	46.5 ± 9 (37 – 58)	32.4 ± 9.8 (25 – 46.8)	5.7 ± 0.4 (5.1 – 6.1)
Admixed American	22%** (24)	44.9 ± 12 (24 – 61)	29.6 ± 4.7 (22 – 38.7)	5.4 ± 0.4 (4.3 – 5.9)	38.5% (62)	39.6 ± 13 (15 – 62)	29.4 ± 4.8 (18.2 – 40)	5.4 ± 0.3 (4.5 – 6.1)
European	57% (61)	46.4 ± 10 (21 – 65)	28.9 ± 6.4 (17.6 – 45.5)	5.3 ± 0.4 (4.4 – 6.4)	55% (89)	46.0 ± 12 (18 – 68)	28.4 ± 5.0 (13.7 – 40.7)	5.3 ± 0.4 (4.0 – 6.3)
Combined	40% (107)	46 ± 11 (20 – 65)	29.4 ± 6.1 (17.6 – 47.6)	5.4 ± 0.42 (4.3 – 6.4)	60% (161)	44 ± 13 (15 – 68)	28.8 ± 5.0 (13.7 – 46.8)	5.4 ± 0.37 (4.0 – 6.3)

Values reported as mean ± SD (range)

G 16.7 2nd AUC was calculated by subtracting the G 16.7 1st AUC from the total G 16.7 AUC.

n = 115 female, *n* = 184 male

Figure 4