Chemostat culturing reduces fecal eukaryotic virus load and delays diarrhea after virome transplantation

Simone Margaard Offersen[1], Signe Adamberg[2], Malene Roed Spiegelhauer[1], Xiaotian Mao[3], Torben Sølbeck Rasmussen[3], Frej Larsen[3], Jingren Zhong[1], Duc Ninh Nguyen[1], Dennis Sandris Nielsen[3], Lise Aunsholt[1,4,5], Thomas Thymann[1], Kaarel Adamberg[2]*, Anders Brunse[1]*

[1]Department of Veterinary and Animal Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Frederiksberg, Denmark. [2]Department of Chemistry and Biotechnology, Tallinn University of Technology, Akadeemia tee 15, Tallinn, Estonia. [3]Section of Food Microbiology, Gut Health, and Fermentation, Department of Food Science, University of Copenhagen, Frederiksberg, Denmark. [4]Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. [5]Department of Neonatology, Copenhagen University Hospital, Rigshospitalet,Copenhagen, Denmark.

*Correspondence to: Kaarel Adamberg (kaarel.adamberg@taltech.ee) for *in vitro* experiments and Anders Brunse (anderss@sund.ku.dk) for *in vivo* experiments.

Abstract

Fecal virome transfer (FVT) shows promise in reducing necrotizing enterocolitis (NEC), likely due to donor bacteriophages preventing the gut dysbiosis preceding disease. However, concurrent transfer of eukaryotic viruses may carry a risk of infection for the recipient. To increase safety, we investigated chemostat propagation as a method to eliminate eukaryotic viruses from donor feces while maintaining a diverse and reproducible bacteriophage community. Donor feces was collected from healthy suckling piglets and inoculated into a fermenter containing growth media supplemented with lactose and milk oligosaccharides (MOs). During continuous medium exchange (20% volume/h), dilution significantly reduced eukaryotic viruses. Viral richness was concurrently reduced although still preserving a stable community of 200-250 bacteriophages. Inclusion of MOs in the medium ensured higher bacterial richness and a bacterial community closer resembling donor feces. Fecal *Lactobacillaceae* bacteria were lost during cultivation but partially replaced by members of the *Bacteroidota* phylum in MO-supplemented cultures, accompanied by phages predicted to have *Parabacteroides* as host. After cultivation, virus-like particles (VLPs) were isolated, and their ability to reduce NEC incidence tested *in vivo*. Preterm piglets were delivered by cesarean section and received either the lactose- or MO-propagated viromes by oral route (n = 14-15/group). These were compared with groups receiving the same dose of donor fecal virome (10^{10} VLPs/kg) or vehicle control. The piglets were subsequently fed infant formula for 96 hours followed by euthanasia and tissue sampling. Both chemostat-propagated viromes effectively mitigated diarrhea compared to the donor virome. The donor virome partially engrafted in recipients and led to higher levels of *Lactobacillaceae* bacteria and *Lactobacillaceae* targeting phages. However, these signatures were lost in recipients of chemostat-propagated viromes, and only minor microbiome effects and no NEC prevention were observed. To conclude, we provide *in vivo* proof-of-concept for chemostat propagation of fecal viruses as a means to deplete eukaryotic viruses and in turn reduce side effects in newborn virome recipients. However, chemostat culture conditions need further optimization to preserve the donor phageome.

Keywords: Fecal Virome Transplantation, Phage Therapy, Bacteriophages, Eukaryotic viruses, Chemostat*,* Fermentation, Preterm, Necrotizing Enterocolitis, Diarrhea.

Introduction

Immediately after birth, microorganisms colonize the gastrointestinal tract, thereby establishing the gut microbiome (GM). Non-digestible milk oligosaccharides (MOs) present in breast milk strongly influence GM assembly by acting as highly specific microbial substrates to promote the growth of bifidobacteria and *Bacteroides*[1]. Preterm birth (< 37 weeks of gestation) is often accompanied by GM perturbations due to factors such as cesarean section, early antibiotic use, immature gastrointestinal tract, and partial formula feeding[2–5]. These disturbances inhibit beneficial anaerobes like bifidobacteria and *Bacteroides* whereby facultative pathobionts like *Enterobacteriaceae* members may take over and initiate disease processes[6]. This can ultimately lead to the severe intestinal disorder necrotizing enterocolitis (NEC), characterized by inflammation and necrosis of intestinal tissue[7]. With global mortality rates ranging from 20-50%, NEC is considered a critical medical and surgical emergency[8].

In recent years, there has been growing interest in studying the gut virome alongside gut bacteria, particularly in the context of severe inflammatory gut disorders like NEC and inflammatory bowel disease (IBD). Specifically, bacteriophages (phages for short) are abundant gut viruses that function to modulate bacterial growth through host-specific lysis or lysogeny[9]. This dynamic interaction and additional preference for intestinal mucosal sites enable phages to alter bacterial compositions and protect the host from bacterial invasion[10–12]. Several studies have associated virome alterations with IBD[13–15] and recently, the link was discovered in a small cohort of NEC patients as well[16]. Thus, the virome may be a novel target for modulating the early life GM and preventing intestinal inflammation in neonates.

Fecal virome transfer (FVT) shows potential for restoring gut health by introducing viruses from donor feces to patients of gut dysbiosis. A recent preclinical study demonstrates that FVT can modify the gut virome of preterm piglets, reducing mucosa-associated *Enterobacteriaceae* levels and providing significant protection against NEC[17]. However, during FVT, fecal eukaryotic viruses are transferred alongside phages, introducing a risk of infections. While screening programs exclude fecal donors with known viral pathogens, many eukaryotic viruses in the gut have unknown functions that could affect neonatal health, potentially leading to both immediate and long-term consequences[18,19]. A follow-up study using the piglet model found that, while a prescreened FVT indeed offered protection against NEC, it was also linked to adverse events, namely an earlier onset of diarrhea, which also led to significant weight loss[20].

Fecal phages and eukaryotic viruses are highly similar in size and structure[21]. As a result, phages

will be damaged if traditional methods for eliminating eukaryotic viruses, such as chemicals or

irradiation, are applied to FVT preparations. However, both types of viruses rely on host cells for

propagation but have very distinct host specificities (bacterial vs. mammalian cells). This distinction

can be leveraged in a *chemostat*: a bioreactor system designed to cultivate microorganisms in a

controlled environment[22–24]. Here, a continuous fecal culture is maintained by continually adding

fresh medium, providing nutrients for bacterial growth, and thus continuously generating hosts for

phage propagation. Concurrently, equivalent volumes of spent media containing microorganisms

and metabolic by-products are removed, leading to a gradual dilution of eukaryotic viruses, whose

host cells are unable to propagate in the chemostat[24,25]. Thus, the method has the potential to create

a steady-state fecal culture from which the phage communities can be extracted and used for

FVT[26,27].

Insights from traditional fecal microbiota transplantations (FMT) for IBD in humans or

experimental NEC suggest that treatment efficacy can vary depending on the donor, a phenomenon

that likely applies to FVT as well[28–30]. In the previous NEC studies, feces from suckling term-born

pigs were used as FVT donor material to best match the preterm piglet recipients[17,20]. Although this

donor-recipient matching approach seems promising, the use of young donors limits the amount of

fecal matter available for donation. The chemostat method overcomes these limitations by

generating a mass-reproducible fecal phage community, potentially providing both a safer and

donor-independent FVT strategy.

To propagate bacteria and phages from young donors, the medium composition should replicate the

nutrients reaching the gut during nursing. Thus, we tested if adding medium MOs would increase

the microbial similarity between the donor feces and chemostat culture. Subsequently, native fecal

virome transfer (FVT) was compared with the chemostat virome transfer propagated without MOs

(CVT) and with MOs (CVT-MO) in the preterm piglet model. We hypothesized that chemostat-

propagated viromes could modulate the GM and reduce NEC lesions while avoiding potential side

effects, such as earlier onset of diarrhea.

Materials and methods

Donor feces collection

111 Gut content was collected from four 10-day-old healthy-appearing piglets from four different

112 conventional farms. After euthanasia, the cecum, colon, and rectum were isolated, and the content

113 was emptied and processed under anaerobic conditions. After diluting 1:1 in 2 × SM buffer (400

114 mM NaCl, 20 mM MgSO4, 100 mM Tris-HCl, pH 7.5) containing 20% sterile glycerol, the content

115 from each pig was homogenized, divided into smaller fecal aliquots, and stored at -70°C.

Chemostat cultivation

Growth medium

A rich tryptone and yeast extract based medium supplemented with mineral salts and vitamins was

prepared as described by Pham et al.[23] with the following modifications: lactalbumin concentration

6 g/L and 0.05 M potassium phosphate buffer were used. Additionally, carbohydrate content varied

as follows: In one setup, a mixture of MOs was added (7 g/L, LAC-MO medium) along with the

original content of lactose (6.4 g/L) and mucin (4 g/L porcine gastric type II mucin, Sigma-Aldrich,

Germany). The total MO level was adjusted to quantities reported in mature porcine milk[31]. This

consisted of 3.5 g/L 3'-sialyllactose (3'SL), 2.8 g/L lacto-N-neotetraose (LNnT), and 0.7 g/L 2'-

fucosyllactose (2'FL), chosen to match porcine milk levels of sialylated-, neutral-, and fucosylated

MOs, respectively[32]. All MOs were donated by DSM-Firmenich (Switzerland). In another setup,

the MO fraction was substituted with lactose, balanced by equimolar levels of C6 monosaccharides.

This resulted in 12.5 g/L lactose (LAC medium) in addition to the 4 g/L mucin.

Cultivation system and culture conditions

The previously described chemostat system[22,24] was used for cultivation of the collected pig feces.

Briefly, the cultivation system consisted of an anaerobic fermenter equipped with sensors for pH,

pO2, and temperature control. The culture volume (V) was kept constant (800 mL) as well as the

temperature (39°C)[33] and pH (6.5). The pH was controlled by adding 3 M NaOH according to the

setpoint. Variable speed pumps enabled a controlled inflow and outflow of medium during

continuous cultivation. The collected fecal aliquots were thawed and processed under anaerobic

conditions where all four donors were pooled in equal amounts and further diluted 1:1 in sterile

0.9% saline (resulting in four times diluted feces). To start the experiments, 4 mL of diluted fecal

pool was inoculated into the fermenter (0.125% w/v raw feces in fermenter volume) and allowed

growth for 24 hours in a fixed medium volume, which corresponds to the end of the exponential

growth phase of the fecal culture (batch phase). Hereafter, continuous medium inflow/outflow was

started (chemostat phase) at a feeding rate (F) of 160 mL per h, resulting in a dilution rate (D) of 0.2

h^{-1} (D=F/V). This phase continued until 75 h, during which the chemostat volume was exchanged

10 times (Fig. 1A). Two and four replicates were carried out with LAC and LAC-MO medium,

respectively. Samples from the outflow were collected on ice three times during the batch phase and

four times during the chemostat phase as shown in Fig. 1A. On-line and at-line parameters used for

experiment control are depicted in Supplementary Fig. S1.

Gas, redox, biomass, and chromatographic analyses

The composition of the gas outflow, including H□, CO□, H□S, CH□, and N□, was evaluated

using an Agilent 490 Micro GC Biogas Analyzer (Agilent 269 Technologies Ltd., USA) connected

to a thermal conductivity detector. Gas flow volume was monitored regularly using a

MilliGascounter (RITTER Apparatebau GMBH & Co, Germany). The redox potential of both the

growth medium and culture supernatant was measured with a pH/Redox meter and an InLab®

Redox electrode (Mettler Toledo, USA). Biomass dry weight was assessed gravimetrically from a

10 mL culture by centrifuging at 6,000 rpm for 20 minutes, followed by washing the biomass with

distilled water and drying it at 105°C for 20 h in an oven.[24]

Chromatographic analyses

Culture samples were centrifuged (11,000 × g for 5 min at 4°C), then stored at -20°C. After

thawing, the supernatants were filtered using Amicon® Ultra-10K Centrifugal Filter Devices with a

3 kDa cut-off, following the manufacturer's instructions (Millipore, USA). Organic acids

(succinate, lactate, formate, acetate, propionate, isobutyrate, butyrate, isovalerate, and valerate) and

ethanol concentrations were measured via high-performance liquid chromatography (HPLC) using

the Waters Alliance 2795 system (Waters, Milford, MA, USA) with a BioRad HPX-87H column

(Hercules, CA, USA). Isocratic elution with 0.005 M H□SO□ was performed at a flow rate of 0.5

mL/min at 35°C, with refractive index (RI) (model 2414) and UV (210 nm; model 2487) detectors

(Waters, USA). Quantification was done using analytical-grade standards, with a detection limit of

0.1 mM.[24]

Pre-processing of samples for separation of bacterial and viral analysis

Fecal inoculum and culture samples were analyzed to investigate microbiome changes over time.

Separation of the viruses and bacteria was done as illustrated in Fig. 1B. Fecal inoculum was diluted

in SM buffer (200 mM NaCl, 10 mM MgSO4, 50 mM Tris-HCl, pH 7.5) until 1:20 and centrifuged

(4,500 × g, 30 min, 4°C). For chemostats, 10 mL cooled outflow culture were sampled per time

point and immediately centrifuged (4,500 × g, 30 min, 4°C). The pellet was stored at -80°C until

bacterial analysis and the supernatant was filtered through a 0.45 µm Filtropur S PES syringe filter

(Sarstedt, Helsingborg, Sweden). Subsequently, the filtrate was concentrated using Vivaspin 20

units with a 300 kDa molecular weight cut-off (MWCO) PES membrane (Sartorius, Göettingen,

Germany), which was centrifuged ($2,500 \times g$, 4°C) until all filtrate volume had passed. Hereafter, 2

mL SM buffer was added and stored at 4°C overnight to release viruses from the membrane. The

concentrated virus samples were stored at -80°C until analysis.

Processing of chemostat culture and feces for separation of viruses

After 10 volume exchanges, the total fermenter culture (800 mL) was drained into a sterile 1,000

mL bottle on ice. Thereafter, the culture was centrifuged ($7,000 \times g$, 60 min, 4°C) and the

supernatant was filtered to remove bacteria using 0.45 µm Filtropour V50 PES vacuum filtration

units (Sarstedt, Helsingborg, Sweden). To concentrate viruses and remove residual medium

components, including MOs, the filtrate was ultracentrifuged following a protocol shown to retain

infective viruses[34]. To further reduce non-viral particle contamination, we employed a higher

MWCO of 300 kDa, compared to the original protocol. Thus, the filtrates were added to Vivaspin

20 units (300 kDa MWCO, PES; Sartorius, Göettingen, Germany) and centrifuged ($6,000 \times g$ at 25°

fixed angle, 4°C). Each unit had 80 mL repeatedly added and concentrated to 0.5-1.0 mL. Hereafter,

SM buffer was added to 10 mL total and stored at 4°C overnight. The eight times concentrated

culture virome from all units were pooled, then aliquoted and stored at -70°C until the piglet

experiment. This procedure was repeated for each chemostat replicate.

Fecal virome transfer (FVT) solution was prepared similarly by pooling the four donors and further

diluting 1:5 in SM buffer. The fecal viromes were isolated as described above (Fig. 1B). After

concentration with Vivaspin 20 units, SM buffer was added to the original volume (1:10 diluted).

Aliquots of FVT and sterile SM buffer were stored at -80°C until use.

Fluorescence microscopy and bacterial cultivation

Epifluorescence microscopy was used to determine the concentration of virus-like particles (VLPs)

in each of the end culture viromes and the FVT. VLPs were stained with SYBR Gold (Cat. no

S11494, Invitrogen, ThermoFisher Scientific) and captured on 0.02 µm aluminum oxide filters

(Whatman™ 6809-6002 Anodisc™, Cytiva, USA). One filter was prepared for viromes from each

chemostat run and the fecal inoculum. The filters were photographed (Photometrics CoolSNAP

camera) using an epifluorescence microscope fitted with a 490 nm filter block. Hereafter, VLPs

were counted and the number per mL was estimated as described previously[20]. To confirm the

absence of living bacteria in each virome solution, 100 uL from aliquots of CVT, CVT-MO, FVT,

and SM buffer (control, CON) was spread on plate count agar. The plates were incubated for 24

hours under both aerobic and anaerobic conditions for 24 h before inspection for bacterial growth.

A plate with no added aliquots and a plate with 100 uL of fecal slurry were included as negative and

positive control, respectively.

Cell stimulation

The human monocyte cell line THP-1 was used to investigate the innate immune response

following challenge with the fecal and chemostat viromes compared with a cocktail of 10 isolated

virulent phages, and an *Escherichia coli* bacterium. A fecal virome transfer (FVT) aliquot was

thawed and used directly. Chemostat virome transfer (CVT) solution consisted of pooled viromes

from both chemostats propagated with LAC medium (replicas 1 and 2). Similarly, chemostat

virome transfer propagated with MOs (CVT-MO) comprised pooled viromes from two of the four

chemostats propagated with LAC-MO medium (replicas 1 and 3). Each fecal or chemostat solution

(FVT, CVT, CVT-MO) was ultracentrifuged (300 kDa MWCO) and adjusted to 5×10^9 VLPs per

mL. The phage solutions were mixed to reach a concentration of 5×10^9 plaque-forming units

(PFU) per mL, as described by Larsen et al.[34]. Details of the phages are listed in Supplementary

Table S1. The *E. coli* strain (ST-2064) was cultured in a liquid Luria-Bertani medium (BD Difco™

LB Broth Lennox, Becton Dickinson, USA) until it reached the exponential phase and adjusted to a

concentration of 10^8 colony forming units (CFU) per mL. Before the challenge, endotoxins were

removed from all virome, phage, and bacterial preparations using Pierce™ High Capacity

Endotoxin Removal Spin Columns (0.5 mL, ThermoFisher Scientific, USA) by following the

manufacturer's instructions. The THP-1 cells were cultivated, stimulated, and RNA was extracted

and RT-qPCR performed exactly as described previously[20]. In brief, THP-1 cells were differentiated

into macrophages and stimulated with 100 µL of FVT, CVT, CVT-MO, phage solution (Phages), or

E. coli solution. RNA was extracted after incubation for 20 h (37°C, 5% CO_2). Seeding and

challenges were performed in triplicates at cell passages 17, 19, and 20.

Piglet Experiment 1

Experimental procedures conducted during the animal experiments were approved by the Danish

Animal Experiments Inspectorate (license number: 2020-15-0201-0052), under the guidelines from

Directive 2010/63/EU of the European Parliament. 60 piglets were delivered preterm from three

sows by cesarean section (day 106, 90% gestation) as described previously [35]. After delivery, the

pigs were transferred to heated incubators and observed or ventilated until the respiration stabilized.

Hereafter, the pigs were fitted with oral catheters for enteral feeding and umbilical arterial catheters

for parenteral nutrition (PN) and access to blood circulation (6 and 4 Fr size, respectively, Portex,

Kent, UK). The pigs were stratified by sex and birth weight and randomly allocated to four groups

receiving fecal virome transfer (FVT), chemostat virome propagated with LAC medium (CVT),

chemostat virome propagated with LACMO medium (CVT-MO), or SM buffer (control, CON).

Immunization and feeding procedures

To substitute colostrum feeding, all pigs were immunized with sows' plasma (16 mL/kg) by

intraarterial infusion over two hours on the first day of life. Hereafter, the pigs received declining

levels of PN over the study period (96-48 mL/kg$^{0.75}$/d, Kabiven, Vamin, Fresenius-Kabi; Bad

Homburg, Germany). Enteral nutrition (EN) was fed every three hours and consisted of a home-

mixed formula (formula 1) as described previously[20]. EN volumes increased gradually, reaching 24,

40, 64, and 96 mL/kg$^{0.75}$/d on days 1, 2, 3, and 4– 5, respectively.

Fecal- and chemostat virome transfers

CVT and CVT-MO were produced as described for the cell stimulation assay above. Just before

use, the treatments were thawed and then administered orally via the orogastric tube. Each virome

treatment (FVT, CVT, CVT-MO) was adjusted to 10^{10} VLP/kg in a total volume of 2-3 mL/kg SM

buffer. The CON group received 2 mL/kg of SM buffer. All groups received their treatment volume

divided into administrations twice daily on days 1 and 2 (Fig. 4A).

In vivo procedures and euthanasia

The pigs were closely monitored by experienced personnel and pigs showing clinical signs of NEC

(abdominal distention, lethargy, or severe bloody diarrhea) were immediately euthanized. Daily

weights and exact times of diarrhea episodes were recorded. On days 3 and 5, blood samples were

obtained for routine hematological profiling. On day 5, all pigs were anesthesized followed by

euthanasia with an intracardiac injection of pentobarbital. As a marker for intestinal permeability,

the urinary lactulose-to-mannitol ratio was measured as previously described[36].

NEC evaluation and tissue sampling

The abdominal organs were excised, separated, and weighed. The gastrointestinal tract was assessed

for NEC-like lesions as previously described[37]. Briefly, levels of hyperemia, hemorrhage,

pneumatosis intestinalis (intramural gas), and necrosis were assessed by a blinded investigator and

converted to a 16-point NEC score. NEC was defined as the presence of hemorrhage, pneumatosis

intestinalis, and/or necrosis. Colon content, two samples from the proximal intestine, as well as

mesenteric lymph nodes from the ileocecal region were snap-frozen for later analysis. The most

severe lesions from the small intestine and colon were both snap-frozen and fixed in 4%

paraformaldehyde. The fixed tissue was later paraffin-embedded, sectioned, and stained with

hematoxylin and eosin. A blinded pathologist assessed each section by microscopy and assigned an

established histopathological score ranging from normal appearance (score 0) to transmural necrosis

(score 8)[37]. Lactase activity was measured in proximal small intestinal tissue by a previously

described colorimetric method[36]. Interleukin (IL) 1β and IL-8 were measured in tissue homogenates

from proximal small intestine and colon lesions by commercial porcine ELISA kits (R&D Systems,

158 Abingdon, Oxfordshire, United Kingdom) according to the manufacturer's instructions[38].

RNA sequencing of mesenteric lymph nodes

Mesenteric lymph node RNA was extracted from 20 mg homogenized tissue (GentleMACS

Dissociator, Miltenyi Biotec, Bergisch Gladbach, Germany) using the RNeasy Mini Kit (Qiagen,

Hilden, Germany) following the protocol for tissue, and with on-column DNase digestion (RNase-

free DNase, Qiagen, Hilden, Germany). Whole-transcriptome shotgun sequencing was performed

by Novogene services (Cambridge, UK) and analyzed as described previously[39]. Briefly, RNA-Seq

libraries were constructed using Novogene NGS RNA Library Prep Set (PT042) and sequenced

using the Illumina NovaSeq 6000 S4 flowcell platform. Fastp software was utilized for trimming

the raw reads to remove adapters and low-quality bases. The raw sequencing data were saved in

FASTQ format, containing both the read sequences and their associated base quality scores.

HISAT2 was employed to align the reads to the porcine reference genome (*Sscrofa11.1*).

FeatureCounts package was then used to create a gene count matrix, which was further analyzed for

differentially expressed genes (DEGs) with DESeq2, using *P*-value cutoff 0.05 and log2 fold

change cutoff 1. The pathway enrichment analysis was performed with a gene set enrichment

analysis (GSEA) through the clusterProfiler package using the Gene Ontology (GO) classification

system for *Sus scrofa*. The false-discovery-rate (FDR) cut-off of 0.05 was applied to identify

significant pathways. A normalized enrichment score (NES) was obtained, with either negative or

positive values indicating the pathways being down- or up-regulated. The enriched pathways were

visualized using ggplot2.

Piglet Experiment 2

53 piglets delivered from two sows were stratified by sex and birth weight and randomly allocated

to two groups receiving CVT-MO solution or SM buffer (CON). The enteral nutrition (EN)

consisted of an ultra-high temperature (UHT) treated ready-to-drink formula (formula 2), previously
reported to have a high capacity for inducing NEC lesions [37]. All other experimental procedures
were exactly as described for *Piglet Experiment 1*.

Bacterial DNA extraction, sequencing, and pre-processing of raw data

The DNAeasy PowerSoil Pro Kit (Qiagen) was used to extract bacterial DNA from the fecal
bacterial pellet and the chemostat aliquot following the manufacturer's instructions. The final
purified DNA concentration was determined on Varioskan Flash (Thermo Fisher Scientific, USA)
using Qubit™ 1x dsDNA high sensitivity kit (Invitrogen). The purified DNA products were stored
at -80°C for later use. The full-length 16S rRNA gene was used to determine the gut bacterial
composition. The 16S rRNA gene was amplified using several forward and reverse primers (Suppl.
Table S2). The first PCR reaction included mixing of 12 μL PCRBIO Ultra Mix (PCRBIO), 6 μL
Sterile Milli Q H2O, 2 μL primer mix, and 5 μL genomic DNA (~5 ng/μL). The PCR cycling was
run on SureCycler 8800 (Agilent Technologies). The first PCR cycles were as follows: 95°C for
2 min; 2 cycles of 95°C for 20 sec, 48°C for 30 sec, 65°C for 10 sec, and 72°C for 45 sec;
followed by a final step at 72°C for 4 min. The second PCR step was done by barcoding the 11 μL
of the first PCR products with 2 μL barcodes in 12 μL PCRBIO Ultra Mix (PCRBIO): 95°C for
2 min; 33 cycles of 95°C for 20 sec, 55°C for 30 sec, and 72°C for 40 sec; followed by a final step
at 72°C for 4 min. The gut bacterial composition was assessed using the Oxford Nanopore
Technologies (ONT) GridION platform. ONT MinKNOW software v22.10.7 collected the data
generated by GridION. The Guppy v6.2.8 base-calling toolkit was used to convert the the raw
FAST4 files to FASTQ format. The Long Amplicon Consensus Analysis (LACA) pipeline
generated the abundance table through raw FASTQ files (https://github.com/yanhui09/laca). The
assigned taxonomy quality was corrected using SILVA database[40].

Viral RNA/DNA extraction, sequencing, and pre-processing of raw data

The viral particle separation was conducted based on an existing protocol[41]. The intestinal content
from pig and chemostat aliquots was suspended in 5 mL autoclaved SM buffer (100 mM NaCl,
8 mM MgSO4·7H2O, 50 mM Tris-HCl with pH 7.5) followed by centrifuging at 4500 x *g* for 30
min at 4 °C. The supernatant was filtered using a 0.45 μm Minisart High Flow PES syringe filter
(Sartorius) to remove bacteria and other larger particles. VLPs in the filtered supernatants were
concentrated and smaller metabolites removed by ultrafiltration using a Centrisart ultrafiltration
devices with a filter cutoff at 100 kDa (Sartorius, UK). To remove free DNA/RNA molecules, 140
μL supernatants were treated with 5 units of Pierce Universal Nuclease (ThermoFisher Scientific)

for 15 min at room temperature. Viral DNA/RNA was extracted with Viral RNA mini kit (Qiagen).
Reverse transcription was conducted using SuperScript VILO Master (Thermo Fisher Scientific)
and subsequently cleaned with DNeasy blood and tissue kit (Qiagen) by following steps 3-8 in the
manufacturer's manual. Multiple displacement amplification was performed to include ssDNA
viruses using the GenomoPhi V3 DNA amplification kit (Cytiva). The library preparation for
sequencing was conducted by using Nextera XT kit (Illumina) as previously described[26]. The
pooled purified PCR product was sequenced on a NovaSeq X plus platform by Novagene. Raw
reads were trimmed (trimmomatic v0.38), deduplicated (seqkit v0.13.2) and assembled (spades
v3.13.0)[42–44]. Assembled contigs were deduplicated at species level (~95% sequence identity) and
non-viral contigs were removed by running geNomad[45] and checkV[46] and selecting contigs that
were either 1) marked "High-quality" or "Complete" by checkV, or 2) had a geNomad score of
>0.9 and >2 hallmark genes. Additionally, all viral contigs should be above 2200 bp, or 10.000 bp if
the geNomad-predicted taxonomies were caudoviral. Viral hosts were predicted using iPHoP[47] and
taxonomies were set by aligning to a custom database. Protein prediction was done with phold
(https://github.com/gbouras13/phold), which was used to predict whether a phage was temperate or
not based on the presence of recombinase/integrase genes. Reads from each sample were mapped to
the viral contigs by msamtools (v1.1.2). The pipeline is described in detail at
https://github.com/frejlarsen/vapline3.

Bioinformatic analysis of bacterial and viral sequences

R version 4.3.0 was used for analysis and presentation of microbiome data. The main packages used
were phyloseq[48], vegan[49], DESeq2[50], ampvis2[51], ggpubr, psych, EnhancedVolcano, and ggplot[48].
The R package microDecon (runs=1, regressions=1) was used to remove the contamination of viral
contigs by read counts in negative controls[35], and 83.0% were kept after removal. For samples in
the chemostat experiment, the media used in the chemostat propagation was used as a control to
remove the viral contigs introduced by the media, using microDecon[52] (runs=1, regressions=1),
leaving 84.2% contigs for analysis of chemostat samples. Cumulative sum scaling normalization
was performed using R metagenomicSeq package. The Bray-Curtis dissimilarity metric was used
for beta diversity analysis and with statistics based on one-sided pairwise PERMANOVA corrected
by FDR. The differential bacteria and viruses were identified by DESeq2 on the summarized
bacterial species level and viral contigs (vOTUs) level based on two-sided Wald test. Alpha
diversity and shared vOTUs were analyzed with the non-parametric two-sided Wilcoxon rank-sum
tests with FDR correction. Eukaryotic viral contigs were manually identified in the vOTU table and

the relative abundance of these contigs was calculated based on total eukaryotic viral taxa and
subsequently visualized in heatmaps.

Statistical analysis of remaining data

Remaining statistical analyses were performed in R version 4.2.2. Survival and diarrhea onset were
analyzed with pairwise Log-Rank tests and Benjamini-Hochberg (BH) adjustment. Continuous
outcomes were analyzed with linear mixed models with group, sex, and birth weight as fixed effects
and litter as a random effect. The group effect was assessed with an ANOVA, followed by Tukey's
test. To ensure the validity of the modeled data, the residuals and fitted values were evaluated for
normality and homogeneity of variance. For some outcomes, this necessitated log-transforming
before modeling. Tissue cytokines did not meet parametric criteria and were analyzed with the
Kruskall-Wallis test followed by Dunn's test with BH correction. Gross pathology and microscopic
lesion scores were analyzed using a proportional ordered logistic regression model (R package
MASS). The analysis included group, sex, litter, and birth weight as fixed effects. Group effects
were assessed using the likelihood ratio test with a BH correction. Incidences of pathologies were
compared with Fisher's exact tests. A repeated measures linear mixed model was employed to
analyze chemostat metabolite levels, with fixed effects for group, time, and their interaction, and a
random intercept for replicate ID. Piglet repeated measures were analyzed using a linear mixed
model, with fixed effects for group, day, their interaction, and covariates (sex, litter, birth weight).
To account for the correlation and variance heterogeneity over time, this model assumed an
unstructured covariance pattern (R package LMMstar).

Results

390   With this study, we aimed to provide a combined *in vitro* and *in vivo* proof-of-concept for phage
391   preservation and concomitant eukaryotic virus reduction through chemostat propagation of fecal
392   matter and subsequent virome transfer into newborn recipients to alleviate NEC and virus-
393   associated diarrhea. Further, we measured the effect of supplementing the chemostat cultures with
394   MOs for better approximation of a milk-shaped fecal inoculum.

Milk oligosaccharides preserved the bacterial richness of a stable fecal chemostat culture

A mix of anaerobically collected fecal samples from four healthy suckling pigs was inoculated into
batch cultures with or without an MO blend representing the major MOs in porcine milk to
supplement lactose as the major carbohydrate source[32,53,54]. Continuous exchange of medium was
initiated after 24 hours of incubation and a constant target pH of 6.5 (mean 6.55 ± 0.04 at 50-75

hours for all chemostats) was maintained by base titration. To check the stability of the chemostat

cultures, base titration curves were inspected (Suppl. Fig. S2). In the batch phase, lactose

conversion to lactic acid resulted in a peak titration at 4-5 hours with a higher peak in the pure

lactose (LAC) cultures. A later pattern of small but prolonged titration occurred only in the MO

(LAC-MO) supplemented cultures, indicating a slower utilization of MOs. After initiation of

continuous volume exchange, titration patterns quickly stabilized with marginally higher acid

production/base consumption in the LAC cultures.

We assessed the bacterial composition of the cultures at seven time points throughout the

experiment to assess culture dynamics (Fig. 1A-B). Whereas all cultures lost a significant

proportion of bacterial richness already in the early batch phase, supplementing the growth medium

with MOs preserved the richness of the culture throughout the following chemostat phase to a

greater extent than pure lactose (Fig. 1C). Taxonomically, donor feces showed >50% relative

abundance of bacteria from the *Lactobacillaceae* family. During the early batch phase, *Escherichia*

coli expanded rapidly, contributing to the fast lactose consumption and acid production. Hereafter,

both *Lactobacillaceae* and *E. coli* were replaced mainly by *Streptococcus* and *Fusobacterium* spp.

Importantly, in the chemostat phase a significant proportion of these were replaced by different

species of the *Bacteroidota* phylum only in the MO cultures indicating that these taxa had a

competitive advantage due to MO utilization capabilities (Fig. 1D). Consequently, in the MO-

supplemented end culture we found a relative enrichment of several taxa, all of which were obligate

anaerobes (Fig. 1E).

Compositional analysis using Bray-Curtis ordination revealed an abrupt change in the early batch

phase, with a partial restoration towards the composition of the fecal inoculum occurring throughout

the chemostat phase visualized along the second principal coordinate. Notably, MO

supplementation augmented this restoration. After five volume exchanges, the bacterial composition

had stabilized and did not develop further for the remainder of the experiment. However, using the

current fermentation conditions including a relatively high dilution rate, the chemostat end culture

did not fully approximate the composition of the fecal inoculum (Fig. 1F).

Metabolically, the culture dynamics throughout the experiment corroborated the bacterial

compositional analysis. As such, high production of lactate, formate, ethanol, succinate, H_2, and

CO_2 in the early batch phase indicated mixed-acid fermentation mainly by *E. coli* (Fig. 2). As the

cultures entered the chemostat phase, lactate was gradually consumed, and short-chain fatty acid

levels (e.g. propionate, butyrate) increased. In the stable end cultures, butyrate levels were equally

high, but importantly, propionate was a major fermentation product only in the MO cultures. Other short-chain fatty acids (e.g. valerate, isovalerate) were also increased by MOs despite lower concentrations. Taken together, the increased proportion of obligate anaerobes in the MO-supplemented chemostat cultures greatly increased the production of propionate and other short-chain fatty acids.

Figure 1. MO supplementation preserves bacterial richness and promotes *Bacteroidota* enrichment in stabilized chemostat cultures. A) The experimental setup of chemostat cultivations of donor feces with

lactose medium (LAC, *n*=2) or lactose and milk oligosaccharide medium (LAC-MO, *n*=4). Donor feces was inoculated into a fermenter followed by batch growth for 24 h. The chemostat culture was then run until 10 volume exchanges. The sampling points for metabolite and microbiome analyses are indicated. B) Processing of chemostat culture to separate virus-like particles from bacteria and the culture residuals. C) Number of observed bacterial species. D) Relative bacterial abundance per replicate summarized at species level. Taxonomic rank specifications: p = phylum, f = family, g = genus. E) Volcano plot visualizing bacterial DESeq2 enrichment analysis with FDR correction based on 16S rRNA gene amplicon sequencing on species level. Horizontal dotted lines show a significance level of *P* < 0.05 and vertical lines show ± 0.6 log2 fold change (FC) cut-off. NS = non-significant. F) Principal coordinate analysis (PCoA) plot visualizing bacterial beta diversity over time, based on Bray-Curtis dissimilarity metrics.

Figure 2. Metabolic transition from mixed-acid to SCFA fermentation enhanced by MO supplementation. Metabolite production in cultures of piglet fecal microbiota during batch phase (0-24 h) and chemostat phase (24-75 h) between replicas (R) of cultivations with lactose (LAC) and lactose-milk oligo-saccharide (LAC-MO) medium. Statistical significance is calculated between LAC-MO and LAC experiments.

Fecal chemostat cultivation established a stable phage community devoid of eukaryotic viruses

When bacteria grow in a bioreactor inoculated with fecal matter, they engage in a dynamic interaction with the vast number of both virulent and temperate fecal phages[55]. To investigate the viral community fluctuations during fermentation, we isolated virus-like particles from the cultures (Fig. 1B) at each sample point followed by metagenomics sequencing. Relative to the fecal inoculum, the viral richnes decreased by 50% during the batch phase and further decreased during the chemostat phase, resulting in an overall decrease to 25% of the initial level (Fig 3A). The dilution occuring at inoculation (1:800 w/v) may explain the initial loss, as it might reduce low-abundant viruses to levels below the sequencing depth. In the chemostat phase, most phages with non- or slow-growing bacterial hosts were likely reduced alongside the eukaryotic viral elimination.

This resulted in a final conserved community of around 200-250 viruses with similar levels for pure

lactose cultures and MO-supplemented cultures. The community compositions changed

dramatically during fermentation until reaching a stable level from five volume exchanges and

onwards with high similarity between replicates (50-75 h, Fig. 3B), closely resembling the bacterial

dynamics although without an obvious MO effect. When examining relative abundances, the

majority of viruses were either classified as *Caudoviricetes* (class of tailed phages), *Microviridae*

(family of small, non-tailed phages), or unknown viruses (Suppl. Fig. S3A). When predicting

bacterial hosts, the fecal inoculum mainly harbored viruses infecting unknown bacteria (Fig. 3C).

Phages targeting *Escherichia* began to increase after 8 hours, suggesting the induction of coli-

phages in response to the *Escherichia* expansion seen at 3 hours (Fig. 1D). In the chemostat phase,

phages targeting *Bacteroides* and *Parabacteroides* started to emerge. These communities were best

maintained during MO supplementation, where especially *Parabacteroides* phages reached

relatively higher levels in end cultures. Interestingly, lifestyle prediction showed that the fecal

inoculum mainly consisted of potential virulent phages, whereas the relative abundance of

temperate phages steadily increased over time in the chemostat with no obvious effect of MO

supplementation (Fig. 3D). Consequently, the transferred phages differed in terms of lifestyle

between FVT and chemostat-propagated virome. DESeq2 analysis after 10 volume exchanges (Fig.

3E) revealed that adding MOs generally entailed a larger upregulation of specific viral OTUs

compared to lactose only (134 vs. 27 vOTUs, respectively).

To uncover the success of eukaryotic viral elimination, eukaryotic vOTU members were manually

identified and assessed in relative abundances over time (Fig. 3F). A total of five eukaryotic viral

families were present in the fecal inoculum and batch phases. After five volume exchanges (50 h),

four of these families were successfully eliminated, however, *Dicistroviridae* gradually increased

during chemostat mode. Conflicting with our dilution hypothesis, we subsequently subtracted all

viral contigs present in the medium and repeated the analysis. This caused   *Dicistroviridae* to

disappear, suggesting that this particular virus was continuously introduced via the medium rather

than propagation. Hence, viral contigs present in the medium were deducted prior to all the *in vitro*

viral analyses presented (Fig. 3). Conclusively, although less diverse than the native fecal virome,

both chemostat settings created stable cultures of phages with an undetectable level of known

eukaryotic viruses. The pronounced level of *Bacteroidota*-specific phages in MO-supplemented

cultures revealed substrate-driven differences in phage host-specificities, worthy of further

investigation.

498
499
Figure 3. Chemostat cultutivation with lactose medium (LAC) or lactose-milk oligosaccharide medium (LAC-
MO) yields selective but stable phage communities while depleting eukaryotic viruses. A) Number of
observed viruses. B) Principal component analysis (PCoA) plot visualizing viral beta diversity over time,
based on Bray-Curtis dissimilarity metrics. C) Relative abundance of viral bacterial hosts per replicate
summarized at genus level. D) Predicted lifestyle of bacteriophages based on presence (temperate) or
absence (virulent) of integrase and/or recombinase genes. E) Vulcano plot visualizing viral OTU differences
between substrates at 75 hours. The DESeq2 enrichment analysis with FDR correction was based on viral

metagenomics sequencing. Horizontal dotted lines show significance level of $P < 0.05$ and vertical lines
show ± 0.6 log2 fold change (FC) cut-off. NS = non-significant. F) Depletion of eukaryotic viral families
presented as heatmaps. The upper panel shows a raw analysis of culture samples. The lower panel shows
the analysis, where the baseline viral content in the medium was deducted from culture samples. Color
intensity indicates the relative abundance of specific eukaryotic viral contigs from total viral eukaryotic
contigs.

Fecal and chemostat-propagated viromes exhibit similar immunogenic potential

Virus-like particles (VLPs) were isolated from the feces (fecal virome transfer, FVT) and the 75-

hour chemostats. CVT comprised of VLPs isolated from chemostats with lactose medium, and

CVT-MO comprised of VLPs isolated from chemostats with lactose and MO medium (Fig. 4A).

Any medium residuals including MOs were removed during this procedure (Fig. 1B). Chemostats

with MOs produced end cultures with similar levels of VLPs compared to propagation with lactose

(1.1-1.7×10^9 and 1.1-1.2×10^9 VLPs/mL, respectively). These liquid cultures had a more than 10-

fold lower VLP level than raw feces (3.7×10^{10} VLPs/g). When preparing the intervention inocula,

the dose was therefore adjusted to match similar absolute VLP levels (Fig. 4B) while carrying

different levels of viral diversity (Fig. 4C). Aerobic and anaerobic cultivation showed no bacterial

growth and nucleic acid extraction yielded extremely low DNA levels (0.0-0.4 ng/µl), with no

amplification of 16S rRNA gene segments. Therefore, the inocula were considered bacteria-free.

Figure 4. Preparation of bacteria-free virome inocula from feces and chemostats with matched virus levels.
A) Inocula were produced by isolating virus-like particles (VLPs) from 1) donor feces to produce fecal virome
transfer (FVT) solution, from 2) chemostat culture with lactose (LAC) medium to produce chemostat virome
transfer (CVT) solution, and from 3) chemostat culture with lactose-milk oligosaccharide (LAC-MO) medium
to produce CVT-MO solution. B) Estimation of average VLP level in donor feces (VLP/g), the chemostat
cultures (VLP/mL), and the dose-adjusted inocula (VLP/mL). C) Number of observed viral species as a
measure of viral alpha diversity in each inoculum and control (CON) solution.

537 A human monocyte cell line was employed to investigate the immunological impact of stimulation

538 with the chemostat viromes. These responses were compared with the dose-adjusted native fecal

539 virome (which includes eukaryotic viruses) and a dose-adjusted pure phage solution (free of

eukaryotic viruses). Interestingly, chemostat viromes induced a lower expression of toll-like receptor (TLR) 7 compared to the native fecal virome (Suppl. Fig. S4). Since TLR-7 recognizes single-stranded RNA, this could reflect the reduced eukaryotic viral load in chemostat viromes. However, the significant expression mounted by the phage solution, consisting of double-stranded DNA phages, complicates the interpretation. Moreover, the typical virus-associated interferons were not markedly affected by the type of viral solution. Surprisingly, the chemostat viromes produced a higher expression of various pro- and anti-inflammatory cytokines, compared to the native fecal virome and with levels similar to the phage solution. This included tumor necrosis factor (TNF) α, IL-1β, IL-4, IL-8, IL-10, and transforming growth factor (TGF) β. Due to the endotoxin removal step and the use of a high MWCO during viral isolation, we consider it unlikely that these effects are driven by bacterial products in chemostat viromes. Given the obvious eukaryotic viral elimination, these data suggest that this innate immune cell line cannot fully distinguish between eukaryotic and prokaryotic viruses. Since fecal-derived eukaryotic viruses mainly infect intestinal cells, an *in vivo* stimulation may reveal a different response pattern.

Chemostat-propagation mitigated side effects after virome transfer into preterm piglets

Previous studies revealed a good NEC protective effect of FVT but with simultaneous induction of early-onset diarrhea and weight loss in recipients[20]. Suspecting eukaryotic viruses to be the main driver, the chemostat method was applied. In *Piglet Experiment 1*, preterm piglets received a total dose of 10^{10} VLPs from feces or chemostat-propagated inoculum (FVT, CVT, CVT-MO) or a control solution shortly after birth (Fig. 5A). Three animals were euthanized early due to apnea or hindleg issues and were subsequently excluded from the experiment (1 CON, 1 FVT, and 1 CVT-MO). The remaining piglets (*n*=14-15/group) were followed for five days and displayed similar survival rates across groups (Fig. 5B). Upon euthanasia, gastrointestinal necropsy revealed an unusually low occurrence of NEC-like lesions in the control group as well as intervened animals (Fig. 5C-D). When investigating pathology incidences, the level of stomach necrosis was reduced in FVT pigs relative to controls (Suppl. Fig. S5A). Apart from this, incidences of hyperemia, hemorrhage, pneumatosis intestinalis, and necrosis were low and comparable between groups within each intestinal segment. Similarly, the histopathological assessment and tissue cytokine levels revealed no obvious differences (Fig. 5E and Suppl. Fig. S5B). Collectively, neither the FVT nor chemostat viromes appeared different from controls regarding NEC. However, the unfortunate lack of NEC-afflicted animals in the control group makes it impossible to draw firm conclusions in terms of treatment efficacy.

Figure 5. Chemostat propagation mitigates FVT-induced diarrhea but does not affect NEC-like lesioning in a low-incidence setting (*Piglet Experiment 1*). A) Cesarean section-delivered piglets received either control solution (CON) or one of the VLP-rich inocula (FVT, CVT, CVT-MO) over four oral treatments. The pigs were fed increasing volumes of enteral formula 1 until euthanasia. B) Survival curve. C) Gross pathology score in stomachs. D) Cumulative gross pathology score in small intestines and colons. E) Cumulative microscopic lesion score in small intestines and colons. F) Time to diarrhea onset. G) Body weight development showing percentage change from the median birth weight (±SD). Lines not sharing the same letter are significantly different ($P < 0.05$). P_{Group} = P-value for group effect, P_{Int} = P-value for group and time interaction. FVT = fecal virome transfer, CVT = chemostat virome transfer, and CVT-MO = CVT propagated with milk oligosaccharides (n = 14-15/group).

The clinical monitoring, however, revealed clear differences in the side effect pattern throughout

the study. Replicating previous results[20], FVT animals debuted with earlier diarrhea, leading to a

reduced weight gain on day four (Fig. 5F-G). Both chemostat viromes were statistically similar to

controls regarding diarrhea onset, suggesting the method successfully alleviated the effect of the
diarrhea-inducing agent (Fig. 5F). By visual inspection, the diarrhea incidence of piglets receiving
lactose or MO-propagated viromes appeared slightly accelerated, resulting in a decline in body
weight by study termination in the CVT group (Fig. 5G). By day five, circulating neutrophil levels
were trending higher in FVT animals compared to controls ($P = 0.08$, Suppl. Fig. S6A). On the
contrary, FVT lymphocyte levels remained lower from day three and onwards. Although
speculative, this implies that FVT may induce migration of lymphocytes to the intestines as well as
granulopoiesis, resulting in a subsequent increase in circulating neutrophils. This response was
mitigated by chemostat propagation. Consistent with previous findings, the FVT side effects were
linked to small intestinal atrophy and reduced lactase activity in the proximal small intestine (Suppl.
Fig. S6B-C). Here, the lactose-propagated virome entailed atrophy while the MO-propagated
virome reduced lactase activity. Additionally, the lactulose-to-mannitol ratio increased only
significantly in FVT animals, suggesting increased intestinal permeability by this treatment alone
(Suppl. Fig. S6D). Conclusively, the side effects associated with fecal virome transfer were
significantly improved, but not fully abolished, using the *in vitro* propagated chemostat viromes.

Gut virus engraftment following fecal virome transfer is lost after chemostat propagation

Having demonstrated a difference in the clinical side effect pattern in response to exogenous native
and chemostat-propagated fecal viruses, we next investigated the changes to the newborn recipient
gut virome using viral metagenomics sequencing. We observed a modest increase in viral richness
in FVT recipients relative to those receiving lactose-propagated fecal virome (Fig. 6A). However,
changes to viral richness did not appear related with the occurrence of NEC (Fig. 6B). Interestingly,
Bray-Curtis ordination plot clearly showed an alteration of the gut viral composition of the FVT
group relative to all remaining groups, which were not different from one another (Fig. 6C). The
binned taxonomic annotation showed mainly *Caudoviricetes* and a significant proportion of
unannotated viruses across all groups (fig. 6D). In pairwise comparisons of single viral OTU
relative abundances between treatment groups and control by DESeq2 analysis, we found a pattern
of modest virus enrichment (32 enriched vs. 16 depleted) in the FVT group but neither of the
chemostat virome groups (CVT: 9 vs. 16; CVT-MO: 17 vs. 15). All of these were annotated as
unknown viruses or the class *Caudoviricetes*. Based on this viral enrichment analysis, we found no
clear evidence of eukaryotic virus shedding to explain the clinical side effects of FVT as virus-
induced. Importantly, when binning the predicted bacterial hosts, gut phages of the FVT recipients
had a much greater specificity for *Lactobacillaceae* accounting for up to 50% relative abundance,

whereas the remaining groups had hardly any *Lactobacillaceae* phages (Fig. 6E). Moreover, when looking into the engraftment potential of the transferred phages, assessed as the percentage of viral OTUs shared between group inoculum and recipient gut sample, only the FVT group showed an increase, whereas recipients of both chemostat-propagated viromes had levels similar to controls (Fig. 6F). Taken together, we documented that transfer of fecal virome leads to an engraftment of viruses short-term and that most of these are phages with specificity for bacteria in the *Lactobacillaceae* family. Chemostat propagation of fecal viruses prior to transplantation disrupted this engraftment.

Figure 6. Native but not chemostat-propagated viromes reshape gut virome and enrich *Lactobacillaceae* phages (*Piglet Experiment* 1). . A-B) Number of observed viral species as a measure of viral alpha diversity between groups (A) and between NEC and healthy appearing colons (B). C) Principal coordinate analysis (PCoA) plot visualizing viral beta diversity based on Bray-Curtis dissimilarity metrics. False discovery rate (FDR) adjusted *P*-values for pairwise comparisons are reported in the adjacent table. D) Mean relative abundance of viral bacterial hosts summarized at genus level. E) Predicted lifestyle of bacteriophages based on presence (temperate) or absence (virulent) of integrase and/or recombinase genes. F) The percentage of shared viral OTUs between recipients of each group and the inoculum material of the group. CON = control, FVT = fecal virome transfer, CVT = chemostat virome transfer, and CVT-MO = CVT propagated with milk oligosaccharides (*n* = 10-13/group). *$P < 0.05$, **$P < 0.01$

Lactobacillaceae expansion following fecal virome transfer is lost after chemostat propagation

The few animals with NEC (here defined as the presence of hemorrhage, pneumatosis intestinalis, and/or necrosis) had a higher bacterial richness in the colon, whereas no differences were seen between treatment groups (Fig. 7A-B). The bacterial composition in recipients of native FVT and MO-propagated fecal virome were slightly different from controls (Fig. 7C). From the taxonomic overview showing the most abundant taxa across groups, the most obvious difference between virome treatment groups and control was an increase in the relative abundance of *Streptococcus,* the same bacterial genus that dominated the chemostat community (Fig. 7D). Additionally, only FVT recipients harboured an increased proportion of *Lactobacillus amylovorus* (Fig. 7D). We performed pairwise DESeq2 analysis to identify all enriched and depleted bacterial species between treatment groups and control. Notably, FVT recipients were the only ones enriched with three different species of *Lactobacillaceae*, and concomitantly depleted in the archetypical NEC pathogen *Clostridium perfringens* (Suppl. Fig. S7 and Suppl. Table S3)[56]. The compositional effects of MO-propagated fecal virome were also evident by DESeq2, where a simultaneous reduction in both *Lactobacillaceae* and *Clostridium perfringens* was observed.

To investigate a potential link between gut microbes and mammalian host response further, we performed bulk RNA-Seq on mesenteric lymph nodes. We observed no major shifts in global gene expression patterns based on a principal component analysis (Fig. 7E). Gene set enrichment analyses did however identify a list of Gene Ontology pathways almost exclusively involved in immune cell activation. Contrary to expectations, these were all enriched in both groups receiving chemostat-propagated virome relative to FVT (Fig. 7F-G). As such, this finding does not support a eukaryotic viral infection by the transferred native fecal virome as an explanation for the observed side effects. However, the collected lymph nodes were draining more distal intestinal segments (near the ileocolic artery), far from the site where donor viruses were first introduced by oral gavage. Since FVT affects the small intestine (Suppl. Fig. S6B-D), sampling from more proximal regions (near the superior mesenteric artery) may have revealed different treatment responses. Additionally, sampling at the time of FVT-associated diarrhea acceleration (48 hours) might uncover group differences not apparent at euthanasia (96 hours), when all groups displayed similar diarrhea levels.

Figure 7. Virome treatments induce modest changes in the gut bacteriome, but have a limited impact on mesenteric lymph node gene expression (*Piglet Experiment 1*).

A-B) Number of observed bacterial species between NEC and healthy appearing colons (A) and between groups (B). C) Principal coordinate analysis (PCoA) plot visualizing bacterial compositional differences as determined by Bray-Curtis dissimilarity metrics. False discovery rate (FDR) adjusted *P*-values for pairwise comparisons are reported in the adjacent table. D) Mean relative bacterial abundance summarized at species or genus (g) level. E) Principal component analysis (PCA) score plot illustrating the global gene expression differences in mesenteric lymph nodes. F-G) Gene set enrichment analysis (GSEA) based on gene ontology classification, showing all significantly enriched pathways (false discovery rate adjusted P-value < 0.05) between CVT and FVT (F) and between CVT-MO and FVT (G). The size of the dots indicates the gene ratio, while the yellow color indicates a lower adjusted P-value. NS = non-significant. CON =

684 control, FVT = fecal virome transfer, CVT = chemostat virome transfer, and CVT-MO = CVT propagated with
685 milk oligosaccharides (n = 10-13/group). *P < 0.05.
686

Milk oligosaccharide-propagated fecal virome did not prevent NEC-like lesioning

The unusually low levels of gut lesions among controls in *Piglet Experiment 1* did not allow for the

investigation of a potential NEC-reducing effect of chemostat-propagated fecal viruses. This

prompted *Piglet Experiment 2*, where two litters of pigs were fed a UHT-treated infant formula

(formula 2) with a demonstrated higher NEC-inducing capacity (Fig. 8A)[37]. To minimize animal

numbers, we focused on investigating the MO-propagated virome, as this treatment had the greatest

microbiome impact of the chemostat-propagated viromes. Both groups displayed similar survival

rates throughout the study period (Fig. 8B). At necropsy, the general level of intestinal gross

pathology was markedly higher compared to *Piglet Experiment 1* (median score of 7 vs. 0).

Chemostat virome recipients exhibited more severe stomach lesions (Fig. 8C and Suppl. Fig. S8A),

not previously noted. The overall enterocolitis level was similar at macroscopic and microscopic

assessments (Fig. 8D-E). However, the chemostat virome group showed a slightly higher incidence

of small intestinal bleeding (P = 0.05, Suppl. Fig. S8B) but with subtly lower IL-8 levels (P = 0.03,

Suppl. Fig. S8D). The visual trends of slightly earlier diarrhea and decreased body weight aligned

with those observed in *Piglet Experiment 1* (Fig. 8F-G). Again, these trends were not statistically

significant and did not entail intestinal atrophy (Suppl. Fig. S8E). However, intestinal permeability

was slightly increased (P = 0.01, Suppl. Fig. S8F). The colonic bacterial analysis showed only

minor changes, consistent with the previous findings (Suppl. Fig. S9). Collectively, the MO-

propagated fecal virome caused gastritis and appeared to mildly compromise small intestinal health.

This highlights that while chemostat propagation effectively mitigates virus-associated diarrhea, the

method needs further optimization to target NEC.

Discussion

Transfer of fecal phages has shown promise in modulating the preterm gut microbiome and

reducing NEC under experimental settings[17,20]. However, as diarrhea occurs simultaneously[20], this

highlights a need for further refinements to increase the level of safety while maintaining efficacy.

Expanding on previous research[24], we investigated whether the elimination of eukaryotic viruses

during chemostat processing could reduce potential side effects of fecal virome transfers for

preterm neonates while preserving a phage composition capable of correcting GM dysbiosis and

preventing NEC.

717
718
Figure 8. MO-propagated virome modestly exacerbates gastric injury and fails to improve NEC (*Piglet*
Experiment 2). A) Overview of study design. Cesarean section-delivered piglets were allocated to one
control group (CON) receiving SM buffer and one treatment group receiving chemostat virome transfer
propagated with milk oligosaccharides (CVT-MO). The pigs were fed increasing volumes of enteral formula 2
until euthanasia after 96 hours. B) Survival curve. C) Gross pathology score in stomachs. D) Cumulative
gross pathology score in small intestines and colons. E) Cumulative microscopic lesion score in small
intestines and colons. F) Time to diarrhea onset. G) Body weight development showing percentage change
from the median birth weight (±SD). P_{Group} = *P*-value for group effect, P_{Int} = *P*-value for group and time
interaction. *$P < 0.05$ ($n = 26$-27/group).

 Choice of parameters for chemostat propagation

We have previously demonstrated the anaerobic chemostat setup as a method to cultivate stable
fecal bacterial and viral communities from adult human and murine donors[22,24]. To adapt this for
cultivating colon and cecal content from 10-day-old suckling piglets, several adjustments were
required. First and foremost, the growth medium substrates were adjusted to closely resemble a
chemostat setup mimicking the proximal colon of breastfed neonates[23], which contained lactose as
the main carbohydrate source along with gastric mucin. However, since most lactose is absorbed
before reaching the colon[57], we considered it important to investigate the effect of MOs as an
additional substrate for cultivation of a milk-shaped fecal microbiome[58]. Despite being less diverse
than human milk, porcine milk contains 30-50 different MOs, several of which are also found in
human milk[32,53,54]. For simplicity, we selected three key MOs found in both species (2'FL, 3'SL,
and LNnT) and known to be degradable by piglet fecal bacteria[59]. Opposite to infants, young pigs
harbor very few bifidobacteria[60]. Therefore, the gradual emergence of fecal *Bacteroidota* species
during chemostat effectively demonstrated the MO-utilizing potential of these gut bacteria
inhabiting healthy suckling piglets[61,62]. MOs allowed the cultivation of around 150 additional
bacterial species and increased the bacterial similarity to better resemble the donor. Optimizing the
MO composition may promote the dominance of more favorable MO-degrading *Bacteroides* spp.
over *B. pyogenes*[63]. Together, this provides a strong proof-of-concept for using MOs to culture
neonatal fecal bacteria *in vitro*.

Despite the restorative effect of MO supplementation, the bacterial composition of the end culture
was fundamentally different from the donor feces, which was primarily due to losing virtually all
lactobacilli. As lactobacilli are fast-growing bacteria that primarily utilize lactose[64], which was
present in ample amounts in both chemostats (especially LAC), the growth inferiority is unlikely
due to substrate limitations. Instead, it may be due to suboptimal pH levels. Studies report varying
colonic pH levels in suckling pigs ranging from 6.3 to 7.2[65–67] and thus an intermediate level of 6.5
was chosen for the experiment. The study by Pham *et al*. investigated pH levels ranging from 5 to 7
and found infant lactobacilli showed optimal growth at pH 5, whereas *Bacteroides* preferred pH 6-
7[23]. Alongside the current findings, this suggests that lower pH levels are needed to support the
growth of both *Lactobacillaceae* and *Bacteroides* (and bifidobacteria) in future neonatal
chemostats.

Chemostat culturing of neonatal feces using the presented parameters resulted in a significant loss
in viral richness, which was not remedied by the addition of MO. Our previous study using adult

fecal material indicated higher phage preservation at slower dilution rates (0.05 h^{-1}) for murine, but

not for human fecal cultures[24]. As the pig GIT resembles humans more than mice[68], we thus decided

on the higher dilution rate (0.2 h^{-1}) to ensure a significant eukaryotic viral elimination. Albeit, the

current study suggests that slower dilution may be required for proper *in vitro* phage propagation of

a neonatal gut virome. The viral inoculation dose of 10^{10} VLPs/kg was chosen to match previous

studies demonstrating NEC prevention (ref to ISME + gut microbes paper). However, if a

transferred phage encounters a suitable host and enters the lytic cycle, phage titers may increase

drastically. Consequently, greater phage richness will, irrespective of the starting dose, increase the

probability of productive phage-bacterial interactions and their overall impact on the gut

microbiome.

Correction of preterm gut dysbiosis

Although the chemostats poorly resembled the piglet gut microbiome, the MO-induced increase in

Bacteroidota-specific phages could beneficially modulate the recipient gut microbiome. Indeed, we

found that a significant proportion of phages in the chemostat virome, including those targeting

Bacteroidota, were induced prophages, as opposed to the fecal inoculum containing almost

exclusively virulent phages. Although less abundant in human infants compared to piglets,

Bacteroides is a part of the normal developing microbiome[11,69,70] and is positively linked to

intestinal health[71]. FVT has repeatedly been shown to increase the similarity between donor and

recipient bacterial compositions, without administering the actual donor bacteria[72,73]. We have

previously observed increased levels of *Bacteroides* spp. in piglet colons after receiving NEC-

reducing FMT or FVT treatments[17,74]. However, the MO-propagated chemostat virome did not

increase *Bacteroides* or *Parabacteroides* spp. in the recipient's gut, nor did it provide NEC

protection when tested in a high-incidence setting in *Experiment 2*. We find it likely that this lack of

engraftment could be due to an absence of bacterial hosts in the preterm piglet recipients.

As Experiment 1 piglets had unusually low levels of intestinal lesions, the NEC-reducing capacity

of this particular donor fecal virome remains unknown, although the FVT concept has been proven

efficacious on previous occasions using similar donor material[17,20]. Importantly, the virome transfer

was capable of reducing Clostridium perfringens while increasing the abundance of

Lactobacillaceae members. Although NEC cases were sparse, these showed the opposite trend

regarding these bacteria, thus indicating a restorative effect of FVT on NEC-associated dysbiosis.

Peculiarly, viral metagenomics data indicated that mainly virulent phages from the FVT inoculum

engrafted in recipients and that these had Lactobacillaceae as hosts. This contrasts a simple lytic

predation of target bacteria as an explanatory mechanism. This virulent classification, however,

only reflects the absence of known recombinase or integrase genes, meaning these phages are

considered potentially virulent based on current annotations. Regardless, the lack of

Lactobacillaceae in chemostats could potentially explain the lack of recipient gut engraftment of the

chemostat-propagated viromes despite transferring similar numbers of virus-like particles. As we

have previously seen Lactobacillaceae abundance linked to NEC protection[30,74], this again

highlights a target for optimizing chemostat conditions to maintain this bacterial family, thereby

enabling efficacy of chemostat propagated feces. Moreover, whether the reduction of C. perfringens

was due to transferred lytic phages or as a result of competitive exclusion by other bacteria remains

unknown. However, we acknowledge that the absence of demonstrated NEC prevention by the

native donor virome is a significant limitation to conclusions regarding efficacy. Without a

protective baseline, we cannot assess whether the virome efficacy was lost during chemostat

propagation. Consequently, we cannot confirm or dismiss the hypothesis that chemostats can

preserve a phage community capable of preventing NEC.

Transfer of chemostat viromes entails later diarrhea

We have previously demonstrated the use of chemostat viromes to improve the health status of mice

during metabolic syndrome and *Clostridioides difficile* infection[26,27]. This initiative aimed to 1)

overcome variations in efficacy between FVT donors and 2) enhance safety by removing donor-

derived eukaryotic viruses[24]. Since the existing studies showed no side effects, the importance of

eukaryotic virus removal for improved FVT safety has remained a theoretical concern. The present

study, however, showed that FVT induces diarrhea consistently and that chemostat propagation

delays diarrhea onset. Regardless, since the viral metagenomics analysis did not identify a specific

eukaryotic viral agent, the exact mechanism for how chemostat propagation reduces side effects

remains elusive[20]. The four eukaryotic viral families found in feces or batch cultures (*Smacoviridae,*

Genomoviridae, Picobirnaviridae, and *Astroviridae*) were not found upregulated in FVT recipients,

nor was any inflammatory signal detected in mesenteric lymph nodes on day five. As diarrhea appeared on days 2-3 alongside reduced systemic lymphocyte counts, earlier analyses may have revealed different trends between groups.

Conclusion

In conclusion, we show that chemostat propagated fecal virome reduces eukaryotic virus abundance drastically and minimizes gastrointestinal side effects when transferred into newborn recipients. Chemostat cultivation is a promising method to generate complex but reproducible fecal viromes, and supplementing the culture with MOs shows promise for approximating milk-shaped fecal microbiomes. However, as neither the chemostat-propagated viromes nor the native donor virome demonstrated NEC prevention, the efficacy of the chemostat approach remains inconclusive. MO composition, pH, and dilution rate may need optimization for better virus engraftment, recipient microbiome restoration, and NEC protection in future studies.

Author's contributions

SMO, SA, TR, DSN, AB, and KA conceived and planned the experiments. SMO, SA, and KA performed the chemostat experiments. SA and KA performed metabolite analyses. SMO, MS, and AB performed the animal experiments. XM carried out bacteriome analyses. XM and FL carried out virome analyses. JZ performed additional lymph node transcriptome analyses. All authors were involved in critical evaluation and interpretation of results. SMO and AB drafted the manuscript with inputs from all the other authors.

Competing interests

The authors declare no competing interests.

Availability of data and materials

All data associated with this study are present in the paper or the Supplementary Materials. All sequencing datasets are available on the BioProject database under accession number PRJNA1121201.

Funding

Funding was provided by the Estonian Ministry of Science and Education (Project number IUT1927) and the Novo Nordisk Foundation (grant ID: NNF-20OC0063874 under the acronym "PrePhage").

References

1. Laursen MF, Pekmez CT, Larsson MW, et al. Maternal milk microbiota and oligosaccharides contribute to the infant gut microbiota assembly. *ISME Commun* 2021; 1: 21.

2. Chernikova DA, Madan JC, Housman ML, et al. The premature infant gut microbiome during the first 6 weeks of life differs based on gestational maturity at birth. *Pediatr Res* 2018; 84: 71–79.

3. Stokholm J, Thorsen J, Chawes BL, et al. Cesarean section changes neonatal gut colonization. *J Allergy Clin Immunol* 2016; 138: 881-889.e2.

4. Ainonen S, Tejesvi M V., Mahmud MR, et al. Antibiotics at birth and later antibiotic courses: effects on gut microbiota. *Pediatr Res* 2022; 91: 154–162.

5. Chi C, Fan Y, Li C, et al. Early gut microbiota colonisation of premature infants fed with breastmilk or formula with or without probiotics: A cohort study. *Nutrients* 2021; 13: 1–10.

6. Litvak Y, Byndloss MX, Tsolis RM, et al. Dysbiotic Proteobacteria expansion: a microbial signature of epithelial dysfunction. *Curr Opin Microbiol* 2017; 39: 1–6.

7. Pammi M, Cope J, Tarr PI, et al. Intestinal dysbiosis in preterm infants preceding necrotizing enterocolitis: a systematic review and meta-analysis. *Microbiome* 2017; 5: 31.

8. Jones IH, Hall NJ. Contemporary Outcomes for Infants with Necrotizing Enterocolitis—A Systematic Review. *J Pediatr* 2020; 220: 86-92.e3.

9. Shah SA, Deng L, Thorsen J, et al. Expanding known viral diversity in the healthy infant gut. *Nat Microbiol* 2023; 8: 986–998.

10. Almeida GMF, Laanto E, Ashrafi R, et al. Bacteriophage adherence to mucus mediates preventive protection against pathogenic bacteria. *MBio* 2019; 10: e01984-19.

11. Lim ES, Zhou Y, Zhao G, et al. Early life dynamics of the human gut virome and bacterial microbiome in infants. *Nat Med* 2015; 21: 1228–1234.

12. Liang G, Zhao C, Zhang H, et al. The stepwise assembly of the neonatal virome is modulated by breastfeeding. *Nature* 2020; 581: 470–474.

13. Liang G, Conrad MA, Kelsen JR, et al. Dynamics of the stool virome in very early-onset inflammatory bowel disease. *J Crohn's Colitis* 2020; 14: 1600–1610.

14. Clooney AG, Sutton TDS, Shkoporov AN, et al. Whole-Virome Analysis Sheds Light on Viral Dark Matter in Inflammatory Bowel Disease. *Cell Host Microbe* 2019; 26: 764-778.e5.

15. Gogokhia L, Buhrke K, Bell R, et al. Expansion of Bacteriophages Is Linked to Aggravated Intestinal Inflammation and Colitis. *Cell Host Microbe* 2019; 25: 285-299.e8.

16. Kaelin EA, Rodriguez C, Hall-Moore C, et al. Longitudinal gut virome analysis identifies

specific viral signatures that precede necrotizing enterocolitis onset in preterm infants. *Nat*
Microbiol 2022; 7: 653–662.

17.   Brunse A, Deng L, Pan X, et al. Fecal filtrate transplantation protects against necrotizing
enterocolitis. *ISME J* 2021; 16: 686–694.

18.   Jankowski R. Viruses and viral epidemics in the metabolic theory of evolution. *Eur Ann*
Otorhinolaryngol Head Neck Dis 2020; 137: 297–301.

19.   Ng RWY, Dharmaratne P, Wong S, et al. Revisiting the donor screening protocol of faecal
microbiota transplantation (FMT): A systematic review. *Gut* 2024; 73: 1029–1031.

20.   Offersen SM, Mao X, Spiegelhauer MR, et al. Fecal virus-like particles are sufficient to
reduce necrotizing enterocolitis. *Gut Microbes* 2024; 16: 2392876.

21.   Garmaeva S, Sinha T, Kurilshikov A, et al. Studying the gut virome in the metagenomic era:
Challenges and perspectives. *BMC Biol* 2019; 17: 84.

22.   Adamberg K, Raba G, Adamberg S. Use of Changestat for Growth Rate Studies of Gut
Microbiota. *Front Bioeng Biotechnol* 2020; 8: 1–12.

23.   Pham VT, Chassard C, Rifa E, et al. Lactate Metabolism Is Strongly Modulated by Fecal
Inoculum, pH, and Retention Time in PolyFermS Continuous Colonic Fermentation Models
Mimicking Young Infant Proximal Colon. *mSystems* 2019; 4: e00264-18.

24.   Adamberg S, Rasmussen TS, Larsen SB, et al. Reproducible chemostat cultures to minimize
eukaryotic viruses from fecal transplant material. *iScience* 2024; 27: 110460.

25.   Wang Y, Hoenig JD, Malin KJ, et al. 16S rRNA gene-based analysis of fecal microbiota
from preterm infants with and without necrotizing enterocolitis. *ISME J* 2009; 3: 944–954.

26.   Mao X, Larsen SB, Zachariassen LSF, et al. Transfer of modified gut viromes improves
symptoms associated with metabolic syndrome in obese male mice. *Nat Commun* 2024; 15:
4704.

27.   Rasmussen TS, Mao X, Forster S, et al. Overcoming donor variability and risks associated
with fecal microbiota transplants through bacteriophage-mediated treatments. *Microbiome*
2024; 12: 119.

28.   Kump P, Wurm P, Gröchenig HP, et al. The taxonomic composition of the donor intestinal
microbiota is a major factor influencing the efficacy of faecal microbiota transplantation in
therapy refractory ulcerative colitis. *Aliment Pharmacol Ther* 2018; 47: 67–77.

29.   Haifer C, Luu LDW, Paramsothy S, et al. Microbial determinants of effective donors in
faecal microbiota transplantation for UC. *Gut* 2022; 90–100.

30.   Hui Y, Vestergaard G, Deng L, et al. Donor-dependent fecal microbiota transplantation
efficacy against necrotizing enterocolitis in preterm pigs. *NPJ Biofilms Microbiomes* 2022; 8:
48.

31.   Difilippo E, Pan F, Logtenberg M, et al. Milk Oligosaccharide Variation in Sow Milk and
Milk Oligosaccharide Fermentation in Piglet Intestine. *J Agric Food Chem* 2016; 64: 2087–
2093.

32.   Wei J, Wang ZA, Wang B, et al. Characterization of porcine milk oligosaccharides over
lactation between primiparous and multiparous female pigs. *Sci Rep* 2018; 8: 4688.

33.   Ingram BYDL, Legge KF. Variations in deep body temperature in the young unrestrained pig
over the 24 hour period. *J Physiol* 1970; 210: 989–998.

34.   Larsen F, Offersen SM, Li VR, et al. Choice of Ultrafilter affects Recovery Rate of
Bacteriophages. *Viruses* 2023; 15: 2051.

35.   Sangild PT, Thymann T, Schmidt M, et al. Invited review: The preterm pig as a model in
pediatric gastroenterology. *J Anim Sci* 2013; 91: 4713–4729.

36.   Østergaard MV, Shen RL, Støy ACF, et al. Provision of Amniotic Fluid during Parenteral
Nutrition Increases Weight Gain with Limited Effects on Gut Structure, Function, Immunity,
and Microbiology in Newborn Preterm Pigs. *J Parenter Enter Nutr* 2016; 40: 552–566.

37.   Offersen SM, Henriksen NL, Brunse A. A weighted and cumulative point system for
accurate scoring of intestinal pathology in a piglet model of necrotizing enterocolitis. *Exp*
Mol Pathol 2024; 140: 104936.

38.   Li Y, Ostergaard M V., Jiang P, et al. Whey protein processing influences formula-induced
gut maturation in preterm pigs. *J Nutr* 2013; 143: 1934–1942.

39.   Wu Z, Tran Nam Tien N, Baek O, et al. Regulation of host metabolism and defense strategies
to survive neonatal infection. *BBA - Mol Basis Dis* 2024; 1870: 167482.

40.   Quast C, Pruesse E, Yilmaz P, et al. The SILVA ribosomal RNA gene database project:
Improved data processing and web-based tools. *Nucleic Acids Res* 2013; 41: 590–596.

41.   Larsen F. VLP Extraction from Fecal Samples V.2. DOI:
dx.doi.org/10.17504/protocols.io.b2qaqdse.

42.   Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence
data. *Bioinformatics* 2014; 30: 2114–2120.

43.   Shen W, Le S, Li Y, et al. SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q file
manipulation. *PLoS One* 2016; 11: 1–10.

44.   Bankevich A, Nurk S, Antipov D, et al. SPAdes: A new genome assembly algorithm and its
applications to single-cell sequencing. *J Comput Biol* 2012; 19: 455–477.

45.   Camargo AP, Roux S, Schulz F, et al. Identification of mobile genetic elements with
geNomad. *Nat Biotechnol* 2023; 42: 1303–1312.

46.   Nayfach S, Camargo AP, Schulz F, et al. CheckV assesses the quality and completeness of
metagenome-assembled viral genomes. *Nat Biotechnol* 2021; 39: 578–585.

47.   Roux S, Camargo AP, Coutinho FH, et al. iPHoP: An integrated machine learning
framework to maximize host prediction for metagenome-derived viruses of archaea and
bacteria. *PLoS Biol* 2023; 21: e3002083.

48.   McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. *PLoS One* 2013; 8: e61217.

49.   Dixon P. VEGAN, a package of R functions for community ecology. *Journal of Vegetation*

Science 2003; 14: 927–930.

50.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* 2014; 15: 1–21.

51.   Andersen KS, Kirkegaard RH, Karst SM, et al. ampvis2: An R package to analyse and
visualise 16S rRNA amplicon data. *bioRxiv* 2018; 10-11 (preprint).

52.   McKnight DT, Huerlimann R, Bower DS, et al. microDecon: A highly accurate read-
subtraction tool for the post-sequencing removal of contamination in metabarcoding studies.
Environ DNA 2019; 1: 14–25.

53.   Salcedo J, Frese SA, Mills DA, et al. Characterization of porcine milk oligosaccharides
during early lactation and their relation to the fecal microbiome. *J Dairy Sci* 2016; 99: 7733–
7743.

54.   Cheng L, Xu Q, Yang K, et al. Annotation of porcine milk oligosaccharides throughout
lactation by hydrophilic interaction chromatography coupled with quadruple time of flight
tandem mass spectrometry. *Electrophoresis* 2016; 37: 1525–1531.

55.   Santiago-Rodriguez TM, Ly M, Daigneault MC, et al. Chemostat culture systems support
diverse bacteriophage communities from human feces. *Microbiome* 2015; 3: 58.

56.   Kiu R, Shaw AG, Sim K, et al. Particular genomic and virulence traits associated with
preterm infant-derived toxigenic Clostridium perfringens strains. *Nat Microbiol* 2023; 8:
1160–1175.

57.   Redel CA, Shulman RJ, Tivey DR. Determinants of lactose digestion in the miniature pig.
Dig Dis Sci 1997; 42: 137–144.

58.   Marcobal A, Barboza M, Froehlich JW, et al. Consumption of Human Milk Oligosaccharides
by Gut-related Microbes. *J Agric Food Chem* 2011; 58: 5334–5340.

59.   Difilippo E, Pan F, Logtenberg M, et al. In Vitro Fermentation of Porcine Milk
Oligosaccharides and Galacto-oligosaccharides Using Piglet Fecal Inoculum. *J Agric Food*
Chem 2016; 64: 2127–2133.

60.   Mikkelsen LL, Bendixen C, Jakobsen M, et al. Enumeration of Bifidobacteria in
Gastrointestinal Samples from Piglets. *Appl Environ Microbiol* 2003; 69: 654–658.

61.   Guevarra RB, Lee JH, Lee SH, et al. Piglet gut microbial shifts early in life: causes and
effects. *J Anim Sci Biotechnol* 2019; 10: 1.

62.   Marcobal A, Barboza M, Sonnenburg ED, et al. Bacteroides in the Infant Gut Consume Milk
Oligosaccharides via Mucus-Utilization Pathways. *Cell Host Microbe* 2011; 10: 507–514.

63.   Fernández Vecilla D, Urrutikoetxea Gutiérrez MJ, Roche Matheus MP, et al. Description of
eight human infections caused by Bacteroides pyogenes in a tertiary hospital of northern
Spain. *Anaerobe* 2023; 82: 1–4.

64.   Gänzle MG, Follador R. Metabolism of oligosaccharides and starch in lactobacilli: A review.
Front Microbiol 2012; 3: 1–15.

65.   Snoeck V, Cox E, Verdonck F, et al. Influence of porcine intestinal pH and gastric digestion

on antigenicity of F4 fimbriae for oral immunisation. *Vet Microbiol* 2004; 98: 45–53.

66. Alizadeh A, Akbari P, Difilippo E, et al. The piglet as a model for studying dietary
components in infant diets: Effects of galacto-oligosaccharides on intestinal functions. *Br J*
Nutr 2016; 115: 605–618.

67. Novotný J, Haraszti K, Krepelková Z, et al. Analysis of Faecal pH in Piglets from Birth to
Weaning. *Folia Vet* 2022; 66: 1–5.

68. Anderschou B, Jensen H, Heyndrickx M, et al. Small intestine vs. colon ecology and
physiology: Why it matters in probiotic administration. *Cell Reports Med* 2023; 4: 101190.

69. Stokholm J, Thorsen J, Blaser MJ, et al. Delivery Mode and Gut Microbial Changes
Correlate with an Increased Risk of Childhood Asthma. *Sci Transl Med* 2021; 12: eaax9929.

70. Frese SA, Parker K, Calvert CC, et al. Diet shapes the gut microbiome of pigs during nursing
and weaning. *Microbiome* 2015; 3: 1–10.

71. Wang X, Cai Z, Wang Q, et al. Bacteroides methylmalonyl-CoA mutase produces propionate
that promotes intestinal goblet cell differentiation and homeostasis. *Cell Host Microbe* 2024;
32: 63–78.

72. Ott SJ, Waetzig GH, Rehman A, et al. Efficacy of Sterile Fecal Filtrate Transfer for Treating
Patients With Clostridium difficile Infection. *Gastroenterology* 2017; 152: 799–811.

73. Rasmussen TS, Mentzel CMJ, Refslund M, et al. Fecal virome transfer improves
proliferation of commensal gut Akkermansia muciniphila and unexpectedly enhances the
fertility rate in laboratory mice. *Gut Microbes* 2023; 15: 2208504.

74. Brunse A, Martin L, Rasmussen TS, et al. Effect of fecal microbiota transplantation route of
administration on gut colonization and host response in preterm pigs. *ISME J* 2019; 13: 720–
733.

1021