Two-dimensional perisaccadic visual mislocalization in rhesus macaque monkeys

Matthias P. Baumann[1,2] and Ziad M. Hafed[1,2]

1. Werner Reichardt Centre for Integrative Neuroscience, University of Tübingen, Tübingen, Germany
2. Hertie Institute for Clinical Brain Research, University of Tübingen, Tübingen, Germany

Correspondence: ziad.m.hafed@cin.uni-tuebingen.de

Abstract

Perceptual localization of brief, high contrast perisaccadic visual probes is grossly erroneous. While this phenomenon has been extensively studied in humans, more needs to be learned about its underlying neural mechanisms. This ideally requires running similar behavioral paradigms in animals. However, during neurophysiology, neurons encountered in the relevant sensory and sensory-motor brain areas for visual mislocalization can have arbitrary, non-cardinal response field locations. This necessitates using mislocalization paradigms that can work with any saccade direction. Here, we first established such a paradigm in three male rhesus macaque monkeys. In every trial, the monkeys generated a visually-guided saccade towards an eccentric target. Once a saccade onset was detected, we presented a brief flash at one of three possible locations ahead of the saccade target location. After an experimentally-imposed delay period, we removed the saccade target, and the monkeys were then required to generate a memory-guided saccade towards the remembered flash location. All three monkeys readily learned the task, and, like humans, they all showed strong backward mislocalization towards the saccade target, which recovered for later flashes from the saccade time. Importantly, we then replicated a well-known property of human perisaccadic mislocalization, as revealed by two-dimensional mislocalization paradigms: that mislocalization is strongest for upward saccades. For horizontal saccades, we additionally found stronger mislocalization for upper visual field flashes, again consistent with humans. Our results establish a robust two-dimensional mislocalization paradigm in monkeys, and they pave the way for exploring the neural mechanisms underlying the dependence of perisaccadic mislocalization strength on saccade direction.

Significance

Visual perception is strongly altered around the time of saccades. Such alteration is often studied by characterizing how brief perisaccadic visual flashes are perceptually localized. While the properties of visual mislocalization have been exhaustively studied, the underlying neural mechanisms remain elusive, and this is due to a lack of suitable behavioral paradigms in animal models. We describe such a paradigm for macaques, which are ideal for exploring sensory-motor neural processes related to perisaccadic mislocalization.

Introduction

Studying visual perception necessarily requires also investigating the interplays between visual sensory processing and eye movements. This is because eye movements, such as saccades, can drastically alter retinal images even if the external environment is completely stable.

Multiple alterations in visual information processing are known to accompany saccades. For example, perisaccadic visual sensitivity is reduced, rendering brief visual presentations completely invisible in some cases (Latour, 1962; Zuber and Stark, 1966; Beeler, 1967). Moreover, when stimulus contrasts are high enough to be detected, these stimuli are very grossly mislocalized (Honda, 1989; Cai et al., 1997; Ross et al., 1997). Such mislocalization depends on various factors related to the visual environment (Lappe et al., 2000). For example, in complete darkness, brief visual flashes are perceived to have appeared ahead of their true positions (Dassonville et al., 1992; Lappe et al., 2000); such mislocalization is often referred to as forward mislocalization, since the error in reported position has a vector of similar direction to the saccade vector direction (Klingenhoefer and Krekelberg, 2017). On the other hand, with the presence of visual references, perisaccadic mislocalization appears like a "compression" towards the saccade target: brief flashes presented ahead of the saccade target experience mislocalization in the opposite direction from the saccade (Ross et al., 1997), sometimes referred to as reverse or backward mislocalization (Klingenhoefer and Krekelberg, 2017); flashes presented in between the initial gaze position and the saccade target are mislocalized forward in the direction of the saccade.

Even though much of the early work on perisaccadic mislocalization has focused on only one dimension (horizontal saccades and perceptual reports on only the horizontal position of flashes), subsequent pioneering work demonstrated that, even with horizontal saccades, mislocalization has a distinct two-dimensional landscape (Kaiser and Lappe, 2004). More importantly, saccade directions themselves alter the two-dimensional mislocalization landscape, with upward saccades causing particularly strong reverse mislocalization for flashes presented ahead of the saccade target (Grujic et al., 2018). This latter result, which was motivated by a difference between upper and lower visual field representations in the superior colliculus (Hafed and Chen, 2016; Zhang et al., 2022), suggests a clear need to study perisaccadic mislocalization, whether perceptually or with neurophysiological experiments, using more than just horizontal saccades.

Here, we aimed to establish a robust and flexible perisaccadic visual mislocalization
paradigm in macaque monkeys, setting the stage for subsequent neurophysiological
experiments. Four previous studies demonstrated perisaccadic mislocalization in monkeys,
but all used only horizontal saccades and with only one-dimensional mislocalization
measures (analyzing the horizontal component of flash positions). In one study, forward
mislocalization was observed in complete darkness (Dassonville et al., 1992), consistent with
human studies (Honda, 1989; Lappe et al., 2000). In another, backward mislocalization was
observed in the absence of complete darkness (even for flashes behind the saccade target
location), inconsistent with so-called "compression" (Jeffries et al., 2007). However, the
flashes used in that study were not the very brief flashes typical of mislocalization
experiments in humans. In the third study, perisaccadic "compression" was observed like in
humans, and under identical experimental conditions between species (Klingenhoefer and
Krekelberg, 2017). And, most recently, evidence of "compression" was again documented,
enabling simultaneous recordings of visual cortical neurons during the behavior (Weng et al.,
2024). Thus, it is indeed plausible to use monkeys for mislocalization experiments.
109
Having said that, the reality of neurophysiological experiments is that saccade direction
could be completely arbitrary. For example, if one were to record superior colliculus saccade
motor bursts, as a potential source of corollary discharge signals (Sommer and Wurtz, 2004,
2006; Berman et al., 2009; Berman and Wurtz, 2010, 2011; Berman et al., 2017) that may
contribute to mislocalization, then it is more likely to record from oblique rather than
horizontal saccade representations. This is also true when probing the representations of
brief perisaccadic flashes themselves. Therefore, and given the dependence of perisaccadic
mislocalization strength on saccade direction (Grujic et al., 2018), true translation of the
classic human mislocalization studies to neurophysiological experiments would only be
possible with a general, and robust, two-dimensional mislocalization paradigm in monkeys.
This is what we document here.
121
122
123
124   **Methods**
125
126   *Experimental animals and ethical approvals*
127   We collected data from three adult, male rhesus macaque monkeys (*Macaca mulatta*). The
128   monkeys (A, F, and M) were aged 11-15 years, and they weighed 10-13 kg.
129
130   All experiments were approved by the ethics committees at the regional governmental
131   offices of the city of Tübingen.
132
133
Laboratory setup and animal procedures
The monkeys were seated in a dark room in front of a computer-controlled display, spanning
approximately 31 deg horizontally and 23 deg vertically. The display was at a distance of ~72
cm from the animals. We tracked eye movements with high precision using the scleral
search coil technique (Robinson, 1963; Fuchs and Robinson, 1966; Judge et al., 1980), and
the experiment was controlled in real-time using a modified version of PLDAPS (Eastman and

Huk, 2012). Stimulus presentation and control was performed using the Psychophysics
Toolbox (Brainard, 1997; Pelli, 1997; Kleiner et al., 2007).
142
The animals were prepared for behavioral and neurophysiological experiments earlier (Tian
et al., 2018; Buonocore et al., 2019; Skinner et al., 2019). This entailed a first surgery to
implant a head fixation apparatus. During data collection, this apparatus was attached to a
coupled device on the monkey chair in order to stabilize head position. Then, a later surgery
was performed to implant a scleral search coil in one eye for eye movement tracking
purposes.
149
150
151 *Experimental procedures*
152 The main task involved generating a primary visually-guided saccade, around which a very
153 brief probe flash was presented; later on in the very same trial, the monkeys generated a
154 "report" saccade aiming at where they remembered the perisaccadic flash to have
155 appeared.
156
Each trial (Fig. 1) started with an initial white fixation spot (of 79.9 cd/m^2 luminance)
presented over a uniform gray background (26.11 cd/m^2). The fixation spot was displaced
from display center (by 7 deg, except for the cardinal saccade directions of monkey M, in
which case it was by 5 deg), and it could appear in one of eight (equally spaced) directions
from this center. After 600-1000 ms, the fixation spot disappeared, and a simultaneous
saccade target appeared at the display center. This target consisted of a vertical Gabor
grating of either 0.5 cycles/deg (cpd) or 8 cycles/deg (cpd) spatial frequency, and it had a
radius of 1.5 deg. The grating had high contrast (100%), and it featured a white central
marker surrounded by a small gray disc, as in our previous experiments (Baumann et al.,
2023). This central marker aided in maintaining saccade accuracy and precision across trials
having different image appearances of the saccade target. The onset of the grating was the
cue to generate a visually-guided saccade.
169
We monitored the monkeys' eye position in real-time and detected the onset of a saccade
when the eye position deviated by 1.5 degrees from the initial fixation spot. Following
saccade detection, a probe flash was presented either immediately or with a delay of 40 or
100 ms (in subsequent post-hoc analyses, we recalculated exact probe times; see below).
The probe flash was displayed for a brief duration of 12 ms (1 frame at a display refresh rate
of 85 Hz) and was positioned 10.5 degrees away from the center of the grating (7.5 deg for
the slightly smaller cardinal saccades of monkey M). The probe flash could appear either
directly along the saccade vector direction (and farther away from the saccade target) or at
an angle of +/-45 deg relative to the saccade vector (again farther away than the saccade
target location). Thus, there were three possible probe flash locations, all of which were
expected to experience backward mislocalization for probe flash times very close to saccade
onset.
182
After the monkeys made the visually-guided saccade, with a maximum allowed reaction time
of 500 ms, they maintained stable fixation on the saccade target for an additional 500-800
ms. At that time, the grating disappeared, cueing the monkeys to make a memory-guided
saccade "reporting" the remembered probe flash location. The monkeys were rewarded if

they made the report saccade within <500 ms, and we allowed a radius of 8 deg tolerance
window around the true flash location for the landing position of the eye. This large
tolerance window was intentional because we wanted to avoid penalizing the monkeys for
their mislocalized percepts; otherwise, the monkeys would have learned, over time, to
constrain their report saccades to the regions gaining them rewards. On successful trials, the
monkeys received juice rewards; on unsuccessful trials (e.g. late initial saccade or broken
fixation), the trials were aborted and juice rewards were withheld.
194
195
196
197

198
199
Figure 1 Perisaccadic visual mislocalization task for macaque monkeys. Each trial started with fixation of an
eccentric spot (bottom-left corner of the display in the shown example). Upon removal of the spot, a saccade
target appeared (Gabor grating), and the monkeys generated a visually-guided saccade towards it (oblique cyan
arrow). We detected saccade onset online, and we presented a brief, single-frame probe flash at one of three
possible delays from saccade detection. The probe flash could appear at one of three possible locations on a
virtual circle (dashed yellow curve and placeholders) around the saccade target: one directly ahead of the
saccade target along the saccade vector direction, and two at +/- 45 deg relative to the saccade vector (the shown
example flash is at the 45 deg counterclockwise position). After an enforced delay, we removed the grating, and
the monkeys reported their perceived flash location with a second, memory-guided, saccade. They had to
maintain their gaze on the reported location for an additional delay time before being rewarded. In baseline
trials, the probe flash appeared 100 ms after saccade detection and remained visible for 400 ms. Thus, it was
clearly visible well after the end of the first saccade, resulting in an expected veridical percept.
212
213
214
215
We also interleaved trials containing what we called a "baseline" condition. In this condition,
which occurred in a total of 6.25% of all trials in a given session, the probe flash was always
presented 100 ms after online saccade detection, and the probe also remained visible for a
full 400 ms. That is, the probe was still clearly visible while the monkeys' were in a stable
fixation position on the grating center after the end of the primary saccade. This allowed
assessing the monkeys' eye movement behavior in the report saccade when no perceptual
mislocalization was to be expected (of course, other errors, such as those associated with
memory-guided saccades, could still occur).

To avoid that the monkeys were solving the task by remembering absolute flash locations
across trials, for example, by guessing their position relative to the edges of the display, we
applied a global constant shift in all stimulus positions, which we changed randomly from
trial to trial, but which we kept constant for all stimulus events within a trial. Specifically, on
every trial, the whole stimulus geometry (initial fixation position, final fixation position, and
the virtual circle on which the probe flashes could occur) was shifted randomly by a constant
known amount, within a range of +/- 0.5 deg both horizontally and vertically. In all analyses,
all positions were re-referenced to the true saccade target location on any given trial. This is
a similar approach to our recent human perisaccadic mislocalization experiments (Baumann
et al., 2024).

Across data collection sessions, we varied the primary saccade direction. Specifically, we
selected eight distinct initial fixation positions on a virtual circle around display center,
allowing us to test eight different saccade vectors. Although our paradigm was designed to
support any starting and end position in theory, we focused on these eight equally spaced
directions for simplicity and for the opportunity to collect many trial repetitions of any given
condition. In practice, we blocked trials of a given saccade direction in a given data collection
session. This was done purely for pragmatic reasons. Specifically, we typically collected
multiple sessions of data from each animal, and we wanted to have large numbers of trials
for each saccade direction. Therefore, it was simpler to specify a given saccade direction for
any given block that we ran. This is also the expected situation during neurophysiological
experiments, in which a given saccade vector would be constant for a given recording site
(for example, in the superior colliculus). However, once the monkeys are proficient in this
task, we believe that randomizing saccade directions within a single session would still be
very easily performed by them (see our note on training trajectory below).

For monkey M, as stated above, the cardinal saccades were only horizontal, and they had an
amplitude of 5 deg (rather than 7 deg). Also, the probe flashes were 7.5 deg away from the
saccade target in these cases (rather than 10.5). Thus, when summarizing this monkey's
results, we separated the oblique conditions (with 7 deg saccades) from the cardinal ones
(see Results). We did this in order to avoid potential confusion with respect to the
quantitative amplitude of mislocalization for the two different saccade sizes that we tested
in this animal. On the other hand, showing qualitatively similar results with different saccade
sizes, both in the same monkey and also across animals (see Results), further confirms our
interpretation of the robustness and flexibility of our visual mislocalization paradigm.

Finally, in the task above, the report by the animals was made through a memory-guided
saccade to the "perceived" flash location. Memory-guided saccades might have systematic
and variable errors in their landing positions (White et al., 1994; Willeke et al., 2019), which
can also depend on saccade direction (Hafed and Chen, 2016). However, as mentioned
above, and as we also show in Results, we varied saccade directions across our experimental
sessions, in order to convince ourselves that what we saw was mislocalization (Grujic et al.,
2018) and not only errors in memory-based reporting. Moreover, the perisaccadic flash
eccentricities were far enough from the saccade target to make the peak mislocalization
errors significantly bigger than any systematic or variable errors expected from memory-
guided saccades (see Results).

271
272   We collected approximately 350-500 trials per session from each monkey, and we analyzed a
273   total of 7582, 16005, and 20902 accepted trials for the saccade direction analysis and a total
274   of 14953, 23830, and 23371 accepted trials for the saccade target appearance analysis from
275   monkeys A, F, and M, respectively (see below for our definitions of saccade direction data
276   analysis and saccade target appearance data analysis).
277
278

279   *Training trajectory*
280   Success in our task was built on a very simple training trajectory that each monkey
281   underwent, and that we describe here briefly. All three monkeys easily mastered the task.
282
All three monkeys were already experts in the individual task components before starting
this project. Specifically, they knew how to perform both visually-guided and memory-
guided saccades. To get the monkeys to report flash position in our current mislocalization
task, we first started with the visually-guided saccade component of the paradigm; the
monkeys simply made a saccade towards the grating image when they were instructed to do
so (by removal of the initial fixation spot and the appearance of the grating). They were also
rewarded classically, immediately after removal of the grating image.
290
We then introduced a second delayed, visually-guided saccade in every trial. Specifically,
upon detection of the first saccade to the grating center, we presented the probe. However,
instead of being flashed briefly, the probe remained visible until trial end. After the first
saccade to the grating center, the monkeys performed a delayed saccade to the probe after
grating removal. The reward now only came after a sequence of two instructed saccades
(both visually-guided). Since the monkeys were already proficient in delayed, visually-guided
saccades, this stage of training was achieved already from the first trial, allowing us to
proceed.
299
We next introduced trials in which we presented the flash like in the main task above, except
for two differences. First, the flash occurred always 100 ms after detecting the visually-
guided saccade. That is, we were sure that it should be minimally mislocalized since it
appeared much later than the saccade time (Ross et al., 1997; Grujic et al., 2018). Second,
the flash was presented for a long period of time (400 ms) on every trial (like in our baseline
condition). This ensured that the flash was minimally mislocalized by the previous saccade,
and that it was still visible during stable fixation of the grating center. Upon removal of the
grating, the monkeys made a relatively accurate memory-guided saccade towards the flash
position, and we immediately rewarded them for having their gaze enter a relaxed virtual
window of 8 deg radius around the true flash position. This made them quickly realize that
the reward was now contingent on reporting the remembered flash position after
completing their initial visually-guided saccade.
312
We then gradually increased the period in which the monkeys had to maintain fixation on
the remembered flash position before getting rewarded. This allowed them to maintain
stable gaze position for a few hundred milliseconds before getting rewarded (Fig. 1). Thus, by
now, the monkeys were proficient in performing an initial visually-guided saccade to an
image, and then making a second memory-guided saccade towards the remembered flash

location (with the flash itself being veridically perceived since it occurred long after saccade
onset, and also since it was visible for a long time). For one monkey (A), the animal had
some difficulty transitioning to the memory-guided component of the task so far. Therefore,
we introduced a few extra sessions in which the probe reappeared when the grating
disappeared. This prompted the monkey to realize that the task was to focus on the probe
flash location.
324
When we were confident that the monkeys were performing this task with high fidelity, we
then reduced the duration of the flash (down to a single display frame). And, finally, we
started introducing the other flash times, in which we expected a large mislocalization error.
As mentioned above, we maintained a relaxed virtual tolerance window radius around flash
position, so that we did not penalize the monkeys for their mislocalized reports. All animals
readily learned the task without much difficulties.
331
332
Data analysis
We detected all saccades (in post-hoc analyses) using the methods that we described
previously (Chen and Hafed, 2013). For the primary saccades, such detection allowed us to
recalculate probe flash times relative to the actual saccade onset in the data. We found that
the probe flashes in the experiment occurred at three distinct times from saccade onset,
consistent with our experimental design. Based on the recalculations, we specifically
classified the probe flash times into the following three categories: probes occurring ~30 ms
after primary saccade onset, probes occurring ~70 ms after primary saccade onset, and
probes occurring ~130 ms after primary saccade onset. That is, the online saccade detection
process described above took approximately 30 ms to trigger the flashes during the data
collection (expected given the dependence of our detection algorithm on eye position
deviating sufficiently from initial fixation, and also given the frame refresh rate of the
display). Figure 2A shows an example of the relationship between primary saccade times
and probe flash onset times from one example session of monkey A. The figure plots radial
eye velocity of the primary saccade (each curve is a saccade), with all trials aligned to probe
flash onset time. There were three distinct time periods separating saccade and flash times
across all trials.
350
Besides checking timing, we also checked primary saccade consistency across conditions.
That is, we wanted to avoid the possibility that any differences in visual mislocalization that
we uncovered (for example, as a function of saccade direction) could be due to systematic
differences in the actual primary saccade metrics and kinematics. Therefore, before
analyzing the behavioral reports of the animals, we first filtered trials according to the eye
movement data of the primary saccades. In one set of comparisons (see Results), we were
interested in investigating the influences of primary saccade direction on perisaccadic visual
mislocalization strength. Therefore, we ensured matched saccade vector and kinematic
properties across the different directions of the primary saccades of our main task (Fig. 1). In
another set of comparisons, we studied the influences of saccade target appearance on
mislocalization strength; here, we matched the primary saccades across the different
saccade target image types (low versus high spatial frequency).
363
364

Figure 2 Controlling probe flash timing and saccade metrics and kinematics across conditions. (A) Radial eye velocity plots of all accepted primary saccades from an example session of monkey A. All traces are aligned to the onset of the probe flash, revealing three distinct perisaccadic flash times, which we categorized by the three different colors. We expected progressively less visual mislocalization for longer times between the saccades and the probe flashes. Baseline trials are not included in this panel. **(B)** Horizontal (top) and vertical (bottom) eye position traces from one example saccade vector direction (up and to the left) out of the same example session shown in **A**. Different colors show the different probe flash times, and the data suggest no systematic differences in the saccades across the flash times. **(C)** We also ensured that landing eye positions (central clouds of dots) overlapped for different saccade directions, again allowing us to investigate the influence of saccade direction on mislocalization strength with overlapping saccade properties (the kinematics were also matched as per the description in the text). The small black dashed circle indicates the extent of the grating towards which the primary instructed saccade was generated, and the big black dashed circle indicates the virtual circle along which the initial fixation positions were placed.

For the vector and kinematic matching in the investigations of primary saccade direction influences, we binned all saccade main sequence data (Zuber et al., 1965; Bahill et al., 1975) into intervals of 0.5 deg in amplitude by 40 deg/s in peak velocity. We included in the analyses only those trials from the main sequence bins that had at least one trial from each of the eight saccade directions (four in the case of monkey M's oblique saccade data; this monkey's cardinal saccades were analyzed separately due to their smaller size). This ensured that we could compare saccades across different directions having similar ranges of amplitudes and peak velocities.

We also noticed that monkey M had a substantial difference in saccade peak velocity between rightward and leftward saccades (~350 versus ~550 deg/s, respectively). Therefore, for this monkey, we decided to report all analyses separately according to the hemifield of the saccade vector (see Results), meaning that we also separated the vector and kinematic matching procedures in this monkey for each hemifield individually. This way, we did not mask any possible quantitative effects of saccade peak velocity on mislocalization amplitude, but the results were always qualitatively the same across hemifields (and across animals).

Figure 2B shows, for the same monkey and session as in Fig. 2A, example accepted saccades. Each panel shows either horizontal (top) or vertical (bottom) eye position aligned on saccade

onset, and the different colors show the different probe flash time classifications of Fig. 2A.
Figure 2C, in turn, shows all start and end eye positions from all trials of all sessions of the
same monkey. These trials passed the vector and kinematic matching (for the investigation
of saccade direction influences on visual mislocalization), and they, therefore, went into
further analysis.

For investigating the influences of saccade target appearance on mislocalization strength, we
applied the same binning strategy above in order to achieve vector and kinematic matching,
but we now compared saccades to either low or high spatial frequency Gabor gratings. We,
thus, pooled across all saccade directions. We included only those trials from the main
sequence bins that contained at least one trial repetition from each of the two saccade-
target image conditions (low and high spatial frequency) of each probe flash time. This
ensured that saccades were matched for both metrics and kinematics across the different
image conditions of the saccade target.

We next analyzed the report saccade endpoints, which constituted our experimental
measure of perisaccadic visual mislocalization. We first focused on the effects of saccade
direction. For each flash time and saccade direction, we measured the final eye position of
the report saccade on every accepted trial, and we then plotted the raw two-dimensional
data (e.g. Fig. 3D in Results) relative to the geometry of the saccade vector and probe flash
location. We also collected summary statistics within each monkey. Specifically, for each
probe flash location and time (for a given saccade direction), we calculated the mean and
SEM of all reported probe flash locations. This allowed us to visualize the pattern of
mislocalization (e.g. Fig. 3E, left in Results).

We quantified mislocalization strength as follows. For each flash location, we used the
baseline condition to know the average reported location by the monkey when perception
was expected to be veridical. Then, for each trial of the same flash location but for one of
the experimental perisaccadic flash times, we measured the Euclidean distance between the
reported location of this trial and the mean reported location of the baseline condition (e.g.
Fig. 3E, right in Results). This allowed us to obtain average mislocalization across trials of a
given flash time relative to any systematic error that might have existed in the report
saccade in the absence of any perisaccadic mislocalization. In other words, because
memory-based reporting is inevitable in mislocalization paradigms, and because memory-
guided saccades exhibit systematic errors, we always assessed mislocalization relative to
such systematic errors. We also inspected the baseline reports themselves, for example to
confirm that they were always more veridical than the reports for perisaccadic flashes, and
to also confirm that differences in them (as a function of, say, saccade direction) were not
sufficient to explain the differences in peak mislocalization effects that we reported in
Results.

To summarize the effects of saccade direction on visual mislocalization strength, in one
analysis, we averaged the Euclidean distance measure described above across all trials that
involved either upward or downward saccades for each probe flash time, respectively.
Upward saccades in this kind of analysis included oblique upward/rightward and oblique
upward/leftward saccades, in addition to purely vertical upward saccades. Similarly,
downward saccades also included the oblique saccades with a downward component. In this

comparison, we averaged across all three probe flash locations of a given saccade direction.
In related analyses, we also measured the mislocalization strength (Euclidean distance to the
reports in the baseline condition) for each saccade direction separately. Again, we pooled all
flash locations per saccade direction together in this analysis, but we always treated flash
times separately. Of course, we had previously inspected all the raw data of individual flash
locations (e.g. Figs. 3D, 4A, B in Results) to justify pooling them in our summary analyses.
458

We also further summarized the difference in mislocalization strength between upward and
downward saccades in a given analysis. To do so, we calculated the mathematical difference
(at any given flash time) between the Euclidean distance measure (to the respective baseline
condition of each flash location) obtained for upward saccades and the Euclidean distance
measure obtained for downward saccades. We then plotted this difference measure across
flash times. Again, this is a similar approach to how we reported changes in mislocalization
strength across different experimental conditions in our recent human work (Baumann et al.,
2024).
467

In a separate comparison, we focused solely on the trials with horizontal primary saccades.
In this case, we compared the Euclidean distance measure for probe flash locations in the
upper versus lower visual fields. Thus, for such horizontal saccades, we compared the
mislocalization strength between the two oblique probe flash locations associated with
these saccades (by definition of the task, one oblique probe flash location would have been
in the upper visual field, and ahead of the saccade target location, and the other would have
been symmetrically in the lower visual field).
475

476   We also performed an investigation of the influences of the saccade target appearance on
477   perisaccadic visual mislocalization. To do so, we averaged the Euclidean distance measures
478   for all saccade directions and probe flash locations, but now separated trials based on the
479   spatial frequency of the saccade target.
480

We statistically compared different conditions using a two-way ANOVA on the Euclidean
distance measure, with the two factors of the ANOVA being probe flash time relative to
saccade onset and either saccade direction, flash location, or saccade target appearance,
depending on the particular comparison that we were making. We always performed these
analyses within each monkey individually, to be able to compare the outcomes across the
individual animals. Although we consistently presented monkey M's leftward and rightward
saccades separately in all figures, we did not distinguish between them in the statistical
analyses, as this separation did not affect the statistical results. We also statistically
compared baseline reports under different comparison conditions. For example, we checked
whether the errors in baseline reports (relative to the true flash location) were different
whether saccades were upward or downward, and we did so by performing (for each
monkey) a t-test. Details are provided in the relevant sections of Results, and the statistics
were performed with IBM SPSS (Version 29.0).

Results

We aimed to establish a robust and flexible visual mislocalization paradigm in rhesus macaque monkeys. We tested this paradigm on three animals, and they all exhibited clear perisaccadic mislocalization, with expected recovery time courses. These animals also all replicated prior human findings of stronger mislocalization with upward saccades (Grujic et al., 2018), as well as stronger mislocalization for upper visual field flashes (with horizontal saccades) (Baumann et al., 2024). We also found subtle effects of saccade target image appearance on mislocalization strength (Baumann et al., 2024), but these effects were not as consistent as with humans, and they were also not as strong as the effects of saccade directions and flash locations in the same animals. We summarize all of these observations below.

Monkeys reported perisaccadic flashes as being closer to the saccade target
We trained our monkeys to report the location of a brief (~12 ms) probe flash presented at different perisaccadic time points (Figs. 1, 2). They did so by directing their gaze, via a memory-guided saccade, to the remembered flash location after an instruction to do so (Fig. 1). Note that such memory-guided reporting of percepts is present even in classic human invocations of this same task (Ross et al., 1997; Lappe et al., 2000). To confirm that there was indeed mislocalization, we interleaved baseline trials in which the flash was long-lasting, and thus clearly visible after primary saccade end (Methods). Therefore, any errors in reporting for this baseline flash condition should have been restricted to only those errors (whether systematic or variable) associated with memory-based behavioral reports (White et al., 1994; Willeke et al., 2019; Willeke et al., 2022), and not directly related to perisaccadic mislocalization.

Figure 3A shows example results from monkey F when generating an oblique upward/rightward primary saccade. The probe flash location in this case was rotated 45 deg counterclockwise from the saccade vector direction. In the figure, the probe flash is indicated by a white square, but it was not actually visible at the time of the report saccade. Each trace in the figure shows eye position from one example trial, and different traces show different trials. We plotted example trials from the baseline condition in green, and example trials from a perisaccadic condition (the one with the probe flash occurring 30 ms after saccade onset) in yellow. As can be seen, the primary saccades (oblique upward/rightward movements and directed towards the grating center) were all the same whether the trials were baseline trials or perisaccadic flash trials; that is, the eye position traces nicely overlapped with each other. This is expected given our eye movement controls mentioned earlier (Methods). Importantly, the report saccades were very different depending on the properties of perisaccadic visual stimulation. In the baseline trials, the report saccades had a veridical direction (upward in this case); for brief perisaccadic flashes occurring 30 ms after saccade onset, the reported locations were strongly shifted backward along the reverse direction of the primary saccade vector (compare the black oblique arrows). Note that the baseline saccades were still memory-guided, just like the report saccades in the perisaccadic test trials. Therefore, the large difference between report saccades in the two conditions in Fig. 3A was attributed to brief, perisaccadic visual stimulation in one of the conditions, and not due to the memory-guided nature of the report saccades. Also note that there was

larger variance in the report saccades in the perisaccadic test condition; this is also true in human perisaccadic mislocalization studies (Baumann et al., 2024). Thus, this monkey experienced strong backward perisaccadic mislocalization for brief flashes presented 30 ms after saccade onset, and this mislocalization happened with an oblique (rather than cardinal) saccade.

Figure 3 Two-dimensional perisaccadic "compression" in rhesus macaque monkeys. (A) Example eye position traces from monkey F when generating an oblique (upward/rightward) primary saccade, followed by a second report saccade. Individual traces show the entire eye movement sequence per trial (Methods), and we selected only trials with probe flashes directly above the saccade target location. The probe flash is schematized with a white square, but it was not actually visible at the time of the second saccade (the grating could also be different across trials; Methods). Green traces show baseline trials in which the probe flash remained visible for a prolonged period after the end of the primary saccade, but it was still invisible at the report saccade time (Methods). The monkey's report saccade was largely accurate, with a small systematic error expected from memory-guided eye movements. For perisaccadic flashes (yellow traces), there was a strong distortion in the opposite direction from the saccade vector (compare black oblique arrows). **(B, C)** Example trials from the same monkey for two other flash locations. There was always reverse mislocalization. **(D)** Across all sessions from the same monkey, and for all probe flash times and the same oblique saccade direction, we plotted the two-dimensional reported locations. On baseline trials (black), the percepts were mostly veridical (with expected small, but systematic, errors associated with memory-guided saccades). For perisaccadic flashes (colored), there was maximal mislocalization at 30 ms, with a gradual temporal recovery. This is consistent with human mislocalization time courses. The gray squares indicate true flash locations. **(E)** The main plot (left) shows mean and SEM summaries of the data of **D**. For longer and longer times, gradual recovery towards baseline performance occurred at each probe flash location. On the right, we magnified the data for one probe flash location, to demonstrate our summary measures in subsequent figures: we characterized mislocalization strength by calculating the Euclidean distance between reports on mislocalization trials to reports on baseline trials (Methods). In separate control analyses, we also checked the errors in baseline reports themselves (from true flash location).

We also checked the other probe flash locations associated with this same saccade direction.
Example traces from the same monkey and session are shown in Fig. 3B, C. Once again,
there was always reverse mislocalization opposite in direction from the saccade vector.
Specifically, in Fig. 3B, the probe flash was directly ahead of the saccade target, and the
reported locations were generally in the same direction as the flash but closer to the
saccade target location (mislocalized backward in direction). Similarly, with the probe flash
location rotated clockwise relative to the saccade vector (Fig. 3C), the reported locations
were now deviated both leftward and downward relative to the baseline reports (resulting in
a mislocalization vector directly opposite the upward and rightward vector of the primary
saccade). Thus, the mislocalization that this monkey experienced perisaccadically was always
backward relative to the saccade vector direction, regardless of flash location.
589
Note also that there were systematic errors in the baseline report saccades themselves in
Fig. 3A-C, which are expected (White et al., 1994; Willeke et al., 2019). For example, in Fig.
3C, a pure rightward report saccade was associated with a slight upward bias in eye position
endpoints, along with saccade hypometry (White et al., 1994; Willeke et al., 2019; Willeke et
al., 2022). However, the perisaccadic mislocalization errors were always larger than these
baseline report errors. Moreover, the perisaccadic mislocalization errors had a clear
systematic relationship in direction relative to the saccade vector direction, as we also
describe in more detail below. Thus, the perisaccadic mislocalization effects of Fig. 3A-C
were not fully explained by systematic errors associated with the reporting process itself.
599
Across all trials and all probe flash times and locations for the same oblique saccade
direction, we found systematic mislocalization that gradually recovered with time from
saccade onset. This is shown in Fig. 3D. Different colors indicate the different probe flash
times, and individual data points are individual trials. There was gradual recovery in
mislocalization towards the baseline report locations when the probe flash time increased.
This systematic trend can also be seen in Fig. 3E (left), in which we plotted the mean and
SEM summary statistics of the same data. In the same figure (Fig. 3E, right), we also
explained our measure of mislocalization strength that we used in subsequent figures.
Specifically, mislocalization strength was defined as the Euclidean distance to the baseline
report location for each probe flash position (Methods).
610
611 Thus, the results so far suggest that, at least for monkey F and oblique (upward/rightward)
612 saccades, we replicated human evidence of reverse perisaccadic visual mislocalization.
613 Importantly, we did so for a non-horizontal saccade direction (Grujic et al., 2018), paving the
614 way for generalized use of this paradigm in neurophysiological experiments.
615
616

Perisaccadic mislocalization was strongest for upward saccades
To demonstrate the robustness of the above results, we explored further properties of
perisaccadic mislocalization. We first focused on the influence of saccade direction. In
previous work, we found that backward mislocalization in humans was particularly strong for
upward saccades (Grujic et al., 2018). If our current paradigm is robust, and if macaques are
indeed a suitable animal model for exploring this property of perisaccadic mislocalization
with neurophysiology, then we should also obtain stronger perisaccadic visual
mislocalization in our animals for upward saccades. This was indeed the case.

Consider the results of Fig. 4A, B, which are formatted similarly to Fig. 3D. Here, we elected
to demonstrate observations from a second monkey, A, in order to additionally demonstrate
that our paradigm worked consistently across multiple animals. In Fig. 4A, the animal
generated purely upward saccades. The mislocalization was backward relative to the saccade
direction, and it was stronger for smaller perisaccadic probe flash times than for later times,
as expected. Importantly, the mislocalization seen in Fig. 4A was substantially stronger than
that observed for purely downward saccades in the same animal. Specifically, in Fig. 4B, we
plotted the data for purely downward saccades in the same format as in Fig. 4A. There was
still reverse mislocalization opposite the saccade direction, but the peak mislocalization
strength (for 30 ms perisaccadic probe flash times) was significantly weaker than in Fig. 4A.
This fact is rendered clearer by noting Fig. 4C. Here, we plotted the average (and SEM)
reported locations for 30 ms perisaccadic probe flashes for the two saccade directions
together in the same plot. To facilitate comparison of the effect sizes in the two cases, we
rotated all data to a schematic representation in which the saccade vector was to the right,
and the probe flashes were further ahead of it. We used exactly the same approach to
visualize results from different saccade directions in our earlier human studies (Grujic et al.,
2018). As can be seen, relative to the baseline reports of each condition (black symbols and
colored connecting arrows) there was clearly stronger perisaccadic mislocalization (in the
backward, reverse direction) for upward saccades. Quantitatively, for purely upward
saccades, the average Euclidean distance measure across all three flash locations in this
figure was 5.8 deg, whereas it was only 3.5 deg for purely downward saccades. Therefore,
there was stronger mislocalization for the upward saccades. Moreover, this difference in
mislocalization strength between upward and downward saccades at peak mislocalization
(30 ms flash times) was clearly larger than any systematic differences in the baseline report
locations between upward and downward saccades: the Euclidean distance between the
baseline report locations (black symbols) for purely upward and purely downward saccades
in Fig. 4C was only 0.51 deg (again averaging the respective distances across all three flash
locations); this was smaller than the difference in mislocalization strengths between the
upward and downward saccades (also see Figs. 5, 6). Thus, we conclude that, at least in this
animal, there was indeed stronger perisaccadic visual mislocalization for upward saccades,
replicating previous human observations (Grujic et al., 2018). We also conclude that this
effect was not explained by differences in baseline reports between the two conditions.
658
659
660
661

662
663
Figure 4 Stronger perisaccadic mislocalization for upward saccades. (A) Similar analysis to that in Fig. 3D but for
purely upward saccades in monkey A. Reverse mislocalization can be seen, consistent with Fig. 3. **(B)** In the same
animal, the strength of reverse mislocalization was weaker for purely downward saccades. This can be seen by
the smaller (than in **A**) deviation in the reports for the 30 ms perisaccadic flashes from the baseline reports. **(C)**
Comparison of the strength of mislocalization for the two saccade directions in **A**, **B**. Here, we rotated the saccade
vectors of **A**, **B** to a single schematic representation (shown here as a rightward saccade), like we did previously
(Grujic et al., 2018). Purple symbols (along with SEM error bars) indicate reports with the 30 ms perisaccadic
flashes, and black symbols (along with SEM error bars) indicate baseline reports. Red and blue connecting arrows
indicate whether the saccade was upward (blue) or downward (red). The blue arrows were always longer than
the red arrows, indicating a stronger mislocalization (relative to baseline) for the upward than for the downward
saccades. Note that for each flash location, there was generally a smaller difference in baseline report locations
between upward and downward saccades than the difference in mislocalization strength (arrow length). Thus,
the stronger mislocalization for upward saccades (relative to baseline reports) was not explained by an equally
large difference in baseline reports across the two saccade directions.

We next summarized the above observations across all three animals. In Fig. 5A, B, we
measured the Euclidean distance of flash localizations (Fig. 3E, right and Methods) as a
function of probe flash time from saccade onset. We did so separately for all upward
saccades and for all downward saccades. In the two shown animals in these two panels, all
upward saccades meant purely upward and two oblique upward saccade directions, and all
downward saccades meant purely downward and two oblique downward saccade directions
(see the inset schematics in Fig. 5A, B). We also included all flash locations per saccade
direction. That is, for each saccade direction and saccade location, we collected the
Euclidean distance, and we then averaged across all measurements. In both animals, at peak
mislocalization (flashes occurring 30 ms after saccade onset in our experiments), there was
stronger perisaccadic visual mislocalization for upward saccades, consistent with human
observations (Grujic et al., 2018). We also reached the same conclusions when treating each
saccade direction individually (Fig. 5C, D).

696
697 We also replicated these results in monkey M. In this monkey, we separated the visual
698 hemifields towards which the primary saccades were directed because this monkey had
699 significantly slower saccades with a rightward component, making it difficult to match
700 kinematics across hemifields (Methods). The monkey again exhibited stronger perisaccadic
701 visual mislocalization for upward saccades (Fig. 5E, F). This can also be seen in Fig. 5G, with

the added observation being that perisaccadic visual mislocalization was stronger for the faster leftward saccades. This is also consistent with human observations (Ostendorf et al., 2007).

Figure 5 Consistently stronger perisaccadic mislocalization for upward saccades. (A) In monkey A, we measured the Euclidean distance of perisaccadic reports (relative to baseline reports) as a function of flash time from saccade onset. The two curves show mislocalization strengths when considering only trials with either upward (blue) or downward (red) saccades. The inset icons clarify the saccade vectors included for each curve (Methods), as well as their relevant flash locations (3 per saccade direction). There was stronger mislocalization for upward saccades. **(B)** Similar results for monkey F. This monkey also showed more rapid recovery dynamics, with time, for the upward saccades. **(C, D)** Same data as in **A**, **B**, but considering each saccade direction individually. In each plot, the angle indicates the direction of the saccade (see inset icon examples), and the radial value indicates the Euclidean distance measured for 30 ms flash times. There was stronger mislocalization for upward saccades. **(E, F)** Similar analyses to **A**, **B**, but for monkey M. In this monkey, we separated saccades with a leftward (**E**) or rightward (**F**) component. In both panels, the monkey still exhibited stronger mislocalization for upward saccades. **(G)** Same as **C**, **D**, but for monkey M. Again, there was stronger mislocalization for upward saccades. Note that there was also stronger mislocalization for leftward saccades. These were the faster eye movements (approximately 550 deg/s versus approximately 350 deg/s peak velocity), suggesting an interaction between saccade speed and mislocalization strength (Ostendorf et al., 2007). **(H)** We summarized the effects of saccade direction by calculating the difference in mislocalization strength between upward and downward saccades. For all monkeys, there was stronger mislocalization for upward saccades, especially at peak mislocalization times. Note that for monkey M, we show two curves for either leftward (l) or rightward (r) saccades. Error bars indicate SEM.

Statistically, all of the above results were robust at the individual monkey level. We
conducted a two-way ANOVA to examine the effects of saccade direction and probe time
relative to saccade onset on the Euclidean distance to the baseline report. There was a
significant main effect of saccade direction for monkeys A ($F_{(1,7576)}=421.07$, $p<0.001$) and
M ($F_{(1,20896)}=812.77$, $p<0.001$), indicating that the direction of the saccade had a
significant impact on perisaccadic mislocalization. Monkey F did not show a significant main
effect for saccade direction ($F_{(1,15999)}=1.92$, $p=0.16$), but this was only because this
monkey exhibited faster recovery times than the other two monkeys (Fig. 5B). Indeed, this
monkey, as well as the other two, showed a significant interaction between saccade
direction and probe flash time relative to saccade onset (monkey A: $F_{(2,7576)}=202.37$,
$p<0.001$; monkey F: $F_{(2,15999)}=265.66$, $p<0.001$; monkey M: $F_{(2,20896)}=333.72$, $p<0.001$),
indicating that the effect of saccade direction depended on the time point. There was also a
significant main effect of probe time relative to saccade onset in all three monkeys (monkey
A: $F_{(2,7576)}=1870.19$, $p<0.001$; monkey F: $F_{(2,15999)}=3371.03$, $p<0.001$; monkey M:
$F_{(2,20896)}=4434.12$, $p<0.001$), suggesting that saccadic mislocalization varied across
different time points (as expected). Post-hoc comparisons using Bonferroni corrections for
probe time relative to saccade onset revealed significant differences between all time points
for all monkeys, with visual mislocalization decreasing progressively from 30 to 130 ms
($p<0.001$). All of these statistical results are summarized graphically in Fig. 5H.

Finally, we also checked the systematic errors in baseline reports themselves. Specifically,
since memory-guided saccades of different directions have different systematic error
magnitudes (Hafed and Chen, 2016), we asked whether the differences in mislocalization
strengths between upward and downward saccades that we saw in Fig. 5 were entirely due
to differences in baseline report errors. For each monkey, we measured the error (from true
flash location) in the baseline report saccade as a function of saccade direction (Fig. 6A). We
did this only for the baseline trials in which perception was expected to be veridical
(Methods). As anticipated (e.g. Figs. 3, 4), there were non-zero absolute errors in the
baseline saccades. However, these errors were significantly smaller than the peak
mislocalization effects (also seen in Figs. 3, 4). For example, monkey F had a peak
mislocalization amplitude of around 4-5 deg (Fig. 5B), but only a baseline report error of
around 1.5-1.6 deg. Most importantly, in all animals, the difference in baseline report errors
between upward and downward saccades (Fig. 6B), although significant (monkey A:
$t_{(647)}=-6.04$, $p<0.001$; monkey F: $t_{(1323)}=-5.03$, $p<0.001$; and monkey M:
$t_{(689)}=-4.11$, $p<0.001$ for leftward saccades and $t_{(704)}=-5.78$, $p<0.001$ for rightward
saccades; independent samples t-tests), was of the opposite sign from the mislocalization
effects comparing upward and downward saccades (purple samples in Fig. 6B and Fig. 5).
Thus, stronger perisaccadic mislocalization for upward saccades was not explained by
systematic differences in baseline report errors.

Figure 6 Different patterns of baseline report errors between upward and downward saccades than for mislocalization. (A) For each monkey, we measured the Euclidean distance between the baseline report saccade endpoints and the true flash locations in the data of Fig. 5. We did this separately for either upward or downward saccades. Error bars denote SEM. Note how there was still some error from the true flash location in the baseline reports in which no mislocalization was to be anticipated (also visible in Figs. 3, 4). This is expected from memory-guided saccades (White et al., 1994; Willeke et al., 2019; Willeke et al., 2022). However, such error was always smaller than peak mislocalization error (also visible in Figs. 3, 4). **(B)** For each monkey, we took the difference in the errors in **A** between upward and downward saccades (bar plots). This difference revealed slightly larger errors for downward memory-guided saccades, consistent with earlier observations (Hafed and Chen, 2016). In contrast, the purple data points are the results from the time of peak mislocalization in Fig. 5. There was a clearly larger difference between upward and downward saccades during peak perisaccadic visual mislocalization than during baseline reports (and also with an opposite sign). Therefore, the results of Fig. 5 were not driven by differences in baseline report errors between upward and downward saccades.

Therefore, our results so far indicate that all three monkeys experienced perisaccadic visual mislocalization, and that they all demonstrated an effect of saccade direction, consistent with human observations (Grujic et al., 2018). A quantitative comparison of effect magnitudes as a function of saccade speeds was also afforded to us by monkey M, and the results were again consistent with human observations (Ostendorf et al., 2007). This reinforces our interpretation of the flexibility and robustness of our paradigm.

Perisaccadic mislocalization was stronger for upper visual field flashes
We next explored the effect of visual field location of the flashes relative to horizontal saccades. In humans, we recently found that, with such horizontal saccades, reverse mislocalization was stronger for upper visual field flashes (ahead of the saccade target location) than for symmetrically located lower visual field ones (Baumann et al., 2024). This was also the case in our monkeys. Specifically, in Fig. 7A-D, we analyzed mislocalization strength for only horizontal saccades in each monkey. This time, we separated the trials based on whether the perisaccadic flash location was above or below the horizon. That is, if the saccade was to the right, our paradigm included three potential flashes: one directly ahead of the saccade target (still on the horizontal axis), and two oblique ones (Fig. 1, Methods). The two oblique flashes were, by definition, either in the upper or lower visual field relative to the saccade direction, and we compared these two oblique flash positions in the current analysis. The results demonstrated that there was indeed generally stronger mislocalization associated with upper rather than lower visual field flashes.

Figure 7 Stronger perisaccadic visual mislocalization for upper visual field probe flashes and horizontal saccades. (A) For monkey A, we only considered horizontal primary saccade directions. We then compared mislocalization strength (Euclidean distance to baseline reports) for probe flashes above or below the saccadic vector (see Methods and inset schematic). There was stronger mislocalization for upper visual field flashes, like in humans (Baumann et al., 2024). **(B)** In monkey F, the effect at peak mislocalization (flashes 30 ms from saccade onset) was much weaker than in monkey A, but with a similar trend. This monkey also had rapid time dynamics, like in Fig. 5, causing a reversal at longer times. **(C, D)** Monkey M replicated the results of monkey A regardless of the direction of the horizontal saccades. **(E)** Summary of the results in **A**-**D** shown here as the difference in mislocalization strength between upper and lower visual field flash locations, and as a function of flash time (like in Fig. 5H). **(F)** Errors in baseline reports from true flash location, like in Fig. 6A. Here, we plotted these errors for either the upward or downward flash conditions. The errors were always smaller in magnitude than the peak mislocalization effects in **A**-**E** (consistent with Figs. 3, 4). **(G)** More importantly, the difference in these errors between upward and downward flashes did not explain the differences in peak mislocalization. This panel is formatted similarly to Fig. 6B: the bars denote the differences in baseline report errors between upper and lower visual field flashes, and the purple symbols show the peak mislocalization differences from **A**-**E**. Differences in baseline reports did not explain differences in peak mislocalization strengths (except for monkey F where there were already no differences in mislocalization strength). Error bars denote SEM.

Statistically, we performed a two-way ANOVA to evaluate the effects of probe location and probe time relative to horizontal saccade onset on the Euclidean distance to the baseline report. There was a significant main effect of probe location for all three monkeys (monkey A: $F_{(1,1632)}=410.01$, $p<0.001$; monkey F: $F_{(1,1214)}=65.2$, $p<0.001$; monkey M: $F_{(1,9687)}=626.66$, $p<0.001$) indicating that the probe location had an impact on the strength

of perisaccadic visual mislocalization. The main effect of probe time relative to saccade onset was also significant (monkey A: $F_{(2,1632)}=552.54$, $p<0.001$; monkey F: $F_{(2,1214)}=714.44$, $p<0.001$; monkey M: $F_{(2,9687)}=1652.95$, $p<0.001$), suggesting that perisaccadic mislocalization varied across different time points. Finally, the interaction between probe location and probe time relative to saccade onset was also significant for all three monkeys (monkey A: $F_{(2,1632)}=21.43$, $p<0.001$; monkey F: $F_{(2,1214)}=21.93$, $p<0.001$; monkey M: $F_{(2,9687)}=70.51$, $p<0.001$), indicating that the effect of probe location was dependent on flash time. These results are summarized in Fig. 7E showing difference measures similar to ones that we used earlier for summarizing saccade direction effects.

We also checked whether systematic differences (between upper and lower visual field flashes) in baseline report errors accounted for the above results. Specifically, just like in Fig. 6A, we plotted the Euclidean distance between baseline reports and true flash location for the upward and downward flash conditions of this analysis. The results are shown in Fig. 7F. Consistent with Fig. 6A, the absolute errors in baseline reports were smaller than the peak mislocalization effects; this is also consistent with the raw example data shown in Figs. 3, 4, demonstrating clear recovery with time from saccade onset (towards the baseline report locations). Importantly, the differences in baseline report errors between upward and downward flashes were not correlated with the differences in peak mislocalization strength (Fig. 7G). Thus, the effects of upper and lower visual field flashes in association with horizontal saccades were likely perisaccadic and not an artifact of report saccade errors.

Perisaccadic mislocalization strength was subtly modulated by the saccade target appearance

Finally, we explored whether mislocalization strength depended on the visual appearance of the saccade target (Methods). In two out of the three monkeys, there was a very subtle difference in mislocalization strength as a function of saccade target appearance; however, this dependence was not the same in both animals. Consider for example, Fig. 8A. The figure shows slightly stronger mislocalization for the low spatial frequency grating as the saccade target. However, Monkey F showed the opposite effect (Fig. 8B). Figure 8C, D shows the data of monkey M, showing no image dependency for the time bin with peak mislocalization, and neither for leftward nor for rightward saccades. These findings for the image dependence are smaller than the dependence for saccade direction.

Statistically, we performed a two-way ANOVA to investigate the effects of saccade target appearance and probe time relative to saccade onset on the Euclidean distance to the baseline report. There was a significant main effect of saccade target appearance for two of the three monkeys (monkey A: ($F_{(1,14947)}=1.15$, $p=0.28$); monkey F: $F_{(1,23824)}=33.05$; monkey M ($F_{(1,21454)}=20.32$, $p<0.001$) indicating that the saccade target had an impact on the strength of perisaccadic mislocalization. However, monkeys' M and F results were not consistent with the previous human results: these monkeys exhibited stronger mislocalization for a high spatial frequency grating. As expected, there was also a significant main effect of probe time relative to saccade onset (monkey A: $F_{(2,14947)}=3867.51$, $p<0.001$; monkey F: $F_{(2,23824)}=5413.83$, $p<0.001$; monkey M: $F_{(2,21454)}=4889.48$, $p<0.001$), suggesting that saccadic mislocalization varied across different time points. Additionally, the interaction between saccade target appearance and probe time relative to

893   saccade onset was significant for all three monkeys (monkey A: F(2,14947)=1.87, p<0.001;
894   monkey F: F(2,23824)=42.36, p<0.001; monkey M: F(2,21454)=4.77, p<0.001).
895
Thus, we cannot confidently conclude that there was an image dependence of
mislocalization strength in our monkeys. This is likely caused, at least in part, by the
differences in the ways monkeys and humans reported their percepts (see Discussion).
Indeed, analyzing the baseline report errors was particularly revealing (Fig. 8F, G): the
difference in magnitudes of baseline report errors between the different target image
appearances (Fig. 8G) was of the same order of magnitude as the difference in the peak
mislocalization strengths for the different saccade target images. This suggests that a
possible image dependence of perisaccadic mislocalization in our monkeys might have been
masked by errors in the memory-guided report saccades.

910
911
Figure 8 Inconsistent effects with respect to saccade target appearance. This figure is formatted identically to
Fig. 7. Here, the comparison was between the different saccade target appearances. Therefore, we pooled across
saccades of all directions (Methods) and only separated trials based on the appearance of the saccade target.
There were subtle differences (**A**-**E**), but the differences were similar to the differences in baseline report
saccades in the absence of perceptual mislocalization (**F, G**). Thus, the image dependence effect in perisaccadic
mislocalization (Baumann et al., 2024) was likely masked by systematic errors in report saccades. All other
conventions are similar to Fig. 7.

Discussion

We established a robust paradigm with which we could probe perisaccadic visual mislocalization in monkeys. Importantly, our paradigm allows mapping the two-dimensional landscape of mislocalization, and it can also be used with arbitrary saccade directions. This latter property enabled us to replicate the observation in humans (Grujic et al., 2018) that backward perisaccadic "compression" of flashes ahead of the saccade target is strongest for upward saccades. We also replicated the human observation (Baumann et al., 2024) of stronger backward mislocalization for upper visual field stimuli during horizontal saccades.

Our work builds on previous demonstrations that perisaccadic visual mislocalization can be studied in macaques (Dassonville et al., 1992; Jeffries et al., 2007; Klingenhoefer and Krekelberg, 2017; Weng et al., 2024). Our work also relates to interesting monkey variants of the classic double-step saccade task (Hallett and Lightstone, 1976; Becker and Jurgens, 1979; Ottes et al., 1984; Vliegen et al., 2005), in which the target of the second saccade in the double-step sequence was presented during the progression of the first eye movement (and thus could be susceptible to some mislocalization) (Van Grootel et al., 2012). However, in the context of visual mislocalization studies, our approach adds paradigm flexibility that we believe is necessary for further success in porting these behavioral measures of perisaccadic mislocalization to neurophysiology. For example, it may not always be possible neurophysiologically to probe only horizontal flash position mislocalization or only horizontal saccades. This is because electrode positions in topographic maps may be in off-axis representations. More importantly, mislocalization has a two-dimensional landscape (Kaiser and Lappe, 2004). Thus, with our work, we provide an additional stepping stone towards direct neurophysiological experiments exploiting essentially identical behavioral tasks to those classically used in humans.

Indeed, we were motivated by the particularly strong backward mislocalization for upward saccades in humans (Grujic et al., 2018), and that we think deserves further investigation. We suspect that this phenomenon might be related to the asymmetry of representation in the superior colliculus between the upper and lower visual fields (Hafed and Chen, 2016; Zhang et al., 2022), but this needs to be tested explicitly. In fact, besides recording activity related either to the saccade vector endpoint or to flash locations, we could make variants of our behavioral paradigm that can be more causal in nature. For example, we could replace the flash presentation with brief electrical microstimulation of the primary visual cortex, or higher-level cortical areas. Similarly, we could replace the visually-guided saccade with an electrically evoked saccade from the superior colliculus. This latter experiment would be particularly fascinating. For example, it is known that beyond a certain strength of electrical microstimulation in the superior colliculus, evoked saccades become more similar to each other under different stimulation parameters (Katnani and Gandhi, 2012). Thus, one could stimulate with two different "motor burst strengths" and ask how perisaccadic flashes are mislocalized under the two conditions (with the evoked saccades being the same). This would causally establish a functional role for sensory tuning in collicular motor bursts (Baumann et al., 2023) in perisaccadic perceptual phenomena that might, at least theoretically (Baumann et al., 2024), rely on saccade-related corollary discharge.

Similarly, we could perturb brain areas not implicated in corollary discharge and identify the
limits of visual contributions to perisaccadic mislocalization. For example, if monkey M's
asymmetry in saccade peak speeds that we alluded to above was due to a region
downstream of the superior colliculus, then it could be that the saccade motor commands
(from the colliculus), and their associated (colliculus-related) corollary discharge, were the
same for either rightward or leftward saccades in this monkey. If so, then the differences
that we saw in the magnitude of perisaccadic mislocalization across hemifields in this animal
would have been intrinsically visual in origin: the monkey would have had similar saccade
commands (at the collicular level) but different speeds of retinal-image shifts.

The above example variants of our paradigm demonstrate why we think that this paradigm
is useful for neurophysiology, and also why we think that it can meaningfully add to a wealth
of previous work exploring perisaccadic modulations in visual processing in the brain
(Duhamel et al., 1992; Nakamura and Colby, 2002; Krekelberg et al., 2003; Kusunoki and
Goldberg, 2003; Sommer and Wurtz, 2006; Churan et al., 2011, 2012; Zirnsak et al., 2014;
Neupane et al., 2016b, a, 2020; Qian et al., 2022). These prior experiments did not exploit a
simultaneous behavioral report by the animals, which our paradigm allows.

Of course, we do acknowledge that we did not explicitly test flash positions between the
initial and final fixation positions associated with the saccade (which should show forward
mislocalization). However, our use of multiple different saccade directions, coupled with the
consistent backward mislocalization seen in all cases, makes us confident that what we saw
was indeed that the prior literature referred to as "compression". Indeed, the vectors of
mislocalization that we observed (e.g. Figs. 3, 4) as a function of time (relative to the saccade
vector direction) were similar to those that we observed earlier in humans using multiple
different saccade directions (Grujic et al., 2018). Thus, we believe that our results are
consistent with those of (Klingenhoefer and Krekelberg, 2017). Importantly, our paradigm
can be easily modified to also include flashes nearer to the initial fixation position than the
saccade target.

We also agree with (Klingenhoefer and Krekelberg, 2017) that there can (and should) be
quantitative differences in the parameters of the observed perisaccadic mislocalization
between monkeys and humans. This is totally expected because factors like foveal
magnification and sensory transduction speeds, and others, can be quantitatively different
between the two species, even when they are qualitatively similar. Such quantitative
differences are inevitable given the different brain sizes. What is important is that the
qualitative equivalence is present, and this motivates using these animals in particular for
invasive neurophysiological investigations.

One caveat with our work is that we relied on memory-guided saccades to report flash
location. Such saccades have systematic errors in their landing positions (White et al., 1994;
Willeke et al., 2019), which could also depend on the visual field location of the flashes
(Hafed and Chen, 2016). Such saccades also have larger dispersions across trials (variable
errors) than visually-guided movements. Still, we chose to use memory-guided saccades
because we did not want to impose any prior assumptions about where the monkeys were
going to see the flashes. For example, if we had used "choice" targets, then we would have
been imposing constraints on the spatial reports of the animals. This is problematic for a

general two-dimensional paradigm like the one that we were trying to develop. Moreover, even manual pointing (via displayed cursors) by humans has systematic and variable errors (Willeke et al., 2019), and such pointing is generally used in mislocalization experiments (Grujic et al., 2018; Baumann et al., 2024). Thus, some amount of reporting errors is inevitable in any paradigm requiring a behavioral response from the subjects.

We also acknowledge that our choice to use memory-guided saccades could have come at the expense of having enough resolution to observe more subtle effects, like a potential dependence of perisaccadic mislocalization strength on the saccade target image appearance. Indeed, in parallel experiments with humans, we did find more systematic effects across most subjects than with the monkeys in the current study (Baumann et al., 2024). Thus, the inconsistency of the image dependence effects that we saw with our monkeys (Fig. 8) could reflect either added contamination by errors in memory-guided saccades, species differences from humans, or the need to test more monkeys with our current paradigm before reaching a conclusion about an image dependence of perisaccadic mislocalization in the animals.

One possibility to avoid the noise associated with memory-guided saccades could be to modify our paradigm slightly: we could adopt the approach of (Klingenhoefer and Krekelberg, 2017; Weng et al., 2024). That is, at the end of every trial, we would display a very dense two-dimensional array of grid positions, and the monkeys would simply choose which grid position matched the perceived flash location. We do not expect this to cause an increased training burden to the task. In any case, for the effects of saccade directions on mislocalization strengths, our paradigm was sufficient.

We also chose to not test presaccadic flashes. This was motivated purely by the pragmatic decision to increase data yield. By presenting flashes contingent on detecting a saccade, we were essentially ensuring obtaining a usable data point for every saccade trial accepted into the analysis. On the other hand, if we had randomized flash times independent of the saccadic reaction time, we would have needed to collect more trials, and then recalculate flash times relative to saccade onset. This should work since it does not alter the paradigm dramatically, and it can also allow obtaining a full time course for perisaccadic mislocalization in monkeys. Moreover, we have previously used both approaches successfully in our experiments on perisaccadic visual suppression, in both humans and monkeys (Hafed and Krauzlis, 2010; Chen and Hafed, 2017; Idrees et al., 2020), and also in our perisaccadic perceptual mislocalization experiments in humans (Grujic et al., 2018; Baumann et al., 2024). Having said that, one has to carefully interpret trials in which the flash happened too long before saccade onset, especially because such a condition gives rise to a complete resetting of saccade plans (Buonocore and Hafed, 2023).

In all, we established a robust, two-dimensional perisaccadic mislocalization paradigm that is particularly useful for neurophysiological experiments with macaque monkeys.

Competing interests statement

The authors declare no competing interests.

Acknowledgements

We thank Tong Zhang, Tatiana Malevich, and Antimo Buonocore for help in data collection. We were funded by the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation) through the SFB 1233 Robust Vision (project number: 276693517) as well as the SPP 2205 Evolutionary Optimization of Neuronal Processing (project number: HA 6749/3-2) and the SPP 2411 Sensing LOOPS: Cortico-subcortical Interactions for Adaptive Sensing (project number: HA 6749/11-1).

References

Bahill AT, Clark MR, Stark L (1975) The main sequence, a tool for studying human eye movements. Mathematical Biosciences 24:191-204.

Baumann MP, Denninger AF, Hafed ZM (2024) Perisaccadic perceptual mislocalization strength depends on the visual appearance of saccade targets. Journal of Neurophysiology.

Baumann MP, Bogadhi AR, Denninger AF, Hafed ZM (2023) Sensory tuning in neuronal movement commands. Proc Natl Acad Sci U S A 120:e2305759120.

Becker W, Jurgens R (1979) An analysis of the saccadic system by means of double step stimuli. Vision Res 19:967-983.

Beeler GW, Jr. (1967) Visual threshold changes resulting from spontaneous saccadic eye movements. Vision Res 7:769-775.

Berman RA, Wurtz RH (2010) Functional identification of a pulvinar path from superior colliculus to cortical area MT. J Neurosci 30:6342-6354.

Berman RA, Wurtz RH (2011) Signals conveyed in the pulvinar pathway from superior colliculus to cortical area MT. The Journal of neuroscience : the official journal of the Society for Neuroscience 31:373-384.

Berman RA, Joiner WM, Cavanaugh J, Wurtz RH (2009) Modulation of presaccadic activity in the frontal eye field by the superior colliculus. J Neurophysiol 101:2934-2942.

Berman RA, Cavanaugh J, McAlonan K, Wurtz RH (2017) A circuit for saccadic suppression in the primate brain. J Neurophysiol 117:1720-1735.

Brainard DH (1997) The Psychophysics Toolbox. Spatial vision 10:433-436.

Buonocore A, Hafed ZM (2023) The inevitability of visual interruption. J Neurophysiol 130:225-237.

Buonocore A, Skinner J, Hafed ZM (2019) Eye Position Error Influence over "Open-Loop" Smooth Pursuit Initiation. J Neurosci 39:2709-2721.

Cai RH, Pouget A, Schlag-Rey M, Schlag J (1997) Perceived geometrical relationships affected by eye-movement signals. Nature 386:601-604.

Chen CY, Hafed ZM (2013) Postmicrosaccadic enhancement of slow eye movements. The
Journal of neuroscience : the official journal of the Society for Neuroscience 33:5375-
5386.
Chen CY, Hafed ZM (2017) A neural locus for spatial-frequency specific saccadic suppression
in visual-motor neurons of the primate superior colliculus. J Neurophysiol 117:1657-
1673.
Churan J, Guitton D, Pack CC (2011) Context dependence of receptive field remapping in
superior colliculus. J Neurophysiol 106:1862-1874.
Churan J, Guitton D, Pack CC (2012) Perisaccadic remapping and rescaling of visual responses
in macaque superior colliculus. PLoS One 7:e52195.
Dassonville P, Schlag J, Schlag-Rey M (1992) Oculomotor localization relies on a damped
representation of saccadic eye displacement in human and nonhuman primates. Vis
Neurosci 9:261-269.
Duhamel JR, Colby CL, Goldberg ME (1992) The updating of the representation of visual
space in parietal cortex by intended eye movements. Science 255:90-92.
Eastman KM, Huk AC (2012) PLDAPS: A Hardware Architecture and Software Toolbox for
Neurophysiology Requiring Complex Visual Stimuli and Online Behavioral Control.
Front Neuroinform 6:1.
Fuchs AF, Robinson DA (1966) A method for measuring horizontal and vertical eye
movement chronically in the monkey. J Appl Physiol 21:1068-1070.
Grujic N, Brehm N, Gloge C, Zhuo W, Hafed ZM (2018) Peri-saccadic perceptual
mislocalization is different for upward saccades. J Neurophysiol.
Hafed ZM, Krauzlis RJ (2010) Microsaccadic suppression of visual bursts in the primate
superior colliculus. J Neurosci 30:9542-9547.
Hafed ZM, Chen CY (2016) Sharper, Stronger, Faster Upper Visual Field Representation in
Primate Superior Colliculus. Curr Biol 26:1647-1658.
Hallett PE, Lightstone AD (1976) Saccadic eye movements towards stimuli triggered by prior
saccades. Vision Res 16:99-106.
Honda H (1989) Perceptual localization of visual stimuli flashed during saccades. Percept
Psychophys 45:162-174.
Idrees S, Baumann MP, Franke F, Munch TA, Hafed ZM (2020) Perceptual saccadic
suppression starts in the retina. Nat Commun 11:1977.
Jeffries SM, Kusunoki M, Bisley JW, Cohen IS, Goldberg ME (2007) Rhesus monkeys
mislocalize saccade targets flashed for 100ms around the time of a saccade. Vision
Res 47:1924-1934.
Judge SJ, Richmond BJ, Chu FC (1980) Implantation of magnetic search coils for
measurement of eye position: an improved method. Vision Res 20:535-538.
Kaiser M, Lappe M (2004) Perisaccadic mislocalization orthogonal to saccade direction.
Neuron 41:293-300.
Katnani HA, Gandhi NJ (2012) The relative impact of microstimulation parameters on
movement generation. J Neurophysiol 108:528-538.
Kleiner M, Brainard D, Pelli DG (2007) What's new in Psychtoolbox-3? (Abstract). Perception
36.
Klingenhoefer S, Krekelberg B (2017) Perisaccadic visual perception. J Vis 17:16.
Krekelberg B, Kubischik M, Hoffmann KP, Bremmer F (2003) Neural correlates of visual
localization and perisaccadic mislocalization. Neuron 37:537-545.

Kusunoki M, Goldberg ME (2003) The time course of perisaccadic receptive field shifts in the
lateral intraparietal area of the monkey. J Neurophysiol 89:1519-1527.
Lappe M, Awater H, Krekelberg B (2000) Postsaccadic visual references generate presaccadic
compression of space. Nature 403:892-895.
Latour PL (1962) Visual threshold during eye movements. Vision Res 2:261-262.
Nakamura K, Colby CL (2002) Updating of the visual representation in monkey striate and
extrastriate cortex during saccades. Proc Natl Acad Sci U S A 99:4026-4031.
Neupane S, Guitton D, Pack CC (2016a) Two distinct types of remapping in primate cortical
area V4. Nat Commun 7:10402.
Neupane S, Guitton D, Pack CC (2016b) Dissociation of forward and convergent remapping in
primate visual cortex. Curr Biol 26:R491-492.
Neupane S, Guitton D, Pack CC (2020) Perisaccadic remapping: What? How? Why? Rev
Neurosci 31:505-520.
Ostendorf F, Fischer C, Finke C, Ploner CJ (2007) Perisaccadic compression correlates with
saccadic peak velocity: differential association of eye movement dynamics with
perceptual mislocalization patterns. J Neurosci 27:7559-7563.
Ottes FP, Van Gisbergen JA, Eggermont JJ (1984) Metrics of saccade responses to visual
double stimuli: two different modes. Vision Res 24:1169-1179.
Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers
into movies. Spatial vision 10:437-442.
Qian N, Goldberg ME, Zhang M (2022) Tuning curves vs. population responses, and
perceptual consequences of receptive-field remapping. Frontiers in computational
neuroscience 16:1060757.
Robinson DA (1963) A Method of Measuring Eye Movement Using a Scleral Search Coil in a
Magnetic Field. IEEE Trans Biomed Eng 10:137-145.
Ross J, Morrone MC, Burr DC (1997) Compression of visual space before saccades. Nature
386:598-601.
Skinner J, Buonocore A, Hafed ZM (2019) Transfer function of the rhesus macaque
oculomotor system for small-amplitude slow motion trajectories. J Neurophysiol
121:513-529.
Sommer MA, Wurtz RH (2004) What the brain stem tells the frontal cortex. II. Role of the SC-
MD-FEF pathway in corollary discharge. J Neurophysiol 91:1403-1423.
Sommer MA, Wurtz RH (2006) Influence of the thalamus on spatial visual processing in
frontal cortex. Nature 444:374-377.
Tian X, Yoshida M, Hafed ZM (2018) Dynamics of fixational eye position and microsaccades
during spatial cueing: the case of express microsaccades. J Neurophysiol 119:1962-
1980.
Van Grootel TJ, Van der Willigen RF, Van Opstal AJ (2012) Experimental test of spatial
updating models for monkey eye-head gaze shifts. PLoS One 7:e47606.
Vliegen J, Van Grootel TJ, Van Opstal AJ (2005) Gaze orienting in dynamic visual double steps.
J Neurophysiol 94:4300-4313.
Weng G, Akbarian A, Clark K, Noudoost B, Nategh N (2024) Neural correlates of perisaccadic
visual mislocalization in extrastriate cortex. Nat Commun 15:6335.
White JM, Sparks DL, Stanford TR (1994) Saccades to remembered target locations: an
analysis of systematic and variable errors. Vision Res 34:79-92.

Willeke KF, Cardenas AR, Bellet J, Hafed ZM (2022) Severe distortion in the representation of
foveal visual image locations in short-term memory. Proc Natl Acad Sci U S A
119:e2121860119.
Willeke KF, Tian X, Buonocore A, Bellet J, Ramirez-Cardenas A, Hafed ZM (2019) Memory-
guided microsaccades. Nat Commun 10:3710.
Zhang T, Malevich T, Baumann MP, Hafed ZM (2022) Superior colliculus saccade motor bursts
do not dictate movement kinematics. Commun Biol 5:1222.
Zirnsak M, Steinmetz NA, Noudoost B, Xu KZ, Moore T (2014) Visual space is compressed in
prefrontal cortex before eye movements. Nature 507:504-507.
Zuber BL, Stark L (1966) Saccadic suppression: elevation of visual threshold associated with
saccadic eye movements. Exp Neurol 16:65-79.
Zuber BL, Stark L, Cook G (1965) Microsaccades and the velocity-amplitude relationship for
saccadic eye movements. Science 150:1459-1460.
1214