Organization of the apical extracellular matrix during tubular organ formation

J. Luke Woodward[1,¶], Jeffrey Matthew[1,¶], Rutuparna Joshi[1], Vishakha Vishwakarma[1], Ying Xiao[2],

and SeYeon Chung[1,*]

1 Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

2 Shared Instrumentation Facility, Louisiana State University, Baton Rouge, LA 70803, USA

¶ These authors contributed equally to this study.

*Author for correspondence: seyeonchung@lsu.edu

14   Abstract

The apical extracellular matrix (aECM) plays a critical role in epithelial tube morphogenesis

during organ formation, but its composition and organization remain poorly understood. Using

the *Drosophila* embryonic salivary gland (SG) as a model, we identify Papss, an enzyme that

synthesizes the universal sulfate donor PAPS, as a critical regulator of tube lumen expansion.

Papss mutants show a disorganized apical membrane, condensed aECM, and disruptions in

Golgi structures and intracellular trafficking. SG-specific expression of wild-type Papss, but not

the catalytically inactive form, rescues the defects in *Papss* mutants, suggesting that defects in

sulfation are the underlying cause of the phenotypes. Additionally, we identify two zona

pellucida (ZP) domain proteins, Piopio (Pio) and Dumpy (Dpy), as key components of the SG

aECM. In the absence of *Papss*, Pio is gradually lost in the aECM, while the Dpy-positive aECM

structure is condensed and dissociates from the apical membrane, leading to a thin lumen.

Mutations in *dpy* or *pio*, or in *Notopleural*, which encodes a matriptase that cleaves Pio to form

the luminal Pio pool, result in a SG lumen with alternating bulges and constrictions, with the loss

of *pio* leading to the loss of Dpy in the lumen. Our findings underscore the essential role of

sulfation in organizing the aECM during tubular organ formation and highlight the mechanical

support provided by ZP domain proteins in maintaining luminal diameter.

Introduction

Epithelial tubes are critical components of many key biological organs, including the
kidney, heart, and lungs. Aberrant tube size leads to common diseases and congenital disorders,
such as polycystic kidney disease, asthma, and lung hypoplasia. Tube morphogenesis is
regulated by multiple factors, including intracellular mechanisms, intercellular interactions, and
interactions between cells and the surrounding extracellular matrix (ECM). Specifically, events
at the apical (luminal) surface of a tube play an important role in regulating tube diameter and
growth, thereby influencing overall tube shape (Myat and Andrew, 2002). During tubular organ
formation, the surrounding epithelium secretes apical ECM (aECM) components into the lumen.
These aECM components form the luminal meshwork and interact with the apical membrane,
playing a critical role in epithelial morphogenesis and regulation of tube dimensions (Jazwinska
et al., 2003; Devine et al., 2005; Luschnig et al., 2006; Wang et al., 2006; Syed et al., 2012;
Dong et al., 2014; Drees et al., 2023; Gill et al., 2016; Cohen, Sparacio, et al., 2020).

The aECM has a distinct composition compared to the basal ECM, with a diverse array of
components (Zheng et al., 2020). ZP domain proteins, which are highly conserved across
species, are of particular interest and are characterized by a conserved 260 amino acid ZP
domain that mediates homo- or heteromeric polymerization into filaments. Initially identified in
the extracellular coat of mammalian oocytes, ZP domain proteins have been shown to play a
critical role in the organization of specialized apical structures in various tissues in species
ranging from worms to humans (Porter & Tamm, 1955; Yan et al., 2012; Legan et al., 1997; Op
De Beeck et al., 2012; Renner et al., 2007; McAllister et al., 1994; Wong et al., 2000; Cohen,
Bermudez, et al., 2020; Plaza et al., 2010; Jovine et al., 2005; Litscher & Wassarman, 2020;
Wilkin et al., 2000; Ghosh & Treisman, 2024). Chitin, a biopolymer of N-acetylglucosamine, is
another major aECM component that is abundant in invertebrates and some vertebrates (Tang et
al., 2015; Mei et al., 2024). In many *Drosophila* and *C. elegans* tissues, chitin forms a network of
fibers that provide mechanical support and maintain the integrity of the epithelial layer (Araujo
et al., 2005; Devine et al., 2005; Moussian et al., 2006; Tonning et al., 2005; Zhang et al., 2005).
Several studies have shown the role of ZP proteins in maintaining the chitin matrix in the aECM.
During *Drosophila* tracheal development, two ZP proteins, Dumpy (Dpy) and Piopio (Pio), build
an elastic structure in the lumen that modulates mechanical stress and maintains the chitin matrix

(Jazwinska et al., 2003; Dong et al., 2014; Drees et al., 2023). In the *Drosophila* corneal lens,

another ZP domain protein, Dusky-like, establishes an external scaffold for chitin retention

(Ghosh & Treisman, 2024). However, many organs in both vertebrates and invertebrates have a

non-chitinous aECM, and the role of ZP proteins in the non-chitinous aECM is less well

understood.

Post-translational modifications of the aECM are essential for proper tube dimensions

during organ formation (Luschnig et al., 2006; Wang et al., 2006; Abrams et al., 2006; Herman

& Horvitz, 1999; Hwang et al., 2003). Sulfation is an abundant post-translational modification

that takes place in the Golgi apparatus, where a sulfate group is covalently attached to amino

acid residues of proteins or side chains of proteoglycans (Gunal et al., 2019). Several studies in

C. elegans have shown key roles of sulfation in neuronal organization, development of

reproductive tissues, and phynotypic plasticitiy, linking this post-translational modification to

multiple cellular processes (Turnbull et al., 2003; Bulow & Hobert, 2004; Kinnunen et al., 2005;

Tecle et al., 2013; Diaz-Balzac et al., 2014; Dierker et al., 2016). A study in a mouse model of

cystic fibrosis also suggest a link between the disease state and proper sulfation levels in organs

(Xia et al., 2005), but the role of sulfation in tubular organ development and aECM organization

remains elusive.

Sulfation reactions require a high-energy form of sulfate, 3'-phosphoadenosine-5'-

phosphosulfate (PAPS), which is produced by the enzyme PAPS synthetase (Papss) (van den

Boom et al., 2012; Gunal et al., 2019). Mammals have two Papss isoforms, Papss1 and Papss2.

Papss1 shows ubiquitous expression, with higher levels in glandular cells and salivary duct cells,

suggesting that high degrees of sulfation are required in these cell types (Uhlen et al., 2005;

Karlsson et al., 2021). *Papss2* shows a more restricted expression in certain tissues such as

cartilage, and mutations in *Papss2* have been associated with skeletal dysplasia in humans

(Faiyaz ul Haque et al., 1998; Ahmad et al., 1998; Kurima et al., 1998; Ford-Hutchinson et al.,

2005). Mouse models with exon deletions in *Papss1* or *Papss2* show extensive defects, including

lethality (*Papss1*) and abnormal craniofacial and limb morphology (*Papss2*) (Groza et al., 2023),

demonstrating that Papss is vital for developmental processes in mammals. *Drosophila* has a

single *Papss* gene, allowing for a straightforward investigation of its function in developmental

processes without redundancy. By studying the role of *Papss* in *Drosophila* development, we can

gain valuable insights into the role of this conserved enzyme in normal development and disease
in humans.

To understand the role of post-translational modification in aECM organization and to
determine the key composition of the non-chitinous aECM in tubular organs, we use the
Drosophila embryonic SG as a model. This single-layered epithelial tube is formed through
highly stereotyped morphogenetic processes (Chung et al., 2014), making it an excellent system
for elucidating the molecular mechanisms underlying aECM composition and modification. SG
cells are specified on the ventral surface of the embryo at stage 10. Once specified, SG cells do
not undergo cell division or cell death because SG-upregulated transcription factors
downregulate the expression of key cell cycle genes and proapoptotic genes (Myat & Andrew,
2000a; Chandrasekaran & Beckendorf, 2003; Fox et al., 2013; Matthew et al., 2024), and thus
the entire morphogenetic process is driven by cell shape change, cell rearrangement, and
collective cell migration (Myat & Andrew, 2000b; Bradley & Andrew, 2001; Bradley et al.,
2003; Chung & Andrew, 2014; Chung et al., 2017). At stage 11, SG cells invaginate to form a
three-dimensional tubular structure, which then elongates and migrates toward its final position
during stages 12-13. The SG continues to migrate and the lumen expands significantly by apical
secretion during stages 14-16, forming the fully developed organ. Despite limited knowledge of
the composition of the aECM in the SG, previous work has demonstrated the critical role of
ADAMTS-A, a secreted metalloprotease, in releasing cells from the aECM to facilitate SG
migration and normal lumen shape (Ismat et al., 2013). Our previous work has also shown that
cell-aECM adhesion via the apical surface protein Cad99C influences cell rearrangement during
SG elongation (Chung & Andrew, 2014).

Here, we identify Papss as being essential for aECM organization and lumen expansion
of the *Drosophila* embryonic SG. Loss of *Papss* results in a disorganized apical membrane and a
condensed aECM, which is associated with disrupted Golgi structures and intracellular
trafficking. *Papss* mutants also show defects in the basal ECM and occasional SG cell death,
leading to invasion of hemocytes into SG epithelia that engulf dying cells. Expressing the wild-
type (WT) Papss in the SG rescues all phenotypes, but not the enzyme-dead form. We identify
two ZP proteins, Pio and Dpy, as key components of the SG aECM. In the SG lumen, these ZP
domain proteins form a filamentous scaffold that is significantly affected by the loss of *Papss*,

with Pio eventually being lost from the aECM and the Dpy-positive aECM structure condensing

and dissociating from the apical membrane. Mutations in *pio* or *dpy*, or in *Notopleural* (*Np*),

which encodes a matriptase that cleaves Pio to form the luminal pool of Pio, exhibit bulging and

constriction of the SG lumen. Notably, mutations in *pio* result in the loss of luminal Dpy signals

in the SG. These data indicate that the Dpy-containing filamentous structure of the aECM is

crucial for maintaining luminal diameter and that the luminal pool of Pio is essential for

organizing this structure during SG morphogenesis.

Results

Papss is required for SG lumen expansion

To identify key enzymes that affect SG morphology by modifying aECM components, we examined the effects of the loss of ~20 enzymes that are highly expressed in the SG and are likely to function in the secretory pathway (Fox et al., 2013). Deficiency lines deleting each enzyme were immunostained with the apical membrane marker Crumbs (Crb) or Stranded at Second (SAS) to assess SG lumen shape. Among the twelve lines showing defects (Table 1), a deficiency line deleting *PAPS synthetase* (*Papss*) and 18 other genes had a very thin SG lumen with irregular apical protrusions (Figure 1A). Papss catalyzes the synthesis of 3'-phosphoadenosine-5'-phosphosulfate (PAPS), the sulfate donor compound in all sulfotransferase reactions (Venkatachalam, 2003). Consistent with a previous report (Jullien et al., 1997), our fluorescence in situ hybridization confirmed that *Papss* transcripts are highly expressed in the SG throughout morphogenesis (Figure 1B), suggesting a high demand for sulfation in this organ.

To test the role of *Papss* and confirm the phenotype, we used a *Papss* EMS mutant line (*Papss^2*; Zhu et al., 2005). The Papss protein contains two critical protein domains, the ATP sulfurylase domain, which adds sulfate to an AMP group of ATP to form adenosine 5'-phosphosulfate (APS), and the APS kinase domain, which phosphorylates APS to form PAPS (Venkatachalam et al., 1998; Mueller & Shafqat, 2013). Sanger sequencing of this mutant identified a point mutation (G->A) that introduces a premature stop codon (Trp271->stop; amino acid number is for the Papss-PA isoform) located near the N-terminus of the ATP sulfurylase domain, resulting in a truncated protein in which most of the domain is not translated (Figure 1-figure supplement S1A-B). Since both protein domains are essential for the enzyme function of Papss, we conclude that this mutant line is potentially null or a strong hypomorphic allele. *Papss* homozygous mutants and transheterozygotes for *Papss* mutant over a deficiency were embryonic lethal. To test the role of Papss in sulfation in the SG, we checked for the presence of sulfated tyrosine using a sulfotyrosine antibody in WT and *Papss* mutant embryos. In WT, we observed strong sulfotyrosine signals both in the apical membrane and as puncta in the cytoplasm of SG cells (Figure 1-figure supplement S2A). The cytoplasmic puncta colocalized with ManII-GFP signals, a medial Golgi marker (Yang et al., 2021), suggesting that proteins undergoing tyrosine sulfation in the Golgi are detected by the antibody (Figure 1-figure supplement S2C). In *Papss*

mutants, the cytoplasmic punctate signals were significantly reduced (Figure 1-figure

supplement 2B), suggesting that sulfation in the Golgi is decreased. The signals at the apical

membrane appeared similar or slightly reduced compared to WT, suggesting that sulfated

molecules, possibly by maternally provided Papss, were still targeted apically, although it is not

clear whether they were properly targeted to their original destination or mislocalized. Together

with a previous study using Alcian Blue staining showing the loss of sulfated biomolecules in the

Papss mutant SG (Zhu et al., 2005), these data suggest sulfation in the SG is dependent on Papss.

Papss homozygous mutant embryos often showed overall defective embryonic

morphology, such as defects in germ band retraction and head involution (39%; N=80/204;

Figure 1-figure supplement 3A-F). The heterozygous *Papss* mutant embryos produced by

crossing *Papss* mutants with WT flies appeared normal (Figure 1-figure supplement 3A-F). A

smaller percentage of transheterozygous embryos showed similar defects (10%; N=13/136), with

many late-stage embryos appearing normal (Figure 1-figure supplement 3C, E and F). We

conclude that transheterozygous embryos with more severe defects did not survive to late stages,

but we cannot rule out the possibility that a secondary mutation at another gene locus contributes

to the overall embryonic defects. Embryos with these morphological defects often showed

defects in SG invagination, forming a SG close to the surface of the embryo (Figure 1-figure

supplement 3D, G and H). We used the embryos that formed the SG in the normal position for

quantification and further analysis.

Using the adherens junction (AJ) marker E-cadherin (Ecad) signals, we analyzed SG

lumen phenotypes in *Papss* homozygous mutant embryos and transheterozygotes. During stages

14-16, the WT SG lumen expanded consistently throughout its length (Figure 1C). In contrast,

Papss mutant embryos exhibited irregularities in SG lumen expansion. At stage 14, when the

WT SG lumen expansion began, more than 40% of *Papss* mutant embryos had either a thinner

lumen (Figure 1D and I; 18%; N=7/38) or a slightly irregular lumen with bulges (Figure 1D and

I; 24%; N=9/38). At stage 16, about half of *Papss* homozygous or mutant embryos or

transheterozygotes over a deficiency showed an overall thin lumen with an irregular apical

surface (Figure 1D and I') and the other half had an irregularly expanded lumen with thin and

expanded regions along its length (Figure 1D and I'). Quantification of the SG lumen length and

diameter at stage 16 showed that whereas the lumen length was comparable between WT and

Papss mutant SGs (Figure 1J), the diameter was significantly decreased or irregular, with a high

degree of variation in *Papss* mutant SGs (Figure 1K and K'). Quantification of the SG lumen

area at stage 16 revealed that *Papss* mutants SGs exhibiting the thin lumen phenotype had a

smaller lumen size compared to WT. Together, these data suggest a role for Papss in SG lumen

expansion primarily by expanding the lumen diameter (Figure 1L).

To determine whether the SG lumen expansion defects are due to the loss of *Papss*, we

performed a rescue experiment by overexpressing Papss in the SG using the SG-specific driver

sage-Gal4 (Chung et al., 2009) in the *Papss* mutant background. *Papss* produces multiple

isoforms, including Papss-PA/B/C/G (629 aa), Papss-PD (657 aa), and Papss-PE/F/H (630 aa),

which differ only slightly in the N-terminus (Flybase; www.flybase.org). Using three additional

probes that detect different subsets of splice forms, we performed fluorescent in situ

hybridization and found that all probes detected signals in the SG at varying levels (Figure 1-

figure supplement 4A and B). We generated UAS lines for Papss-PD and Papss-PE and found

that overexpression of either isoform in the SG of WT embryos resulted in a normally expanded

SG lumen (data for Papss-PE overexpression are shown in Figure 1F and I'). Given the high

levels of *Papss* mRNA in the WT SG, the Papss protein is likely abundant in the SG, which may

explain why further overexpression of Papss does not lead to any additional effects. Since our

initial analysis showed identical rescue features in the SG for the generated UAS lines, we used

the UAS-Papss-PD and UAS-Papss-PE lines interchangeably for further experiments (hereafter,

when overexpressed, Papss-PD and Papss-PE will be referred to as Papss). When Papss was

overexpressed in the SG of *Papss* mutant embryos, only a small fraction (4%; N=1/26) showed a

thin lumen phenotype along the entire length of the SG (Figure 1G and I'). The majority of

embryos (73%) showed either a normal, fully expanded lumen along the entire length of the SG,

with occasional slight irregularities at the apical membrane (42%; N=11/26), or a irregularly

expanded lumen, where some parts of the SG lumen were expanded and others were thin (31%;

N=8/26) (Figure 1G and I'). SGs with normally expanded lumens had luminal geometry

comparable to that of WT embryos (Figure 1J-L). Interestingly, some of the embryos had an

over-expanded SG lumen, often with a non-uniform diameter (24%; N=6/24; Figure 1G and I').

To test whether the enzyme activity of Papss is required for its role in SG lumen expansion, we

also generated UAS lines with mutations in the critical residues of the ATP sulfurylase (F593P)

and APS kinase (K193A) domains using the Papss-PD isoform (See Materials and Methods for

details; amino acid number is for Papss-PD). Overexpressing a mutant version of Papss

containing both of these mutations in the SG did not rescue the SG phenotypes as WT Papss did,

but it showed the same range of defects as *Papss* mutants (Figure 1H, I' and J-L). These data

suggest that the sulfation activity of Papss is required for SG lumen expansion. Overall, although

a range of phenotypes was present, the thin SG lumen phenotype was almost completely rescued

by SG-specific expression of WT Papss, but not the enzyme-dead form.

Loss of *Papss* disrupts the apical domain and aECM organization

To investigate the cellular basis of the thin SG lumen phenotype in *Papss* mutants, we examined

the localization of key apical membrane proteins and junctional markers. In WT SGs, the apical

polarity protein Crb localized to the subapical region (SAR) between the apical surface (free

apical area) and the AJs, and we observed clear Crb signals along the apical cell boundary of

each SG cell (Figure 2A and B). In contrast, the Crb signals in *Papss* mutants appeared less

prominently localized to the SAR in SG cells (Figure 2C). We measured the Crb signal intensity

in the SAR and the apical-medial region of SG cells. The ratio of the Crb signal intensity at SAR

to apical-medial intensity was significantly lower in *Papss* mutants than in WT (Figure 2D),

indicating disorganization of the apical domain. The AJ marker Ecad and the septate junction

(SJ) marker Discs large (Dlg) showed no apparent disruption in localization compared to WT

(Figure 2-figure supplement 1A-E). These data suggest that loss of *Papss* primarily affects the

apical membrane in the SG, while leaving the junctional regions intact.

Next, we investigated the organization of the aECM, using wheat germ agglutinin (WGA),

which marks several glycan modifications including O-linked β-N-acetylglucosamine (O-

GlcNAc). WGA has been shown to label the apical membrane and luminal structure in the

Drosophila embryonic trachea (Tonning et al., 2005; Chung et al., 2009). Similarly, we observed

relatively weak WGA signals in the apical membrane and the lumen of WT SGs (Figure 2E; n=25

SGs). Notably, the luminal signals showed multiple thin filamentous organizations traversing the

lumen (Figure 2E), suggesting heterogeneity of the luminal structure. In *Papss* mutants, WGA

signals were increased in both the lumen and apical membrane, with the apical membrane signals

appearing more diffuse (Figure 2F; n=28 SGs). Interestingly, the luminal WGA signals in *Papss*

mutants accumulated strongly on one side along the entire length of the lumen, whereas the thin
filamentous WGA signals in WT SGs appeared to show no bias and were mostly centered in the
lumen. In regions with a thin SG lumen, WGA was present in a highly condensed rod-like
structure; in expanded regions, WGA signals showed either a similar condensed rod-like structure
favoring one side of the SG lumen or a slightly condensed filamentous structure often broadly
associated with the apical membrane (Figure 2F). Transmission electron microscopy (TEM)
analysis also revealed highly condensed and more electron-dense luminal structures in the SG of
Papss mutants compared to WT (Figure 2H and I; original, raw TEM images of several WT and
Papss mutant samples are shown in Figure 2-figure supplement 2A and B). SG-specific
overexpression of Papss in the *Papss* mutant background resulted in a partial rescue; the luminal
WGA structures were less condensed than *Papss* mutants, but they were not as thin as those in WT
SGs (Figure 2G). Taken together, these data suggest a role for Papss in organizing the SG aECM
with loss of *Papss* resulting in a tightly packed or condensed aECM that is loosely associated with
the apical membrane.

Loss of *Papss* disrupts Golgi organization and mislocalizes intracellular trafficking
markers

In addition to the apical membrane and luminal signals, we detected WGA signals around the
nucleus, consistent with WGA binding to glycoproteins on the nuclear envelope (Rutherford et al.,
1997), and in several large, punctate cytoplasmic structures in WT SG cells (Figure 3A and E).
The cytoplasmic WGA puncta colocalized with the medial-Golgi marker ManII-GFP, suggesting
that WGA binds glycoproteins in the Golgi (Figure 3J and L). The number and intensity of the
punctate WGA signals were significantly reduced in *Papss* mutants (Figure 3B, F, I and I').
Overexpression of WT Papss, but not the mutant form of Papss, in the SG restored the number of
WGA cytoplasmic puncta (Figure 3C, D, G, H, I and I'). The intensity of the WGA signals was
not restored in either case (Figure 3I').

Interestingly, we observed highly dispersed ManII-GFP signals in *Papss* mutants SGs
(Figure 3K and L'), suggesting that Golgi structures may be disrupted upon *Papss* loss. To examine
the Golgi structures in more detail, we performed TEM analysis in WT and *Papss* mutant SG cells.

In contrast to the mammalian Golgi, the *Drosophila* Golgi lacks the characteristic ribbon-like

structure and is composed of a series of individual cisternae arranged in a random, disordered

fashion (Kondylis & Rabouille, 2009). While the Golgi in WT SG cells showed randomly arranged

cisternae, the Golgi in *Papss* mutants showed abnormal arrangement, either stacked or

compressed, with small electron-dense puncta along these structures (Figure 3M and N),

suggesting a functional defect and possible protein aggregation. Other organelles, such as the

endoplasmic reticulum (ER), appeared relatively normal. SG-specific expression of GFP-KDEL,

GFP fused to the ER retention motif KDEL, or immunostaining for prolyl-4-hydroxylase-alpha

SG1, an ER-resident enzyme (Abrams et al., 2006), showed no significant differences between

WT and *Papss* mutant SGs (Figure 3-figure supplement 1 A-D). TEM analysis also revealed

similar ER structures in WT and *Papss* mutant SG cells (Figure 3-figure supplement 1E and F).

These data suggest that loss of *Papss* disrupts the Golgi, but not the ER.

To determine whether the phenotypes observed in *Papss* mutants are due to defects in

Golgi structure and function, we examined the morphology of the SG in null mutants of two key

Golgi components, GM130 and Grasp65 (Zhou et al., 2014). Both mutants were homozygous-

viable. Although the GM130 null allele (*GM130^{Δ23}*) was extremely weak and difficult to grow, the

embryos we managed to collect appeared normal, and the SG exhibited only mild irregularity

(Figure 3-figure supplement 2A; n=5/5). It is possible that only relatively normal embryos survived

until late stages of embryogenesis. However, *Grasp65* null mutants (*Grasp65^{102}*) exhibited a

highly irregular SG lumen with condensed luminal WGA signals (Figure 3-figure supplement 2B

and C; n=10/10). These data suggest that disrupted Golgi structures in *Papss* mutants may

contribute to the SG defects.

The aberrant Golgi structures observed in *Papss* mutants prompted us to investigate the

impact on intracellular trafficking components. We examined the localization of Rab11, a marker

for recycling endosomes, Rab7, a marker for late endosomes, and Sec15, an exocyst marker. As

previously shown (Chung & Andrew, 2014; Le & Chung, 2021), Rab11 and Sec15 were enriched

along the apical domain of WT SG cells (Figure 3O and Q). In *Papss* mutants SGs, we observed

an increase in apical Rab11 signal intensity, with some signals mislocalized to the basolateral

region (Figure 3P and S). In contrast, Sec15 signals were significantly reduced at the apical

membrane (Figure 3R and T). Unlike Rab11 or Sec15, Rab7 localized in large punctate structures

in the cytoplasm of WT SG cells (Figure 3-figure supplement 3A). In *Papss* mutants, the number

of Rab7 puncta was significantly reduced, although the intensity of Rab7 puncta was not

significantly different from WT (Figure 3-figure supplement 3B-D). TEM analysis revealed a

number of electron-dense vesicles in the apical domain of SG cells in WT (Figure 3U). In contrast,

Papss mutant SGs showed fewer electron-dense vesicles, but instead we observed several larger,

empty vesicle-like structures (Figure 3V and W). Collectively, these data suggest that defective

sulfation affects Golgi structures and multiple routes of intracellular trafficking.

Loss of *Papss* results in abnormal cell death and disruption of the basal ECM

In *Papss* mutant embryos, we often observed one or more acellular regions within the SG

epithelium during late embryogenesis that lacked any markers for nuclei, cell membranes, or

junctions. To test whether these acellular regions were caused by cell death, we stained embryos

with an antibody against *Drosophila* cleaved death caspase-1 (DCP-1; Sarkissian et al., 2014).

Indeed, DCP-1 signals were often found within the acellular spaces (Figure 4A and B),

suggesting that loss of *Papss* leads to abnormal cell death in the SG. TEM analysis also captured

a dying cell in the *Papss* mutant SG supporting the occurrence of cell death (Figure 4C). We

found that *Papss* mutant SGs have a significantly higher number of these acellular spaces

compared to WT SGs (Figure 4D). Quantifying the number of SG cells at stage 16 revealed that

Papss homozygous embryos and transheterozygotes have significantly fewer cells than WT

(Figure 4E). Overexpression of WT Papss, but not the mutant form of Papss, in the *Papss* mutant

background decreased the number of acellular regions within the SG (Figure 4D) and restored

the number of SG cells (Figure 4E). This suggests that loss of sulfation leads to conditions that

hinder SG cell survival. Blocking apoptosis in the *Papss* mutant SG by overexpressing the

baculoviral protein p35, a caspase inhibitor of apoptosis, also restored the SG cell number

(Figure 4E), suggesting that apoptosis is a key mechanism underlying SG cell loss in *Papss*

mutants. Importantly, the SGs in these embryos still showed thin (25%; N=2/8) or irregularly

expanded (75%; N=6/8) lumens, suggesting that SG lumen expansion defects in *Papss* mutants

are independent of abnormal cell death.

Using srpHemo3x-mCh (Gyoergy et al., 2018), we labeled hemocytes, which are

analogous in function to macrophages in vertebrates. Hemocyte signals were often present in the

acellular regions of the SG in *Papss* mutants, suggesting that hemocytes may aid in apoptotic cell

clearance from the SG epithelium (Figure 4F and G). In normal epithelia, the basal ECM acts as

a physical barrier, preventing hemocytes from migrating into the epithelial layer. To examine the

integrity of the basal ECM in *Papss* we used an antibody against Nidogen (Ndg), a glycoprotein

with important linker functions in the basement membrane (Wolfstetter & Holz, 2012). Ndg-

positive basal ECM was detected in a regular, thin layer around WT SGs at stage 16 (Figure 4H).

However, in *Papss* mutants, Ndg signals were irregular and often disconnected, and hemocytes

invaded SG cells through the gaps in the Ndg-positive basal ECM (Figure 4I). Overall, our data

suggest that loss of *Papss* leads to aberrant cell death in the SG and disrupts both the apical and

basal ECM of the SG.

349   The ZP domain proteins Pio and Dpy of the SG aECM are affected by the loss of *Papss*

To better understand how aECM composition and organization are affected by *Papss* loss, we set

out to identify SG aECM components. Since chitin, a major aECM component in many *Drosophila*

tissues, is absent from the SG lumen (Figure 5A), we focused on ZP domain proteins. Of the 18

ZP domain proteins in *Drosophila*, at least six are expressed in the embryonic SG, including *pio*,

dpy, *quasimodo* (*qsm*), *cypher* (*cyr*), *dusky* (*dy*), and *CG17111* (Tomancak et al., 2002; Jazwinska

& Affolter, 2004; Weiszmann et al., 2009). We assessed the localization of Pio and Dpy, key

aECM components critical for *Drosophila* tracheal morphogenesis (Jazwinska et al., 2003; Drees

et al., 2023), and Qsm, which plays a critical role in remodeling Dpy in the aECM during

Drosophila flight muscle development (Chu & Hayashi, 2021).

To test Pio localization in the SG, we used a transgenic line expressing an mCherry-Pio

(mCh-Pio) fusion protein with the N-terminal mCh tag (Drees et al., 2023; Figure 5B), as well as

a Pio antibody generated with the peptide corresponding to the sequence within the ZP domain of

Pio, between the ZP-N and ZP-C domains (Jazwinska et al., 2003; Figure 5B). mCh-Pio signals

were observed in cytoplasmic puncta, at the apical membrane, and in the SG lumen. These signals,

especially the luminal signals, became stronger at stages 15-16 (Figure 5C), suggesting that Pio is

transported apically and secreted into the SG lumen. Pio antibody signals showed distinct localization, predominantly in the SG lumen and at the apical membrane (Figure 5E and F). At stages 11 and 12, the Pio antibody revealed low levels of cytoplasmic punctate signals along with strong luminal signals, but from stage 13, Pio signals were predominantly localized to the SG lumen (Figure 5E and F; Figure 5-figure supplement 1A). Interestingly, compared to the diffuse luminal mCh-Pio signals (Figure 5C), Pio antibody signals in the SG lumen were heterogeneous with filamentous structures (Figure 5E-6F''), resembling luminal WGA signals. Pio antibody signals were absent at all stages in the *pio* null mutant, in which the entire ZP domain is deleted (Drees et al., 2023; Figure 5-figure supplement 1B and C), confirming antibody specificity. The Pio protein is known to be cleaved, at one cleavage site after the ZP domain by the furin protease and at another cleavage site within the ZP domain by the matriptase Notopleural (Np) (Drees et al., 2019; Drees et al., 2023; Figure 5B). The latter cleavage site falls within the peptide sequence used to generate the Pio antibody (Figure 5B). A possible explanation for the different signals between mCh-Pio and Pio antibody signals is that the Pio antibody preferentially recognizes the C-terminal ZP domain fragment after furin and Np cleavage, whereas diffuse, luminal mCh-Pio signals are either the secreted Pio protein cleaved only by furin or the N-terminal fragment after Np cleavage.

Dpy-YFP signals from the Dpy-YFP knock-in line (Lye et al., 2014; YFP insertion into the *dpy* locus near the middle of the protein) showed heterogeneous, filamentous structures in the SG lumen throughout SG morphogenesis (Figure 5-figure supplement 2A) that colocalized with Pio antibody and WGA signals (Figure 5E-F"; Figure 5-figure supplement 2C). It should be noted that because homozygous Dpy-YFP embryos showed a bulging and constricted SG lumen phenotype with highly condensed Dpy-YFP signals throughout the lumen (Figure 5-figure supplement 2B), we used embryos with one copy of the transgene to analyze Dpy localization. Qsm localization was analyzed using the mCh-Qsm knock-in line, in which mCh is inserted into the *qsm* locus a few amino acids downstream of the N-terminal end of the protein (Chu & Hayashi, 2021). Unlike Pio and Dpy, mCh-Qsm signals were not detected in the SG lumen. Instead, they showed a dynamic intracellular localization ranging from cytoplasmic signals with apical membrane enrichment (stages 11-14) and apical vesicular signals (stages 14-15) to prominent signals at the apical membrane and SJs (stage 16) (Figure 5-figure supplement 3A). These data suggest that Qsm is not a component of the SG aECM. However, it is possible that the mCh-Qsm signals only detect

the cleaved N-terminal fragment of Qsm, while the C-terminal fragment may localize to a different

subcellular location. Taken together, our data suggest that Pio and Dpy form filamentous aECM

structures in the SG.

We next examined Pio, mCh-Pio, Dpy-YFP, and mCh-Qsm signals in *Papss* mutant SGs

at stages 14 and 16. At stage 14, Pio antibody and Dpy-YFP signals appeared condensed and

enriched within the SG lumen, with reduced association with the apical membrane compared to

WT (Figure 5G and G'). Interestingly, Pio antibody signals were barely detectable in the SG

lumen at stage 16 (Figure 5H-H'''), likely due to either protein degradation or epitope masking

caused by aECM condensation and protein misfolding. In contrast, Dpy-YFP signals remained in

the lumen at all stages and were condensed and localized to one side of the lumen at stage 16

(Figure 5H-H'''), similar to WGA signals (Figure 2E). mCh-Pio signals were depleted from the

expanded region of the lumen but weak signals colocalized with condensed Dpy-YFP signals

(Figure 5D). We did not observe any significant differences in the localization or intensity of

mCh-Qsm signals in *Papss* mutants (Figure 5-figure supplement 3B and C). These data suggest

that the luminal pool of Pio and Dpy, which form filamentous structure, is significantly affected

by the loss of *Papss*, resulting in the loss of Pio and the condensation and dissociation of the

Dpy-containing aECM structure from the apical membrane.

ZP domain matrix is required for uniform SG lumen diameter

We next investigated the role of Pio and Dpy in SG morphogenesis by analyzing SG lumen

morphology in *pio* and *dpy* mutants. Immunostaining with Ecad revealed that *pio* null mutants

formed relatively normal SGs up to stage 15, but stage 16 SGs showed dramatically abnormal

luminal morphology characterized by multiple constrictions, with a slight expansion of the lumen

between constrictions (Figure 6A, B and F; Figure 6-figure supplement 1B). The constricted

regions were severely narrowed to the point of almost no luminal space, and some luminal regions

also showed a flat protrusion on one side of the SG lumen (Figure 6C). Notably, these abnormal

luminal shapes were associated with a progressive loss of luminal WGA and Dpy-YFP signals

(Figure 6B, D). In *pio* mutants, WGA signals were comparable to WT at stages 13 and 14 (Figure

6-figure supplement 1A-B), but appeared slightly reduced and diffuse in the SG lumen at late stage

14, and this reduction became more pronounced at stage 15 (Figure 6A and B). Dpy-YFP signals were more severely affected in *pio* mutants. Unlike the strong, filamentous luminal signals in the control (Figure 5-figure supplement 1A), Dpy-YFP signals were reduced and diffuse in the SG lumen and were detected in the cytoplasm of *pio* mutant SG cells from stage 12 to stage 14, with a conspicuous absence in the lumen at stage 15 (Figure 6-figure supplement 1C). By stage 16, both WGA and Dpy-YFP signals were completely absent from the lumen in *pio* mutants, whereas the signals were still present at the apical membrane and in the cytoplasm (Figure 6B and D). We conclude that WGA primarily detects Dpy in the SG lumen and that Pio is required for Dpy secretion to the SG lumen.

To test a role for Pio cleavage and localization in shaping the SG lumen, we analyzed embryos mutant for the matriptase Np, which cleaves Pio. In *Np* mutants, Pio and WGA signals were observed in both the apical membrane and lumen up to stage 14, similar to WT embryos (Figure 6E). However, at stage 16, Pio signals were observed closely associated with the apical membrane, with no luminal Pio signals present (Figure 6E), suggesting that Np is required for this luminal pool of Pio. Importantly, *Np* mutants exhibited the same bulging and constricting SG luminal phenotype at stage 16 as *pio* mutants (Figure 6E and F), indicating that the filamentous luminal Pio structure plays a crucial role in maintaining a uniform SG diameter.

dpy mutants (*dpy^{ov1}*; a spontaneous mutant with a transposable element insertion into intron 11) exhibited a more severe bulging and constriction phenotype in the SG lumen compared to *pio* mutants. Until stage 14, SGs in *dpy* mutants often showed a curved shape, but otherwise had a relatively normal morphology and uniform diameter (Figure 6-figure supplement 1D). At stages 15 and 16, SG lumen lost regularity and uniformity, often exhibiting multiple bulges and constrictions. (Figure 6G and G'). In constricting regions, we observed condensed Pio and WGA signals in the lumen, whereas in bulging regions, weak Pio and WGA signals were detected along the apical boundaries (Figure 6G and G'). These data suggest a disrupted luminal scaffold that correlates with the bulging and constricting luminal morphology.

To further investigate the role of Np in cleaving Pio to organize the aECM during SG morphogenesis, we overexpressed WT (Np^{WT}) and mutant (Np^{mut}) forms of Np using the UAS-Np-Strep and UAS-Np.S990A-GFP lines (Drees et al., 2019). It should be noted that the endogenous Np remains when either form of Np is overexpressed in the SG. Overexpression of

Np^{WT} increased the intensity of luminal mCh-Pio signals (Figure 6H), whereas overexpression of

Np^{mut} resulted in the absence of the mCh-Pio signals in the lumen but strong mCh-Pio signals close

to the apical membrane and in the cytoplasmic puncta (Figure 6I), supporting the role of Np in

cleaving Pio to generate the luminal pool of Pio. In Np^{WT}-overexpressing SGs, the overall intensity

of Pio and Dpy-YFP signals also appeared increased at both the apical membrane and in the lumen

compared to WT (Figure 6H). These embryos did not show obvious changes in the SG

morphology. In Np^{mut}-overexpressing SGs, Pio signals appeared to be increased at the apical

membrane, and various levels of filamentous luminal signals were detected (Figure 6I-I''').

Importantly, the regions of the lumen lacking luminal Pio signals was always slightly expanded

compared to the region with filamentous luminal Pio signals, supporting the idea that the luminal

Pio pool is important for a uniform lumen diameter (Figure 6I' and I''; n=11/13). Dpy-YFP signals

could not be analyzed properly with Np^{mut} overexpression because of the GFP tag in the Np^{mut}

transgene. However, we detected two distinct pools of fluorescent signals in the GFP channel. The

first pool was cytoplasmic punctate signals that colocalized with mCh-Pio, which we confirmed

are Np^{mut}-GFP signals in a separate staining, and the second pool was thin filamentous luminal

signals that overlapped with Pio antibody signals, which we suspect are Dpy-YFP signals (Figure

6I). These data suggest that overexpressed Np^{mut} failed to cleave mCh-Pio and Pio, leading to their

accumulation at and near the apical membrane, but the endogenous Np still cleaved some pool,

resulting in low levels of filamentous luminal Pio structures. Taken together, our data suggest that

Dpy and Pio form a luminal scaffold necessary for a uniform SG tube diameter and that Pio

cleavage by Np is essential for forming this scaffold.

Discussion

Our study highlights the critical role of Papss, a key enzyme that produces the universal sulfate donor PAPS, in aECM organization and lumen expansion in the *Drosophila* embryonic SG. *Papss* mutants show defective SG lumen expansion, accompanied by disruption of the apical domain of SG cells and the aECM. The loss of *Papss* leads to defects in Golgi and intracellular trafficking, abnormal cell death in the SG, and defective basal ECM, emphasizing the importance of sulfation in multiple developmental processes. We also identify the ZP domain proteins Dpy and Pio as key components of the SG aECM, where Dpy is a scaffolding aECM structure that provides mechanical support and Pio is required for proper organization of this structure.

Sulfation is critical for aECM organization and SG lumen expansion

During *Drosophila* SG morphogenesis, secreted cargos are modified in the Golgi, secreted apically, and deposited in the lumen to form the aECM meshwork. Our data show that sulfation is essential for the proper organization of the aECM and the formation of a normal SG luminal shape. Consistent with a high demand for sulfation in the SG, transcripts for *Papss* and several other key enzymes involved in sulfation reactions are also upregulated in the SG, including *slalom* (*sll*), which encodes the *Drosophila* PAPS transporter (Kamiyama et al., 2003; Luders et al., 2003), and *pipe* (*pip*) and *tyrosylprotein sulfotransferase* (*Tpst*), which encode sulfotransferases. Notably, Alcian Blue-stained materials are absent from the SG in *Papss*, *sll*, or *pip* mutants (Zhu et al., 2005), suggesting that sulfation in the SG depends entirely on these enzymes. Our data show that the irregularly expanded SG lumen phenotype in *Papss* mutants is associated with the condensed aECM structures and reduced association of the aECM with the apical membrane (Figures 1 and 2). Inappropriate charge balance due to defective sulfation of proteoglycans and glycoproteins, as well as protein misfolding or failure to bind to appropriate partners, may contribute to aggregation of aECM components and condensation of the aECM structure. The ubiquitously expressed human *Papss1* gene shows upregulation in several glandular organs and the salivary duct (Uhlen et al., 2005; Karlsson et al., 2021), suggesting a potentially conserved role for this enzyme in modifying secreted molecules and modifying aECM in these organs.

The defects in SG lumen expansion in *Papss* mutants are linked to defects in intracellular trafficking, characterized by aberrant Golgi structures and mislocalized or reduced levels of several vesicle markers (Figure 3). This suggests that the loss of *Papss* has a broader impact on the intracellular trafficking system, beyond the sulfation of specific substrate molecules. Mutants of the key Golgi component Grasp65 show a mildly irregular SG lumen (Figure 3-figure supplement 3), supporting the important role of the Golgi in establishing proper lumen dimensions during organ formation. Consistent with the idea that the secretory pathway plays a key role in trafficking aECM components to the lumen, several mutants in intracellular trafficking genes, such as *CrebA*, *Sec61β*, and *Sar1*, which encode a major transcription factor that boosts secretory capacity, a Sec61 translocon component, and a COPII vesicle formation component, respectively, have been shown to have a slightly thinner SG lumen (Fox et al., 2013; Fox & Andrew, 2015). However, the defects in *Papss* mutants differ from those in general secretion mutants. While the latter exhibit a uniformly thin lumen, altered localization of several subcellular organelles including mitochondria and the ER, and reduced Golgi signals (Fox & Andrew, 2015), *Papss* mutants show an irregularly expanded lumen and primarily affect the Golgi, while other organelles remain largely intact (Figure 3 and Figure 3-figure supplement 1). Some of the abnormally modified targets appear to be aggregated in the defective Golgi in *Papss* mutants, but many are still transported apically and secreted into the lumen, resulting in a highly condensed luminal aggregate.

The role of ZP domain proteins in non-chitinous aECM in the SG

We are beginning to understand the structural components and biological functions of the non-chitinous aECM during tubular organ formation. For example, in the non-chitinous aECM of the *C. elegans* vulva and excretory duct, distinct aECM components, including zona pellucida (ZP) domain proteins, chondroitin proteoglycans, and glycoproteins, work together to form the lumens of these tubular organs (Gill et al., 2016; Cohen, Sparacio, et al., 2020). Our work reveals that two ZP proteins, Dpy and Pio, play essential roles in organizing the aECM in the *Drosophila* SG, highlighting the important role of ZP proteins in forming the non-chitinous luminal matrix during tubular organ formation. Dpy, a gigantic protein (>2 MDa) with hundreds of epidermal growth factor (EGF) repeats interspersed with a Dpy module, forms long filamentous structures

(Wilkin et al., 2000). With these filamentous structures, Dpy plays an essential role in the

organization and attachment of apical structures in several tissues, including the thoracic cuticle

(Wilkin et al., 2000), embryonic trachea (Jazwinska et al., 2003), pupal wings (Bokel et al.,

2005), genitalia (Smith et al., 2020), and muscle tendon cells (Chu & Hayashi, 2021). In the SG,

aberrantly formed or lost luminal Dpy structures correlate with abnormal luminal morphology

(Figures 5, 6 and Figure 5-figure supplement 2). We propose that the Dpy filamentous structure

provides mechanical support in the SG aECM and that proper organization of these structures is

critical for the formation of a uniform tube diameter. WGA signals always colocalize with Dpy-

YFP signals in WT and *Papss* mutants, and both signals are absent from the lumen in *pio*

mutants. WGA may primarily detect Dpy in the SG lumen by binding sugar residues that are

attached to numerous EGF modules of the Dpy protein. A caveat to the analysis of Dpy using the

Dpy-YFP line is that it is a non-functional fusion protein. An antibody against Dpy or a new

functional fusion protein will be required to confirm Dpy localization. Even with this caveat, our

data with the Pio antibody suggest that Dpy forms the filamentous luminal structure with Pio. Pio

is required to secrete and maintain Dpy in the lumen of the SG (Figure 6). This is consistent with

a role for Pio in Dpy secretion in the *Drosophila* trachea (Drees et al., 2023) and suggests a

conserved role for Pio in multiple tissues. Since the ZP domains of Pio and Dpy have been

shown to bind directly to each other in vitro (Drees et al., 2023), it is likely that the binding of

Pio and Dpy is required for the formation of filamentous luminal structures. This binding may

require cleavage of Pio by the matriptase Np, which generates the filamentous luminal pool of

Pio (C-terminal ZP fragment of Pio). *pio* and *Np* mutants show similar bulging and constricting

SG lumen morphology (Figure 6), suggesting that the generation of the C-terminal ZP fragment

of Pio by Np is critical for the formation of the filamentous luminal structure with Dpy.

Papss plays an essential role in the formation of the filamentous Dpy/Pio structure

(Figure 5). It will be important to learn if the defects of the Pio-Dpy matrix in *Papss* mutants is a

direct consequence of the loss of sulfation on these proteins or if it's an indirect outcome. The ZP

domain can mediate homomeric or heteromeric polymerization into filaments, and its structure is

stabilized by disulfide bonds formed between conserved Cys residues responsible for the

polymerization of ZP modules (Jovine et al., 2002). Defective disulfide bond formation due to

defective sulfation may destabilize Pio-Dpy interactions. Additionally, hundreds of EGF

modules in the Dpy protein are highly susceptible to O-glycosylation. Sulfation of the glycan

groups attached to Dpy may be critical for its ability to form a filamentous structure. In the

absence of sulfation, the glycan groups on Dpy may not properly interact with the surrounding

materials in the lumen, resulting in an aggregated and condensed structure.

It will be interesting to determine the role of other ZP proteins highly expressed in the SG.

Qsm, for example, has been found in the same intracellular vesicles as Dpy in *Drosophila* tendon

cells and has been shown to physically interact with Dpy through their ZP domains (Chu &

Hayashi, 2021). Qsm has been shown to remodel Dpy filaments that link tendon cells to the pupal

cuticle, increasing their tensile strength (Chu & Hayashi, 2021). In the SG, we find that Qsm is

mainly localized to cytoplasmic vesicles and AJ/SJ regions rather than being present in the SG

lumen like Pio and Dpy. It should be noted that like Pio, Qsm has a furin cleavage site near the

transmembrane domain and another putative protease cleavage site in the ZP domain. Since the

mCh-Qsm line contains the mCh insertion near the N-terminal end of the protein, the mCh-Qsm

signals are likely those of the cleaved N-terminal fragment after cleavage by these proteases and

will need to be confirmed using western blot. Further investigation will be needed to determine

the role of Qsm in junctional regions of SG cells. Also, the roles of other SG-upregulated ZP

domain proteins remain to be elucidated.

In conclusion, our analysis of the *Drosophila Papss* gene, a single ortholog of human

Papss1 and Papss2, reveals its multiple roles during SG development. We expect that these

findings will provide important insights into the function of these enzymes in normal

development and disease in humans. Our findings on the key role of two ZP proteins, Pio and

Dpy, as major components of the SG aECM also provide valuable information on the

organization of the non-chitinous aECM during organ formation.

Materials and Methods

Fly stocks and genetics

The fly stocks used in this study are listed in Table S2. All the crosses were performed at 25°C.

Generation of transgenic lines

UAS lines for the Papss-PD (657 aa) and Papss-PE/F/H (630 aa; hereafter referred to as Papss-

PE) isoforms were generated. Papss-PD and Papss-PE differ only slightly in the N-terminus.

Two UAS clones, UFO09742 (DGRC Stock 1643418;

https://dgrc.bio.indiana.edu//stock/1643418 ; RRID:DGRC_1643418) and UFO09743 (DGRC

Stock 1643419 ; https://dgrc.bio.indiana.edu//stock/1643419 ; RRID:DGRC_1643419), were

obtained from the *Drosophila* Genomics Resource Center. UFO09742 was generated from the

RE15281 cDNA clone encoding the open reading frame (ORF) for the Papss-PD isoform.

UFO09743 was generated using the RE03925 cDNA clone, which encodes the same ORF for the

Papss-PE isoform. Each clone contains a C-terminal Flag-HA tag. Purified DNA for the UAS

clones (500 ng/μl) was sent to GenetiVision (Houston, TX) for injection. GenetiVision

performed PhiC31-mediated insertions using the attP40 site.

UAS-Papss mutant lines for the APS kinase and ATP sulfurylase domains were generated

by site-directed mutagenesis using the UAS-Papss-PD construct (UAS-Papss-PD^{K193A}, UAS-

Papss-PD^{F593P}, and UAS-Papss-PD$^{K193A, F593P}$). K161 and F550 in human Papss (K193 and F593

residues of *Drosophila* Papss-PD*)* were identified as being critical for the catalytic efficiency of

the APS kinase and ATP sulfurylase domains, respectively (Zhang et al., 2022; Zhang et al.,

2023), and were mutated to alanine and proline, respectively. The UAS-Papss-PD construct was

amplified using the NEB Q5 master mix and the following primers.

Primers used to make *UAS-Papss-PD^{K193A}*

Primer forward: 5'-CCGGGACGTGGCGGGTCTATACAAG-3'

Primer reverse: 5'-GTTTCGCACACATCCAG-3'

Primers used to make *UAS-Papss-PD^F593P*

Primer forward: 5'-GATCCTGCCCCCGCGTGTTGCTG-3'

Primer reverse: 5'-TCCATGCTGTCTAGTC-3'

Following amplification, a kinase, ligase, and DpnI (KLD) reaction was performed and the

mutant construct was transformed into DH5α competent cells. To screen for colonies containing

a correct mutation, plasmid DNAs were isolated and the ORF of *Papss* was sequenced using the

following primers.

Primers for sequencing the *Papss* ORF

Forward primer 1: 5'-TGACCGAGCAAAAGCACCAT-3'

Forward primer 2: 5'-CGAGGTGGCCAAGTTGTTTG-3'

Forward primer 3: 5'-CCCGAGTTCTACTTTCAGCGT-3'

Forward primer 4: 5'-GCCGGTGCGAACTTCTACAT-3'

Reverse primer: 5'-ATACCAGGTACGCCTCCAGT-3'

To ensure no secondary mutations are present, the Papss ORF containing the correct mutations

was ligated with the original backbone of the UAS-Papss construct. The construct containing

both the K193A and F593P mutations was generated by combining DNA fragments containing

each mutation, which were obtained by digesting the respective K193A and F593P constructs,

and ligating them to a UAS backbone. Midi-prepped DNA samples were sent to GenetiVision

(Houston, TX) for a PhiC31 injection.

Immunostaining and confocal microscopy

Drosophila embryos were collected on a grape juice agar plate. All embryos were dechorionated

by incubation in 50% bleach for three minutes at room temperature. After dechorionation,

embryos were fixed in either saturated heptane or standard formaldehyde fixative (7 ml heptane,

5.6 ml ddH$_2$O, 700 μl 37% formaldehyde, 700 μl 10x PBS). Saturated heptane was prepared by

mixing heptane:37% formaldehyde 1:1 and shaking vigorously. For immunofluorescence
staining, embryos were fixed in saturated heptane for forty minutes at room temperature,
devitellinized by vortexing in 80% ethanol, and then washed three times in absolute ethanol. For
staining using the horseradish peroxidase (HRP) reaction for visualization, embryos were fixed
in formaldehyde fixative and devitellinized by vortexing in 100% methanol. Embryos were then
stained with primary and secondary antibodies in PBSTB (1X PBS, 0.2% bovine serum albumin,
and 0.1% Triton X-100). All antibodies used are listed in Table S3. Embryos were then mounted
in Aqua-Poly/Mount (Polysciences, Inc). Confocal images were taken using a Leica TCS SP8
confocal microscope with 63x, NA 1.4 and 40x, NA 1.3 objectives. HRP images were taken
using a Leica DM2500 microscope using a 20x, 0.8 NA objective.

Fluorescence in-situ hybridization

In situ hybridization in WT embryos was performed as described in Knirr et al., 1999. Embryos were
fixed in a standard formaldehyde fixation, followed by devitellinzation by 100% methanol. To make the
Papss mRNA probe that detects all splice forms, a 799 bp portion of the *Papss* ORF was amplified
using the *UAS-Papss-PD* construct as a template. To make isoform-specific mRNA probes, unique
sequence segments common to specific *Papss* isoforms were identified using Flybase sequences and
used to design primers for PCR amplification, using WT genomic DNA as a template. The primers are
listed below.

Papss isoform	Primer sequence	Product size
All isoforms	Forward Primer: 5'- ACAAGGATGCGGGTCTGAAG-3' Reverse Primer: 5'- GATTTGGTGACACTATAGCGCCTTAACTC GTTGGGAG-3'	799 bp
ABCFG	Forward primer: 5'-	106 bp

	TTCGCAGTTTTGTTTGGAAT-3' Reverse primer: 5'-<u>GATTTAGGTGACACTATAG</u>TGCTCTCCGG AACTTCACTCAT-3'	
EFH	Forward primer: 5'-TTCGCTTTCGACGTTCGTCG-3' Reverse primer: 5'-<u>GATTTAGGTGACACTATAG</u>CTTCTTTTGA TATGGGGATAAAA-3'	228 bp
DH	Forward primer: 5'-TTCACCTGCCCAACGTTTCG-3' Reverse primer: 5'-<u>GATTTAGGTGACACTATAG</u>CAAGCGCTGT TGTTATTGTTAA-3'	106 bp

*SP6 RNA polymerase promoter sequence underlined

Digoxigenin (DIG)-labeled RNA probe was generated by transcribing amplified and purified

DNA using SP6 RNA polymerase and DIG RNA labeling mix (Sigma). Following hybridization,

the RNA probe was detected using a sheep α-DIG antibody (Enzo; ENZ-ABS302-0001;

1:1,000). The secondary antibody, α-sheep IgG conjugated with biotin (Invitrogen; 1:500), was

visualized using a tyramide reaction with Cy3 (Perkin Elmer Life Sciences, NEL753001kt;

1:50). The embryos were mounted in Aqua-Poly/Mount. Confocal images were taken using a

Leica TCS SP8 confocal microscope with 63x, NA 1.4 and 40x, NA 1.3 objectives.

Genomic DNA sequencing of the Papss open reading frame region in *Papss*[2] mutant

Papss[2] homozygous mutant embryos were sorted for GFP signals in the balancer chromosome, and

genomic DNA was extracted from them. The open reading frame of *Papss* was amplified using the

isolated genomic DNA as a template, with the following primers.

675 Primer forward *Papss*: 5'-CTACAGGTGGCGACGAATGT-3'

676 Primer reverse *Papss*: 5'-GCGGCAGGTTCTGGTAGTAG-3'

677 Sanger sequencing of the amplified DNA segment of *Papss* was performed using the primers

678 below.

679 Forward primer 1: 5'-ACAGGTGGCGACGAATGTG-3'

680 Forward primer 2: 5'-TGACCGAGCAAAAGCACCAT-3'

681 Forward primer 3: 5'-CGAGGTGGCCAAGTTGTTTG-3'

682 Forward primer 4: 5'-CCCGAGTTCTACTTTCAGCGT-3'

683 Forward primer 5: 5'-GCCGGTGCGAACTTCTACAT-3'

684 Reverse primer: 5'-GCGGCAGGTTCTGGTAGTAG-3'

Quantification of SG cell number

687 Confocal images immunolabeled with CrebA for stage 16 SGs were used. 3D images for SGs were

688 generated using Imaris (Andor), and SG nuclei were counted using cell ID numbers. Data were

689 displayed using GraphPad Prism, and statistical significance was determined by one-way ANOVA.

Quantification of SG lumen area

692 Confocal images immunolabeled with Ecad for stage 16 SGs were used. Z-stacks were merged to

693 encompass the entirety of the SG lumen. Using Fiji (NIH), images were set to scale according to image

694 properties listed in LasX files. The polygon selection tool was used to outline the entirety of the SG

695 lumen, excluding the SG duct. SG lumen areas were determined for WT, *Papss* thin, and *Papss*

696 irregularly expanded respectively. Statistical significance was determined using a nonparametric t-test as

697 the data did not fit a Gaussian distribution. Sample numbers are: n=16, WT; n=13, *Papss^2* with a thin

698 lumen; n=11, *Papss^2* with an irregularly expanded lumen; n=8, *Papss^2/Df* with a thin lumen; n=8,

699 *Papss^2/Df* with a irregularly expanded lumen; n=10, *sage>Papss* with a normally expanded lumen; n=8,

700 *sage>Papss-PE; Papss^2* with a normally expanded lumen; n=10, *sage>Papss-PD; Papss^2* with a

701 normally expanded lumen; n=11, *sage>Papss-PD^{K193A,F593P}; Papss^2* with a thin lumen; n=7,

702 *sage>Papss-PD^{K193A,F593P}; Papss^2* with a irregularly expanded lumen.

Quantification of SG lumen length and diameter

Confocal images of stage 16 SGs immunolabeled with an antibody for the cell junction marker, E-cad,

were used. Several Z-sections, spanning approximately half of the circumferential/radial axis of the SG

lumen, were merged to create a projection which covered the entire length of the lumen. The line tool in

Fiji was used to measure the length of the SG lumen. To measure the diameter of the SG lumen, the

lumen was divided into four equal segments using the lumen length obtained above, as there are

sometimes minor variations in width value along the length of the lumen. The line tool in Fiji was then

used to measure the width along each quarter of the lumen, and an average value was obtained per SG.

Each measurement obtained in Fiji was adjusted for scale according to the image properties in the raw

confocal file. Sample numbers are: n=8, WT; n=8, *Papss^2* with a thin lumen; n=10, *Papss^2* with an

irregularly expanded lumen; n=8, *Papss^2/Df* with a thin lumen; n=8, *Papss^2/Df* with a irregularly

expanded lumen; n=10, *sage>Papss-PE* with a normally expanded lumen; n=8, *sage>Papss-PE; Papss^2*

with a normally expanded lumen; n=10 *sage>Papss-PD; Papss^2* with a normally expanded lumen;

n=11, *sage>Papss-PD^{K193A,F593P}; Papss^2* with a thin lumen; n=7, *sage>Papss-PD^{K193A,F593P}; Papss^2* with

a irregularly expanded lumen.

Quantification of Crb signal intensity

Confocal images of stage 16 SGs immunolabeled with antibodies against the sub-apical region (SAR)

marker, Crb, and the AJ marker, Ecad, were used. A maximum intensity projection of the apical region

of several SG cells with the highest Ecad and Crb signals was created using seven z-sections. Regions

were manually drawn along the inner or outer boundary of the Ecad signals of each cell to mark the

SAR and apical-medial regions. The mean gray value of the apical-medial Crb signals in each cell (the

region delineated by the inner boundary of the Ecad signals) was quantified using Fiji. The mean gray

value of the SAR Crb signals was calculated using the integrated densities and areas calculated with Fiji

for regions between inner and outer boundaries. To subtract the background, the average mean gray

value of Crb signals for five regions outside the embryo was subtracted from the mean gray values of the

SAR and apical-medial Crb signals in each SG cell. The ratio of the SAR to the apical-medial Crb level

was compared. Seven and six embryos were used for the WT and *Papss* mutants, respectively, and four

to eight cells per SG. SuperPlots were used to address variability in the data sets of each SG. Statistical

significance was determined using a Welch's t-test.

Quantification of the number and signal intensity of WGA puncta

High-resolution confocal images of stage 16 SGs immunostained with CrebA (SG nuclei), Ecad, and

WGA were obtained using an Olympus SpinSR10 Sora spinning disk confocal microscope with a 100x,

NA 1.5 objective. Each image was taken under identical imaging conditions. Three-dimensional (3D)

reconstruction of the WGA puncta was obtained using the Imaris software (Andor), and the number of

puncta per SG was counted by Imaris. The Imaris machine learning tool was used to exclude non-puncta

WGA signals within the SG and non-SG WGA puncta. After 3D reconstruction, the pixel intensity of

each WGA punctum was also obtained and used for further analysis.

Quantification of WGA and ManII-GFP signal intensity

Confocal images of stage 16 SGs immunostained for WGA and ManII-GFP were used. Single z-sections

of each image were imported into Fiji, and a line intensity profile was generated using the Line Tool and

Color Profiler pluggin. Lines were drawn across regions spanning two ManII-GFP/WGA puncta, and

the signal intensities for each pixel along the line for both ManII-GFP and WGA channels were

obtained.

Quantification for Rab11 and Sec15 apical region intensity

Single z-sections halfway through the proximal-distal axis from confocal images for stage 16 SGs

immunolabeled for Rab11 or Sec15 and Ecad were used for quantification. Threshold images for Sec15

or Rab11, which display pixels as black or white based on whether they are above a value for signal

intensity, were generated in Adobe Photoshop. In addition, Sec15 and Rab11 images were converted

using the RGB stack function in Fiji. A stack of images including the threshold and RGB stack

converted images for Sec15 or Rab11 and the Ecad images was generated using the images to stack

function in Fiji. Using the threshold images for Sec15 and Rab11, a boundary for the apical region

encompassing five SG cells, and the mean gray value intensity was calculated from the RGB stack

converted Rab11or Sec15 images. The graph was generated in GraphPad Prism, and statistical

significance was calculated using Welch's t-test.

Quantification for Rab7 puncta number and intensity

Single z-sections halfway through the proximal-distal axis from confocal images for stage 16 SGs

immunolabeled for Rab7 and Ecad were used for quantification. Threshold images for Rab7, which

display pixels as black or white based on whether they are above a value for signal intensity, were

generated in photoshop. Rab7 images were converted using the RGB stack function in Fiji. A stack of

images including the threshold and RGB stack converted images for Rab7 and Ecad images were

generated using the images to stack function in Fiji. Using the threshold images for Rab7, a boundary

was drawn around all puncta present in five cells for each sample, and the mean gray value intensity and

area was calculated for the RGB stack Rab7 converted image. The average mean gray value of punctate

intensity and the average number of punta in each cell were calculated using Fiji. Punta with an area

greater than or equal to 0.100 μm^2 were used for quantification. Statistical significance was calculated

by Welch's t-test using GraphPad Prism.

Quantification of the number of dying cells

777   Confocal images of stage 16 SGs that were immunolabeled with antibodies against CrebA (to mark the

778   SG nucleus), and DCP-1 (a cell death marker) were used. The number of regions within the SG with

779   DCP-1 signals and/or blank spots where CrebA signals should have been present were counted

780   manually.

Transmission electron microscopy (TEM)

Stage 15 embryos were fixed in 2% glutaraldehyde in 0.2 M phosphate buffer pH 7.4 and rinsed

4 times in 0.1 M phosphate buffer pH 7.2 containing 0.02 M glycine. Embryos were post-fixed in

2% osmium tetroxide for 1 hour in the dark, rinsed 3 times in dH$_2$O, stained en bloc in 1% uranyl

acetate for 1 hour in the dark, and rinsed 3 times in dH$_2$O. Samples were dehydrated in a series

of ethanol and then propylene oxide twice. Embryos were then infiltrated in propylene oxide:

Epon resin series and embedded in Epon at 60ºC for 24 hours. Semi-thin sections were

performed followed by Toluidine Blue staining to obtain a salivary gland section. Ultra-thin

sections for TEM were cut on a Leica EM UC7 microtome. TEM sections were mounted on

carbon-coated copper grids (EMS FCF-150-CU), stained with 2% uranyl acetate and Reynolds

lead citrate, and the ultra-thin sections were viewed on a JEOL JEM-1400, 120kV transmission

electron microscope and imaged with a Gatan digital camera. Sections from at least five

independent embryos were analyzed for wild type and *Papss* mutants.

 Figure legends

Figure 1. *Papss* mutants show defects in SG lumen expansion. (A) Stage 16 SGs
immunostained for Stranded at Second (SAS). (B) In situ hybridization using an antisense probe
for *Papss* mRNA in WT embryos. (C-H) Stage 14 and stage 16 SGs immunostained for Ecad
and CrebA. Yellow asterisks indicate regions of irregular, non-uniform expansion (D, E, H) or
over-expansion (G) of the SG lumen. (I, I') Quantification of different SG lumen phenotypes. (J)
Quantification of SG lumen length. n=7-11 SGs for each genotype. See the Materials and
Methods for the sample size of each genotype. One-way ANOVA test with multiple comparison
(*, p<0.05; ns, non-significant). (K, K') Quantification of the SG lumen diameter (K) and the
coefficient of variation for each SG (K'). The diameter is measured in three different regions of
the lumen. The average diameter, as well as the individual data points, are shown. The same
samples are used for the quantifications in J-K'. One-way ANOVA test with multiple
comparison (***, p<0.001; ns, non-significant) (L) Quantification of SG lumen area. n=7-16
SGs for each genotype. One-way ANOVA test with multiple comparison (***, p<0.001; ****,
p<0.0001; ns, non-significant). Horizontal lines indicate mean values.

Figure 2. Mutations in *Papss* result in mislocalized Crb and disruption of the aECM
architecture. (A) Cartoon showing the subcellular localization of Crb in SG cells. (B, C) Stage
16 SGs immunostained for Crb and Ecad. Magnified images are shown for yellow dotted boxed
regions. (D) Quantification of the Crb signal intensity ratio between the SAR and apical-medial
regions of SG cells. (n=7, WT; n=6, *Papss²* (4-8 cells per SG)). Student's T-test with Welch's
correction (***, p<0.001). (E-G) Stage 16 SGs stained for Ecad and wheat germ agglutinin
(WGA). Magnified images are shown for the boxed regions. Arrows, WGA localization at the
apical membrane in WT (yellow) and *Papss* mutant (cyan) SGs. Yellow arrowhead in E, WGA
localizes as thin filaments in the lumen of WT SG. Cyan arrowheads, WGA localizes as a highly
condensed structure in the thin luminal region of the *Papss* SG. White arrowheads, WGA
localizes as a condensed, filamentous structure in the expanded luminal region of the *Papss* SG.
Yellow arrowheads in G, WGA localizes as a mildly condensed filament in the lumen of
sage>Papss-PE; Papss² SG. (H, I) TEM images of stage 15 SGs. SGs are pseudo-colored in
magenta. Magnified images of the luminal areas (boxed regions) are shown in the bottom.
Yellow asterisk in H, electron-dense luminal material in WT SG. Cyan asterisks in I, condensed

electron-dense luminal material in the *Papss* mutant SG. Green asterisk, a large luminal area

lacking electron-dense material in the *Papss* mutant SG.

Figure 3. Loss of *Papss* results in disorganized Golgi structures and defects in intracellular

trafficking components. (A-D) Stage 16 SGs stained with WGA. Magnified images are shown

for the boxed regions. Yellow arrows, WGA signals as cytoplasmic puncta. Cyan arrows, WGA

signals in the nuclear envelope. (E-H) Three-dimensional (3D) reconstruction of cytoplasmic

WGA puncta in stage 16 SGs. (I, I') Quantification of the number of WGA puncta (I) and signal

intensity (I') within SG cells. n=5 SGs for all genotypes, except WT in G (n=6). One-way

ANOVA test with multiple comparisons (**, $p<0.01$; ***, $p<0.001$; ns, non-significant).

Horizontal lines indicate mean values. (J, K) Stage 16 SGs stained for GFP (for ManII-GFP) and

WGA. Magnified images are shown for the boxed regions. Yellow arrows in J, ManII-GFP and

WGA puncta colocalize in WT SG. Asterisks in K, dispersed ManII-GFP signals in *Papss*

mutants. Cyan arrows in K, little colocalization of ManII-GFP and WGA in the *Papss* SG. (L,

L') Line intensity profile showing the signal intensity of ManII-GFP (green) and WGA

(magenta) along the yellow and cyan dotted lines in J and K, respectively. (M, N) TEM images

of the Golgi. Yellow arrows in M, Golgi structures in WT SG cells. Magenta arrows in N, Golgi

structures in the *Papss* SG. Cyan arrows in N, electron-dense structures in the Golgi of *Papss*

mutant SG cells. (O, P) Stage 16 SGs immunostained for Rab11 and Ecad. Magnified images are

shown for boxed regions. Rab11 signals at the apical membrane are increased in *Papss* mutants

(cyan arrowheads in P) compared to WT (yellow arrowheads in O), with occasional large

punctate signals mislocalized to the basolateral domain (cyan arrow in P). (Q, R) Stage 16 SGs

immunostained for Sec15 and Ecad. Compared to strong apical Sec15 signals in the WT SG

(yellow arrowheads in Q), Sec15 signals are reduced in *Papss* mutants (cyan arrowheads in R).

(S, T) Quantification of the intensity of Rab11 (S) and Sec15 (T) at the apical membrane regions

in WT and *Papss* mutant SGs. Welch's t-test (**, $p<0.05$). Horizontal lines indicate mean values.

(U, V) TEM images for the apical region of SG cells. Yellow arrowheads, electron-dense

vesicles in both genotypes. Red arrowheads, empty, larger vesicle-like structures in *Papss*

mutants. (W) Quantification of the electron-dense vesicle numbers per cell slice. Welch's t-test

(**, $p<0.05$). Horizontal lines indicate mean values.

Figure 4. *Papss* mutants show premature SG cell death. (A, B) Stage 16 SGs immunostained

for CrebA, DCP-1 and Ecad. Cross-sectional images are shown for the yellow dashed line in B.

Yellow arrowheads, DCP-1 signal is detected in the SG in *Papss* mutants. White dashed lines,

the basal boundary of the SG. (C) TEM image of stage 15 SG cells, including a dying cell

(yellow arrow). Cyan arrows, nuclei of normal living cells. (D) Quantification of the number of

acellular regions in stage 16 SGs. (n=8 SGs, WT; n=10, *Papss²;* n=15, *sage>Papss-PD; Papss²*;

n=8, *sage>Papss-PD^{K193A,F593P}; Papss²*) (E) Quantification of the number of SG cells at stage 16

(n=10 SGs, WT; n=10, *Papss²*; n=8, *Papss²/Df*; n=8, *sage>p35; Papss²*; n=10, *sage>Papss-PE;*

Papss²; n=8, *sage>Papss-PD^{K193A,F593P}; Papss²*). One way ANOVA test (****, p<0.0001; ns,

non-significant). (F, G) Stage 16 SGs immunostained for CrebA, Ecad, and mCherry (for

Hemo3x-mCh). Cross-section images are shown for the yellow dashed lines. White dashed lines,

the basal boundary of the SG. Yellow arrowhead in F, hemocyte signals outside of the control

SG. White arrowhead in G, hemocyte signals are present inside the SG in *Papss* mutants. (H, I)

Stage 16 SGs immunostained for DAPI, mCh (for Hemo3x-mCh), and Ndg. Magnified views are

shown for boxed regions. Yellow dashed lines, the basal boundary of the SG. (H) In WT,

hemocytes are found outside of the SG (yellow arrowheads), with Ndg forming a barrier at the

basal boundary of the SG (yellow arrows). (I) In *Papss* mutants, hemocytes invade the SG

epithelia (white arrowheads), where Ndg signals are irregular or absent (white arrows).

Figure 5. Loss of *Papss* disrupts the localization of Dpy and Pio in the aECM. (A) Stage 16

SGs stained for a chitin-binding protein (CBP) and Ecad. (B) Cartoon of the mCh-Pio fusion

protein showing the region for the antibody and protease cleavage sites. (C, D) Stage 14 and

stage 16 SGs immunostained for mCh (for mCh-Pio) and GFP (for Dpy-YFP). The mCh-Pio

(white arrowheads) and Dpy-YFP (cyan arrowheads) signals are largely absent from the

expanded region of the lumen. Along one side of the lumen, weak mCh-Pio signals colocalize

with condensed Dpy-YFP signals (yellow arrows). (E-H''') Stage 14 and stage 16 SGs

immunostained for Pio, GFP, and Ecad. Magnified images of boxed regions in E-H and shown in

E'-H''. F'' and H''' are cross-sectional images of the SG lumen for regions indicated by white

lines F' and H''. Yellow arrowheads, Pio and Dpy localized to the apical membrane. Cyan

arrowheads, Dpy and Pio localized to the SG lumen. Note a higher proportion of Dpy and Pio

localized to the lumen in the *Papss* mutant SG at stage 14 (cyan arrowheads in G'). White

arrowheads in H'-H''', Dpy accumulates towards one side of the lumen in the stage 16 *Papss*

SG. Yellow asterisks, Pio is absent from the lumen at stage 16 in the *Papss* SG.

Figure 6. Pio is essential for uniform SG lumen diameter by maintaining Dpy in the lumen.

(A, B) SGs at stages 13-16 stained for WGA and Ecad. Yellow arrowheads in A, filamentous

luminal WGA signals. Cyan arrowheads in B, WGA signals at the apical membrane. Yellow

asterisks, loss of lumina WGA signals. White arrowheads, constricted lumen. (C) Stage 16 SGs

immunostained for Ecad and CrebA. Cross sectional images for the yellow and green dotted

lines are shown on the right. Yellow arrowhead, flat protrusion of the lumen. Green arrowhead,

constricted lumen. (D) Stage 16 SGs stained for Ecad and GFP (for Dpy-YFP). Projection

images contain merged z-sections, from the apical surface of the SG cells to midway through the

SG. Single section, one z-section at the approximate midpoint of the lumen width. Yellow

arrowhead, area of constricted lumen. Yellow asterisk, luminal area with no Dpy-YFP signals.

(E) Single z-sections of confocal images of SGs at stages 15 and 16 stained for Ecad, Pio and

WGA. White arrowheads, area of constricted lumen. Yellow asterisks, no luminal signals of

WGA and Pio. (F) Quantification of the number of constrictions in the SG lumen at stage 16

(n=7 for both genotypes). (G, G') SGs at stages 14-16 stained for Ecad, WGA and Pio. Higher

magnification of the yellow boxed regions is shown in G'. (H, I) Stage 16 SGs overexpressing

Np^{WT} (H) or Np^{mut} (I) that are immunostained for GFP (for Dpy-YFP and Np-GFP), Pio, and

mCh (for mCh-Pio). Np^{WT}-overexpressing SGs show strong, uniform mCh-Pio signals in the

lumen (cyan asterisks in H), whereas Np^{mut}-overexpressing SGs shows no detectable signals in

the lumen (cyan asterisk in I). In Np^{WT}-overexpressing SGs, the luminal mCh-Pio signals are so

bright that the cytoplasmic, punctate mCh-Pio signals are rarely detectable alongside the luminal

signals. Increasing the brightness to visualize the cytoplasmic punctate signals shows very strong

luminal mCh-Pio signals (inset in H). Np^{WT} overexpression shows high levels of Pio (magenta

arrowheads in H) and dpy-YFP (green arrowheads in H) signals in both the apical membrane and

lumen. In contrast, Np^{mut} overexpression shows strong Pio signals in the apical membrane and

various levels of Pio in the lumen. (I', I'') Higher magnification of the yellow boxed regions.

The lumen diameter of the region with undetectable luminal Pio levels is slightly larger (magenta

asterisk) compared to the region with clear luminal Pio signals (magenta arrowheads).

Figure 7. A proposed model. (A) Cartoon diagram showing normal lumen expansion in WT SG

and defective luminal morphology in *Papss*, *pio*, and *Np* mutants. (B) Organization of the ZP

domain protein-containing aECM in the SG. In WT, the ZP-C fragment of Pio (which is cleaved

by Np and furin proteases) forms a complex with Dpy to build a filamentous scaffold within the

lumen. In *Papss* mutants, this fragment is absent from the lumen at stage 16, and the Dpy-

containing aECM structure becomes aggregated and highly condensed. The absence of luminal

Pio in *pio* and *Np* mutants results in the loss of luminal Dpy and lumen constrictions.

Acknowledgements

We thank the members of the Chung laboratory for their comments and suggestions. We thank

M. Behr, S. Hayashi, the Bloomington stock center (NIH P40OD018537), and the Kyoto

Drosophila stock center at Kyoto Institute of Technology for fly stocks. We thank M. Affolter

for the Pio antibody. Many antibodies were obtained from the Developmental Studies

Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa,

Department of Biology, Iowa City, IA 52242. cDNAs for *Papss* were obtained from the

Drosophila Genomics Resource Center (NIH 2P40OD010949). Confocal and electron

microscopy were performed at the Shared Instrumentation Facility (SIF) at Louisiana State

University. The model in Figure 7 was created in https://BioRender.com. This work was

supported by the National Science Foundation (MCB 2141387) and National Institutes of Health

(R03DE034704) to SC.

Table S1. A targeted Df screen revealed Papss as a key enzyme for SG lumen expansion.

SG-upregulated enzymes	Predicted roles	Df lines tested	SG lumen shape
CG17633	Metallocarboxypeptidase	Exel6024	Disrupted embryo morphology
CG30283	Serine-type endopeptidase	Exel6076	Disrupted embryo morphology
Papss	PAPS synthetase	Bsc445	Thin SG lumen with apical protrusions
PH4αSG1/SG2	Prolyl 4-hydroxylase	Exel6216	Non-uniform lumen diameter
CG16758	Purine nucleotide phosphorylase	Bsc116	Non-uniform lumen diameter
CG8708	N-acetylgalactosaminide β-1,3-galactosyltransferase	Exel7095	Irregular apical membrane
CG7920	Acetate CoA-transferase	Bsc620	Irregular apical membrane
Pgant6	Polypeptide N-acetylgalactosaminyltransferase	Exel6091	Irregular apical membrane
GlcAT-P	Glucuronosyltransferase	Bsc817	Irregular apical membrane
CG34290	Serine-type endopeptidase	Bsc638	Irregular apical membrane
Peroxidasin	Bromide peroxidase	Bsc119	Irregular apical membrane
ScsβG	Succinyl Co-A synthetase β subunit	Bsc387	Short and rounded SG

939 **Table S2. Fly strains used**

Fly strain	Sources and References	RRID
Oregon R (wild type)		
sage-Gal4	Chung et al., 2009	
fkh-Gal4	Henderson and Andrew, 2000	
Papss²	Bloomington Stock Center	BDSC_26171
UAS-p35	Bloomington Stock Center	BDSC_5072
UAS-Papss-PD	This work	
UAS-Papss-PE	This work	
UAS-Papss-PD^{K193A, F593F}	This work	
Dpy-YFP	Kyoto stock center	DGGR_115238
SrpHemo3x-mCherry	Gyoergy et al., 2018	BDSC_78358
Dp-YFP, mCh-Qsm	Chu and Hayashi, 2021	
mCh-Qsm	Chu and Hayashi, 2021	
Dpy-YFP, mCh-Pio	Drees et al., 2023	
UAS-ManII-GFP	Bloomington stock center	BDSC_65248
UAS-GFP-KDEL	Bloomington stock center	BDSC_9898
pio^{17c}	Drees et al., 2023	
Np^{C2}	Drees et al., 2023	
dpy^{ov1}	Bloomington stock center	BDSC_276
Grasp65^{102}	Zhou et al., 2014	BDSC_65257
GM130^{Δ23}	Zhou et al., 2014	BDSC_65255

UAS-Np.WT-Strep	Drees et al., 2019	
UAS-Np.S990A-GFP	Drees et al., 2019	

942 **Table S3. Antibodies Used**

Antibody	Source	RRID	Dilution
α-Ecad (rat)	DSHB (DCAD2)	AB_528120	1:50
α-CrebA (rat)	Andrew lab (John Hopkins University); re-made in this study		1:3,000
α-CrebA (rabbit)	Andrew lab (John Hopkins University)	AB_10805295	1:5,000
α-CrebA (guinea pig)	This study		1:5,000
α-GFP (mouse)	Invitrogen (A11120)	AB_221568	1:500
α-GFP (chicken)	Invitrogen (A10262)	AB_2534023	1:1,000
α-RFP (rabbit)	Invitrogen (R10367)	AB_10563941	1: 2,000
α-β-galactosidase (rabbit)	Invitrogen (A11132)	AB_221539	1:500
α-β-galactosidase (mouse)	Invitrogen (MA5-15222)	AB_10980770	1:500
α-mCherry (rat)	Invitrogen (M11217)	AB_2536611	1:1,000
α-mCherry (rabbit)	Invitrogen (PA534974)	AB_2552323	1:1,000
α-DCP-1 (rabbit)	Cell signaling technology (9578)	AB_2721060	1:200
α-Crb (mouse)	DSHB (Cq4)	AB_528181	1:10
α-Rab11 (rabbit)	Andrew lab (John Hopkins University)		1:500
α-Rab7 (mouse)	DSHB (Rab7)	AB_2722471	1:10

α-Sec15 (guinea pig)	Bellen lab (Baylor College of Medicine)		1:2,000
α-α-Spectrin (mouse)	DSHB (3A9)	AB_528473	1:2
α-Discs large (mouse)	DSHB (4F3)	AB_528203	1:500
α-Pio (rabbit)	Affolter lab (Jaźwińska et al., 2003)		1:500
α-Nidogen (rabbit)	Holz lab (Wolfstetter et al., 2009)		1:2,000
α-Sulfotyrosine (mouse)	Sigma (ZMS1096-25UL)		1:50
Alexa Fluor 488/568/647-coupled secondary antibodies	Invitrogen		1:500

References

Abrams, E. W., Mihoulides, W. K., & Andrew, D. J. (2006). Fork head and Sage maintain a
uniform and patent salivary gland lumen through regulation of two downstream target
genes, PH4alphaSG1 and PH4alphaSG2. *Development*, *133*(18), 3517-3527.
https://doi.org/10.1242/dev.02525

Ahmad, M., Faiyaz Ul Haque, M., Ahmad, W., Abbas, H., Haque, S., Krakow, D., Rimoin, D.
L., Lachman, R. S., & Cohn, D. H. (1998). Distinct, autosomal recessive form of
spondyloepimetaphyseal dysplasia segregating in an inbred Pakistani kindred. *Am J Med*
Genet, *78*(5), 468-473. https://www.ncbi.nlm.nih.gov/pubmed/9714015

Araujo, S. J., Aslam, H., Tear, G., & Casanova, J. (2005). mummy/cystic encodes an enzyme
required for chitin and glycan synthesis, involved in trachea, embryonic cuticle and CNS
development--analysis of its role in Drosophila tracheal morphogenesis. *Dev Biol*,
288(1), 179-193. https://doi.org/10.1016/j.ydbio.2005.09.031

Bokel, C., Prokop, A., & Brown, N. H. (2005). Papillote and Piopio: Drosophila ZP-domain
proteins required for cell adhesion to the apical extracellular matrix and microtubule
organization. *J Cell Sci*, *118*(Pt 3), 633-642. https://doi.org/10.1242/jcs.01619

Bradley, P. L., & Andrew, D. J. (2001). ribbon encodes a novel BTB/POZ protein required for
directed cell migration in Drosophila melanogaster. *Development*, *128*(15), 3001-3015.
https://doi.org/10.1242/dev.128.15.3001

Bradley, P. L., Myat, M. M., Comeaux, C. A., & Andrew, D. J. (2003). Posterior migration of
the salivary gland requires an intact visceral mesoderm and integrin function. *Dev Biol*,
257(2), 249-262. https://doi.org/10.1016/s0012-1606(03)00103-9

Bulow, H. E., & Hobert, O. (2004). Differential sulfations and epimerization define heparan
sulfate specificity in nervous system development. *Neuron*, *41*(5), 723-736.
https://doi.org/10.1016/s0896-6273(04)00084-4

Chandrasekaran, V., & Beckendorf, S. K. (2003). senseless is necessary for the survival of
embryonic salivary glands in Drosophila. *Development*, *130*(19), 4719-4728.
https://doi.org/10.1242/dev.00677

Chu, W. C., & Hayashi, S. (2021). Mechano-chemical enforcement of tendon apical ECM into
nano-filaments during Drosophila flight muscle development. *Curr Biol*, *31*(7), 1366-
1378 e1367. https://doi.org/10.1016/j.cub.2021.01.010

975 Chung, S., & Andrew, D. J. (2014). Cadherin 99C regulates apical expansion and cell
976  rearrangement during epithelial tube elongation. *Development*, *141*(9), 1950-1960.
977  https://doi.org/10.1242/dev.104166

978 Chung, S., Hanlon, C. D., & Andrew, D. J. (2014). Building and specializing epithelial tubular
979  organs: the Drosophila salivary gland as a model system for revealing how epithelial
980  organs are specified, form and specialize. *Wiley Interdiscip Rev Dev Biol*, *3*(4), 281-300.
981  https://doi.org/10.1002/wdev.140

982 Chung, S., Kim, S., & Andrew, D. J. (2017). Uncoupling apical constriction from tissue
983  invagination. *Elife*, *6*. https://doi.org/10.7554/eLife.22235

984 Chung, S., Vining, M. S., Bradley, P. L., Chan, C. C., Wharton, K. A., Jr., & Andrew, D. J.
985  (2009). Serrano (sano) functions with the planar cell polarity genes to control tracheal
986  tube length. *PLoS Genet*, *5*(11), e1000746. https://doi.org/10.1371/journal.pgen.1000746

987 Cohen, J. D., Bermudez, J. G., Good, M. C., & Sundaram, M. V. (2020). A C. elegans Zona
988  Pellucida domain protein functions via its ZPc domain. *PLoS Genet*, *16*(11), e1009188.
989  https://doi.org/10.1371/journal.pgen.1009188

990 Cohen, J. D., Sparacio, A. P., Belfi, A. C., Forman-Rubinsky, R., Hall, D. H., Maul-Newby, H.,
991  Frand, A. R., & Sundaram, M. V. (2020). A multi-layered and dynamic apical
992  extracellular matrix shapes the vulva lumen in Caenorhabditis elegans. *Elife*, *9*.
993  https://doi.org/10.7554/eLife.57874

994 Devine, W. P., Lubarsky, B., Shaw, K., Luschnig, S., Messina, L., & Krasnow, M. A. (2005).
995  Requirement for chitin biosynthesis in epithelial tube morphogenesis. *Proc Natl Acad Sci*
996  *U S A*, *102*(47), 17014-17019. https://doi.org/10.1073/pnas.0506676102

997 Diaz-Balzac, C. A., Lazaro-Pena, M. I., Tecle, E., Gomez, N., & Bulow, H. E. (2014). Complex
998  cooperative functions of heparan sulfate proteoglycans shape nervous system
999  development in Caenorhabditis elegans. *G3 (Bethesda)*, *4*(10), 1859-1870.
1000  https://doi.org/10.1534/g3.114.012591

1001 Dierker, T., Shao, C., Haitina, T., Zaia, J., Hinas, A., & Kjellen, L. (2016). Nematodes join the
1002  family of chondroitin sulfate-synthesizing organisms: Identification of an active
1003  chondroitin sulfotransferase in Caenorhabditis elegans. *Sci Rep*, *6*, 34662.
1004  https://doi.org/10.1038/srep34662

Dong, B., Miao, G., & Hayashi, S. (2014). A fat body-derived apical extracellular matrix enzyme is transported to the tracheal lumen and is required for tube morphogenesis in Drosophila. *Development*, *141*(21), 4104-4109. https://doi.org/10.1242/dev.109975

Drees, L., Konigsmann, T., Jaspers, M. H. J., Pflanz, R., Riedel, D., & Schuh, R. (2019). Conserved function of the matriptase-prostasin proteolytic cascade during epithelial morphogenesis. *PLoS Genet*, *15*(1), e1007882. https://doi.org/10.1371/journal.pgen.1007882

Drees, L., Schneider, S., Riedel, D., Schuh, R., & Behr, M. (2023). The proteolysis of ZP proteins is essential to control cell membrane structure and integrity of developing tracheal tubes in Drosophila. *Elife*, *12*. https://doi.org/10.7554/eLife.91079

Faiyaz ul Haque, M., King, L. M., Krakow, D., Cantor, R. M., Rusiniak, M. E., Swank, R. T., Superti-Furga, A., Haque, S., Abbas, H., Ahmad, W., Ahmad, M., & Cohn, D. H. (1998). Mutations in orthologous genes in human spondyloepimetaphyseal dysplasia and the brachymorphic mouse. *Nat Genet*, *20*(2), 157-162. https://doi.org/10.1038/2458

Ford-Hutchinson, A. F., Ali, Z., Seerattan, R. A., Cooper, D. M., Hallgrimsson, B., Salo, P. T., & Jirik, F. R. (2005). Degenerative knee joint disease in mice lacking 3'-phosphoadenosine 5'-phosphosulfate synthetase 2 (Papss2) activity: a putative model of human PAPSS2 deficiency-associated arthrosis. *Osteoarthritis Cartilage*, *13*(5), 418-425. https://doi.org/10.1016/j.joca.2004.12.011

Fox, R. M., & Andrew, D. J. (2015). Changes in organelle position and epithelial architecture associated with loss of CrebA. *Biol Open*, *4*(3), 317-330. https://doi.org/10.1242/bio.201411205

Fox, R. M., Vaishnavi, A., Maruyama, R., & Andrew, D. J. (2013). Organ-specific gene expression: the bHLH protein Sage provides tissue specificity to Drosophila FoxA. *Development*, *140*(10), 2160-2171. https://doi.org/10.1242/dev.092924

Ghosh, N., & Treisman, J. E. (2024). Apical cell expansion maintained by Dusky-like establishes a scaffold for corneal lens morphogenesis. *Sci Adv*, *10*(34), eado4167. https://doi.org/10.1126/sciadv.ado4167

Gill, H. K., Cohen, J. D., Ayala-Figueroa, J., Forman-Rubinsky, R., Poggioli, C., Bickard, K., Parry, J. M., Pu, P., Hall, D. H., & Sundaram, M. V. (2016). Integrity of Narrow

Epithelial Tubes in the C. elegans Excretory System Requires a Transient Luminal Matrix. *PLoS Genet*, *12*(8), e1006205. https://doi.org/10.1371/journal.pgen.1006205

Groza, T., Gomez, F. L., Mashhadi, H. H., Munoz-Fuentes, V., Gunes, O., Wilson, R., Cacheiro, P., Frost, A., Keskivali-Bond, P., Vardal, B., McCoy, A., Cheng, T. K., Santos, L., Wells, S., Smedley, D., Mallon, A. M., & Parkinson, H. (2023). The International Mouse Phenotyping Consortium: comprehensive knockout phenotyping underpinning the study of human disease. *Nucleic Acids Res*, *51*(D1), D1038-D1045. https://doi.org/10.1093/nar/gkac972

Gunal, S., Hardman, R., Kopriva, S., & Mueller, J. W. (2019). Sulfation pathways from red to green. *J Biol Chem*, *294*(33), 12293-12312. https://doi.org/10.1074/jbc.REV119.007422

Gyoergy, A., Roblek, M., Ratheesh, A., Valoskova, K., Belyaeva, V., Wachner, S., Matsubayashi, Y., Sanchez-Sanchez, B. J., Stramer, B., & Siekhaus, D. E. (2018). Tools Allowing Independent Visualization and Genetic Manipulation of Drosophila melanogaster Macrophages and Surrounding Tissues. *G3 (Bethesda)*, *8*(3), 845-857. https://doi.org/10.1534/g3.117.300452

Herman, T., & Horvitz, H. R. (1999). Three proteins involved in Caenorhabditis elegans vulval invagination are similar to components of a glycosylation pathway. *Proc Natl Acad Sci U S A*, *96*(3), 974-979. https://doi.org/10.1073/pnas.96.3.974

Hwang, H. Y., Olson, S. K., Brown, J. R., Esko, J. D., & Horvitz, H. R. (2003). The Caenorhabditis elegans genes sqv-2 and sqv-6, which are required for vulval morphogenesis, encode glycosaminoglycan galactosyltransferase II and xylosyltransferase. *J Biol Chem*, *278*(14), 11735-11738. https://doi.org/10.1074/jbc.C200518200

Ismat, A., Cheshire, A. M., & Andrew, D. J. (2013). The secreted AdamTS-A metalloprotease is required for collective cell migration. *Development*, *140*(9), 1981-1993. https://doi.org/10.1242/dev.087908

Jazwinska, A., & Affolter, M. (2004). A family of genes encoding zona pellucida (ZP) domain proteins is expressed in various epithelial tissues during Drosophila embryogenesis. *Gene Expr Patterns*, *4*(4), 413-421. https://doi.org/10.1016/j.modgep.2004.01.003

Jazwinska, A., Ribeiro, C., & Affolter, M. (2003). Epithelial tube morphogenesis during

Drosophila tracheal development requires Piopio, a luminal ZP protein. *Nat Cell Biol*,

5(10), 895-901. https://doi.org/10.1038/ncb1049

Jovine, L., Darie, C. C., Litscher, E. S., & Wassarman, P. M. (2005). Zona pellucida domain

proteins. *Annu Rev Biochem*, *74*, 83-114.

https://doi.org/10.1146/annurev.biochem.74.082803.133039

Jovine, L., Qi, H., Williams, Z., Litscher, E., & Wassarman, P. M. (2002). The ZP domain is a

conserved module for polymerization of extracellular proteins. *Nat Cell Biol*, *4*(6), 457-

461. https://doi.org/10.1038/ncb802

Jullien, D., Crozatier, M., & Kas, E. (1997). cDNA sequence and expression pattern of the

Drosophila melanogaster PAPS synthetase gene: a new salivary gland marker. *Mech Dev*,

68(1-2), 179-186. https://doi.org/10.1016/s0925-4773(97)00136-6

Kamiyama, S., Suda, T., Ueda, R., Suzuki, M., Okubo, R., Kikuchi, N., Chiba, Y., Goto, S.,

Toyoda, H., Saigo, K., Watanabe, M., Narimatsu, H., Jigami, Y., & Nishihara, S. (2003).

Molecular cloning and identification of 3'-phosphoadenosine 5'-phosphosulfate

transporter. *J Biol Chem*, *278*(28), 25958-25963.

https://doi.org/10.1074/jbc.M302439200

Karlsson, M., Zhang, C., Mear, L., Zhong, W., Digre, A., Katona, B., Sjostedt, E., Butler, L.,

Odeberg, J., Dusart, P., Edfors, F., Oksvold, P., von Feilitzen, K., Zwahlen, M., Arif, M.,

Altay, O., Li, X., Ozcan, M., Mardinoglu, A.,…Lindskog, C. (2021). A single-cell type

transcriptomics map of human tissues. *Sci Adv*, *7*(31).

https://doi.org/10.1126/sciadv.abh2169

Kinnunen, T., Huang, Z., Townsend, J., Gatdula, M. M., Brown, J. R., Esko, J. D., & Turnbull, J.

E. (2005). Heparan 2-O-sulfotransferase, hst-2, is essential for normal cell migration in

Caenorhabditis elegans. *Proc Natl Acad Sci U S A*, *102*(5), 1507-1512.

https://doi.org/10.1073/pnas.0401591102

Kondylis, V., & Rabouille, C. (2009). The Golgi apparatus: lessons from Drosophila. *FEBS Lett*,

583(23), 3827-3838. https://doi.org/10.1016/j.febslet.2009.09.048

Kurima, K., Warman, M. L., Krishnan, S., Domowicz, M., Krueger, R. C., Jr., Deyrup, A., &

Schwartz, N. B. (1998). A member of a family of sulfate-activating enzymes causes

murine brachymorphism. *Proc Natl Acad Sci U S A*, *95*(15), 8681-8685. https://doi.org/10.1073/pnas.95.15.8681

Le, T. P., & Chung, S. (2021). Regulation of apical constriction via microtubule- and Rab11-dependent apical transport during tissue invagination. *Mol Biol Cell*, *32*(10), 1033-1047. https://doi.org/10.1091/mbc.E21-01-0021

Legan, P. K., Rau, A., Keen, J. N., & Richardson, G. P. (1997). The mouse tectorins. Modular matrix proteins of the inner ear homologous to components of the sperm-egg adhesion system. *J Biol Chem*, *272*(13), 8791-8801. https://doi.org/10.1074/jbc.272.13.8791

Litscher, E. S., & Wassarman, P. M. (2020). Zona pellucida genes and proteins and human fertility. *Trends Dev Biol*, *13*, 21-33. https://www.ncbi.nlm.nih.gov/pubmed/33335361

Luders, F., Segawa, H., Stein, D., Selva, E. M., Perrimon, N., Turco, S. J., & Hacker, U. (2003). Slalom encodes an adenosine 3'-phosphate 5'-phosphosulfate transporter essential for development in Drosophila. *EMBO J*, *22*(14), 3635-3644. https://doi.org/10.1093/emboj/cdg345

Luschnig, S., Batz, T., Armbruster, K., & Krasnow, M. A. (2006). serpentine and vermiform encode matrix proteins with chitin binding and deacetylation domains that limit tracheal tube length in Drosophila. *Curr Biol*, *16*(2), 186-194. https://doi.org/10.1016/j.cub.2005.11.072

Lye, C. M., Naylor, H. W., & Sanson, B. (2014). Subcellular localisations of the CPTI collection of YFP-tagged proteins in Drosophila embryos. *Development*, *141*(20), 4006-4017. https://doi.org/10.1242/dev.111310

Matthew, J., Vishwakarma, V., Le, T. P., Agsunod, R. A., & Chung, S. (2024). Coordination of cell cycle and morphogenesis during organ formation. *Elife*, *13*. https://doi.org/10.7554/eLife.95830

McAllister, K. A., Grogg, K. M., Johnson, D. W., Gallione, C. J., Baldwin, M. A., Jackson, C. E., Helmbold, E. A., Markel, D. S., McKinnon, W. C., Murrell, J., & et al. (1994). Endoglin, a TGF-beta binding protein of endothelial cells, is the gene for hereditary haemorrhagic telangiectasia type 1. *Nat Genet*, *8*(4), 345-351. https://doi.org/10.1038/ng1294-345

Mei, Z., Kuzhir, P., & Godeau, G. (2024). Update on Chitin and Chitosan from Insects: Sources, Production, Characterization, and Biomedical Applications. *Biomimetics (Basel)*, *9*(5). https://doi.org/10.3390/biomimetics9050297

Moussian, B., Tang, E., Tonning, A., Helms, S., Schwarz, H., Nusslein-Volhard, C., & Uv, A. E. (2006). Drosophila Knickkopf and Retroactive are needed for epithelial tube growth and cuticle differentiation through their specific requirement for chitin filament organization. *Development*, *133*(1), 163-171. https://doi.org/10.1242/dev.02177

Mueller, J. W., & Shafqat, N. (2013). Adenosine-5'-phosphosulfate--a multifaceted modulator of bifunctional 3'-phospho-adenosine-5'-phosphosulfate synthases and related enzymes. *FEBS J*, *280*(13), 3050-3057. https://doi.org/10.1111/febs.12252

Myat, M. M., & Andrew, D. J. (2000a). Fork head prevents apoptosis and promotes cell shape change during formation of the Drosophila salivary glands. *Development*, *127*(19), 4217-4226. https://doi.org/10.1242/dev.127.19.4217

Myat, M. M., & Andrew, D. J. (2000b). Organ shape in the Drosophila salivary gland is controlled by regulated, sequential internalization of the primordia. *Development*, *127*(4), 679-691. https://doi.org/10.1242/dev.127.4.679

Op De Beeck, K., Vermeire, S., Rutgeerts, P., & Bossuyt, X. (2012). Antibodies to GP2, the major zymogen granule membrane glycoprotein, in inflammatory bowel diseases. *Gut*, *61*(1), 162-164; author reply 164-165. https://doi.org/10.1136/gut.2010.233148

Plaza, S., Chanut-Delalande, H., Fernandes, I., Wassarman, P. M., & Payre, F. (2010). From A to Z: apical structures and zona pellucida-domain proteins. *Trends Cell Biol*, *20*(9), 524-532. https://doi.org/10.1016/j.tcb.2010.06.002

Porter, K. R., & Tamm, I. (1955). Direct visualization of a mucoprotein component of urine. *J Biol Chem*, *212*(1), 135-140. https://www.ncbi.nlm.nih.gov/pubmed/13233216

Renner, M., Bergmann, G., Krebs, I., End, C., Lyer, S., Hilberg, F., Helmke, B., Gassler, N., Autschbach, F., Bikker, F., Strobel-Freidekind, O., Gronert-Sum, S., Benner, A., Blaich, S., Wittig, R., Hudler, M., Ligtenberg, A. J., Madsen, J., Holmskov, U.,…Mollenhauer, J. (2007). DMBT1 confers mucosal protection in vivo and a deletion variant is associated with Crohn's disease. *Gastroenterology*, *133*(5), 1499-1509. https://doi.org/10.1053/j.gastro.2007.08.007

Rutherford, S. A., Goldberg, M. W., & Allen, T. D. (1997). Three-dimensional visualization of

the route of protein import: the role of nuclear pore complex substructures. *Exp Cell Res*,

232(1), 146-160. https://doi.org/10.1006/excr.1997.3487

Sarkissian, T., Timmons, A., Arya, R., Abdelwahid, E., & White, K. (2014). Detecting apoptosis

in Drosophila tissues and cells. *Methods*, *68*(1), 89-96.

https://doi.org/10.1016/j.ymeth.2014.02.033

Smith, S. J., Davidson, L. A., & Rebeiz, M. (2020). Evolutionary expansion of apical

extracellular matrix is required for the elongation of cells in a novel structure. *Elife*, *9*.

https://doi.org/10.7554/eLife.55965

Syed, Z. A., Bouge, A. L., Byri, S., Chavoshi, T. M., Tang, E., Bouhin, H., van Dijk-Hard, I. F.,

& Uv, A. (2012). A luminal glycoprotein drives dose-dependent diameter expansion of

the Drosophila melanogaster hindgut tube. *PLoS Genet*, *8*(8), e1002850.

https://doi.org/10.1371/journal.pgen.1002850

Tang, W. J., Fernandez, J., Sohn, J. J., & Amemiya, C. T. (2015). Chitin is endogenously

produced in vertebrates. *Curr Biol*, *25*(7), 897-900.

https://doi.org/10.1016/j.cub.2015.01.058

Tecle, E., Diaz-Balzac, C. A., & Bulow, H. E. (2013). Distinct 3-O-sulfated heparan sulfate

modification patterns are required for kal-1-dependent neurite branching in a context-

dependent manner in Caenorhabditis elegans. *G3 (Bethesda)*, *3*(3), 541-552.

https://doi.org/10.1534/g3.112.005199

Tomancak, P., Beaton, A., Weiszmann, R., Kwan, E., Shu, S., Lewis, S. E., Richards, S.,

Ashburner, M., Hartenstein, V., Celniker, S. E., & Rubin, G. M. (2002). Systematic

determination of patterns of gene expression during Drosophila embryogenesis. *Genome

Biol*, *3*(12), RESEARCH0088. https://doi.org/10.1186/gb-2002-3-12-research0088

Tonning, A., Hemphala, J., Tang, E., Nannmark, U., Samakovlis, C., & Uv, A. (2005). A

transient luminal chitinous matrix is required to model epithelial tube diameter in the

Drosophila trachea. *Dev Cell*, *9*(3), 423-430. https://doi.org/10.1016/j.devcel.2005.07.012

Turnbull, J., Drummond, K., Huang, Z., Kinnunen, T., Ford-Perriss, M., Murphy, M., &

Guimond, S. (2003). Heparan sulphate sulphotransferase expression in mice and

Caenorhabditis elegans. *Biochem Soc Trans*, *31*(2), 343-348.

https://doi.org/10.1042/bst0310343

Uhlen, M., Bjorling, E., Agaton, C., Szigyarto, C. A., Amini, B., Andersen, E., Andersson, A. C., Angelidou, P., Asplund, A., Asplund, C., Berglund, L., Bergstrom, K., Brumer, H., Cerjan, D., Ekstrom, M., Elobeid, A., Eriksson, C., Fagerberg, L., Falk, R.,…Ponten, F. (2005). A human protein atlas for normal and cancer tissues based on antibody proteomics. *Mol Cell Proteomics*, *4*(12), 1920-1932. https://doi.org/10.1074/mcp.M500279-MCP200

van den Boom, J., Heider, D., Martin, S. R., Pastore, A., & Mueller, J. W. (2012). 3'-Phosphoadenosine 5'-phosphosulfate (PAPS) synthases, naturally fragile enzymes specifically stabilized by nucleotide binding. *J Biol Chem*, *287*(21), 17645-17655. https://doi.org/10.1074/jbc.M111.325498

Venkatachalam, K. V. (2003). Human 3'-phosphoadenosine 5'-phosphosulfate (PAPS) synthase: biochemistry, molecular biology and genetic deficiency. *IUBMB Life*, *55*(1), 1-11. https://doi.org/10.1080/1521654031000072148

Venkatachalam, K. V., Akita, H., & Strott, C. A. (1998). Molecular cloning, expression, and characterization of human bifunctional 3'-phosphoadenosine 5'-phosphosulfate synthase and its functional domains. *J Biol Chem*, *273*(30), 19311-19320. https://doi.org/10.1074/jbc.273.30.19311

Wang, S., Jayaram, S. A., Hemphala, J., Senti, K. A., Tsarouhas, V., Jin, H., & Samakovlis, C. (2006). Septate-junction-dependent luminal deposition of chitin deacetylases restricts tube elongation in the Drosophila trachea. *Curr Biol*, *16*(2), 180-185. https://doi.org/10.1016/j.cub.2005.11.074

Weiszmann, R., Hammonds, A. S., & Celniker, S. E. (2009). Determination of gene expression patterns using high-throughput RNA in situ hybridization to whole-mount Drosophila embryos. *Nat Protoc*, *4*(5), 605-618. https://doi.org/10.1038/nprot.2009.55

Wilkin, M. B., Becker, M. N., Mulvey, D., Phan, I., Chao, A., Cooper, K., Chung, H. J., Campbell, I. D., Baron, M., & MacIntyre, R. (2000). Drosophila dumpy is a gigantic extracellular protein required to maintain tension at epidermal-cuticle attachment sites. *Curr Biol*, *10*(10), 559-567. https://doi.org/10.1016/s0960-9822(00)00482-6

Wolfstetter, G., & Holz, A. (2012). The role of LamininB2 (LanB2) during mesoderm differentiation in Drosophila. *Cell Mol Life Sci*, *69*(2), 267-282. https://doi.org/10.1007/s00018-011-0652-3

Wong, S. H., Hamel, L., Chevalier, S., & Philip, A. (2000). Endoglin expression on human microvascular endothelial cells association with betaglycan and formation of higher order complexes with TGF-beta signalling receptors. *Eur J Biochem*, *267*(17), 5550-5560. https://doi.org/10.1046/j.1432-1327.2000.01621.x

Xia, B., Royall, J. A., Damera, G., Sachdev, G. P., & Cummings, R. D. (2005). Altered O-glycosylation and sulfation of airway mucins associated with cystic fibrosis. *Glycobiology*, *15*(8), 747-775. https://doi.org/10.1093/glycob/cwi061

Yan, B., Zhang, Z. Z., Huang, L. Y., Shen, H. L., & Han, Z. G. (2012). OIT3 deficiency impairs uric acid reabsorption in renal tubule. *FEBS Lett*, *586*(6), 760-765. https://doi.org/10.1016/j.febslet.2012.01.038

Yang, K., Liu, M., Feng, Z., Rojas, M., Zhou, L., Ke, H., & Pastor-Pareja, J. C. (2021). ER exit sites in Drosophila display abundant ER-Golgi vesicles and pearled tubes but no megacarriers. *Cell Rep*, *36*(11), 109707. https://doi.org/10.1016/j.celrep.2021.109707

Zhang, L., Song, W., Li, T., Mu, Y., Zhang, P., Hu, J., Lin, H., Zhang, J., Gao, H., & Zhang, L. (2023). Redox switching mechanism of the adenosine 5'-phosphosulfate kinase domain (APSK2) of human PAPS synthase 2. *Structure*, *31*(7), 826-835 e823. https://doi.org/10.1016/j.str.2023.04.012

Zhang, P., Zhang, L., Hou, Z., Lin, H., Gao, H., & Zhang, L. (2022). Structural basis for the substrate recognition mechanism of ATP-sulfurylase domain of human PAPS synthase 2. *Biochem Biophys Res Commun*, *586*, 1-7. https://doi.org/10.1016/j.bbrc.2021.11.062

Zhang, Y., Foster, J. M., Nelson, L. S., Ma, D., & Carlow, C. K. (2005). The chitin synthase genes chs-1 and chs-2 are essential for C. elegans development and responsible for chitin deposition in the eggshell and pharynx, respectively. *Dev Biol*, *285*(2), 330-339. https://doi.org/10.1016/j.ydbio.2005.06.037

Zhou, W., Chang, J., Wang, X., Savelieff, M. G., Zhao, Y., Ke, S., & Ye, B. (2014). GM130 is required for compartmental organization of dendritic golgi outposts. *Curr Biol*, *24*(11), 1227-1233. https://doi.org/10.1016/j.cub.2014.04.008

Zhu, X., Sen, J., Stevens, L., Goltz, J. S., & Stein, D. (2005). Drosophila pipe protein activity in the ovary and the embryonic salivary gland does not require heparan sulfate glycosaminoglycans. *Development*, *132*(17), 3813-3822. https://doi.org/10.1242/dev.01962

Figure 1

Figure 2

Figure 3

Figure 4

A — CrebA DCP-1 Ecad — WT — 10 μm

B — *Papss²* — DCP-1

C — *Papss²* — 2 μm

D — Number of acellular regions in the SG (St. 16)

WT Papss² sage>Papss-PD; Papss² sage>Papss-PD^Y193A,F383P; Papss²

E — Number of SG cells

WT Papss² Papss²/Df sage>p35; Papss² sage>Papss-PE; Papss² sage>Papss-PD^Y193A,F383P; Papss²

F — CrebA Hemo3x-mCh Ecad — control — 10 μm — Hemo3x-mCh

G — *Papss²*

H — DAPI Hemo3x-mCh Ndg — control — 10 μm — 5 μm — Hemo3x-mCh — Ndg

I — *Papss²*

Figure 5

Figure 6

Figure 7

A

WT

st. 14 → Uniform lumen expansion → st. 16

Papss st. 16 — Failed/irregular lumen expansion

pio st. 16 — Constrictions of the lumen

Np st. 16 — Constrictions of the lumen

B

Organization of the ZP protein-containing aECM →

Dpy-Pio(ZP-C) scaffold

Pio(ZP-C) missing or masking Dpy aggregation

No Pio No luminal Dpy

No luminal Pio No luminal Dpy

Legend:
Dpy — Pio — NP — Trypsin-like serine protease
● EGF-like repeats
▭ transmembrane domain
▭ ZP domain (Dpy)
▭ ZP domain (Pio)