Diminished gap junction coupling under diabetogenic conditions does not drive loss of functional β-cell sub-populations

Claire H Levitt[a], Dominic Isaacs[a], Maria S Hansen[b], Vira Kravets[a], Jennifer K Briggs[a], Richard KP Benninger[a,b,1]

[a] Department of Bioengineering, University of Colorado, Anschutz Medical campus, Aurora, CO. USA

[b] Barbara Davis center for childhood diabetes, University of Colorado, Anschutz Medical campus, Aurora, CO. USA

[1] To whom correspondence should be addressed: richard.benninger@cuanschutz.edu Tel: (303) 724-6388. Fax: (303) 724-5800.

1775 Aurora court, M20-4306D, mailstop B140, University of Colorado Anschutz Medical campus, Aurora, CO. 80045.

Running Title: β-cell sub-populations under diabetogenic conditions

Key Words: Optogenetics, Islet Dynamics; Optical Microscopy; Diabetes; Gap junctions

Classification: Systems Biology

Manuscript word count: 4432

Abstract word count: 272

Number of figures: 4

Abstract

Within the islets of Langerhans, gap junction coupling is important for synchronizing oscillatory free-calcium activity ($[Ca^{2+}]$) and regulating pulsatile insulin release. In islets from multiple models of diabetes, gap junction coupling is disrupted, and $[Ca^{2+}]$ synchronization and pulsatile insulin is lost. Functional sub-populations have been identified within the islet that are linked to driving synchronized $[Ca^{2+}]$ and insulin release. These sub-populations can be disrupted under conditions associated with diabetes such as glucolipotoxicity and inflammatory environments, and their loss may drive islet dysfunction. Here we investigate how loss of gap junction coupling influences functional subpopulations under diabetogenic environments. We treated islets with a cocktail of pro-inflammatory cytokines and specifically recovered gap junction coupling via co-treatment with a Cx36 mimetic peptide S293 that was previously shown to protect gap junction permeability and synchronized $[Ca^{2+}]$ dynamics. We performed calcium imaging and ChR2 stimulation, analyzed islet $[Ca^{2+}]$ dynamics and the presence of functional sub-populations including hubs and first-responders. 1h or 24h cytokine-treatment disrupted gap junction coupling, which was fully recovered by S293 peptide co-treatment. Treatment with pro-inflammatory cytokines decreased the recruitment of $[Ca^{2+}]$ upon ChR2 stimulation, increased the time between first and last responding cells upon glucose stimulation, and reduced the number and consistency of hub cells. When preserving gap junction coupling by S293 during cytokine treatment, the presence and consistency of these sub-populations was only marginally improved. We therefore concluded that restoration of gap junctions alone is not sufficient to recover functional sub-populations and their influence over islet function upon diabetogenic conditions associated with diabetes. Thus, preventing a disruption to intrinsic β-cell properties that define functional subpopulations is likely important for preserving these sub-populations during diabetes.

Introduction

Diabetes Mellitus (DM) affects approximately 537 million people globally [1,2], and is characterized by the dysfunction and/or death of insulin-secreting β-cells in the islets of Langerhans in the pancreas. β-cells in the islet secrete insulin in response to elevations in blood glucose or other nutrients. At elevated glucose, β-cells show increased ATP/ADP following the metabolism of glucose, which drives membrane depolarization via closure of ATP-sensitive K+ (K_{ATP}) channels and activation of voltage gated calcium (Ca_V) channels. Subsequent oscillations of intracellular free-calcium ($[Ca^{2+}]$) across the islet drive the release of insulin in discrete pulses.

The dynamics of $[Ca^{2+}]$ are synchronized across β-cells in the islet, which is facilitated by connexin-36 (Cx36) gap junctions [3-5]. Gap junctions synchronize the β-cell response to glucose despite heterogeneity in β-cell metabolism and excitability [6,7]. Under low glucose conditions, gap junction-mediated hyper-polarizing currents suppress spontaneous elevations in $[Ca^{2+}]$ from excitable β-cells [8,9]. At high glucose, gap junction-mediated depolarizing currents coordinate both first-phase $[Ca^{2+}]$ and oscillatory second phase $[Ca^{2+}]$ responses [3]. In Cx36 knockout models, synchronous $[Ca^{2+}]$ responses are disrupted [3,5,9], both first phase insulin secretion and second phase insulin pulses are diminished and glucose tolerance is impaired [5,10]. Importantly, islet gap junction coupling is disrupted under conditions associated with diabetes including by glucolipotoxicity [2,11,12] and pro-inflammatory cytokines [13] as well as in mouse models of diabetes [14-16] and human [11,12,17]. Synchronous $[Ca^{2+}]$ activity is also disrupted under conditions associated with diabetes [11,13,17,18] likely as a result of disrupted Cx36 gap junction coupling.

β-cells are heterogeneous in gene expression [19-21], glucose sensitivity [6,19], insulin secretion [22], and cell-to-cell communication [23]. As a result, β-cells exhibit functional heterogeneity, and display distinct insulin secretory patterns. Previous studies found that ~75% of β-cells exhibited a fixed response to glucose and ~25% fluctuated between states of responsive and unresponsiveness [24]. The characterization of β-cell heterogeneity has led to the identification of functional subpopulations which are thought to play distinct roles in governing the spatiotemporal $[Ca^{2+}]$ and insulin secretory dynamics of the islet [25,26]. For example, following optogenetic ChR2 stimulation, a population of β-cells can recruit and elevate $[Ca^{2+}]$ in a large number of neighboring β-cells within the islet [27]. Similarly, 'first-responder cells', that are first to show elevated $[Ca^{2+}]$ upon elevated glucose stimulation, can

recruit and coordinate less excitable cells to respond earlier [28]. This first-responder

subpopulation is distinct from previously identified 'hub cells' that exhibit a high degree of co-

activity with neighboring β-cells [29] and have been shown to play a critical role in driving the

synchronized oscillatory dynamics. Importantly, upon both glucolipotoxic environments and

pro-inflammatory cytokines, the proportion of highly connected 'hub cells' declines [29]. This

finding suggests that a selective loss of functional sub-populations may be sufficient to drive

islet dysfunction in diabetes. However, the mechanisms underlying this loss of cells are unclear.

In this study, we seek to understand how functional sub-populations of cells are

disrupted under conditions associated with diabetes. We specifically ask whether the loss of

gap junction coupling that is associated with diabetes is a primary cause for the loss of these

sub-populations and their influence over islet function. We first test how functional sub-

populations are influenced by pro-inflammatory cytokine treatment. We then test for a role of

gap junction coupling, by using a Cx36 mimetic peptide [12] that was previously shown to

specifically protect gap junction permeability and retain glucose-stimulated [Ca^{2+}] dynamics

under conditions associated with diabetes.

Materials and Methods

Mice/Animal Care: Islets expressing ChR2-YFP in β-cells was achieved by crossing Pdx^{PB}-Cre (Jackson Laboratory) and ROSA26-lox-STOP-lox-ChR2-YFP mice (Jackson Laboratory). Islets expressing GCaMP6s in β-cells was achieved by crossing MIP-Cre^{ER} mice (Jackson Laboratory) and ROSA26-lox-STOP-lox-GCaMP6s mice (Jackson Laboratory). Mice were genotypes via qPCR (Transnetyx). All animal experiments were approved by the University of Colorado Institutional Animal Care and Use Committee (IACUC Protocol number: 000024). Mice were housed in a temperature-controlled environment with a 12 hr light/dark cycle and continuous access to both food and water. For mice with β-cell specific GCaMP6s expression, tamoxifen (solubilized in corn oil) was administered via IP injection for 5 consecutive days (50 mg/kg of body weight) to facilitate CreER-mediated recombination.

Islet Isolation and Culture: Islets were isolated from mice under ketamine/xylazine anesthesia and euthanized via exsanguination. Collagenase (Type V, Sigma) was delivered to the pancreases at a concentration of 2.5 mg/mL via the bile duct. After inflation, the pancreas was surgically removed and digested (5 min at 37 °C). Islets were then hand-picked and cultured in RPMI medium (Gibco) containing 10% fetal bovine serum, 100 U/mL penicillin, 100 ug/mL streptomycin, and 11 mM glucose. Islets were incubated at 37°C, 5% CO2.

Islet Treatment: Islets were treated overnight with a '0.1x' cocktail of proinflammatory cytokines (5 ng/mL IL-1β , 10 ng/mL TNF-α, and 100 ng/mL IFN-γ), as previously used [13] To restore gap junction coupling, islets were co-treated with 30 μM mimetic peptide S293 for 1-6 hours for experiments using optogenetic stimulation, and 24 hours for experiments measuring gap junction permeability (FRAP) and calcium activity ([Ca^{2+}]).

Fluorescent Recovery After Photobleaching and Analysis Gap junction permeability was assessed by Fluorescence Recovery After Photobleaching (FRAP) [4]. Isolated islets were immobilized onto glass bottom dishes (MatTek) using cell tissue adhesive (CellTak, Corning) and incubated at 37°C, 5% CO2 overnight. Islets were then loaded with 12.5 mM cationic dye Rhodamine-123 for 30 minutes at 37°C in Hank's Balanced Salt Solution (HBSS) at 5mM glucose. Islets were washed and imaged at room temperature in HBSS using a Zeiss LSM 800 confocal microscope with a x40 1.2NA water immersion objective. Rhodamine-123 was excited

at 488 nm. Half the islet area was selected using a rectangular ROI and photobleached for 25 frames at 40% laser power and achieved ~50% bleaching on average. Images were collected every 5 seconds for 6 to 10 minutes.

Image analysis of Rh123 dye diffusion after photobleaching was conducted using MATLAB and Image J. Fluorescent recovery was measured within a region of interest encompassing the entire photobleached area and individual cells. The initial fluorescence intensity was considered the point directly after photobleaching (I_P). Fluorescence recovery was the final fluorescence (I_F) once the diffusion of dye reached a plateau. The recovery was calculated as a percentage, relative to the initial intensity before photobleaching (I_0) as follows:

$$\% \, Recovery = 100 * \frac{I_F - I_P}{I_0 - I_P}$$

The rate of fluorescent recovery was calculated based on the linearized exponential fluorescent recovery curve of the bleached area over time (I_i) normalized to the total recovery. Calculations were conducted as follows:

$$Rate \, of \, recovery = \, -Ln(1 - \frac{I_i - I_P}{I_0 - I_P})$$

Channelrhodopsin-2 Single Cell Entrainment and Analysis: ChR2-YFP expressing islets were incubated with pro-inflammatory cytokines, with or without S293 mimetic peptide, or S293 peptide alone for 6 hours after isolation in RPMI medium (Gibco) containing 10% fetal bovine serum, 100 U/mL penicillin, 100 ug/mL streptomycin, and 11 mM glucose at 37°C, 5% CO2 prior to study.

Islets were loaded with 2μM of calcium indicator Rhodamine-2 (Rhod-2) in BMHH imaging solution (125mM NaCl, 5.7mM KCl, 2.5mM CaCl2, 1.2mM MgCl2, 10mM HEPES, 2mM glucose, and 0.1% BSA, pH 7.4) for 35 minutes at 37°C. Islets were then washed and imaged on a glass bottom dish in a 5mM glucose imaging solution using a LSM 800 confocal microscope kept at 37°C .

ChR2-YFP was excited at 488 nm. A rectangular ROI was drawn within the bounds of a single cell, so as to avoid stimulating neighboring cells that express ChR2-YFP. The cell was activated using a photobleaching macro (intensity 5% laser power) for 8 stimulations at a frequency of 0.1 Hz. Rhod-2 was simultaneously excited at 560 nm and the corresponding Ca^{2+} response across the whole islet was measured based on fluorescent intensity of Rhod-2 that was coincident to ChR2 stimulation.

MATLAB was used to analyze Channelrhodopsin-2 recruitment data, as previously
described [27]. Islets were selected by hand drawing a region of interest around the islet.
Background was identified as regions that did not show any calcium response. Activity was
based on the fluorescent intensity of the Ca^{2+} indicator Rhod-2 upon ChR2 stimulation and
assessed using individual pixel time-courses. A pixel was considered 'active' if its intensity was
twice that of the background. A threshold of 0.15 was used to determine which regions of the
islet were activate due to ChR2 stimulation compared to the islet average. The percent of the
islet area that was 'active' was calculated as a proportion of the entire islet area.

GCaMP6 Imaging of Calcium Dynamics and Analysis: Isolated islets expressing β-cell specific
GCaMP6 were incubated overnight with proinflammatory cytokines with or without S293
mimetic peptide, or S293 peptide alone. 30 minutes prior to imaging, islets were fasted at 2mM
glucose in BMHH imaging solution. Islets were then transferred to a glass bottom dish and
imaged on an LSM 800 confocal microscope. GCaMP6 fluorescence was excited at 488 nm
with images collected at 1 frame/sec for approximately 80 minutes. 3-5 minutes after the start
of imaging, glucose was raised from 2mM to 11mM. For experiments measuring consistency
between first responders and hub cells, glucose was subsequently lowered to 2mM after
achieving second phase oscillations for 10-15 minutes. After the islet had returned to basal
activity without oscillations (~15 minutes), the glucose concentration was increased again to
11mM. Samples were imaged at either x20NA or x40NA water immersion and held at 37°C.
Calcium dynamics were analyzed using MATLAB, first phase response time,
identification of first and last responders, and characterization of highly connected hub cells
was determined as previously described [28]. Briefly, to identify the response time, we
determined the time at which individual β-cells reached their half maximum amplitude following
stimulation from low to high glucose. This response time was rank ordered for all cells in the
islet. The earliest 10% were classed as first responders and the latest 10% classed as last
responders. ΔT was calculated as the difference in time between the means of the last
responders and first responders. The percent consistency was determined as a proportion of
first 5 responding cells of the initial stimulation that were also first responders during repeated
stimulation.
Highly correlated β-cell cells were identified using the calcium time-course of the high
glucose second phase. A highly correlated β-cell is a β-cell whose calcium activity strongly

parallels many other β-cell in the islet. To quantify the degree of correlation, the cross-
correlation value obtained using the xcorr function in MATLAB across all measured β-cell
calcium time traces was compared among each cell. "Links" were derived using a threshold of
correlation above 0.9. The top 60% of linked cells were considered hubs, (as defined in [Cite
Johnston et al].
200
201   *Statistical analysis:* 2-sided t-test was used to compare 2 groups, whereas ANOVA with
202   Tukey's multiple comparisons test was used to compare multiple treatment groups, where *
203   indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$, **** indicates $p < 0.0001$.
204

Results

Successful restoration of gap junction coupling by Cx36 S293 mimetic peptide under pro-inflammatory cytokines: To test whether functional sub-populations are disrupted by a cocktail of pro-inflammatory cytokines (0.5 ng/mL IL1-β, 1 ng/mL TNF-α, and 10 ng/mL IFN-γ), and whether sub-population loss depends on loss of gap junction coupling, we followed an approach outlined in Figure 1A. To assess gap junction permeability and confirm a disruption to gap junction coupling by pro-inflammatory cytokines and recovery by S293 mimetic peptide we used FRAP (Figure 1.B), as previously established [4]. Islets treated with pro-inflammatory cytokines showed a reduction in gap junction permeability, as indicated by a slower recovery rate. During co-treatment with the S293 mimetic peptide islets showed a significant increase in the recovery rate compared to cytokine treatment alone (Figure 1.C, p = 0.0471). No significant difference was observed between untreated control and co-treatment groups (p = 0.9958) or untreated control vs S293 mimetic peptide alone (p = 0.9979). Thus co-treatment with S293 peptide fully restores gap junction permeability, consistent with previous studies that use the S293 mimetic peptide to specifically recover gap junction coupling in diabetogenic environments [12].

Restoration of gap junction coupling provides a partial recovery to the cytokine-mediated disruption of ChR2-mediated recruitment of [Ca^{2+}]: With the conditions established to restore gap junction coupling under a diabetogenic environment, we asked whether functional subpopulations of β-cells are also restored.. Previous studies applied the optogenetic construct ChannelRhodopsin-2 (ChR2) and single cell stimulation to discover a subpopulation of β-cells with elevated metabolic activity that can recruit large numbers of neighboring cells when activated [27]. We therefore first examined how pro-inflammatory cytokines influenced these sub-populations and then tested the effects of gap junction coupling restoration on these sub-populations.

Islets from Pdx1^{PB}-Cre;Rosa-LSL-ChR2-YFP mice that express ChR2 in β-cells were loaded with the Ca^{2+} dye Rhod2. ChR2 expressing cells were subjected to focal stimulation in single β-cells and the spatial Ca^{2+} response was observed in neighboring cells (Figure 2.A). We quantified the proportion of β-cells in the imaging plane that showed Ca^{2+} elevations coincident with ChR2 stimulation. This was repeated in ChR2-expressing islets treated with a cocktail of pro-inflammatory cytokines (Figure 2.B) and/or Cx236 S293 peptide mimetic.

Under all treatments we observed heterogeneity in the recruitment of elevated [Ca^{2+}]

following single cell stimulation (Figure 2.C). In untreated control islets, on average ~42% of

cells within the imaging plane showed elevated [Ca^{2+}] following single cell stimulation, with a

substantial variation (Figure 2.C,D) as previously observed [27]. Treatment with pro-

inflammatory cytokines (for 1 hour) led to a significant decrease in the proportion of cells

showing elevated [Ca^{2+}] following single cell stimulation (~15%) compared to untreated controls

(p < 0.0001, Figure 2.C,D). Co-treatment with the S293 mimetic peptide led to a moderate

increase in elevated [Ca^{2+}] following single cell stimulation (~25%) (Figure 2.C,D), however this

was still significantly lower than in untreated control (p = 0.0026). Treatment with S293 peptide

alone showed a similar proportion of cells with elevated [Ca^{2+}] (~35%) to untreated control

islets (Figure 2.C,D).

Untreated control islets showed broad heterogeneity in the recruitment of [Ca^{2+}] (Figure

2.C), as previously observed [27]. This included a population of cells that recruited elevated

[Ca^{2+}] across the median recruited area of the untreated control islet (>30%, Figure 2.C), as

also previously observed [27]. Under cytokine treated (1 hour), this 'high recruiting' population

of cells was almost completely absent (~4%, Figure 2.E). Co-treatment with S293 mimetic

peptide partially recovered this population of cells (~10%, Figure 2.E), but again much less

than in either untreated controls or S293 peptide treatment alone.

Therefore, treatment with pro-inflammatory cytokines reduced the ChR2-mediated

recruitment of cells to increase [Ca^{2+}], with an abolishment of 'high-recruiting' cells. Restoration

of gap junction coupling only partially restored ChR2-mediatred recruitment of cells to increase

[Ca^{2+}] with only a small recovery of 'high-recruiting' cells.

Restoration of gap junction coupling does not recover the cytokine-mediated disruption to first

responder cells and the 1st phase [Ca^{2+}] response: We next examined the impact of pro-

inflammatory cytokines and the restoration of gap junction coupling on first responder cells and

the first phase [Ca^{2+}] response. Islets from β-cell specific GCaMP6 mice were used to assess

changes in [Ca^{2+}] following glucose elevation. We measured the [Ca^{2+}] response of individual

cells and ranked their response time after glucose stimulation. First responder cells were

identified as the 10% of cells which showed the earliest response, and last responders as the

10% of cells that showed the latest response, with the spread in response time (Δt) being the

time between first and last responders (Figure 3.A). In control untreated islets the response

time was fairly narrow ($\Delta t=175s$). Upon treatment with pro-inflammatory cytokines for 1 hour,

there was no significant impact on the time between first and last responders (Figure 3.A-B).

However following treatment with pro-inflammatory cytokines for 24 hours there was a

significant increase in the time between first and last responders ($\Delta t=380s$) compared to

untreated control islets (p=0.0036, Figure 3.A-B). Using this 24 hour treatment, co-treatment

with S293 peptide did not show a decrease in Δt compared to cytokines alone, indicating

restoration of gap junction coupling was not sufficient to recover the narrow spread of the first

phase response (Figure 3.C). S293 peptide alone showed a similar time between first and last

responders as untreated controls. Under 1 hour cytokine treatment, co-treatment with S293

had no impact (Supplemental 1).

Clusters of first responders were previously found to be consistent upon repeated

glucose stimulations [28]; where ~35% of first responder cells remained first responders and

~75% of first responders remained within a 1 cell distance separation. We tested whether this

consistency was affected by both pro-inflammatory cytokines and restoration of gap junction

coupling. We identified the first 5 responding cells during an initial glucose stimulation and

determined what proportion of these 5 cells remained first responders during a repeated

glucose stimulation (Figure 3.D). Upon cytokine treatment, there was a small but non-

significant change in the average consistency of first responders upon repeated glucose

stimulation compared to untreated controls (34% vs. 38% respectively, p=0.9769) (Figure 3.E).

Restoration of gap junction coupling via co-treatment with S293 mimetic peptide had no impact

on the consistency of first responders relative to cytokine only treatment or untreated controls

(Figure 3.E) (p=0.8758).

Therefore, treatment with pro-inflammatory cytokines reduced the coordination of the

first phase $[Ca^{2+}]$ response. Restoration of gap junction coupling does not provide any recovery

of first-responder cells, nor the coordination of the first phase $[Ca^{2+}]$ response.

Restoration of gap junction coupling provides a partial recovery to the cytokine-mediated

disruption of highly connected β-cell hubs: Finally, we examined the impact of pro-inflammatory

cytokines and the restoration of gap junction coupling on highly connected β-cell hubs. β-cell

hubs have been demonstrated to maintain coordinated $[Ca^{2+}]$ dynamics and are distinct from

first responder cells [25,28,29]. Importantly previous work indicated a decrease in β-cell hubs

upon diabetogenic conditions [13,29]. Following Ca^{2+} imaging we analyzed individual cell time-

courses and formed a functional network and identified the most highly connected cells as β-
cell hubs (Figure 4.A, see Methods). In control islets the number of highly synchronized
functional connections and highly synchronized hubs was high and consistent with prior
measurements [29,30]. Upon treatment with pro-inflammatory cytokines for 1 hour, there was
no significant impact the mean degree of correlation (Figure 4.A-B). However following
treatment with pro-inflammatory cytokines for 24 hours, there was a significant decrease in the
mean degree of correlation (p = 0.0375, Figure 4. A-C). There was also a decrease in proportion
of highly synchronized functional connections or "links" (p=0.0750, Figure 4.D) and highly
connected β-cell hubs (p = 0.2087, Figure 4.E). Co-treatment with S293 peptide did not show
a significant change in either functional connections or β-cell hubs compared to cytokines alone
(Figure 4.D,E). This indicates that restoration of gap junction coupling was not sufficient to
recover highly synchronized functional connections and highly connected β-cell hubs. S293
peptide alone showed similar functional connections and β-cell hubs as untreated controls.
Under 1 hour cytokine treatment, co-treatment with S293 had no impact (Supplemental 2).

Similar to our analysis of first-responder cell consistency, an average of ~50% of β-cell
hubs remained as hubs following repeated glucose stimulation (Figure 4.F). We tested whether
this consistency is affected by both pro-inflammatory cytokines and restoration of gap junction
coupling. Upon cytokine treatment, there was a decrease in the consistency of hubs (~32%)
compared to untreated control islets (~53%). Restoration of gap junction coupling via co-
treatment with S293 mimetic peptide showed a trend to improved the consistency of hubs
compared to cytokine treatment alone (~67% vs ~32%, p=0.4285, Figure 4.G), with similar
measurements to S293 peptide treatment alone, albeit lacking statistical significance.

Therefore, treatment with pro-inflammatory cytokines reduced the presence of highly
synchronized functional connections and β-cell hubs. Restoration of gap junction coupling and
islet-wide synchronization has little influence on hubs but maintains the consistency of those
remaining hub cells.

Discussion

Cx36 gap junctions have previously been identified as an important regulator of β-cell coordination and islet function through the electrical coupling of neighboring β-cells. Pharmacological or genetic improvements in Cx36 gap junctions has been shown to recover $[Ca^{2+}]$ and insulin secretion responses in models of diabetes invitro, and delay onset of diabetes in vivo, indicating a protective role that gap junctions play in islet function [12,31]. Functional sub-populations (which emerge as a result of β-cell heterogeneity) have been identified that can drive the coordinated islet electrical response [27]. The goal of this study was to examine the influence of gap junctions on functional sub-populations of β-cells in a cytokine-mediated diabetogenic environment. Towards this, in the presence of a cocktail of pro-inflammatory cytokines, we recovered Cx36 gap junction coupling using a mimetic peptide which we previously developed [12]. We then applied both optogenetics and calcium imaging to measure cell recruitment and calcium dynamics to identify and characterize three functional sub-populations that have previously been described [28] including first responders and hubs. We found that while gap junction coupling could be successfully recovered under pro-inflammatory conditions, as indicated by FRAP, optogenetic stimulation, and/or analysis of calcium dynamics revealed that functional subpopulations remained largely impaired. Thus, while Cx36 gap junction coupling is critical for overall islet function, loss of cell-intrinsic properties likely drives the impairment of functional sub-populations under diabetogenic conditions.

Impairment of functional sub-populations under diabetogenic conditions: Increased levels of proinflammatory cytokines within the islet microenvironment, including IL1-β and TNF-α, is a key feature of diabetes and has been attributed to driving islet dysfunction [16,32,33]. Prior studies have found that functional sub-populations are also disrupted in diabetes, such as reduced numbers of β-cell hubs under both pro-inflammatory cytokines and gluco-lipo-toxic conditions [29],reduced consistency of wave initiator / leader cells under high-fat diet conditions [15] and reduced presence of hubs and leaders in islets from human subjects with type2 diabetes [34]. In this study we extended these findings to show both the cells that recruit a large proportion of the neighbors to show elevated $[Ca^{2+}]$ (high-responders), and the first responder cells that respond earliest to glucose were also both disrupted by pro-inflammatory cytokines. This points to a general occurrence for functional sub-populations to be disrupted under diabetogenic conditions. Notably, each of these functional sub-population is associated with

driving a specific aspect of islet electrical response, which were also disrupted under the pro-inflammatory cytokines. For example, β-cell hubs coordinate oscillatory dynamics across the islet, and the synchronization of these oscillations was disrupted by pro-inflammatory cytokines. Similarly, first-responders coordinate and enhance the initial first-phase response to glucose elevation and the coordination of this response is disrupted by pro-inflammatory cytokines. Thus disruption to these functional sub-populations may be contributing to islet dysfunction under diabetogenic conditions.

Influence of gap junction disruption to functional subpopulations under diabetogenic conditions: Functional sub-populations such as β-cell hubs, first-responders and those that show high ChR2-recruitment of [Ca^{2+}] all exert their influence to coordinate and recruit other β-cells via Cx36 gap junction coupling. Thus, a natural hypothesis is that pro-inflammatory cytokines disrupt these sub-populations and their action, because of diminished gap junction coupling. By using the S293 peptide mimetic we can recover functional gap junction coupling equivalent to that in control untreated islets. However, unexpectedly, we did not observe substantial recoveries in the presence and action of functional sub-populations when gap junctions were recovered: we observed only partial recovery to ChR2-recruitment, and no recovery to first responders or β-cell hubs. One simple explanation is non-specific effects of the S293 peptide, however prior studies did not observe non-specific effects on [Ca^{2+}]. These findings indicate that the disruptions to functional sub-populations under pro-inflammatory cytokines are not primarily due to disruptions to gap junction coupling.

While initially unexpected, our findings are consistent with the fact that gap junction coupling is not per se a defining feature of these sub-populations. For example, cells with high ChR2-recruitment do not show increased gap junction coupling [27], β-cell hubs do not show increased gap junction coupling [30], and first responders show lower gap junction coupling[28]. Thus, gap junction coupling does not drive the formation of these sub-populations, rather they are defined by cell-intrinsic properties such as elevated glucose metabolism or electrical differences, as well as other environmental properties such as communication with α- or δ-cells [35]. Dysfunction to these cell intrinsic properties or altered communication with α- or δ-cells is therefore likely driving their disruption. However, we note that some amount of gap junction coupling will still be required for these cells to exert their influence across the islet [3].

Enhancing gap junction coupling can recover islet function in diabetogenic conditions
[12,16,36]. This has been demonstrated via both genetic and pharmacological approaches in
models of diabetes invitro and invivo. While promising, our findings indicate that targeting gap
junction coupling alone will be insufficient to fully recover islet function, given that functional β-
cell sub-populations will still be disrupted. For example, caloric restriction is able to improve
islet dysfunction that results from a high-fat diet; yet while gap junction coupling recovered,
wave initiators / leaders did not [15]. Inclusion of the GLP1R agonist Exendin-4 with pro-
inflammatory cytokines recovered gap junction coupling and coordinated $[Ca^{2+}]$ dynamics [37],
while also impacting insulin secretion pathways via cAMP signaling. Thus, ensuring intrinsic
factors and/or communication with α- and δ-cells are recovered will likely be required. Future
work should examine which of these factors is important to both define functional sub-
populations and preserve their action on islet function under diabetogenic conditions.

Limitations and Future Work: Our findings indicate that protection of gap junctions alone is not
sufficient to recover cytokine-mediated loss of functional subpopulations. However other
factors within the islet microenvironment during diabetes can also impact islet function,
including gluco- and lipo-toxicity, amyloid plaque formation, and other pro-inflammatory
cytokines such as IL6. Thus, extending our findings to other invitro treatments and mouse
models of diabetes will be needed in future work. Another important factor is that our study
examined a 24 hours treatment with pro-inflammatory cytokines and/or S293 peptide mimetic.
This compares to a 1-6 hour treatment time in previous studies that support the recovery of
gap junctions and islet function by S293 [12]. While we observed a recovery of gap junction
coupling, it is possible that under more acute treatments we will observe some recovery in
functional sub-populations and a further enhancement in gap junction coupling: Cx36 turnover
has a half-life of ~3 hours [38]. Nevertheless a longer term treatment is likely more reflective of
the islet microenvironment in diabetes.

Summary: Overall, in thus study we demonstrate that proinflammatory cytokines disrupt the
presence and action of functional β-cell sub-populations such as hubs, first responders and
cell-recruitment of $[Ca^{2+}]$. Preventing the decline in gap junction coupling was not sufficient for
recovering these functional β-cell sub-populations. This correlates with gap junction coupling
not being a factor that defines β-cell heterogeneity within the islet, and rather cell intrinsic

properties or other aspects of intra-islet communication instead being disrupted by the pro-

inflammatory environment.

Figure Descriptions

Figure 1. Restoration cytokine-mediated disruptions to gap junction coupling, as measured by FRAP. (A) Isolated mouse islets were treated with a 0.1x cocktail of proinflammatory cytokines and co-incubated with 30 μM S293 peptide for 1 hour before imaging. (B) Representative image of Rhodamine123 loaded islet pre- and post-photobleaching, along with Fluorescence Recovery After Photobleaching (FRAP) time-course after photobleaching, showing fluorescence recovery in the bleached region. (C) Rate of recovery was calculated as a proxy for gap junction permeability in untreated, cytokine, cytokine plus S293 peptide, and S293 peptide alone treatments. Data represents Mean $\pm$ S.E.M. with each islet represented by a data point (from 6 mice), and each measurement is normalized to the mean control value per mouse. * indicates $p < 0.05$.

Figure 2. Impact of gap junction restoration on the cytokine-mediated disruption to ChR2-mediated recruitment of $[Ca^{2+}]$. (A) A ChR2-expressing islet (left), where a single ChR2-YFP expressing cell was stimulated at 1 Hz (i) and calcium response was recorded across the islet which exhibited both regions of recruited (ii) and unrecruited (iii) cells in untreated conditions (middle). The result $[Ca^{2+}]$ activity was then quantified (right). (B) As in A, under cytokine-treated conditions. (C) Histogram indicating the proportion of the islet recruited by a single cell stimulation under each treatment condition. (D) Mean proportion of the islet recruited by a single cell stimulation under each treatment condition. Data represents Mean $\pm$ S.E.M. with each islet represented by a data point (from 15 mice). (E) The proportion of cells that were able to recruit an area larger than 30% of the islet under the cytokine and S293 conditions indicated. ** indicates $p < 0.01$, **** indicates $p < 0.0001$.

Figure 3. Impact of gap junction restoration on the cytokine-mediated disruption to the temporal and spatial consistency of first responders. (A) Time-course of $[Ca^{2+}]$ in first responder cells (red) and last responder cells (green) following stimulation by high (11mM) glucose, under control or cytokine-treated conditions (for 1h or 24h). (B) The mean time spread between the first and last responders (ΔT) in islets treated with cytokines for 24 hours or 1 hour, compared to untreated. (C) The mean time spread in islets treated for 24 hours, with cytokines and/or S293 peptide recovery of gap junction coupling. (D) Time-course of $[Ca^{2+}]$ in first responder cells (red) and last responder cells (green), subjected to repeated stimulation from low to high

glucose (E) The percent consistency of first responders in untreated cells (~40%) or those with cytokines and/or S293 peptide recovery of gap junction coupling. ** indicates p < 0.01, *** indicates p < 0.001.

Figure 4. Impact of gap junction restoration on the cytokine-mediated disruption of highly connected β-cell hubs. Representative connectivity map (above) and time-courses (below) of islets under control or cytokine-treated conditions (for 1h or 24h). (B) The mean correlation coefficient in islets treated with cytokines for 24 hours or 1 hour, compared to untreated. (C) Mean correlation coefficient in islets treated for 24 hours, with cytokines and/or S293 peptide recovery of gap junction coupling. (D) As in C for the average number of links between cells. (E) As in C for the proportion of hub cells among the entire islet cell population. (F) Time-course of [Ca^{2+}] in β-cell hubs that are consistent or not consistent, together with representative connectivity maps. (G) The percent consistency of hubs in untreated cells (>50%) or those with cytokines and/or S293 peptide recovery of gap junction coupling. * indicates p < 0.05, ** indicates p < 0.01, *** indicates p < 0.001.

Supplement 1. First responder cells under 1h cytokine treatment with/without gap junction restoration. (A) The mean time spread in islets treated for 1 hour, with cytokines and/or S293 peptide recovery of gap junction coupling. (B) The percent consistency of first responders in untreated cells (~40%) or those with cytokines and/or S293 peptide recovery of gap junction coupling.

Supplement 2. Highly connected β-cell hubs under 1h cytokine treatment with/without gap junction restoration. (A) Mean correlation coefficient in islets treated for 1 hour, with cytokines and/or S293 peptide recovery of gap junction coupling. (B) As in A for the average number of links between cells. (C) As in A for the proportion of hub cells among the entire islet cell population. (D) As in A for percent consistency of hubs.

References

1.	Federation, I.D. *Diabetes around the world in 2021*. 2022 [cited 2024.

2.	Allagnat, F., D. Martin, D.F. Condorelli, G. Waeber, and J.-A. Haefliger, *Glucose represses connexin36 in insulin-secreting cells.* Journal of cell science, 2005. **118**(22): p. 5335-5344.

3.	Benninger, R.K., M. Zhang, W.S. Head, L.S. Satin, and D.W. Piston, *Gap junction coupling and calcium waves in the pancreatic islet.* Biophysical journal, 2008. **95**(11): p. 5048-5061.

4.	Farnsworth, N.L., A. Hemmati, M. Pozzoli, and R.K. Benninger, *Fluorescence recovery after photobleaching reveals regulation and distribution of connexin36 gap junction coupling within mouse islets of Langerhans.* The Journal of physiology, 2014. **592**(20): p. 4431-4446.

5.	Ravier, M.A., M. Guldenagel, A. Charollais, A. Gjinovci, D. Caille, G. Sohl, C.B. Wollheim, K. Willecke, J.-C. Henquin, and P. Meda, *Loss of connexin36 channels alters β-cell coupling, islet synchronization of glucose-induced Ca2+ and insulin oscillations, and basal insulin release.* Diabetes, 2005. **54**(6): p. 1798-1807.

6.	Pipeleers, D.G., *Heterogeneity in pancreatic β-cell population.* Diabetes, 1992. **41**(7): p. 777-781.

7.	Stožer, A., M. Gosak, J. Dolenšek, M. Perc, M. Marhl, M.S. Rupnik, and D. Korošak, *Functional connectivity in islets of Langerhans from mouse pancreas tissue slices.* PLoS computational biology, 2013. **9**(2): p. e1002923.

8.	Benninger, R.K., W.S. Head, M. Zhang, L.S. Satin, and D.W. Piston, *Gap junctions and other mechanisms of cell–cell communication regulate basal insulin secretion in the pancreatic islet.* The Journal of physiology, 2011. **589**(22): p. 5453-5466.

9.	Speier, S., A. Gjinovci, A. Charollais, P. Meda, and M. Rupnik, *Cx36-mediated coupling reduces β-cell heterogeneity, confines the stimulating glucose concentration range, and affects insulin release kinetics.* Diabetes, 2007. **56**(4): p. 1078-1086.

10.	Head, W.S., M.L. Orseth, C.S. Nunemaker, L.S. Satin, D.W. Piston, and R.K. Benninger, *Connexin-36 gap junctions regulate in vivo first-and second-phase insulin secretion dynamics and glucose tolerance in the conscious mouse.* Diabetes, 2012. **61**(7): p. 1700-1707.

11.	Hodson, D.J., R.K. Mitchell, E.A. Bellomo, G. Sun, L. Vinet, P. Meda, D. Li, W.-H. Li, M. Bugliani, and P. Marchetti, *Lipotoxicity disrupts incretin-regulated human β cell connectivity.* The Journal of clinical investigation, 2013. **123**(10): p. 4182-4194.

12.	St. Clair, J.R., M.J. Westacott, J. Miranda, N.L. Farnsworth, V. Kravets, W.E. Schleicher, J.M. Dwulet, C.H. Levitt, A. Heintz, and N.W. Ludin, *Restoring connexin-36 function in diabetogenic environments precludes mouse and human islet dysfunction.* The Journal of Physiology, 2023. **601**(18): p. 4053-4072.

13.	Farnsworth, N.L., R.L. Walter, A. Hemmati, M.J. Westacott, and R.K. Benninger, *Low level pro-inflammatory cytokines decrease connexin36 gap junction coupling in mouse and human islets through nitric oxide-mediated protein kinase Cδ.* Journal of Biological Chemistry, 2016. **291**(7): p. 3184-3196.

14.	Carvalho, C.d.F., R.B. Oliveira, A. Britan, J. Santos-Silva, A.C. Boschero, P. Meda, and C.B. Collares-Buzato, *Impaired β-cell-β-cell coupling mediated by Cx36 gap junctions in prediabetic mice.* American Journal of Physiology-Endocrinology and Metabolism, 2012. **303**(1): p. E144-E151.

15.	Corezola do Amaral, M.E., V. Kravets, J.M. Dwulet, N.L. Farnsworth, R. Piscopio, W.E. Schleicher, J.G. Miranda, and R.K. Benninger, *Caloric restriction recovers impaired β-*

cell-β-cell gap junction coupling, calcium oscillation coordination, and insulin secretion
in prediabetic mice. American Journal of Physiology-Endocrinology and Metabolism,
2020. **319**(4): p. E709-E720.
16.   Farnsworth, N.L., R.A. Piscopio, and R.K. Benninger, *Modulation of gap junction*
coupling within the islet of langerhans during the development of type 1 diabetes.
Frontiers in Physiology, 2022. **13**: p. 913611.
17.   Westacott, M.J., N.L. Farnsworth, J.R. St. Clair, G. Poffenberger, A. Heintz, N.W. Ludin,
N.J. Hart, A.C. Powers, and R.K. Benninger, *Age-dependent decline in the coordinated*
[Ca2+] and insulin secretory dynamics in human pancreatic islets. Diabetes, 2017.
66(9): p. 2436-2445.
18.   Ravier, M., J. Sehlin, and J. Henquin, *Disorganization of cytoplasmic Ca 2+ oscillations*
and pulsatile insulin secretion in islets from ob/ob mice. Diabetologia, 2002. **45**: p. 1154-
1163.
19.   Heimberg, H., A. De Vos, A. Vandercammen, E. Van Schaftingen, D. Pipeleers, and F.
Schuit, *Heterogeneity in glucose sensitivity among pancreatic beta-cells is correlated to*
differences in glucose phosphorylation rather than glucose transport. The EMBO
journal, 1993. **12**(7): p. 2873-2879.
20.   Roscioni, S.S., A. Migliorini, M. Gegg, and H. Lickert, *Impact of islet architecture on β-*
cell heterogeneity, plasticity and function. Nature Reviews Endocrinology, 2016. **12**(12):
p. 695-709.
21.   Wang, Y.J., J. Schug, K.-J. Won, C. Liu, A. Naji, D. Avrahami, M.L. Golson, and K.H.
Kaestner, *Single-cell transcriptomics of the human endocrine pancreas.* Diabetes, 2016.
65(10): p. 3028-3038.
22.   Wojtusciszyn, A., M. Armanet, P. Morel, T. Berney, and D. Bosco, *Insulin secretion from*
human beta cells is heterogeneous and dependent on cell-to-cell contacts. Diabetologia,
2008. **51**: p. 1843-1852.
23.   Benninger, R.K., T. Hutchens, W.S. Head, M.J. McCaughey, M. Zhang, S.J. Le
Marchand, L.S. Satin, and D.W. Piston, *Intrinsic islet heterogeneity and gap junction*
coupling determine spatiotemporal Ca2+ wave dynamics. Biophysical journal, 2014.
107(11): p. 2723-2733.
24.   Giordano, E., D. Bosco, V. Cirulli, and P. Meda, *Repeated glucose stimulation reveals*
distinct and lasting secretion patterns of individual rat pancreatic B cells. The Journal of
clinical investigation, 1991. **87**(6): p. 2178-2185.
25.   Benninger, R.K. and V. Kravets, *The physiological role of β-cell heterogeneity in*
pancreatic islet function. Nature Reviews Endocrinology, 2022. **18**(1): p. 9-22.
26.   Rutter, G.A., A. Gresch, L. Delgadillo Silva, and R.K. Benninger, *Exploring pancreatic*
beta-cell subgroups and their connectivity. Nature metabolism, 2024: p. 1-15.
27.   Westacott, M.J., N.W. Ludin, and R.K. Benninger, *Spatially organized β-cell*
subpopulations control electrical dynamics across islets of Langerhans. Biophysical
journal, 2017. **113**(5): p. 1093-1108.
28.   Kravets, V., J.M. Dwulet, W.E. Schleicher, D.J. Hodson, A.M. Davis, L. Pyle, R.A.
Piscopio, M. Sticco-Ivins, and R.K. Benninger, *Functional architecture of pancreatic*
islets identifies a population of first responder cells that drive the first-phase calcium
response. PLoS Biology, 2022. **20**(9): p. e3001761.
29.   Johnston, N.R., R.K. Mitchell, E. Haythorne, M.P. Pessoa, F. Semplici, J. Ferrer, L.
Piemonti, P. Marchetti, M. Bugliani, and D. Bosco, *Beta cell hubs dictate pancreatic islet*
responses to glucose. Cell metabolism, 2016. **24**(3): p. 389-401.

30. Briggs, J.K., A. Gresch, I. Marinelli, J.M. Dwulet, D.J. Albers, V. Kravets, and R.K.
Benninger, *β-cell intrinsic dynamics rather than gap junction structure dictates*
subpopulations in the islet functional network. Elife, 2023. **12**: p. e83147.
31. Farnsworth, N.L., R.A. Piscopio, W.E. Schleicher, D.G. Ramirez, J.G. Miranda, and R.K.
Benninger, *Modulation of gap junction coupling within the islet of langerhans during the*
development of type 1 diabetes. Frontiers in Physiology, 2022. **13**: p. 913611.
32. Akash, M.S.H., K. Rehman, and S. Chen, *Role of inflammatory mechanisms in*
pathogenesis of type 2 diabetes mellitus. Journal of cellular biochemistry, 2013. **114**(3):
p. 525-531.
33. Alexandraki, K., C. Piperi, C. Kalofoutis, J. Singh, A. Alaveras, and A. Kalofoutis,
Inflammatory process in type 2 diabetes: The role of cytokines. Annals of the New York
Academy of Sciences, 2006. **1084**(1): p. 89-117.
34. Gosak, M., R. Yan-Do, H. Lin, P.E. MacDonald, and A. Stožer, *Ca2+ oscillations, waves,*
and networks in islets from human donors with and without type 2 diabetes. Diabetes,
2022. **71**(12): p. 2584-2596.
35. Jin, E., J.K. Briggs, R.K. Benninger, and M.J. Merrins, *Glucokinase activity controls*
subpopulations of β-cells that alternately lead islet Ca2+ oscillations. bioRxiv, 2024.
36. Klee, P., F. Allagnat, H. Pontes, M. Cederroth, A. Charollais, D. Caille, A. Britan, J.-A.
Haefliger, and P. Meda, *Connexins protect mouse pancreatic β cells against apoptosis.*
The Journal of clinical investigation, 2011. **121**(12): p. 4870-4879.
37. Farnsworth, N.L., R. Walter, R.A. Piscopio, W.E. Schleicher, and R.K. Benninger,
Exendin‐4 overcomes cytokine‐induced decreases in gap junction coupling via protein
kinase A and Epac2 in mouse and human islets. The Journal of physiology, 2019. **597**(2):
p. 431-447.
38. Wang, H.Y., Y.-P. Lin, C.K. Mitchell, S. Ram, and J. O'Brien, *Two-color fluorescent*
analysis of connexin 36 turnover: relationship to functional plasticity. Journal of cell
science, 2015. **128**(21): p. 3888-3897.
610

Figure 1

Figure 2

Figure 3

Figure 4

Supplement 1

Supplement 2