Diminished gap junction coupling under diabetogenic conditions does not drive loss of functional β-cell sub-populations

Claire H Levitt[a], Dominic Isaacs[a], Maria Hansen[b], Vira Kravets[a], Jennifer K Briggs[a], Richard KP Benninger[a,b,1]

[a] Department of Bioengineering, University of Colorado, Anschutz Medical campus, Aurora, CO. USA

[b] Barbara Davis center for childhood diabetes, University of Colorado, Anschutz Medical campus, Aurora, CO. USA

[1] To whom correspondence should be addressed: richard.benninger@cuanschutz.edu

1775 Aurora court, M20-4306D, mailstop B140, University of Colorado Anschutz Medical campus, Aurora, CO. 80045.

Running Title: Effect of gap junction coupling under diabetogenic conditions

Key Words: Optogenetics, Islet Dynamics; Optical Microscopy; Diabetes; Gap junctions

Classification: Systems Biology

Manuscript word count: 5560

Abstract word count: 282

Number of figures: 5

Abstract

Within the islets of Langerhans, gap junction coupling is important for synchronizing oscillatory

free-calcium activity ($[Ca^{2+}]$) and regulating pulsatile insulin release. In islets from multiple

models of diabetes, gap junction coupling is disrupted, and $[Ca^{2+}]$ synchronization and pulsatile

insulin is lost. Functional sub-populations have been identified within the islet that are linked to

driving synchronized $[Ca^{2+}]$ and insulin release. These sub-populations can be disrupted under

conditions associated with diabetes such as glucolipotoxicity and inflammatory environments,

and their loss may drive islet dysfunction. Here we investigated how loss of gap junction

coupling influences functional subpopulations under diabetogenic environments. We treated

islets with a cocktail of pro-inflammatory cytokines, and protected gap junction coupling via co-

treatment with a Cx36 peptide S293 that was previously shown to specifically prevent a decline

in gap junction permeability and synchronized $[Ca^{2+}]$ dynamics. We performed calcium imaging

and ChR2 stimulation, analyzed islet $[Ca^{2+}]$ dynamics and the presence of functional sub-

populations including hubs and first-responders. 1h or 24h cytokine-treatment disrupted gap

junction coupling, which was fully prevented by S293 peptide co-treatment. Treatment with pro-

inflammatory cytokines decreased the recruitment of $[Ca^{2+}]$ upon ChR2 stimulation, increased

the time between first and last responding cells upon glucose stimulation, and reduced the

number and consistency of hub cells. When preserving gap junction coupling by S293 during

cytokine treatment, the presence and consistency of these sub-populations was only marginally

improved. We therefore concluded that while gap junction coupling is important for functional

sub-populations to exert their influence on islet function, restoration of gap junctions alone is

not sufficient to recover functional sub-populations upon diabetogenic conditions. Thus,

preventing a disruption to intrinsic β-cell properties that define functional subpopulations is

likely important for preserving these sub-populations during diabetes.

Introduction

Diabetes Mellitus (DM) affects approximately 537 million people globally [1,2], and is characterized by the dysfunction and/or death of insulin-secreting β-cells in the islets of Langerhans in the pancreas. β-cells in the islet secrete insulin in response to elevations in blood glucose or other nutrients. At elevated glucose, β-cells show increased ATP/ADP following the metabolism of glucose, which drives membrane depolarization via closure of ATP-sensitive K+ (K_{ATP}) channels and activation of voltage gated calcium (Ca_V) channels. Subsequent oscillations of intracellular free-calcium ($[Ca^{2+}]$) across the islet drive the release of insulin in discrete pulses.

The dynamics of $[Ca^{2+}]$ are synchronized across β-cells in the islet, which is facilitated by connexin-36 (Cx36) gap junctions [3-5]. Gap junctions synchronize the β-cell response to glucose despite heterogeneity in β-cell metabolism and excitability [6,7]. Under low glucose conditions, gap junction-mediated hyper-polarizing currents suppress spontaneous elevations in $[Ca^{2+}]$ from excitable β-cells [8,9]. At high glucose, gap junction-mediated depolarizing currents coordinate and enhance both first-phase $[Ca^{2+}]$ and oscillatory second phase $[Ca^{2+}]$ responses [3]. In Cx36 knockout models, synchronous $[Ca^{2+}]$ responses are disrupted [3,5,9], both first phase insulin secretion and second phase insulin pulses are diminished and glucose tolerance is impaired [5,10]. Importantly, islet gap junction coupling is disrupted under conditions associated with diabetes including by glucolipotoxicity [2,11,12] and pro-inflammatory cytokines [13] as well as in mouse models of diabetes [14-16] and human [11,12,17]. Synchronous $[Ca^{2+}]$ activity is also disrupted under conditions associated with diabetes [11,13,17,18], likely as a result of disrupted Cx36 gap junction coupling.

β-cells are heterogeneous in gene expression [19-21], glucose sensitivity [6,19], insulin secretion [22], and cell-to-cell communication [23]. As a result, β-cells exhibit functional heterogeneity, and display distinct insulin secretory patterns. This functional heterogeneity is evidenced in previous studies that found ~75% of β-cells exhibited a fixed response to glucose, whereas ~25% fluctuated between states of responsive and unresponsiveness [24]. The characterization of β-cell heterogeneity has led to the identification of functional subpopulations which have been shown to play distinct roles in governing the spatiotemporal $[Ca^{2+}]$ and insulin secretory dynamics of the islet [25,26]. For example, following optogenetic Channelrhodopsin-2 (ChR2) stimulation, a population of β-cells can recruit and elevate $[Ca^{2+}]$ in a large number of neighboring β-cells within the islet [27]. Similarly, 'first-responder cells', that are first to show elevated $[Ca^{2+}]$ upon elevated glucose stimulation, can recruit and coordinate less excitable

cells to respond earlier [28]. This first-responder subpopulation is distinct from previously
identified highly-connected 'hub cells' that exhibit a high degree of co-activity with neighboring
β-cells [29] and have been shown to play a critical role in driving the synchronized oscillatory
dynamics. Importantly, upon both glucolipotoxic environments and pro-inflammatory cytokines,
the proportion of 'hub cells' declines [29]. This finding suggests that a selective loss of functional
sub-populations may be sufficient to drive islet dysfunction in diabetes. However, the
mechanisms underlying this loss of cells are unclear.

In this study, we sought to understand how functional sub-populations of cells are
disrupted under conditions associated with diabetes. Given that gap junction coupling
influences overall islet coordination, we specifically asked whether the loss of gap junction
coupling that is associated with diabetes is a primary cause for the loss of these sub-
populations and their influence on islet function. We first tested how functional sub-populations
are influenced by pro-inflammatory cytokine treatment. We then tested a role of gap junction
coupling, by using a Cx36 peptide [12] that was previously shown to specifically protect gap
junction permeability and retain glucose-stimulated [Ca^{2+}] dynamics under conditions
associated with diabetes.

Materials and Methods

Mice/Animal Care: Islets expressing ChR2-YFP in β-cells were obtained by crossing Pdx^{PB}-Cre (JAX #014647) and ROSA26-lox-STOP-lox-ChR2-YFP mice (JAX #024109). Islets expressing GCaMP6s in β-cells were obtained by crossing MIP-Cre^{ER} mice (JAX #024709) and ROSA26-lox-STOP-lox-GCaMP6s mice (JAX #028866). Mice were genotyped via qPCR (Transnetyx). All animal experiments were approved by the University of Colorado Institutional Animal Care and Use Committee (IACUC Protocol number: 000024). Mice were housed in a temperature-controlled environment with a 12 hr light/dark cycle and continuous access to both food and water. For mice with β-cell specific GCaMP6s expression, tamoxifen (solubilized in corn oil) was administered via IP injection for 5 consecutive days (50 mg/kg of body weight) to facilitate CreER-mediated recombination. All experiments use both male and female mice.

Islet Isolation and Culture: Islets were isolated from mice under ketamine/xylazine anesthesia and euthanized via exsanguination. Collagenase (Type V, Sigma) was delivered to the pancreases at a concentration of 2.5 mg/mL via the bile duct. After inflation, the pancreas was surgically removed and digested (5 min at 37 °C). Islets were then hand-picked and cultured in RPMI medium (Gibco) containing 10% fetal bovine serum, 100 U/mL penicillin, 100 ug/mL streptomycin, and 11 mM glucose. Islets were incubated at 37°C, 5% CO2.

Islet Treatment: Islets were treated overnight with a cocktail of proinflammatory cytokines (1 ng/mL TNF-α, 5 ng/mL IL-1β, and 10 ng/mL IFN-γ), which is equivalent to a '0.1x' cocktail as previously used [13]. To protect gap junction coupling, islets were co-treated with 30 μM peptide S293 for 1-6 hours for experiments using optogenetic stimulation, and 24 hours for experiments measuring gap junction permeability (FRAP) and calcium activity ([Ca^{2+}]).

Fluorescent Recovery After Photobleaching and Analysis Gap junction permeability was assessed by Fluorescence Recovery After Photobleaching (FRAP) [4]. Isolated islets were immobilized onto glass bottom dishes (MatTek) using cell tissue adhesive (CellTak, Corning) and incubated at 37°C, 5% CO2 overnight. Islets were then loaded with 12.5 mM cationic dye Rhodamine-123 (Rh123) for 30 minutes at 37°C in Hank's Balanced Salt Solution (HBSS) at 5mM glucose. Islets were washed and imaged at room temperature in HBSS using a Zeiss LSM 800 confocal microscope with a x40 1.2NA water immersion objective. Rh123 was excited

at 488 nm. Half the islet area was selected using a rectangular ROI and photobleached for 25
frames at 40% laser power, achieving ~50% bleaching on average. Images were collected
every 5 seconds for 6 to 10 minutes.

Image analysis of Rh123 dye diffusion after photobleaching was conducted using
MATLAB and Image J. Fluorescent recovery was measured within a region of interest
encompassing the entire photobleached area and individual cells. The initial fluorescence
intensity was considered the point directly after photobleaching (I_P). Fluorescence recovery
was the final fluorescence (I_F) once the diffusion of dye reached a plateau. The recovery was
calculated as a percentage, relative to the initial intensity before photobleaching (I_0) as follows:

142
$$\% \, Recovery = 100 * \frac{I_F - I_P}{I_0 - I_P}$$

143   The rate of fluorescent recovery was calculated based on the linearized exponential fluorescent
144   recovery curve of the bleached area over time (I_i) normalized to the total recovery. Calculations
145   were conducted as follows:

148
$$Rate \, of \, recovery = -Ln(1 - \frac{I_i - I_P}{I_0 - I_P})$$

While Rhodamine 123 also loads the mitochondria, dependent on mitochondrial membrane
potential, this loading does not influence the measurements of gap junction permeability [12].

Channelrhodopsin-2 Single Cell Entrainment and Analysis: ChR2-YFP expressing islets were
incubated with pro-inflammatory cytokines, with or without S293 peptide, or S293 peptide alone
for 6 hours after isolation in RPMI medium (Gibco) containing 10% fetal bovine serum, 100
U/mL penicillin, 100 ug/mL streptomycin, and 11 mM glucose at 37°C, 5% CO2 prior to study.

Islets were loaded with 2μM of calcium indicator Rhodamine-2 (Rhod-2) in BMHH
imaging solution (125 mM NaCl, 5.7 mM KCl, 2.5 mM CaCl2, 1.2 mM MgCl2, 10 mM HEPES,
2 mM glucose, and 0.1% BSA, pH 7.4) for 35 minutes at 37°C. Islets were then washed and
imaged on a glass bottom dish (MatTek) in a 5mM glucose imaging solution using a LSM 800
confocal microscope kept at 37°C .

ChR2-YFP was excited at 488 nm. A rectangular ROI was drawn within the bounds of a
single cell to avoid stimulating neighboring cells that express ChR2-YFP. The cell was activated
using a photobleaching macro (intensity 5% laser power) for 8 stimulations at a frequency of
0.1 Hz. Rhod-2 was simultaneously excited at 560 nm and the corresponding Ca^{2+} response

across the whole islet was measured based on fluorescent intensity of Rhod-2 that was
coincident to ChR2 stimulation.

MATLAB was used to analyze ChR2 recruitment data, as previously described [27], as
well as entrainment. Islets were selected by hand drawing a region of interest around the islet.
Background was identified as regions that did not show any calcium response. Activity was
based on the fluorescent intensity of the Ca^{2+} indicator Rhod-2 upon ChR2 stimulation and
assessed using individual pixel time-courses. A pixel was considered 'active' if its intensity was
twice that of the background. A threshold of 0.15 was used to determine which regions of the
islet were activate due to ChR2 stimulation compared to the islet average. The percent of the
islet area that was 'active' was calculated as a proportion of the entire islet area and
subsequently referred to as "Area Active". At elevated glucose, the mean Ca^{2+} time trace was
averaged across the entire islet and segmented into pre and during stimulation. After
detrending, a Fast Fourier Transform (FFT) was performed for both segments and the DC
component subtracted. The natural oscillation frequency was identified from the pre stimulation
segment as the frequency with peak power. The driving frequency was confirmed at ~0.1Hz as
the frequency with peak power during stimulation.

GCaMP6 Imaging of Calcium Dynamics and Analysis: Isolated islets expressing β-cell specific
GCaMP6 were incubated overnight with proinflammatory cytokines with or without S293
peptide, or S293 peptide alone. 30 Minutes prior to imaging, islets were fasted at 2 mM glucose
in BMHH imaging solution. Islets were then transferred to a glass bottom dish and imaged on
an LSM 800 confocal microscope. GCaMP6 fluorescence was excited at 488 nm with images
collected at 1 frame/sec for approximately 80 minutes. 3-5 Minutes after the start of imaging,
the glucose concentration was raised from 2 mM to 11 mM. For experiments measuring
consistency between first responders and hub cells, glucose was subsequently lowered to 2
mM after achieving second phase oscillations for 10-15 minutes. After the islet had returned
to basal activity without oscillations (~15 minutes), the glucose concentration was increased
again to 11 mM. Samples were imaged at either x20 air or x40 water immersion and held at
37°C.

Calcium dynamics were analyzed using MATLAB. First phase response time,
identification of first and last responders, and characterization of highly connected hub cells
was determined as previously described [28]. Briefly, to identify the response time, we

determined the time at which individual β-cells reached their half maximum amplitude following

stimulation from low to high glucose. This response time was rank ordered for all cells in the

islet. The earliest 10% were classed as first responders and the latest 10% classed as last

responders, as previously defined [29]. ΔT was calculated as the difference in time between

the means of the last responders and first responders to capture the changes in spread

between first and last responders between treatment groups relative to total time of the first

phase. The percent consistency was determined as a proportion of top 10% of responding cells

of the initial stimulation that were also first responders or nearest neighbors to first responders

during repeated stimulation.

Highly correlated β-cell cells were identified using the calcium time-course at high

glucose during the second phase. A highly correlated β-cell is a β-cell whose calcium activity

strongly parallels many other β-cell in the islet. To quantify the degree of correlation, the degree

of cross-correlation value obtained was obtained using the xcorr() function in MATLAB across

all measured β-cell calcium time traces was compared among each cell. "Links" were derived

using a threshold of correlation above 0.9. Hub cells have previously been defined as those

possessing the "majority of connections" [29-30] from which we derived the criteria of >60% of

functional "links" determined by mean degree of correlation above the defined threshold relative

to all cells measured in the islet**.**

Statistical analysis: All statistical analysis was performed in Prism (GraphPad). A 2-sided t-test

was used when comparing parameters of 2 treatment groups such as between Control versus

Cytokine treated islets. Ordinary one-way ANOVA with Tukey's *post hoc* analysis was used to

compare across multiple treatment groups, where * indicates $p < 0.05$, ** indicates $p < 0.01$,

*** indicates $p < 0.001$, **** indicates $p < 0.0001$. In all cases, error bars represent the standard

error of the mean (S.E.M.).

Results

Successful protection of gap junction coupling by Cx36 S293 peptide under pro-inflammatory
cytokines: To test whether functional sub-populations are disrupted by a cocktail of pro-
inflammatory cytokines (0.5 ng/mL IL1-β, 1 ng/mL TNF-α, and 10 ng/mL IFN-γ), and whether
sub-population loss depends on loss of gap junction coupling, we followed an approach
outlined in Figure 1A. Fluorescence Recovery After Photobleaching (FRAP) was used to
assess gap junction permeability and confirm a disruption to gap junction coupling by pro-
inflammatory cytokines (Figure 1.B) and protection of gap junction function by the Cx36 S293
peptide, as previously described [4]. Islets treated with pro-inflammatory cytokines showed a
significant reduction in gap junction permeability, as indicated by a slower recovery rate (Figure
1C, p=0.0300). During co-treatment with the S293 peptide, islets showed a 92.2%±38.6%
increase in the recovery rate compared to cytokine treatment alone. No significant difference
was observed between untreated control and co-treatment groups (p=0.958) or untreated
control vs S293 peptide alone (p=0.918). Thus, co-treatment with S293 peptide fully protects
gap junction permeability, consistent with previous studies that use the S293 peptide to protect
or recover gap junction coupling in diabetogenic environments [12].

Protecting gap junction coupling provides a partial recovery to the cytokine-mediated disruption
of ChR2-mediated recruitment of [Ca^{2+}] at low glucose: With the conditions established to
protect gap junction coupling under a diabetogenic environment, we asked whether functional
subpopulations of β-cells are also protected. Previous studies applied the optogenetic construct
ChannelRhodopsin-2 (ChR2) and single cell stimulation to discover a subpopulation of β-cells
with elevated metabolic activity that can recruit large numbers of neighboring cells when
activated at low (5 mM) glucose levels [27]. We therefore examined how pro-inflammatory
cytokines influenced these sub-populations and then tested the effects of preventing a decline
in gap junction coupling on these sub-populations.

Islets from Pdx1^{PB}-Cre;Rosa-LSL-ChR2-YFP mice that express ChR2 in β-cells were
loaded with the Ca^{2+} dye Rhod2. ChR2 expressing cells were subjected to focal stimulation in
single β-cells and the spatial Ca^{2+} response was observed in neighboring cells (Figure 2.A).
We quantified the proportion of β-cells in the imaging plane that showed Ca^{2+} elevations
coincident with ChR2 stimulation, which was repeated in ChR2-expressing islets treated with
the cocktail of pro-inflammatory cytokines for 1 hour (Figure 2.B) and/or Cx36 S293 peptide.

Under all treatments we observed heterogeneity in the recruitment of elevated [Ca^{2+}]

following single cell stimulation (Figure 2.C). In untreated control islets, 41.3%±4.7% of cells

within the imaging plane showed elevated [Ca^{2+}] following single cell stimulation, with a

substantial variation (Figure 2.C,D) as previously observed [27]. Treatment with pro-

inflammatory cytokines led to a significant decrease in the proportion of cells showing elevated

[Ca^{2+}] following single cell stimulation (15.4%±2.3%) compared to untreated controls (Figure

2.C,D, p<0.0001,). Co-treatment with the S293 peptide led to a moderate increase in elevated

[Ca^{2+}] following single cell stimulation (24.8%±2.6%), however this was still significantly lower

than in untreated control (Figure 2.C,D, p=0.0026). Treatment with S293 peptide alone showed

a similar proportion of cells with elevated [Ca^{2+}] (35.0%±4.5 %) to untreated control islets

(Figure 2.C,D, p=0.6680).

Untreated control islets showed broad heterogeneity in the recruitment of [Ca^{2+}] (Figure

2.C), as previously observed [27]. This included a population of cells that recruited elevated

[Ca^{2+}] across the islet that was above the median value in untreated control islet (>40%, Figure

2.C), consistent with earlier findings [27]. Under cytokine treatment (1 hour), this 'high-

recruiting' population of cells was almost completely absent (~4%, Figure 2.E). Co-treatment

with S293 peptide partially recovered this population of cells (~10%, Figure 2.E), but again less

than in either untreated controls or S293 peptide treatment alone. The trend in these findings

were similar when we applied other thresholds for the islet area recruited upon single-cell

stimulation (Supplemental 3.A).

Therefore, treatment with pro-inflammatory cytokines reduced the ChR2-mediated

recruitment of cells to increase [Ca^{2+}], with an abolishment of 'high-recruiting' cells. Protecting

gap junction coupling only partially recovers ChR2-mediatred recruitment of cells to increase

[Ca^{2+}], with only a small number of 'high-recruiting' cells low glucose.

Protecting gap junction coupling recovers cytokine-mediated disruption of ChR2-mediated

recruitment of [Ca^{2+}] at high glucose: Although protecting gap junction coupling did not

significantly improve single-cell mediated islet recruitment at low glucose, we sought to assess

whether the action of ChR2 was influenced by cytokine-mediated disruption at high glucose.

Islets generally respond to changes from low to high glucose by displaying a robust first phase

and sustained second phase within a 30-minute period. We therefore exposed islets to a high

glucose environment for 30 minutes during which intrinsic islet [Ca^{2+}] oscillations were

observed, confirming that the islets had entered the second phase. We then performed single-

cell stimulation of ChR2-expressing β-cells as we performed at low glucose (Figure 3.A-B). In

untreated control islets $62.3\pm1.4\%$ of the islet within the imaging plane showed elevated $[Ca^{2+}]$

that was coincident with the ChR2 stimulation following single cell stimulation at 0.1 Hz (Figure

3.C). This indicates how connected individual cells are across the islet. Treatment with pro-

inflammatory cytokines led to a significant decrease in this $[Ca^{2+}]$ elevation ($39.7\pm3.4\%$,

$p<0.01$). Co-treatment with the S293 peptide led to a full recovery in elevated $[Ca^{2+}]$

($62.6\pm3.8\%$, $p<0.001$), indicating recovery in the overall islet connectivity.

We next investigated the extent to which islet cells were entrained, by analyzing the

spectral power of the 'natural oscillation frequency' of $[Ca^{2+}]$ oscillations that are observed prior

to ChR2 stimulation and analyzing the spectral power of the $[Ca^{2+}]$ oscillations at the frequency

of the ChR2 stimulation (Figure 3.D, Supplemental 1.A). In untreated control islets the spectral

power in the 'natural oscillation frequency' decreased upon single-cell ChR2 stimulation (Figure

3.E, $p<0.0001$). This was matched by an increase in the spectral power of the ChR2 driving

frequency upon single-cell ChR2 stimulation (Supplemental 1.B, $p<0.0001$). Following

treatment with pro-inflammatory cytokines the spectral power in the natural oscillation

frequency also decreased, although this decrease was lessened compared to in control islets

(Figure 3.F), indicating a less robust response to single-cell stimulation, as expected. Co-

treatment with pro-inflammatory cytokines and S293 peptide increased the change in spectral

power of the natural oscillation frequency compared that upon pro-inflammatory cytokines

alone, and similar to that in control islets. This shows a robust recovery in the response to

ChR2-mediated single-cell stimulation. Similarly, co-treatment with pro-inflammatory cytokines

and S293 peptide showed a slightly increased change in spectral power at the ChR2 driving

frequency, as compared to treatment with pro-inflammatory cytokines alone (Supplemental

1.C, $p = 0.2274$). These experiments were conducted over 10-20 islets per condition (from 3

ChR2-expressing mice), and our analysis indicates that treatment condition accounts for 19.9%

of total variation indicating the observed differences in oscillation at high-glucose are attributed

more to treatment rather than biological variability among samples.

Therefore, protecting gap junction coupling was able to fully recover islet-wide

connectivity and entrainment at elevated glucose that resulted from ChR2 stimulation of

individual cells.

Protecting gap junction coupling does not recover the cytokine-mediated disruption to first
responder cells and the 1st phase [Ca^{2+}] response: We next examined the impact of pro-
inflammatory cytokines and the protection of gap junction coupling on first responder cells and
the first phase [Ca^{2+}] response. Islets from Mip-Cre^{ER};ROSA26-lox-STOP-lox-GCaMP6s mice
that express GCaMP6 specifically in β-cells were used to assess changes in [Ca^{2+}] following
glucose elevation. We measured the [Ca^{2+}] response of individual cells and ranked their
response time after glucose stimulation. First responder cells were identified as the 10% of
cells which showed the earliest response, and last responders as the 10% of cells that showed
the latest response, with the spread in response time (Δt) being the time between first and last
responders (Figure 4.A). In control untreated islets the response time was consistently short
(Δt=179.5s$\pm$32.8s). Upon treatment with the pro-inflammatory cytokine cocktail for 1 hour, there
was no significant impact on the time between first and last responders (Figure 4.A-B,
p=0.3897). However following treatment with pro-inflammatory cytokines for 24 hours there
was a significant increase in the time between first and last responders compared to untreated
control islets (p=0.0043, Figure 4.A-B). Using this 24-hour treatment, co-treatment with S293
peptide did not show a decrease in Δt compared to cytokines alone (Δt=374$\pm$62s vs. 310$\pm$31s
respectively), indicating protection of gap junction coupling was not sufficient to recover a
narrow spread of the first phase response (Figure 4.C, p=0.72). Analysis of the response time
for first responders and last responder cells alone showed that treatment with pro-inflammatory
cytokines significantly increased the delay of last responding cells rather than affecting the
timing of first-responding cells, consistent with a reduced first-responder action (Supplemental
4. A,B). These findings were similar when other definitions for first-responder cells were used
(Supplemental 3.B). S293 peptide alone showed a similar time between first and last
responders as untreated controls. Under 1 hour cytokine treatment, co-treatment with the S293
peptide had no impact (Supplemental 2.A).
Clusters of first responders were previously found to be consistent upon repeated
glucose stimulations [28]; where ~35% of first responder cells remained first responders and
~75% of first responders remained within a 1 cell distance separation. We tested whether this
consistency was affected by both pro-inflammatory cytokines and protection of gap junction
coupling. We identified the top 10% of responding cells during an initial glucose stimulation and
determined what proportion of these cells remained first responders or nearest neighbors to
first responders during a repeated glucose stimulation (Figure 4.D). Upon cytokine treatment,

there was a slight, but not significant, increase in the consistency of first responders upon

repeated glucose stimulation compared to untreated controls (57.9% vs. 38% respectively, p =

0.2835, Figure 4.E). Upon protection of gap junction coupling via co-treatment with S293

peptide there was a slight, but not significant, decrease in the consistency of first responders

relative to cytokine only treatment (39.5%, p = 0.3264, Figure 4.E), and similar to that measured

in untreated controls. The consistency of first responders after 1 hour of cytokine treatment

also did not significantly improve upon co-treatment with the S293 peptide (Supplemental 2.B).

Therefore, treatment with pro-inflammatory cytokines disrupted the first-phase timing,

which was impacted by the protection of gap junction coupling. However pro-inflammatory

cytokines increased the consistency of the first phase $[Ca^{2+}]$ response, which was reversed

upon protection of gap junction coupling.

Protection of gap junction coupling provides a partial recovery to the cytokine-mediated

disruption of of highly connected β-cell hubs: Finally, we examined the impact of pro-

inflammatory cytokines and the protection of gap junction coupling on highly connected β-cell

hubs. β-cell hubs have been demonstrated to maintain coordinated $[Ca^{2+}]$ dynamics [25,28,29],

and previous work indicated a decrease in β-cell hubs upon diabetogenic conditions [13,29].

Following Ca^{2+} imaging we analyzed individual cell time-courses and formed a functional

network and identified the most highly connected cells as β-cell hubs (Figure 5.A, see

Methods). In control islets the number of highly synchronized functional connections and highly

synchronized hubs was high and consistent with prior measurements [29,30]. Upon treatment

with pro-inflammatory cytokines for 1 hour, there was no significant impact the mean degree of

correlation (Figure 5.B, p=0.3708). However following treatment with pro-inflammatory

cytokines for 24 hours, there was a significant decrease in the mean degree of correlation

(Figure 5.B, p=0.008 and Figure 5.C, p = 0.0375). There was also a significant decrease in

proportion of highly synchronized functional connections or "links" (Figure 5.D, p=0.0094) and

modest decrease in highly connected β-cell hubs (Figure 5.E, p=0.3759). Co-treatment with

S293 peptide did not show a significant change in neither functional connection nor β-cell hubs

compared to cytokines alone (Figure 5.C-E, p=0.9832 and p=0.8101 respectively). While the

proportion of hub-like cells across the entire β-cell population slightly increased upon co-

treatment with the S293 peptide compared to cytokine-treated islets, this was not statistically

significant. This indicates that protection of gap junction coupling was not sufficient to recover

highly synchronized functional connections and highly connected β-cell hubs. S293 peptide

alone showed similar functional connections and β-cell hubs as untreated controls. These
findings were similar when other definitions for highly-connected cells were used
(Supplemental 3.C). Under 1 hour cytokine treatment, co-treatment with S293 had no impact
on mean correlation or number of links per islet (Supplemental 2.C,D).

Approximately 43% of β-cell hubs remained as hubs following repeated glucose
stimulation under control conditions (Figure 5.F). We tested whether this consistency is affected
by both pro-inflammatory cytokines and protection of gap junction coupling. Upon cytokine
treatment, there was a decrease in the consistency of hubs (31%) compared to untreated
control islets. Protection of gap junction coupling via co-treatment with S293 peptide showed a
trend toward improved consistency of hubs compared to cytokine treatment alone (66%,
p=0.2109, Figure 5.G), with similar measurements to S293 peptide treatment alone, albeit
lacking statistical significance. Under 1 hour cytokine treatment, co-treatment with S293 did not
increase the population of hub cells as a percent, per islet (Supplement 2.E,F).

Therefore, treatment with pro-inflammatory cytokines reduced the presence of highly
synchronized functional connections and β-cell hubs. Restoration of gap junction coupling, and
subsequent islet-wide synchronization has little influence on hubs but maintains the
consistency of those remaining hub cells.

Discussion

Cx36 gap junctions have previously been identified as important regulators of β-cell coordination and islet function through the electrical coupling of neighboring β-cells. Pharmacological or genetic improvements in Cx36 gap junctions has been shown to recover $[Ca^{2+}]$ and insulin secretion responses in models of diabetes *in vitro*, and delay onset of diabetes *in vivo*, indicating a protective role that gap junctions play in islet function [12,31]. Functional sub-populations, which emerge as a result of β-cell heterogeneity, have been identified, and these sub-populations can drive the coordinated islet electrical response [27]. The goal of this study was to examine the influence of gap junctions on functional sub-populations of β-cells in a cytokine-mediated diabetogenic environment. Towards this, in the presence of a cocktail of pro-inflammatory cytokines, we protected Cx36 gap junction coupling using a peptide which we previously developed [12]. We then applied both optogenetics and calcium imaging to measure cell recruitment and calcium dynamics to identify and characterize three functional sub-populations that have previously been described [28] including first responders and hubs. Gap junction coupling could be successfully protected under pro-inflammatory conditions, and this was accompanied by recovery in islet-wide entrainment by ChR2 stimulation. However, optogenetic stimulation, and/or analysis of calcium dynamics revealed that functional subpopulations remained largely impaired. We conclude that while Cx36 gap junction coupling is critical for overall islet coordination and emergence of functional sup-populations, protection of gap junctions is not sufficient to recover the loss of these subpopulations upon cytokine-mediated disruption. Therefore cell-intrinsic properties are likely more influential in driving the impairment of functional sub-populations under diabetogenic conditions.

Impairment of functional sub-populations under diabetogenic conditions: Increased levels of proinflammatory cytokines within the islet microenvironment, including IL1-β and TNF-α, is a key feature of diabetes and has been attributed to driving islet dysfunction [16,32,33]. Prior studies have found that functional sub-populations are also disrupted in diabetes, such as reduced numbers of β-cell hubs under both pro-inflammatory cytokines and glucolipotoxic conditions [29], reduced consistency of wave initiator/leader cells under high-fat diet conditions [15] and reduced presence of hubs and leaders in islets from human subjects with type2 diabetes [34]. In this study we extended these findings to show both the cells that recruit a large proportion of the neighbors to show elevated $[Ca^{2+}]$ upon stimulation at low glucose ('high-

recruiting'), and the response time of first responder cells that respond earliest to glucose were

both disrupted by pro-inflammatory cytokines. This points to a general occurrence for functional

sub-populations to be disrupted under diabetogenic conditions. Notably, each of these

functional sub-population is associated with driving a specific aspect of islet electrical response,

which were also disrupted under the pro-inflammatory cytokines. For example, β-cell hubs

coordinate oscillatory dynamics across the islet, and the synchronization of these oscillations

was disrupted by pro-inflammatory cytokines. First responders coordinate and enhance the

initial first-phase response to glucose elevation, which was also disrupted by pro-inflammatory

cytokines. Thus, disruption to these functional sub-populations may be contributing to islet

dysfunction under diabetogenic conditions.

Influence of gap junction disruption to functional subpopulations under diabetogenic conditions:

Functional sub-populations, such as β-cell hubs, first-responders, and those that show high

ChR2-recruitment of $[Ca^{2+}]$ at low glucose, all exert their influence to coordinate and recruit

other β-cells via Cx36 gap junction coupling. Therefore, we hypothesized that pro-inflammatory

cytokines disrupt these sub-populations and their action because of diminished gap junction

coupling. By using the S293 peptide we could recover functional gap junction coupling

equivalent to that in control untreated islets indicated by FRAP. However, we did not observe

substantial recoveries in the presence and action of functional sub-populations when gap

junctions were protected: we observed only partial recovery to ChR2-recruitment at low

glucose, a slight recovery of β-cell hubs and no recovery to first responders. One simple

explanation is non-specific effects of the S293 peptide, however prior studies did not observe

non-specific effects on $[Ca^{2+}]$. Further, we observed a recovery in islet-wide entrainment of

$[Ca^{2+}]$ dynamics upon ChR2 stimulation at elevated glucose, which is expected to be more

dependent on gap junction electrical coupling and thus confirms the effects of the S293 peptide.

These findings therefore indicate that the disruptions to functional sub-populations under pro-

inflammatory cytokines are not primarily due to disruptions to gap junction coupling.

While initially unexpected, our findings are consistent with the fact that gap junction

coupling is not per se a defining feature of these sub-populations. For example, cells with high

ChR2-recruitment at low glucose do not show increased gap junction coupling [27], β-cell hubs

do not show increased gap junction coupling [30], and first responders show lower gap junction

coupling [28]. Thus, gap junction coupling does not drive the formation of these sub-

populations, rather they are defined by cell-intrinsic properties such as elevated glucose

metabolism or electrical differences, as well as other environmental properties such as the potential influence of α- or δ-cells [35]. Dysfunction to these cell intrinsic properties or altered communication with α- or δ-cells is therefore likely driving their disruption. However, as discussed above, we note that some amount of gap junction coupling will still be required for these cells to exert their influence across the islet [3].

Enhancing gap junction coupling can recover islet function in diabetogenic conditions [12,16,36]. Indeed, we observed a recovery in overall islet entrainment at elevated glucose by ChR2 stimulation. This enhanced function has also been demonstrated via both genetic and pharmacological approaches in models of diabetes *in vitro* and *in vivo*. While promising, our findings indicate that targeting gap junction coupling alone will be insufficient to fully recover islet function, given that functional β-cell sub-populations will still be disrupted. For example, caloric restriction is able to improve islet dysfunction that results from a high-fat diet; yet, while gap junction coupling recovered, wave initiators/leaders did not [15]. Inclusion of the GLP1R agonist Exendin-4 with pro-inflammatory cytokines recovered gap junction coupling and coordinated [Ca^{2+}] dynamics [37], while also impacting insulin secretion pathways via cAMP signaling. Thus, ensuring intrinsic factors and/or communication with α- and δ-cells are recovered will likely be required. Future work should examine which of these factors is important to both define functional sub-populations and preserve their action on islet function under diabetogenic conditions.

Limitations and Future Work: Our findings indicate that protection of gap junctions alone is not sufficient to recover cytokine-mediated loss of functional subpopulations. However, other factors within the islet microenvironment during diabetes can also impact islet function, including gluco- and lipotoxicity, amyloid plaque formation, and other pro-inflammatory cytokines such as IL6. Thus, extending our findings to other invitro treatments and mouse models of diabetes will be needed in future work. Another important factor is that our study examined a 24-hour treatment with pro-inflammatory cytokines and/or S293 peptide. This compares to a 1-6 hour treatment time in previous studies that support the recovery of gap junctions and islet function by S293 [12]. While we observed a protection of gap junction coupling, it is possible that under more acute treatments we will observe some recovery in functional sub-populations and a further enhancement in gap junction coupling, given that Cx36 turnover has a half-life of ~3 hours [38]. After 24 hours of cytokine treatment, observed loss of functional subpopulations may therefore result from effects of prolonged cytokine exposure in

addition to disrupted gap junctions, further supporting our conclusion that recovery of gap
junctions alone is not sufficient in restoring functional subpopulations under diabetogenic
conditions. Nevertheless, a longer-term treatment is likely more reflective of the islet
microenvironment in diabetes and should be further studied. Lastly, we acknowledge that our
study primarily focuses on β-cells despite potential influences of other existing cell types
including α- and δ-cells on β-cell behavior. Further studies addressing intercellular crosstalk is
necessary to fully understand the global effects of cytokine treatment and the role of gap
junctions.

Summary: Overall, we demonstrate that proinflammatory cytokines disrupt the presence and
action of functional β-cell sub-populations such as hubs and first responders, as well as the
cell-recruitment of $[Ca^{2+}]$. Preventing the decline in gap junction coupling was not sufficient for
recovering these functional β-cell sub-populations. This correlates with gap junction coupling
not being a factor that defines β-cell heterogeneity within the islet, but rather cell intrinsic
properties or other aspects of intra-islet communication instead being disrupted by the pro-
inflammatory environment.

 Figure Descriptions

Figure 1. Restoration of cytokine-mediated disruptions to gap junction coupling, as measured

by Fluorescence Recovery After Photobleaching (FRAP). (A) Isolated mouse islets were

treated with a cocktail of pro-inflammatory cytokines (0.5 ng/mL IL1-β, 1 ng/mL TNF-α, and 10

ng/mL IFN-γ) and co-incubated with 30 μM S293 peptide for 1 hour before imaging. (B)

Representative image of Rhodamine123 loaded islet pre- and post-photobleaching, along with

FRAP time-course after photobleaching, showing fluorescence recovery in the bleached

region. (C) Rate of recovery was calculated as a proxy for gap junction permeability in

untreated, cytokine, cytokine plus S293 peptide, and S293 peptide alone treatments. Data

represents Mean $\pm$ S.E.M. with each islet represented by a data point (from 6 mice), and each

measurement is normalized to the mean control value per mouse. * indicates $p < 0.05$.

Figure 2. Impact of gap junction restoration on the cytokine-mediated disruption to ChR2-

mediated recruitment of $[Ca^{2+}]$. (A) A ChR2-expressing islet (left), where a single ChR2-YFP

expressing cell was stimulated at 0.1 Hz (i) and calcium response was recorded across the

islet which exhibited both regions of recruited (ii) and unrecruited (iii) cells in untreated

conditions (middle). The result $[Ca^{2+}]$ activity was then quantified (right). (B) As in A, under

cytokine-treated conditions. (C) Histogram indicating the proportion of the islet recruited by a

single cell stimulation under each treatment condition. (D) Mean proportion of the islet recruited

by a single cell stimulation under each treatment condition. Data represents Mean $\pm$ S.E.M.

with each islet represented by a data point (from 15 mice). (E) The proportion of cells that were

able to recruit an area larger than 40% of the islet under the cytokine and S293 conditions

indicated. ** indicates $p < 0.01$, **** indicates $p < 0.0001$.

Figure 3. Impact of gap junction restoration *on* ChR2 single cell stimulation and islet

entrainment at high glucose. (A) Representative calcium responses, where a single ChR2-YFP

expressing cell was stimulated at 0.1 Hz at elevated glucose (i), and calcium response was

recorded across the islet which exhibited both regions of recruited/entrained (ii) and

unrecruited/not entrained (iii) cells. The result $[Ca^{2+}]$ activity was then quantified (right). (B) As

in A, under cytokine-treated conditions. (C) Mean proportion of the islet recruited by a single

cell at high glucose. Data presented as Mean $\pm$ S.E.M. with each islet. (D) Power spectra of

the $[Ca^{2+}]$ oscillation in control islets pre-ChR2 stimulation (top) and during-stimulation

(bottom). The dominant natural oscillation frequency (NOF) is highlighted in blue and the ChR2

driving frequency is highlighted in orange (E). Mean normalized power of the NOF, pre and

post ChR2 stimulation under each treatment condition. (F) Ratio of the spectral power of the

NOF pre/post stimulation. Data represents Mean ± S.E.M. with each experimental islet

represented by a data point. All data represents n = 3 mice, 2-3 islets per condition per mouse.

* indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$, **** indicates $p < 0.0001$.

Figure 4. Impact of gap junction restoration on the cytokine-mediated disruption to the temporal

and spatial consistency of first responders. (A) Time-course of $[Ca^{2+}]$ in first responder cells

(red) and last responder cells (green) following stimulation by high (11mM) glucose, under

control or cytokine-treated conditions (for 1h or 24h). (B) The mean time spread between the

first and last responders (ΔT) in islets treated with cytokines for 24 hours or 1 hour, compared

to untreated. Data represents Mean ± S.E.M. with each islet represented by a data point (n =

4 mice). (C) The mean time spread in islets treated for 24 hours, with cytokines and/or S293

peptide recovery of gap junction coupling. (D) Time-course of $[Ca^{2+}]$ in first responder cells

(red) and last responder cells (green), subjected to repeated stimulation from low to high

glucose (E) The percent consistency of first responders in untreated cells (~40%) or those with

cytokines and/or S293 peptide recovery of gap junction coupling. Data represents Mean ±

S.E.M. with each islet represented by a data point (n = 9 mice). ** indicates $p < 0.01$, ***

indicates $p < 0.001$.

Figure 5. Impact of gap junction restoration on the cytokine-mediated disruption of highly

connected β-cell hubs. (A) Representative connectivity map (above) and time-courses (below)

of islets under control or cytokine-treated conditions (for 1h or 24h). (B) The mean correlation

coefficient in islets treated with cytokines for 24 hours or 1 hour, compared to untreated. (C)

Mean correlation coefficient in islets treated for 24 hours, with cytokines and/or S293 peptide

recovery of gap junction coupling. (D) As in C for the average number of links between cells.

(E) As in C for the proportion of hub cells among the entire islet cell population. (F) Time-course

of $[Ca^{2+}]$ in β-cell hubs that are consistent or not consistent, together with representative

connectivity maps. (G) The percent consistency of hubs in untreated cells (>50%) or those with

cytokines and/or S293 peptide recovery of gap junction coupling. Data represents Mean ±

S.E.M. with each islet represented by a data point (n = 10 mice). * indicates p < 0.05, **

indicates p < 0.01, *** indicates p < 0.001.

Supplement 1. Spectral analysis of ChR2 single cell stimulation at high glucose. (A) Power

spectra of the [Ca^{2+}] oscillation in control islets post-ChR2 stimulation. The natural oscillation

frequency (NOF) is highlighted in blue and the ChR2 driving frequency is highlighted in orange

(B). Mean normalized power of the ChR2 driving frequency, pre and post ChR2 stimulation

under each treatment condition. Data presented as Mean ± S.E.M. with each experimental islet

represented by a data point. (C) Ratio of the spectral power of the ChR2 driving frequency

post/pre stimulation averaged across all islets. Data presented as Mean ± S.E.M. with each

experimental islet represented by a data point. All data represents n = 3 mice, 2-3 islets per

condition per mouse. * indicates p < 0.05, ** indicates p < 0.01, *** indicates p < 0.001, ****

indicates p < 0.0001.

Supplement 2. First responder and hub cells under 1h cytokine treatment with/without gap

junction restoration. (A) The mean time spread in islets treated for 1 hour, with cytokines and/or

S293 peptide recovery of gap junction coupling. (B) The percent consistency of first responders

in untreated cells (~40%) or those with cytokines and/or S293 peptide recovery of gap junction

coupling. (C) Mean correlation coefficient in islets treated for 1 hour, with cytokines and/or S293

peptide recovery of gap junction coupling. (D) As in C for the average number of links between

cells. (E) As in C for the proportion of hub cells among the entire islet cell population. (F) As in

C for percent consistency of hubs. Data represents Mean ± S.E.M. with each islet represented

by a data point (n = 4 mice).

Supplement 3. Threshold-dependent variation in defined subpopulations. (A) The proportion

of cells that were able to recruit an area larger than 30%, 40% and 50% of the islet upon single

cell ChR2-stimulation. Data represents mean area across all islets per condition. (B) The mean

time spread between the first and last responders (ΔT) defined as the earliest and latest 10%

of cells, earliest and latest 20% of cells and earliest 5 cells and latest 5 cells. (C) Population of

hub cells per islet and corresponding highest number of links for correlation thresholds of 0.9,

0.8, and 0.5. Data represents Mean ± S.E.M. with each islet represented by a data point (n =

4 mice).

Supplement 4. Time of first and last responder cells after 24h cytokine treatment with/without

gap junction restoration. (A) The exact time of the 10% of fastest responding β-cells to reach

first phase after stimulation with high glucose (B) The exact time of the 10% of slowest

responding β-cells to reach first phase after stimulation with high glucose. Data represents

Mean $\pm$ S.E.M. with each cell represented by a data point (n = 9 mice) ** indicates $p < 0.01$, ***

indicates $p < 0.001$, **** indicates $p < 0.0001$.

References

1. Federation, I.D. *Diabetes around the world in 2021*. 2022 [cited 2024.
2. Allagnat, F., D. Martin, D.F. Condorelli, G. Waeber, and J.-A. Haefliger, *Glucose represses connexin36 in insulin-secreting cells.* Journal of cell science, 2005. **118**(22): p. 5335-5344.
3. Benninger, R.K., M. Zhang, W.S. Head, L.S. Satin, and D.W. Piston, *Gap junction coupling and calcium waves in the pancreatic islet.* Biophysical journal, 2008. **95**(11): p. 5048-5061.
4. Farnsworth, N.L., A. Hemmati, M. Pozzoli, and R.K. Benninger, *Fluorescence recovery after photobleaching reveals regulation and distribution of connexin36 gap junction coupling within mouse islets of Langerhans.* The Journal of physiology, 2014. **592**(20): p. 4431-4446.
5. Ravier, M.A., M. Guldenagel, A. Charollais, A. Gjinovci, D. Caille, G. Sohl, C.B. Wollheim, K. Willecke, J.-C. Henquin, and P. Meda, *Loss of connexin36 channels alters β-cell coupling, islet synchronization of glucose-induced Ca2+ and insulin oscillations, and basal insulin release.* Diabetes, 2005. **54**(6): p. 1798-1807.
6. Pipeleers, D.G., *Heterogeneity in pancreatic β-cell population.* Diabetes, 1992. **41**(7): p. 777-781.
7. Stožer, A., M. Gosak, J. Dolenšek, M. Perc, M. Marhl, M.S. Rupnik, and D. Korošak, *Functional connectivity in islets of Langerhans from mouse pancreas tissue slices.* PLoS computational biology, 2013. **9**(2): p. e1002923.
8. Benninger, R.K., W.S. Head, M. Zhang, L.S. Satin, and D.W. Piston, *Gap junctions and other mechanisms of cell–cell communication regulate basal insulin secretion in the pancreatic islet.* The Journal of physiology, 2011. **589**(22): p. 5453-5466.
9. Speier, S., A. Gjinovci, A. Charollais, P. Meda, and M. Rupnik, *Cx36-mediated coupling reduces β-cell heterogeneity, confines the stimulating glucose concentration range, and affects insulin release kinetics.* Diabetes, 2007. **56**(4): p. 1078-1086.
10. Head, W.S., M.L. Orseth, C.S. Nunemaker, L.S. Satin, D.W. Piston, and R.K. Benninger, *Connexin-36 gap junctions regulate in vivo first-and second-phase insulin secretion dynamics and glucose tolerance in the conscious mouse.* Diabetes, 2012. **61**(7): p. 1700-1707.
11. Hodson, D.J., R.K. Mitchell, E.A. Bellomo, G. Sun, L. Vinet, P. Meda, D. Li, W.-H. Li, M. Bugliani, and P. Marchetti, *Lipotoxicity disrupts incretin-regulated human β cell connectivity.* The Journal of clinical investigation, 2013. **123**(10): p. 4182-4194.
12. St. Clair, J.R., M.J. Westacott, J. Miranda, N.L. Farnsworth, V. Kravets, W.E. Schleicher, J.M. Dwulet, C.H. Levitt, A. Heintz, and N.W. Ludin, *Restoring connexin-36 function in diabetogenic environments precludes mouse and human islet dysfunction.* The Journal of Physiology, 2023. **601**(18): p. 4053-4072.
13. Farnsworth, N.L., R.L. Walter, A. Hemmati, M.J. Westacott, and R.K. Benninger, *Low level pro-inflammatory cytokines decrease connexin36 gap junction coupling in mouse and human islets through nitric oxide-mediated protein kinase Cδ.* Journal of Biological Chemistry, 2016. **291**(7): p. 3184-3196.
14. Carvalho, C.d.F., R.B. Oliveira, A. Britan, J. Santos-Silva, A.C. Boschero, P. Meda, and C.B. Collares-Buzato, *Impaired β-cell-β-cell coupling mediated by Cx36 gap junctions in prediabetic mice.* American Journal of Physiology-Endocrinology and Metabolism, 2012. **303**(1): p. E144-E151.
15. Corezola do Amaral, M.E., V. Kravets, J.M. Dwulet, N.L. Farnsworth, R. Piscopio, W.E. Schleicher, J.G. Miranda, and R.K. Benninger, *Caloric restriction recovers impaired β-cell-β-cell gap junction coupling, calcium oscillation coordination, and insulin secretion in prediabetic*

mice. American Journal of Physiology-Endocrinology and Metabolism, 2020. **319**(4): p. E709-E720.

16. Farnsworth, N.L., R.A. Piscopio, and R.K. Benninger, *Modulation of gap junction coupling within the islet of langerhans during the development of type 1 diabetes.* Frontiers in Physiology, 2022. **13**: p. 913611.

17. Westacott, M.J., N.L. Farnsworth, J.R. St. Clair, G. Poffenberger, A. Heintz, N.W. Ludin, N.J. Hart, A.C. Powers, and R.K. Benninger, *Age-dependent decline in the coordinated [Ca2+] and insulin secretory dynamics in human pancreatic islets.* Diabetes, 2017. **66**(9): p. 2436-2445.

18. Ravier, M., J. Sehlin, and J. Henquin, *Disorganization of cytoplasmic Ca 2+ oscillations and pulsatile insulin secretion in islets from ob/ob mice.* Diabetologia, 2002. **45**: p. 1154-1163.

19. Heimberg, H., A. De Vos, A. Vandercammen, E. Van Schaftingen, D. Pipeleers, and F. Schuit, *Heterogeneity in glucose sensitivity among pancreatic beta-cells is correlated to differences in glucose phosphorylation rather than glucose transport.* The EMBO journal, 1993. **12**(7): p. 2873-2879.

20. Roscioni, S.S., A. Migliorini, M. Gegg, and H. Lickert, *Impact of islet architecture on β-cell heterogeneity, plasticity and function.* Nature Reviews Endocrinology, 2016. **12**(12): p. 695-709.

21. Wang, Y.J., J. Schug, K.-J. Won, C. Liu, A. Naji, D. Avrahami, M.L. Golson, and K.H. Kaestner, *Single-cell transcriptomics of the human endocrine pancreas.* Diabetes, 2016. **65**(10): p. 3028-3038.

22. Wojtusciszyn, A., M. Armanet, P. Morel, T. Berney, and D. Bosco, *Insulin secretion from human beta cells is heterogeneous and dependent on cell-to-cell contacts.* Diabetologia, 2008. **51**: p. 1843-1852.

23. Benninger, R.K., T. Hutchens, W.S. Head, M.J. McCaughey, M. Zhang, S.J. Le Marchand, L.S. Satin, and D.W. Piston, *Intrinsic islet heterogeneity and gap junction coupling determine spatiotemporal Ca2+ wave dynamics.* Biophysical journal, 2014. **107**(11): p. 2723-2733.

24. Giordano, E., D. Bosco, V. Cirulli, and P. Meda, *Repeated glucose stimulation reveals distinct and lasting secretion patterns of individual rat pancreatic B cells.* The Journal of clinical investigation, 1991. **87**(6): p. 2178-2185.

25. Benninger, R.K. and V. Kravets, *The physiological role of β-cell heterogeneity in pancreatic islet function.* Nature Reviews Endocrinology, 2022. **18**(1): p. 9-22.

26. Rutter, G.A., A. Gresch, L. Delgadillo Silva, and R.K. Benninger, *Exploring pancreatic beta-cell subgroups and their connectivity.* Nature metabolism, 2024: p. 1-15.

27. Westacott, M.J., N.W. Ludin, and R.K. Benninger, *Spatially organized β-cell subpopulations control electrical dynamics across islets of Langerhans.* Biophysical journal, 2017. **113**(5): p. 1093-1108.

28. Kravets, V., J.M. Dwulet, W.E. Schleicher, D.J. Hodson, A.M. Davis, L. Pyle, R.A. Piscopio, M. Sticco-Ivins, and R.K. Benninger, *Functional architecture of pancreatic islets identifies a population of first responder cells that drive the first-phase calcium response.* PLoS Biology, 2022. **20**(9): p. e3001761.

29. Johnston, N.R., R.K. Mitchell, E. Haythorne, M.P. Pessoa, F. Semplici, J. Ferrer, L. Piemonti, P. Marchetti, M. Bugliani, and D. Bosco, *Beta cell hubs dictate pancreatic islet responses to glucose.* Cell metabolism, 2016. **24**(3): p. 389-401.

30. Briggs, J.K., A. Gresch, I. Marinelli, J.M. Dwulet, D.J. Albers, V. Kravets, and R.K. Benninger, *β-cell intrinsic dynamics rather than gap junction structure dictates subpopulations in the islet functional network.* Elife, 2023. **12**: p. e83147.

31. Farnsworth, N.L., R.A. Piscopio, W.E. Schleicher, D.G. Ramirez, J.G. Miranda, and R.K.
Benninger, *Modulation of gap junction coupling within the islet of langerhans during the*
development of type 1 diabetes. Frontiers in Physiology, 2022. **13**: p. 913611.
32. Akash, M.S.H., K. Rehman, and S. Chen, *Role of inflammatory mechanisms in pathogenesis of*
type 2 diabetes mellitus. Journal of cellular biochemistry, 2013. **114**(3): p. 525-531.
33. Alexandraki, K., C. Piperi, C. Kalofoutis, J. Singh, A. Alaveras, and A. Kalofoutis, *Inflammatory*
process in type 2 diabetes: The role of cytokines. Annals of the New York Academy of Sciences,
2006. **1084**(1): p. 89-117.
34. Gosak, M., R. Yan-Do, H. Lin, P.E. MacDonald, and A. Stožer, *Ca2+ oscillations, waves, and*
networks in islets from human donors with and without type 2 diabetes. Diabetes, 2022.
71(12): p. 2584-2596.
35. Jin, E., J.K. Briggs, R.K. Benninger, and M.J. Merrins, *Glucokinase activity controls*
subpopulations of β-cells that alternately lead islet Ca2+ oscillations. bioRxiv, 2024.
36. Klee, P., F. Allagnat, H. Pontes, M. Cederroth, A. Charollais, D. Caille, A. Britan, J.-A. Haefliger,
and P. Meda, *Connexins protect mouse pancreatic β cells against apoptosis.* The Journal of
clinical investigation, 2011. **121**(12): p. 4870-4879.
37. Farnsworth, N.L., R. Walter, R.A. Piscopio, W.E. Schleicher, and R.K. Benninger, *Exendin-4*
overcomes cytokine-induced decreases in gap junction coupling via protein kinase A and Epac2
in mouse and human islets. The Journal of physiology, 2019. **597**(2): p. 431-447.
38. Wang, H.Y., Y.-P. Lin, C.K. Mitchell, S. Ram, and J. O'Brien, *Two-color fluorescent analysis of*
connexin 36 turnover: relationship to functional plasticity. Journal of cell science, 2015.
128(21): p. 3888-3897.
729

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Supplemental Figure S1

Supplemental Figure S2

Supplemental Figure S3

A

B

Supplemental Figure S4