I: TITLE PAGE

Impact of Acute Endurance Exercise on Alternative Splicing in Skeletal Muscle.

Alexander Ahn, MS[1]

ORCID: 0009-0009-7865-2061

Ann Arbor, Michigan, USA

May 6, 2025

Co-authors:

i. Jeongjin J. Kim, PhD[1] (ORCID: 0000-0002-6959-8544)
ii. Aaron L. Slusher, PhD[1] (ORCID: 0000-0003-2082-4827)
iii. Jeffrey Y. Ying, BS[1] (no ORCID)
iv. Eric Y. Zhang[1] (ORCID: 0009-0000-7975-0303)

Corresponding author: Andrew T. Ludlow, PhD[1]

- ORCID: 0000-0003-0423-4800; Ann Arbor, Michigan, USA
- 830 North University Avenue, Ann Arbor, Michigan 48109-1048, USA
- atludlow@umich.edu
- +1 (734) 763-4720

Affiliation:

1. School of Kinesiology, University of Michigan, Ann Arbor, Michigan, USA

II: ABSTRACT

Purpose: Alternative splicing (AS) is a highly conserved post-transcriptional mechanism, generating mRNA variants to diversify the proteome. Acute endurance exercise appears to transiently perturb AS in skeletal muscle, but transcriptome-wide responses are not well-defined. We aimed to better understand differential AS (DAS) and differential isoform expression (DIE) in skeletal muscle by comparing short-read (SRS) and long-read RNA sequencing (LRS) data.

Methods: Publicly accessible SRS of clinical exercise studies were extracted from the Gene Expression Omnibus. Oxford Nanopore LRS was performed on mouse gastrocnemius before and following treadmill exercise (30m running, n=5 mice/group, 20 total, 10 weeks old). Differential gene expression (DGE) and DIE were analyzed and validated using RT-PCR and immunoblots.

Results: Both SRS and LRS illustrated significant DGE in skeletal muscle post-exercise, including 89 RNA-binding proteins (RBPs). rMATS analysis of SRS revealed that exon-skipping and intron-retaining events were the most common. Swan analysis of LRS revealed several common genes across post-exercise cohorts with significant DAS but no DGE: 13 exercise-associated genes, including *mSirt2* (24.5% shift at 24hr post-exercise [24pe], p=0.005); 61 RBPs, including *mHnrnpa3* (28.5% at 24pe, p=0.02), *mHnrnpa1* (30.6% at 24pe, p=0.004), and *mTia1* (53.6% at 24pe, p=0.004).

Conclusions: We illustrated that acute endurance exercise can elicit changes in AS-related responses and RBP expression in skeletal muscle, especially at 24pe. SRS is a powerful tool for analyzing DGE but lacks isoform detection, posing a major gap in knowledge of "hidden" genes with no transcriptional but significant DIE and protein expression changes. Additionally, LRS can uncover previously unknown transcript diversity and mechanisms influencing endurance exercise adaptations and responses.

Keywords

Endurance exercise, alternative RNA splicing, RNA-binding proteins, splicing factors, skeletal

muscle.

Abbreviations

A3SS, alternative 3' splice site;

A5SS, alternative 5' splice site;

AS, alternative (RNA) splicing;

ATSS, alternative transcription start site;

ATTS, alternative transcription termination site;

DAS, differential alternative splicing;

DGE, differential gene expression;

DIE, differential isoform expression;

hsAA, *homo sapiens* aged active;

hsAS, *homo sapiens* aged sedentary;

hsYS, *homo sapiens* young sedentary;

LRS, long-read RNA sequencing;

mmYS, *mus musculus* young sedentary;

MXE, mutually exclusive exons;

RBP, RNA-binding protein;

RI, retained intron;

SE, skipped exon;

SRS, short-read RNA sequencing.

III: BODY

1 INTRODUCTION

The positive impact of exercise on one's health is undeniable. With skeletal muscle

accounting for approximately 40% of body mass in adults, it is the most directly affected tissue

from both the beneficial effects of physical activity as well as the deleterious effects of

sedentarism. However, a large portion of the population fails to meet the minimum amount of

exercise to reduce health risks. The U.S. Centers for Disease Control and Prevention reported

overall prevalence of physical inactivity was 25.3% in 2020,[1] while causing 9% of premature

death globally.[2] Why people fail to meet the minimum amount of exercise are varied but some

are simply unable to even if desired: for example, muscle dystrophy, motoneuron-related

sclerosis diseases, and obesity-induced osteoarthritis. To fully harness the health-promoting

power of endurance exercise, we must continue to uncover the fundamental mechanisms of

molecular responses and adaptations elicited by the exercise stimulus.

Endurance exercise promotes disease-free longevity via the enhancement of skeletal

muscle oxidative capacity among other important responses and adaptations.[3] Exercise is a

potent cellular stressor (e.g., mechanical, metabolic, oxidative stress, etc.) that triggers a series of

responses and adaptations.[4–7] Part of the adaptive process in skeletal muscle is changes in gene

expression regulation that regulates the protein landscape and alters structural and functional

outcomes. Presently, reports have focused on changes in transcript abundance and differential

gene expression (DGE).[8,9] For example, endurance exercise training increases mitochondrial

density, muscle-specific angiogenesis, and improves calcium handling in slow-twitch skeletal

myofibers by altering the expression of the genes encoding peroxisome proliferator-activated

receptor gamma coactivator 1 alpha (*PPARGC1A*, PGC-1α),[10] vascular endothelial growth factor

(*VEGF*),[11] and troponin T (*TNNT1*, ssTnT),[12] respectively. While *PPARGC1A*, *VEGF*, and

TNNT1 are all differentially expressed, there is also evidence that exercise impacts the

expression of different transcript variants of each of these genes,[12–15] indicating that alternative

RNA splicing (AS) is impacted by exercise.[16,17] To date, however, only one group has

investigated mRNA isoform expression using short-read RNA sequencing (SRS) in skeletal

muscle from individuals with varying levels of physical activity and observed associations

between activity level and mRNA isoform expression.[17,18]

RNA splicing and AS are RNA processing steps that offer additional regulatory layers in

gene expression. RNA splicing is mediated by a dynamic RNA-protein complex, the

spliceosome, that assembles on primary transcripts (i.e., precursor messenger RNA, pre-mRNA)

to remove introns and splice exons together to form the final protein-coding transcript (i.e.,

mature messenger RNA, mRNA).[19] In conjunction with the spliceosome, RNA-binding proteins

(RBPs) provide tissue-/context-specific regulation of splicing.[20] While most exons of a pre-

mRNA undergo constitutive splicing to form full-length variants, many undergo AS: the choice

of including or excluding certain exons and/or introns in the mRNA.[21,22] AS is a highly

conserved co-/post-transcriptional mechanism that functions to generate multiple RNA transcript

variants: these transcript variants may be intended for either translation to respective protein

isoforms,[23,24] non-coding RNAs,[25,26] or degradation via RNA decay mechanisms.[27,28] Therefore,

a single gene can form several protein isoforms depending on the demands of the cells in a

tissue.[23] For instance, a significant change in AS can orchestrate a switch in the expression of

certain protein isoforms. These differential use of isoforms between conditions is often referred

as isoform switching (or differential isoform expression, DIE),[13,14] which facilitates a functional

change in cellular responses to stimuli. The significance of AS is evident as nearly all

mammalian genes undergo AS and generate two or more protein isoforms, which vastly expands

the protein-coding capacity of the genome. AS also greatly impacts the function of cells in a

tissue, as well as their responses to homeostatic perturbations.[23]

Interestingly, transcriptome-wide impact of endurance exercise on AS and isoform

generation is currently unknown. Previously, Tonevitsky et al.[29] observed that acute aerobic

exercise with maximal intensity in trained skiers resulted in an increased gene expression from a

set of RBPs (DEAD-box helicase 17, *DDX17*; DEAD-box helicase 46, *DDX46*; heterogeneous

nuclear ribonucleoprotein R, *HNRNPR*; pre-mRNA processing factor 4 kinase, *PRPF4B*; and SR

protein kinase 2, *SRPK2*) that regulate the formation of the precatalytic spliceosome. Similarly,

our laboratory also observed that exercise resulted in reduced protein expression of splicing

factor 3B subunit 4 (SF3B4) from young sedentary mouse gastrocnemius in response to acute

treadmill running.[30] These findings indicate that a single bout of endurance exercise is sufficient

to potentially instigate differential AS (DAS) and DIE of many other genes through differential

expression and regulation of RBPs. As such, we hypothesized that exercise results in isoform

switching of a multitude of genes, including isoform switching of RBPs that regulate splicing.

Therefore, we set out to globally profile AS events (mRNA variants/DIE and exon-level events)

in skeletal muscle from a single bout of submaximal endurance exercise.

2 MATERIALS AND METHODS

2.1 Short-read RNA sequencing

The human SRS data had formerly been published by two different studies:

Pattamapraponont et al.[31] and Rubenstein et al.[32] The raw FASTQ files were downloaded from

publicly accessible NCBI Gene Expression Omnibus database (RRID: SCR_005012) (accession

numbers GSE87749 and GSE151066, respectively). Briefly, Pattamapraponont et al. consisted of

10 young (19-25 years) and healthy (BMI of 23.1 $\pm$ 2.8) Danish male participants with a

sedentary lifestyle (hsYS), and performed a single bout of intense (80% VO_2max) cycling

(cycle ergometer) for 15 minutes; muscle biopsies of the vastus lateralis were obtained pre-

(hsYS-pre) and 4 hours post-exercise (hsYS-4pe).[31] Rubenstein et al. studied two cohorts of

older individuals (65-90 years): 10 (8 males, 2 females) an active/endurance trained group (hsAA)

with a BMI of 24.2 $\pm$ 4.2; nine (7 males, 2 females) and a sedentary (hsAS) group with a BMI

of 28.7 $\pm$ 3.3. The participants performed a single bout of endurance exercise (cycling at 60-70%

of heart rate reserve) for 40 minutes. Muscle biopsies of the vastus lateralis were obtained pre-

(hsAA-pre & hsAS-pre) and 3 hours post-exercise (hsAA-3pe & hsAS-3pe).[32] To maintain

consistency for the initial processing and statistical analyses of the SRS data, only single-end

FASTQ files were used from the Pattamaprapanont et al. study; similarly, only males at pre- and

3 hours post-exercise time points with paired-end FASTQ files were used from the Rubenstein et

al. to maintain comparability between the two data sets (Figure 1A).

2.2 Animals

All animal experiments were approved by the University of Michigan Institutional

Animal Care and Use Committee (IACUC) and conducted as per the institutional guidelines.

C57BL/6-Tg(TERT)C10Hode/J (approximately 8 weeks old) were purchased from The Jackson

Laboratory (Bar Harbor, Maine, USA, RRID: IMSR_JAXX:006140). The mice were housed at

25°C on a 12:12-hour light-dark cycle, fed *ad libitum* laboratory mouse chow (Prolab® RMH

3000, 5P00, LabDiet by Purina, Nestlé, Vevey, Switzerland), and given free access to water. The

description of the running protocol was previously published.[30] Animals were randomly assigned

to each pre-/post-exercise time points. All original animals were included without exclusion nor

attrition. Investigators were not blinded to the groupings. Sample size and power analysis were

determined based on our previous study.[30] Briefly, 20 mice underwent acute treadmill running at

60% of maximal speed, approximately 75% of mouse VO_2max according to Fernando et al.,[33]

for 30 minutes (Figure 1B,S1M). Euthanasia practices were followed as suggested by the

American Veterinarian Medical Association guidelines (2020 edition): briefly, the animals were

anesthetized with isoflurane, followed by cervical dislocation, and cardiac removal.

Gastrocnemius muscles were collected at pre- (mmYS-pre; 3 males, 2 females), immediately

post-exercise (mmYS-ipe; 3 males, 2 females), 1 hour post-exercise (mmYS-1pe; 2 males, 3

females), and 24 hours post-exercise (mmYS-24pe; 3 males, 2 females). Gastrocnemius muscles

were flash frozen in liquid nitrogen and stored at –80°C for subsequent tissue processing and

RNA isolation.

2.3 Tissue processing, RNA isolation, and poly(A)-mRNA isolation

The gastrocnemius tissues were first powdered with a mortar and pestle in liquid nitrogen,

then either processed immediately or stored at –80°C, as described in Slusher et al.[30] Total RNA

from powdered tissues were isolated using the RNeasy Plus Universal Mini Kit (Qiagen, Cat#

73404) according to manufacturer instructions, and stored at –80°C. Isolated RNA was

quantified using the NanoDrop™ 2000 spectrophotometer and assessed using the Qubit™ RNA

IQ Assay Kits (Invitrogen, Cat# Q33221) with the Qubit 4 Fluorometer. Subsequently, 1.2 µg of

total RNA was used to isolate poly(A)-containing RNA with the NEBNext® Poly(A) mRNA

Magnetic Isolation Module (New England BioLabs, Cat# E7490S) accordingly. Poly(A)-mRNA

isolates were then quantified with Agilent Bioanalyzer (RRID: SCR_018043), NanoDrop™

2000 spectrophotometry, and Qubit Fluorometry.

2.4 Long-read RNA sequencing

1 ng of poly(A)-mRNA isolates were used to synthesize barcoded PCR-cDNA libraries

(Oxford Nanopore Technologies®, ONT; Cat# SQK-PCB109) according to the manufacturer

instructions. Subsequently, the prepared PCR-cDNA libraries were quantified, multiplexed at

equal molar concentrations, loaded onto a MinION™ R9.4.1 Flow Cell (ONT, Cat# FLO-

MIN106) and sequenced (long-read RNA sequencing, LRS) for 72 hours using the MinION™

Mk1C sequencer (ONT, Cat# MIN-101C) (Figure 1B). During the sequencing period, the flow

cell was "washed" with the Flow Cell Wash Kit (ONT, Cat# EXP-WSH004) and reloaded with

fresh multiplexed libraries as needed.

2.5 Data preprocessing

For the SRS retrieved data, the quality of the raw FASTQ files were checked and

trimmed to the minimum read length of 25 bases. Subsequently, RSEM-STAR[34,35] was used to

align and quantify the trimmed reads based on the human GENCODE release 42 (GRCh38.p13).

Matrices of "counts" (estimation of expected read counts), "abundance" (TPM-normalized

counts), and "length" (effective transcript lengths) were generated for statistical analyses (hsYS:

Figure S1A,E,I; hsAS: Figure S1B,F,J; and hsAS: Figure S1C,G,K).

For the LRS generated data, raw FASTQ (converted/basecalled from FAST5 with Guppy

MinKNOW v22.03.4; ONT, RRID: SCR_023196) files, which passed the MinKNOW quality

check with minimum q-score threshold of 8, were used for read alignment. Minimap2 v2.14[36]

(RRID: SCR_018550) and SAMtools v1.13[37] (RRID: SCR_002105) were used to align the

passed FASTQ files into SAM format; the mouse GENCODE release M29 (GRCm39) genome

sequence assembly was used for the alignment step. TranscriptClean[38] was used to sort and

"clean" the aligned SAM files. TALON[39] was used to quantify reads and generate an abundance

matrix (coverage = 0.75, identity = 0.8) in GTF format for statistical analyses (Figure S1D,H,L).

2.6 Differential gene expression and differential alternative splicing analyses

For the SRS-retrieved data, TXImport[40] was used to import the multidimensional RSEM-

STAR quantification matrix arrays at the gene-level to the R environment. Subsequently,

DESeq2[41] was used to perform gene-level differential expression (differential gene expression,

DGE) analyses with $\log_2$ fold changes ($\log_2$FC) and respective statistics (p-values adjusted for

FDR with default Benjamini-Hochberg method, BH; p-adj) on the quantification matrix arrays,

and enriched for significantly up-/down-regulated genes (Figure 2A–E). IsoformSwitch-

AnalyzeR[42] was initially used (Figure S2) to identify genes with isoform-switching events with

absolute difference in isoform fraction > 10% and p-adj < .05. Splicing type resulting in the

isoform switches were annotated as skipped exon (SE), alternative 5' splice site (A5SS),

alternative 3' splice site (A3SS), mutually exclusive exons (MXE), multiple exon-skipping

(MES), retained introns (RI), alternative transcription start site (ATSS), and alternative

transcription termination site (ATTS). Additionally, rMATS-turbo v4.1.2[43,44] was used to

directly analyze exon-level splicing events with difference in percent spliced in ($\Delta\Psi$ or DPSI)

and respective statistics on the trimmed FASTQ files, and enriched for genes with significant

splicing events annotated as SE, A5SS, A3SS, MXE, and RI. The transcript-level rMATS data

frames were filtered based on the absolute $\Delta\Psi$ > 10% and p-adj < .05 for the significant splicing

overlap analyses and for visualizing the proportions of splicing profile (Figure 4).

For the LRS-generated data, Swan v2.0[45] was used to generate data frames with $\log_2$FC

and respective significance statistics at the gene- and transcript-levels. Based on known genes

that were detected in at least half of the samples, the counts matrix was generated for the

downstream differential analyses. The corresponding DGE (gene-level) $\log_2$FC and respective

statistics from Swan were used for data visualizations (Figure 2F,G,H). Similarly, DIE analyses

generated absolute difference in percentage isoform use ($\Delta\Pi$ or DPI) to determine isoform-

switching data, whereby respective statistics from Swan were filtered based on $\Delta\Pi > 10\%$ and p-

adj $< .05$.

Functional enrichment analysis was performed on significantly up- and down-regulated

genes from the DGE analyses ($|\log_2$FC$| > 1.5$ and p-adj $< .05$) using either Database for

Annotation, Visualization, and Integrated Discovery (DAVID)[46] or PANTHER[47] to generate top

10 gene ontology (GO) terms. Enrichment analysis was only performed on data sets with greater

than 80 genes observed to be significant according to experiment specific statistical levels. With

the splicing and DIE data sets from both SRS and LRS data, multiple multivariate overlap

analyses across varying conditions and data sets were performed to identify any common genes

with potential biologically significant splicing events. Only overlapping genes with p-adj $< .05$

from bivariate hypergeometric tests were considered significant. All code scripts used for

bioinformatical data analyses are available in the Deep Blue data repository (DOI will be

provided upon publication) or upon request.

An in-house-derived RNA-binding protein (RBP) list was constructed. We generated this

list using two sources: the census of human RBPs[48] and a gene cards list search for RBPs using

human genes. From there we combined the two lists, generating a list of ~1700 human genes

without redundancy. Then we matched the human gene IDs to rodent gene IDs and came up with

1130 rodent genes that are conserved RBPs. This list was used to analyze RBP-specific

enrichment in the data sets.

2.7 Gel-based and droplet digital RT-PCR

256   1 µg of total RNA was used to synthesize cDNA using the SuperScript™ IV reverse

transcriptase (ThermoFisher Scientific, Cat# 18090010), as previously described in Slusher et

al.[30] All cDNAs were diluted 1:40 (20 µL of cDNA□+□780 µL nuclease-free water; ie, 1.25

ng/µL of RNA input) before use for both droplet digital PCR (ddPCR) and gel-based PCR

validations, and were stored at□−20°C thereafter. All cDNA samples were assayed for 18S

rRNA to ensure equal loading across the cohorts using ddPCR (Figure S5A, F: 5'-GTA ACC

CGT TGA ACC CCA TT-3', R: 5'-CCA TCC AAT CGG TAG TAG C-3'). Subsequently,

several genes found significant from the LRS DGE analysis were selected for validation using

RT-ddPCR (Figure 2I, *mUbe2d1*, exon 4 F: 5'-TCC TCA CTG TCC ACT TTC CG-3', exon 5 R:

5'-TCG ATA CAG TCA AAG CGG GC-3'; Figure 2J, *mFbxo32*, exon 2 F: 5'-TCA GCA GCC

TGA ACT ACG AC-3', exon 4 R: 5'-GGC AGT CGA GAA GTC CAG TC-3'; Figure 2K,

mFos, exon 2 F: 5'-TTT ATC CCC ACG GTG ACA GC-3', exon 2 R: 5'-TCT ACT TTG CCC

CTT CTG CC-3'). Several genes found significant from the LRS isoform-switching analysis

were also selected for validation using both RT-ddPCR and gel-based RT-PCR (Figure 5C,

mSirt2, exon 1 F: 5'-CTC TTC TTG TTT CCG CTG CC-3', exon 3 R: 5'-CTC CCT CAG TGT

CCG AGT CT-3'; Figure 6F, *mHnrnpa3*, exon 3 F: 5'-GGT GGA TGC TGC AAT GTG TG-3',

intron 3 R: 5'-ACC AAA CTG AAA CAT ATC CAA GT-3', exon 4 R: 5'-TTT CCC ACT CTG

CCT GTC TT-3'; Figure 6H, *mHnrnpa1*, exon 7 F: 5'-GGG GAT GGC TAT AAT GGA TTT

GGC-3', exon 9 R: 5'-GCA AAG TAC TGG CCT CCA CC-3'; Figure 6I, *mTia1*, exon 4 F: 5'-

GCA ACA ACC CCT AGC AGT CA-3', exon 6 R: 5'-CGC TGC TTT GAT GTC TTC GG-3').

2.7 Protein lysate preparation, antibody validation, and western blot analyses

To examine protein expression level of one of the prominent RBPs, heterogeneous

nuclear ribonucleoprotein A3 (hnRNP-A3), HEK 293 fetal kidney cells (ATCC, Cat# CLR-1573,

RRID: CVCL_0045) were used as positive controls to verify cross-species specificity of the

hnRNP-A3 antibody for the correct molecular weight at 42 kDa. HEK 293 blots were normalized

with GAPDH (conditions listed below). Additionally, C2C12 murine myoblasts (ATCC, Cat#

CRL-1772, RRID: CVCL_0188) were transfected with 30 nM of either scrambled (siCTL) or

mouse *Hnrnpa3*-targeting (exon 10) siRNAs (ThermoFisher Scientific, Cat# 4390771) in

Lipofectamine™ RNAiMAX transfection reagent (Invitrogen, Cat# 13778075) for 24 and 48

hours to knock-down hnRNP-A3 and thus validate the protein specificity of hnRNP-A3 antibody.

All cell lines used were authenticated by the manufacturer (ATCC) and visually verified in-

house. The cell pellets were resuspended in an SDS-based lysis buffer, sonicated, and

centrifuged to extract the total protein lysate. Total protein lysates were prepared from

gastrocnemius, as described in Slusher et al.,[30] were used to quantify protein expression level of

hnRNP-A3 in the skeletal muscle tissues. Protein concentrations were quantified using the Pierce

660 nm Protein Assay Kit (ThermoFisher Scientific, Cat# 1861426). These protein lysates from

cell lines and the animal-derived tissues were then prepared using 2X Laemmli loading buffer

(Bio-Rad, Cat#1610737) and boiled for 10 minutes at 95°C.

All protein lysates from cells and tissues were resolved by SDS-polyacrylamide gel

electrophoresis at 16 mAmp per gel for 1 hour, transferred to Mini 0.2 μm polyvinylidene

fluoride membranes using the Trans-Blot® Turbo™ Transfer System (Bio-Rad, Cat# 1704150)

with the rapid 7-minute mixed molecular weight transfer protocol. To verify cross-species

specificity of the hnRNP-A3 antibody, protein lysates of wild-type HEK 293 was used. To

quantify bands from the wild-type HEK 293 samples, specifically, loading control was

normalized with GAPDH (Cell Signaling Technology, Cat# 2118, RRID: AB_561053; 1:1000

dilution in 5% bovine serum albumin with 1X PBST) (Figure S6A). To validate protein

specificity of the hnRNP-A3 antibody, protein lysates of C2C12 myoblasts treated with either

siCTL or *Hnrnpa3* siRNA were used. To quantify bands from the C2C12 myoblasts, specifically,

loading control was normalized with total protein quantification of bands at 40-45 kDa using

Pierce™ Reversible Protein Stain Kit (ThermoFisher Scientific, Cat# 24585) (Figure S6B,C).

HnRNP-A3 from these cell lysates was detected with a rabbit polyclonal antibody (Abcam, Cat#

ab78300, RRID: AB_2041662; 1:1000 dilution factor in 5% bovine serum albumin with 1X

PBST and 0.02% sodium azide). To quantify bands from the skeletal muscle tissue, specifically,

loading control of protein lysates from the skeletal muscle tissue was normalized with total

protein quantification of bands at 40-45 kDa using Ponceau S stain (Sigma-Aldrich, Cat# P3504)

and detected for hnRNP-A3 with the validated polyclonal antibody (Figure 6G,S6D). All blots

were imaged with Bio-Rad ChemiDoc XRS+ Molecular Imager (RRID: SCR_019690) and

quantified with Bio-Rad Image Lab software. Values from either the reversible Pierce-/Ponceau

S-stained blots (tissue-based lysates) or the loading control blots (cell-based lysates) were

averaged and used to normalize the target protein quantification. All images that were used to

quantify hnRNP-A3 expression levels are included in Figure S6D.

2.8 Statistical analysis

Unless otherwise noted, the unpaired Student's *t*-test was used for two group

comparisons using a statistical package (GraphPad Prism 10, RRID: SCR_002798). One-way

ANOVAs were used to determine significance for multiple group comparisons of the western

blot data, whereby the groups (time points) had 1 level each. Subsequently, *post hoc* adjustments

were conducted with Tukey's HSD. Statistical significance was defined as a $p < .05$.

3 RESULTS

3.1 RNA-Seq identified differential gene expression in acutely exercised skeletal

muscle with and without training

We analyzed gene expression from public SRS data and compared these findings to our

acute exercise LRS study. DGE of vastus lateralis in hsYS-pre and hsYS-4pe was analyzed with

DESeq2 (Figure 2A): out of 13727 total detected genes, 19 were significantly up-regulated

($\log_2$FC > 1.5, p-adj < .05) including *PPARGC1A* (i.e., PGC-1α), nuclear receptor subfamily 4

group A member 3 (*NR4A3*), and heat shock protein family A member 1A/B (*HSPA1A/B*,

HSP70/72), while 10 were significantly down-regulated ($\log_2$FC < −1.5, p-adj < .05). There were

insufficient numbers of DGE genes, therefore it was not appropriate to conduct functional

enrichment analyses. DGE of vastus lateralis in hsAS-pre and hsAS-3pe was analyzed (Figure

2B): out of 14082 total detected genes, 201 were significantly up-regulated ($\log_2$FC > 1.5, p-adj

< .05) including *PPARGC1A*, *NR4A3*, and *HSPA1B*, while 9 were significantly down-regulated

($\log_2$FC < −1.5, p-adj < .05). We took all significantly up-regulated genes and performed

functional enrichment analysis (DAVID, statistical over-enrichment test). The hsAS-pre:3pe

revealed 204 terms (p-adj < .05; Figure 2C). Similarly, DGE of vastus lateralis in hsAA-pre and

hsAA-3pe was analyzed (Figure 2D): out of 14315 total detected genes, 112 were significantly

up-regulated ($\log_2$FC > 1.5, p-adj < .05) including *PPARGC1A*, *NR4A3*, nuclear receptor

subfamily 4 group A member 1 (*NR4A1*), and *HSPA1B*, while 4 were significantly down-

regulated ($\log_2$FC < −1.5, p-adj < .05). We took all significantly up-regulated genes and

performed functional enrichment analysis (DAVID, statistical over-enrichment test). The hsAA-

pre:3pe revealed 149 terms (p-adj < .05; Figure 2E). GO terms that were common between the

hsAS and hsAA groups were skeletal muscle cell differentiation (GO:0035914) and negative

regulation of apoptotic process (GO:0043066). The major insights we are able to draw from

these data are that there is a different response to acute aerobic exercise based on training status:

DGE analyses illustrated different up-regulated genes post-exercise in the older active

individuals with few GO terms in common between the two older cohorts.

DGE of gastrocnemius in mmYS-pre and mmYS-ipe, mmYS-pre and mmYS-1pe, and

mmYS-pre and mmYS-24pe were analyzed ($|\log_2FC| > 1.5$, p-adj $< .05$). At ipe, 6323 total genes

were detected, and we observed 23 differentially expressed genes (6 down-regulated and 17 up-

regulated); at 1pe, 5459 total genes were detected, and we observed 13 differentially expressed

genes (4 down-regulated and 9 up-regulated); and at 24pe, 6125 total genes were detected, and

we observed 31 differentially expressed genes (16 down-regulated and 15 up-regulated). Since

this was a conservative statistical cutoff, we relaxed the parameters to include more genes in our

analysis ($|\log_2FC| > 1.0$, p-adj $< .05$; Figure 2F, G and H). With these new statistical parameters,

at ipe, we detected 46 genes that reached statistical significance (13 down-regulated and 33 up-

regulated); at 1pe, we detected 32 genes that reached statistical significance (8 down-regulated,

24 up-regulated); and at 24pe, we detected 49 genes that reached statistical significance (26

down-regulated, 23 up-regulated). To validate our data sets we selected one gene at each time

point for RT-PCR validation. Ubiquitin-conjugating enzyme E2 D1 (*mUbe2d1*; LRS data:

$\log_2FC = 0.75$, p-adj $= 0.0002$), F-box protein 32 (i.e., atrogin-1, *mFbxo32*; LRS data: $\log_2FC =$

2.7, p-adj $= 0.002$), and fos proto-oncogene (*mFos*; LRS data: $\log_2FC = -3.9$, p-adj $= 0.0002$) at

ipe, 1pe, and 24pe, respectively. Using RT-ddPCR, we validated all three selected genes at each

timepoint (*mUbe2d1* was significantly up-regulated at ipe [22.6% increase, t-test $p = 0.03$);

mFbxo32 was up-regulated at 1pe [144% increase, t-test $p = 0.004$); and *mFos* was down-

regulated at 24pe [55.9% decrease, t-test $p = 0.01$]; Figure 2I,J,K, respectively). Genes

commonly observed to be differentially expressed in acute endurance exercise studies such as

mHspa8 (i.e., HSP70)[49] were observed in this analysis. Interestingly, we observed that exercise

impacted on genes related to protein degradation, such as *mFbxo32* and *mUbe2d1*, indicating

that exercise seemed to enhance clearing damaged proteins.[50,51]

We did not observe the common acute exercise-induced genes such as NRFs or PGC-1α

in our LRS data like we did using the publicly available human SRS data. A potential reason for

this is the low expression of these transcription factors, as well as the timing of muscle sampling

post-exercise in our mouse study (1 hour versus 3-4 hours post exercise in the humans).

3.2 Differentially expressed RNA-binding proteins in acutely exercised skeletal

muscle with and without training

Using a list of 1130 RBP genes generated in house (see methods above for details on how

this list was curated), select RBP genes were included in the heatmap if it was significantly

differentially expressed within a cohort. Significant up- and down-regulated RBPs ($log_2FC > 0.5$,

$log_2FC < -0.5$, raw $p < .05$) in response to an acute bout of exercise from all four cohorts

(mmYS, hsYS, hsAS, and hsAA) were queried, identifying 89 genes across all four groups

(Figure 3A–D, *p*-adj < .05 indicated with asterisks). The mmYS cohort had 3 RBPs that were

significantly down-regulated with *p*-adj < .05 in response to acute exercise (Figure 3A). We

show an additional 17 RBP genes (10 up-regulated, 7 down-regulated, raw *p* < .05) with less

conservative statistical parameters (Figure 3A). The hsYS cohort had a total of 9 and 1 RBPs

were significantly up- and down-regulated, respectively, in response to acute exercise (Figure

3B). The hsAS cohort had a total of 20 and 1 RBPs were significantly up- and down-regulated,

respectively, in response to acute exercise (Figure 3C). The hsAA cohort had a total of 50 and 4

RBPs were significantly up- and down-regulated, respectively, in response to acute exercise

(Figure 3D). Finally, we overlapped the RBPs to determine conserved and age-related changes in

RBP expression between cohorts (Figure 3E). We did not observe any overlapping genes across

the mice and human cohorts. This could indicate that there is not a well-conserved response to

endurance exercise and changes in the mRNA expression of RBPs or this could indicate

differences in the exercise protocols and muscles sampled between the cohorts (gastrocnemius

for the mice versus quadriceps for the humans). There were 6 overlapping RBPs that were

significant between the aged cohorts using hypergeometric testing (6.0-fold over-enrichment, $p =$

0.0002). There were also 5 overlapping RBPs that were significant between the YS and AA

cohorts using hypergeometric testing (10.5-fold over-enrichment, $p = 4.4e^{-5}$). There were also 2

overlapping RBPs that were significant between hsYS and hsAS cohorts using hypergeometric

testing (11.3-fold over-enrichment, $p = 0.02$).

Cytoplasmic polyadenylation element binding protein 4 (CPEB4) was a common up-

regulated RBP gene among the human cohorts. CPEB4 is a protein mainly found in the

cytoplasm that is a sequence-specific RBP and binds to uridine-rich sequence elements

(consensus sequence 5'-UUU UUA U-3') in the 3'-untranslated region of mRNAs in the

cytoplasm regulating their stability/degradation and their translatability. Specifically, during

energetic stress this protein blocks global translation while preserving translation of essential

proteins.[52] CPEB4 is involved in muscle regeneration and metabolism and in response to

exercise and caloric restriction.[53,54] Overall, these data indicate that acute endurance exercise

results in a mild response in terms of gene expression changes of RBPs in young organisms;

however, that the response of gene expression changes of RBPs is much more robust in older

individuals.

3.3 SRS identified differential splicing events in acutely exercised young and aged skeletal muscle with and without training

Isoform-switching and DIE analyses in human skeletal muscles following acute exercise were first attempted with IsoformSwitchAnalyzeR using only the SRS data. Firstly, DIE revealed the most common AS event type of each human cohort (Figure S2A,B,C). Isoform switching was also determined: compared to the respective baselines, 10 genes had significant isoform-switching events in hsYS, 15 in hsAS, and 21 in hsAA from an acute bout of exercise (Figure S2D,E). These minimal changes were attributed to the read counts and paired-end versus non-paired end reads of the SRS data sets. Subsequently we choose to utilize an exon splicing data analysis pipeline to glean additional alternative splicing information from the SRS data.

Splicing events at the exon level were analyzed with rMATS, using only the SRS data. The most common event across the three human cohorts was SE followed by RI. The hsYS cohort revealed 67 SE events (Figure 4A); the hsAS revealed 136 SE events (Figure 4B); the hsAA revealed 136 SE events (Figure 4C); all with statistical significance (p-adj < .05). When comparing the ratio of splicing event types between the younger humans and the older humans, it is interesting to note that exercise induced more A5SS and A3SS in older individuals compared to younger, though this could potentially be due to the differences in the exercise protocols between the studies as well. Also, these data illustrate a considerable difference in number of splicing events in skeletal muscle between the younger and aged cohorts (younger cohort had 127 total events compared to greater than 320 events in both aged cohorts). Overall, exercise impacts differential exon usage, especially SE events, whereby age and exercise appear to interact to modify differential exon usage to a greater extent.

To determine if there were specific pathways that were enriched by post-exercise splicing

events as determined by rMATS, we took all the genes with statistically significant differential

splicing events ($\Delta\Psi > 10\%$, p-adj $< .05$) and performed functional enrichment analyses (DAVID

and PANTHER, statistical over-enrichment test, GO molecular function complete, Fisher's exact

test). 115 genes from the hsYS-pre:4pe revealed 1 term, protein binding GO:0005515. The same

test was repeated for hsAS-pre:3pe (52 enriched terms, p-adj $< .05$; Figure S3A) and for hsAA-

pre:3pe (30 enriched terms, p-adj $< .05$; Figure S3B). Protein binding was a term that was in

common across all three groups. GO terms that were common between the hsAS and hsAA

groups were catalytic activity (GO:00033824), binding (GO:0005488), and molecular adaptor

activity (GO:0060090). The major insights we are able to draw from these data are that there is a

different response based on age and training status, which is also displayed above in Figure 3

exploring which RBPs are differentially expressed. These data are in alignment with previous

research.[16,55]

The genes with statistically significant differential splicing events were also compared to

observe any overlaps (Figure 4D). Out of 115, 359, and 287 known genes from the hsYS, hsAS,

and hsAA cohorts, respectively (unique gene IDs; some genes had multiple splicing events, but

each gene ID was only entered once in the overlap analysis), 11 genes overlapped between hsYS

and hsAS, 8 genes between hsYS and hsAA, 40 genes between hsAS and hsAA, and 3 genes

across all. A bivariate hypergeometric test between hsAA and hsAS cohorts revealed significant

over-enrichment compared to expectations (40 genes in common, 1.984-fold over-enrichment, p

$= 2.62e^{-5}$), the other overlapping genes were not statistically significant beyond random chance

from the hypergeometric testing. The three genes with overlapping splicing events between the

human cohorts were ras homolog family member T2 (*RHOT2*), mitochondrial arginyl-tRNA

synthetase 2 (*RARS2*), and AMP-activated protein kinase non-catalytic subunit gamma-2

(*PRKAG2*, γ2-AMPK).

RHOT2 is an outer mitochondrial membrane GTPase that plays a role in mitochondrial

trafficking and fusion-fission dynamics, with little information on the impact of exercise on this

gene or protein product. RARS2 is a mitochondrial tRNA synthetase that plays an important role

in the translation of mitochondrially encoded proteins. There is little research about the role of

this gene related to exercise or skeletal muscle, but mutations in this gene are associated with

neurological disease and lactic acidosis.[56,57] The final overlapping gene, *PRKAG2*, is a part of a

well-known enzyme, AMPK, associated with exercise responses. *PRKAG2* encodes for the non-

catalytic γ2 subunit with 35 transcript variants that can form 12 different protein isoforms, 7

transcripts that are degraded by nonsense-mediated decay, and the remaining either lack on open

reading frame or contain a retained intron that lacks a protein product. This subunit is important

in binding AMP and regulating activity levels of the catalytic subunits.[58] Further mutations in

this gene cause heart issues and metabolic issues in humans.[59,60] In terms of the functional role of

the AS variants that give rise to protein isoforms of γ2-AMPK, little information is published in

the literature. This represents an area ripe for future investigation in the field of exercise

physiology and metabolic health. When we searched to see if the event was the same splicing

event across the cohorts, we found that it was an MXE event in the hsYS cohort, while A3SS in

the older hsAS and hsAA cohorts. This indicates that exercise impacts the splicing of this gene,

but the events shift from an MXE event to an A3SS either due to aging or due to differences in

the exercise protocols. This will have to be determined in future studies. While the γ2-AMPK is

not the major gamma subunit studied or expressed in skeletal muscle, its complex regulation

raises the possibility that this minor subunit plays important functional roles, and warrants future

investigation.

We next wanted to determine how many RBPs had significant exon splicing events post

exercise in the human cohorts. Using our RBPs list (described above) we observed 12 RBPs, 32

RBPs, and 20 RBPs had significant splicing events in hsYS, hsAS, and hsAA, respectively.

Notable RBPs with significant splicing events in the hsYS cohort include heterogeneous nuclear

ribonucleoprotein A2/B1 (hnRNP-A2/B1),[61] RNA-binding Fox-1 homolog 2 (RBFOX2),[62]

RNA-binding motif protein 3 (RBM3),[63] and cold-inducible RNA-binding protein (CIRBP),[64] all

of which have been related to muscle phenotypes in the literature. In the hsAS cohort, some

interesting RBPs with splicing events include several ATP-dependent RNA helicases (DDX17,

DDX23, and DDX5), components of the RNA exosome (EXOSC2, EXOSC3, EXOSC8), and

SRPK2, among others. In the hsAA cohort, CLK4, RBM39, DDX17, and SRPK2 are commonly

studied RBPs with significant splicing events post-exercise. To determine if there were any

common RBPs across the cohorts, we overlapped the significant RBPs. This analysis revealed no

overlapping genes between all three cohorts, and the only pairwise comparison with any

overlapping genes was hsAA with hsAS, which revealed *SRPK2* and *DDX17*. These genes are

interesting as they are both ATP-dependent enzymes with the potential to trigger downstream

impacts on splicing regulation and they may be impacted by acute endurance exercise.

3.4 LRS identified differential gene expression, differential alternative splicing,

and isoform switching in young mice during recovery from acute exercise

Significant DGE and DAS (SE and RI only) of gastrocnemius in mmYS were analyzed

using Swan ($\log_2$FC $\neq$ 0, $\Delta\Pi$ > 10%, raw $p < .05$) to contrast how expressional changes of

transcription and AS differed across the three time points in recovery from acute exercise. Based

on mean normalized counts, mmYS-ipe compared to mmYS-pre illustrated DGE only (313,

63.9%), DAS only (159, 32.4%), both DGE and DAS (18, 3.7%) profiles (Figure 5A); mmYS-

1pe compared to mmYS-pre illustrated DGE only (254, 68.6%) DAS only (39, 10.5%) both

DGE and DAS (77, 20.8%) profiles (Figure 5B); and mmYS-24pe compared to mmYS-pre

illustrated DGE only (254, 65.5%), DAS only (50, 12.9%), both DGE and DAS (84, 21.6%)

profiles (Figure 5C). These data indicate that exercise not only perturbs transcript expression but

also impacts DAS in combination with transcriptional changes and independent of transcriptional

changes. This reveals a new source of exercise-induced responses and adaptations that need to be

explored for their functional relevance in the physiology of skeletal muscle and in their roles in

exercise-induced health benefits.

Isoform switching (i.e., differential isoform expression, DIE) in gastrocnemius of mmYS-

ipe, mmYS-1pe, and mmYS-24pe compared to mmYS-pre was analyzed with Swan. A total of

609, 1010, and 806 transcripts had significant isoform switching events ($\Delta\Pi > 10\%$, p-adj $< .05$)

at each timepoint, respectively (Figure 5D). We investigated our isoform switching data for

known exercise associated genes. We generated a list of 60 well-established exercise-associated

genes and overlapped with the list of genes that had significant isoform-switching events at each

time point. We observed isoform switching in 18, 24, and 21 of those 60 genes in ipe, 1pe, and

24pe, respectively. According to hypergeometric testing each of these overlapping gene sets is

highly over enriched compared to random expectations (ipe, 20.3-fold over-enrichment, $p = 5.2e^-$

^{19}; 1pe, 17.6-fold over-enrichment, $p = 4.6e^{-24}$; and 24pe, 17.4-fold over-enrichment, $p = 6.5e^{-21}$).

Genes including sirtuin 2 (*mSirt2*), myosin heavy chain IIb (*mMyh4*, MHC-IIb), *mNr4a1*, and

myocyte-specific enhancer factor 2C (*mMef2c*) were observed to have significant DIE across all

three time points post exercise (there were 13 genes in common across all three groups).

We used RT-PCR to validate the splicing event observed in *mSirt2*. The LRS data

showed that there was an exon skipping event of exon 2 of Ensembl transcript *mSirt2*-201

resulting in a 22% increase in *mSirt2*-202 at 24pe (Figure 5F). We designed primers for exons 1

and 3 of *mSirt2*-201 and we were able to quantify exon skipping. When exon 2 of *mSirt2*-201 is

skipped this produces *mSirt2*-202. We were able to show that there were no changes in transcript

levels of *mSirt2*-201 at 24pe compared to pre (14.6%, p = n.s.); however, there was a significant

increase in *mSirt2*-202 at the same time points (24.5% p = 0.005) validating the isoform-

switching event post-exercise (Figure 5G). We queried if *SIRT2* had a significant exon-level

splicing event post-exercise in the rMATS data of the human cohorts, but did not observe any

significant events. This could be due to the different timing post-exercise sample collection (3-4

hours for the humans compared to 24 hours for the mice). There is limited data on the functional

differences between *mSirt2* transcripts 201 and 202. Future studies would be required to

determine the function and regulation of the exon 2 skipping event. Overall, our *mSirt2* data

indicate that genes commonly associated with exercise responses and adaptations undergo

isoform-switching events that may be important to investigate their physiological relevance in

future mechanistic follow-up studies.

Furthermore, 144 (10.1%) overlapped between mmYS-ipe and mmYS-1pe, 103 (7.2%)

between mmYS-ipe and mmYS-24pe, 230 (16.2%) between mmYS-1pe and mmYS-24pe, and

262 (18.4%) across all three groups (Figure 6A). When unique gene identifiers of significant

isoform-switching genes at each time point (as described above) were queried for enriched

pathways using PANTHER, this revealed several common GO terms including RNA binding

(top 10 GO terms are shown in Figure S4A,B,C). When the 262 common genes across time

points post-exercise were independently queried for enriched pathways, revealing enrichment of

GO terms including RNA binding (GO:0003723), mRNA 3'-untranslated region (UTR) binding

(GO:0003730), transmembrane signaling receptor activity (GO:0004888), and molecular

transducer activity (GO:0060089) (top 10 GO terms are shown in Figure 6B). Within these 262

common genes, 61 were RBPs as identified by overlapping the 262 gene identifiers with a list of

1130 RBP genes generated in house (Figure 6C) and hypergeometric testing revealed that this

was significantly over-enriched compared to expectations (24.4-fold over-enrichment, $p = 5.1e^{-}$

^{69}).

Several top candidate RBP genes, heterogeneous nuclear ribonucleoprotein A3

(*mHnrnpa3*), A1 (*mHnrnpa1*), and T-cell intracellular antigen 1 (*mTia1*) were chosen for

validation due to the abundance of each gene, and ability to design RT-PCR primers to detect the

splicing event. The LRS data showed that there was a significant isoform-switching event of $\Delta\Pi$

= 10% increase in the protein-coding transcripts at 24pe (Figure 6E). We designed primers for

exons 3 and 4 of *mHnrnpa3* (to detect transcripts 203 and 204) and intron 3 to quantify intron

retention of *mHnrnpa3*-205 (non-coding RNA) (Figure 6D). RT-ddPCR showed that there was

no change in transcript levels of *mHnrnpa3*-205 at 24pe compared to pre (*t*-test p = n.s.);

however, there was a significant increase in protein-coding *mHnrnpa3* at the same time points

(28.5%, *t*-test $p = 0.02$) validating the isoform-switching event post-exercise (Figure 6F).

Western blots showed that there were significant increases of hnRNP-A3 at the protein level

post-exercise (pre:ipe, 23.1%, *p*-adj = n.s.; pre:1pe, 95.6%, *p*-adj = 0.008; pre: 24pe, 84.9%, *p*-

adj = 0.02; Figure 6G). The LRS data also showed that there was a significant DIE event in

mHnrnpa1, resulting in an overall 24% increase in *mHnrnpa1*-202 (SE) and *mHnrnpa1*-203

(nonsense-mediated decay) transcripts at 24pe (Figure 6H). RT-PCR showed significant

increases in *mHnrnpa1*-202 and *mHnrnpa1* nonsense-mediated decay products at 24pe (30.6%,

t-test $p = 0.004$; Figure 6I). The LRS detected a significant DIE event in *mTia1* as well, resulting

in an overall 40% increase in *mTia1*-202 (SE) transcripts at 24pe (Figure 6J), validated by RT-

PCR of the 202 transcripts at 24pe (53.6%, *t*-test $p = 0.004$; Figure 6K).

4 DISCUSSION

The present study supports that, in addition to DGE in recovery from endurance exercise, there is a robust AS response across the transcriptome in skeletal muscles from both humans and mice. Several insights from our data include the following: 1) post-exercise recovery of skeletal muscle impacts AS; 2) the pathways enriched in splicing events appear to be related to splicing regulation (i.e., RBPs) among several other pathways; 3) common exercise-associated genes are impacted by AS post-exercise, such as *PRKAG2* (γ2-AMPK) and *SIRT2*; 4) several of the common RBPs observed to have either DGE or DAS post-exercise are related to 3'-untranslated region binding (CPEB4, hnRNP-A1, and TIA1) and transcript stability; 5) there appears to be a difference in how AS is regulated by aging with older individuals having high RBP expression and more splicing changes; and 6) our mouse LRS data revealed that more splicing events may be significant 24 hours post-exercise, compared to transcriptional events being highly enriched at earlier timepoints.

Our data indicate that while some RBPs and spliceosome proteins undergo DGE in response to an acute bout of endurance exercise in untrained skeletal muscle, far more RBPs and spliceosome proteins undergo DAS and isoform switching in response to exercise. Our rMATS analysis of SRS data revealed that young individuals have splicing events in response to endurance exercise, but both sedentary and trained older individuals have a greater number of significant events compared to younger individuals. While SRS is excellent for detecting DGE and exon splicing events, it lacks the resolution to accurately detect transcript variants and/or isoform-switching events due to multi-read alignment and mapping ambiguity.[65] Finally, our data clearly points out the strengths of LRS for the detection of full-length transcript variants and

opens an entirely new area of adaptive potential in skeletal muscle to explore in response to

endurance exercise.

Recent literature has indicated changes in the expression of transcript variants with aging

in skeletal muscle.[16,55,66–69] Since endurance exercise is a potent anti-aging intervention (i.e., by

maintaining V̇O$_2$max), we surmised that the response of skeletal muscle gene expression at the

level of AS would differ by age and training status. Our data clearly indicate that this is likely the

case (Figure 3,4). First, since previous research has shown significant RNA and protein level

differences in RBPs with age, we explored a biased DGE analysis of RBPs. This revealed that

acute endurance exercise in young individuals had a minimal impact on DGE of RBPs in

comparison to the effect acute endurance exercise in muscles of older individuals (Figure 3).

Interestingly, older trained individuals had fewer differentially expressed RBPs in response to

acute exercise compared to acute exercise in sedentary individuals (Figure 3). This finding is in

line with recently published data that indicated lower RBP expression levels in masters athletes

compared to age-matched non-athletes.[70,71]

Becoming or remaining physically active, as well as performing regular exercise, shifts to

play a more important role in health as one ages than during their youth. Recent findings have

indicated that AS becomes dysregulated with aging in several tissues, including skeletal

muscle.[16,71,72] When aged skeletal muscles were compared to younger muscles, proteins of the

spliceosome and several RBPs were overrepresented, leading to a change in the expressed

variants of mRNA.[16] Interestingly, when octogenarian masters athletes were compared to age-

matched non-athletes, the athletes had lower protein expression of the spliceosomal and several

RBPs.[70] The authors went on to hypothesize that the underrepresentation of the spliceosomal

proteins was related to the higher functioning mitochondria and the lower need for energy to be

expended for splice-isoform generation.[70] Thus, further research is needed to link exercise, AS

regulation, and mRNA variant generation to the adaptive process in skeletal muscle of both

younger and older adults.

Next, we used rMATS to explore the type and number of AS events at the exon level

from the SRS from humans (Figure 4). To our knowledge, rMATS has not previously been used

to explore how exercise may modify AS, allowing us to make several novel insights. First, it is

apparent that SE is the dominant form of AS event category in response to an acute bout of

exercise while A5SS and A3SS are potentially subjected to other factors, such as age. The

second most abundant type of splicing event across all data sets was RI (Figure 4). Both SE and

RI events have important implications on the transcriptome and proteome that should be

explored in future studies, but these data provide novel mechanistic insights into how acute

endurance exercise is modifying gene expression and potentially skeletal muscle responses and

adaptations to exercise. Second, we observed that there were fewer splicing events in the muscles

of younger individuals compared to older individuals in response to exercise. However, we

interpret this cautiously due to technical limitations and differences with how the samples were

collected and the way the sequencing was performed between the data sets. Future studies will be

needed to test this important question more carefully. A third observation is that older skeletal

muscles utilized more A5SS and A3SS compared to younger muscles in response to a bout of

endurance exercise. This could be due to the different responses of signal transduction pathways

to modify the splicing machinery in older muscles compared to younger muscles. Alternatively,

this could indicate greater energetic stress in older muscles leading to different signaling

pathways being activated, resulting in the utilization of weaker splice sites. This could have

important ramifications on the transcriptome and proteome following exercise in older muscles

and should be explored in more detail in future research.[73,74]

While SRS is a powerful technology and our data above illustrate its ability to detect

DGE and exon level splicing events, it lacks the ability to detect full-length transcripts, which

have great implications on the expression of protein isoforms. Thus, we wanted to determine if

additional insights could be made concerning how exercise mechanistically impacts skeletal

muscle by using LRS. First, we were able to separate DGE and DAS using the LRS data (Figure

5A,B,C). This revealed that a significant portion of genes only undergo DAS, meaning previous

post-exercise DGE analyses would not have detected these changes even though they would

have likely impacted the post-exercise protein levels in muscle. Next, we analyzed isoform

switching, specifically in genes that were not differentially expressed following exercise. We

observed a robust isoform-switching response at all three time points in recovery from endurance

exercise (Figure 5D). Further analysis illuminated a significant enrichment of RBPs that undergo

isoform-switching events during recovery from endurance exercise (Figure 6A,B,C). In other

words, RBPs and splicing factors undergo regulatory alteration by genes encoding these proteins

via AS. Such regulatory process is referred to as either auto-regulation (whereby a RBP regulates

the splicing of its own transcripts) or cross-regulation (whereby a RBP regulates the splicing of

other RBP-coding transcripts). This idea of auto- and cross-regulation as a means of regulating

functional levels of RBPs is not a novel concept in the field of RNA biology, as previously

demonstrated in neuronal (NOVA1)[75] and cancer (PTBP1/2 and SRSF3)[76] cells.

Interestingly, several of the isoform-switching events lasted at least 24 hours into

recovery from exercise, indicating a lasting (i.e., not transient) re-wiring of the transcriptome that

likely impacts the proteome as we have shown is the case for hnRNP-A3 (Figure 6D–G).

Splicing and AS are regulated by a multitude of factors including (but not limited to)

transcriptional rate, chromatin structure, the spliceosome, and RBPs.[4,5] The potential for exercise

to modify the activity of the spliceosome, as well as post-translational modification of RBPs

which may impact binding to their respective target RNAs (and thus modify their splicing), is

particularly important. RBPs bind to RNAs and impact their entire lifecycle: from splicing to

localization, to translation, and degradation.[18] Specifically, RBPs bind to *cis*-regulatory elements

(i.e., intronic/exonic splicing enhancers and silencers) to regulate the inclusion of specific exons,

resulting in the expression of different transcript variants and protein isoforms. Differential use

of isoforms between conditions (isoform switching) can have profound biological impacts, even

if they are only transiently induced by exercise, due to the difference in functions of the isoforms.

Functional consequences of isoform switches can include gain or loss of protein domains or

signal peptides, loss of protein coding sequence, and change in propensity for nonsense-mediated

decay.[13] Isoform switches are implicated in many diseases, and it has already been observed

following exercise (PGC-1α)[15]; however, the full extent and impact of isoform switching post-

exercise is far from understood.

It is highly likely, but not yet tested, that exercise remodels the binding patterns of RBPs

either through post-translational modifications of the RBPs themselves or through isoform

switching and AS changes to RBPs themselves, which would modify the target genes of RBPs.

Metabolic stress responsive pathways, such as AMPK, are already known to regulate post-

translation modifications of RBPs.[21] Interestingly, many RBPs tend to regulate their protein

abundance through AS and isoform switching. Our isoform switching data indicate an

enrichment of events in RBPs post-exercise. This indicates that exercise may remodel the

expression of protein isoforms by modifying the function of RBPs. Our findings not only support

the preceding study from our laboratory that focused on SF3B4 and SRSF2,[30] but others that

have observed similar phenomenon in a separate setting: Henrich et al.[77] illustrated 9 weeks of

microgravity conditions leading to AS-driven diversification of the transcriptome in mouse

skeletal muscle, including RBPs (e.g., *mMbnl1* and *mRbfox1*). This mechanism is important in

shaping the transcriptome and proteome in response to specific cellular stresses. Bringing this

idea to the study of skeletal muscle adaptations in response to exercise is an innovative and novel

molecular mechanism of adaptation (Figure 7).

The present study is not without limitations. The human SRS data utilized for this study

included on average 20 million reads per sample, which is insufficient/minimum threshold to

yield biologically true isoform-switching analyses with IsoformSwitchAnalyzeR. Hence, there is

a growing need for employing LRS technologies to better examine alternative splicing and

isoform switching in the field of exercise biology. Additionally, the mice cohort had mixed sexes

unlike the human cohorts utilized for analyses, which may further limit the comparability. Indeed,

protein analysis was only conducted for one of the RBPs, hnRNP-A3, but not at the global level

for this study. To better validate the significance of DAS and isoform-switching events, future

studies will incorporate proteomics data with LRS data via proteogenomic approaches.

In summary, the present data support recent evidence of divergence in the skeletal muscle

AS-related responses and RBP expression by age and training status. Moreover, LRS is the

preferrable method of transcriptomics study when exploring AS and isoform-switching events,

whereas utilizing SRS may require cost prohibitive read depth per sample for many groups (at

least 100 million reads per sample) to proficiently conduct the isoform switching analyses.

Additionally, our acute endurance exercise mouse model demonstrated that, even without any

changes in overall gene expression, the protein expression of hnRNP-A3 was increased by nearly

two-fold in response to exercise for at least 24 hours via isoform switching with the power of

LRS. Therefore, our study points out that AS following exercise in skeletal muscle is an

underexplored and untapped area, ripe for future studies to increase our mechanistic

understanding of exercise.

IV: DATA AVAILABILITY STATEMENT

The human SRS data were extracted from deposits by Pattamaprapanont et al.[31] and Rubenstein

et al.[32] in the NCBI Gene Expression Omnibus database with the accession numbers GSE87749

and GSE151066, respectively. The mouse LRS data are deposited in the NCBI Gene Expression

Omnibus database with the accession number GSE279359. All data used to generate the figures

in this manuscript will be deposited at the Deep Blue Data repository with a permanent DOI

(DOI will be provided upon publication).

V: CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest. The authors

declare no competing interests.

VI: AUTHOR CONTRIBUTIONS

AA, ALS, JJK, and ATL conceived and/or designed the research. AA, ALS, JJK performed the

research and acquired the data, AA, JJK, JYY, EYZ, and ATL analyzed and interpreted the data.

All authors were involved in drafting and revising the manuscript.

VII: ACKNOWLEDGMENTS

This research was supported by the National Institutes of Health and National Cancer Institute

under award number 5R00CA197672-04, Michigan Integrative Musculoskeletal Health Core

Center (MiMHC) P30 grant from NIH/NIAMS under award number P30 AR069620, and the

University of Michigan's School of Kinesiology Marie Hartwig Research Fund.

VIII: REFERENCES

1.   CDC. Adult Physical Inactivity. Centers for Disease Control and Prevention. November 20, 2023. Accessed February 3, 2024. https://www.cdc.gov/physicalactivity/data/inactivity-prevalence-maps/index.html

2.   Lee IM, Shiroma EJ, Lobelo F, et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet*. 2012;380(9838):219-229. doi:10.1016/S0140-6736(12)61031-9

3.   Holloszy JO, Smith EK, Vining M, Adams S. Effect of voluntary exercise on longevity of rats. *Journal of Applied Physiology*. 1985;59(3):826-831. doi:10.1152/jappl.1985.59.3.826

4.   Tomasek JJ, Gabbiani G, Hinz B, Chaponnier C, Brown RA. Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol*. 2002;3(5):349-363. doi:10.1038/nrm809

5.   Ferraro E, Giammarioli AM, Chiandotto S, Spoletini I, Rosano G. Exercise-Induced Skeletal Muscle Remodeling and Metabolic Adaptation: Redox Signaling and Role of Autophagy. *Antioxid Redox Signal*. 2014;21(1):154-176. doi:10.1089/ars.2013.5773

6.   Powers SK, Talbert EE, Adhihetty PJ. Reactive oxygen and nitrogen species as intracellular signals in skeletal muscle. *J Physiol*. 2011;589(Pt 9):2129-2138. doi:10.1113/jphysiol.2010.201327

7.   Egan B, Sharples AP. Molecular responses to acute exercise and their relevance for adaptations in skeletal muscle to exercise training. *Physiological Reviews*. 2023;103(3):2057-2170. doi:10.1152/physrev.00054.2021

8.   Pillon NJ, Gabriel BM, Dollet L, et al. Transcriptomic profiling of skeletal muscle

adaptations to exercise and inactivity. *Nat Commun*. 2020;11(1):470. doi:10.1038/s41467-019-

13869-w

9.   Amar D, Lindholm ME, Norrbom J, Wheeler MT, Rivas MA, Ashley EA. Time

trajectories in the transcriptomic response to exercise - a meta-analysis. *Nat Commun*.

2021;12(1):3471. doi:10.1038/s41467-021-23579-x

10.   Mathai AS, Bonen A, Benton CR, Robinson DL, Graham TE. Rapid exercise-induced

changes in PGC-1α mRNA and protein in human skeletal muscle. *Journal of Applied Physiology*.

2008;105(4):1098-1105. doi:10.1152/japplphysiol.00847.2007

11.   Gustafsson T, Puntschart A, Kaijser L, Jansson E, Sundberg CJ. Exercise-induced

expression of angiogenesis-related transcription and growth factors in human skeletal muscle.

American Journal of Physiology-Heart and Circulatory Physiology. 1999;276(2):H679-H685.

doi:10.1152/ajpheart.1999.276.2.H679

12.   Gahlmann R, Troutt AB, Wade RP, Gunning P, Kedes L. Alternative splicing generates

variants in important functional domains of human slow skeletal troponin T. *Journal of*

Biological Chemistry. 1987;262(33):16122-16126. doi:10.1016/S0021-9258(18)47705-8

13.   Norrbom J, Sällstedt EK, Fischer H, Sundberg CJ, Rundqvist H, Gustafsson T.

Alternative splice variant PGC-1α-b is strongly induced by exercise in human skeletal muscle.

American Journal of Physiology-Endocrinology and Metabolism. 2011;301(6):E1092-E1098.

doi:10.1152/ajpendo.00119.2011

14.   Jensen L, Pilegaard H, Neufer PD, Hellsten Y. Effect of acute exercise and exercise

training on VEGF splice variants in human skeletal muscle. *Am J Physiol Regul Integr Comp*

Physiol. 2004;287(2):R397-402. doi:10.1152/ajpregu.00071.2004

15.   Zhang T, Choi SJ, Wang ZM, et al. Human Slow Troponin T (TNNT1) Pre-mRNA

Alternative Splicing Is an Indicator of Skeletal Muscle Response to Resistance Exercise in Older

Adults. *The Journals of Gerontology: Series A*. 2014;69(12):1437-1447.

doi:10.1093/gerona/glt204

16.   Ubaida-Mohien C, Gonzalez-Freire M, Lyashkov A, et al. Physical Activity Associated

Proteomics of Skeletal Muscle: Being Physically Active in Daily Life May Protect Skeletal

Muscle From Aging. *Front Physiol*. 2019;10:312. doi:10.3389/fphys.2019.00312

17.   Innocenzi E, Cariati I, De Domenico E, et al. Aerobic Exercise Induces Alternative

Splicing of Neurexins in Frontal Cortex. *J Funct Morphol Kinesiol*. 2021;6(2):48.

doi:10.3390/jfmk6020048

18.   Gaugel J, Jähnert M, Neumann A, Heyd F, Schürmann A, Vogel H. Alternative splicing

landscape in mouse skeletal muscle and adipose tissue: Effects of intermittent fasting and

exercise. *The Journal of Nutritional Biochemistry*. 2025;137:109837.

doi:10.1016/j.jnutbio.2024.109837

19.   Wahl MC, Will CL, Lührmann R. The Spliceosome: Design Principles of a Dynamic

RNP Machine. *Cell*. 2009;136(4):701-718. doi:10.1016/j.cell.2009.02.009

20.   Tao Y, Zhang Q, Wang H, Yang X, Mu H. Alternative splicing and related RNA binding

proteins in human health and disease. *Sig Transduct Target Ther*. 2024;9(1):1-33.

doi:10.1038/s41392-024-01734-2

21.   Anczukow O, Allain FHT, Angarola BL, et al. Steering research on mRNA splicing in

cancer towards clinical translation. *Nat Rev Cancer*. Published online October 9, 2024:1-19.

doi:10.1038/s41568-024-00750-2

22.   Tao Y, Zhang Q, Wang H, Yang X, Mu H. Alternative splicing and related RNA binding

proteins in human health and disease. *Sig Transduct Target Ther*. 2024;9(1):1-33.

doi:10.1038/s41392-024-01734-2

23.   Aebersold R, Agar JN, Amster IJ, et al. How many human proteoforms are there? *Nat*

Chem Biol. 2018;14(3):206-214. doi:10.1038/nchembio.2576

24.   Manuel JM, Guilloy N, Khatir I, Roucou X, Laurent B. Re-evaluating the impact of

alternative RNA splicing on proteomic diversity. *Frontiers in Genetics*. 2023;14. Accessed

February 9, 2024.

https://www.frontiersin.org/journals/genetics/articles/10.3389/fgene.2023.1089053

25.   Dinger ME, Amaral PP, Mercer TR, et al. Long noncoding RNAs in mouse embryonic

stem cell pluripotency and differentiation. *Genome Res*. 2008;18(9):1433-1445.

doi:10.1101/gr.078378.108

26.   Khan MR, Avino M, Wellinger RJ, Laurent B. Distinct regulatory functions and

biological roles of lncRNA splice variants. *Molecular Therapy - Nucleic Acids*. 2023;32:127-143.

doi:10.1016/j.omtn.2023.03.004

27.   Saudemont B, Popa A, Parmley JL, et al. The fitness cost of mis-splicing is the main

determinant of alternative splicing patterns. *Genome Biol*. 2017;18(1):208. doi:10.1186/s13059-

017-1344-6

28.   García-Moreno JF, Romão L. Perspective in Alternative Splicing Coupled to Nonsense-

Mediated mRNA Decay. *Int J Mol Sci*. 2020;21(24):9424. doi:10.3390/ijms21249424

29.   Tonevitsky EA, Trushkin EV, Shkurnikov MU, Akimov EB, Sakharov DA. Changed

Profile of Expression of Splicing Regulator Genes in Response to Exercise. *Bull Exp Biol Med*.

2009;147(6):733-736. doi:10.1007/s10517-009-0593-0

30.   Slusher AL, Kim JJ, Ribick M, Ludlow AT. Acute Exercise Regulates hTERT Gene

Expression and Alternative Splicing in the hTERT-BAC Transgenic Mouse Model. *Med Sci*

Sports Exerc. 2022;54(6):931-943. doi:10.1249/mss.0000000000002868

31.   Pattamaprapanont P, Garde C, Fabre O, Barrès R. Muscle Contraction Induces Acute

Hydroxymethylation of the Exercise-Responsive Gene Nr4a3. *Front Endocrinol (Lausanne)*.

2016;7:165. doi:10.3389/fendo.2016.00165

32.   Rubenstein AB, Hinkley JM, Nair VD, et al. Skeletal muscle transcriptome response to a

bout of endurance exercise in physically active and sedentary older adults. *American Journal of*

Physiology-Endocrinology and Metabolism. 2022;322(3):E260-E277.

doi:10.1152/ajpendo.00378.2021

33.   Fernando P, Bonen A, Hoffman-Goetz L. Predicting submaximal oxygen consumption

during treadmill running in mice. *Can J Physiol Pharmacol*. 1993;71(10-11):854-857.

doi:10.1139/y93-128

34.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. *BMC Bioinformatics*. 2011;12(1):323. doi:10.1186/1471-2105-12-

323

35.   Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner.

Bioinformatics. 2013;29(1):15-21. doi:10.1093/bioinformatics/bts635

36.   Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*.

2018;34(18):3094-3100. doi:10.1093/bioinformatics/bty191

37.   Danecek P, Bonfield JK, Liddle J, et al. Twelve years of SAMtools and BCFtools.

GigaScience. 2021;10(2):giab008. doi:10.1093/gigascience/giab008

38.   Wyman D, Mortazavi A. TranscriptClean: variant-aware correction of indels, mismatches

and splice junctions in long-read transcripts. *Bioinformatics*. 2019;35(2):340-342.

doi:10.1093/bioinformatics/bty483

39.   Wyman D, Balderrama-Gutierrez G, Reese F, et al. A technology-agnostic long-read

analysis pipeline for transcriptome discovery and quantification. Published online March 24,

2020:672931. doi:10.1101/672931

40.   Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences. Published online February 29, 2016.

doi:10.12688/f1000research.7563.2

41.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biology*. 2014;15(12):550. doi:10.1186/s13059-014-0550-

8

42.   Vitting-Seerup K, Sandelin A. IsoformSwitchAnalyzeR: analysis of changes in genome-

wide patterns of alternative splicing and its functional consequences. *Bioinformatics*.

2019;35(21):4469-4471. doi:10.1093/bioinformatics/btz247

43.   Shen S, Park JW, Lu Z xiang, et al. rMATS: Robust and flexible detection of differential

alternative splicing from replicate RNA-Seq data. *Proc Natl Acad Sci U S A*.

2014;111(51):E5593-E5601. doi:10.1073/pnas.1419161111

44.   Wang Y, Xie Z, Kutschera E, Adams JI, Kadash-Edmondson KE, Xing Y. rMATS-turbo:

an efficient and flexible computational tool for alternative splicing analysis of large-scale RNA-

seq data. *Nat Protoc*. Published online February 23, 2024:1-22. doi:10.1038/s41596-023-00944-

2

45.   Reese F, Mortazavi A. Swan: a library for the analysis and visualization of long-read

transcriptomes. *Bioinformatics*. 2021;37(9):1322-1323. doi:10.1093/bioinformatics/btaa836

46.   Sherman BT, Hao M, Qiu J, et al. DAVID: a web server for functional enrichment

analysis and functional annotation of gene lists (2021 update). *Nucleic Acids Research*.

2022;50(W1):W216-W221. doi:10.1093/nar/gkac194

47.   Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albou LP, Mi H. PANTHER:

Making genome-scale phylogenetics accessible to all. *Protein Science*. 2022;31(1):8-22.

doi:10.1002/pro.4218

48.   Gerstberger S, Hafner M, Tuschl T. A census of human RNA-binding proteins. *Nat Rev*

Genet. 2014;15(12):829-845. doi:10.1038/nrg3813

49.   Noble EG, Shen GX. Impact of Exercise and Metabolic Disorders on Heat Shock

Proteins and Vascular Inflammation. *Autoimmune Dis*. 2012;2012:836519.

doi:10.1155/2012/836519

50.   Parker BL, Kiens B, Wojtaszewski JFP, Richter EA, James DE. Quantification of

exercise-regulated ubiquitin signaling in human skeletal muscle identifies protein modification

cross talk via NEDDylation. *The FASEB Journal*. 2020;34(4):5906-5916.

doi:10.1096/fj.202000075R

51.   Lord SO, Lai Y. Exercise mediates ubiquitin signalling in human skeletal muscle. *FASEB*

Bioadv. 2022;4(6):402-407. doi:10.1096/fba.2021-00142

52.   Afroz T, Skrisovska L, Belloc E, Guillén-Boixet J, Méndez R, Allain FHT. A fly trap

mechanism provides sequence-specific RNA recognition by CPEB proteins. *Genes Dev*.

2014;28(13):1498-1514. doi:10.1101/gad.241133.114

53.   Zeng W, Zhang W, Tse EHY, et al. Restoration of CPEB4 prevents muscle stem cell

senescence during aging. *Developmental Cell*. 2023;58(15):1383-1398.e6.

doi:10.1016/j.devcel.2023.05.012

54.   Kang JS, Kim MJ, Kwon ES, et al. Identification of novel genes associated with exercise

and calorie restriction effects in skeletal muscle. *Aging (Albany NY)*. 2023;15(11):4667-4684.

doi:10.18632/aging.204793

55.   Ubaida-Mohien C, Lyashkov A, Gonzalez-Freire M, et al. Discovery proteomics in aging

human skeletal muscle finds change in spliceosome, immunity, proteostasis and mitochondria.

Kaeberlein M, Rosen CJ, Kaeberlein M, Anderson R, eds. *eLife*. 2019;8:e49874.

doi:10.7554/eLife.49874

56.   Lühl S, Bode H, Schlötzer W, et al. Novel homozygous RARS2 mutation in two siblings

without pontocerebellar hypoplasia – further expansion of the phenotypic spectrum. *Orphanet*

Journal of Rare Diseases. 2016;11(1):140. doi:10.1186/s13023-016-0525-9

57. Mathew T, Avati A, D'Souza D, Therambil M. Expanding spectrum of RARS2 gene

disorders: Myoclonic epilepsy, mental retardation, spasticity, and extrapyramidal features.

Epilepsia Open. 2018;3(2):270-275. doi:10.1002/epi4.12108

58. Cheung PC, Salt IP, Davies SP, Hardie DG, Carling D. Characterization of AMP-

activated protein kinase gamma-subunit isoforms and their role in AMP binding. *Biochem J*.

2000;346(Pt 3):659-669.

59. Yavari A, Bellahcene M, Bucchi A, et al. Mammalian γ2 AMPK regulates intrinsic heart

rate. *Nat Commun*. 2017;8:1258. doi:10.1038/s41467-017-01342-5

60. Yavari A, Stocker CJ, Ghaffari S, et al. Chronic Activation of γ2 AMPK Induces Obesity

and Reduces β Cell Function. *Cell Metab*. 2016;23(5):821-836. doi:10.1016/j.cmet.2016.04.003

61. Wheeler JR, Whitney ON, Vogler TO, et al. RNA-binding proteins direct myogenic cell

fate decisions. Black DL, Manley JL, eds. *eLife*. 2022;11:e75844. doi:10.7554/eLife.75844

62. Wiedner HJ, Blue RE, Sadovsky M, et al. RBFOX2 regulated EYA3 isoforms partner

with SIX4 or ZBTB1 to control transcription during myogenesis. *iScience*. 2023;26(11):108258.

doi:10.1016/j.isci.2023.108258

63. Hettinger ZR, Confides AL, Vanderklish PW, Dupont□Versteegden EE. The transcript

interactome of skeletal muscle RNA binding protein motif 3 (RBM3). *Physiol Rep*.

2023;11(3):e15596. doi:10.14814/phy2.15596

64. Chazarin B, Ziemianin A, Evans AL, et al. Limited Oxidative Stress Favors Resistance to

Skeletal Muscle Atrophy in Hibernating Brown Bears (Ursus Arctos). *Antioxidants (Basel)*.

2019;8(9):334. doi:10.3390/antiox8090334

65.   Slotkin RK. The case for not masking away repetitive DNA. *Mob DNA*. 2018;9:15.

doi:10.1186/s13100-018-0120-9

66.   Solovyeva EM, Ibebunjo C, Utzinger S, et al. New insights into molecular changes in

skeletal muscle aging and disease: Differential alternative splicing and senescence. *Mechanisms*

of Ageing and Development. 2021;197:111510. doi:10.1016/j.mad.2021.111510

67.   Rodríguez SA, Grochová D, McKenna T, et al. Global genome splicing analysis reveals

an increased number of alternatively spliced genes with aging. *Aging Cell*. 2016;15(2):267-278.

doi:10.1111/acel.12433

68.   Tumasian RA, Harish A, Kundu G, et al. Skeletal muscle transcriptome in healthy aging.

Nat Commun. 2021;12(1):2014. doi:10.1038/s41467-021-22168-2

69.   Lee BP, Pilling LC, Emond F, et al. Changes in the expression of splicing factor

transcripts and variations in alternative splicing are associated with lifespan in mice and humans.

Aging Cell. 2016;15(5):903-913. doi:10.1111/acel.12499

70.   Ubaida-Mohien C, Spendiff S, Lyashkov A, et al. Unbiased proteomics, histochemistry,

and mitochondrial DNA copy number reveal better mitochondrial health in muscle of high-

functioning octogenarians. Grano M, Isales C, Piasecki M, Degens H, eds. *eLife*.

2022;11:e74335. doi:10.7554/eLife.74335

71.   Roberts MD, Ruple BA, Godwin JS, et al. A novel deep proteomic approach in human

skeletal muscle unveils distinct molecular signatures affected by aging and resistance training.

Aging (Albany NY). 2024;16(8):6631-6651. doi:10.18632/aging.205751

72.   Angarola BL, Anczuków O. Splicing alterations in healthy aging and disease. *Wiley*

Interdiscip Rev RNA. 2021;12(4):e1643. doi:10.1002/wrna.1643

73.   Bartz J, Jung H, Wasiluk K, Zhang L, Dong X. Progress in Discovering Transcriptional

Noise in Aging. *International Journal of Molecular Sciences*. 2023;24(4):3701.

doi:10.3390/ijms24043701

74.   Perez-Gomez A, Buxbaum JN, Petrascheck M. The aging transcriptome: read between

the lines. *Current Opinion in Neurobiology*. 2020;63:170-175. doi:10.1016/j.conb.2020.05.001

75.   Dredge BK, Stefani G, Engelhard CC, Darnell RB. Nova autoregulation reveals dual

functions in neuronal splicing. *EMBO J*. 2005;24(8):1608-1620. doi:10.1038/sj.emboj.7600630

76.   Guo J, Jia J, Jia R. PTBP1 and PTBP2 impaired autoregulation of SRSF3 in cancer cells.

Sci Rep. 2015;5(1):14548. doi:10.1038/srep14548

77.   Henrich M, Ha P, Wang Y, et al. Alternative splicing diversifies the skeletal muscle

transcriptome during prolonged spaceflight. *Skelet Muscle*. 2022;12:11. doi:10.1186/s13395-

022-00294-9

IX: FIGURE LEGENDS

Figure 1. Layout of RNA-Seq study. (A) Publicly accessible human SRS data (hsYS, hsAS, and hsAA) were downloaded from NCBI Gene Expression Omnibus, preprocessed, and analyzed for differential expression. (B) Young C57BL/6 mice (mmYS) underwent an acute bout (30 minutes) of treadmill running, sacrificed, and their gastrocnemius skeletal muscle tissues were harvested for RNA sequencing cDNA library prep. Following LRS, the data were preprocessed and analyzed for differential expression. Biologically significant genes from (A) and (B) were retrieved for subsequent pathway and biased analyses.

Figure 2. Differential gene expression (DGE) of hsYS (A), hsAS (B,C), hsAA (D,E), and mmYS (F,G,H) in the skeletal muscle comparing pre- and post-acute exercise, with LRS validated using droplet digital RT-PCR (RT-ddPCR) validations of LRS DGE data (I,J,K). (A) Volcano plot of hsYS-pre and hsYS-4pe DGE. (B) Volcano plot of hsAS-pre and hsAS-3pe DGE. (C) Top 10 GO terms from significantly up-regulated genes of hsAS-pre and hsAS-3pe DGE. (D) Volcano plot of hsAA-pre and hsAA-3pe DGE. (E) Top 10 GO terms from significantly up-regulated genes of hsAA-pre and hsAA-3pe DGE. (F) Volcano plot of mmYS-pre and mmYS-ipe DGE. (G) Volcano plot of mmYS-pre and mmYS-1pe DGE. (H) Volcano plot of mmYS-pre and mmYS-24pe DGE. (I) DGE of mmYS-pre and mmYS-ipe was validated with *mUbe2d1* using RT-ddPCR. (J) DGE of mmYS-pre and mmYS-1pe was validated with *mFbxo32* using RT-ddPCR. (K) DGE of mmYS-pre and mmYS-24pe was validated with *mFos* using RT-ddPCR. *, $p < 0.05$ compared to mmYS-pre. **, $p < 0.01$ compared to mmYS-pre. All experiments were conducted with $n = 5$ per cohort.

Figure 3. 89 RBPs with significant DGE ($|\log_2 FC| > 0.5$; raw $p < .05$ for mouse, p-adj $< .05$ for human) in response to an acute bout of exercise in skeletal muscle were visualized on the

individual subject basis (raw counts converted to relative Z-scores) in mmYS (A), hsYS (B),

hsAS (C), and hsAA (D), and the RBP genes were overlapped across the cohorts to find common

RBPs (E).

Figure 4. Differential splicing events in acutely exercised young and aged skeletal muscle with

and without training. (A) Level of potential splicing change detected between hsYS-pre and

hsYS-4pe, categorized into 5 common event types (SE, RI, MXE, A5SS, and A3SS). (B) Level

of potential splicing change detected between hsAS-pre and hsAS-3pe, categorized into 5

common event types. (C) Level of potential splicing change detected between hsAA-pre and

hsAA-3pe, categorized into 5 common event types. (D) Euler overlap of genes with differences

in potential splicing events detected pre- and post-acute exercise across the three human SRS

cohorts.

Figure 5. Splicing and isoform switching in acutely exercised young sedentary skeletal muscle

in a temporal manner. (A) Expressional changes in genes with significant (raw $p < .05$) DGE and

DAS between mmYS-pre and mmYS-ipe. (B) Expressional changes in genes with significant

DGE and DAS between mmYS-pre and mmYS-1pe. (C) Expressional changes in genes with

significant DGE and DAS between mmYS-pre and mmYS-24pe. (D) Isoform-switching genes

with statistical significance ($\Delta\Pi > 10\%$, p-adj $< .05$) but without significant DGE between pre-

and post-acute exercise time points. (E) Euler overlap of exercise-associated genes with

statistical significant isoform switching but without significant DGE between pre- and post-acute

exercise time points. (F) DIE of *mSirt2*, an exercise-associated gene, between its 201 and 202

transcript variants between pre- and 24 hours post-exercise according to LRS data. (G) DIE of

mSirt2-202 transcript variant between pre and 24pe was validated using gel-based RT-PCR. **,

$p < 0.01$ compared to mmYS-pre. All experiments were conducted with $n = 5$ per cohort.

Figure 6. Splicing and isoform switching of RBPs in acutely exercised young sedentary skeletal

muscle in a temporal manner. (A) Euler overlap of all genes with statistically significant isoform

switching but without significant DGE between pre- and post-acute exercise time points found in

Figure 5D. (B) Top 10 GO terms from the 262 common genes across the three post-acute

exercise time points found in Figure 6A using PANTHER. (C) Overlap of the 262 common

genes with in-house list of 1130 RBP genes, highlighting 61 RBP genes without significant DGE.

(D) Overview of mouse *Hnrnpa3* gene loci and location of primers used for RT-ddPCR, and the

exon/intron structures of transcript variants 203, 204, and 205. (E) DIE of *mHnrnpa3*, an RBP

gene, between its protein-coding (all reads not assigned to 205) and non-coding (205, retained

intron) transcript variants between pre- and 24 hours post-exercise according to LRS data. (F)

DIE of *mHnrnpa3* protein-coding transcripts between pre and 24pe was validated using RT-

ddPCR. (G) Western blot validation of significant isoform switching of hnRNP-A3 at the protein

level, normalized to total protein at 40-45 kDa band detection then normalized to mmYS-pre for

relative quantification in arbitrary units (AU), with representative image of western blots. (H)

DIE of *mHnrnpa1*, an RBP gene, between its 201 and 202/nonsense-mediated decay transcript

variants between pre- and 24 hours post-exercise according to LRS data. (I) DIE of *mHnrnpa1*-

202/nonsense-mediated decay transcript variants between pre and 24pe was validated using gel-

based RT-PCR. Full RT-PCR gel shown in S5B. (H) DIE of *mTia1*, an RBP gene, between its

201 and 202 transcript variants between pre- and 24 hours post-exercise according to LRS data.

(K) DIE of *mTia1*-202 transcript variant between pre and 24pe was validated using gel-based

RT-PCR. *, *p*-adj < 0.05 compared to mmYS-pre. **, *p*-adj < 0.01 compared to mmYS-pre. ***,

p-adj < 0.001 compared to mmYS-pre. All experiments were conducted with $n = 5$ per cohort.

Figure 7. Current model on the effects of acute endurance exercise on the AS regulatory

mechanism with key *trans*-factors/RBPs (e.g., hnRNP-A1, -A3, and TIA1).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

(A) hsYS-pre:4pe Splicing Events

RI, 20
MXE, 14
SE, 67
A5SS, 13
A3SS, 22

(B) hsAS-pre:3pe Splicing Events

RI, 97
MXE, 33
SE, 136
A3SS, 101
A5SS, 43

(C) hsAA-pre:3pe Splicing Events

RI, 68
MXE, 21
SE, 136
A3SS, 63
A5SS, 40

(D) Overlaps of Differential Splicing Across Human Cohorts

hsAA−pre:3pe
287

hsYS−pre:4pe
115

hsAS−pre:3pe
359

Figure 5.

(A) mmYS-pre:ipe DGE vs DAS

(B) mmYS-pre:1pe DGE vs DAS

(C) mmYS-pre:24pe DGE vs DAS

Color

● Both DGE and DAS

● Only DAS

● Only DGE

(D) mmYS Genes with Only DAS

(E) Overlaps of Exercise-Associated Genes with Only DAS in mmYS

(F) *mSirt2* in mmYS-pre:24pe LRS Data

(G) *mSirt2* in mmYS-pre:24pe RT-PCR

Figure 6.

(A) Overlaps of mmYS Only DAS Genes

(B) Enrichment of Overlapping Only DAS Genes

(C) Enrichment of RBPs in Genes with Significant Isoform Switching Post-Exercise

(D) Mouse *Hnrnpa3* Gene and AS Variants

(E) *mHnrnpa3* in mmYS-pre:24pe LRS Data

(F) *mHnrnpa3* in mmYS-pre:24pe RT-ddPCR

(G) hnRNP-A3 Protein Levels in mmYS

(H) *mHnrnpa1* in mmYS-pre:24pe LRS

(I) *mHnrnpa1* Differential Splicing

(J) *mTia1* in mmYS-pre:24pe LRS

(K) *mTia1* Differential Splicing

Figure 7.