1 **Identifying genomic adaptation to local climate using a**

2 **mechanistic evolutionary model**

Abstract

1. Identifying genomic adaptation is key to understanding species' evolutionary responses to environmental changes. However, current methods to identify adaptive variation have two major limitations. First, when estimating genetic variation, most methods do not account for observational uncertainty in genetic data because of finite sampling and missing genotypes. Second, many current methods use phenomenological models to partition genetic variation into adaptive and non-adaptive components.

2. We address these limitations by developing a hierarchical Bayesian model that explicitly accounts for observational uncertainty and underlying evolutionary processes. The first layer of the hierarchy is the data model that captures observational uncertainty by probabilistically linking RAD-sequence data to genetic variation. The second layer is a process model that represents how evolutionary forces, such as local adaptation, mutation, migration, and drift, maintain genetic variation. The third layer is the parameter model, which incorporates our knowledge about biological processes. For example, because most loci in the genome are expected to be neutral, the environmental sensitivity coefficients are assigned a regularized prior centered at zero. Together, the three models provide a rigorous probabilistic framework to identify local adaptation in wild organisms.

3. Analysis of simulated RAD-seq data shows that our statistical model can reliably infer adaptive genetic variation. To show the real-world applicability of our method, we re-analyzed RAD-seq data (~105k SNPs) from Willow Flycatchers (*Empidonax traillii*) in the USA. We found 30 genes close to loci that showed a statistically significant association with temperature seasonality. Gene ontology suggests that several of these genes play a crucial role in egg mineralization, feather development, and the ability to withstand extreme temperatures.

4. Moreover, the data and process models can be modified to accommodate a wide range of genetic

datasets (*e.g.*, pool and low coverage genome sequencing) and demographic histories (*e.g.,* range

shifts) to study climatic adaptation in a wide range of natural systems.

Data and code availability: The data and code files are provided with the submission. Upon acceptance,

the data and code files will be deposited in Zenodo.

Keywords: Bayesian statistics, climate adaptation, Environmental association analysis, metapopulation,

and uncertainty quantification.

Introduction

Many species face a looming threat of extinction as global temperature rises (Thomas *et al.* 2004;

Urban 2015). This risk is particularly concerning for species with limited dispersal capacity that are slow

to track their climatic niches (Carroll *et al.* 2015). Alternatively, some species may adapt to the local

climate, potentially alleviating the risk of extinction. Natural selection favors individuals with heritable

phenotypes best suited to survive and reproduce in local climates, creating a geographical mosaic in

frequencies of adaptive alleles congruent with climatic gradients (Hedrick, Ginevan & Ewing 1976;

Hedrick 1986; Hedrick 2006). Therefore, to understand the evolutionary responses of a species to a

changing climate and incorporate this knowledge to inform evolutionary management policies (Smith *et al.*

2014), we need robust methodologies to identify genes (and their function) that confer local adaptation in

natural populations.

With recent advances in high-throughput sequencing and global environmental sensing

technologies (Chuvieco 2020; Satam *et al.* 2023), new computational approaches have enabled researchers

to identify adaptive genetic variation in wild populations. These approaches, collectively known as

environmental association analysis (Rellstab *et al.* 2015; Hoban *et al.* 2016), have gained widespread

popularity, which can be partly attributed to declining sequencing costs, improvements in genomic tools,

detailed global climatic maps, and availability of computational resources. These advances have allowed

researchers to study climatic adaptation in non-model organisms at a fine genomic resolution and large

spatial extent for a reasonable cost (Frichot *et al.* 2013; De Villemereuil & Gaggiotti 2015; Fitzpatrick &

Keller 2015; Wagner, Chávez-Pesqueira & Forester 2017).

Broadly, environmental association analysis involves three steps. First, sequencing data from

spatially referenced individuals are used to estimate spatial variation in allele frequencies. Next, a statistical

model is used to partition this genetic variation into adaptive and non-adaptive components. The adaptive

variation corresponds to the response of an allele to changes in prevailing climatic conditions. The non-

adaptive variation describes the allelic variation (or neutral variation) that stems from evolutionary forces

other than local adaptation, including gene flow among populations, mutation, and drift. Finally, a statistical

criterion is used to select loci that exhibit strong adaptive genetic variation relative to non-adaptive

variation.

Despite the success of environmental association studies, current methods face several conceptual

and statistical challenges that limit the reliability of the inferences. Many commonly used environmental

association methods do not account for uncertainty in estimates of allele frequencies (see Coop *et al.* (2010)

and Foll and Gaggiotti (2008) for exceptions), which can introduce biases in downstream analyses. Due to

practical constraints, biogeographers collect and sequence only a finite number of individuals in a

population. The genotype of these individuals, inferred from RAD (restriction site-associated DNA)

sequencing (Baird *et al.* 2008), for instance, contains imprecise information regarding the underlying allele

frequencies that result from sampling a small subset of individuals within a population. Additionally,

because of low coverage and alignment errors, the true genotypes of some individuals may be missing

(Huang & Knowles 2016), which further reduces sample size and increases uncertainty. Failure to account

for these sources of uncertainty results in overly precise estimates of genetic variation, which can lead to

an increased number of false positives (i.e., incorrectly inferring non-adaptive loci as important).

Alternatively, researchers discarding data due to noisy estimates of genetic variation may lose valuable

(although imprecise) information, thereby increasing the rates of false negatives (i.e., inability to identify

an adaptive locus).

Another major difficulty is that many environmental association methods characterize genetic

variation using models, such as generalized linear models (Rellstab *et al.* 2015). Although the underlying

relationships between data and parameters (e.g., environment and allele frequency) in phenomenological

models are based on evolutionary thinking, these relationships are often weakly motivated by evolutionary

theory (but not always; see De Villemereuil and Gaggiotti (2015) and Günther and Coop (2013) for

exceptions). For example, some classical environmental association models assume an S-shape response

curve to relate environmental variables and allele frequency (Joost *et al.* 2007; Stucki *et al.* 2017). This

response curve has several attractive characteristics: it is bounded between zero and one, and has an incline

in the middle that emulates the response of an allele to local climatic adaptation. However, the S-shaped

response curve is not unique in these characteristics. A biogeographer can construct alternate equally

plausible response curves (*e.g.*, cdf functions) and, consequently, there is no easy way to choose one over

the other *a priori*.

In contrast, mechanistic models of evolution are based on the first principles of birth-death

processes, and, therefore, a biogeographer can evaluate model assumptions based on the constraints

imposed by the population demography and natural history of the species (Rice 2004). Mechanistic models

of evolution in statistical inferences offers a unique advantage—they constrain the flow of information from

data to model parameters using a theoretical understanding. This may allow biogeographers to create

bespoke statistical models tailored to match the demographic history of the species (Wikle 2003; Hooten

& Hefley 2019).

Over the past two decades, theoretical and computational advances in Bayesian statistics have

fueled a surge in the development of probabilistic hierarchical models (Beaumont & Rannala 2004; Stadler

et al. 2024), offering researchers new avenues to address the aforementioned challenges. These hierarchical

models can account for uncertainty in estimates of genetic variation (stemming from finite sampling and

sequencing) (Gompert & Buerkle 2011) and then propagate this uncertainty downstream to mechanistic

process models of evolution, enabling a wide range of inferences ranging from constructing phylogenies

(Suchard *et al.* 2018) to identifying migration corridors (Hanks & Hooten 2013; Marcus *et al.* 2021) to

estimating growth rates of emerging viral diseases (Stadler 2010). In fact, these statistical insights are also

beginning to permeate the latest generation of genotype-environment association methods (Coop *et al.*

2010; De Villemereuil & Gaggiotti 2015).

Hierarchical models help us do joint inferences that allows us to learn parameters using data,

process, and parameter models (Berliner 1996). In the context of environmental association analysis, the

data model describes how the unobserved allele frequencies could have led to genetic data. The process

model describes the biological processes determining spatial variation in allele frequencies. The parameter

model describes the knowledge the biogeographer has about the parameters before the data are collected,

based on past research.

This structure of the hierarchical model offers several advantages in identifying genomic adaptation

to climate. First, the data model enables a biogeographer to quantify observational uncertainty in estimates

of genetic variation, particularly in cases where sample sizes are small and genotypes are missing. Second,

these imprecise estimates of genetic variation can then be linked to a process model of evolution. This

evolution model is based on the demographic history of the species and provides a mechanistic explanation

of how to partition genetic variation into adaptive (response curve) and non-adaptive (neutral) components.

Lastly, the parameter model can be used to incorporate knowledge from past research. For example, the

theory of molecular evolution suggests that the overwhelming majority of variation in genomes is non-

adaptive and stems from the interplay between mutation, drift, and migration, while only a small fraction

of the variation stems from local adaptation (Kimura 1983). To incorporate this prior knowledge, a

biogeographer can assign a low prior probability that the response curve has a finite slope. This prior

effectively shrinks (regularizes) the adaptive genetic variation to zero for most loci, providing a systematic

way to evaluate the relative contribution of climate adaptation and neutral processes in determining genetic

variation.

Because of these features of the data, process, and parameter models, hierarchical models provide

a rigorous probabilistic framework to identify genomic adaptation in wild organisms informed by

theoretical principles of evolutionary biology using noisy genetic data. In what follows, we present a new

demographic Bayesian model and test its robustness by analyzing synthetic data with known parameter

values. We use the model to analyze RAD-seq data from Willow Flycatchers (*Empidonax traillii*) and

identify candidate genes that may be responsible for local climatic adaptation. Although we apply our model

to Willow Flycatchers, the statistical insights are general and broadly applicable to other species.

140   *Willow Flycatchers*

The Willow Flycatcher is a migratory songbird that breeds mainly in the United States and Southern

Canada. The species is categorized into four distinct subspecies—the Pacific Northwestern form (*E. t.*

brewsteri), Western Central form (*E. t. adastus*), Eastern form (*E. t. traillii*), and Southwestern form (*E. t.*

extimus). Among them, the Southwestern form has experienced a precipitous decline in abundance, likely

due to the loss of riparian habitat along streams and waterways (Sedgwick 2000), which provide respite

during extreme temperatures. In 1995, the Southwestern form was listed as federally endangered

subspecies (Unitt 1987) due to its genetic, ecological, and song distinctiveness (Theimer *et al.* 2016; Ruegg

et al. 2018; Mahoney *et al.* 2020).

Previous landscape genomic work identified highly significant correlations between allele

frequencies in genes linked to thermal tolerance and the intensity of summer heat waves in the southwest

(Ruegg *et al.* 2018). Therefore, re-analyzing the Willow Flycatcher genome-wide genetic dataset, initially

examined using traditional phenomenological models, offers an ideal opportunity to apply a new

environmental association analysis that mechanistically accounts for genetic variation. More broadly, the

results have important implications for implementing management practices designed to improve the

genetic health of the endangered subspecies.

157   **Data**

A total of 175 Willow flycatchers were sampled at 23 one-degree squares across the continental

United States (Fig. 1). The sampling effort varied from 2-21 individuals per location. DNA was extracted

from blood and tissue samples using the QiagenTM DNeasy Blood and Tissue extraction kit and quantified

using the Qubit® dsDNA HS Assay kit (Thermo Fisher Scientific). Sequencing was conducted across three

lanes of 100 bp paired-end reads on an Illumina HiSeq 2500 at the UC Davis Genome Center. We filtered

SNPs using the tradeoff between discarding SNPs with low coverage and discarding individuals with

missing genotypes using the R package genoscapeRtools (Anderson 2019), resulting in approximately 105k

SNPs, of which 3 percent of the loci had a missing genotype (Ruegg *et al.* 2018). At each sampling location,

climate data were obtained from WorldClim, which averaged the climate between 1960 and 1990 (Hijmans

et al. 2005). Due to the high correlation between some of the top-ranked climatic variables identified in

Ruegg *et al.* (2018), we limited our analysis to the four least correlated climate variables (standardized) to

reduce collinearity: BIO 4 (temperature seasonality), BIO 5 (maximum temperature of the warmest month),

BIO 11 (mean temperature of the coldest quarter), and BIO 17 (Precipitation in the driest quarter). For more

details about the sampling design, sequencing, and bioinformatic analysis, refer to Ruegg *et al.* (2018).

Model Formulation

To identify putative loci under selection due to local adaptation, we specified a Bayesian model

with three hierarchical levels corresponding to the data, process, and parameter models. These three levels

are organized such that the output of the parameter model is the input for the process model, whose output

is the input for the observer model (Pagel & Schurr 2012). To facilitate a broad understanding of how the

statistical model works, Figure 2 illustrates the three levels of the hierarchical model, showing how these

levels are connected. In Figure 3, we provide a Directed Acyclic Graph (DAG) to illustrate the relationship

between data and parameters. In the following section, we give details on the assumptions underlying the

model's construction (also see the Supplementary Information for a concise summary).

Data model

We assume that Willow Flycatchers were sampled at K sites across the United States, and each bird

was genotyped at L independent genetic (RAD) markers (see sampling and sequencing above). We

characterized the genotype of the ith sampled bird at site k using g_{ilk}, which corresponds to the number of

copies of the reference allele at locus l. Assuming individuals have biallelic loci, g_{ilk} is a discrete variable

that takes values zero, one, or two. Using these individual genotype data, we define a population-level

genotype variable

$$y_{lk} = \sum_{i=1}^{N_{lk}} g_{ilk}, \tag{1}$$

190   that corresponds to the number of reference alleles for N_{lk} birds that were genotyped at locus l. In the data,

191   N_{lk} is always less than or equal to the number of birds sampled at a site because, for some individuals, the

192   genotype at a locus may be missing due to sequencing errors or low coverage. Assuming the birds were

193   randomly sampled, we model the variation in reference allele frequency (p_{lk}) using a binomial distribution,

$$y_{lk} \sim \text{Binomial}(2N_{lk}, p_{lk}). \tag{2}$$

194   Alternatively, we can also use

$$g_{ilk} \sim \text{Binomial}(2, p_{lk}), \tag{3}$$

195   as a data model to describe the statistical relationship between allele frequency and an individual's

196   genotype. However, because of computational efficiency, we use equation (2). Note that the observer model

197   accounts for the uncertainty in genetic variation that stems from finite sampling and missing genotypes.

198   *Process model*

199   We consider an island-metapopulation model proposed by Wright (1931) to model evolutionary

200   dynamics. We rely on this model because it provides a parsimonious explanation of how evolutionary

201   processes maintain genetic variation and allows for reasonable statistical inferences even when the process

202   model is misspecified due to complexity in evolutionary dynamics (see the Model Testing section below).

203   We assume that the demes in the metapopulation correspond to the sites where the birds were

sampled. Each deme has a population size of N_e, and the migration rate between any pair of demes is equal.

We assume that the variation in allele frequencies across demes is maintained by directional and non-

directional evolutionary forces. Directional forces—such as local adaptation to climate, mutation, and

migration—result in changes in the mean value of allele frequency. Non-directional forces—such as genetic

drift—do not change the mean value of allele frequency but create sampling variance due to finite

population size. Because of this variance, the frequency of an allele cannot be determined exactly. Instead,

allele frequency is characterized probabilistically using a probability distribution, $\psi(p_{lk}, t)$ (Rice 2004;

Blanquart, Gandon & Nuismer 2012). Using the Fokker-Planck equation, one can show that the probability

distribution of the frequency of the reference allele changes as follows:

$$\frac{\partial \psi(p_{lk}, t)}{\partial t} = -\frac{\partial}{\partial p_{lk}}[\psi(p_{lk}, t)M(p_{lk})] + \frac{1}{2}\frac{\partial^2}{\partial^2 p_{lk}}[\psi(p_{lk}, t)V(p_{lk})], \tag{4}$$

213   where

$$M(p_{lk}) = \underbrace{s_{lk}p_{lk}q_{lk}}_{\text{Selection}} + \underbrace{u_{2l}q_{lk} - u_{1l}p_{lk}}_{\text{Mutation}} + \underbrace{m(\bar{p}_l - p_{lk})}_{\text{Migration}} \tag{5}$$

214   is the rate of directional change in the allele frequency and

$$V(p_{lk}) = \frac{p_{lk}q_{lk}}{2N_e} \tag{6}$$

is the variance in the allele frequency due to non-directional (*i.e.*, drift) effects. In equations (4)-(6),

$q_{lk}(= 1 - p_{lk})$ is the frequency of the alternate allele, s_{lk} is the environmentally regulated selection

coefficient, u_{2l} and u_{1l} are the locus-specific forward and backward mutation rates, m is the migration rate,

and $\bar{p}_l$ is the average allele frequency of the immigrants. We consider multiplicative selection dynamics

(*i.e.*, no dominance), such that the relative fitness of individuals with genotype (g_{ilk}) zero, one, and two is

$1 + 2s_{lk}$, $1 + s_{lk}$, and 1, respectively. Our model can also be modified to accommodate alternate forms of

selection dynamics, such as frequency-dependent selection, underdominance, and heterozygote advantage.

At the time of sampling, the allele frequency distribution is assumed to be in quasi-equilibrium

with climate conditions. To obtain this equilibrium distribution, also known as the stationary distribution

(ψ_s), we set $\partial\psi(p_{lk}, t)/\partial t = 0$. The stationary distribution of the allele frequency can be expressed in terms

of evolutionary parameters as follows (Blanquart, Gandon & Nuismer 2012):

$$\psi_s(p_{lk}|\tilde{s}_{lk}, \mu_l, \kappa_l) = \frac{e^{\tilde{s}_{lk}p_{lk}}}{{}_1F_1(\mu_l\kappa_l, \kappa_l, \tilde{s}_{lk})} \text{Beta}(p_{lk}|\mu_l, \kappa_l), \tag{7}$$

226   where $\tilde{s}_{lk} = 4N_e s_{lk}$, ${}_1F_1$ is the hypergeometric confluent function, and $\mu_l = (u_{2l} + m\bar{p}_l)/(u_{1l} + u_{2l} +$

227     $m)$ and $\kappa_l = 4N_e(u_{1l} + u_{2l} + m)$ are the mean and precision parameters of the beta distribution. To

228   incorporate climate adaptation, we assume that the selection coefficient, $\tilde{s}_{lk}$, is regulated by local

229   environmental conditions. For simplicity, we assume a linear relationship between the selection coefficient

230   and *E* standardized environmental variables:

$$\tilde{s}_{lk} = \alpha_l + \sum_{j=1}^{E} \beta_{lj}e_{jk}. \tag{8}$$

However, based on the natural history of the species, a biogeographer may consider alternative functional

forms to describe the relationship between the selection coefficient and environmental variables. In

equation (8), e_{jk} is the jth environmental variable (standardized) in deme k, β_{lj} is the selection coefficient's

sensitivity to variation in the jth environmental variable, and α_l can be interpreted either as the selection

coefficient at mean environmental conditions or a fixed value of selection coefficient arising from adaptive

processes other than local climatic adaptation.

The stationary distribution in equation (7) has several notable features. The response curve of an

adaptive allele (the relationship between climate and mean allele frequency),

$$\mathbb{E}[p_{lk}] = \mu_l \frac{{}_1F_1(1 + \mu_l\kappa_l, 1 + \kappa_l, \tilde{s}_{lk})}{{}_1F_1(\mu_l\kappa_l, \kappa_l, \tilde{s}_{lk})}, \tag{9}$$

239 is bounded between zero and one, and its shape is determined by evolutionary parameters with biological

240 interpretation (Blanquart, Gandon & Nuismer 2012). When selection is absent, the beta distribution

241 captures the non-adaptive genetic variation,

$$\psi_s(p_{lk}|0, \mu_l, \kappa_l) = \text{Beta}(p_{lk}| \mu_l, \kappa_l). \tag{10}$$

242 Therefore, to account for the joint contribution of adaptive and non-adaptive evolutionary forces in

243 determining genetic variation, we use the stationary distribution to statistically model evolutionary

244 dynamics,

$$p_{lk} \sim \frac{e^{\tilde{s}_{lk}p_{lk}}}{{}_1F_1(\mu_l\kappa_l, \kappa_l, \tilde{s}_{lk})} \text{Beta}(\mu_l, \kappa_l). \tag{11}$$

245 For fast and memory-efficient implementation of the statistical model, we combine the data (Eq.

246 2) and process (Eq. 11) models by marginalizing over p_{lk},

$$y_{lk} \sim \int_0^1 \text{Binomial}(y_{lk}|2N_{lk}, p_{lk})\psi_s(p_{lk}|\tilde{s}_{lk}, \mu_l, \kappa_l) \, dp_{lk}, \tag{12}$$

247 which results in an integrated likelihood that relates genotype counts to evolutionary dynamics as follows:

$$y_{lk} \sim \frac{{}_1F_1(\mu_l\kappa_l + y_{lk}, 2N_{lk} + \kappa_l, \tilde{s}_{lk})}{{}_1F_1(\mu_l\kappa_l, \kappa_l, \tilde{s}_{lk})} \text{BetaBinomial}(2N_{lk}, \mu_l, \kappa_l). \tag{13}$$

Note that the integrated likelihood in equation (13) accounts for the uncertainty that stems from noisy

genetic data and the stochastic nature of evolutionary dynamics.

Parameter model

Finally, we assign priors to parameters in equation (13) based on our prior scientific knowledge.

We assign $\mu_l \sim \text{Uniform}(0,1)$ and $\kappa_l \sim \text{Normal}^+(0,5)$ priors in equation (13). Due to degeneracy of the

likelihood (Eq. 11, see Fig. S1), the mean of the beta distribution (μ_l) and selection coefficient at the mean

environment (α_l) are non-identifiable (see Supplementary Information for more details). We alleviate this

by fixing $\alpha_l = 0$ and re-interpret μ_l as the mean of the baseline allele frequency distribution (or neutral

distribution) resulting from evolutionary forces other than adaptation to variation in local climate around

the mean. Next, because we expect that most of the genetic variation arises due to neutral processes (Kimura

1983), we use a regularized horseshoe shrinkage parameter model (Piironen & Vehtari 2017) for sensitivity

coefficients, β_{lj}. The horseshoe prior shrinks most of the β_{lj} to zero, allowing a fraction of the coefficients

to take non-zero values. The horseshoe model achieves this using global (τ) and local ($\tilde{\lambda}_{lj}$) shrinkage

parameters and is expressed as

$$\beta_{lj} \sim \text{Normal}(0, \tau \tilde{\lambda}_{lj}), \tag{14}$$

263   where

$$\tilde{\lambda}_{lj} = \frac{c \lambda_{lj}}{\sqrt{c^2 + \tau^2 \lambda_{lj}^2}}. \tag{15}$$

264   To control global shrinkage, we specify a half-Cauchy prior,

$$\tau \sim \text{Cauchy}^+\left(0, \frac{f}{\sqrt{KLE}}\right), \tag{16}$$

where f is a free variable, which we set to 20. Based on the recommendations of Piironen and Vehtari

(2017), we tested a range of values of f and found that the statistical results are robust to orders of

magnitude of variation in f (see Model Testing section below). Note that the global sparsity (controlled by

τ) increases with the number of SNPs (L) and environmental variables (E). On the other hand, the local

shrinkage parameter, $\tilde{\lambda}_{lj}$, allows some coefficients to escape shrinkage because of a heavy-tail prior,

$\lambda_{lj} \sim \text{Cauchy}^+(0,1)$. However, these large-valued coefficients are weakly shrunk with an implied

regularization of $\text{Normal}(0, c)$ with $c^2 \sim \text{InvGamma}(2,4)$. We adopt an inverse-gamma prior because the

prior means for both c^2 (which controls adaptive dynamics via sensitivity coefficients) and κ_l (which

governs neutral dynamics) are similar in magnitude, but the heavy right tail of the inverse-gamma

distribution allows adaptive loci to attain large sensitivity coefficients. These features of the local and global

shrinkage parameters effectively induce a spike and slab prior (Mitchell & Beauchamp 1988), which

prevent numerical pathologies when large effect loci are weakly identifiable.

278   **Model Testing**

To assess the robustness of our statistical method, we fit the model to synthetic data that are similar

in characteristics to the Willow Flycatcher genomic data, and the generative process only partially

resembles the process model considered in our statistical model. This simulation allows us to test the

sensitivity of our method to model misspecifications and test its applicability to biologically realistic

datasets.

We generated the synthetic genotype data in three stages (see Supplementary code for simulation

details). In the first stage, we simulated selection dynamics. We considered diploid individuals with a

genome size of one thousand independent loci. Each locus in the genome was assigned a random non-zero

α_l. We randomly selected fifteen loci that contribute to local adaptation. Each of the fifteen loci was

randomly paired with one of the four environmental variables, and the corresponding selection coefficient

(s_{lk}) was calculated using equation (8). We used real environmental values (standardized) to calculate s_{lk}

to preserve the correlation structure between environmental variables.

In the second stage, we simulated metapopulation dynamics with demes equal to the number of

sampling sites in the real data. We sampled random variables from the stationary distribution of allele

frequencies by simulating the following stochastic differential equation (Korolev *et al.* 2010):

$$dp_{lk} = \left[\underbrace{s_{lk}p_{lk}q_{lk}}_{\text{Selection}} + \underbrace{u_{2l}q_{lk} - u_{1l}p_{lk}}_{\text{Mutation}} + \sum_{n=1}^{K} \underbrace{e^{-\rho d_{nk}}(p_{ln} - p_{lk})}_{\text{Migration}} \right] dt + \underbrace{\sqrt{p_{lk}q_{lk}/2N_e}}_{\text{Drift}} \, dB, \qquad (17)$$

where s_{lk} is the selection coefficient obtained from stage one, B is the standard Brownian motion, d_{nk} is

the distance between demes n and k, and ρ^{-1} is the dispersal length scale. The above equation is an Itô

representation of the Fokker-Planck equation presented earlier (Eq. 4), with a small modification: migration

rate depends on the distance between demes (Fig. S2).

In the third stage, we simulated genotypes. In each deme, we simulated genotypes for the same

number of individuals as in the real data. For each individual, we generated a genotype at a locus by

sampling from the distribution Binomial$(2, p_{lk})$ (Eq. 3), where p_{lk} is the reference allele frequency

obtained from the second stage. We randomly selected three percent of the loci and treated them as missing.

To test if the statistical model can identify loci that contribute to local adaptation in synthetic data,

we fit the model using Stan programming language (Carpenter *et al.* 2017) to obtain a posterior sample of

sensitivity coefficients, β_{lj}, using Hamiltonian Monte Carlo (Neal 2011). For each pair of loci and

environmental variables, we computed the probability that the posterior distribution of the sensitivity

coefficient included zero. If this probability is less than a threshold value of 0.05 (p_{th}), we inferred that the

locus may have contributed to local adaptation for the corresponding environmental variable.

To conduct robustness checks, we generated ten synthetic datasets for nine parameter regimes (a

total of 90 datasets; see Supplementary Information for parameter values used to generate synthetic

datasets). These parameter combinations represent three mutation and migration regimes—high, medium,

and low—relative to local adaptation. For each regime, we calculated the false-negative and false-positive

rates. We found that our model performs best in the low mutation-migration regime, where the genetic

variation is more sensitive to climatic variation (Tables S1 and S2). However, this regime also yields higher

rates of false positives. In contrast, in the high mutation-migration regime, neutral evolutionary forces play

a larger role, and consequently, the statistical model results in high false negative rates.

For illustration, the Manhattan plot in Fig. 4A shows the result from a simulation corresponding to

the low mutation and medium migration regime. The y-axis corresponds to the negative log probability that

the posterior distribution of sensitivity coefficients included zero (Wang *et al.* 2022). The points above $y =$

2.99 (the black line) have $p_{th} < 0.05$. These points are characterized by a ring and solid center. The color

of the center represents the true environmental variable associated with the loci, and the color of the ring

represents the statistically inferred adaptive environmental variable. Note that, in this simulation run, the

model incorrectly identified the corresponding environmental variable for two of the nine loci (points above

the black line with a green ring and purple center). This feature can be explained by a negative correlation

(-0.72) between environmental variables, BIO 4 (green) and BIO 11 (purple). As a result, the statistical

model used one of the environmental variables as a substitute for another (Rellstab *et al.* 2015). We also

re-analyzed the data in Fig. 4A by varying f over four orders of magnitude (2^0 to 2^{14}). We find that our

results are nearly identical to results in Fig. 4A (see Fig. S3 and S4), suggesting that our analysis is robust

to a wide variation in choice of priors for τ.

Next, we re-analyzed the synthetic datasets using latent factor mixed model (LFMM; Frichot *et al.*

2013) which was used to analyze Willow Flycatcher data in Ruegg *et al.* (2018). We conducted simulations

using two versions of the synthetic dataset. In the first version, we used individual genotypes, and, in the

second version, we used raw allele frequencies (*i.e.*, $p_{lk} = 0.5\, y_{lk}/N_{lk}$) as input. Our results suggest that

LFMM yielded much higher false-negative and false-positive rates (Tables S3-S6).

335   **Data Analysis**

Analyzing genomic data from Willow Flycatchers revealed 47 significant loci, all except one were

associated with BIO 4 (Fig. 4B and Fig. S5). A closer inspection of these loci suggests that many of them

cluster together on the genome (indicated by vertical gray lines in Fig. 4B). These clustering patterns are

unlikely to happen by random chance. Alternatively, the observed clustering patterns can be explained by

gene hitchhiking (Smith & Haigh 1974). The allele frequencies at a locus under selection and its

neighboring neutral loci rise or fall in unison due to physical linkage on the chromosome. Consequently,

when a gene is under selection, neutral polymorphic sites close to the gene experience environmentally

regulated pseudo-selection (Barton 1998).

Using the annotated genome of Willow Flycatchers, 36 of the candidate loci were found within or

nearby (within 25kb) 30 named genes (Table S7). Twenty-three of these genes are characterized, and 20

have functional roles in chicken (*Gallus gallus*) that span 8 gene ontology categories. The majority of genes

cluster in 4 categories: 5 genes are involved in catalytic activity, 5 genes have binding functionality, 3 have

transcription regulatory activity, and 3 have transporter activity. A closer investigation into several genes

shows that EDIL3 plays a role in the egg mineralization process in Aves (Le Roy *et al.* 2021), PCDH1 is

involved in feather development (Lin, Wang & Redies 2013), and GRIK2 acts as a thermoreceptor

conferring sensitivity to cold temperatures in mice (Cai *et al.* 2024).

The above analysis required 32 hours to complete on a machine with a 24-core 3.6 GHz processor.

By subsampling the genomic data, we also estimated the model run time for $E = 4$ and $L = 2^8, 2^9, ...$,

and 2^{17} (Fig. S6).

Discussion

Adaptation to climate is ubiquitous and will continue to play a major role in maintaining

biodiversity in the Anthropocene (Thompson 2013). However, many commonly used methods for

identifying genomic adaptation are limited because they often lack the ability to handle noisy genetic data

and do not formally account for the demographic history of the species. Our mechanistic Bayesian model

addresses these limitations. The important aspects of our approach comprise the following features.

First, we proposed a data model that probabilistically links RAD-seq data to genetic variation (Eq.

2). This probabilistic link allowed us to quantify uncertainty in genetic variation that arises due to finite

sampling and missing genotypes. Consequently, the data model properly accounts for uncertainty in the

estimation of genetic variation and faithfully propagates the available information in raw genetic data to

the process model for downstream analysis (Hobbs & Hooten 2025).

Second, we used a metapopulation process model to partition estimated genetic variation into

adaptive and non-adaptive components (Blanquart, Gandon & Nuismer 2012). The metapopulation model

can be implemented using a beta distribution to characterize neutral variation (Eq. 10) and hypergeometric

confluent functions to model adaptive variation (Eq. 9). Our process model is motivated by theoretical

principles from evolutionary biology. Thus, we can evaluate the model based on the underlying

assumptions.

Finally, previous work shows that most of the genetic variation in wild populations is maintained

by non-adaptive evolutionary forces (Kimura 1983). We incorporated this prior knowledge into our

statistical model using a regularized horseshoe parameter model, which shrinks most of the selection

coefficients to zero. The level of shrinkage is controlled by a global shrinkage parameter that introduces

sparsity in sensitivity coefficients (Piironen & Vehtari 2017) (Eq. 14). Thus, the parameter model allowed

us to partition genetic variation into adaptive (driven by environmental variation) and non-adaptive (driven

by neutral processes) components.

To test our statistical model, we conducted simulations to assess the model's performance on ninety

synthetic RAD-seq datasets generated by simulating bird genomes of 1000 loci (Eq. 17), which emulate the

characteristics of genetic data from Willow Flycatchers. Our statistical method performed best in the low

mutation-migration regime, and the performance decreased in the high mutation-migration regime (Table

S1 and S2). These results are consistent with our theoretical expectations. When the mutation and migration

rates are low, genetic variation is more sensitive to variation in climate. However, in the high mutation-

migration regime, neutral processes play a bigger role in maintaining genetic variation. As a result, it is

more challenging to identify weakly adaptive loci.

Figure 4A shows results from a simulation from the low mutation and medium migration regime.

The model correctly identified nine out of fifteen adaptive loci. However, we found that out of the nine

correctly identified loci, two loci were incorrectly paired with their corresponding environmental variable

due to a -0.72 correlation between BIO 4 and BIO 11. This highlights that biogeographers may need to

exercise caution when interpreting statistical results or use uncorrelated predictors. The statistically inferred

environmental variable might be correlated with the true environmental variable responsible for local

adaptation (Rellstab *et al.* 2015). Nevertheless, the synthetic simulations suggest that our statistical model

can be used with genetic datasets from wild populations.

Indeed, we identified 30 genes located within the 25 kb region flanking 47 significant loci in RAD-

seq data from Willow Flycatchers, most of which were associated with temperature seasonality (Fig. 4B

and S5). Some of these genes include EDIL3, PCDH1, and GRIK2, which play a role in the egg

mineralization process (Le Roy *et al.* 2021), feather development (Lin, Wang & Redies 2013), and act as a

thermoreceptor conferring sensitivity to cold temperatures (Cai *et al.* 2024), respectively. This suggests that

temperature fluctuations may be a key driver of local adaptation in the species, providing evidence that

standing genetic variation in Willow Flycatchers could alleviate or buffer extinction risk due to increasing

temperature variability predicted by climate projections (Olonscheck *et al.* 2021).

Although these genes differ from those identified in Ruegg *et al.* (2018), these differences are not

surprising because our hierarchical model is fundamentally different from traditional environmental

association analysis in terms of quantifying uncertainty while estimating genetic variation and using an

evolutionary process model to explain sources of genetic variation. Our synthetic data simulations confirm

that these differences may play an important role in inferences; re-analyzing synthetic data using LFMM

resulted in substantially higher rates of false positives and false negatives (Table S3-S6).

In addition to the conceptual advantages offered by our mechanistic statistical model, synthetic

simulations highlight several practical scenarios where our hierarchical model may be better suited to

analyze genomic data. Some of these scenarios include low sample size (between 1-4 individuals), wide

dispersion in the number of individuals sampled at various locations, and a large fraction of missing

genotypes. We can address these data scenarios by quantifying genetic variation probabilistically and

propagating the corresponding uncertainty in downstream analyses. However, there are some

computational tradeoffs: our statistical analysis required 32 hours to analyze 105k SNPs, and, as a result,

our approach may require substantial computational resources to analyze genomic datasets with millions of

SNPs.

Despite these computational requirements, the hierarchical structure of the model provides new

opportunities to improve statistical inferences. For example, the hierarchical model is flexible and can

integrate a wide range of process and data models, allowing biogeographers to better understand the

evolutionary responses of a species to a changing climate while dramatically reducing the cost of analysis.

The cost of sequencing varies along three axes—the number of sequenced sites in the genome, the depth of

sequencing effort, and the number of sampled individuals. Typically, in RAD sequencing, a small

proportion of sites in the genome are sequenced deeply to identify true genotypes, which are subsequently

used to estimate allele frequencies (Baird *et al.* 2008). Although RAD sequencing is a cost-effective

protocol to obtain a reduced representation of the genome, it fails to characterize a large proportion of

genetic variation in the genome that could be potentially adaptive (Lowry *et al.* 2017). New sequencing

protocols, such as low-depth whole genome sequencing (Alex Buerkle & Gompert 2013) and pool

sequencing (Gautier *et al.* 2013), are emerging as attractive alternatives because they provide a wider

genomic coverage without increasing cost or sacrificing statistical power. The key idea behind these

approaches is to redistribute the same resources to sequence a larger sample of individuals and a greater

proportion of the genome (Lou *et al.* 2021). This reduction in sequencing effort (per individual per locus)

increases uncertainty in the estimates of individual genotypes. But, instead of discarding SNPs due to low

coverage, one can model the true genotype as a parameter and propagate the corresponding uncertainty to

inform population-level allele frequencies. In effect, these approaches sacrifice certainty in individual

genotypes to gain genomic coverage while keeping the cost and genetic information constant. To leverage

information from these sequencing protocols, the data model in the Bayesian hierarchy can be modified to

account for genotype uncertainty in estimating allele frequencies.

Another potential avenue to improve inference is to construct alternate process models

incorporating a wider range of demographic histories to correct for model misspecification. For example,

most species, including Willow Flycatchers, are geographically structured, and, as such, the migration rates

depend on the distance between demes. Closer demes exchange more migrants than demes that are far apart.

This may create genetic patterns, such as isolation by distance (Fig. S2) and genetic swamping, that cannot

be adequately captured by an unstructured metapopulation model (Eq. 11). Because of these

misspecifications, our method less sensitive in identifying adaptive loci in weak selection and high

migration regimes. One possible resolution to this problem is to use the stationary distribution

corresponding to the structured evolutionary dynamics as the process model (see Constable & McKane

2015 for an analytical solution for the stationary distribution of a structured metapopulation). Our process

model may also be limited in non-equilibrium settings, such as range shifts due to rapidly changing climate

or land use practices. In these cases, the quasi-stationary assumption in the process model is unlikely to

hold because of transient evolutionary dynamics and eco-evolutionary feedback. Consequently, one may

want to consider a probabilistic process model that jointly describes spatiotemporal changes in abundance

and genetic variation (Schurr *et al.* 2012; Polechová & Barton 2015). To inform these joint process models,

researchers will require temporal and spatial sampling of abundances and genomes, which can be obtained

from population surveys (Pardieck *et al.* 2020) and museum collections (Payne & Sorenson 2002),

respectively. These improvements in data and process models may allow biogeographers to forecast

species' response to global change using mechanistic models of ecology and evolution.

Figure 1: Breeding range of Willow Flycatchers (dark grey region) and 23 sampling locations. The size of

the points is proportional to the number of sampled individuals in our study, which varied between 2 and

21 individuals.

Figure 2: A conceptual diagram describing the hierarchical structure of the Bayesian model. The

data model (*top*) links the individuals' genotype (g_{ilk}) to allele frequency (p_{lk}) at a locus l in patch

k (Eq. 3). The process model (*middle*) describes the stationary distribution of allele frequency

resulting from evolutionary forces, such as mutation, migration, drift, and local adaptation. (Eq. 8

and 11). Finally, the parameter model (*bottom*) allows us to regularize sensitivity coefficients (β_{lj})

using prior knowledge from molecular evolutionary theory (Eqs. [14], [15], and [16]).

Figure 3: Directed Acyclic Graph (DAG) of the hierarchical Bayesian model. The squares represent data

(genotype counts and environmental variables), the circles represent parameters, the dashed arrows

represent deterministic relationships, and the solid arrows represent stochastic relationships.

Figure 4: Manhattan plot showing negative log probability $(-\log(p_{th}))$ that the posterior distribution of sensitivity coefficients (β_{li}) includes zero for synthetic (A) and real data (B). In both plots, points above the black horizontal line have p_{th} less than 0.05. (A) For synthetic data, we denote points above the black line using an outer ring and a center. The color of the center (ring) corresponds to the true (statistically inferred) environmental variable responsible for local adaptation. (B) For real data, we denote points above the black line with a solid center, with its color corresponding to the statistically inferred environmental variable responsible for local adaptation (also see Fig. S5). The vertical gray lines represent physically linked loci with statistically significant sensitivity coefficients (see Supplementary Information for parameter values used to simulate synthetic data in Fig. 4A)

References

Alex Buerkle, C. & Gompert, Z. (2013) Population genomics based on low coverage sequencing: How low

should we go? *Molecular Ecology,* **22,** 3028-3035.

Anderson, E. (2019) genoscapeRtools: Tools for building migratory bird genoscapes. *R package version*

0.1. 0. https://rdrr. io/github/eriqande/genoscapeRtools.

Baird, N.A., Etter, P.D., Atwood, T.S., Currey, M.C., Shiver, A.L., Lewis, Z.A., Selker, E.U., Cresko, W.A.

& Johnson, E.A. (2008) Rapid SNP discovery and genetic mapping using sequenced RAD markers.

PLoS ONE, **3,** e3376.

Barton, N.H. (1998) The effect of hitch-hiking on neutral genealogies. *Genetics Research,* **72,** 123-133.

Beaumont, M.A. & Rannala, B. (2004) The Bayesian revolution in genetics. *Nature Reviews Genetics,* **5,**

251-261.

Berliner, L.M. (1996) Hierarchical Bayesian Time Series Models. pp. 15-22. Springer Netherlands,

Dordrecht.

Blanquart, F., Gandon, S. & Nuismer, S. (2012) The effects of migration and drift on local adaptation to a

heterogeneous environment. *Journal of Evolutionary Biology,* **25,** 1351-1363.

Cai, W., Zhang, W., Zheng, Q., Hor, C.C., Pan, T., Fatima, M., Dong, X., Duan, B. & Xu, X.S. (2024) The

kainate receptor GluK2 mediates cold sensing in mice. *Nature Neuroscience,* **27,** 679-688.

Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., Brubaker, M., Guo, J.,

Li, P. & Riddell, A. (2017) Stan: A probabilistic programming language. *Journal of Statistical*

Software, **76,** 1–32.

Carroll, C., Lawler, J.J., Roberts, D.R. & Hamann, A. (2015) Biotic and climatic velocity identify

contrasting areas of vulnerability to climate change. *PLoS ONE,* **10,** e0140486.

Chuvieco, E. (2020) *Fundamentals of Satellite Remote Sensing: An Environmental Approach*. CRC press,

Boca Raton.

Constable, G.W. & McKane, A.J. (2015) Stationary solutions for metapopulation Moran models with

mutation and selection. *Physical Review E,* **91,** 032711.

Coop, G., Witonsky, D., Di Rienzo, A. & Pritchard, J.K. (2010) Using environmental correlations to

identify loci underlying local adaptation. *Genetics,* **185,** 1411-1423.

De Villemereuil, P. & Gaggiotti, O.E. (2015) A new FST-based method to uncover local adaptation using

environmental variables. *Methods in Ecology and Evolution,* **6,** 1248-1258.

Fitzpatrick, M.C. & Keller, S.R. (2015) Ecological genomics meets community-level modelling of

biodiversity: Mapping the genomic landscape of current and future environmental adaptation.

Ecology Letters, **18,** 1-16.

Foll, M. & Gaggiotti, O. (2008) A genome-scan method to identify selected loci appropriate for both

dominant and codominant markers: A Bayesian perspective. *Genetics,* **180,** 977-993.

Frichot, E., Schoville, S.D., Bouchard, G. & François, O. (2013) Testing for associations between loci and

environmental gradients using latent factor mixed models. *Molecular Biology and Evolution,* **30,**

1687-1699.

Gautier, M., Foucaud, J., Gharbi, K., Cézard, T., Galan, M., Loiseau, A., Thomson, M., Pudlo, P.,

Kerdelhué, C. & Estoup, A. (2013) Estimation of population allele frequencies from next-

generation sequencing data: pool-versus individual-based genotyping. *Molecular Ecology,* **22,**

3766-3779.

Gompert, Z. & Buerkle, C.A. (2011) A hierarchical Bayesian model for next-generation population

genomics. *Genetics,* **187,** 903-917.

Günther, T. & Coop, G. (2013) Robust identification of local adaptation from allele frequencies. *Genetics,*

195, 205-220.

Hanks, E.M. & Hooten, M.B. (2013) Circuit theory and model-based inference for landscape connectivity.

Journal of the American Statistical Association, **108,** 22-33.

Hedrick, P.W. (1986) Genetic polymorphism in heterogeneous environments: A decade later. *Annual*

Review of Ecology and Systematics, **17,** 535-566.

Hedrick, P.W. (2006) Genetic polymorphism in heterogeneous environments: The age of genomics. *Annual*

Review of Ecology and Systematics, **37,** 67-93.

Hedrick, P.W., Ginevan, M.E. & Ewing, E.P. (1976) Genetic polymorphism in heterogeneous

environments. *Annual Review of Ecology and Systematics, 7,* 1-32.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very high resolution interpolated

climate surfaces for global land areas. *International Journal of Climatology: A Journal of the Royal*

Meteorological Society, **25,** 1965-1978.

Hoban, S., Kelley, J.L., Lotterhos, K.E., Antolin, M.F., Bradburd, G., Lowry, D.B., Poss, M.L., Reed, L.K.,

Storfer, A. & Whitlock, M.C. (2016) Finding the genomic basis of local adaptation: pitfalls,

practical solutions, and future directions. *The American Naturalist,* **188,** 379-397.

Hobbs, N.T. & Hooten, M.B. (2025) *Bayesian Models: A Statistical Primer for Ecologists, 2nd Edition.*

Princeton University Press, Princeton, New Jersey.

Hooten, M.B. & Hefley, T.J. (2019) *Bringing Bayesian Models to Life.* CRC Press, Boca Raton, Florida,

USA.

Huang, H. & Knowles, L.L. (2016) Unforeseen consequences of excluding missing data from next-

generation sequences: Simulation study of RAD sequences. *Systematic Biology,* **65,** 357-365.

Joost, S., Bonin, A., Bruford, M.W., Després, L., Conord, C., Erhardt, G. & Taberlet, P. (2007) A spatial

analysis method (SAM) to detect candidate loci for selection: Towards a landscape genomics

approach to adaptation. *Molecular Ecology,* **16,** 3955-3969.

Kimura, M. (1983) *The Neutral Theory of Molecular Evolution.* Cambridge University Press, Cambridge

UK.

Korolev, K.S., Avlund, M., Hallatschek, O. & Nelson, D.R. (2010) Genetic demixing and evolution in

linear stepping stone models. *Reviews of Modern Physics,* **82,** 1691-1718.

Le Roy, N., Stapane, L., Gautron, J. & Hincke, M.T. (2021) Evolution of the avian eggshell

biomineralization protein toolkit–new insights from multi-omics. *Frontiers in Genetics,* **12,**

672433.

Lin, J., Wang, C. & Redies, C. (2013) Expression of multiple delta-protocadherins during feather bud

formation. *Gene Expression Patterns,* **13,** 57-65.

Lou, R.N., Jacobs, A., Wilder, A.P. & Therkildsen, N.O. (2021) A beginner's guide to low-coverage whole

565   genome sequencing for population genomics. *Molecular Ecology,* **30,** 5966-5993.

566   Lowry, D.B., Hoban, S., Kelley, J.L., Lotterhos, K.E., Reed, L.K., Antolin, M.F. & Storfer, A. (2017)

567   Breaking RAD: An evaluation of the utility of restriction site-associated DNA sequencing for

568   genome scans of adaptation. *Molecular Ecology Resources* **17,** 142–152.

569   Mahoney, S.M., Reudink, M.W., Pasch, B. & Theimer, T.C. (2020) Song but not plumage varies

570   geographically among Willow Flycatcher *Empidonax traillii* subspecies. *Journal of Avian Biology,*

571   **51**.

572   Marcus, J., Ha, W., Barber, R.F. & Novembre, J. (2021) Fast and flexible estimation of effective migration

573   surfaces. *Elife,* **10,** e61927.

574   Mitchell, T.J. & Beauchamp, J.J. (1988) Bayesian variable selection in linear regression. *Journal of the*

575   *American Statistical Association,* **83,** 1023-1032.

576   Neal, R.M. (2011) MCMC using Hamiltonian dynamics. *Handbook of Markov Chain Monte Carlo* (eds S.

577   Brooks, A. Gelman, G. Jones & X.-L. Meng), pp. 2. CRC Press, New York.

578   Olonscheck, D., Schurer, A.P., Lücke, L. & Hegerl, G.C. (2021) Large-scale emergence of regional changes

579   in year-to-year temperature variability by the end of the 21st century. *Nature Communications,* **12,**

580   7237.

581   Pagel, J. & Schurr, F.M. (2012) Forecasting species ranges by statistical estimation of ecological niches

582   and spatial population dynamics. *Global Ecology and Biogeography,* **21,** 293—304.

583   Pardieck, K.L., Ziolkowski, D., Lutmerding, M., Aponte, V. & Hudson, M.R. (2020) North American

584   Breeding Bird Survey Dataset 1966—2019 (ed. U.S.G.S.).

585   Payne, R.B. & Sorenson, M.D. (2002) Museum collections as sources of genetic data. *Bonner Zoologische*

586   *Beiträge,* **51,** 97-104.

587   Piironen, J. & Vehtari, A. (2017) Sparsity information and regularization in the horseshoe and other

588   shrinkage priors. *Electronic Journal of Statistics,* **11,** 5018–5051.

589 Polechová, J. & Barton, N.H. (2015) Limits to adaptation along environmental gradients. *Proceedings of*
590   *the National Academy of Sciences,* **112,** 6401-6406.

591 Rellstab, C., Gugerli, F., Eckert, A.J., Hancock, A.M. & Holderegger, R. (2015) A practical guide to
592   environmental association analysis in landscape genomics. *Molecular Ecology,* **24,** 4348-4370.

593 Rice, S.H. (2004) *Evolutionary Theory: Mathematical and Conceptual Foundations*. Sinauer Associates,
594   MA.

595 Ruegg, K., Bay, R.A., Anderson, E.C., Saracco, J.F., Harrigan, R.J., Whitfield, M., Paxton, E.H. & Smith,
596   T.B. (2018) Ecological genomics predicts climate vulnerability in an endangered southwestern
597   songbird. *Ecology Letters,* **21,** 1085-1096.

598 Satam, H., Joshi, K., Mangrolia, U., Waghoo, S., Zaidi, G., Rawool, S., Thakare, R.P., Banday, S., Mishra,
599   A.K. & Das, G. (2023) Next-generation sequencing technology: current trends and advancements.
600   *Biology,* **12,** 997.

601 Schurr, F.M., Pagel, J., Cabral, J.S., Groeneveld, J., Bykova, O., O'Hara, R.B., Hartig, F., Kissling, W.D.,
602   Linder, H.P., Midgley, G.F., Schröder, B., Singer, A. & Zimmermann, N.E. (2012) How to
603   understand species' niches and range dynamics: a demographic research agenda for biogeography.
604   *Journal of Biogeography,* **39,** 2146-2162.

605 Sedgwick, J.A. (2000) Willow Flycatcher (*Empidonax traillii*). *The Birds of North America,* **533**.

606 Smith, J.M. & Haigh, J. (1974) The hitch-hiking effect of a favourable gene. *Genetics Research,* **23,** 23-35.

607 Smith, T.B., Kinnison, M.T., Strauss, S.Y., Fuller, T.L. & Carroll, S.P. (2014) Prescriptive evolution to
608   conserve and manage biodiversity. *Annual Review of Ecology, Evolution, and Systematics,* **45,** 1-
609   22.

610 Stadler, T. (2010) Sampling-through-time in birth–death trees. *Journal of Theoretical Biology,* **267,** 396-
611   404.

612 Stadler, T., Magnus, C., Vaughan, T., Barido-Sottani, J., Bošková, V., Huisman, J. & Pečerska, J. (2024)
613   *Decoding Genomes: From Sequences to Phylodynamics*. ETH Zurich, Zurich.

Stucki, S., Orozco-terWengel, P., Forester, B.R., Duruz, S., Colli, L., Masembe, C., Negrini, R., Landguth,

E., Jones, M.R. & Consortium, N. (2017) High performance computation of landscape genomic

models including local indicators of spatial association. *Molecular Ecology Resources,* **17,** 1072-

1089.

Suchard, M.A., Lemey, P., Baele, G., Ayres, D.L., Drummond, A.J. & Rambaut, A. (2018) Bayesian

phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evolution,* **4,** vey016.

Theimer, T.C., Smith, A.D., Mahoney, S.M. & Ironside, K.E. (2016) Available data support protection of

the Southwestern Willow Flycatcher under the Endangered Species Act. *The Condor:*

Ornithological Applications, **118,** 289-299.

Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C., Erasmus, B.F.,

De Siqueira, M.F., Grainger, A. & Hannah, L. (2004) Extinction risk from climate change. *Nature,*

427, 145-148.

Thompson, J.N. (2013) *Relentless Evolution*. University of Chicago Press, Chicago.

Unitt, P. (1987) *Empidonax traillii extimus*: an endangered subspecies. *Western Birds,* **18,** 137-162.

Urban, M.C. (2015) Climate change. Accelerating extinction risk from climate change. *Science,* **348,** 571-

573.

Wagner, H.H., Chávez-Pesqueira, M. & Forester, B.R. (2017) Spatial detection of outlier loci with Moran

eigenvector maps. *Molecular Ecology Resources,* **17,** 1122-1135.

Wang, J., Yu, J., Lipka, A.E. & Zhang, Z. (2022) Interpretation of Manhattan plots and other outputs of

genome-wide association studies. *Genome-Wide Association Studies* (eds D. Torkamaneh & F.

Belzile), pp. 63-80. Springer, New York.

Wikle, C.K. (2003) Hierarchical Bayesian models for predicting the spread of ecological processes.

Ecology, **84,** 1382—1394.

Wright, S. (1931) Evolution in Mendelian populations. *Genetics,* **16,** 97-159.