Comparison of genomic assembly and annotation based on two clones of avian pathogenic *Escherichia coli*

Yufei Zhao, John Elmerdahl Olsen, Louise Poulsen and Henrik Christensen[*]

Department of Veterinary and Animal Sciences, University of Copenhagen, Denmark

*Corresponding author. Email: hech@sund.ku.dk

Keywords: open reading frame, false annotations, genome assembly

Comparison of genomic assembly and annotation based on two

clones of avian pathogenic *Escherichia coli*

Yufei Zhao, John Elmerdahl Olsen, Louise Poulsen and Henrik Christensen

Abstract

Methods for assembly and annotation of whole genomic sequences were compared

for six strains of avian pathogenic *Escherichia coli* (APEC). Two vertically transferred

E. coli clones, represented by three isolates all belonging to pulse field genome

electrophoresis (PFGE) type 65- sequence type (ST)95 and three isolates belonging

to PFGE type 47- ST131, were selected for Illumina short read sequencing. There

was no significant difference between SPAdes and CLC Genomic Workbench for

benchmark parameters to assemble the short reads. The six strains were also

sequenced by long read sequencing (Nanopore) and these reads were hybrid

assembled with the short reads. Unicycler provided a lower number of contigs and

higher N50 compared to Flye. No significant differences between total length of

genomes were obtained from the four assemblers. At least 2.1 and 0.9% of coding

gene sequences (CDSs) annotated with RAST and PROKKA, respectively were

wrongly annotated. The errors were most often associated to CDS of shorter length

(< 150 nt) with functions such as transposases, mobile genetic elements or being

hypothetical. The investigation points out the importance of controlling automatic

annotations and suggest further work to improve annotations in strains not belonging

to the K12 or B lineages.

1. Introduction

High quality whole genomic sequences are needed for accurate annotation,

identification of plasmids, to trace bacterial clones and more general to provide the

basic information for modern investigation of bacteriology. The Illumina sequencing

platforms can generate high accuracy (>99.99%) raw reads with a relatively low cost

[1] and enable extraction of information of multi locus sequence type (MLST),

prediction of virulence associated genes (VAGs), antibiotic resistance genes (ARGs)

and provide clonal comparison by single nucleotide polymorphism (SNP) analysis [2].

However, the pair-end raw reads generated are often not longer than 250 to 300

base pairs, which makes it difficult to fully close a bacterial genome leaving several

incomplete contiguous sequences (contigs). The fragmented contigs may hinder the

determination of whether VAGs and ARGs are located on chromosomes or mobile

genetic elements. Another problem is the lack of knowledge of contig orientation and

orientation of annotation of structural repetitive regions and larger rearrangement

such as transposons, insertion sequences and genetic cassette. The repeated

sequences are often larger than the sequencing length of the Illumina reads making

assembly impossible [1, 3]. The long read sequencing platforms, such as Pacific

Biosciences (PacBio) and Oxford Nanopore Technologies (ONT) can generate

single DNA reads with a median length of 10k bp and up to 100k bp [1, 4], and thus

can be a great improvement to correct sequences generated with Illumina

technology with respect to the repetitive and other complex regions on genomes [5].

Although the raw read accuracy can go up to 99.5% with the new R10.4.1 flow cell

and V14 kit according to ONT (https://nanoporetech.com/), hybrid assemblies with a

combination of the accuracy of short read sequence and the completeness of long

read sequencing still provide the most reliable and cost-effective approach to

reconstruct bacterial genomes [6, 7, 8].

Two of the most common pipelines for annotation are RAST [9, 10] and Prokka [11].

RAST predicts open reading frames (ORFs) based on GLIMMER 3.0 [12], combined

with protein families "subsystems" as well as other tools and iterations. GLIMMER

3.0 is based on interpolated Markov model (IMM) search where ORFs are identified

in reverse from the stop codon and back towards the start codon. Further

optimization is by optimization for ribosomal binding sites using a position weight

matrix improving the prediction of translation initiation sites. Prokka [11] uses

Prodigal [13] to identify ORFs based on a dynamic programming approach

optimizing start and start and stop codon identification using prediction of ribosome

binding sites, GC contents, hexamer distributions as well as iterations of the process.

The aim of the study was to investigate the influence of assembly and annotation

methods (RAST, Prokka) using *E. coli* strains of the avain pathogenic type (APEC)

as model organisms. The comparison of closely related strains allowed the

identification of annotation errors associated with both RAST and Prokka.

2. Material and methods

2.1 Sample collection and molecular typing

Six APEC strains were chosen for the current study. They represented two APEC clones of sequence type (ST) 95, pulse field gel electrophoresis type 65 and ST131, PFGE type (47), respectively [14] (Table 1).

2.2 DNA extraction

E. coli isolates were cultured from stocks stored frozen in glycerol at - 80 °C. Cultivation was initiated on blood agar (Tryptose blood agar base Difco, Brøndby, Denmark) added 5% bovine blood with subsequent incubation at 37 °C for 24 h. A single colony from each plate was cultured in Luria-Bertani Broth (LB) at 37 °C overnight. The DNA was extracted with DNeasy Blood&Tissue Kit (QIAGEN) and Maxwell RSC Cultured Cells DNA kit (Promega, Madison, USA) according to manufacturer's instructions. The purity and concentration of extracted DNA were evaluated by Nanodrop spectrophotometer (Thermo Fisher Scientific) and Qubit fluorometer (Thermo Fisher Scientific). The integrity of DNA was determined by gel-electrophoresis. DNA extracted from the isolates was used for both Illumina and ONT sequencing.

2.3 Library preparation and sequencing

DNA Libraries for Illumina sequencing were prepared using Illumina DNA Prep (Illumina, SanDiego, CA) and sequence reactions were prepared with Miseq Reagent kit v2 (Illumina, SanDiego, CA). Samples were sequenced by the Illumina MiSeq platform using 2 × 250 bp paired end sequencing strategy. In parallel, strains were sequenced by ONT on a MinION Mk1B sequencer (ONTechnology, Oxford). Libraries of selected DNA samples were prepared by using Rapid Barcoding Kit 96 V14 (SQK-RBK114, ONT), and were loaded on one MinION Flow Cell (R.10.4.1 FLO-MIN114, ONT) for sequencing for 48 h with an accurate model (260 bps). MinKNOW software v5.4.3 was used to perform the sequencing and data acquisition. Post-run base-calling and demultiplexing were performed with Guppy v6.4.8 (https://nanoporetech.com/document/Guppy-protocol, accessed 20 February 2025).

2.4 Raw reads quality control and trimming

Quality control, adapter removal and short reads filtering of Illumina reads were performed on fastp v0.12.4 for Linux with default parameter settings [15]. For ONT long reads, the short and low-quality reads were filtered by Filtlong v0.2.1 (https://github.com/rrwick/Filtlong) and quality control of long reads were assessed using SeqKit v2.5.1 [16].

2.5 Genome assembly and annotation

SPAdes v3.13.1 was used with a '--careful' parameter for genome assemblies [17] for short reads generated by Illumina. Contigs smaller than 1000 bp were removed from the assembled genomes. CLC Genomic Workbench 22 (QIAGEN) was used for short reads assemblies to compare the assembly quality with SPAdes. Hybrid assemblies were performed by two workflows: Unicycler v0.5.0 was used directly with Illumina short reads and corresponding Nanopore long reads [4, 18]. The other approach used Flye v2.9.2 [19] to do a long-read assembly first, then the draft assemblies were polished with medaka v1.8.1 [20] and further improved with short reads by Pilon v1.2.2 [21]. Quast v5.0.2 [22] (Gurevich et al. 2013) and BUSCO v5.5.0 [23] were used to evaluate the quality of assemblies. Annotation was performed with Prokka v1.14.6 with '--force --centre X –compliant' parameters [11] and the online toolbox Rapid Annotation using Subsystem Technology (RAST) [10].

2.6 Comparison of annotations

To control the differences among annotations, gene presence and absence matrices were established for RAST using Seed Viewer, and for Prokka using Roary. A Python script was written by chatGPT and used to parse the differential genes from matrices and extract corresponding sequences from original FASTA files. The sequences of differential genes were integrated into one Excel file. The sequences were aligned by BLASTn against the whole genome sequence of strains which did not annotate these genes within each clone. Different genes annotated with more than 95% identity were considered as annotation errors if they were annotated to the same DNA sequence. To illustrate selected regions, comparison of BLASTn alignments of differential region were performed using Clinker (https://cagecat.bioinformatics.nl/tools/clinker). The Venn diagrams showing

relationships between Prokka and RAST annotations were generated on

https://bioinformatics.psb.ugent.be/webtools/Venn/.

3. Results and discussion

3.1 Statistics of short read and hybrid assemblies

For the six strains, the average of read length of Illumina sequencing was 197.3 bp

(range from 162.8 -262.4 bp) and 3909 bp (range from 2710 - 5129 bp) of ONT

reads (Table 2). The average number of reads was 622,721 bp for Illumina and

109,933 bp for ONT. The coverage depth for all Illumina raw reads exceeded 30×

with an average of 54×, while the ONT coverages ranged from 64× with an average

coverage of 94× (Table 2). These parameters satisfy the common quality standards

used for WGS assembly [24, 25, 26].

QUAST and BUSCO were used to test the assembly performance of the commonly

used software CLC Genomic Workbench and SPAdes for short reads assembly and

Flye, and Unicycler for hybrid assemblies (Table 3). The number of contigs did not

differ significantly between CLC Genomic Workbench and SPAdes (Fisher's exact

test, P = 0.819). For hybrid assemblies, Unicycler provided lower number of contigs

than Flye in both clones (P = 0.0003, Fisher's exact test). NG50 is the shortest contig

length of an assembly which is 50% of the reference genome or longer. N50 did not

differ significantly between SPAdes and CLC (Fisher's exact test, P = 0.768),

however, longer N50 was obtained by Unicycler than from Flye (Fisher's exact test,

P = 0.0002). The total length of genomes did not differ significantly between CLC

Genomics Workbench, SPAdes, Flye and Unicycler.

Considering the performance of Unicycler in giving lower number of contigs and

higher NG50 values, it was decided to be the best assembler for downstream

annotation.

3.2 Comparison of annotations for hybrid assemblies

Annotations were compared for RAST and Prokka (Table 3). Identical number of

rRNA genes (n = 22) were predicted by both annotation methods in all strains. The

number of tRNA genes differed slightly. One tmRNA was annotated in all strains by

Prokka compared to none by RAST. Different numbers of coding DNA sequences

(CDS) were found for the hybrid genomes when annotated by RAST and Prokka.

Overall, Prokka annotated less (187-327) CDSs compared to RAST (Table 4). RAST

predicted that strains within clone PFGE65-ST95 differed by 88 CDS and strains of

PFGE47-ST131 by 235 CDS. Meanwhile, Prokka predicted 47 different CDS in

PFGE65-ST95 and 200 in PFGE47-ST131 (Fig. 1 A C).

The reasons for the different numbers of CDS annotated by RAST and Prokka were

analysed. Annotation comparison was performed based on presence and absence

matrix generated from both RAST and Prokka. It was then surprising to find

divergent annotations between strains for exactly the same DNA sequences. For

RAST annotations of strains of the PFGE65-ST95 clone, 71 out of the 88 genes

annotated differently between the three strains got 100% identity in complementary

genomes, when controlled on sequence level showing that they were error prone

annotations. Another 14 genes got more than 97% identity for at least one strain by

BLASTn leaving three real gene differences in 729_270514_x_12 since the strain did

not harbor the 2312 bp plasmid with three CDSs (Fig. 1 A B).

Similar problematic annotations were obtained among PFGE65-ST95 genomic

sequences annotated by Prokka, in which 13 of 47 genes annotated differently within

the three strains of PFGE65-ST95 got hits of 100% identity. Another 32 genes got

more than 97% identity and one gene got 94.7% identity. These 46 genes were

regarded as error prone annotations with Prokka (Fig. 1 A-B).

In PFGE47-ST131, with RAST annotation, 120 out of 235 genes annotated

differently got 100% identity in whole genome sequences which were not annotated

for these genes and three gene differences were identified the same way with more

than 98% identity. All 123 annotations were considered error prone. Ninety-eigth real

gene differences were related to contig 3 identified as a plasmid where

723_010814_2_1 and 723_220514_1_6 carried a 6668-bp *ColRNAI* related plasmid

and 729_141114_1_15 carried a 100,533-bp Incl-1 plasmid. Another 14 real

differences originated in phage related insertions into the chromosome of

729_141114_1_15 (Fig. 1 C – D).

For PFGE47-ST131 genomic sequences annotated with Prokka, 25 of 200 different
genes got 100% identity in reciprocal sequences which were not annotated for these
genes and 20 different genes got more than 97% identity. These 45 annotations
were considered error prone. Similar to RAST annotation, 141 gene differences were
caused by different genes on contig 3 (unique IncI-1 plasmid), and 14 annotation
differences related to prophage regions in the chromosome of strain
729_141114_1_15 (Fig. 1 C – D).

Fig. 2 shows examples of inaccurately annotated genes and their flanking regions
visualized by Clinker. The *neuO* gene (polysialic acid O-acetyltransferase) was only
annotated in strain 729_270514_x_12 and not in the other two strains although they
had identical DNA sequences in this region (A). In 729_141114_1_15, the duplicate
annotation for the *pdeL* gene as *pdeL* and *pdeC* in strain 729_141114_1_15
compared to the other two strains was found to be an error since all three strains
shared 100% sequence identity (B). The last example is with two copies of the
transposase IS*Cep1* gene in two strains which were only annotated as one in
729_141114_1_15. Also here regions compared on the figure shared 100%
sequence identity.

In all, 2.1 and 0.9% of CDSs annotated with RAST and PROKKA, respectively were
error prone. The errors were most often associated to CDSs of shorter length (< 150
nt) with functions annotated as transposases, mobile genetic elements or
hypothetical. These estimates are only based on reciprocal differences between the
annotations and may be underestimated and not included CDSs wrongly annotated
in all strains compared.

The divergence associated with annotations of identical DNA sequence regions falls
back on the annotating methods. The methods and procedures in RAST and Prokka
are quite different. RAST predicts ORFs based on GLIMMER 3.0 where ORFs are
predicted by IMM search in reverse from the stop codon and back towards the start
codon followed by optimization for ribosomal binding sites improving the prediction of
translation initiation sites. Optimizations from previous versions of the program
included in particular the prediction of translation initiation sites and one reason for

inaccurate ORF prediction can be that further optimization of this step is needed.

Prokka uses Prodigal [13] to identify ORFs based on a dynamic programming

approach optimizing start and start and stop codon identification by prediction of

ribosome binding sites, GC contents, hexamer distributions as well as iterations of

the process. A lower limit of 90 bp is used in the program to score start-stop codon

pairs [13] and this may be some of the reason for the errors observed for shorter

genes in the current study. Another reason for errors with Prodigal may be the use of

a window of 5 kb to identify start and stop codons. In the current comparison it can

be anticipated that sequence divergence between two strains in such a window may

lead to different annotations at local level even though sequenced are identical for

this region. Hyatt et al. [13] found that Glimmer 3 predicted around 4% more ORFs in

E. coli K12 compared to Prodigal which is in line with the current study. A problem

when comparing annotations is the lack of a golden standard. Such a standard could

have been obtained from *E. coli* strain K12 variant MG1655 where annotation has

been refined for decades for instance by EcoGene [27, 28]. Unfortunately this

resource seems no longer maintained and accessible. Considering the high genomic

variation between reference and field strain of *E. coli* a full comparison would not

have been possible. Further optimization of the search parameters in the annotation

programs may be needed and targeted better towards *E. coli* and further curation of

annotations needed.

274   4. Conclusion

There was no significant difference between SPAdes and CLC Genomic Workbench

for benchmark parameters to assemble short reads. However, for hybrid assemblies,

Unicycler provided lower number of contigs and higher N50 than Flye. In the

comparisons, at least 2.1 and 0.9% of genes annotated with RAST and PROKKA,

respectively were false annotations stressing the importance of controlling the

annotations. This investigation showed that strains with closely related genomes at

DNA level obtained differences in annotation for local regions with identical DNA

sequences. The problem was identified with the most commonly used annotation

programs RAST and Prokka. The errors are related to the statistical methods used

for prediction of ORFs where differences in up or downstream regions may influence

the prediction.

CRediT authorship contribution statement

Yufei Zhao: Formal analysis, Investigation, Writing – original draft. Henrik

Christensen: Strains, Formal analysis, Investigation, Writing – original draft. Louise

Ladefoged Poulsen: Strains, Writing – review & editing. John Elmerdahl Olsen:

Investigation, Resources, Conceptualization, Writing – review & editing, Supervision.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Funding information

This study was partially funded by a State Scholarship Fund to Yufei Zhao from

China Scholarship Council (CSC No. 202206350041).

Repositories

GenBank under accession numbers in Table 1.

Acknowledgements

Xiao Fei and Yaovi Mahuton Gildas Hounmanou are thanked for help with the

analysis of Nanopore sequencing data.

References

[1] De Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J et al.

Comparison of long-read sequencing technologies in the hybrid assembly of

complex bacterial genomes. *Microb Genom* 2019; 5: e000294.

[2] Christensen H, Bachmeier J, Bisgaard M. New strategies to prevent and control

avian pathogenic *Escherichia coli* (APEC). *Avian Pathol* 2021; 50:370-381.

[3] Therrien DA, Konganti K, Gill JJ, Davis BW, Hillhouse AE, Michalik J et al.

Complete whole genome sequences of *Escherichia coli* surrogate strains and

comparison of sequence methods with application to the food industry.

Microorganisms 2021; 9: 608.

[4] Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome

assemblies from short and long sequencing reads. *PLoS Comput Biol* 2017; 13:

e1005595.

[5] Maguire M, Kase JA, Roberson D, Muruvanda T, Brown EW, Allard M et al.

Precision long-read metagenomics sequencing for food safety by detection and

assembly of Shiga toxin-producing *Escherichia coli* in irrigation water. *PloS One*

2021; 16: e0245172.

[6] Khezri A, Avershina E, Ahmad R. Hybrid assembly provides improved resolution

of plasmids, antimicrobial resistance genes, and virulence factors in *Escherichia coli*

and *Klebsiella pneumonia* clinical isolates. *Microorganisms* 2021; 9:2560.

[7] Craddock HA, Motro Y, Zilberman B, Khalfin B, Bardenstein S, Moran-Gilad J.

Long-Read sequencing and hybrid assembly for genomic analysis of clinical *Brucella*

melitensis isolates. *Microorganisms* 2022; 10:619.

[8] Sanderson H, McCarthy MC, Nnajide CR, Sparrow J, Rubin JE, Dillon JR, White

AP. Identification of plasmids in avian-associated Escherichia coli using nanopore

and illumina sequencing. *BMC Genomics* 2023; 24: 698. doi: 10.1186/s12864-023-

09784-6.

[9] Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K,

Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL,

Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C,

Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. The RAST Server: rapid

annotations using subsystems technology. *BMC Genomics* 2008; 9: 75. doi:

10.1186/1471-2164-9-75.

[10] Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T et al. The SEED

and the rapid annotation of microbial genomes using subsystems technology (RAST).

Nucleic Acids Res 2014; 42(D1): D206-D214.

[11] Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;

30: 2068-2069.

[12] Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes

and endosymbiont DNA with Glimmer. *Bioinformatics* 2007; 23: 673-9. doi:

10.1093/bioinformatics/btm009.

[13] Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:

prokaryotic gene recognition and translation initiation site identification. *BMC*

Bioinformatics 2010;11:119. doi: 10.1186/1471-2105-11-119.

[14] Poulsen LL, Thøfner I, Bisgaard M, Christensen JP, Olsen RH, Christensen H.

Longitudinal study of transmission of *Escherichia coli* from broiler breeders to

broilers. *Vet Microbiol* 2017; 207:13-18.

[15] Chen S. Ultrafast one-pass FASTQ data preprocessing, quality control, and

deduplication using fastp. *iMeta* 2023; 2: e107.

[16] Shen W, Le S, Li Y, Hu F. SeqKit: a cross-platform and ultrafast toolkit for

FASTA/Q file manipulation. *PloS One* 2016; 11: e0163962.

[17] Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes

de novo assembler. *Curr Protoc Bioinformatics* 2020; 70: e102.

[18] Antipov D, Korobeynikov A, McLean JS, Pevzner PA. hybridSPAdes: an

algorithm for hybrid assembly of short and long reads. *Bioinformatics* 2016; 32:

1009-1015.

[19] Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads

using repeat graphs. *Nat Biotechnol* 2019; 37: 540-546.

[20] Oxford Nanopore Technologies 2022. Medaka. Available online:

https://github.com/nanoporetech/medaka (accessed on 17 October 2023).

[21] Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S et al. Pilon:

an integrated tool for comprehensive microbial variant detection and genome

assembly improvement. *PloS One* 2014; 9: e112963.

[22] Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* 2013; 29: 1072-1075.

[23] Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO update:

novel and streamlined workflows along with broader and deeper phylogenetic

coverage for scoring of eukaryotic, prokaryotic, and viral genomes. *Mol Biol Evol*

2021; 38: 4647-4654.

[24] Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, da Costa MS, Rooney

AP, Yi H, Xu XW, De Meyer S, Trujillo ME. Proposed minimal standards for the use

of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 2018; 68:

461-466. doi: 10.1099/ijsem.0.002516.

[25] European Food Safety Authority (EFSA). EFSA statement on the requirements

for whole genome sequence analysis of microorganisms intentionally used in the

food chain. *EFSA J* 2024; 22 :e8912. doi: 10.2903/j.efsa.2024.8912.

[26] Riesco R, Trujillo ME. Update on the proposed minimal standards for the use of

genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 2024; 74:

006300. doi: 10.1099/ijsem.0.006300.

[27] Zhou J, Rudd KE. EcoGene 3.0. *Nucleic Acids Res* 2013; 41(Database issue):

D613-24. doi: 10.1093/nar/gks1235.

[28] Rudd KE. EcoGene: a genome sequence database for Escherichia coli K-12.

Nucleic Acids Res. 2000 Jan 1;28(1):60-4. doi: 10.1093/nar/28.1.60.

Figure legends

Fig. 1. Venn diagrams illustrating the deviating annotations between Prokka and

RAST. For PFGE65-ST95, (A) shows the number of different genes predicted before

BLASTn control and (B) shows the numbers after control. For PFGE47-ST131, (C)

shows the number of different genes predicted before BLASTn control and (D)

shows the numbers after control.

Fig. 2. Examples of genetic comparisons of inaccurately annotated genes from

Prokka and their flanking regions. An additional annotation is given for the *neuO*

gene in 729_270514_x_12 (A). In 729_141114_1_15, duplicate annotation for the

pdeL gene as *pdeL* and *pdeC* was found (B). Two copies of the transposase IS*Cep1*

gene in two strains were only annotated as one in strain 729_141114_1_15 (C). The

regions compared all shared 100% DNA sequence identity without gaps.

Table 1. Strains investigated, genome length, genetic elements from six hybrid assembled genomes.

Strain	Origin	Organ/lesion	PFGE-ST	Hybrid assembly Contig number	Contig length (bp)	INDSC acc. no. Hybrid assembly	Short read assembly	Matching genomic elements
729_021014_1_4	Broiler parent	Salpingitis	65-95	1	5,232,657	CP160125	JBELNO01	Chromosome
				2	193,604	CP160126		Plasmid [a]
				3	6655	CP160127		Plasmid
				4	2676	CP160128		Plasmid
				5	2312	CP160129		Plasmid
729_270514_x_12	Broiler parent	Salpingitis	65-95	1	5,232,714	CP160121	JBELNS01	Chromosome
				2	193,604	CP160122		Plasmid [a]
				3	6655	CP160123		Plasmid
				4	2677	CP160124		Plasmid
729_xx_1_3_yolk	Broiler	Yolk sac infection	65-95	1	5,232,647	CP160116	JBELNM01	Chromosome
				2	193,604	CP160117		Plasmid [a]
				3	6655	CP160118		Plasmid
				4	2676	CP160119		Plasmid
				5	2312	CP160120		Plasmid
723_010814_2_1	Broiler parent	Salpingitis	47-131	1	4,866,047	CP160112	JBELNC01 (SSQU01)	Chromosome
				2	201,035	CP160113		Plasmid [a]
				3	6668	CP160114		Plasmid
				4	1552	CP160115		Plasmid
729_141114_1_15	Broiler parent	Salpingitis	47-131	1	4,887,791	CP160108	JBELNB01	Chromosome
				2	201,054	CP160109		Plasmid [a]
				3	100,533	CP160110		Plasmid
				4	1552	CP160111		Plasmid
723_220514_1_6	Broiler	Liver/septicaemia	47-131	1	4,866,598	CP160104	JBELNH01	Chromosome
				2	200,992	CP160105		Plasmid [a]
				3	6668	CP160106		Plasmid
				4	1552	CP160107		Plasmid

[a] ColV-like

Table 2. Statistical analysis of reads.

ST	Strain	Illumina				ONT			
		Mean read length (bp)	Number of reads	Total bases (Mbp)	Coverage (x)*	Mean read length (bp)	Number of reads	Total bases (Mbp)	Coverage (x)
95	729_021014_1_4	184.9	515,309	154	41.1	5129	137,504	664	151.9
	729_270514_x_12	190.5	613,533	186	50.3	4138	84,953	336	75.7
	729_xx_1_3_yolk	162.8	594,174	150	41.7	4260	95,064	387	87.2
131	723_010814_2_1	262.4	800,953	283	90.5	2710	109,803	283	64.1
	729_141114_1_15	176.2	593,386	167	45.0	3832	138,333	508	114.2
	723_220514_1_6	207.2	618,972	186	55.2	3386	93,939	301	68.5
	Average	197.3	622,721	188	54.0	3909	109,933	413	93.6

*coverage = (number of read × read length) / total genome size. In which total genome size was estimated to 5.2 Mbase.

Table 3. The quality of assemblies generated by CLC Genomic Workbench and SPAdes for short-read only assemblies and Flye and Unicycler for hybrid assemblies.

ST	Strain	Number of contigs				NG50 (bp)				Total length (bp)				Completeness (%)			
		CLC	SPAdes	Flye	Unicycler	CLC	SPAdes	Flye	Unicycler	CLC	SPAdes	Flye	Unicycler	CLC	SPAdes	Flye	Unicycler
95	729_021014__1_4	110	99	5	5	169,077	196,246	5,232,681	5,232,657	5,310,368	5,312,171	5,444,670	5,437,904	100	100	100	99.8
	729_270514_x_12	100	100	5	4	211,910	204,287	5,232,730	5,232,714	5,313,797	5,314,927	5,449,369	5,435,650	100	99.8	100	99.8
	729_xx_1_3_yolk	143	132	6	5	135,314	164,290	5,232,625	5,232,647	5,307,620	5,316,910	5,446,817	5,437,894	100	100	100	99.8
Average		117.7	110.3	5.3	4.7	172,100.3	188,274.3	5,232,678,7	5,232,672.7	5,310,595.0	5,314,669.3	5,446,952,0	5,437,149.3	100.0	99.9	100.0	99.8
131	723_010814_2_1	46	45	16	4	354,765	353,089	1,074,415	4,866,047	5,033,610	5,022,129	5,076,852	5,075,302	100	100	100	99.8
	729_141114_1_15	88	88	5	4	192,789	224,060	4,949,514	4,887,791	5,110,669	5,115,410	5,164,053	5,190,930	100	99.8	100	99.8
	723_220514_1_6	51	50	8	4	354,390	353,089	2,855,420	4,866,598	5,021,087	5,022,103	5,083,114	5,075,810	100	100	100	99.8
Average		61.7	61.0	9.7	4.0	300,648.0	310,079.3	2,959,783.0	4,873,478.7	5,055,122.0	5,053,214.0	5,108,006,3	5,114,014.0	100.0	99.9	100.0	99.8

Table 4. Statistical analysis of annotations.

ST	Strain	CDS		tRNA		tmRNA		Total *RNA		Number of CDS differences	
		RAST	Prokka	RAST	Prokka	RAST	Prokka	RAST	Prokka	RAST	Prokka
95	729_021014_1_4	5476	5149	90	92	0	1	112	115	88	47
	729_270514_x_12	5450	5263	90	92	0	1	112	115		
	729_xx_1_3_yolk	5442	5148	90	92	0	1	112	115		
131	723_220514_1_6	4967	4699	84	86	0	1	106	109	235	200
	723_010814_2_1	4958	4698	84	86	0	1	106	109		
	729_141114_1_15	5094	4830	85	86	0	1	107	109		

*22 rRNA genes in all strains for both annotation methods.

Fig. 1A

Fig. 1B

Fig. 1C

Fig. 1D

Fig. 2A

Fig. 2B

Fig. 2C