Comparison of genomic assembly and annotation based on two

clones of avian pathogenic *Escherichia coli*

Yufei Zhao[1], John Elmerdahl Olsen[1], Louise Poulsen[1] and Henrik Christensen[1*]

[1] Department of Veterinary and Animal Sciences, University of Copenhagen,

Denmark

*Corresponding author. Email: hech@sund.ku.dk

Abstract

Methods for assembly and annotation of whole genomic sequences were compared

for six strains of avian pathogenic *Escherichia coli* (APEC). Two vertically transferred

E. coli clones, represented by three isolates all belonging to pulse field genome

electrophoresis (PFGE) type 65- sequence type (ST)95 and three isolates belonging

to PFGE type 47- ST131, were selected for Illumina short read sequencing. There

was no significant difference between SPAdes and CLC Genomic Workbench for

benchmark parameters to assemble the short reads. The six strains were also

sequenced by long read sequencing (Nanopore) and these reads were hybrid

assembled with the short reads. Unicycler provided a lower number of contigs and

higher N50 compared to Flye. No significant differences between total length of

genomes were obtained from the four assemblers. At least 2.1 and 0.9% of coding

gene sequences (CDSs) annotated with RAST and PROKKA, respectively were

wrongly annotated. The errors were most often associated to CDS of shorter length

(< 150 nt) with functions such as transposases, mobile genetic elements or being

hypothetical. The investigation points out the importance of controlling automatic

annotations and suggest further work to improve annotations in strains not belonging

to the K12 or B lineages.

Introduction

High quality whole genomic sequences are needed for accurate annotation,

identification of plasmids, to trace bacterial clones and more general to provide the

basic information for modern investigation of bacteriology. The Illumina sequencing

platforms can generate high accuracy (>99.99%) raw reads with a relatively low cost

(De Maio et al. 2019) and enable extraction of information of multi locus sequence

type (MLST), prediction of virulence associated genes (VAGs), antibiotic resistance

genes (ARGs) and provide clonal comparison by single nucleotide polymorphism

(SNP) analysis (Christensen et al. 2021). However, the pair-end raw reads

generated are often not longer than 250 to 300 base pairs, which makes it difficult to

fully close a bacterial genome leaving several incomplete contiguous sequences

(contigs). The fragmented contigs may hinder the determination of whether VAGs

and ARGs are located on chromosomes or mobile genetic elements. Another

problem is the lack of knowledge of contig orientation and orientation of annotation of

structural repetitive regions and larger rearrangement such as transposons, insertion

sequences and genetic cassette. The repeated sequences are often larger than the

sequencing length of the Illumina reads making assembly imporrible (De Maio et al.

2019, Therrien et al. 2021). The long read sequencing platforms, such as Pacific

Biosciences (PacBio) and Oxford Nanopore Technologies (ONT) can generate

single DNA reads with a median length of 10k bp and up to 100k bp (De Wick et al.

2017, Maio et al. 2019), and thus can be a great improvement to correct sequences

generated with Illumina technology with respect to the repetitive and other complex

regions on genomes (Maguire et al. 2021). Although the raw read accuracy can go

up to 99.5% with the new R10.4.1 flow cell and V14 kit according to ONT

(https://nanoporetech.com/), hybrid assemblies with a combination of the accuracy of

short read sequence and the completeness of long read sequencing still provide the

most reliable and cost-effective approach to reconstruct bacterial genomes (Khezri et

al. 2021, Craddock et al. 2022, Sanderson et al. 2023).

Two of the most common pipelines for annotation are RAST (Aziz et al. 2008,

Overbeek et al. 2014) and Prokka (Seemann 2014). RAST predicts open reading

frames (ORFs) based on GLIMMER 3.0 (Delcher et al. 2007), combined with protein

families "subsystems" as well as other tools and iterations. GLIMMER 3.0 is based

on interpolated Markov model (IMM) search where ORFs are identified in reverse

from the stop codon and back towards the start codon. Further optimization is by

optimization for ribosomal binding sites using a position weight matrix improving the

prediction of translation initiation sites. Prokka (Seemann 2014) uses Prodigal (Hyatt

et al. 2010) to identify ORFs based on a dynamic programming approach optimizing

start and start and stop codon identification using prediction of ribosome binding

sites, GC contents, hexamer distributions as well as iterations of the process.

The aim of the study was to investigate the influence of assembly and annotation

methods (RAST, Prokka) using *E. coli* strains of the avain pathogenic type (APEC)

as model organisms. The comparison of closely related strains allowed the

identification of annotation errors associated with both RAST and Prokka.

Methods

Sample collection and molecular typing

Six APEC strains were chosen for the current study. They represented two APEC

clones of sequence type (ST) 95, pulse field gel electrophoresis type 65 and ST131,

PFGE type (47), respectively (Poulsen et al. 2017) (Table 1).

DNA extraction

E. coli isolates were cultured from stocks stored frozen in glycerol at - 80 °C.

Cultivation was initiated on blood agar (Tryptose blood agar base Difco, Brøndby,

Denmark) added 5% bovine blood with subsequent incubation at 37 °C for 24 h. A

single colony from each plate was cultured in Luria-Bertani Broth (LB) at 37 °C

overnight. The DNA was extracted with DNeasy Blood&Tissue Kit (QIAGEN) and

Maxwell RSC Cultured Cells DNA kit (Promega, Madison, USA) according to

manufacturer's instructions. The purity and concentration of extracted DNA were

evaluated by Nanodrop spectrophotometer (Thermo Fisher Scientific) and Qubit

fluorometer (Thermo Fisher Scientific). The integrity of DNA was determined by gel-

electrophoresis. DNA extracted from the isolates was used for both Illumina and

ONT sequencing.

Library preparation and sequencing

DNA Libraries for Illumina sequencing were prepared using Illumina DNA Prep

(Illumina, SanDiego, CA) and sequence reactions were prepared with Miseq

Reagent kit v2 (Illumina, SanDiego, CA). Samples were sequenced by the Illumina

MiSeq platform using 2 × 250 bp paired end sequencing strategy. In parallel, strains

were sequenced by ONT on a MinION Mk1B sequencer (Oxford Nanopore

Technology, Oxford). Libraries of selected DNA samples were prepared by using

Rapid Barcoding Kit 96 V14 (SQK-RBK114, Oxford Nanopore Technology), and

were loaded on one MinION Flow Cell (R.10.4.1 FLO-MIN114, Oxford Nanopore

Technology) for sequencing for 48 h with an accurate model (260 bps). MinKNOW

software v5.4.3 was used to perform the sequencing and data acquisition. Post-run

base-calling and demultiplexing were performed with Guppy v6.4.8.

Raw reads quality control and trimming

Quality control, adapter removal and short reads filtering of Illumina reads were

performed on fastp v0.12.4 for Linux with default parameter settings (Chen et al.

2023). For ONT long reads, the short and low-quality reads were filtered by Filtlong

v0.2.1 (https://github.com/rrwick/Filtlong) and quality control of long reads were

assessed using SeqKit v2.5.1 (Shen et al. 2016).

Genome assembly and annotation

SPAdes v3.13.1 was used with a '--careful' parameter for genome assemblies

(Prjibelski et al. 2020) for short reads generated by Illumina. Contigs smaller than

1000 bp were removed from the assembled genomes. CLC Genomic Workbench 22

(QIAGEN) was used for short reads assemblies to compare the assembly quality

with SPAdes. Hybrid assemblies were performed by two workflows: Unicycler v0.5.0

was used directly with Illumina short reads and corresponding Nanopore long reads

(Antipov et al. 2016, Wick et al. 2017). The other approach used Flye v2.9.2

(Kolmogorov et al. 2019) to do a long-read assembly first, then the draft assemblies

were polished with medaka v1.8.1 (Oxford Nanopore Technologies. 2022)

(https://github.com/nanoporetech/medaka) and further improved with short reads by

Pilon v1.2.2 (Walker et al. 2014). Quast v5.0.2 (Gurevich et al. 2013) and BUSCO

v5.5.0 (Manni et al. 2021) were used to evaluate the quality of assemblies.

Annotation was performed with Prokka v1.14.6 with '--force --centre X –compliant'

parameters (Seemann 2014) and the online toolbox Rapid Annotation using

Subsystem Technology (RAST) (Overbeek et al. 2014).

Comparison of annotations

To control the differences among annotations, gene presence and absence matrices were established for RAST using Seed Viewer, and for Prokka using Roary. A Python script was written by chatGPT and used to parse the differential genes from matrices and extract corresponding sequences from original FASTA files. The sequences of differential genes were integrated into one Excel file. The sequences were aligned by BLASTn against the whole genome sequence of strains which did not annotate these genes within each clone. Different genes annotated with more than 95% identity were considered as annotation errors if they were annotated to the same DNA sequence. To illustrate selected regions, comparison of BLASTn alignments of differential region were performed using Clinker (https://cagecat.bioinformatics.nl/tools/clinker). The Venn diagrams showing relationships between Prokka and RAST annotations were generated on https://bioinformatics.psb.ugent.be/webtools/Venn/.

Results and discussion

Statistics of short read and hybrid assemblies

For the six strains, the average of read length of Illumina sequencing was 197.3 bp (range from 162.8 -262.4 bp) and 3909 bp (range from 2710 - 5129 bp) of ONT reads (Table 2). The average number of reads was 622,721 bp for Illumina and 109,933 bp for ONT. The coverage depth for all Illumina raw reads exceeded 30✕ with an average of 54✕, while the ONT coverages ranged from 64✕ with an average coverage of 94✕ (Table 2). These parameters satisfy the common quality standards used for WGS assembly (Chun et al. 2018, EFSA 2024, Riesco and Trujillo, 2024).

QUAST and BUSCO were used to test the assembly performance of the commonly used software CLC Genomic Workbench and SPAdes for short reads assembly and Flye, and Unicycler for hybrid assemblies (Table 3). The number of contigs did not differ significantly between CLC Genomic Workbench and SPAdes (Fisher's exact test, P = 0.819). For hybrid assemblies, Unicycler provided lower number of contigs than Flye in both clones (P = 0.0003, Fisher's exact test). NG50 is the shortest contig length of an assembly which is 50% of the reference genome or longer. N50 did not differ significantly between SPAdes and CLC (Fisher's exact test, P = 0.768),

however, longer N50 was obtained by Unicycler than from Flye (Fisher's exact test,

P = 0.0002). The total length of genomes did not differ significantly between CLC

Genomics Workbench, SPAdes, Flye and Unicycler.

Considering the performance of Unicycler in giving lower number of contigs and

higher NG50 values, it was decided to be the best assembler for downstream

annotation.

Comparison of annotations for hybrid assemblies

Annotations were compared for RAST and Prokka (Table 3). Identical number of

rRNA genes (n = 22) were predicted by both annotation methods in all strains. The

number of tRNA genes differed slightly. One tmRNA was annotated in all strains by

Prokka compared to none by RAST. Different numbers of coding DNA sequences

(CDS) were found for the hybrid genomes when annotated by RAST and Prokka.

Overall, Prokka annotated less (187-327) CDSs compared to RAST (Table 4). RAST

predicted that strains within clone PFGE65-ST95 differed by 88 CDS and strains of

PFGE47-ST131 by 235 CDS. Meanwhile, Prokka predicted 47 different CDS in

PFGE65-ST95 and 200 in PFGE47-ST131 (Fig. 1 A C).

The reasons for the different numbers of CDS annotated by RAST and Prokka were

analysed. Annotation comparison was performed based on presence and absence

matrix generated from both RAST and Prokka. It was then surprising to find

divergent annotations between strains for exactly the same DNA sequences. For

RAST annotations of strains of the PFGE65-ST95 clone, 71 out of the 88 genes

annotated differently between the three strains got 100% identity in complementary

genomes, when controlled on sequence level showing that they were error prone

annotations. Another 14 genes got more than 97% identity for at least one strain by

BLASTn leaving three real gene differences in 729_270514_x_12 since the strain did

not harbor the 2312 bp plasmid with three CDSs (Fig. 1 A-B).

Similar problematic annotations were obtained among PFGE65-ST95 genomic

sequences annotated by Prokka, in which 13 of 47 genes annotated differently within

the three strains of PFGE65-ST95 got hits of 100% identity. Another 32 genes got

more than 97% identity and one gene got 94.7% identity. These 46 genes were

regarded as error prone annotations with Prokka (Fig. 1 A-B).

In PFGE47-ST131, with RAST annotation, 120 out of 235 genes annotated

differently got 100% identity in whole genome sequences which were not annotated

for these genes and three gene differences were identified the same way with more

than 98% identity. All 123 annotations were considered error prone. Ninety-eigth real

gene differences were related to contig 3 identified as a plasmid where

723_010814_2_1 and 723_220514_1_6 carried a 6668-bp *ColRNAI* related plasmid

and 729_141114_1_15 carried a 100,533-bp Incl-1 plasmid. Another 14 real

differences originated in phage related insertions into the chromosome of

729_141114_1_15 (Fig. 1 C – D).

For PFGE47-ST131 genomic sequences annotated with Prokka, 25 of 200 different

genes got 100% identity in reciprocal sequences which were not annotated for these

genes and 20 different genes got more than 97% identity. These 45 annotations

were considered error prone. Similar to RAST annotation, 141 gene differences were

caused by different genes on contig 3 (unique Incl-1 plasmid), and 14 annotation

differences related to prophage regions in the chromosome of strain

729_141114_1_15 (Fig. 1 C – D).

Fig. 2 shows examples of inaccurately annotated genes and their flanking regions

visualized by Clinker. The *neuO* gene (polysialic acid O-acetyltransferase) was only

annotated in strain 729_270514_x_12 and not in the other two strains although they

had identical DNA sequences in this region (A). In 729_141114_1_15, the duplicate

annotation for the *pdeL* gene as *pdeL* and *pdeC* in strain 729_141114_1_15

compared to the other two strains was found to be an error since all three strains

shared 100% sequence identity (B). The last example is with two copies of the

transposase IS*Cep1* gene in two strains which were only annotated as one in

729_141114_1_15. Also here regions compared on the figure shared 100%

sequence identity.

In all, 2.1 and 0.9% of CDSs annotated with RAST and PROKKA, respectively were

error prone. The errors were most often associated to CDSs of shorter length (< 150

nt) with functions annotated as transposases, mobile genetic elements or

hypothetical. These estimates are only based on reciprocal differences between the

annotations and may be underestimated and not included CDSs wrongly annotated

in all strains compared.

The divergence associated with annotations of identical DNA sequence regions falls

back on the annotating methods. The methods and procedures in RAST and Prokka

are quite different. RAST predicts ORFs based on GLIMMER 3.0 where ORFs are

predicted by IMM search in reverse from the stop codon and back towards the start

codon followed by optimization for ribosomal binding sites improving the prediction of

translation initiation sites. Optimizations from previous versions of the program

included in particular the prediction of translation initiation sites and one reason for

inaccurate ORF prediction can be that further optimization of this step is needed.

Prokka uses Prodigal (Hyatt et al. 2010) to identify ORFs based on a dynamic

programming approach optimizing start and start and stop codon identification by

prediction of ribosome binding sites, GC contents, hexamer distributions as well as

iterations of the process. A lower limit of 90 bp is used in the program to score start-

stop codon pairs (Hyatt et al. 2010) and this may be some of the reason for the

errors observed for shorter genes in the current study. Another reason for errors with

Prodigal may be the use of a window of 5 kb to identify start and stop codons. In the

current comparison it can be anticipated that sequence divergence between two

strains in such a window may lead to different annotations at local level even though

sequenced are identical for this region. Hyatt et al. (2010) found that Glimmer 3

predicted around 4% more ORFs in E. coli K12 compared to Prodigal which is in line

with the current study. A problem when comparing annotations is the lack of a

golden standard. Such a standard could have been obtained from *E. coli* strain K12

variant MG1655 where annotation has been refined for decades for instance by

EcoGene (Rudd et al. 2000, Zhou et al. 2013). Unfortunately this resource seems no

longer maintained and accessible. Considering the high genomic variation between

reference and field strain of *E. coli* a full comparison would not have been possible.

Further optimization of the search parameters in the annotation programs may be

needed and targeted better towards *E. coli* and further curation of annotations
needed.
269
Conclusion
There was no significant difference between SPAdes and CLC Genomic Workbench
for benchmark parameters to assemble short reads. However, for hybrid assemblies,
Unicycler provided lower number of contigs and higher N50 than Flye. In the
comparisons, at least 2.1 and 0.9% of genes annotated with RAST and PROKKA,
respectively were false annotations stressing the importance of controlling the
annotations.
277
278 **Conflicts of interest**
279 The authors declare that there are no conflicts of interest.
280

281 **Funding information**
282 This study was partially funded by a State Scholarship Fund to Yufei Zhao from
283 China Scholarship Council (CSC No. 202206350041).
284
Acknowledgements
Xiao Fei and Yaovi Mahuton Gildas Hounmanou are thanked for help with the
analysis of Nanopore sequencing data.

References
Antipov D, Korobeynikov A, McLean JS, Pevzner PA. hybridSPAdes: an algorithm
for hybrid assembly of short and long reads. *Bioinformatics* 2016; 32: 1009-1015.
292
Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes
S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA,
McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R,
Vassieva O, Vonstein V, Wilke A, Zagnitko O. The RAST Server: rapid annotations
using subsystems technology. *BMC Genomics* 2008; 9: 75. doi: 10.1186/1471-2164-
9-75.

Chen S. Ultrafast one-pass FASTQ data preprocessing, quality control, and

deduplication using fastp. *iMeta* 2023; 2: e107.

Christensen H, Bachmeier J, Bisgaard M. New strategies to prevent and control

avian pathogenic *Escherichia coli* (APEC). *Avian Pathol* 2021; 50:370-381.

Chun J, Oren A, Ventosa A, Christensen H, Arahal DR, da Costa MS, Rooney AP, Yi

H, Xu XW, De Meyer S, Trujillo ME. Proposed minimal standards for the use of

genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 2018; 68:

461-466. doi: 10.1099/ijsem.0.002516.

Craddock HA, Motro Y, Zilberman B, Khalfin B, Bardenstein S, Moran-Gilad J. Long-

Read sequencing and hybrid assembly for genomic analysis of clinical *Brucella*

melitensis isolates. *Microorganisms* 2022; 10:619.

De Maio N, Shaw LP, Hubbard A, George S, Sanderson ND, Swann J et al.

Comparison of long-read sequencing technologies in the hybrid assembly of

complex bacterial genomes. *Microb Genom* 2019; 5: e000294.

Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes and

endosymbiont DNA with Glimmer. *Bioinformatics* 2007; 23: 673-9. doi:

10.1093/bioinformatics/btm009.

European Food Safety Authority (EFSA). EFSA statement on the requirements for

whole genome sequence analysis of microorganisms intentionally used in the food

chain. *EFSA J* 2024; 22 :e8912. doi: 10.2903/j.efsa.2024.8912.

Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* 2013; 29: 1072-1075.

Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal:

prokaryotic gene recognition and translation initiation site identification. *BMC*

Bioinformatics 2010;11:119. doi: 10.1186/1471-2105-11-119.

Jørgensen SL, Stegger M, Kudirkiene E, Lilje B, Poulsen LL, Ronco T et al. Diversity

and population overlap between avian and human *Escherichia coli* belonging to

sequence type 95. *mSphere* 2019; 4: e00333-18.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads

using repeat graphs. *Nat Biotechnol* 2019; 37: 540-546.

Köstlin-Gille N, Härtel C, Haug C, Göpel W, Zemlin M, Müller A et al. Epidemiology

of early and late onset neonatal sepsis in very low birthweight infants: data from the

German Neonatal Network. *Pediatr Infect Dis J* 2021; 40: 255-259.

Khezri A, Avershina E, Ahmad R. Hybrid assembly provides improved resolution of

plasmids, antimicrobial resistance genes, and virulence factors in *Escherichia coli*

and *Klebsiella pneumonia* clinical isolates. *Microorganisms* 2021; 9:2560.

Maguire M, Kase JA, Roberson D, Muruvanda T, Brown EW, Allard M et al.

Precision long-read metagenomics sequencing for food safety by detection and

assembly of Shiga toxin-producing *Escherichia coli* in irrigation water. *PloS One*

2021; 16: e0245172.

Manges AR, Geum HM, Guo A, Edens TJ, Fibke CD, Pitout JDD. Global

extraintestinal pathogenic *Escherichia coli* (ExPEC) lineages. *Clin Microbiol Rev*

2019; 32: e00135-18.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO update: novel

and streamlined workflows along with broader and deeper phylogenetic coverage for

scoring of eukaryotic, prokaryotic, and viral genomes. *Mol Biol Evol* 2021; 38: 4647-

4654.

Nolan LK, Barnes HJ, Vaillancourt JP, Abdul-Aziz T, Logue CM. 2020. Colibacillosis,

Diseases of Poultry, 14th ed, pp. 770-830. Wiley.

Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T et al. The SEED and

the rapid annotation of microbial genomes using subsystems technology (RAST).

Nucleic Acids Res 2014; 42(D1): D206-D214.

Oxford Nanopore Technologies. 2022. Medaka. Available online:

https://github.com/nanoporetech/medaka (accessed on 17 October 2023).

Poulsen LL, Thøfner I, Bisgaard M, Christensen JP, Olsen RH, Christensen H.

Longitudinal study of transmission of *Escherichia coli* from broiler breeders to

broilers. *Vet Microbiol* 2017; 207:13-18.

Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes de

novo assembler. *Curr Protoc Bioinformatics* 2020; 70: e102.

Riesco R, Trujillo ME. Update on the proposed minimal standards for the use of

genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 2024; 74:

006300. doi: 10.1099/ijsem.0.006300.

Sanderson H, McCarthy MC, Nnajide CR, Sparrow J, Rubin JE, Dillon JR, White AP.

Identification of plasmids in avian-associated Escherichia coli using nanopore and

illumina sequencing. *BMC Genomics* 2023; 24: 698. doi: 10.1186/s12864-023-

09784-6.

Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014; 30:

2068-2069.

Shen W, Le S, Li Y, Hu F. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q
file manipulation. *PloS One* 2016; 11: e0163962.

Therrien DA, Konganti K, Gill JJ, Davis BW, Hillhouse AE, Michalik J et al. Complete
whole genome sequences of *Escherichia coli* surrogate strains and comparison of
sequence methods with application to the food industry. *Microorganisms* 2021; 9:
608.

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S et al. Pilon: an
integrated tool for comprehensive microbial variant detection and genome assembly
improvement. *PloS One* 2014; 9: e112963.

Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome
assemblies from short and long sequencing reads. *PLoS Comput Biol* 2017; 13:
e1005595.

Zhou J, Rudd KE. EcoGene 3.0. *Nucleic Acids Res* 2013; 41(Database issue):
D613-24. doi: 10.1093/nar/gks1235.

Figure legends

Fig. 1. Venn diagrams illustrating the deviating annotations between Prokka and

RAST. For PFGE65-ST95, (A) shows the number of different genes predicted before

BLASTn control and (B) shows the numbers after control. For PFGE47-ST131, (C)

shows the number of different genes predicted before BLASTn control and (D)

shows the numbers after control.

Fig. 2. Examples of genetic comparisons of inaccurately annotated genes from

Prokka and their flanking regions. An additional annotation is given for the *neuO*

gene in 729_270514_x_12 (A). In 729_141114_1_15, duplicate annotation for the

pdeL gene as *pdeL* and *pdeC* was found (B). Two copies of the transposase IS*Cep1*

gene in two strains were only annotated as one in strain 729_141114_1_15 (C). The

regions compared all shared 100% DNA sequence identity without gaps.

Table 1. Strains investigated, genome length, genetic elements from six hybrid assembled genomes.

| Strain | Origin | Organ/lesion | PFGE-ST | Hybrid assembly | | | INDSC acc. no. | | |
				Contig number	Contig length (bp)	Hybrid assembly	Short read assembly	Matching genomic elements
729_021014_1_4	Broiler parent	Salpingitis	65-95	1	5,232,657	CP160125	JBELNO01	Chromosome
				2	193,604	CP160126		Plasmid *
				3	6655	CP160127		Plasmid
				4	2676	CP160128		Plasmid
				5	2312	CP160129		Plasmid
729_270514_x_12	Broiler parent	Salpingitis	65-95	1	5,232,714	CP160121	JBELNS01	Chromosome
				2	193,604	CP160122		Plasmid *
				3	6655	CP160123		Plasmid
				4	2677	CP160124		Plasmid
729_xx_1_3_yolk	Broiler	Yolk sac infection	65-95	1	5,232,647	CP160116	JBELNM01	Chromosome
				2	193,604	CP160117		Plasmid *
				3	6655	CP160118		Plasmid
				4	2676	CP160119		Plasmid
				5	2312	CP160120		Plasmid
723_010814_2_1	Broiler parent	Salpingitis	47-131	1	4,866,047	CP160112	JBELNC01 (SSQU01)	Chromosome
				2	201,035	CP160113		Plasmid *
				3	6668	CP160114		Plasmid
				4	1552	CP160115		Plasmid
729_141114_1_15	Broiler parent	Salpingitis	47-131	1	4,887,791	CP160108	JBELNB01	Chromosome
				2	201,054	CP160109		Plasmid *
				3	100,533	CP160110		Plasmid
				4	1552	CP160111		Plasmid
723_220514_1_6	Broiler	Liver/septicaemia	47-131	1	4,866,598	CP160104	JBELNH01	Chromosome
				2	200,992	CP160105		Plasmid *
				3	6668	CP160106		Plasmid
				4	1552	CP160107		Plasmid

*ColV-like

Table 2. Statistical analysis of reads.

ST	Sample	Illumina				ONT			
		Mean read length (bp)	Number of reads	Total bases (Mbp)	Coverage (x)*	Mean read length (bp)	Number of reads	Total bases (Mbp)	Coverage (x)
95	729_021014__1_4	184.9	515,309	154	41.1	5129	137,504	664	151.9
	729_270514_x_12	190.5	613,533	186	50.3	4138	84,953	336	75.7
	729_xx_1_3_yolk	162.8	594,174	150	41.7	4260	95,064	387	87.2
131	723_010814_2_1	262.4	800,953	283	90.5	2710	109,803	283	64.1
	729_141114_1_15	176.2	593,386	167	45.0	3832	138,333	508	114.2
	723_220514_1_6	207.2	618,972	186	55.2	3386	93,939	301	68.5
	Average	197.3	622,721	188	54.0	3909	109,933	413	93.6

*coverage = (number of read × read length) / total genome size. In which total genome size was estimated to 5.2 Mbase.

Table 3. The quality of assemblies generated by CLC Genomic Workbench and SPAdes for short-read only assemblies and Flye and Unicycler for hybrid assemblies.

ST	Strain	Number of contigs				NG50 (bp)				Total length (bp)				Completeness (%)			
		CLC	SPAdes	Flye	Unicycler	CLC	SPAdes	Flye	Unicycler	CLC	SPAdes	Flye	Unicycler	CLC	SPAdes	Flye	Unicycler
95	729_021014__1_4	110	99	5	5	169,077	196,246	5,232,681	5,232,657	5,310,368	5,312,171	5,444,670	5,437,904	100	100	100	99.8
	729_270514_x_12	100	100	5	4	211,910	204,287	5,232,730	5,232,714	5,313,797	5,314,927	5,449,369	5,435,650	100	99.8	100	99.8
	729_xx_1_3_yolk	143	132	6	5	135,314	164,290	5,232,625	5,232,647	5,307,620	5,316,910	5,446,817	5,437,894	100	100	100	99.8
Average		117.7	110.3	5.3	4.7	172,100.3	188,274.3	5,232,678,7	5,232,672.7	5,310,595.0	5,314,669.3	5,446,952,0	5,437,149.3	100.0	99.9	100.0	99.8
131	723_010814_2_1	46	45	16	4	354,765	353,089	1,074,415	4,866,047	5,033,610	5,022,129	5,076,852	5,075,302	100	100	100	99.8
	729_141114_1_15	88	88	5	4	192,789	224,060	4,949,514	4,887,791	5,110,669	5,115,410	5,164,053	5,190,930	100	99.8	100	99.8
	723_220514_1_6	51	50	8	4	354,390	353,089	2,855,420	4,866,598	5,021,087	5,022,103	5,083,114	5,075,810	100	100	100	99.8
Average		61.7	61.0	9.7	4.0	300,648.0	310,079.3	2,959,783.0	4,873,478.7	5,055,122.0	5,053,214.0	5,108,006,3	5,114,014.0	100.0	99.9	100.0	99.8

Table 4. Statistical analysis of annotations.

ST	Strain	CDS		tRNA		tmRNA		Total *RNA		Number of CDS differences	
		RAST	Prokka	RAST	Prokka	RAST	Prokka	RAST	Prokka	RAST	Prokka
95	729_021014_1_4	5476	5149	90	92	0	1	112	115	88	47
	729_270514_x_12	5450	5263	90	92	0	1	112	115		
	729_xx_1_3_yolk	5442	5148	90	92	0	1	112	115		
131	723_220514_1_6	4967	4699	84	86	0	1	106	109	235	200
	723_010814_2_1	4958	4698	84	86	0	1	106	109		
	729_141114_1_15	5094	4830	85	86	0	1	107	109		

*22 rRNA genes in all strains for both annotation methods.

Fig. 1A

Fig. 1B

Fig. 1C

Fig. 1D

Fig. 2A

Fig. 2B

Fig. 2C