A reference genome for *Trichogramma kaykai*: A tiny desert-dwelling parasitoid wasp

with competing sex-ratio distorters

Jack Culotta & Amelia RI Lindsey*

Department of Entomology, University of Minnesota, Saint Paul, MN, USA 51108

*corresponding: alindsey@umn.edu

JC ORCID: 0000-0002-2533-0802

ARIL ORCID: 0000-0003-3354-7419

ABSTRACT

The tiny parasitoid wasp *Trichogramma kaykai* inhabits the Mojave Desert of the southwest

United States. Populations of this tiny insect variably host up to two different sex-distorting

genetic elements: (1) the endosymbiotic bacterium *Wolbachia* which induces the

parthenogenetic reproduction of females, and (2) a B-chromosome, "Paternal Sex Ratio" (PSR),

which converts would-be female offspring to PSR-transmitting males. We report here the

genome of a *Wolbachia*-infected *Trichogramma kaykai* isofemale colony KSX58. Using Oxford

Nanopore sequencing we produced a final genome assembly of 203 Mbp with 45x coverage,

consisting of 213 contigs with an N50 of 1.9 Mbp. The assembly is quite complete, with 91.41%

complete BUSCOs recovered: a very high score for Trichogrammatids that have been

previously characterized for having high levels of core gene losses. We also report a complete

mitochondrial genome for *T. kaykai,* and an assembly of the associated *Wolbachia*, strain *w*Tkk.

We identified copies of the parthenogenesis-inducing genes *pifA* and *pifB* in a remnant

prophage region of the *w*Tkk genome. The *Trichogramma kaykai* assembly is the highest quality

genome assembly for the genus to-date and will serve as a great resource for understanding

the evolution of sex and selfish genetic elements.

INTRODUCTION

Trichogramma wasps (Hymenoptera: Trichogrammatidae) are some of the smallest animals on the planet (Polilov 2015). The genus contains more than 200 described species: all parasitoids that complete their development within the eggs of other insects (Burks et al. 2024; Pinto 2006). Trichogrammatid research has largely focused on (1) their application as biological control agents of insect pests (Knutson 1998; Cherif et al. 2021), (2) innovations associated with extreme miniaturization (Polilov 2012), and (3) sex allocation, especially due to relationships with sex-distorting elements (Stouthamer et al. 1990; Stouthamer and Kazmer 1994; Russell and Stouthamer 2010). The most common sex-ratio distorter is the intracellular, maternally transmitted bacterium *Wolbachia,* a common associate of many arthropods and nematodes (Kaur et al. 2021). In *Trichogramma*, most *Wolbachia* strains are "parthenogenesis-inducing" (PI), and enable the asexual reproduction of females (i.e., "thelytokous parthenogenesis") (Stouthamer et al. 1990; Stouthamer et al. 1993; Ma and Schwander 2017).

To date all instances of microbe-mediated PI are in animals with haplodiploid sex determination (Ma and Schwander 2017; Verhulst et al. 2023). Under haplodiploidy (and without PI-*Wolbachia*) males typically develop from unfertilized (i.e., haploid) eggs, and females are typically derived from fertilized, diploid, eggs (De La Filia et al. 2015). PI-*Wolbachia* diplodize the unfertilized eggs, resulting in a female (Stouthamer and Kazmer 1994). In one species with PI-*Wolbachia*, *Trichogramma kaykai* (Figure 1A-B), a second sex-distorter is sometimes present: a supernumerary B-chromosome, "Paternal Sex Ratio" (PSR) (van Vugt et al. 2003; Stouthamer et al. 2001). PSR achieves the opposite outcome of *Wolbachia*'s PI: haploid males with PSR mate, and any fertilized eggs develop into more PSR-transmitting males (Van Vugt et al. 2009). PSR facilitates destruction of the paternal genome (except for itself), resulting in a haploid embryo (the maternal copy) and the untouched PSR chromosome. In populations where *Wolbachia* and PSR are present, a curious pattern of reproduction is present: males are derived

from fertilized eggs (with PSR-containing sperm), and females are derived from unfertilized

eggs (with PI-*Wolbachia*) (Figure 1C). Unlike many other PI-*Wolbachia* systems where PI is

accompanied by a decay of sexual function (Stouthamer et al. 2010; Russell and Stouthamer

2011; Jeong and Stouthamer 2005; Stouthamer and Mak 2002; Gottlieb and Zchori-Fein 2001),

Trichogramma kaykai are easily cured of their *Wolbachia* in the lab, and readily return to a fully

functional sexual form (Hohmann and Luck 2000; Hohmann et al. 2001; Miura and Tagami

2004; Russell et al. 2016). The PSR chromosome ensures males and sexual reproduction are

maintained.

As host to PI-*Wolbachia* and PSR, *Trichogramma kaykai* is a valuable model for understanding

the evolution of sex ratios and interactions between selfish genetic elements. This species was

described in 1997 (Pinto et al.) and is native to the deserts of the Southwest United States

(Figure 1D). We report a reference genome for an isofemale colony of *Trichogramma kaykai*

from the Mojave Desert, plus the genome of its PI-*Wolbachia* strain, *w*Tkk. To our knowledge,

there are currently no *Trichogramma kaykai* PSR chromosomes in culture, but this reference

genome will aid in future efforts to understand how this selfish element alters chromosome

dynamics and sex ratios.

Figure 1. *Trichogramma kaykai* biology. (A) Three *Trichogramma kaykai* females ovipositing into host

moth eggs (*Ephestia kuehniella*). **(B)** An exemplary specimen of *T. kaykai* (female). **(C)** Sex in *T. kaykai* is

determined based on haplodiploidy, mediated by the presence or absence of *Wolbachia* (maternally

transmitted) and the PSR chromosome (paternally transmitted). **(D)** The sample collection site for KSX58

and predicted geographic range of *Trichogramma kaykai*.

MATERIALS & METHODS

Species Origin and Sampling Strategy

Genome sequencing and assembly was performed for *Trichogramma kaykai* line "KSX58", an

isofemale laboratory culture. A single unmated *Wolbachia*-infected, thelytokous female was

reared out of a parasitized *Apodemia mormo* egg collected off an *Eriogonum inflatum* stem and

used to initiate an isofemale line. The founding female was collected in May 2010 in Kelso, CA,

USA, by R. Stouthamer and J. Russell (Figure 1D). The colony has since been maintained in 5

ml glass culture tubes stopped with cotton, and kept at 25°C with a 12:12 light:dark cycle.

Wasps are hosted every 12 days on sterilized *Ephestia kuehniella* eggs adhered to cardstock

alongside a streak of honey. *Wolbachia* infection status was confirmed by PCR with *Wolbachia*

specific "Wspec" primers (Werren and Windsor 2000), and *Trichogramma* species was

confirmed by molecular identification (Stouthamer et al. 1999), both as detailed previously

(Lindsey and Stouthamer 2017). To collect wasps for DNA extraction, freshly emerged females

were allowed to crawl up into a sterile tube attached to the colony culture vial. The pool of

wasps was flash frozen in liquid nitrogen and stored at -80°C for further processing.

Geographic Range Map

Locations of *Trichogramma kaykai* are centered around the Southern Mojave Desert (Pinto et

al. 1997; Russell et al. 2018; Tulgetske and Stouthamer 2012; Russell et al. 2016; Van Vugt et

al. 2009; van Vugt et al. 2003). The predicted northern and southern boundaries of this species'

range were estimated from these observations. As it is assumed *Trichogramma kaykai* is

restricted to desert habitat, the eastern and western borders of range are indicated by the

Southern Mojave Desert and Northern Sonoran Desert. The map was generated in ArcGIS

Online (www.arcgis.com).

Sequencing Methods and Sample Preparation

DNA was extracted from 25 mg of whole insect tissues using the MagAttract High Molecular

Weight kit (Qiagen), following manufacturer's instructions. The DNA was concentrated to 25 uL

using Sergi Lab Supplies magnetic beads and went through the PacBio SRE kit to deplete

fragments shorter than 10kb. The sample was barcoded and library prepped with the ONT SQK-

NBD114.24 kit. The libraries were sequenced on a P2 Solo instrument using PromethION

10.4.1 flow cells. Every 24 hours the libraries were recovered and flowcells were flushed with

nuclease (EXP-WSH004 kit) and reloaded.

Nuclear Genome Assembly, Curation, and Quality Control

Samples were originally basecalled within Minknow using 'super accuracy' mode with

5mC_5hmC modified base calling. Reads were then re-basecalled with dorado v.0.7.2 using

basecall model dna_r10.4.1_e8.2_400bps_sup\@v5.0.0. Reads at least 5kb in length were

maintained, processed with 'dorado correct', and used for generating an assembly with Hifiasm

v.0.19.9 and default parameters. The genome was manually curated, and cytoplasmic genomes

were identified through tblastn results implemented in Blobtools v.1.1.1 (Challis et al. 2020).

Assemblies were assessed with Compleasm v.0.2.6 (Huang and Li 2023) with the hymenoptera

lineage flag ('-l hymenoptera').

Genomic Methylation

Methylation and hydroxymethylation of genomic DNA at 5' cytosines (5mC and 5hmC) in a

cytosine-guanine dinucleotide (CpG) context was determined from the basecalling information

stored in the unmapped modBAM files (Flack et al. 2024). These were aligned to the final

assembly using Minimap v.2.17 (Li 2016), converted to bedMethyl format with Modkit v.0.4.1

(https://github.com/nanoporetech/modkit), and the 5mC and 5hmC percentages were calculated

with an AWK script.

***Trichogramma* Phylogeny**

A whole-genome phylogeny was reconstructed with SANS v.2.4_10, which uses a pangenomic

approach to calculate splits in a phylogenetic tree (Rempel and Wittler 2021). SANS parameters

included '--filter strict' with an output Newick tree file and 100 bootstrap replicates. Taxa

included the available *Trichogramma* genomes (for *Trichogramma brassicae*, which is

represented by two assemblies, only GCA_902806795.1 was used; Table 1), and an outgroup

species from a closely related family (Cruaud et al. 2024), *Phymastichus coffea* (Hymenoptera:

Eulophidae) GCF_024137745.1. Tree topology was configured in FigTree v.1.4.4

(https://github.com/rambaut/figtree/) and annotated in Inkscape (https://www.inkscape.org).

Table 1. Available *Trichogramma* genome assemblies.

Species	Accession	Size (bp)	Contig/ Scaffold Count*	N50 (bp)*	*Wolbachia* Genome	Citation
Trichogramma brassicae	GCA_902806795.1	235,386,796	1,570 (C)	556,663	No	Ferguson et al. (2020)
Trichogramma brassicae	GCA_030522885.1	203,810,232	87,792 (S)	18,131	No	Guinet et al. (2023)
Trichogramma dendrolimi	GCA_034770305.1	215,209,100	316 (S)	1,412,680	No	Zhang et al. (2023)
Trichogramma evanescens	GCA_902732785.1	213,671,129	146,286 (S)	38,173	No	N/A
Trichogramma pretiosum	GCA_000599845.3	187,641,947	925 (S)	1,825,723	*w*Tpre (Lindsey et al. 2016)	Lindsey et al. (2018a)
Trichogramma kaykai	Processing	203,423,343	213 (C)	1,898,390	*w*Tkk	This study

*If assembly is scaffolded, metrics reported are for scaffolds and an (S) is indicated in the count column. If

there are only contigs, those metrics are reported and (C) is indicated in the count column.

Repeat Assembly Techniques

We identified and masked repetitive sequences in each genome. First a custom *de novo* repeat

library was crated with RepeatModeler v.2.0.5 (Flynn et al. 2020) with the -LTRStruct parameter

included. Then this library was used to mask the genome with RepeatMasker v.4.1.1 (Tarailo-

Graovac and Chen 2009) with the -s (sensitive mode) parameter included.

Gene Finding Methods

To annotate the *T. kaykai* genome, a soft masked genome was used for gene model prediction with Galba v.1.0.11 (Brůna et al. 2023), using the RefSeq annotations for *Trichogramma brassicae* (GCA_902806795.1), *Trichogramma pretiosum* (GCA_000599845.3), *Nasonia vitripennis* (GCA_009193385.2), *Copidosoma floridanum* (GCF_000648655.2), *Phymastichus coffea* (GCF_024137745.1) and *Ceratosolen solmsi marchali* (GCF_000503995.2) as references. The Galba pipeline was executed using Singularity with parameters to output a gff3 file. Summary statistics for the resulting gff3 file were computed with GAG v.2.0.1 (Geib et al. 2018). Split genes (those encoded across the ends of two contigs) were manually re-assigned gene identifiers as per NCBI best practices.

Synteny Analysis

We identified conserved regions and mapped synteny between the *T. kaykai* and *T. pretiosum* genomes (Table 1) using D-GENIES webtool (https://dgenies.toulouse.inra.fr/run) (Cabanettes and Klopp 2018) employing Minimap v.2.28 (Li 2016), the "many repeats" flag, and the "hide noise" option.

Mitogenome

A single circular contig was identified as the mitochondrial genome based on GC content, size, and coverage. Mitogenome annotation was completed with MITOS2 v.2.1.9 (Bernt et al. 2013; Donath et al. 2019) and the circular mitogenome was started at Cox1 per convention with rearrangement in SnapGene v.7.2. MITOS2 parameters were the RefSeq63 Metazoa reference and the invertebrate mitochondrial translation code. Manual curation of the control region and inferences of gene structure were made based on comparisons to other *Trichogramma* mitochondrial genomes (Chen et al. 2018).

Wolbachia **Strain *w*Tkk Genome**

Four contigs were identified as a *Wolbachia* genome based on cumulative size and Blobtools

results. Genome completeness was analyzed against the rickettsiales_odb10 database with

Compleasm v.0.2.6 (Huang and Li 2023). Prophage regions and mobile elements were

identified with VirSorter2 v.2.2.4 (Guo et al. 2021) and mobileOG-db v.1.0.1 (Brown et al. 2022),

implemented in proksee (Grant et al. 2023)(https://proksee.ca/) with default parameters. To

identify putative parthenogenesis-inducing genes (*pifs*) (Fricke and Lindsey 2024), we leveraged

annotation and orthology data generated by Prokka v.1.14.6 (Seemann 2014) and OrthoFinder

v.2.5.4 (Emms and Kelly 2019), implemented in the *Wolbachia* Phylogeny Pipeline (WHOP;

https://github.com/gerthmicha/WHOP). Phylogenetic analysis was performed based on the

clustering results from WHOP/OrthoFinder results. Single-copy orthologs were aligned with

MAFFT L-INS-i v.7.487 (Katoh and Standley 2013), recombining genes were eliminated with

PhiPack v.1.1 (Bruen and Bruen 2005), and alignments were concatenated for phylogenetic

reconstruction in IQtree v.2.2.3 (Nguyen et al. 2015), run with model optimization and 1000

ultrafast bootstrap replicates.

RESULTS & DISCUSSION

Sequencing and Assembly

We generated 10.5 billion base pairs of nanopore sequencing data: a total of 1,543,039 reads

with a read N50 of 13,472 (Supplementary Table S1). A draft assembly from reads longer than

5,000 base pairs was generated with HiFiasm which produced a 204.6 Mbp assembly contained

in 226 contigs. A combination of coverage, GC%, and blast hits from BlobTools results were

used to identify non-nuclear contigs and curate the assembly. After removing spurious and

contaminant contigs (Supplemental Table S2), and extracting the *Wolbachia w*Tkk and

mitochondrial genomes, the final assembly was 203.4 Mbp in 213 contigs, with an average of

45x coverage (Table 2). The *Trichogramma kaykai* assembly falls in the middle of the size

range for the genus, (187.6 Mbp in *T. pretiosum* to 235.4 Mbp in one of the *T. brassicae* (Table

1). Additionally, this size closely aligns with a flow cytometry-based estimate of 216 Mbp for a

different colony of *T. kaykai*, "LC19-1" (van Vugt et al. 2005). The GC% of *Trichogramma*

genomes appears to be highly conserved, with all at 40%.

Quality assessments indicate that the genome assembly is quite complete, with 91.41% of

Hymenopteran BUSCO loci present as complete coding sequences (Table 2). These metrics

are on par with other well-assembled *Trichogramma* genomes (e.g., *T. pretiosum,* Hymenoptera

BUSCO: C:92.7%[S:90.9%,D:1.8%],F:0.8%,M:6.5%,n=5991)(Lindsey et al. 2018a). Comparative

genomics of *T. pretiosum* relative to other hymenopterans indicated that these wasps have

undergone a large number of core gene losses and have highly accelerated rates of protein

evolution (Lindsey et al. 2018a) so we do not expect BUSCO scores close to 100% even for a

"perfect" assembly.

Table 2. *Trichogramma kaykai* genome assembly statistics.

Metric	Draft	Final
Contigs	226	213
Total Length (bp)	204,642,443	203,423,343
Min contig length	5,287	6,150
Average contig length	905,498	955,039
Max contig length	6,625,044	6,625,044
N50	1,898,390	1,898,390
L50	34	34
%GC	39.60	39.63
Compleasm*	91.44% [S:90.32%, D: 1.12%] F: 0.78%, M:7.78%, n=5991	91.41% [S:90.29%, D: 1.12%] F: 0.82%, M:7.78%, n=5991

*Standard BUSCO annotation: Complete BUSCOs (C) [Complete and single-copy BUSCOs (S),

Complete and duplicated BUSCOs (D)], Fragmented BUSCOs (F), Missing BUSCOs (M), Total BUSCO

groups searched (n). Hymenoptera dataset used for determining completeness.

Genome Methylation

We determined 5' methylation at cytosines in a CpG context based on the direct sequencing

basecalls. Less than 1% of CpGs were methylated: 0.67% of CpGs had 5mC (methyl)

modifications and 0.18% had 5hmC (hydroxymethyl) modifications. While this is a low level of

methylation as compared to vertebrates, this is not atypical for insects (Hunt et al. 2013).

Importantly, this level of methylation closely mirrors the number of methylated CpG sites

identified in *T. pretiosum* using bisulfite sequencing (Lindsey et al. 2018a; Wu et al. 2020).

Analysis of Repetitive DNA

Trichogramma kaykai is sister to all other *Trichogramma* species with published genomes

(Figure 2A). Across the genus, repetitive content appears to be relatively conserved. Repetitive

sequences account for between 17.9% - 29.39% of the total genome lengths (Table 1, Figure

2B, Supplemental Table S3). This is in contrast to the outgroup species, *Phymastichus coffea*

(Hymenoptera: Eulophidae), that has a 421 Mbp genome with more than half (57.21%)

attributed to repetitive sequences (Figure 2B, Supplemental Table S3). Across *Trichogramma*,

the majority of repetitive sequences are unclassified. In *T. kaykai*, 4% of the genome is derived

from retroelements, <1% from DNA transposons, and around 3% of the genome is simple and

low complexity repeats (Table 2). We then assessed the level of synteny between *T. kaykai* and

T. pretiosum by cross-mapping similar genomic sequences with D-GENIES (Figure 3). A large

proportion (60.34%) of the *T. kayaki* genome shares 50-75% identity with *T. pretiosum*, and

there are high levels of synteny across the two assemblies (Figure 3).

A

B

millions of base pairs

T. kaykai

T. pretiosum

T. dendrolimi

T. evanescens

T. brassicae

Phymastichus coffea

8000000.0

unmasked   unclassified   retroelements   DNA transposons   other

Figure 2. Comparative genomics of *Trichogramma*. (A) Whole genome phylogeny of five

Trichogramma species and outgroup *Phymastichus coffea* (Hymenoptera: Eulophidae). Double slashes

indicate branches that were shortened to half their length for ease of visualization. **(B)** Repetitive content

of each genome, corresponding to the taxa in (A). "Other" includes rolling circles, simple repeats, and low

complexity repeats.

Table 2. Interspersed repeats in *Trichogramma kaykai*.

Name	Number	Length (bp)	Percent (%)
Retroelements	8,591	8,132,788	4.00%
Penelope class	203	76,919	0.04 %
LINE class	5,232	5,084,665	2.50 %
L2/CR1/Rex	674	423,279	0.21 %
R1/LOA/Jockey	1,929	1,977,765	0.97 %
R2/R4/NeSL	2,056	2,301,966	1.13 %
LTR class	3,359	3,048,123	1.50 %
BEL/Pao	484	436,375	0.21 %
Ty1/Copia	497	368,036	0.18 %
Gypsy/DIRS1	2,378	2,243,712	1.10 %
DNA transposons	2,812	1,863,150	0.92 %
hobo-Activator	497	176,867	0.09 %
Tc1-IS630-Pogo	370	118,645	0.06 %
Rolling-circles	797	350,441	0.17 %
Unclassified	110,502	29,820,645	14.66 %
Total interspersed repeats		39,816,583	19.57 %
Simple repeats	147,687	5,403,933	2.66 %
Low complexity	15,687	704,136	0.35 %
Bases masked		46,275,093	22.75 %

Figure 3. Synteny is highly conserved between *T. kaykai* and *T. pretiosum*. Dot plot indicating

syntenic regions between *Trichogramma* genomes. Dots are colored according to percent identity.

Genome Annotation

We annotated the *T. kaykai* genome using a set of protein sequences from other wasps in the

superfamily Chalcidoidea as a reference and identified 20,798 genes (Table 3). These genes

corresponded to 24,714 transcripts, with a mean of four exons per mRNA (Table 3). Compared

to other *Trichogramma* species, this is a larger number of annotated genes (e.g., 13,395 in *T.*

pretiosum, 16,905 in *T. brassicae*). However, this could be due to differences in annotation

pipelines, and or, lineage-specific differences in the patterns of gene gain and loss.

270   **Table 3. Annotation metrics for the *Trichogramma kaykai* genome.**

Metric	Value*
Number of genes	20,798
Number of mRNAs	24,714
Number of exons	116,714
Number of introns	92,000
Mean exons per mRNA	4
Total gene length	102,108,152 bp
Longest gene	160,105 bp
Mean gene length	4,910 bp
Longest CDS	54,789 bp
Mean CDS length	1,462 bp
Longest exon	14,469 bp
Mean exon length	310 bp

*Base pairs = bp

Mitogenome

We identified the mitogenome based on GC content (14.81%), size (16,399 bp), and coverage

(3708x). Annotation revealed all expected mitochondrial tRNAs and coding genes (Figure 4).

MITOS2 annotated a single large rRNA of only 712 bp and three regions (387, 49, and 38 bp)

as small rRNAs. Comparison to other Trichogrammatid mitochondrial genomes indicated that

the large rRNA annotation had been truncated on the 5' end, and the small rRNA annotation

had been fragmented (Figure 4), which is likely due to the extreme divergence of these

mitochondrial sequences. A 878 bp region between the tRNAs for tryptophan (W) and

methionine (M) corresponds to the putative control region identified in other Trichogrammatid

mitochondrial genomes (Chen et al. 2018).

Figure 4. Mitochondrial genome of *Trichogramma kaykai*. Genes were annotated with MITOS2 (Bernt

et al. 2013). Putative regions of rRNAs that were not correctly annotated by MITOS2 are indicated with

stripes. The control region and the putative full length rRNAs were identified based on homology and

gene order of other *Trichogramma* mitochondria (Chen et al. 2018). Transfer RNAs (tRNAs) are denoted

by IPUC-IUB amino acid codes.

Parthenogenesis-Inducing *Wolbachia* Strain *w*Tkk

We assembled a near-complete *Wolbachia* genome of the *w*Tkk strain: ~1.12Mbp contained in

four contigs, sequenced at 55X coverage (Table 4). Phylogenetic reconstruction revealed that

*w*Tkk is in the "Supergroup B" clade of *Wolbachia*, and is sister to *w*Tpre, which infects

Trichogramma pretiosum (Lindsey et al. 2016)(Figure 5A). The *w*Tkk and *w*Tpre genomes are

similar in size: the *w*Tpre assembly (a single scaffold) is just slightly larger at 1,133,709 bp

(Lindsey et al. 2016). We queried the *w*Tkk proteins to identify the recently identified

parthenogenesis inducing factors, *pifA* and *pifB* (Fricke and Lindsey 2024). We identified a

single copy of each gene in the *w*Tkk genome, encoded next to each other within a remnant

prophage region (Figure 5B), as is typical of many other *Wolbachia* loci that induce host

reproductive manipulations (Fricke and Lindsey 2024; LePage et al. 2017; Lindsey et al. 2018b;

Shropshire et al. 2018; Perlmutter et al. 2019; Bordenstein and Bordenstein 2016). The *w*Tkk

PifA protein was 67% identical to the PifA from *w*Tpre, and 30% identical to the PifA from *w*Lcla

(another PI-*Wolbachia* infecting the parasitoid wasp *Leptopilina clavipes* [Hymenoptera:

Figitidae])(Pannebakker et al. 2004). In contrast, PifB proteins were more conserved: *w*Tkk and

*w*Tpre PifB were 93% identical, and *w*Tkk and *w*Lcla PifB shared 56% amino acid identity.

Figure 5. Parthenogenesis-inducing *Wolbachia* strain *w*Tkk. (A) Maximum likelihood-based

phylogeny of *Wolbachia* strains and Rickettsiales outgroups based on 78 core, single-copy, protein

coding genes (a total of 30,477 aligned amino acid sites). **(B)** Gene models for a predicted remnant

prophage region that contains the parthenogenesis factors *pifA* and *pifB.* Three tandem CDS were

annotated as *mutL*, which is likely a pseudogenization of *mutL* due to nonsense mutations and

fragmentation of the coding region into multiple open reading frames. Abbreviations: insertion element

(IS), parthenogenesis inducing factor (*pif*), coding sequence (CDS).

Table 4. *Wolbachia* strain *w*Tkk genome assembly and annotation.

Metric	*w*Tkk
Contigs	4
Length (bp)	1,119,794
%GC	33%
Compleasm*	93.96% [S:93.96%, D: 0%] F: 0.55%, M:5.49%, n=364
CDS	1,265
rRNAs	3
tRNAs	34

*Standard BUSCO annotation: Complete BUSCOs (C) [Complete and single-copy BUSCOs (S),

Complete and duplicated BUSCOs (D)], Fragmented BUSCOs (F), Missing BUSCOs (M), Total BUSCO

groups searched (n). Rickettsiales dataset used for determining completeness.

SUMMARY

We report here a high-quality assembly for the parasitoid wasp *Trichogramma kaykai* along with

genomes for its mitochondrion and associated *Wolbachia* strain, *w*Tkk. There are five other

Trichogramma genomes currently available on NBCI: one each from the *Trichogramma* species

pretiosum, dendrolimi, and *evanescens*, and two assemblies for *Trichogramma brassicae*.

These species are some of the more commonly available *Trichogramma* sold as biological

control agents of lepidopteran pests (Knutson 1998; Cherif et al. 2021). The *Trichogramma*

kaykai assembly reported here is arguably the highest quality assembly available for the genus,

as it has the lowest number of contigs and the highest N50 (Table 1). To date, all *Trichogramma*

species assayed for karyotype have a haploid genome of five chromosomes (2n=10) (Gokhman

and Quicke 1995; Van Vugt et al. 2009; Gokhman 2020; Gokhman et al. 2017; Farsi et al.

2020). While chromosome number and approximate genome size are conserved, there do

appear to be species-specific differences in chromosome morphometrics (e.g., centromere

location, arm lengths, chromosome sizes) (Farsi et al. 2020; Gokhman et al. 2017; Gokhman

2020).

Of the *Trichogramma* genome sequencing efforts, one other reports a *Wolbachia* genome:

strain *w*Tpre, from *T. pretiosum* (Lindsey et al. 2016). The two *Trichogramma*-infecting strains,

*w*Tpre and *w*Tkk, are closely related members of the "Supergroup B" clade which contains a

suite of other arthropod-infecting strains, including other parthenogenesis-inducers from a range

of host insects (Scholz et al. 2020; Lindsey et al. 2016). While *Wolbachia* are maternally

transmitted, across longer evolutionary time scales there is a significant amount of horizontal

transfer, and often sister strains infect distantly related hosts (Bailly-Bechet et al. 2017; Scholz

et al. 2020). However, the PI-*Wolbachia* infecting *Trichogramma* appear to have a single origin

(Poorjavad et al. 2012; Schilthuizen and Stouthamer 1997; Almeida and Stouthamer 2017).

These PI-*Wolbachia* still undergo host switching (i.e., there is no co-cladogenesis) (Huigens et

al. 2000; Huigens et al. 2004; Almeida and Stouthamer 2017), but that this clade of *Wolbachia*

seem restricted to a single host genus makes them an interesting case study for host adaptation

and the evolution of their PI effector proteins.

In addition to the PI-*Wolbachia* present in *T. kaykai*, the PSR chromosome found in some males

offers another opportunity to understand the evolution of sex ratio distortion (Zhang and Ferree

2024). One other such PSR chromosome has been described: in the parasitoid wasp *Nasonia*

vitripennis (Hymenoptera: Pteromalidae)(Werren 1991; Nur et al. 1988). The PSR

chromosomes from these two wasp species appear to have independent origins, albeit a very

similar paternal genome elimination phenotype (van Vugt et al. 2003; Zhang and Ferree 2024).

Curiously, both PSR chromosomes seem to have originated from hybridization events in which

chromosomal regions with abundant repetitive elements were transferred in via a close relative

(McAllister and Werren 1997; van Vugt et al. 2005; Van Vugt et al. 2009). In contrast to *T.*

kaykai, *Nasonia* are not known to host any PI symbionts (Beukeboom and Van De Zande 2010).

However, some *Nasonia vitripennis* do host male-killing bacteria: *Arsenophonus nasoniae*

(Gherna et al. 1991; Ferree et al. 2008). The PSR chromosomes are likely playing a key role in

male-rescue which balances the male-eliminating cytoplasmic factors in both systems (either

elimination by conversion to female via PI-*Wolbachia* in *Trichogramma*, or, elimination via death

via *Arsenophonus* in *Nasonia*). The *Wolbachia*-infected line of *T. kaykai* reported here will

enable the long-term maintenance of PSR chromosomes in the lab, and in the future, we hope

to re-collect PSR-containing males from the native range to better understand the evolution of

these selfish genetic elements.

Data Availability

This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the

BioProject accession PRJNA1150630. BioSample accessions for *Trichogramma kaykai* and

Wolbachia strain *w*Tkk are SAMN43292057 and SAMN43292058, respectively. Sequencing

reads are deposited under SRR30339640. Genome assemblies and annotations for the

Trichogramma kaykai nuclear genome, *Trichogramma kaykai* mitochondrial genome, and

Wolbachia strain *w*Tkk are currently processing. Supplemental materials are available on the

GSA Figshare portal and include: (A) Table S1. Nanopore sequencing statistics, (B) Table S2.

Details on draft assembly curation, (C) Table S3. Comparison of interspersed repeats between

Trichogramma kaykai and related species, (D) File S1. *Trichogramma kaykai g*enome

annotations in GFF3 format, and (E) File S2. Analysis notebook with bioinformatics workflows

and scripts. A voucher of the *Trichogramma kaykai* KSX58 colony is available at the University

of California Riverside Insect Collection: UCRC_ENT00496298.

Acknowledgements

We thank Chris Faulk and Carrie Walls of Decorative Genomics (https://decogenomics.com/) for

providing sequencing services. Many thanks to Richard Stouthamer for gifting ARIL the KSX58

colony. We acknowledge that the initial collection of insects to start a colony occurred on the

traditional land of the Mojave People, and we are deeply grateful to the people who have

stewarded the land throughout generations.

Funder Information

Research reported in this publication was supported by the National Institute of General Medical

Sciences of the National Institutes of Health under award number R35GM150991 to ARIL.

Conflicts of Interest Statement

The authors declare no conflict of interest.

Author Contributions

ARIL provided samples, funding, analytical guidance, and performed some analyses. JC

performed molecular work and bioinformatic analyses. JC and ARIL co-wrote the manuscript

and created figures. Both authors read and approved the manuscript prior to submission.

CITATIONS

Almeida, R., and R. Stouthamer, 2017 Phylogeny of the *Trichogramma* endosymbiont

       Wolbachia, an alpha-proteobacteria (Rickettsiae). Braz. J. Biol. 78 (03): 421-428.

Bailly-Bechet, M., P. Martins-Simões, G. J. Szöllősi, G. Mialdea, M.-F. Sagot *et al.*, 2017 How

       long does *Wolbachia* remain on board? Mol. Biol. Evol. 34 (5): 1183-1193.

Bernt, M., A. Donath, F. Jühling, F. Externbrink, C. Florentz *et al.*, 2013 MITOS: improved de

       novo metazoan mitochondrial genome annotation. Mol. Phylogen. Evol. 69 (2): 313-319.

Beukeboom, L. W., and L. Van De Zande, 2010 Genetics of sex determination in the

       haplodiploid wasp *Nasonia vitripennis* (Hymenoptera: Chalcidoidea). Journal of Genetics

       89: 333-339.

Bordenstein, S. R., and S. R. Bordenstein, 2016 Eukaryotic association module in phage WO

       genomes from *Wolbachia*. Nat. Commun. 7.

Brown, C. L., J. Mullet, F. Hindi, J. E. Stoll, S. Gupta *et al.*, 2022 mobileOG-db: a manually

       curated database of protein families mediating the life cycle of bacterial mobile genetic

       elements. Appl. Environ. Microbiol. 88 (18): e00991-00922.

Bruen, T., and T. Bruen, 2005 PhiPack: PHI test and other tests of recombination. McGill

       University, Montreal, Quebec: 1-8.

Brůna, T., H. Li, J. Guhlin, D. Honsel, S. Herbold *et al.*, 2023 Galba: genome annotation with

       miniprot and AUGUSTUS. BMC Bioinformatics 24 (1): 327.

Burks, R. A., J. D. Pinto, and A. R. I. Lindsey, 2024 *Trichogrammatidae, Chapter 60*.

       Wallingford, UK: CABI.

Cabanettes, F., and C. Klopp, 2018 D-GENIES: dot plot large genomes in an interactive,

       efficient and simple way. PeerJ 6: e4958.

Challis, R., E. Richards, J. Rajan, G. Cochrane, and M. Blaxter, 2020 BlobToolKit–interactive

       quality assessment of genome assemblies. G3: Genes, Genomes, Genetics 10 (4):

       1361-1374.

Chen, L., P.-Y. Chen, X.-F. Xue, H.-Q. Hua, Y.-X. Li *et al.*, 2018 Extensive gene

rearrangements in the mitochondrial genomes of two egg parasitoids, *Trichogramma*

japonicum and *Trichogramma ostriniae* (Hymenoptera: Chalcidoidea:

Trichogrammatidae). Scientific reports 8 (1): 7034.

Cherif, A., R. Mansour, and K. Grissa-Lebdi, 2021 The egg parasitoids Trichogramma: from

laboratory mass rearing to biological control of lepidopteran pests. Biocontrol Sci.

Technol. 31 (7): 661-693.

Cruaud, A., J. Y. Rasplus, J. Zhang, R. Burks, G. Delvare *et al.*, 2024 The Chalcidoidea bush of

life: evolutionary history of a massive radiation of minute wasps. Cladistics 40 (1): 34-63.

De La Filia, A. G., S. A. Bain, and L. Ross, 2015 Haplodiploidy and the reproductive ecology of

Arthropods. Current Opinion in Insect Science 9: 36-43.

Donath, A., F. Jühling, M. Al-Arab, S. H. Bernhart, F. Reinhardt *et al.*, 2019 Improved annotation

of protein-coding genes boundaries in metazoan mitochondrial genomes. Nucleic Acids

Res. 47 (20): 10543-10552.

Emms, D. M., and S. Kelly, 2019 OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 20: 1-14.

Farsi, F., J. Nozari, and V. Hosseininaveh, 2020 Karyotype analysis of *Trichogramma*

embryophagum Htg.(Hymenoptera: Trichogrammatidae) using a new method and

estimate its karyotype symmetry. Caryologia 73 (3).

Ferguson, K. B., T. Kursch-Metz, E. C. Verhulst, and B. A. Pannebakker, 2020 Hybrid genome

assembly and evidence-based annotation of the egg parasitoid and biological control

agent *Trichogramma brassicae*. G3: Genes, Genomes, Genetics 10 (10): 3533-3540.

Ferree, P. M., A. Avery, J. Azpurua, T. Wilkes, and J. H. Werren, 2008 A bacterium targets

maternally inherited centrosomes to kill males in *Nasonia*. Curr. Biol. 18 (18): 1409-

1414.

Flack, N., L. Hughes, J. Cassens, M. Enriquez, S. Gebeyehu *et al.*, 2024 The genome of

Przewalski's horse (*Equus ferus przewalskii*). G3: Genes, Genomes, Genetics: jkae113.

Flynn, J. M., R. Hubley, C. Goubert, J. Rosen, A. G. Clark *et al.*, 2020 RepeatModeler2 for

automated genomic discovery of transposable element families. Proc. Natl. Acad. Sci.

117 (17): 9451-9457.

Fricke, L. C., and A. R. Lindsey, 2024 Identification of parthenogenesis-inducing effector

proteins in *Wolbachia*. Genome Biol. Evol.: evae036.

Geib, S. M., B. Hall, T. Derego, F. T. Bremer, K. Cannoles *et al.*, 2018 Genome Annotation

Generator: a simple tool for generating and correcting WGS annotation tables for NCBI

submission. Gigascience 7 (4): giy018.

Gherna, R. L., J. H. Werren, W. Weisburg, R. Cote, C. R. Woese *et al.*, 1991 *Arsenophonus*

nasoniae gen. nov., sp. nov., the causative agent of the son-killer trait in the parasitic

wasp *Nasonia vitripennis*. Int. J. Syst. Evol. Microbiol. 41 (4): 563-565.

Gokhman, V. E., 2020 Chromosomes of parasitic wasps of the superfamily Chalcidoidea

(Hymenoptera): An overview. Comparative Cytogenetics 14 (3): 399.

Gokhman, V. E., F. F. Pereira, and M. A. Costa, 2017 A cytogenetic study of three parasitic

wasp species (Hymenoptera, Chalcidoidea, Eulophidae, Trichogrammatidae) from Brazil

using chromosome morphometrics and base-specific fluorochrome staining.

Comparative Cytogenetics 11 (1): 179.

Gokhman, V. E., and D. L. Quicke, 1995 The last twenty years of parasitic Hymenoptera

karyology: an update and phylogenetic implications. J. Hymenoptera Res. 4: 41-63.

Gottlieb, Y., and E. Zchori-Fein, 2001 Irreversible thelytokous reproduction in *Muscidifurax*

uniraptor. Entomol. Exp. Appl. 100 (3): 271-278.

Grant, J. R., E. Enns, E. Marinier, A. Mandal, E. K. Herman *et al.*, 2023 Proksee: in-depth

characterization and visualization of bacterial genomes. Nucleic Acids Res. 51 (W1):

W484-W492.

Guinet, B., D. Lepetit, S. Charlat, P. N. Buhl, D. G. Notton *et al.*, 2023 Endoparasitoid lifestyle

promotes endogenization and domestication of dsDNA viruses. Elife 12: e85993.

Guo, J., B. Bolduc, A. A. Zayed, A. Varsani, G. Dominguez-Huerta *et al.*, 2021 VirSorter2: a

multi-classifier, expert-guided approach to detect diverse DNA and RNA viruses.

Microbiome 9: 1-13.

Hohmann, C. L., and R. F. Luck, 2000 Effect of temperature on the development and thermal

requirements of *Wolbachia*-infected and antibiotically cured *Trichogramma kaykai* Pinto

and Stouthamer (Hymenoptera: Trichogrammatidae). Anais da Sociedade Entomológica

do Brasil 29 (3): 497-505.

Hohmann, C. L., R. F. Luck, and R. Stouthamer, 2001 Effect of *Wolbachia* on the survival and

reproduction of *Trichogramma kaykai* Pinto & Stouthamer (Hymenoptera:

Trichogrammatidae). Neotrop. Entomol. 30 (4): 607-612.

Huang, N., and H. Li, 2023 compleasm: a faster and more accurate reimplementation of

BUSCO. Bioinformatics 39 (10): btad595.

Huigens, M. E., R. P. de Almeida, P. A. H. Boons, R. F. Luck, and R. Stouthamer, 2004 Natural

interspecific and intraspecific horizontal transfer of parthenogenesis-inducing *Wolbachia*

in *Trichogramma* wasps. Proc. R. Soc. Lond. B 271 (1538): 509-515.

Huigens, M. E., R. F. Luck, R. H. G. Klaassen, M. F. P. M. Maas, M. J. T. N. Timmermans *et al.*,

2000 Infectious parthenogenesis. Nature 405 (6783): 178-179.

Hunt, B. G., K. M. Glastad, V. Y. Soojin, and M. A. Goodisman, 2013 The function of intragenic

DNA methylation: insights from insect epigenomes. Integr. Comp. Biol.: ict003.

Jeong, G., and R. Stouthamer, 2005 Genetics of female functional virginity in the

parthenogenesis-*Wolbachia* infected parasitoid wasp *Telenomus nawai* (Hymenoptera:

Scelionidae). Heredity 94 (4): 402-407.

Katoh, K., and D. M. Standley, 2013 MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol. Biol. Evol. 30 (4): 772-780.

Kaur, R., J. D. Shropshire, K. L. Cross, B. Leigh, A. J. Mansueto *et al.*, 2021 Living in the

endosymbiotic world of *Wolbachia*: A centennial review. Cell Host & Microbe 29(6): 879-

893.

Knutson, A., 1998 *The Trichogramma manual: a guide to the use of Trichogramma for biological*

control with special reference to augmentative releases for control of bollworm and

budworm in cotton: Texas Agricultural Extension Service, the Texas A&M University

System.

LePage, D. P., J. A. Metcalf, S. R. Bordenstein, J. On, J. I. Perlmutter *et al.*, 2017 Prophage WO

genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. Nature

543 (7644): 243-247.

Li, H., 2016 Minimap and miniasm: fast mapping and de novo assembly for noisy long

sequences. Bioinformatics 32 (14): 2103-2110.

Lindsey, A. R. I., Y. D. Kelkar, X. Wu, D. Sun, E. O. Martinson *et al.*, 2018a Comparative

genomics of the miniature wasp and pest control agent *Trichogramma pretiosum*. BMC

Biol. 16 (1): 54.

Lindsey, A. R. I., D. W. Rice, S. R. Bordenstein, A. W. Brooks, S. R. Bordenstein *et al.*, 2018b

Evolutionary genetics of cytoplasmic incompatibility genes *cifA* and *cifB* in prophage WO

of *Wolbachia*. Genome Biol. Evol. 10 (2): 434-451.

Lindsey, A. R. I., and R. Stouthamer, 2017 The effects of outbreeding on a parasitoid wasp fixed

for infection with a parthenogenesis-inducing *Wolbachia* symbiont. Heredity 119.6: 411-

417.

Lindsey, A. R. I., J. H. Werren, S. Richards, and R. Stouthamer, 2016 Comparative genomics of

a parthenogenesis-inducing *Wolbachia* symbiont. G3: Genes|Genomes|Genetics 6 (7):

2113-2123.

Ma, W. J., and T. Schwander, 2017 Patterns and mechanisms in instances of endosymbiont-

induced parthenogenesis. J. Evol. Biol. 30 (5): 868-888.

533 McAllister, B. F., and J. H. Werren, 1997 Hybrid origin of a B chromosome (PSR) in the parasitic

534    wasp *Nasonia vitripennis*. Chromosoma 106 (4): 243-253.

535 Miura, K., and Y. Tagami, 2004 Comparison of life history characters of arrhenotokous and

536    *Wolbachia*-associated thelytokous *Trichogramma kaykai* Pinto and Stouthamer

537    (Hymenoptera: Trichogrammatidae). Ann. Entomol. Soc. Am. 97 (4): 765-769.

538 Nguyen, L.-T., H. A. Schmidt, A. Von Haeseler, and B. Q. Minh, 2015 IQ-TREE: a fast and

539    effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol. Biol.

540    Evol. 32 (1): 268-274.

541 Nur, U., J. H. Werren, D. G. Eickbush, W. D. Burke, and T. H. Eickbush, 1988 A" selfish" B

542    chromosome that enhances its transmission by eliminating the paternal genome.

543    Science 240 (4851): 512-514.

544 Pannebakker, B. A., L. P. Pijnacker, B. J. Zwaan, and L. W. Beukeboom, 2004 Cytology of

545    *Wolbachia*-induced parthenogenesis in *Leptopilina clavipes* (Hymenoptera: Figitidae).

546    Genome 47 (2): 299-303.

547 Perlmutter, J. I., S. R. Bordenstein, R. L. Unckless, D. P. LePage, J. A. Metcalf *et al.*, 2019 The

548    phage gene wmk is a candidate for male killing by a bacterial endosymbiont. PLoS Path.

549    15 (9).

550 Pinto, J. D., 2006 A review of the New World genera of Trichogrammatidae (Hymenoptera). J.

551    Hymenoptera Res. 15 (1): 38-163.

552 Pinto, J. D., R. Stouthamer, and G. R. Platner, 1997 A new cryptic species of *Trichogramma*

553    (Hymenoptera: Trichogrammatidae) from the Mojave Desert of California as determined

554    by morphological, reproductive and molecular data. Proc. Entomol. Soc. Wash. 99 (2):

555    238-247.

556 Pololov, A. A., 2012 The smallest insects evolve anucleate neurons. Arthropod Struct. Dev. 41

557    (1): 29-34.

Polilov, A. A., 2015 Small is beautiful: features of the smallest insects and limits to

miniaturization. Annu. Rev. Entomol. 60: 103-121.

Poorjavad, N., S. H. Goldansaz, T. Machtelinckx, L. Tirry, R. Stouthamer *et al.*, 2012 Iranian

Trichogramma: ITS2 DNA characterization and natural *Wolbachia* infection. BioControl

57: 361-374.

Rempel, A., and R. Wittler, 2021 SANS serif: alignment-free, whole-genome-based phylogenetic

reconstruction. Bioinformatics 37 (24): 4868-4870.

Russell, J. E., L. Nunney, M. Saum, and R. Stouthamer, 2018 Host and symbiont genetic

contributions to fitness in a *Trichogramma–Wolbachia* symbiosis. PeerJ 6: e4655.

Russell, J. E., M. Saum, V. Burgess, K. Bollavaram, and T. Donnell, 2016 Influence of

parthenogenesis-inducing *Wolbachia* infection and sexual mode on *Trichogramma*

kaykai (Hymenoptera: Trichogrammatidae) fitness. Ann. Entomol. Soc. Am.: saw093.

Russell, J. E., and R. Stouthamer, 2010 Sex ratio modulators of egg parasitoids. Egg

parasitoids in agroecosystems with emphasis on Trichogramma: 167-190.

Russell, J. E., and R. Stouthamer, 2011 The genetics and evolution of obligate reproductive

parasitism in *Trichogramma pretiosum* infected with parthenogenesis-inducing

Wolbachia. Heredity 106 (1): 58-67.

Schilthuizen, M., and R. Stouthamer, 1997 Horizontal transmission of parthenogenesis-inducing

microbes in *Trichogramma* wasps. Proc. R. Soc. Lond. B 264 (1380): 361-366.

Scholz, M., D. Albanese, K. Tuohy, C. Donati, N. Segata *et al.*, 2020 Large scale genome

reconstructions illuminate *Wolbachia* evolution. Nat. Commun. 11 (1): 5235.

Seemann, T., 2014 Prokka: rapid prokaryotic genome annotation. Bioinformatics 30 (14): 2068-

2069.

Shropshire, J. D., J. On, E. M. Layton, H. Zhou, and S. R. Bordenstein, 2018 One prophage WO

gene rescues cytoplasmic incompatibility in *Drosophila melanogaster*. Proc. Natl. Acad.

Sci. 115 (19): 4987-4991.

Stouthamer, R., J. A. J. Breeuwer, R. F. Luck, and J. H. Werren, 1993 Molecular-identification of

microorganisms associated with parthenogenesis. Nature 361 (6407): 66-68.

Stouthamer, R., J. G. Hu, F. van Kan, G. R. Platner, and J. D. Pinto, 1999 The utility of internally

transcribed spacer 2 DNA sequences of the nuclear ribosomal gene for distinguishing

sibling species of *Trichogramma*. BioControl 43 (4): 421-440.

Stouthamer, R., and D. J. Kazmer, 1994 Cytogenetics of microbe-associated parthenogenesis

and its consequences for gene flow in *Trichogramma* wasps. Heredity 73: 317-327.

Stouthamer, R., R. F. Luck, and W. D. Hamilton, 1990 Antibiotics cause parthenogenetic

Trichogramma (Hymenoptera, Trichogrammatidae) to revert to sex. Proc. Natl. Acad.

Sci. 87 (7): 2424-2427.

Stouthamer, R., and F. Mak, 2002 Influence of antibiotics on the offspring production of the

Wolbachia-infected parthenogenetic parasitoid *Encarsia formosa*. J. Invertebr. Pathol. 80

(1): 41-45.

Stouthamer, R., J. E. Russell, F. Vavre, and L. Nunney, 2010 Intragenomic conflict in

populations infected by Parthenogenesis Inducing *Wolbachia* ends with irreversible loss

of sexual reproduction. BMC Evol. Biol. 10: 12.

Stouthamer, R., M. van Tilborg, J. H. de Jong, L. Nunney, and R. F. Luck, 2001 Selfish element

maintains sex in natural populations of a parasitoid wasp. Proc. R. Soc. Lond. B 268

(1467): 617-622.

Tarailo-Graovac, M., and N. Chen, 2009 Using RepeatMasker to identify repetitive elements in

genomic sequences. Current protocols in bioinformatics 25 (1): 4.10. 11-14.10. 14.

Tulgetske, G. M., and R. Stouthamer, 2012 Characterization of intersex production in

Trichogramma kaykai infected with parthenogenesis-inducing *Wolbachia*.

Naturwissenschaften 99 (2): 143-152.

van Vugt, J., S. de Nooijer, R. Stouthamer, and H. de Jong, 2005 NOR activity and repeat

sequences of the paternal sex ratio chromosome of the parasitoid wasp *Trichogramma*

kaykai. Chromosoma 114 (6): 410-419.

van Vugt, J. F. A., M. Salverda, J. H. de Jong, and R. Stouthamer, 2003 The paternal sex ratio

chromosome in the parasitic wasp Trichogramma kaykai condenses the paternal

chromosomes into a dense chromatin mass. Genome 46 (4): 580-587.

Van Vugt, J. J., H. de Jong, and R. Stouthamer, 2009 The origin of a selfish B chromosome

triggering paternal sex ratio in the parasitoid wasp *Trichogramma kaykai*. Proc. R. Soc.

Lond. B 276 (1676): 4149-4154.

Verhulst, E. C., B. A. Pannebakker, and E. Geuverink, 2023 Variation in sex determination

mechanisms may constrain parthenogenesis-induction by endosymbionts in haplodiploid

systems. Current Opinion in Insect Science: 101023.

Werren, J. H., 1991 The paternal-sex-ratio chromosome of *Nasonia*. Am. Nat. 137 (3): 392-402.

Werren, J. H., and D. M. Windsor, 2000 *Wolbachia* infection frequencies in insects: Evidence of

a global equilibrium? Proc. R. Soc. Lond. B 267 (1450): 1277-1285.

Wu, X., A. R. I. Lindsey, P. Chatterjee, J. H. Werren, R. Stouthamer *et al.*, 2020 Distinct

epigenomic and transcriptomic modifications associated with *Wolbachia*-mediated

asexuality. PLoS Path. 16 (3): e1008397.

Zhang, X., and P. M. Ferree, 2024 PSRs: Selfish chromosomes that manipulate reproductive

development, pp. 66-73 in *Semin. Cell Dev. Biol.* Elsevier.

Zhang, X., Z. Jiang, X. Jiao, Y. Yu, Z. Wang *et al.*, 2023 Genome Assembly and Comparative

Analysis of the Egg Parasitoid Wasp *Trichogramma dendrolimi* Shed Light on the

Composition and Evolution of Olfactory Receptors and Venoms. Insects 14 (2): 144.