1 **Title: Efficient pheromone navigation via antagonistic detectors**

2 **Authors:** Xuan Wan[1,2], Tingtao Zhou*[3,4], Vladislav Susoy*[5], Alessandro Groaz[1,2], Core Francisco Park[5], John
3 F. Brady[3,4], Aravinthan D.T. Samuel[5], Paul W. Sternberg†[1,2].
4 *These authors contributed equally to this work.
5 † corresponding author
6
Author affiliations:
1. Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA
91125, USA.
2. Tianqiao and Chrissy Chen Institute for Neuroscience, California Institute of Technology, Pasadena,
CA 91125, USA.
3. Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA
91125, USA.
4. Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA
91125, USA.
5. Department of Physics and Center for Brain Science, Harvard University, Cambridge, MA, USA
17
18 **Corresponding author:** Paul W. Sternberg

19 **Address:** Tianqiao and Chrissy Chen Institute for Neuroscience, California Institute of Technology,
20 Pasadena, CA 91125, USA.

21 **Email:** pws@caltech.edu

22 **Author information:**

Name	Email	ORCID
Xuan Wan	wanxuan@caltech.edu	0000-0002-6165-6340
Tingtao Zhou	tingtaoz@caltech.edu	0000-0002-1766-719X
Vladislav Susoy	vladislav_susoy@fas.harvard.edu	0000-0002-0669-0087
Alessandro Groaz	agroaz@caltech.edu	0000-0002-3873-7409
Core Francisco Park	corefranciscopark@g.harvard.edu	0000-0002-9542-2913
John F. Brady	jfbrady@caltech.edu	0000-0001-5817-9128
Aravinthan D.T. Samuel	samuel@g.harvard.edu	0000-0002-1672-8720
Paul W. Sternberg	pws@caltech.edu	0000-0002-7699-0173

23
Abstract

Chemotaxis to a moving potential mate emitting a volatile sex pheromone poses a navigation challenge
that requires rapid, precise responses to maximize reproductive success. Volatile chemicals form
gradients that differ from soluble compounds, potentially making navigation based on comparisons
between spatially separated sensors unreliable for small-bodied animals. Contrary to this model of
simple spatial comparison, *C. elegans* employs an antagonistic strategy, comparing inputs from head
(AWA) and tail (PHD) sensory neurons with distinct response properties. Despite sharing a receptor, SRD-
1, these detectors play different roles: AWA head neurons promote forward movement and acceleration,
while tail PHD neurons induce reversals and deceleration. In increasing pheromone gradients AWA
dominates; whereas decreasing gradients inactivate AWAs, allowing PHDs to fine-tune the response and
correct the trajectory. Head AWAs are essential for mate-searching, while tail PHDs are crucial for
complex tasks. Using a minimal-parameter computational model that recapitulates key findings, we infer

the interplay between head and tail signals in adaptive navigation. This study reveals a sexually
dimorphic dual-detector system that integrates antagonistic sensory inputs from head and tail neurons
to enable adaptive navigation strategies essential for efficient free-moving target location in dynamic
environments.

40
41 **Keywords:** sensory integration, navigation, chemotaxis modeling, sex pheromone, sexual dimorphism
42
Introduction
For most animals, finding a mate is an essential task and is an especially significant challenge for
organisms that rely on volatile pheromones for mate detection, as the potential mate often moves in a
complex environment and establishes a fleeting, low concentration pheromone gradient. To obtain a
mechanistic understanding of how animals can rise to this challenge, we studied navigation in *C. elegans*
as it has a highly accessible nervous system.

49
Self-fertilizing *C. elegans* hermaphrodites emit a volatile pheromone when sperm are depleted [1-5].
Volatile chemosensory signals are fundamental to facilitating complex behaviors such as strategic
navigation, active sampling, and sequential decision-making across diverse species; they are crucial for
reproductive success [6-8]. As sex pheromones guide both innate and learned mate-searching activities in
many animals, they provide an opportunity to study the neurobiological underpinnings of sexually
dimorphic behaviors [9-18]. For example, the differential responses of males and females to identical
stimuli, such as sex pheromones, raise questions about the sensory and neural circuit bases of sexual
dimorphism in behavior [19-22].

58
C. elegans produces both soluble and volatile sex attractants. Soluble nematode-specific ascarosides are
modulators of social/sexual behaviors and developmental regulation [23-41]. In contrast, this study focuses
on a volatile, non-ascaroside sex pheromone (hereafter referred to as "the volatile sex pheromone") that
remains chemically unidentified but plays a critical role in sexual communication by signaling fertility and
sexual maturity, as it can only be produced by hermaphrodites with no sperm due to depletion or
dysfunction [3,42]. As hermaphrodites with no sperm can be considered females we classify this
pheromone as a female sex pheromone. The only known receptor for volatile sex pheromone is the G
protein-coupled receptor (GPCR) SRD-1, which is expressed in the sex-shared amphid neuron AWA in
males only, implying underlying sexually dimorphic sensory processing by that neuron [4].

68
The complexity of neural circuit questions makes studies in simple models such as *C. elegans* highly
useful to rigorously define neural algorithms used by animals. The neural circuitry underlying
chemoattraction in *C. elegans* has been mapped with increasing precision, suggesting that sensory
neurons process environmental stimuli with remarkable reliability [43]. Besides of the sensitive sensory
neurons, it also has a network of interneurons that plays critical roles in processing sensory information
and guiding behavioral responses that collectively facilitate navigation and avoidance behaviors: AIA
amplifies olfactory signals, AIB controls reversals, AIY mediates response to gradients, AIZ modulates
turns, and AVA initiates backward movement [44-52]. These circuits facilitate essential survival behaviors,
employing mechanisms such as temporal integration [43], rapid desensitization [53], and the modulation of
behavioral responses based on previous experiences and current conditions [44,48,54-65]. In short, the
probabilistic nature of behavioral outcomes in *C. elegans* can be traced to the dynamic state of the
neural network beyond sensory neurons, potentially explaining the variability in behavior outcomes from
identical stimuli in the same individual [54,66].

82
Previous studies have shown that *C. elegans* employs sensory detectors on opposite ends of its body for
rapid avoidance responses to aversive stimuli like SDS, H2O2, and glycerin [67-69], where quick response is
prioritized over accuracy. In contrast, our study uncovers a novel volatile attractant chemotaxis
mechanism in which dual detectors are instead used for target searching, a process that demands both
rapid responses and high precision to successfully locate the target. Here, we describe a unique
attractant dual-sampling algorithm that utilizes controlled expression of a single receptor to achieve
sexually dimorphic chemosensory responses and an innate, adaptive navigation strategy for efficient
volatile chemical chemotaxis.
91
In this study, we discover that the tail neuron, PHD, in addition to AWA, expresses SRD-1 and responds to
the sex pheromone as well. We demonstrate that PHD is specifically required for navigation during
challenging tasks. Using optogenetics, we compare the contributions of head and tail neurons to
behavioral responses. To understand their interaction, we investigate the neural circuitry connecting
them, and revisiting their response properties, we find that head and tail neurons detect distinct aspects
of the pheromone gradient and concentration. Based on these findings, we propose a computational
model to help elucidate navigation strategies.
99
100 **Main**
101
Volatile pheromone receptor SRD-1 is expressed in PHD, a male-specific neuron in the tail
Male *C. elegans* sense the volatile female sex pheromone via SRD-1 [4,42], a GPCR. In hermaphrodites, it is
expressed only in ASIs in head region and PHAs in tail region. Whereas In males, SRD-1 is expressed in
the AWA and ASI sensory neurons in head region. In males, SRD-1 was reported to be expressed in a pair
of tail neurons [70,71], which we now have identified as the newly discovered pair of PHD neurons using
several transcriptional reporters, based on their location, morphology, and marker gene expression
(Figure 1A extrachromosomal array strain (*srd-1*p::mCherry),2A and S1B integrated cGAL-UAS strain (*srd-*
*1*p::cGAL; UAS::GFP), and S1C marker for Ray B neurons cGAL-UAS strain *pkd-2*p::cGAL; UAS::GFP; *srd-*
*1*p::mcherry). We further verified SRD-1 expression in PHD neurons using a split cGAL driver with *lin-48d*
and *srd-1* promoter, where *lin-48* is known to be expressed in PHD neurons [72]. The expression of SRD-1 in
PHD neurons was further supported by recent male-specific neuronal transcriptomics data, which
revealed heterogeneous SRD-1 expression within these neurons [73]. Under normal conditions, only
approximately 40% of PHD neurons express SRD-1 [73]. Our approach also generated a PHD-specific driver
for targeted manipulation of PHD neurons for subsequent experiments (See Figure 2). In *C. elegans*
males, AWA is required for chemotaxis towards the sex pheromone [4,42]. While previous studies have
linked male tail-specific neuron PHD to various aspects of male behavior [72,74], their specific function
remains unclear.
119
AWA neurons are known to respond to multiple volatile attractants [75-79], whereas PHA neurons are
involved in repellent avoidance behaviors [67,76,80], and ASI mediates multiple sensory responses, including
repulsion [81]. The sex-specific expression pattern of SRD-1 suggests that the differing behavioral responses
to the sex pheromone—strong attraction in males and weak repulsion in hermaphrodites—may result
from the sexually dimorphic expression of the chemoreceptor SRD-1. Additionally, the dual-site
expression of SRD-1 in both head and tail neurons suggests that males may integrate SRD-1-mediated
sensory inputs from the anterior and posterior regions to navigate effectively toward the volatile
pheromone.
128
Male-specific PHD is activated by the volatile pheromone

We next tested if PHD neurons activate in response to the sex pheromone (Figures 1C). We observed

that PHD is activated by the sex pheromone from both *C. elegans fog-2* hermaphrodites and *C. remanei*

females. To test if this response was mediated via SRD-1, we repeated the experiment in *srd-1(eh1)* loss-

of-function mutant males. *srd-1* mutant males showed no PHD activation upon sex pheromone

stimulation (Figure 1C). Therefore, male-specific PHD neurons sense volatile sex pheromone via SRD-1.

These findings, combined with the known role of PHD in mate-related behaviors [72], suggest its potential

involvement in pheromone-guided navigation. Calcium imaging demonstrates that SRD-1 expression in

both head (AWA) and tail (PHD) neurons is not merely anatomical but functionally mediates pheromone

responses, providing direct evidence that males employ dual detectors to sense pheromone sources.

PHD neurons enhance mate-searching efficiency in complex environments

To test the hypothesis that these spatially separated detectors for pheromone actually aid navigation, we

used four types of chemoattraction assays designed to represent varying levels of difficulty males might

encounter in natural conditions, comparing intact males to those with HisCl inhibited-PHD neurons

(Figure 2). We achieved specific expression of the HisCl effector (a chemogenetic silencer) in PHD under

the control of a split cGAL-UAS line: *lin-48d*p::NLS::cGAL (DBD); *srd-1*p::NLS::cGAL (AD) (Figure 2A), and

our experimental design included wild-type males and PHD-HisCl lines without histamine treatment as

controls. Our data showed no significant differences between the two control groups in their response to

sex pheromones under all test conditions (Figure 2C-G). Therefore, any observed effects can be

attributed to the treatment-induced inactivation of PHD neurons.

We first challenged PHD-inhibited males to locate a sex pheromone source placed at the same distance

(3 cm) from the source, using two different diameters (6 and 8 cm)agar-filled Petri dishes (Figure 2C, D).

All males located the source within 30 mins. PHD-inhibited males performed as effectively as controls at

both distances, indicating that tail neuron input is not essential for simple, close-range navigation tasks.

By contrast, when we increased the distance and placed males 5 cm from the source in an 8 cm-Petri

dish (Figure 2E), all test subgroup males were unable to locate the source.

We then tested males' ability to locate a more dilute odor. We diluted the sex pheromone concentration

10-fold and observed the responses of both wild-type, HisCl-untreated control, and PHD-inhibited males

(Figure 2F, S2A). PHD-inhibited males exhibited notable difficulties in identifying the diluted pheromone

source, unlike the two control group males. This differential response underscores the essential role of

male tail PHD neurons in navigating toward weak pheromone signals.

To understand the effects of increased distance versus stimulus dilution, we conducted Finite Element

Simulations of the pheromone concentration field (Figure S3). In our assay setup, the stimulus was

placed on the lid of the plate that was half-filled with agar, creating a bi-layer distribution of the

pheromone. As illustrated in Figure S3A, the volatile component diffuses rapidly in the air, while the agar

absorbs and then slowly releases the volatile component. The concentration field simulation revealed

that in experiments where males were positioned far away from the pheromone source, they

encountered areas of low absolute concentration without a clear gradient. Conversely, when males were

tested nearer to a diluted pheromone source (3 cm), their performance, while reduced relative to

controls, was better than in the 5 cm experiment, likely because the diluted source still produced a weak

but discernible Gaussian gradient despite its overall low concentration (Figure S3B). These findings

suggest that the concentration gradient plays a crucial role in guiding males to the pheromone source.

Since *C. elegans* navigate through three-dimensional (3D) soil and rotting fruit in their natural habitat, we

examined the efficacy of PHD-inhibited males in a 3D search paradigm using gellan gum gel (see

Methods), which likely better mimics natural conditions than a 2D Petri dish surface (Figure 2G, H). In
this endpoint assay, we recorded the number of worms that reach the target spot every 5 minutes.
Worms must stop moving at the exact 3D location (depth included) of the pheromone to count as finding
it. Control and test droplets are placed at the same depth to control for vertical bias, avoiding
confounding from a preference for movement toward the surface or bottom. PHD-inhibited males had
reduced proficiency in identifying pheromone sources within 3D spaces, underscoring the importance of
PHD neuron inputs for environmental navigation that mirrors natural conditions (Figure 2G, S2A).
185
Analysis of the distribution of males at multiple timepoints after the addition of stimulus revealed a
uniformly low chemotaxis index for the PHD-inhibited male group across all intervals in weak signal and
3D exploration tasks (Figure S2B). In our 2D experiments involving weak signals, most control males
successfully located the target within 10 minutes. However, in 3D exploration tasks, the peak arrival time
at the target extended to a significantly longer 20 minutes (Figure S2C). Notably, we observed distinct
impairments in the navigational behaviors of PHD-inhibited males between weak signal detection and 3D
spatial exploration tasks. In weak signal detection, the low chemotaxis index resulted from an equal
distribution of males reaching both the test and control spots. Conversely, in 3D searching tasks, the low
chemotaxis index stemmed from fewer males reaching both spots (Figure S2B-D).
195
Optogenetic dissection of SRD-1 positive neurons and PHD neuron's roles in male locomotion
To examine the effects of head versus tail SRD-1 positive neurons on sex pheromone-induced behavior,
we used the red-shifted Chrimson variant of channelrhodopsin [82]. We first evaluated the UAS::Chrimson
effector driven by a pan-neuronal promoter, demonstrating the system's efficacy (Figure S4A): worms
temporarily ceased movement during light stimulation, promptly resuming normal behavior once the
light was switched off.
202
We then activated all SRD-1 positive neurons with Chrimson expressed under the *srd-1* regulatory
control, using an *srd-1*p cGAL driver (Figures 3A ATR- and S4C ATR+). The absence of ATR served as a
control to negate any potential effects of light alone, confirming that the observed changes in velocity
were attributable from targeted neuron activation. Under ATR+ conditions (Figure 3A), males exhibited
continuous forward movement throughout the 12.5s light stimulation period, transitioning to rapid
steering or reduced speed after stimulation ceased, but not in ATR- males (Figure S4C).
209
To further confirm the role of SRD-1-positive neuron stimulation in modulating locomotor behaviors, we
investigated the effects using extended stimulation protocols (30s or 60s light-on intervals followed by
10s light-off intervals) (Figure 3C). During prolonged stimulation, forward movement was sustained,
while turning behaviors remained consistently suppressed. Also, the cessation of stimulation was
associated with reduced movement speed and an increase in turning rate, consistent with observed
dynamics under shorter stimulation conditions. Using the *odr-10*p cGAL driver to activate AWA neurons
in both sexes showed no behavioral change, suggesting AWA activation alone does not drive forward
motion, or the *odr-10* promoter provides insufficient expression levels (Figure S4B).
218
We next targeted PHD neurons exclusively. Using a split cGAL driver under the control of *lin-48d* and *srd-*
1 promoters, we achieved PHD-specific Chrimson expression (Figures 3B ATR+ and S4D ATR-). Under the
ATR+ condition, PHD activation in males elicited a decline in average velocity throughout the red light
stimulation period, after which males resumed movement. However, this was not observed in ATR-
males. This result contrasts with the overall movement patterns observed upon and after the activation
of all SRD-1 positive neurons (See Figure 3A). Previous studies also suggest that PHD neurons have a
potential role in maintenance of backward locomotion during the mating partner body scanning [72]. We

also observed that repeated activation of PHD neurons also resulted in a gradual reduction in velocity
(See Figure 3B right panel), suggesting a persistent slowdown effect on locomotor behavior after
activation.
229
Since we also observed behavioral responses to light stimulation in hermaphrodites, we performed a
quantitative analysis to distinguish the robustness of these responses in hermaphrodites and males
(Figure 3). Hermaphrodite responses were more variable, as also seen in the average velocity (Figure 3c)
and Fourier analyses (Figure 3d). Fourier analysis of total velocity during SRD-1-positive PHD neuron
stimulation revealed prominent peaks at 0.05 Hz and 0.025 Hz, corresponding to the 20 s and 40 s
stimulation protocols, respectively (Figure 3d). The amplitude and sharpness of the spectral peak at 0.05
Hz (or 0.025 Hz) quantifies the strength of the signal, i.e., the consistency and robustness of the
behavioral response to periodic stimulation, and it is consistently stronger in the males' response than
the hermaphrodites' by about a factor of two. Hence, we concluded that males display a more robust
response. The more variable response in hermaphrodites also manifested as larger inter-individual
differences, as shown by comparing the error bars (shaded area) in Figure 3c in the left two panels. Both
these analyses suggest a sex-specific effect of SRD-1-positive neuron activation. Upon PHD neuron
stimulation, the male also showed a large error bar. Recent male neuron transcriptomics data from the
Murphy group revealed that only ~40% of PHD neurons express SRD-1 under normal conditions [73], and
this partial penetrance of SRD-1 expression could explain the behavioral variability we observe in males,
as individual worms would differ in their functional PHD neuron complement.
246
247 **Head and tail neurons mediate distinct micro-behavioral responses to optogenetic activation**
248 To confirm and further investigate how activation of SRD-1 positive neurons in the head versus the tail
249 affects behavior, we optically targeted either the head or tail of freely moving males expressing *srd-1*
250 Chrimson cGAL-UAS.
251
Initial observations showed that males predominantly moved forward in the absence of stimulation
(ATR– controls, Figure 4, S4 and S5), so a short observation window could lead to a false positive result
on forward movement. To evaluate the duration and persistence of this response, red light is applied to
head region for 30s (Figures 4A). As shown in Figure 3, tail stimulation primarily induced reversals, a
rapid transition from forward to backward movement. For the tail region stimulation, we chose short
and repeated pulses alternating every 2s over a 15-cycle period to capture multiple reversals (Figure 4A).
258
After postprocessing, we divided the recording into 2-second segments. For each 2-second segment, we
tabulated the behavioral motif displayed by the male or the first behavioral transition that occurred
during that segment. We used two complementary behavioral classification frameworks to capture
distinct aspects of locomotion dynamics. : Four-Motif Analysis (Figure S5): Focuses on core locomotion
states (forward, backward, forward-to-backward transition, backward-to-forward transition) to highlight
macro-level movement changes during head/tail stimulation. Six-Motif Analysis (Figure 4): Expands
categorization to include nuanced behaviors like self-exploratory (where the worm attempts to probe its
own body) and hesitation (marked by frequent direction changes within the 2-second window), which
better capture micro-behavioral variability.
268
A short light pulse delivered to the tail region induced a transition from forward to backward movement,
accompanied by increased directional hesitation (Figure 4B). During these hesitant periods, males
frequently switched between forward and backward locomotion. However, we observed substantial
individual variability in response to tail stimulation. While most worms reliably responded to light
stimulation and for 7 pulses, they responded an average of 4.3 times. A subset completely ignored the

stimulus. This behavioral heterogeneity may arise from the variable expression of SRD-1 in PHD neurons
[73].
276
277   With 30 seconds of head stimulation, we noted behaviors analogous to those from activating all SRD-1
278   positive neurons, including persistent forward movement during light exposure followed by rapid
279   directional adjustment (Figures 4C, S5C). We also observed that after head stimulation, males exhibited
280   self-exploratory behavior (Figure 4C), using their tails to probe against their bodies, a behavior that can
281   persist for several minutes.
282
Circuit that combines head and tail inputs to control locomotion direction
The male connectome [83] suggests locomotion command neurons are the likely locus of integration of the
head and tail inputs. Specifically, PHDs have strong connections to PVN (ROI1) and EF (ROI2)
interneurons [83] (See Figure 1B), which connect to pre-motor command neurons. The direction of
locomotion is controlled by reciprocal inhibition of forward and reverse command neurons [44,50,84,85]. To
investigate the circuitry between sex pheromone sensory neurons and these motor command neurons,
we specifically activated SRD-1 positive neurons with Chrimson and monitored pan-neuronal calcium
dynamics using GCaMP6s (Figure 5A-C, see Method). SRD-1 positive neurons, including PHD, AWA, and
ASI neurons in males, had robust responses to light stimulation (Figure 5C). In addition, two groups of
neurons within the male tail region—presumably a pair of PVN neurons in one region of interest (ROI1)
and a single EF neuron in ROI2 (Figure 5B-C, S6)—had a strong positive correlation with PHD activity in
ATR+ and not in ATR- males (Figures S7). Here, we used normalized activity for regions of interest (ROIs)
to facilitate comparison across populations of neurons with varying baseline fluorescence, while raw
activity was analyzed for single neurons to preserve their intrinsic activity dynamics and avoid potential
distortions from normalization. In the male head region, we observed a more negative correlation with
light stimulation. About 4 neurons in the head region are positively correlated with AWA neuron activity
and about 10 neurons are negatively correlated (Figure S7).
300
301   We observed a strong negative correlation between the activity of the reversal command neurons (AVA)
302   and SRD-1-positive neurons in the head region (AWA and ASI). Although this inverse relationship was
303   also present in the ATR-untreated control group, it was less pronounced. In ATR-untreated worms, the
304   decrease in AVA activity may have been influenced by factors such as movement of the microscope
305   objective or non-specific light responses.
306
AWA and PHD have distinct response properties
It has been suggested that AWA neurons adjust their response based on fold-changes in odor
concentration, reducing turning behavior with minimal increases in odor concentration [53]. We used
prolonged, behaviorally relevant high concentration sex pheromone stimulation with a thin layer micro-
diffusion (TLMD) unit (Figure S8A, see Method), facilitating pan-neuronal calcium imaging (Figure 5D and
E) [4]. As shown in Figure 5D, upon pheromone application, AWA and ASI neurons showed immediate
calcium influx followed by rapid desensitization to a low, steady state within 1.5 mins, maintaining this
level for at least 5 mins. This pattern indicates robust desensitization to even relatively high pheromone
concentration. PHD neurons exhibited sustained activation for at least 9.5 mins with a gradual decline,
indicative of their threshold-sensing properties (Figure 5E). 62.5% of imaged males exhibited a rapid
response to pheromone exposure and sustained high levels of activity, whereas the remainder displayed
a slow response, with delayed activation (2.5 mins post-stimulus) with a 40% reduced calcium signal
amplitude (Figure 5E, cropped to 250 second; D, full length 600 second). The fast and slow responses of
PHD neurons are particularly intriguing. Recent male neuron transcriptomics data revealed a
heterogenetic expression of SRD-1 in male PHD neurons, which may result in this individual variation.

This pattern may also be caused by PHDs being actively inhibited by other neurons. According to the
connectome, PHDs are post-synaptic to several neurons [83], which may modulate their activity and
contribute to the observed variability in response dynamics.
325
During the initial phase of neuronal activity, before the activation of PHD neurons, pheromone-
stimulated calcium imaging revealed a negative correlation between reversal command neurons (AVA
and AVE) and head sensory neuron activities (AWA and ASI) (Figure 5D). This negative correlation
suggests that the inhibition of reversal command neurons by head neurons is pivotal for forward
movement during gradient ascent. As demonstrated by the set of optogenetic experiments (see Figure 3
and 4), gradient-sensing neurons in the head region dominate PHD input and primarily drive forward
movement, suggesting that head neuron activity may override tail neuron activity.
333
In the subsequent phase of neuronal activity, AWA and ASI activity became hyperpolarized, and all PHD
neurons were activated. This allowed us to examine the direct correlation between PHD neurons and
reversal command neurons (AVA and AVE). We observed that reversal command neurons exhibited a
negative correlation with head sensory neurons and a positive correlation with tail sensory neurons
(PHD) (Figure 5F), suggesting PHD-mediated activation of reversal command neurons, consistent with
the observed reversal behavior during optogenetic activation (see Figures 3, 4).
340
341   **PHD and AWA neurons drive AVA activity at distinct concentration thresholds**
342
To confirm the correlation between reversal command neurons and anterior and posterior SRD-1-
positive neurons in different sex pheromone concentrations, and to determine the concentration
threshold for behavior changes, we fabricated a head and tail multi-inlet microfluidic chip (Figure 6A).
We utilized immobilized worms within microfluidic chips to precisely control the delivery of varying
concentration stimulations to the head or tail independently. This setup enabled accurate stimulus
application to distinct regions and facilitates switching between a control buffer and three different
stimulus concentrations.
350
C. elegans has five pairs of premotor command interneurons, two of which promote forward locomotion
and three of which promote reverse locomotion. These groups are interconnected and control distinct
motor neurons [44,50,84,85]. AVA neuron activity peaks at the start of a reorientation [44,49,50,52,84] and AVA
interneurons integrate opposing signals from head and tail sensory neurons, guiding behavior [68,86]. AVA's
diverse activities across time scales position it as a key neuron in coordinating forward and backward
movements [68,86]. Therefore, monitoring AVA neuron activity provides a useful indicator of the animal's
forward and reversal behaviors.
358
Using diacetyl as a baseline attractant to validate our device (Figure 6B-b), we observed that all tested
concentrations led to decreased AVA neuron activity, suggesting an enhancement of forward movement.
Diacetyl at higher concentrations more effectively suppresses AVA neuron activity. We then tested
pheromones at varying concentrations applied to both ends and separately to the head and tail (Figure
6B-c). Simultaneous stimulus of both ends immediately decreased AVA activity, indicating forward
movement. Remarkably, head-only stimulation caused a greater reduction in AVA activity compared with
stimuli to both ends, whereas tail-only stimulation resulted in increased AVA activity (Figure 6B-d,e); this
difference suggests antagonism of head and tail input, consistent with our optogenetic and sex
pheromone stimulation calcium imaging showing that head neurons activation promotes forward
movement and tail PHD neuron activation triggers reversal (see Figures 3, 4 and 6Ba). These increases
and decreases in AVA activity only last for one minute.

370
We performed area-under-curve (AUC) analysis on the AVA neural activity data (See Method). Notably,
SRD-1 positive neurons in the head and tail show different preferences for sex pheromone
concentrations. Higher pheromone concentrations more effectively suppress AVA activity when applied
to both ends or exclusively to the head (Figure 6Bc-d). In contrast, AVA activation peaked at pheromone
dilutions of 100–500X when applied only to the tail (Figure 6C). But, at dilutions of 500X, 1000X, and
2000X, AVA neurons displayed predominantly random oscillations when pheromones were applied to
the head or both ends (Figure S10D). These findings suggest that higher pheromone concentrations
promote forward movement more effectively, with head neurons strongly inhibiting AVA activity and
weaker tail neuron activation contributing to sustained forward movement and increased speed. At low
concentrations, the inhibition from head neurons diminishes, and tail neuron activation becomes more
prominent, reducing forward movement. We also noted significant spontaneous oscillations in AVA
neuron signals across individuals, a finding shown in similar studies using microfluidic chips [66,87,88]. These
fluctuations are likely due to the inherent spontaneous oscillation nature of AVA neurons, individual
animals' internal status, and the restrictive environment of the microfluidic channels, which may
discomfort the worms and trigger attempts to escape, causing variable AVA neuron signal patterns.
386
Command neurons integrate antagonistic inputs from the head and tail
The integration of inputs from SRD-1-positive neurons in the head and tail regions reveals a sophisticated
signal-processing algorithm that optimizes navigation. Optogenetic stimulation data (Figures 3, 4)
demonstrate that activation of tail-region neurons induces reversal behavior and slows locomotion,
while stimulation of head-region neurons suppresses turning, promotes forward locomotion, and
increases speed. Notably, head-region activation alone mimics the effects of activating all SRD-1-positive
neurons, highlighting the dominant role of anterior SRD-1-positive neurons in guiding behavior. This
aligns with findings from AWA-defective *odr-7* mutants and AWA laser-ablated males [4,42], where AWA
neurons are critical for locating pheromone sources, while ASI and PHD neurons are insufficient on their
own. Conversely, the absence of tail-region PHD neurons impairs pheromone localization under natural,
complex conditions, though it has minimal effects in ideal laboratory settings with high pheromone
concentrations, reduced noise, and simpler navigation challenges (Figure 2C, D). Together, these data
suggest a dual-detector system in males integrating head and tail neurons' sensory input, with tail
neuron activation acting as a corrective mechanism during navigational errors to enhance chemotaxis
efficiency.
402
403 Pheromone stimulation calcium imaging data further elucidates this integration (Figure 5D). By fitting
404 AVA neuronal activity into a linear model incorporating AWA and PHD activity, the best-fit equation has
405 AVA activity = −0.7 × AWA activity + 0.1 × PHD activity − 0.2, indicating that AVA responses are primarily
406 driven by inhibitory input from AWA neurons in the head, with a smaller excitatory contribution from
407 PHD neurons in the tail (Figure 5F).
408
Microfluidics chip-based experiments (Figure 6D) further validated this integration mechanism. The area
under the curve (AUC) of AVA neuron activation during head-specific and tail-specific stimulation closely
matches the trends observed with dual stimulation. This suggests that the neuronal response results
from a combination of antagonistic inputs from head and tail regions, providing a computational
framework for navigating pheromone gradients. This dual-system integration reflects a sex-specific
chemoreceptor-driven behavioral adaptation for efficient navigation in complex environments.
415
A simple computational model for spatially separated detectors navigation in *C. elegans*

Sensory neuron inputs are typically conserved through the connectome and determine the behavioral
outcomes [53]. Summarizing the experimental observations of the strong correlations between the AWA,
PHD and AVA neurons (Figures 5, 6), we propose a simple model to demonstrate the roles of head and
tail sensory inputs. We focused on macroscopic navigation behavior and neglected details of microscopic
traits and changes in worm body curvature in motion.

C. elegans exhibits a remarkable ability to navigate complex environments using chemical cues. Two
established directional navigation strategies have been observed [76,89-94]: (1) Klinokinesis, a random walk
biased toward the stimulus, such as in bacterial chemotaxis, requires comparing concentrations over
time to adjust random turning that lacks directionality towards the correct heading. (2) Klinotaxis, a
more refined navigation, utilizes directional turns to achieve more accurate gradient navigation [89,95-98].
Here, we modeled kinesis and taxis separately.

During kinesis searching, we modeled locomotion as a mixture of active Brownian motion [99] and run-
and-tumble motion [100] (see Methods). Our first key assumption is that sensory inputs from the head and
tail are converted into confidence variables Q^H and Q^T, which then determine the worm's dynamical
state (Figure 7A). As the activity of reversal command neurons AVA and AVE is found to correlate
negatively with AWA activation but positively with PHD neuron activation, we propose a simple form to
qualitatively capture these correlations:

$$\gamma = \gamma_0 e^{-\lambda_K(Q^H - Q^T)}, \; U = U_0 + U_1(Q^H - Q^T + \eta).$$

Here $\gamma_0 = 0.067/s$ is the baseline tumbling rate. $U_0 = 0.064$ mm/s is the baseline average speed, and
$U_1 = 0.03$ mm/s is the magnitude of speed fluctuation. λ_K represents the strength of klinokinesis, and η
represents a Gaussian noise in the velocity. The worm's dynamic state is characterized by its velocity and
tumble rate (U, γ). Loss of AWA or PHD input disrupts Q^H or Q^T computation, leading to impaired speed
and turning regulation. The equations for (U, γ) connect the non-directional kinesis movements to
confidence variables from head and tail inputs, which are subcircuit responses to stimuli on the head
and tail sensory neurons. These mechanisms are also known as orthokinesis (for U) and klinokinesis (for
γ).

Our model is based on the idea that sensory information coming from the worm's head and tail is
translated into a signal that is a measure of 'confidence' (Q), and these signals directly determine the
worm's behavior and movements (Figure 7A). AWA neurons become more active as sex pheromone
concentration increases [4]. In our microfluidics chip-based experiment, we found that input from head
neurons suppressed AVA neuron activity (Figures 5, 6B-d). Therefore, in our model, the confidence signal
based on head input (Q^H) will rise with increasing pheromone concentration until it reaches saturation
(see blue line in Figure 7A). Interestingly, high pheromone concentrations applied to the tail suppress
AVA neuron activity, in contrast to the activation observed at low concentrations (Figures 5, 6B-e).
Furthermore, continuous activation of the PHD neuron will result in a reduction in baseline velocity and
the sprint behavior exhibited upon proximity to the target, suggesting that Q^T should decrease below
zero to capture these observations (see red line in Figure 7A). We propose using the $Q^H - Q^T$ model
(see purple line in Figure 7A) to represent the negative correlation between AVA/AVE command neurons
and AWA activation, and the positive correlation with PHD neuron activation. Importantly, only $Q^H -$
Q^T accurately captures the observed increase in speed towards the end of navigation, which wouldn't be
captured by simply adding the two signals together (compare purple and green lines in Figure 7A).

To determine the dynamics of Q^H and Q^T given external concentration field evolution, we adopt the
simplest dynamics model with a single memory timescale:

$$\frac{dQ^H}{dt} = -k_1 Q^H + k_2^H f_H\left(C, \frac{dC}{dt}\right),$$

$$\frac{dQ^T}{dt} = -k_1 Q^T + k_2^T f_T(C).$$

Here $\frac{1}{k_1}$ is the timescale of memory, while the values of $k_2^{H,T}$ represent the sensitivity of SRD-1 positive neurons. f_H and f_T represent the sensory inputs, and they are chosen to qualitatively reflect the observations in this work and AWA calcium dynamics in [53] (see Methods). Key assumptions are: (1) the long decay time of PHD activity level during constant sex pheromone stimulation, as seen in Figure 5F, indicates that PHD sensory input mainly depends on the absolute concentration. Hence, f_T is only a function of C. (2) PHD response is positive at intermediate concentration ranges but becomes strongly suppressed at high concentrations, as shown in Figure 6C. (3) The relatively short decay time of AWA activity level during constant stimulation, as shown in Figure 2 of ref [53], indicates that AWA sensory response mainly reflects the detected concentration change, with a mild increase as absolute concentration increases. For simplicity, in this model worms will make directional (taxis) turns when they (1) detect a high enough concentration above a threshold value, and (2) if they currently experience a decrease in concentration. Such turns are assumed to be perfectly accurate toward the source direction. Based on our experimental observations, head-inhibited worms will not perform taxis turns.

We performed simulations for navigation on a small Petri dish with the concentration simulation for 30 mins (Figure 7B). Each simulation contains 10000 worms, initialized with random orientations. When both head and tail sensory inputs are missing, $Q^H = Q^T = 0$, the worms perform a standard active random walk with constant mean velocity and tumble rate (Figure 7C). Only a small fraction – about 10% of worms – were able to find the pheromone source by luck, as seen in the grey curve of Figure 7D and SI video S2.

Adding tail sensory input alone significantly enhanced the chance of finding the pheromone source, almost doubling the probability (Figure 7D light blue curve). Adding head sensory input boosts the overall success rate to nearly 100%, simply due to the dominating effect of taxis turns (SI video S1). Overall, with tail input, the trajectories are biased closer to the pheromone source, as shown in Figure 7E and S11. In each simulation, worms start from a small region around the center of the plate and the pheromone source is located at the red point on the left side. We denote the direction from the pheromone source to the plate center as the x direction and measured the average x position of all worms during the entire simulation. This is denoted as <x> on top of each panel in Figure 7E. We notice that <x> is negative whenever tail input is present but becomes positive when tail input is absent. Hence, tail input enables worms to move on average to the left side of the plate, i.e., they behave more attracted to the pheromone source with tail inputs. We compute the maximum distance from the pheromone source that reaches worm reaches during the search, denoted as d_{max}. For successful worms with head inputs, the addition of tail input accelerates their search so that their d_{max} is slightly smaller compared with those without tails. In the absence of either head or tail inputs, worms only perform random walks and stay near the starting position; adding tail input effectively incentivizes them to explore larger areas and be more biased to the source, manifested in significantly larger d_{max} values. Overall, these simulations suggest that tail input significantly improves the success rate of mate-searching.

Discussion
Computational modeling suggests tail input suppresses mistakes
Building on our computational modeling of pheromone diffusion, we find that tail input is essential for suppressing navigation errors arising from the rapid diffusion of volatile attractants. Specifically, our finite element method (FEM) simulations reveal that the pheromone concentration field deviates from Gaussian

distributions, and based on our simulations of the dynamical model, the resulting pheromone distributions
can lead to artificial increases in head confidence Q^H for worms in low concentration regions due to the
finite plate size, leading to erroneous behaviors without tail input. Compared to water-soluble chemicals
studied previously, air-borne pheromones diffuse rapidly, creating a nonlinear process where the agar
initially absorbs pheromones to a high concentration before emitting them back into the air. This non-
Gaussian behavior is characteristic of chemicals transported across air and agar phases and must be
considered in chemotaxis models.

The errors caused by artificial increase in head confidence can be mitigated by the tail. Tail input corrects
these errors by increasing the tumble rate, ensuring more efficient directional adjustments. Moreover,
suppressing PHD activity at high concentrations $Q^T \rightarrow -1$ significantly improves navigation, supported by
GCaMP data showing reduced PHD activity under 1x concentration stimulation (Figure 4). Consequently,
we adopted a functional form for PHD response: $\lim_{C \rightarrow 0} Q^T \rightarrow 0$, $\lim_{C \rightarrow C_0} Q^T \rightarrow 1$, and $\lim_{C \rightarrow \infty} Q^T \rightarrow -1$. These
dynamic properties suggest that the interplay between head and tail neurons is critical for robust
navigation strategies, offering an extensible framework for modeling chemotaxis in complex environments.

Critically, our simulations also revealed that the concentration differences between anterior and
posterior positions are negligible due to the small size of the worms. This fundamental constraint
necessitates an alternative navigation strategy: rather than measuring spatial differences along the body
axis, *C. elegans* employs antagonistic comparison between functionally specialized head (AWA) and tail
(PHD) neurons. This computational innovation allows robust navigation where traditional gradient
sensing would fail.

Sexually dimorphic decision-making in *C. elegans* stems from sensory neuron inputs
The distinct responses of males and females to the same sex pheromone prompt questions about the
neural foundations of behavioral sexual dimorphism [3,4,42]. Our findings indicate that sexual dimorphism
in naïve *C. elegans* males' response to sex pheromone stimulation stems from variations in receptor
expression. This leads to differential sensory neuron activity propagating through neural circuits,
culminating in distinct behaviors. Although existing data from other organisms suggest that sex-specific
pheromone responses arise from sexually dimorphic neural circuits rather than differential sensory
perception [9,101,102], incomplete characterizations of neural networks in more complex model organisms
may bias our interpretation of this phenomenon. *C. elegans* exemplifies an olfactory system where
sophisticated modulation of a single receptor's expression allows both efficient chemotaxis and the
execution of distinct sex-specific behaviors. Therefore, the behavioral outcomes in naïve animal during
navigation can be predicted from activity of the key sex pheromone sensory neurons.

Decision in uncertainty: how worms navigate complex gradients with imperfect sensory input and
inaccurate steering angle control
As a complementary approach to experimental observations of the worms' turning behaviors, we
conducted mathematical analysis for the optimal turn angle and compared it for consistency with our
observed data. We modeled an idealized worm with precise short-term memory of its trajectory and the
ability to precisely detect concentration gradients in its path by traveling along the same direction for a
short duration. We assume that the ideal worm's sensory systems are infallible and the worm's neural
circuitry can process this information, such that sampling along two distinct directions will be theoretically
sufficient to identify the gradient's direction. In practice, however, a non-ideal worm's memory and
sensory perception are inaccurate due to inherent noise, complicating the task of gradient estimation. This

challenge mirrors a regression problem where the objective is to select an optimal turning angle for a second sample that minimizes the uncertainty in determining the gradient's direction. We assumed a steady concentration field that does not vary with time during the sampling of two directions of a non-ideal worm.

Our analysis for a non-ideal worm, grounded in a least-squared-error regression framework, suggests that the optimal choice for a second sampling direction consistently exceeds 90 degrees regardless of the initial sampling orientation relative to the actual gradient direction. With an initial orientation distributed uniformly randomly, the preferable turning angle is approximately 120 degrees (Figure S12). This direction is chosen to be away from the gradient if the initial angle is qualitatively aligned (less than 90 degrees from the gradient) and towards the gradient otherwise. Given the worm's lack of precise knowledge of gradient direction, it faces a choice: turn 120 degrees in either direction or, as a compromise, nearly 180 degrees (Figure S12). Observations suggest that worms likely opt for turns close to 180 degrees in early search stages, indicating a strategic approach to maximize information gain about the gradient's direction. Worms thus intricately balance their behavioral strategies with their sensation, motor control, and cognitive limitations.

A navigation strategy based on an integrative dual-detector mechanism

Dual detectors to control directional movement allow simultaneous comparison of stimulus rather than a single organ comparison in time. In *C. elegans*, the phasmid neurons in the tail, similar to the amphid chemosensory organs in the head, perform a chemosensory function [67]. Indeed, amphid and phasmid neurons influence the worm's backward movements when exposed to repellents, integrating sensory inputs from both the head and tail to promote effective escape behaviors [67,69,103]. The AVA interneurons consolidate signals from the head and tail sensory neurons to steer behavioral choices [103]. Our study provides the first evidence of dual detectors in attractant navigation. Repellents demand rapid avoidance (reversal), while attractant sensing requires complex, targeted navigation strategies.

For repellent avoidance behaviors, worms employ a simple strategy in which both head and tail inputs trigger reversal [67,69,103], distinguishing it from the more complex mechanisms underlying attractant navigation. In attractant navigation, head- and tail- located neurons utilize the same receptor but drive distinct and opposing sensorimotor responses: 1) AWA promotes forward movement and speeding up while PHD induces reverse movement and slowing down. 2) Behavioral output is typically dominated by AWA activation, while PHD input provides antagonistic modulation of AWA-driven behavior. 3) The head neuron exhibits gradient-sensing properties and rapid desensitization in constant concentrations, while the tail neuron displays threshold-sensing properties and remains activated under constant concentrations. 4) PHD neurons are activated at lower pheromone concentration, thus preventing males from making false taxis turns induced by rapid temporal evolution of volatile pheromone concentration. PHD neurons are inactive at high pheromone concentration, promoting rapid acceleration (sprints) as the male approaches the pheromone source. We computationally modeled the response, proposing that males have a measure of confidence that they are proceeding in the right direction; the absence of PHD input disrupts directional confidence, preventing the acceleration response. This demonstrates a fundamental aspect of neural circuitry: different neurons have evolved to handle distinct environmental stimuli.

Contrary to prior hypotheses [104-107], *C. elegans* does not rely solely on simple head-tail concentration comparisons but instead employs a more complex antagonistic mechanism involving head and tail sensory neurons, each with distinct preferred concentrations for optimal activation. Our proposed mechanism focuses not on the spatial separation of receptors but on comparing signals from two distinct sensory neuron types, AWA and PHD, with different response properties. This comparison enables the animal to integrate various aspects of pheromone gradient dynamics and make adaptive behavioral decisions. While

604 the mechanism relies on comparing outputs from these neurons, their differences stem more from their
605 response characteristics rather than their physical locations. This strategy would be particularly beneficial
606 for navigating a volatile gradient, such as a shallow and low-concentration pheromone gradient, where
607 there is minimal difference in concentration between the worm's head and tail, making head-tail absolute
608 concentration comparisons very unreliable.
609
Besides our work, other studies have shown that numerous chemosensory GPCRs are also co-expressed
in the head amphid and tail phasmid neuron classes [67,108-111], which raises the possibility that this
antagonistic detector mechanism may represent a general strategy in *C. elegans* navigation. This
mechanism, integrating distinct neuronal responses to environmental stimuli, allows precise adaptation
to dynamic and complex sensory landscapes. Beyond pheromone navigation, the principles uncovered
here likely extend to other volatile chemosensory contexts, providing a broader framework for
understanding how animals solve the universal challenge of searching for moving targets within noisy and
variable environments.
618
619 **STAR Methods**

620 **Key Resource Table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
E. coli	*Caenorhabditis* Genetics Center (CGC)	WormBase: OP50; WormBase: WBStrain00041969
Experimental models: Organisms/strains		
him-5 (e1490) V	CGC	CB4088 (male enriched mutant)
srd-1(eh1)	CGC	CB5414 (*srd-1* mutant)
syEx1972 [srd-1P::mCherry]; syEx1974 [unc-119::gfp]	This study	PS10451 (*srd-1* reporter; pan-neural marker)
syEx1973 [rab-3P::mCherry]; syIs912[srd-1P::Gal4(sk)::VP64]; syIs300 [15xUAS::GFP]	This study	PS10448 (*srd-1* driver; GFP effector; pan-neural marker)
syEx1972 [srd-1P::mCherry]; syIs913 [pkd-2P::Gal4(sk)::VP64]; syIs300 [15xUAS::GFP]	This study	PS10449 (*srd-1* reporter; *pkd-2* driver; GFP effector)
syIs912[srd-1P::Gal4(sk)::VP64]	This study	PS9477 (*srd-1* driver)
syIs914[srd-1P::Gal4(sk)::VP64]	This study	PS9478 (*srd-1* driver)
syIs912[srd-1P::Gal4(sk)::VP64]; syIs300[UAS::GFP]	This study	PS9473 (*srd-1* driver; GFP effector)
syIs913[pkd-2P::Gal4(sk)::VP64]; syIs300 [15xUAS::GFP]	This study	PS9681 (*pkd-2* driver; GFP effector)
syIs838[lin-48dP::NLS::cGAL (DBD)::gp41-1::N-intein::let-858 3'UTR, srd-1P::NLS::gp41-1::C-intein::cGAL(AD)::let-858 3'UTR]	This study	PS10007 (PHD driver)

syIs838[lin-48dP::NLS::cGAL (DBD)::gp41-1::N-intein::let-858 3'UTR, srd-1P::NLS::gp41-1::C-intein::cGAL(AD)::let-858 3'UTR]; syIs300[15xUAS::GFP]	This study	PS9573 (PHD driver; GFP effector)
syIs888 [gpa-14P::NLS::cGAL (DBD)::gp41-1::N-intein::let-858 3'UTR, rig-3P::NLS::gp41-1::C-intein::cGAL(AD)::let-858 3'UTR]	This study	PS10179 (AVA driver)
syIs888[gpa-14P::NLS::cGAL (DBD)::gp41-1::N-intein::let-858 3'UTR, rig-3P::NLS::gp41-1::C-intein::cGAL(AD)::let-858 3'UTR]]; syIs300[15xUAS::GFP]	This study	PS10087 (AVA driver; GFP effector)
syIs564 [odr-10P::Gal4(sk)::VP64]; syIs300 [15xUAS::GFP]	This study	PS8293 (AWA driver; GFP effector)
syIs340 [15xUAS::hChR (H134R)::EYFP::let-858 3'UTR]; him-5 (e1490) V	This study	PS9782 (Chrimson effector; *him-5*)
syIs340 [15xUAS::hChR (H134R)::EYFP::let-858 3'UTR]; him-5 (e1490) V	This study	PS7043 (Chrimson effector)
syIs371[15xUAS::HisCL::SL2::GFP::let-858 3'UTR]	This study	PS7199 (HisCL effector)
syIs612[15xUAS::GCaMP7b::SL2::mKate2]	This study	PS9046 (GCaMP7b effector)
syIs912[srd-1P::Gal4(sk)::VP64]; syIs612[15xUAS::GCaMP7b::SL2::mKate2]	This study	PS9680 (*srd-1* driver; GCaMP7b effector)
syIs912[srd-1P::Gal4(sk)::VP64]; syIs371[15xUAS::HisCL::SL2::GFP::let-858 3'UTR]	This study	PS10005 (*srd-1* driver; HisCl effector)
syIs912[srd-1P::Gal4(sk)::VP64]; syIs340 [15xUAS::hChR (H134R)::EYFP::let-858 3'UTR]; him-5 (e1490) V	This study	PS9990 (*srd-1* driver; Chrimson effector)
syIs838[lin-48dP::NLS::cGAL(DBD)::gp41-1::N-intein::let-858 3'UTR, srd-1P::NLS::gp41-1::C-intein::cGAL(AD)::let-858 3'UTR]; syIs371[15xUAS::HisCL::SL2::GFP::let-858 3'UTR]	This study	PS10189 (PHD driver; HisCL effector)
syIs838[lin-48dP::NLS::cGAL(DBD)::gp41-1::N-intein::let-858 3'UTR, srd-1P::NLS::gp41-1::C-intein::cGAL(AD)::let-858 3'UTR]; syIs340 [15xUAS::hChR(H134R)::EYFP::let-858 3'UTR]; him-5(e1490)V.	This study	PS10187 (PHD driver; Chrimson effector)
syIs888 [gpa-14P::NLS::cGAL (DBD)::gp41-1::N-intein::let-858 3'UTR, rig-3P::NLS::gp41-1::C-intein::cGAL(AD)::let-858 3'UTR]; syIs612[15xUAS::GCaMP7b::SL2::mKate2]	This study	PS10181 (AVA driver; GCaMP7b effector)
syIs564 [odr-10P::Gal4(sk)::VP64]; syIs340 [15xUAS::hChR (H134R)::EYFP::let-858 3'UTR]	This study	PS9387 (AWA driver; Chrimson effector)
hpIs675 [rgef-1P::GCaMP6s::3xNLS::mNeptune + lin-15(+)]; lin-15(n765) X; him-5 (e1490) V	This study	ZM9627 (pan-neuronal nucleus localized mNeptune and GCaMP6s)
hpIs675 [rgef-1P::GCaMP6s::3xNLS::mNeptune + lin-15(+)]; lin-15(n765) X; him-5 (e1490) V; srd-1(eh1)	This study	PS10450 (pan-neuronal nucleus localized mNeptune and GCaMP6s in *srd-1(eh1)* background)

lite-1(ce314); JAC66[ift-20P:BFP; eat-4P::cyOFP1; unc-17P::mScarlet: acr-5P::BFP; hpIs675 [rgef-1P::GCaMP6s::3xNLS::mNeptune + lin-15(+)]; lin-15(n765) X	This study	ADS1046 (pan-neuronal nucleus localized mNeptune, GCaMP6s, and neuron ID markers)
lite-1(ce314); JAC66[ift-20P:BFP; eat-4P::cyOFP1; unc-17P::mScarlet: acr-5P::BFP; hpIs675 [rgef-1P::GCaMP6s::3xNLS::mNeptune + lin-15(+)]; lin-15(n765) X; syIs912[srd-1P::Gal4(sk)::VP64]; syIs340 [15xUAS::hChR(H134R)::EYFP::let-858 3'UTR]; him-5(e1490).	This study	PS10153 (pan-neuronal nucleus localized mNeptune, GCaMP6s, neuron ID markers; *srd-1* driver; Chrimson effector)
Kp1368 [myo-2P::NLS::mCherry]; syIs371[15xUAS::HisCL::SL2::GFP::let-858 3'UTR]	This study	PS8720 (muscle driver; HisCl effector)
syIs334 [rabP::GAL4::gp41::N::let-858 3' UTR]; syIs340 [15xUAS::hChR(H134R)::EYFP::let-858 3'UTR]; him-5(e1490).	This study	PS8026 (pan-neural driver; Chrimson effector)
Software and algorithms		
Fiji	Schindelin et al., 2012 [112]	https://imagej.net/software/fiji/
MaMuT 0.27	Wolff et al., 2018 [113]	https://imagej.net/plugins/mamut/
Target track	Park et al., 2018 [114]	https://www.nature.com/articles/s41592-023-02096-3
Ilastik software	Berg et al., 2019 [115]	https://www.ilastik.org
Other		
cGAL-UAS bipartite expression toolkit	Nava et al., 2023 [116]	https://www.pnas.org/doi/full/10.1073/pnas.2221680120

Worm Strains

623 Strains of *C. elegans* used in this study originated from the N2 (Bristol) wild-type lineage and were
624 maintained under standard laboratory conditions using Nematode Growth Medium (NGM) plates with *E.*
625 *coli* OP50 as the food source. Detailed information regarding the strains is provided in Key Resource
626 Table.

Microscopy and Imaging

628 Animals were immobilized on a 4% ultrapure agarose pad using 3-5 μL of 2-5 mM levamisole. The
629 imaging was conducted using an inverted laser scanning confocal microscope, Zeiss LSM 980, integrated
630 with a Zeiss Axio Observer 7 with Definite Focus 2. Subsequent image processing was executed via FiJi
631 (ImageJ) software.

Inhibition of PHD Neurons by the HisCl method and Modified Chemoattraction Assay

Inhibition of PHD neurons was achieved through the histamine-gated chloride channel method, as
outlined in the study [117], with slight alterations. Nematode Growth Medium containing histamine (NGM-

HA) was prepared by incorporating histamine dihydrochloride from Sigma Aldrich (1 M stock solution)

into NGM agar at approximately 65 °C just prior to plate pouring. Control plates without histamine were

prepared simultaneously from the same batch of NGM. Plates supplemented with 10 mM histamine

remained effective for a minimum of two weeks when refrigerated at 4 °C, verified by preliminary testing

with a control strain possessing the *myo-2* promoter-driven HisCl effector in the cGAL-UAS system. L4-

stage males were selected and isolated a day in advance of the assays. Histamine treatments were

conducted without the presence of food to eliminate confounding factors. Histamine-induced inhibition

occurred within ten mins of exposure. Once removed from histamine, the animals typically require one

to two hours to fully recover.

The chemoattraction assay plates were prepared using 1.5% agar, 25 mM NaCl, 1.5 mM Tris-base, and

3.5 mM Tris-Cl, and conducted according to the protocol described in the literature [4], but included

modifications to accommodate different plate dimensions, adjustments in the spatial distance between

the stimulus source and the starting point of the animals, and alterations in the sex pheromone

concentration, as depicted in the associated figures (see Figure 1 and S1).

Stimulus droplets are placed on the lid of the plate, and the location of the worms is recorded every 5

mins up to 30 mins post-stimulation to assess their response to the introduced stimulus. In our research,

we utilized two sizes of plates for the assays, with effective worm-moving surfaces measuring 5.5cm and

8.4cm in diameter. For simplicity and clarity in labeling, these were referred to as 6-cm and 8-cm plates,

respectively. In our study, we employed two types of 2D chemoattraction assays distinguished by the

setup and calculation of the Chemoattraction Index (C.I.). For Type 1 assays, constrained by the smaller

plate size, a control buffer spot was not included. Here, the C.I. was derived from the ratio of worms that

reached the test spot to the total number of worms introduced. Conversely, in Type 2 assays, the C.I. was

determined by subtracting the number of worms that reached the control spot from those that arrived

at the test spot, divided by the total number of worms. In each chemoattraction assay, we placed 20

worms at a designated starting point. To ensure the reliability and robustness of our data, we repeated

each assay 20 times across three distinct days.

Chemoattraction Assay

Male chemotaxis was quantified using a chemoattraction assay as previously described (Wan et al.,

2024). Briefly, 2 µL of pheromone extract or M9 buffer (control) was applied to designated spots on the

assay plate, each supplemented with 1 M sodium azide to immobilize responding males. 20 one-day-old

adult males were placed at the origin point, and chemoattraction index (C.I.) were calculated after 30

minutes using the formula: C.I. = $(E - C) / (E + C + N)$, where E, C, and N represent the number of worms

at the experimental, control, or neither spot, respectively. PHD-inhibited worms prepared via the

"Inhibition of PHD Neurons by HisCl Method" .

3D chemoattraction assay employing 0.3g of gellan gum in 100ml chemoattraction assay buffer to create

a navigable gel environment. This composition ensures that worms can navigate through the gel freely,

without becoming immobilized by water or causing structural damage to the gel through movement. The

gellan gum solution is adjusted to match the salt concentration of the chemoattraction assay plates

utilized in this research.

For plate preparation, 22 ml of the gellan gum solution is poured into a 6-cm plate immediately after the

gellan gum is dissolved by heating in a microwave. The solution is allowed to cool to room temperature

before introducing the worms. To embed the worms within the gel, 0.5ml of the gellan gum gel is

removed from the center of the plate by pipetting. Then, 20 worms are picked by worm picker and placed. This cavity is then refilled with 0.7 ml of gellan gum gel, ensuring the worms are fully embedded within the gel. Unlike traditional chemoattraction assays, the stimulus is applied directly onto the surface of the gel rather than on the lid of the plate (see Figure 2B). This approach ensures that the pheromone remains accessible to the worms and provides a localized and consistent gradient for them to follow. In this endpoint assay, trajectories are not recorded; instead, the number of worms reaching the target spot is documented every 5 minutes for up to 30 minutes post-stimulation to evaluate their responses to the introduced stimulus. This scoring method focuses on the culmination of navigational behavior, capturing the effectiveness of the worms' ability to detect and respond to the pheromone gradient over time. To ensure the reliability and robustness of our data, we repeated each assay 20 times across three distinct days. PHD-inhibited worms were prepared via the "Inhibition of PHD Neurons by HisCl Method". We remark that this C.I. quantification is not a projection onto the 2D plane, since worms at the target will stay still afterwards but those not reaching the target will move in all directions. For example, a worm at the same (x,y) coordinate as the pheromone location but vertically below the stimulus droplet (so the z coordinate is different from the droplet) will continue to move around and will not be counted when computing the C.I. To control for vertical bias, the control solution and pheromone are placed on the same z-plane, preventing preference for movement toward the surface or bottom. Therefore, the C.I. reliably identifies worms at the target location in 3D space.

Optogenetics Perturbation Velocity Analysis

The optogenetics perturbation assay was conducted using Chrimson, a red-shifted channelrhodopsin, following protocols outlined by the literature [82,118,119], with slight modifications. All-trans-retinal (Sigma-Aldrich) was incorporated at a concentration of 500 μM into 6-cm diameter plates containing 20mL of chemoattraction assay solution. One-day-old adult animals were placed on these plates and subjected to a red LED stimulus using a WormLab Tracking system (MBF Biosciences). Prior to stimulation, allow 5 worms a 5-minute acclimation period on the assay plate. The LED stimulus was administered in a pattern of 5 seconds baseline record before stimulation, and 10-60 seconds on, followed by 7.5-30 seconds off, at 100% intensity across 6-8 cycles. Implement a 30-second inter-stimulus interval for PHD activation. This provides sufficient time for the worm to return to a stable forward movement state. Worm tracking was conducted using the same WormLab system.

Dissecting Micro-behaviors Under Optogenetic Perturbation

Experiments used 8-cm NGM agar plates seeded with a thin OP50 layer to prevent starvation-induced behaviors. Males were transferred from ATR+ and ATR-culture plates to an assay plate. The plate was mounted on a motorized stage controlled by a joystick. Individual males were tracked and recorded with a 10x 0.3NA Nikon Plan Fluor objective at 10Hz. Illumination was provided by dim, red-filtered light using a HOYA 52 mm R(25A) filter to minimize disturbance to the worms. For the stimulation, we used a Sola Light Engine (lumencor) light source and a 607/36 nm BrightLine excitation filter. The microscope's field diaphragm was adjusted such that only a small area was illuminated when the excitation light was on. We used the SOLA-SE2 software to control the light source. Prior to stimulation, allow a single worm for a 5-minute acclimation period on the assay plate. The male was tracked manually using the motorized stage and the following stimulation regime was applied: 30 seconds without light, followed by twenty 2-second tail pulses (2-second intervals), another 30-second dark period, then 30 seconds of head stimulation, and a final 30-second interval without light.

After postprocessing, we divided the recording into 2-second segments. For each 2-second segment, we tabulated the behavioral motif displayed by the male or the first behavioral transition that occurred during that segment. The following behavioral motifs and transitions were included: forward, backward, transition from forward to backward, transition from backward to forward movement, hesitation (marked by frequent direction changes within the 2-second window), and self-exploration (where the worm attempts to probe its own body). This analysis provides a detailed account of the nuanced responses of male *C. elegans* to optogenetic perturbation, offering insights into the micro-behavioral dynamics elicited by targeted neural stimulation.

Our analysis employs two analyses: The four-motif analysis (Figure S6) focuses on core locomotion states (forward, backward) and transitions (forward to backward, backward to forward). In the four-motif analysis, transitions (e.g., forward-to-backward, backward-to-forward) were classified based on the initial and final directions observed within the 2-second window. For example, a segment with rapid oscillations between forward and backward movement was categorized as "forward-to-backward" if the net directionality shifted from forward to backward, or vice versa.

The six-motif analysis (Figure 4) expands categorization to include self-exploratory and high-frequency direction transitions (renamed from "hesitation"), which capture other behaviors that cannot be easily classified as the "four-motif". The high-frequency direction transitions ("hesitation"), motif specifically identifies intervals with more than 2 direction changes within a 2-second window, representing a distinct exploratory strategy that cannot be unambiguously assigned to single transitions (forward to backward or backward to forward).

Sex Pheromone-Induced Pan-neuronal Calcium Imaging

For sex pheromone extraction, we processed 100 virgin *fog-2* female *C. elegans* by isolating them in 1 ml of NGM buffer within an Eppendorf tube, following a protocol slightly modified from the previous study [4]. These females were washed five times to clear bacterial contamination, then incubated in 100 μl of NGM or M9 buffer for 6 hours. A parallel extraction process utilized for *C. remanei* involved similar steps but with only 5 females.

In Figure 1C, to evaluate PHD activation upon exposure to the volatile pheromone, we performed calcium imaging with a custom-built spinning disk confocal microscope setup [49,74]. Utilizing the ZM9627 strain, which marks all neuronal nuclei with GCaMP6s and mNeptune [74] and incorporates *him-5(e1490)* mutation to enrich for males. We also crossed ZM9627 with the CB5414 *srd-1(eh1)* line. Preparation for imaging involved isolating L4 males 15-20 hours in advance, placing them on 10% agar NGM buffer slides.

The Thin Layer Micro-Diffusion (TLMD) method was adapted from Wan et al. [4] (see Figure S9A). Neuronal excitation was detected by measuring changes in fluorescence intensity using a disposable worm-mounting and a chemical-loading device called the thin-layer micro-diffusion (TLMD) unit. The TLMD unit consisted of a 4% agarose pad (0.2 mm thick, 1.3 mm radius) sandwiched between two autoclave tapes. Before mounting, well-fed transgenic worms were transferred to an empty worm plate for 5 minutes to remove residual bacteria. A 0.2 μL of 1 mM levamisole was then applied to the agarose pad to partially immobilize the worms, which were positioned uniformly with their heads facing outward. A clean 22 × 40 mm coverslip was gently placed over the TLMD unit. At the 30-second time point, 8 μL of the test chemical solution was loaded. Due to capillary action, the solution rapidly filled the space between the

slide and coverslip. Chemical diffusion began immediately (within milliseconds) as the micro-gel pad
became surrounded by the loaded solution. Approximately 5 ul of 0.2 um latex beads (Polysciences, Inc.)
were added to aid immobilization.

Imaging was conducted with a 40x 0.95 NA objective, achieving 200 Hz to record 10 volumes per second,
with each comprising 20 optical slices. Pheromone or control buffer introduction occurred 30 seconds
after initiating imaging, continuing for an additional 3.5-9.5 mins. Subsequent processing and registration
allowed for tracking PHD neuron nuclei using semi-automated methods in MAMUT [112]. Intensities from
GCaMP and mNeptune channels were extracted for analysis within defined ROIs, calculating neuronal
activity ratios and smoothing data with Savitzky-Golay filters. The activity was quantified by normalizing
GCaMP fluorescence against mNeptune. We used the ratio R=F(GCaMP)/F(mNeptune), after applying
Savitzky-Golay filtering to each channel (polynomial order 1 and length 13) Relative intensity values
($\Delta R/R_o$) were calculated as $(R-R_0)/R_0$, where raw fluorescence intensity ratio (R) was extracted from the
region of interest (ROI) of the target neuron in the calcium imaging data, and R_o is the baseline
fluorescence, defined as the mean fluorescence during the quiescent period. Analyses were performed
using custom code.

In Figure 5, our study employed the same custom-built spinning disk confocal microscope setup
mentioned above and a sparse labeling markers pan-neuronal calcium imaging strain, ADS1046, which
features a dual expression of the calcium indicator GCaMP6s and the red fluorescent protein mNeptune
in all neuronal nuclei, alongside the *lite-1(ce314)* mutation to eliminate undesired light-induced neural
activity. This strain also has sparse labeling markers (P*ift-20*:BFP, P*eat-4*:cyOFP1, P*unc-17*:mScarlet, P*acr-*
5:BFP) for neuron identification. Intracellular calcium dynamics were quantified by tracking the calcium
indicator GCaMP6s signals, which enhances green fluorescence intensity within cells. Concurrently,
mNeptune signals were captured to serve as a reference. The intensity ratio of GCaMP to mNeptune
(GFP/RFP) at each time point was plotted. To aid neuron identification, after the recording session, we
acquired additional volumes using the following combination of excitation lasers and emission filters:
30% 640 nm 617/73 nm (200 Hz), 40% 561 nm 578 nm (200 Hz), 35% 488 nm 578 nm (200 Hz), 15% 488
nm (200 Hz), 95% 405 nm 440 or 450 nm (50Hz).

We fit the activity of AVA neurons by a linear combination of AWA and PHD neurons by minimizing the
least-squared error (Figure 5F).

Neuron Tracking
Each neuron was automatically tracked through all 3D volumes with targettrack. We first applied a strong
low pass filter to blur the whole image and thresholded it to find the general location of the head or tail.
After centering each volume by the region found above, we cropped a rectangular ROI that covered the
whole region for 99% of the recording. Then, we manually identified AWA, ASI, AVA, AVE, PHD neurons in
each recording and annotated an average of 5 frames per recording. We used targettrack [114] to
automatically segment and track each neuron in every volume. We trained the 3DCN of targettrack for
10K gradient steps with a batch size of 3. The quality of the tracking was manually verified after tracking.

The xyz coordinates were used to extract pixel intensities around neuronal nuclei. These changes were
then standardized against the baseline pre-stimulation background, resulting in the relative intensity
values ($\Delta R/R0$) being displayed. Data representation is in the form of mean ± standard error of the mean
(s.e.m.), processed using Graphpad Prism 6 and Microsoft Excel for comprehensive analysis. Custom
Matlab code was utilized for graphing, with the $\Delta R/R_0$ average depicted by a black line and the s.e.m. by

a gray-shaded envelope. The duration of stimulation is visually encoded on the plots with a pink color
bar.

Pan-neuronal calcium imaging with optogenetic activation

This experiment used the same sparse labeling makers, pan-neuronal calcium imaging strain mentioned
previously. For the experiments, males were prepared on 10% agar NGM buffer slides. Using a spinning
disc confocal setup, we navigated the challenge of simultaneous neuronal imaging and optogenetic
stimulation of Chrimson, which responds to both 488 nm and 578 nm light, by implementing a strategy
to exploit the GCaMP signal delay post-stimulation, as outlined in Lu et al. (2022) [120]. This involved
interlacing five 10-second imaging/stimulation intervals with 50-second breaks, applying 12% and 35%
intensities for the 488 nm and 561 nm lasers, respectively, while acknowledging potential mechanical
stimulation from objective movement as a confounding variable. During each 5-second
imaging/activation period the piezo-controlled objective was moving, whereas it was not moving
between these periods. We therefore cannot rule out mechanical stimulation that coincided with the
light stimulation as a confounding factor. To aid neuron identification, after the recording session, we
acquired additional volumes using the following combination of excitation lasers and emission filters:
30% 640 nm 617/73 nm (200 Hz), 40% 561 nm 578 nm (200 Hz), 35% 488 nm 578 nm (200 Hz), 15% 488
nm (200 Hz), 95% 405 nm 440 or 450 nm (50Hz).

The same pre-processing and neuron activity tracking via Targettrack was performed as outlined in
"Neuron Tracking". For each neuron, we created 3 ground truth points, 100 volumes apart. Activities of
PHD neurons in ATR+ and ATR- males were extracted and averaged. We calculated the correlation
between the mean PHD activity in ATR+ males and activity traces of other neurons in both ATR+ and ATR-
males. A subset of 12 neurons was identified across all males and we used Generalized Procrustes
analysis [121] and thin plate splines [122] to register all male tail neurons to a common coordinate space.
These landmark neurons were: PHDR, PHDL, PLMR, PCAR, PCAL, HOA, PVY, PDC, PVX, VD12, VA10, and
VA11. A visual inspection identified two regions that contain neurons highly correlated with PHD
(correlation cutoff >0.55). Here we refer to these regions as ROI1 and ROI2 (see Figure 5 and S6). AWA,
ASI, AVA and AVE are identified based on sparse labeling makers, positioning and cell body size. The
relative intensity values (ΔR/R0) are plotted following the methodology outlined in "Sex Pheromone-
Induced Pan-neuronal Calcium Imaging in *C. elegans*" with minor modifications. We normalized the
ΔR/R0 values for each neuron in the ATR+ and ATR- groups to a range of 0 to 1. The dashed line indicated
the starting point of each light stimulation.

Microfluidic chip design and manufacturing

The design of our microfluidic devices is accomplished using SolidWorks software, tailored to the specific
requirements of our experiments. The fabrication process utilizes both photolithography and
polydimethylsiloxane (PDMS) soft lithography, as detailed in the literature [51,123]. Initially, 3-inch silicon
wafers were prepared by coating with a layer of SU-8 photoresist. This layer was subsequently patterned
through photolithography, transforming the wafer into a mold for PDMS casting. For constructs intended
for male worm imaging, the SU-8 2015 photoresist was spun at 1450 rpm to achieve a thickness of 24-25
μm. Following PDMS casting, the microfluidic devices were bonded to glass coverslips using a plasma
cleaner, employing settings of 50mW power for 30 seconds to ensure a robust bond. This fabrication
procedure is designed to produce precise microfluidic devices, enabling the simultaneous and individual
delivery of stimulus to the head and tail regions of *C. elegans* males, while also facilitating concurrent
calcium imaging. This chip allows for accurate delivery of stimulus to both the head and tail regions

separately, with 4 inlets dedicated to each region and an additional outlet for waste. An optimized worm
trapper adaptor from a previous study [124], secures males for imaging while ensuring both ends remain
exposed to the stimulus channels. Designed stoppers within the tail inlet channels prevent the worms
from escaping during injection, without disrupting fluid flow. This approach allows for the detailed study
of neural responses to localized stimuli in a controlled environment.

Sex pheromone preparation for microfluidics chip-based experiments

To obtain large quantities of crude sex pheromone for microfluidics chip-based experiments, we
modified the standard extraction protocol to avoid the time-consuming isolation of L4 stage *fog-2*
females. We used 5-day-old *daf-22* mutant *C. elegans* hermaphrodites, which lack ascaroside synthesis,
to ensure the absence of confounding neuronal responses from this water-soluble sex pheromone
stimulus. Bleach-synchronized *daf-22* mutant worms were washed five times with M9 buffer. L1 stage
arrest was induced by rotating worms in M9 buffer for 12 hours before seeding on culture plates (8-cm
NGM with OP50, approximately 500 L1 worms/plate). Adult worms were collected after three days (at
the onset of egg-laying). Adults were washed from culture plates with M9 buffer and separated from
eggs using the precipitation method (repeated 5-7 times until eggs were absent). Separated adults were
transferred to new OP50-seeded plates, and this washing process was repeated for 6 days to exhaust
self-sperm. Sex pheromone extraction was performed as described in "sex pheromone-induced pan-
neuronal calcium imaging" with the following modification: instead of 100 worms/mL, we used 50 µl of
packed worm volume per 1 mL of M9 buffer (see Figure S10 A-C). Three batches of extracted pheromone
were tested for quality using a chemoattraction assay and combined to create a 100 ml homogenous
stimulus for microfluidics chip-based experiments.

Microfluidics chip-based experiment calcium imaging and data analysis
Young adult *C. elegans* males (PS10181) were loaded into a custom microfluidic chip that allows selective
exposure of head and tail to odor solutions of choice. Solution administration was directed by an array of
8 memory-shape alloy valves (Takasago Fluidics Inc., Japan) individually addressed by an ESP32
microcontroller. Imaging was performed using a Leica DM6000 B microscope equipped with a X-Cite
XLED1 Multi-Triggering LED illumination system (Excelitas Technologies) and iXon EMCCD detectors
(Oxford Instruments Andor). Image acquisition and microfluidic stimulation administration were
controlled and synchronized by a custom Python graphic interface using the Pycro-Manager interface for
the Micro-Manager microscope control software [125]. Time-lapse image segmentation was performed
using the pixel classification function of the Ilastik software and intensity values of GCaMP and RFP were
extracted using a custom Jupyter notebook. The calcium imaging traces were calculated as the ratio of
GCaMP6s/RFP fluorescence to correct for out-of-plane imaging artifacts. Because of the high background
activity of the AVA neuron, baseline values for the calculation of the $\Delta R/R_0$ metric were estimated using
the peakutils Python package. The relative intensity values ($\Delta R/R_0$) are plotted following the
methodology outlined in "Sex Pheromone-Induced Pan-neuronal calcium imaging in *C. elegans*," with
minor modifications. To quantify the effect of suppression/excitation on AVA neuron, we performed
area-under-curve (AUC) analysis. We first compute the baseline AVA activity by averaging over the 3 mins
before stimulation given at the most dilute concentration and subtract this baseline value from all
signals. Then we integrate the area under curve for each worm data for 2 mins after the stimulation. For
suppression, we only account for time intervals when the signal is lower than the baseline value; for
excitation, we only account for time intervals when the signal is higher than the baseline value so that to
naturally terminate the integration of area when the signal disappears. In Figure 6D, we compute the

linear combination of AVA neuron activities with only head or tail stimulation (AVA_head+AVA_tail) and
compare that with the AVA neuron activity with simultaneous head and tail stimulation (AVA_dual).

Concentration field simulation
The concentration evolution inside the petri dishes was simulated using the Finite Element method. The
simulated plates consist of a top layer of 4mm thick air and a bottom layer of 9mm thick agar. The
diffusion coefficient of the sex pheromone was estimated 1.6 mm^2/s in the air and $10^{-3}\ mm^2/s$ in the
agar. The initial condition was set as concentration at the surface of the droplet $c_0 = 0.01\ mol/m^3$ while
$c_0 = 0$ elsewhere. No-flux boundary condition was applied to the walls of the plate. For the dynamical
simulations of worm searching, we take the 2D cross-section of the concentration field at z=-0.5 mm
inside the agar (the air-agar interface is at z=0).

Navigation simulation
The navigation processes were simulated using an in-house extension of the HOOMD-Blue package.
Mathematically, worm locomotion dynamics are described by the following equations:
901
$$d\boldsymbol{r} = U\boldsymbol{q}dt, \boldsymbol{q} = (\cos\boldsymbol{\theta}, \sin\boldsymbol{\theta}), d\boldsymbol{\theta} = \sqrt{2D_R}dW_S + \gamma dW_T.$$
Here, $\boldsymbol{r}$ is the position of the worm, U the velocity magnitude, $\mathbf{q}$ the orientation direction, D_R the rotary
diffusivity of the orientation angle θ. dW_S and dW_T denote the standard Brownian motion on a circle
$-\pi < \theta < \pi$ and a Poisson jump process respectively. $\boldsymbol{\gamma}$ is the intensity of the jump process, i.e., the
tumbling rate. It has been observed [126] that during tumbles (large angle turns), the turning angle
distribution displays a peak near π. Empirically, we represent the turning angle distribution as a mixture
of uniform distribution and a wrapped Gaussian distribution:
908
$$T(\theta, \theta') = \alpha T_0 + (1 - \alpha)T_1.$$
909   Here the turning kernel $T(\theta, \theta')$ is the probability of turning from θ' direction to θ direction. $T_0 =$
910    $1/(2\pi)$ is the uniform distribution and $T_1 = e^{-\frac{(\theta-\pi)^2}{2\sigma^2}}$ is a wrapped Gaussian distribution.
911   We adopt the following functional forms to reflect the above observed sensory neuron dynamics:
912
$$f_H = \frac{\dot{C}}{C_0}\frac{1+tanh(log(C/C_0))}{2},$$
913
$$f_T =.$$
914   In the simulations, we fix $D_R = 0.02$, η $= 1$, $k_1 = 1/60$, $k_2^{H,T} = 1$.

Figure 1: *srd-1* sexual dimorphism in expression patterns and male PHD neuron excitation triggered by sex pheromone depends on the chemoreceptor SRD-1. (A) Expression pattern of *srd-1*::mCherry in male tail. *srd-1* is expressed in a pair of PHD cells located in tail region of male *C. elegans*. PHD neurons are defined based on their location in the tail ganglia, and its process morphology, with P*unc-119*::rfp indicating all neuronal cell bodies. SRD-1 is not expressed in Ray B neurons (Figure S1) as reported in the literature before the existence of PHD was known[70,71]. **(B)** Sexual dimorphic expression of *srd-1* in both head and tail: *srd-1* GFP reporter expression in AWAs, ASIs and male-specific neuron PHDs in males and in ASIs and PHAs in hermaphrodites. *rab-3P::mCherry* marks all neurons; an integrated cGAL-UAS line *(srd-1P*::GAL4; UAS::GFP*)* reports *srd-1* expression. The right panel shows the expression pattern summary of *srd-1* in both males and females, along with the predicted neural circuits it might be involved in, based on their connectome and functional connectome insights. **(C)** Sex pheromone-triggered activation of PHD neurons, exhibiting SRD-1-dependent excitation, was observed through pan-neuronal calcium imaging with *rgef-1P*::GCaMP6s; a pan-neural marker (*rgef-1P*::mNeptune) indicated all neuron cell body locations (sample size: 5-7 worms per condition). Shaded areas represent the s.e.m. White indicates the pre-stimulation stage and yellow shading indicates the time window of buffer application, while pink denotes the period of pheromone application. The y-axis represents $\Delta R/R_0$, where R is defined as the fluorescence ratio of GCaMP to mNeptune.

Figure 2: Exploring the role of PHD neurons in mate-searching efficiency. (A) Verification of SRD-1
expression in the PHD neuron using a split cGAL driver with *lin-48d* and *srd-1* promotor. This approach also
generated a PHD-specific driver for targeted manipulation of the PHD neuron. Male tail image of cGAL-
UAS line: *lin-48d*p::NLS::cGAL (DBD); *srd-1p*::NLS::cGAL (AD), 15XUAS::GFP. **(B)** Schematic representation
of the chemoattraction assay and the calculation of the Chemoattraction Index (C.I.). **(C)-(G)** Impact of
PHD inhibition on male mate-searching efficacy under varied task complexities. Males with HisCl induced
PHDs inhibition (split cGAL-UAS bipartite system line: *lin-48dP*-NLS-cGAL (DBD); *srd-1P*-NLS-cGAL (AD)
driver; UAS:: HisCl effector) can effectively locate sex pheromones in basic navigation tasks as control lines
(WT and HisCl-absent sample); however, their ability to detect a 10X diluted sex pheromone and navigate
through 3D environments is significantly impaired. The sample size for each assay consisted of 120 worms,
20 worms per trials, 6 trails. ***$P < 0.001$, not significant (ns): $P > 0.05$. Two-tailed unpaired t-test. Error
bars represent the s.e.m. **(H)** Schematic diagram of 2D (type 1 and 2) and 3D searching (type 3).

956
957
958
959

Figure 3: Contrasting locomotor responses to optogenetics activation of SRD-1 positive neurons and PHD neurons. (A)(B) Schematic diagram of the optogenetic experiment: Using *srd-1* or PHD-specific drivers

with Chrimson effector in cGAL-UAS lines, we precisely targeted the head or tail regions of freely moving males to evaluate locomotion effects. **(A)** Sex-specific locomotor responses to SRD-1 positive neuron activation in *C. elegans*. SRD-1 positive neurons (*srd-1*P-NLS-cGAL driver) were activated in both sexes using Chrimson (UAS::Chrimson effector). The stimulation protocol consisted of 12.5-second light-on intervals followed by 7.5-second light-off intervals, forming a 20-second cycle. Light stimulation is represented by the pink bar in the figure. Sample size: 120 worms. **(B)** PHD neuron activation effects on male locomotion: triggers slowdown and reversal. Targeting PHD neurons with a split cGAL-UAS bipartite system line: *lin-48dP*-NLS-cGAL (DBD); *srd-1P*-NLS-cGAL (AD) driver). The stimulation protocol consisted of 10-second light-on intervals followed by 30-second light-off intervals, forming a 40-second cycle. Light stimulation is represented by the pink bar in the figure. The sample size for the SRD-1 positive neuron and PHD neuron activation assay included 120 worms per sex. (See ATR- control in Figure S4) **(a, b)** A representative trajectory sample and its corresponding time series of instantaneous velocity magnitude, total velocity V(t), are shown. The red line indicates periods of light stimulation. The yellow circle labeled all the turning events. **(c)** V(t) averaged over 30 samples. **(d)** Fourier analysis of V(t) from Figure 2A and B demonstrates that optogenetic activation of SRD-1 positive neurons and PHD neurons produces expected peaks at 0.05 Hz and 0.025 Hz, corresponding to the 20-second and 40-second stimulation intervals, respectively. Red shade indicates light stimulation. (c), (d) Shaded areas represent the std. **(C)** Persistent enhancement of forward movement upon activation of SRD-1 positive neurons, sustained for up to 60 seconds. The stimulation protocol involved 30- or 60-second light-on intervals followed by 10-second light-off intervals. Sample size: 100 (30s stimulation) and 110 (60s stimulation) worms.

Figure 4: Six-Motif Analysis: Characterizing micro-behavioral responses to optogenetic stimulation of SRD-1 positive neurons in male *C. elegans*' anterior and posterior regions. (A) Summary of all characterized micro-behavioral and their portion among all behaviors. With *srd-1* drivers and Chrimson effectors in cGAL-UAS lines combined with region-specific light sources, we precisely targeted the head or tail regions of freely moving males to analyze locomotion effects and characterize specific micro-behaviors. **(B)-(C)** Region-specific stimulation of SRD-1 positive neurons results in varied behavioral responses, underscoring the distinct roles of these neuron groups in locomotion. **(A)-(C)** The sample size for ATR+ group consisted of 18 worms. The sample size for ATR- group consisted of 12 worms. Green arrows indicate decreases in the ATR+ group during stimulation relative to pre-stimulation, post-stimulation, and ATR- group levels during stimulation. Pink arrows represent increases relative to the same conditions. ***$P < 0.001$; **$P < 0.005$; *$P < 0.05$.

Figure 5: neural activation and correlation patterns following *srd-1* positive neuron optogenetic stimulation and pheromone exposure. (A) Schematic representation of the optogenetic activation protocol, consisting of five 10-second imaging/stimulation intervals interspersed with 50-second breaks. Neuronal activation of AWA, ASI and AVA neurons after optogenetic activation of SRD-1 positive neurons on head region; sample sizes: control and experimental groups (n=8 each). **(B)** Representative activation patterns were observed in male tail neurons upon optogenetic stimulation of SRD-1 positive neurons. Through visual analysis, two regions, designated as ROI1 and ROI2, were identified as containing neurons with strong correlations to PHD activity (correlation coefficient >0.55) (See Figure S6 and S7). The sample sizes: control (n=9) and experimental (n=12). **(C)** Top row: PHD neurons exhibited increased activity upon light stimulation of all SRD-1 positive neurons in the tail region in ATR+ males, but not in ATR- males. Second row: traces of neuronal activities of neurons correlated with PHD ATR+, in two regions (ROI1 and ROI2) which often contain neurons strongly correlated with PHD. The sample sizes: control (n=9) and experimental (n=12) (See Figure S6 and S7). Third and Forth row: neuronal activation in male head regions after optogenetic activation of all SRD-1 positive neurons; sample sizes: control and experimental groups (n=8 each, See Figure S9). **(D)** Calcium responses of PHD neurons over 9.5 mins of pheromone stimulation, with data grouped by response initiation time into two categories. The sample sizes are 13 worms. **(E)** Calcium responses in PHD, AWA, ASI, and reversal commanding neurons AVA and AVE upon 250 seconds of pheromone stimulation in both tail and head regions. Sample sizes for both regions are control (n=5) and experimental (n=8). (B), (C), and (E) Shaded areas represent the s.e.m. **(F)** Fitting AVA neural activity data using a linear combination of AWA and PHD neural activity, focusing on the period both AWA and PHD neurons activated. The best-fit equation for AVA activity is AVA activity=−0.7×AWA activity+0.1×PHD activity−0.2. This model indicates a weighted influence of AWA and PHD neuron activities on AVA neuronal response.

Figure 6: Dissecting reversal command neuron responses across different sex pheromone concentrations

by microfluidics chip-based experiment (A) Schematic of the head and tail microfluidic chip designed for
precise, independent delivery of target concentration stimuli to the head and tail regions separately. This
chip allows for controlled, localized stimulation, enabling the delivery of different stimuli to the head and
tail simultaneously. The chip includes four inlets on both the head and tail sides with a male-size-adapted
worm trapper connected to a worm injection inlet. Two stoppers within the trapper intersect the tail inlet
region, preventing worm escape into the tail inlet or outlet during injection. All inlets converge into a wider
outlet for efficient waste and used worm collection. **(B) (a)** Schematic diagram of stimulation program and
predicted behavioral outcomes. **(b)-(e)** Effects of diacetyl and sex pheromone stimulation on AVA reversal
command interneuron activity in *C. elegans* males. The calcium indicator GCaMP6s was used to monitor
calcium dynamics in the AVA neuron. The AVA-specific split cGAL driver construct consisted of *gpa-*
14P::NLS::cGAL (DBD)::gp41-1::N-intein::*let-858* 3'UTR and *rig-3P*::NLS::gp41-1::C-intein::cGAL(AD)::*let-*
858 3'UTR, paired with the GCaMP effector construct 15xUAS::GCaMP7b::SL2::mKate2. Diacetyl and dual-
sided sex pheromone stimulation reduce AVA activity, as does head-specific pheromone stimulation,
whereas tail-specific pheromone stimulation activates AVA neurons. Shaded areas represent the standard
error of the mean (s.e.m.). The right panel shows area-under-curve (AUC) analysis (see Method), indicating
that dual-sided diacetyl and sex pheromone stimulation, as well as head-specific pheromone stimulation,
exert the strongest inhibitory effects on AVA at the highest tested concentration. Tail-specific pheromone
stimulation produces peak AVA activation at the lowest tested concentration (100-fold dilution). **(C)**
Further testing of three additional diluted concentrations reveals that tail-specific stimulation elicits the
highest activation of the AVA neuron at an intermediate concentration (500-fold dilution), with activation
levels decreasing at both higher and lower concentrations. Each condition (head, tail, or dual location) at
each concentration was tested with a sample size of 8 worms. **(D)** The area under the curve (AUC) for AVA
neuron activation resulting from head-specific stimulation combined with tail-specific stimulation closely
matches the trend observed with dual stimulation alone. The hot pink star indicates the concentration
that resulted in the highest inhibitory or activating effect on AVA neurons.

Figure 7: Simple dynamic model for a navigation mechanism with antagonistic detectors. (A) Schematic plot illustrating how confidence change rates depend on absolute concentration C in the model. C_0 is the reference concentration at the pheromone source. For head input Q^H, a fixed concentration change rate is assumed for the sensory input. At low C, both head (Q^H) and tail (Q^T) inputs accumulate slowly. At

intermediate C, tail input accumulates faster than head input. However, as C further increases, tail input
starts to accumulate slower and is eventually suppressed, while head input signal is monotonically
stronger as C increases until saturation at very high C. The high C regime corresponds to a large $Q^H - Q^T$
which leads to the sprint behavior. **(B)** The left panel illustrates the simulated concentration field within
an 8-cm Petri dish 10 mins after stimulus introduction. The experimental setup involved placing stimulus
on the lid of a Plate half-filled with agar (9mm thickness), establishing a dual-layer distribution for the
volatile component. The black arrowhead marks the layer position corresponding to the details provided
in the bottom panel. The right panel captures the evolution of the concentration field at the air and agar
interface (8.5mm) over a 30-minute duration. **(C-E)** Simulation results to compare the roles of head and
tail inputs. **(C)** Typical trajectories of successful and failed subgroups under different conditions of head/tail
inhibition. **(D)** Fraction of worms that have found source during 30min search task under each condition.
Tail activation results in slightly better performance for head-activated worms (compare purple and green
curves) and about twice better for head-inhibited worms (compare blue and grey curves). No taxis happen
under head inhibition. **(E)** The upper panel shows the simulation results comparing four head/tail input-
enabled conditions, showing the spatial distributions of worms averaged over the duration of the
experiments, i.e., 30 mins. The red dot shows the location of the pheromone source, and all worms start
at an average distance of 15 mm to the right side of the pheromone source. The mean position along the
horizontal direction is displayed as <x>. The bottom panels show the histogram of the maximum distance
from the pheromone source that worms reached. The Y-axis is worm counts, and the X-axis is the
maximum distance. For each of the successful and failed attempts across four different head or tail input-
enabled conditions we plot the histogram from simulated data sample size of 10,000. The mean of
maximum distance and total variance are indicated in each panel.

1100
1101
1102

1103

1104

Graphical abstract:

Spatially separated pheromone detection drives adaptive navigation in *C. elegans* males.

The **upper panel** illustrates how *C. elegans* males utilize spatially separated pairs of sensory neurons, AWA (head) and PHD (tail), that detect the same volatile sex pheromone molecule via the receptor SRD-1. Despite sharing this receptor, head neurons detect gradient changes and activate when concentration increases, suppressing turning, and promoting forward movement and acceleration. In contrast, tail neurons detect absolute concentration and activate at preferred moderate concentrations, inducing reversals and deceleration. The **bottom panel** shows that upon initial pheromone introduction, simultaneous activation of head and tail neurons cancels each other out, causing males to transition from random walking to a stationary anchoring state, preventing them from chasing the concentration wave initially. As pheromone concentration increases, head neurons dominate, driving forward locomotion and acceleration. At the highest concentrations, tail neurons become less active while head neurons are highly activated, leading to rapid, goal-directed movement (sprinting). Incorrect directional choices result in AWA deactivation and persistent PHD activity, inducing pausing and local exploration. Repeated unsuccessful attempts trigger area-restricted searching.

1120

 Supplementary Figure and Legends

Figure S1: Localization of SRD-1 expression in the male tail, excluding ray B neurons. (A) Overview of
srd-1 expression patterns in both sexes. **(B)** Sexual dimorphic expression of *srd-1* in both head and tail:
srd-1 GFP reporter expression in AWAs, ASIs and male-specific neuron PHDs in males and in ASIs and
PHAs in hermaphrodites. *rab-3P::mCherry* marks all neurons; an integrated cGAL-UAS line *(srd-1P*::GAL4;
UAS::GFP*)* reports *srd-1* expression. **(C)** Visualization of *pkd-2* expression in the male tail region using a
cGAL-UAS system: *pkd-2P*-NLS-cGAL driver combined with a UAS::gfp effector. PKD-2 is known to be
expressed in CEM neurons in the head and Ray B neurons in the tail. The figure presents two
representative males. Absence of overlap between PKD-2 (GFP) and SRD-1 (mCherry) expressions in the
male tail region, utilizing *pkd-2p*-NLS-cGAL driver and UAS::*gfp* effector in combination with *srd-*
1p::mCherry, demonstrating distinct expression domains. **(D)** Excitation of GCaMP6s reporter in PHD
neurons observed in pan-neuronal calcium imaging upon sex pheromone stimulation. PLMs are
spontaneously activated before stimulation, and PHDs are activated after applying sex pheromone.

1134

Figure S2: Chemotaxis performance of *him-5* males in weak signal and 3D searching. (A) The distribution of worms at various intervals following stimulus shows a consistently low chemotaxis index in PHD-inhibited males. (B) Presents the number of worms reaching a pheromone target spot at various time points during weak signal searching and 3D searching assays. The analysis of worm distribution at multiple time points after the introduction of the stimulus indicated a consistently low number of the PHD-inhibited males arrived at the test spot across all intervals, highlighting their compromised chemotactic ability. **The sample size for each assay consisted of 120 worms, 20 worms per trials, 6 trails. (C)** Different navigational deficits in PHD-inhibited males across different exploration tasks: in weak signal tasks, their even distribution between test and control spots led to a low chemotaxis index; in 3D tasks, the low index resulted from fewer males reaching the target. The sample size for each assay consisted of 120 worms, 20 worms per trials, 6 trails. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, not significant (ns): $P > 0.05$. Two-tailed unpaired t-test. Error bars represent the s.e.m.

1167

Figure S3: Visualization and dynamics of pheromone concentration fields in chemoattraction assays.
(A) Captures the evolution of the concentration field at the air and agar interface (8.9mm) over a 30-minute duration. The color scales for 1min, 3min, and 6min intervals are consistent, as are those for 10min, 20min, and 30min intervals, as indicated on the right side of the panel, illustrating the dynamic changes in pheromone distribution. (B) Compares the evolution of the concentration field over 30 mins between diluted stimulation and large distance setups. The concentration field simulation shows that in scenarios where males are positioned far from the pheromone source, they experience areas of low absolute concentration lacking a clear gradient. In contrast, proximity to a diluted pheromone source results in a Gaussian distribution with a discernible gradient, despite the overall low concentration level, indicating different challenges in pheromone detection and navigation based on distance and concentration.

Figure S4: Effects of Pan-Neural and AWA Neuron Activation on *C. elegans* Locomotion and ATR- Control of SRD-1 Positive Neuron and PHD Neuron Activation. (A) Demonstrates global neuronal activation via the Chrimson system in *C. elegans*, employing a pan-neuronal promoter (*rab-3p*-NLS-cGAL driver) to drive UAS::Chrimson effector expression. This approach achieved effective neuronal stimulation across the nervous system in both male and female worms, leading to a temporary cessation of movement during light exposure, with immediate resumption of normal behavior upon cessation of the stimulus. The frequency of movement cessation closely followed the stimulation protocol, showing a dominant frequency at 0.05Hz, indicating the system's effectiveness in modulating worm locomotion through pan-neural activation. **(B)** Explores the specific activation of AWA neurons and its impact on locomotion in both sexes of *C. elegans*, utilizing a targeted cGAL-UAS system (*odr-10p*-NLS-cGAL driver for AWA specificity) to activate AWA neurons. Contrary to the pan-neural activation results, selective stimulation of AWA neurons did not alter the average velocity of the worms during a 20-second period of light exposure. This outcome suggests that AWA neuron activation alone may not be sufficient to initiate forward movement, or it may reflect limitations in the activation efficacy of the *odr-10* promoter used. Notably, the frequency analysis of this experimental setup did not exhibit the expected peak at 0.05Hz. The figure includes (a)(b) a representative trajectory sample alongside its total velocity time series V(t), with the red line marking periods of light stimulation; (c) the averaged V(t) across 30 samples, highlighted by a red shade during light stimulation; and (d) the Fourier analysis of V(t), illustrating the differential response patterns to pan-neural versus AWA-specific activation. The sample size for the pan-neural activation assay included 50 worms per sex, while the AWA activation assay involved 110-120 worms per sex. Error bars represent the std. **(C)** Locomotor responses to SRD-1 positive neuron activation in both sexes of *C. elegans*. SRD-1 positive neurons (*srd-1*P-NLS-cGAL driver) were activated in both sexes using Chrimson (UAS::Chrimson effector). The control, without all-trans retinal (ATR), had no change in behavior, eliminating any light effects. **(D)** Locomotor responses to PHD neuron activation in both sexes of *C. elegans*. PHD neurons (split cGAL-UAS line: *lin-48d*P-NLS-cGAL (DBD); *srd-1*P-NLS-cGAL (AD) driver) were activated in both sexes using Chrimson (UAS::Chrimson effector). The control, without all-trans retinal (ATR), also had no change in behavior, eliminating any light effects. **(C) (D)** The sample size for the SRD-1 positive neuron and PHD neuron activation assay included 120 worms per sex (See ATR+ control in Figure 3).

Figure S5: Four-Motif Analysis: Differential locomotion responses to optogenetic stimulation of SRD-1
positive neurons in male *C. elegans*. Utilizes the *srd-1* Chrimson cGAL-UAS system for precise
optogenetic stimulation of SRD-1 positive neurons in the head and tail regions of male *C. elegans*,
revealing distinct locomotive responses. Tail-targeted light pulses induced transitions from forward to
backward movement and hesitancy in movement direction. Continuous head stimulation resulted in
sustained forward motion and exploratory post-stimulation behaviors. Experiments included ATR+ (n=18)
and ATR- (n=12) groups, with sequential recording periods before, during, and after stimulation in both
regions, to assess the nuanced impact on locomotion. **(A)** Tail activation effects on locomotion. A short
light pulse targeting the tail induced a notable transition from forward to backward movement and
increased hesitancy in movement direction during stimulation. **(B)** Head activation and subsequent
behaviors. Continuous head stimulation for 30 seconds prompted sustained forward movement during
light exposure and immediate directional adjustments post-stimulation. After head stimulation, males
engaged in self-exploratory behaviors, using their tails for body probing. Each worm was recorded for 30
seconds before tail stimulation, followed by a 2-second alternating light pulse targeting the tail over 12
cycles. Post-tail stimulation, there was another 30-second recording period before initiating head
stimulation, then the head stimulation lasted for 30 seconds. The worm was then recorded for an
additional 30 seconds after the light stimulation concluded. Error bars represent the s.e.m.

1235

1236

1237

1238

1239

Activation Patterns in Male Tail Neurons Following Optogenetic Stimulation of PHD Neurons

1240 **Figure S6: Activation patterns observed in male tail neurons upon optogenetic stimulation of PHD**
1241 **neurons.** Visual analysis identified two regions, designated ROI1 and ROI2 in multiple males, which
1242 contain neurons with strong correlations to PHD activity in ATR+ group, but not ATR- group (correlation
1243 coefficient >0.55). Sample sizes: control (n=9) and experimental (n=12).

1244

1245

1246

1247

1248 **Figure S7: Neural activity correlations with PHD optogenetic stimulation in tail regions.** Yellow and

1249 purple indicate positive correlations with target neuron activity in ATR- and ATR+ samples, respectively.

1250 Green denote negative correlations in ATR- and ATR+ samples, respectively. The dashed lines indicate

1251 the thresholds for positive and negative correlations. Sample sizes: control (n=9) and experimental

1252 (n=12).

1253

1254

A — Thin Layer Micro-Diffusion (TLMD) unit

B — PHD Neurons GCaMP Signal in Response to Volatile Sex Pheromone Stimulation

1255

Figure S8: TLMD unit for neuronal activation and calcium response measurement. (A) Depicts the Thin Layer Micro-Diffusion (TLMD) unit setup for detecting changes in intracellular calcium signaling via GCaMP6s fluorescence intensity. The device consists of a 4% agarose pad placed between autoclave tapes, used with 0.2μl of 1mM levamisole to partially immobilize well-prepared transgenic worms. Test chemicals are introduced at 30 seconds, with rapid diffusion facilitated by capillary action. **(B)** Calcium responses of PHD neurons during 9.5 mins of pheromone stimulation, with individual animals distinguished by color coding. The sample sizes is 13 worms. Shaded areas represent the s.e.m.

Activation Patterns in Male Head Neurons Following Optogenetic Stimulation of AWA Neurons

ATR-   **ATR+**

ATR - Positive Correlated   ATR + Positive Correlated

ATR - Negative Correlated   ATR + Negative Correlated

1263

1264

1265   **Figure S9: Neural activity correlations with AWA and ASI optogenetic stimulation in head regions.**
1266   Yellow and purple indicate positive correlations with target neuron activity in ATR- and ATR+ samples,
1267   respectively. Green denotes negative correlations in ATR- and ATR+ samples, respectively. The dashed
1268   lines indicate the thresholds for positive and negative correlations. Sample sizes: control and
1269   experimental groups (n=8 each).

Figure S10: Sex pheromone preparation for microfluidics chip-based experiment. (A) Modified Sex
Pheromone Extraction Protocol for microfluidics chip-based experiments. To obtain large quantities of
crude sex pheromone for microfluidics chip-based experiments, we streamlined the standard protocol.
We used *daf-22* mutant *C. elegans* hermaphrodites, eliminating confounding ascaroside signals. Worms
were synchronized and repeatedly washed to deplete self-sperm, and the sex pheromone extraction
ratio was optimized. **(B)** Ratio Optimization: We tested four ratios (100µl/ml, 50µl/ml, 25µl/ml,
12.5µl/ml) for their attractiveness to males using a chemoattraction assay. The 50µl/ml ratio exhibited
attractiveness comparable to 100µl/ml, with attractiveness declining at lower concentrations. Therefore,
50µl/ml was chosen for sex pheromone extraction. **(C)** Quality Control and Homogenization: Three
independent batches of extracted pheromone were quality-controlled using the chemoattraction assay,
all exceeding a predetermined quality threshold suggested by previous studies (C.I. = 0.8). Finally, the
quality-controlled batches were combined to create a homogenous stimulus for microfluidics chip-based
experiments. (B) and (C) Error bars represent the s.e.m. **(D)** Effects of sex pheromone stimulation on AVA
reversal command interneuron activity in *C. elegans* males at Higher Dilutions. The calcium indicator
GCaMP6s was used to monitor calcium dynamics in the AVA neuron. The AVA-specific split cGAL driver
construct consisted of gpa-14P::NLS::cGAL (DBD)::gp41-1::N-intein::let-858 3'UTR and rig-3P::NLS::gp41-
1::C-intein::cGAL(AD)::let-858 3'UTR, paired with the GCaMP effector construct
15xUAS::GCaMP7b::SL2::mKate2. The upper panel shows the calcium-shaded areas representing the
s.e.m. This set of microfluidics chip-based experiments used 500X, 1000X, and 2000X fold diluted
pheromones. The right panel shows area-under-curve (AUC) analysis (see Method). The bottom panel
displays oscillation frequency distributions, revealing spontaneous oscillations in AVA neuron activity
upon exposure to 500X, 1000X, and 2000X diluted pheromones. Heatmaps represent the frequency
responses of AVA neurons under varying stimulation conditions. Each panel corresponds to specific
activation sites (head, tail, and dual activation) and serial pheromone dilutions (set 1: 1X, 10X, 100X; set
2: 500X, 1000X, 2000X). The heatmaps depict the oscillation frequency dynamics of AVA activity over 12
mins for each stimulus program, with color coding indicating the distribution and intensity of oscillation
frequencies. Each condition (head, tail, or dual location) at each concentration was tested with a sample
size of 8 worms.

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335 **Figure S11:** Each panel shows simulation snapshots comparing head/tail input-enabled conditions. In
1336 each snapshot with a time point, 100 randomly chosen worms from 10,000 stimulation events are
1337 displayed as spheres, colored by their instantaneous confidence $Q^H - Q^T$.

1338

1339 **Figure S12:** Ideal model for effective direction sampling to minimize directional uncertainty. δθ is the angle
1340 between the first sampled direction and the concentration gradient. The left panel shows the best angle
1341 to turn to sample the second direction. The right panel shows the directional uncertainty associated with
1342 the best turn angle. Averaging over an assumed uniform distribution of δθ results in a best turn angle close
1343 to 120 degrees.

1344

Author contributions
X.W. and P.W.S. conceived the study and designed the experiments. X.W. and V.S. performed the
behavioral assays, imaging, and optogenetics experiments. T.Z. and X.W. designed the computational
model, performed the simulation and data analysis. X.W. and A.G. performed the microfluidics chip-
based experiment. C.F.P. provided a whole-brain imaging data analysis platform. X.W., T.Z., and P.W.S.
wrote the initial draft of the manuscript. All authors discussed the results and contributed to the final
version of the manuscript. P.W.S., A.D.T.S., and P.W.S. supervised the project.

1352

ACKNOWLEDGMENTS:
We are grateful to David Aguirre, Wilber Palma, Stephanie Nava, and Zachary Blumenfeld for their
assistance during the early stages of this project. We acknowledge Natalie Sim and Elena Ruiz for data
analysis. We also thank Helena Casademunt and Nicholas Markarian for their discussion and suggestions
during the manuscript preparation. This work was supported by: R01 NS113119 (PWS, ADTS), Tianqiao
and Chrissy Chen Institute for Neuroscience senior postdoc fellowship, Tianqiao and Chrissy Chen
Institute for Neuroscience postdoc innovator grant and Tianqiao and Chrissy Chen Institute for
Neuroscience Systems Neuroscience Awards. PWS is Bren Professor of Biology. The *Caenorhabditis*
Genetics Center provided some strains.

DISCLOSURES:
The authors have no conflicts of interest to disclose.

Data availability

Source data are provided with this paper.

Code availability

All code for data analysis is available at https://github.com/edmondztt/pheromone-traj-analysis.

Reference list:

Simon, J. M. & Sternberg, P. W. Evidence of a mate-finding cue in the hermaphrodite nematode *Caenorhabditis elegans*. *Proceedings of the National Academy of Sciences* **99**, 1598-1603 (2002).

Lipton, J., Kleemann, G., Ghosh, R., Lints, R. & Emmons, S. W. Mate searching in *Caenorhabditis elegans*: a genetic model for sex drive in a simple invertebrate. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **24**, 7427-7434, doi:10.1523/JNEUROSCI.1746-04.2004 [doi] (2004).

Leighton, D. H., Choe, A., Wu, S. Y. & Sternberg, P. W. Communication between oocytes and somatic cells regulates volatile pheromone production in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* **111**, 17905-17910, doi:10.1073/pnas.1420439111 (2014).

Wan, X. *et al.* SRD-1 in AWA neurons is the receptor for female volatile sex pheromones in *C. elegans* males. *EMBO Rep* **20**, doi:10.15252/embr.201846288 (2019).

Chasnov, J. R., So, W. K., Chan, C. M. & Chow, K. L. The species, sex, and stage specificity of a *Caenorhabditis* sex pheromone. *Proc Natl Acad Sci U S A* **104**, 6730-6735, doi:10.1073/pnas.0608050104 (2007).

Lockery, S. R. The computational worm: spatial orientation and its neuronal basis in *C. elegans*. *Current opinion in neurobiology* **21**, 782-790 (2011).

Schone, H. *Spatial orientation: The spatial control of behavior in animals and man*. Vol. 588 (Princeton University Press, 2014).

Grasso, F. W. & Basil, J. A. How lobsters, crayfishes, and crabs locate sources of odor: current perspectives and future directions. *Current opinion in neurobiology* **12**, 721-727 (2002).

Gomez-Diaz, C. & Benton, R. The joy of sex pheromones. *EMBO reports* **14**, 874-883, doi:10.1038/embor.2013.140 [doi] (2013).

Moncho-Bogani, J., Lanuza, E., Lorente, M. J. & Martinez-Garcia, F. Attraction to male pheromones and sexual behaviour show different regulatory mechanisms in female mice. *Physiol Behav* **81**, 427-434, doi:10.1016/j.physbeh.2004.01.014 (2004).

Moncho-Bogani, J., Martinez-Garcia, F., Novejarque, A. & Lanuza, E. Attraction to sexual pheromones and associated odorants in female mice involves activation of the reward system and basolateral amygdala. *Eur J Neurosci* **21**, 2186-2198, doi:10.1111/j.1460-9568.2005.04036.x (2005).

Kurtovic, A., Widmer, A. & Dickson, B. J. A single class of olfactory neurons mediates behavioural responses to a *Drosophila* sex pheromone. *Nature* **446**, 542-546 (2007).

Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirián, L. & Dickson, B. J. Neural circuitry that governs *Drosophila* male courtship behavior. *Cell* **121**, 795-807 (2005).

Datta, S. R. *et al.* The *Drosophila* pheromone cVA activates a sexually dimorphic neural circuit. *Nature* **452**, 473-477 (2008).

Miyamoto, T. & Amrein, H. Suppression of male courtship by a *Drosophila* pheromone receptor. *Nature neuroscience* **11**, 874-876 (2008).

Ruta, V. *et al.* A dimorphic pheromone circuit in *Drosophila* from sensory input to descending output. *Nature* **468**, 686-690 (2010).

Hofman, M. A. & Swaab, D. F. The sexually dimorphic nucleus of the preoptic area in the human brain: a comparative morphometric study. *Journal of anatomy* **164**, 55 (1989).

Wei, D., Talwar, V. & Lin, D. Neural circuits of social behaviors: Innate yet flexible. *Neuron* **109**, 1600-1620 (2021).

Auer, T. O. & Benton, R. Sexual circuitry in *Drosophila*. *Current opinion in neurobiology* **38**, 18-26 (2016).

20   Yang, C. F. & Shah, N. M. Representing sex in the brain, one module at a time. *Neuron* **82**, 261-
278 (2014).
21   De Vries, G. J. & Simerly, R. B. in *Hormones, brain and behavior*   137-XXIX (Elsevier, 2002).
22   Cachero, S., Ostrovsky, A. D., Jai, Y. Y., Dickson, B. J. & Jefferis, G. S. Sexual dimorphism in the fly
brain. *Current biology* **20**, 1589-1601 (2010).
23   Peckol, E. L., Troemel, E. R. & Bargmann, C. I. Sensory experience and sensory activity regulate
chemosensory receptor gene expression in *Caenorhabditis elegans*. *Proceedings of the National*
Academy of Sciences of the United States of America **98**, 11032-11038,
doi:10.1073/pnas.191352498 [doi] (2001).
24   Ludewig, A. H. *et al.* Pheromone sensing regulates *Caenorhabditis elegans* lifespan and stress
resistance via the deacetylase SIR-2.1. *Proc Natl Acad Sci U S A* **110**, 5522-5527,
doi:10.1073/pnas.1214467110 (2013).
25   Golden, J. W. & Riddle, D. L. A pheromone influences larval development in the nematode
Caenorhabditis elegans. *Science* **218**, 578-580, doi:10.1126/science.6896933 (1982).
26   Butcher, R. A., Fujita, M., Schroeder, F. C. & Clardy, J. Small-molecule pheromones that control
dauer development in *Caenorhabditis elegans*. *Nature chemical biology* **3**, 420-422 (2007).
27   Jeong, P. Y. *et al.* Chemical structure and biological activity of the *Caenorhabditis elegans* dauer-
inducing pheromone. *Nature* **433**, 541-545, doi:10.1038/nature03201 (2005).
28   Butcher, R. A., Ragains, J. R., Kim, E. & Clardy, J. A potent dauer pheromone component in
Caenorhabditis elegans that acts synergistically with other components. *Proc Natl Acad Sci U S A*
105, 14288-14292, doi:10.1073/pnas.0806676105 (2008).
29   Aprison, E. Z. & Ruvinsky, I. Counteracting Ascarosides Act through Distinct Neurons to
Determine the Sexual Identity of *C. elegans* Pheromones. *Curr Biol* **27**, 2589-2599 e2583,
doi:10.1016/j.cub.2017.07.034 (2017).
30   Aprison, E. Z. & Ruvinsky, I. Sex Pheromones of *C. elegans* Males Prime the Female Reproductive
System and Ameliorate the Effects of Heat Stress. *PLoS Genet* **11**, e1005729,
doi:10.1371/journal.pgen.1005729 (2015).
31   Kaplan, F. *et al.* Ascaroside expression in *Caenorhabditis elegans* is strongly dependent on diet
and developmental stage. *PLoS One* **6**, e17804, doi:10.1371/journal.pone.0017804 (2011).
32   Srinivasan, J. *et al.* A modular library of small molecule signals regulates social behaviors in
Caenorhabditis elegans. *PLoS biology* **10**, e1001237 (2012).
33   Srinivasan, J. *et al.* A blend of small molecules regulates both mating and development in
Caenorhabditis elegans. *Nature* **454**, 1115-1118 (2008).
34   Ludewig, A. H. & Schroeder, F. C. Ascaroside signaling in *C. elegans*. *WormBook*, 1-22,
doi:10.1895/wormbook.1.155.1 (2013).
35   Choe, A. *et al.* Ascaroside signaling is widely conserved among nematodes. *Curr Biol* **22**, 772-780,
doi:10.1016/j.cub.2012.03.024 (2012).
36   von Reuss, S. H. *et al.* Comparative metabolomics reveals biogenesis of ascarosides, a modular
library of small-molecule signals in *C. elegans*. *J Am Chem Soc* **134**, 1817-1824,
doi:10.1021/ja210202y (2012).
37   Choe, A. *et al.* Sex-specific mating pheromones in the nematode Panagrellus redivivus. *Proc Natl*
Acad Sci U S A **109**, 20949-20954, doi:10.1073/pnas.1218302109 (2012).
38   Hsueh, Y. P., Mahanti, P., Schroeder, F. C. & Sternberg, P. W. Nematode-trapping fungi eavesdrop
on nematode pheromones. *Curr Biol* **23**, 83-86, doi:10.1016/j.cub.2012.11.035 (2013).
39   Pungaliya, C. *et al.* A shortcut to identifying small molecule signals that regulate behavior and
development in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A* **106**, 7708-7713,
doi:10.1073/pnas.0811918106 (2009).

40   Izrayelit, Y. *et al.* Targeted metabolomics reveals a male pheromone and sex-specific ascaroside
biosynthesis in *Caenorhabditis elegans*. *ACS chemical biology* **7**, 1321-1325 (2012).
41   Fagan, K. A. *et al.* A Single-Neuron Chemosensory Switch Determines the Valence of a Sexually
Dimorphic Sensory Behavior. *Curr Biol* **28**, 902-914 e905, doi:10.1016/j.cub.2018.02.029 (2018).
42   Chasnov, J. R., So, W. K., Chan, C. M. & Chow, K. L. The species, sex, and stage specificity of a
Caenorhabditis sex pheromone. *Proceedings of the National Academy of Sciences of the United*
States of America **104**, 6730-6735, doi:0608050104 [pii] (2007).
43   Kato, S., Xu, Y., Cho, C. E., Abbott, L. & Bargmann, C. I. Temporal responses of *C. elegans*
chemosensory neurons are preserved in behavioral dynamics. *Neuron* **81**, 616-628 (2014).
44   Luo, L. *et al.* Dynamic encoding of perception, memory, and movement in a *C. elegans*
chemotaxis circuit. *Neuron* **82**, 1115-1128 (2014).
45   Wang, L. *et al.* A gustatory neural circuit of *Caenorhabditis elegans* generates memory-
dependent behaviors in Na+ chemotaxis. *Journal of Neuroscience* **37**, 2097-2111 (2017).
46   Satoh, Y. *et al.* Regulation of experience-dependent bidirectional chemotaxis by a neural circuit
switch in *Caenorhabditis elegans*. *Journal of Neuroscience* **34**, 15631-15637 (2014).
47   Kunitomo, H. *et al.* Concentration memory-dependent synaptic plasticity of a taste circuit
regulates salt concentration chemotaxis in *Caenorhabditis elegans*. *Nature communications* **4**,
2210 (2013).
48   Ji, N. *et al.* Corollary discharge promotes a sustained motor state in a neural circuit for
navigation. *Elife* **10**, e68848 (2021).
49   Venkatachalam, V. *et al.* Pan-neuronal imaging in roaming *Caenorhabditis elegans*. *Proceedings*
of the National Academy of Sciences **113**, E1082-E1088 (2016).
50   Zhen, M. & Samuel, A. D. *C. elegans* locomotion: small circuits, complex functions. *Current*
opinion in neurobiology **33**, 117-126 (2015).
51   Chalasani, S. H. *et al.* Dissecting a circuit for olfactory behaviour in *Caenorhabditis elegans*.
Nature **450**, 63-70 (2007).
52   Gray, J. M., Hill, J. J. & Bargmann, C. I. A circuit for navigation in *Caenorhabditis elegans*.
Proceedings of the National Academy of Sciences **102**, 3184-3191 (2005).
53   Larsch, J. *et al.* A circuit for gradient climbing in *C. elegans* chemotaxis. *Cell reports* **12**, 1748-
1760 (2015).
54   Dobosiewicz, M., Liu, Q. & Bargmann, C. I. Reliability of an interneuron response depends on an
integrated sensory state. *Elife* **8**, e50566 (2019).
55   Luo, L. *et al.* Bidirectional thermotaxis in *Caenorhabditis elegans* is mediated by distinct
sensorimotor strategies driven by the AFD thermosensory neurons. *Proceedings of the National*
Academy of Sciences **111**, 2776-2781 (2014).
56   Colbert, H. A. & Bargmann, C. I. Environmental signals modulate olfactory acuity, discrimination,
and memory in *Caenorhabditis elegans*. *Learning & memory (Cold Spring Harbor, N.Y.)* **4**, 179-191
(1997).
57   Wen, J. Y. M. *et al.* Mutations that prevent associative learning in *C. elegans*. *Behavioral*
neuroscience **111**, 354 (1997).
58   Saeki, S., Yamamoto, M. & Iino, Y. Plasticity of chemotaxis revealed by paired presentation of a
chemoattractant and starvation in the nematode *Caenorhabditis elegans*. *The Journal of*
experimental biology **204**, 1757-1764 (2001).
59   Nuttley, W. M., Atkinson-Leadbeater, K. P. & Van Der Kooy, D. Serotonin mediates food-odor
associative learning in the nematode *Caenorhabditis*elegans. *Proceedings of the National*
Academy of Sciences of the United States of America **99**, 12449-12454,
doi:10.1073/pnas.192101699 [doi] (2002).

60 Zhang, Y., Lu, H. & Bargmann, C. I. Pathogenic bacteria induce aversive olfactory learning in
Caenorhabditis elegans. *Nature* **438**, 179-184 (2005).
61 Vellai, T., McCulloch, D., Gems, D. & Kovács, A. L. Effects of sex and insulin/insulin-like growth
factor-1 signaling on performance in an associative learning paradigm in *Caenorhabditis elegans*.
Genetics **174**, 309-316 (2006).
62 Kauffman, A. *et al. C. elegans* positive butanone learning, short-term, and long-term associative
memory assays. *J Vis Exp*, doi:10.3791/2490 (2011).
63 Sakai, N. *et al.* A sexually conditioned switch of chemosensory behavior in *C. elegans*. *PLoS One*
8, e68676, doi:10.1371/journal.pone.0068676 (2013).
64 Stein, G. M. & Murphy, C. T. *C. elegans* positive olfactory associative memory is a molecularly
conserved behavioral paradigm. *Neurobiol Learn Mem* **115**, 86-94,
doi:10.1016/j.nlm.2014.07.011 (2014).
65 Hong, M. *et al.* Early Pheromone Experience Modifies a Synaptic Activity to Influence Adult
Pheromone Responses of *C. elegans*. *Curr Biol* **27**, 3168-3177 e3163,
doi:10.1016/j.cub.2017.08.068 (2017).
66 Gordus, A., Pokala, N., Levy, S., Flavell, S. W. & Bargmann, C. I. Feedback from network states
generates variability in a probabilistic olfactory circuit. *Cell* **161**, 215-227 (2015).
67 Hilliard, M. A., Bargmann, C. I. & Bazzicalupo, P. *C. elegans* responds to chemical repellents by
integrating sensory inputs from the head and the tail. *Current Biology* **12**, 730-734 (2002).
68 Gat, A. *et al.* Integration of spatially opposing cues by a single interneuron guides decision
making in *C. elegans*. *bioRxiv*, 2023.2001. 2023.525194 (2023).
69 Quintin, S., Aspert, T., Ye, T. & Charvin, G. Distinct mechanisms underlie H2O2 sensing in *C.*
elegans head and tail. *Plos one* **17**, e0274226 (2022).
70 Troemel, E. R., Chou, J. H., Dwyer, N. D., Colbert, H. A. & Bargmann, C. I. Divergent seven
transmembrane receptors are candidate chemosensory receptors in *C. elegans*. *Cell* **83**, 207-218
(1995).
71 Yi, W., Ross, J. M. & Zarkower, D. Mab-3 is a direct tra-1 target gene regulating diverse aspects of
C. elegans male sexual development and behavior. *Development* **127**, 4469-4480 (2000).
72 Molina-García, L. *et al.* Direct glia-to-neuron transdifferentiation gives rise to a pair of male-
specific neurons that ensure nimble male mating. *Elife* **9**, e48361 (2020).
73 Morillo, K. S., St. Ange, J., Weng, Y., Kaletsky, R. & Murphy, C. T. Single-Nucleus Neuronal
Transcriptional Profiling of Male *C. elegans* Uncovers Regulators of Sex-Specific and Sex-Shared
Behaviors. *bioRxiv*, 2024.2012. 2012.628226 (2024).
74 Susoy, V. *et al.* Natural sensory context drives diverse brain-wide activity during *C. elegans*
mating. *Cell* **184**, 5122-5137. e5117 (2021).
75 Sengupta, P., Chou, J. H. & Bargmann, C. I. odr-10 encodes a seven transmembrane domain
olfactory receptor required for responses to the odorant diacetyl. *Cell* **84**, 899-909,
doi:10.1016/s0092-8674(00)81068-5 (1996).
76 Bargmann, C. I. Chemosensation in *C. elegans*. *WormBook : the online review of C.elegans*
biology, 1-29, doi:10.1895/wormbook.1.123.1 [doi] (2006).
77 Ferkey, D. M., Sengupta, P. & L'Etoile, N. D. Chemosensory signal transduction in *Caenorhabditis*
elegans. *Genetics* **217**, iyab004 (2021).
78 Im Choi, J. *et al.* Odor-dependent temporal dynamics in Caenorhabitis elegans adaptation and
aversive learning behavior. *PeerJ* **6**, e4956 (2018).
79 Bargmann, C. I., Hartwieg, E. & Horvitz, H. R. Odorant-selective genes and neurons mediate
olfaction in *C. elegans*. *Cell* **74**, 515-527 (1993).
80 Tran, A. *et al. C. elegans* avoids toxin-producing Streptomyces using a seven transmembrane
domain chemosensory receptor. *Elife* **6**, e23770 (2017).

81   Zhou, Y. *et al.* Potential Nematode Alarm Pheromone Induces Acute Avoidance in *Caenorhabditis*
elegans. *Genetics* **206**, 1469-1478, doi:10.1534/genetics.116.197293 (2017).
82   Klapoetke, N. C. *et al.* Independent optical excitation of distinct neural populations. *Nature*
methods **11**, 338-346 (2014).
83   Cook, S. J. *et al.* Whole-animal connectomes of both *Caenorhabditis elegans* sexes. *Nature* **571**,
63-71 (2019).
84   Cohen, N. & Denham, J. E. Whole animal modeling: piecing together nematode locomotion.
Current Opinion in Systems Biology **13**, 150-160 (2019).
85   Kato, S. *et al.* Global brain dynamics embed the motor command sequence of *Caenorhabditis*
elegans. *Cell* **163**, 656-669 (2015).
86   Meng, J. *et al.* A tonically active master neuron modulates mutually exclusive motor states at
two timescales. (2023).
87   Choi, M.-K., Liu, H., Wu, T., Yang, W. & Zhang, Y. NMDAR-mediated modulation of gap junction
circuit regulates olfactory learning in *C. elegans*. *Nature Communications* **11**, 3467 (2020).
88   Marvin, J. S. *et al.* An optimized fluorescent probe for visualizing glutamate neurotransmission.
Nature methods **10**, 162-170 (2013).
89   Albrecht, D. R. & Bargmann, C. I. High-content behavioral analysis of *Caenorhabditis elegans* in
precise spatiotemporal chemical environments. *Nature methods* **8**, 599-605 (2011).
90   Cheung, B. H., Cohen, M., Rogers, C., Albayram, O. & De Bono, M. Experience-dependent
modulation of *C. elegans* behavior by ambient oxygen. *Current Biology* **15**, 905-917 (2005).
91   Bretscher, A. J. *et al.* Temperature, oxygen, and salt-sensing neurons in *C. elegans* are carbon
dioxide sensors that control avoidance behavior. *Neuron* **69**, 1099-1113,
doi:10.1016/j.neuron.2011.02.023 (2011).
92   Ryu, W. S. & Samuel, A. D. Thermotaxis in *Caenorhabditis elegans* analyzed by measuring
responses to defined thermal stimuli. *Journal of Neuroscience* **22**, 5727-5733 (2002).
93   Gabel, C. V. *et al.* Neural circuits mediate electrosensory behavior in *Caenorhabditis elegans*.
Journal of Neuroscience **27**, 7586-7596 (2007).
94   Sawin, E. R., Ranganathan, R. & Horvitz, H. R. *C. elegans* locomotory rate is modulated by the
environment through a dopaminergic pathway and by experience through a serotonergic
pathway. *Neuron* **26**, 619-631 (2000).
95   Pierce-Shimomura, J. T., Morse, T. M. & Lockery, S. R. The fundamental role of pirouettes in
Caenorhabditis elegans chemotaxis. *Journal of Neuroscience* **19**, 9557-9569 (1999).
96   Iino, Y. & Yoshida, K. Parallel use of two behavioral mechanisms for chemotaxis in *Caenorhabditis*
elegans. *Journal of Neuroscience* **29**, 5370-5380 (2009).
97   Miller, A. C., Thiele, T. R., Faumont, S., Moravec, M. L. & Lockery, S. R. Step-response analysis of
chemotaxis in *Caenorhabditis elegans*. *Journal of Neuroscience* **25**, 3369-3378 (2005).
98   Zariwala, H. A., Miller, A. C., Faumont, S. & Lockery, S. R. Step response analysis of thermotaxis in
Caenorhabditis elegans. *Journal of Neuroscience* **23**, 4369-4377 (2003).
99   Schimansky-Geier, L., Mieth, M., Rosé, H. & Malchow, H. Structure formation by active Brownian
particles. *Physics Letters A* **207**, 140-146 (1995).
100   Berg, H. C. *E. coli in Motion*. (Springer, 2004).
101   Ibarra-Soria, X., Levitin, M. O., Saraiva, L. R. & Logan, D. W. The olfactory transcriptomes of mice.
PLoS genetics **10**, e1004593 (2014).
102   Kohl, J., Ostrovsky, A. D., Frechter, S. & Jefferis, G. S. A bidirectional circuit switch reroutes
pheromone signals in male and female brains. *Cell* **155**, 1610-1623 (2013).
103   Gat, A. *et al.* Integration of spatially opposing cues by a single interneuron guides decision-
making in *C. elegans*. *Cell Reports* **42** (2023).

104 Page, J. L., Dickman, B. D., Webster, D. R. & Weissburg, M. J. Staying the course: chemical signal
spatial properties and concentration mediate cross-stream motion in turbulent plumes. *Journal*
of Experimental Biology **214**, 1513-1522 (2011).
105 Gomez-Marin, A., Duistermars, B., Frye, M. A. & Louis, M. Mechanisms of odor-tracking: multiple
sensors for enhanced perception and behavior. *Frontiers in cellular neuroscience* **4**, 1374 (2010).
106 Borst, A. & Heisenberg, M. Osmotropotaxis in *Drosophila* melanogaster. *Journal of comparative*
physiology **147**, 479-484 (1982).
107 Budick, S. A. & Dickinson, M. H. Free-flight responses of *Drosophila* melanogaster to attractive
odors. *Journal of experimental biology* **209**, 3001-3017 (2006).
108 Taylor, S. R. *et al.* Molecular topography of an entire nervous system. *Cell* **184**, 4329-4347.e4323,
doi:10.1016/j.cell.2021.06.023 (2021).
109 Yemini, E. *et al.* NeuroPAL: a multicolor atlas for whole-brain neuronal identification in *C.*
elegans. *Cell* **184**, 272-288. e211 (2021).
110 Vidal, B. *et al.* An atlas of *Caenorhabditis elegans* chemoreceptor expression. *PLoS Biol* **16**,
e2004218, doi:10.1371/journal.pbio.2004218 (2018).
111 Pereira, L. *et al.* Timing mechanism of sexually dimorphic nervous system differentiation. *Elife* **8**,
doi:10.7554/eLife.42078 (2019).
112 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature methods* **9**,
676-682 (2012).
113 Wolff, C. *et al.* Multi-view light-sheet imaging and tracking with the MaMuT software reveals the
cell lineage of a direct developing arthropod limb. *Elife* **7**, e34410 (2018).
114 Park, C. F. *et al.* Automated neuron tracking inside moving and deforming *C. elegans* using deep
learning and targeted augmentation. *Nature Methods* **21**, 142-149 (2024).
115 Berg, S. *et al.* Ilastik: interactive machine learning for (bio) image analysis. *Nature methods* **16**,
1226-1232 (2019).
116 Nava, S. *et al.* A cGAL-UAS bipartite expression toolkit for *Caenorhabditis elegans* sensory
neurons. *Proceedings of the National Academy of Sciences* **120**, e2221680120 (2023).
117 Pokala, N., Liu, Q., Gordus, A. & Bargmann, C. I. Inducible and titratable silencing of
Caenorhabditis elegans neurons in vivo with histamine-gated chloride channels. *Proc Natl Acad*
Sci U S A **111**, 2770-2775, doi:10.1073/pnas.1400615111 (2014).
118 Cao, M., Chai, C., Liu, J. & Sternberg, P. Application of the red-shifted channel rhodopsin
Chrimson for the *Caenorhabditis elegans* cGAL bipartite system. *MicroPubl Biol* **2018**,
doi:10.17912/2jgw-fj52 (2018).
119 Schild, L. C. & Glauser, D. A. Dual color neural activation and behavior control with chrimson and
CoChR in *Caenorhabditis elegans*. *Genetics* **200**, 1029-1034 (2015).
120 Lu, Y. *et al.* Extrasynaptic signaling enables an asymmetric juvenile motor circuit to produce
symmetric undulation. *Current Biology* **32**, 4631-4644. e4635 (2022).
121 Gower, J. C. Generalized procrustes analysis. *Psychometrika* **40**, 33-51 (1975).
122 Wood, S. N. Thin plate regression splines. *Journal of the Royal Statistical Society Series B:*
Statistical Methodology **65**, 95-114 (2003).
123 Zhou, T. *et al.* AI-aided geometric design of anti-infection catheters. *Science Advances* **10**,
eadj1741 (2024).
124 Chronis, N., Zimmer, M. & Bargmann, C. I. Microfluidics for in vivo imaging of neuronal and
behavioral activity in *Caenorhabditis elegans*. *Nature methods* **4**, 727-731 (2007).
125 Pinkard, H. *et al.* Pycro-Manager: open-source software for customized and reproducible
microscope control. *Nature methods* **18**, 226-228 (2021).

126   Appleby, P. A. The role of multiple chemotactic mechanisms in a model of chemotaxis in *C.*
elegans: different mechanisms are specialised for different environments. *Journal of*
computational neuroscience **36**, 339-354 (2014).

1653