Resolving the Taxonomic Status of the Marbled Toad (Bufonidae: *Incilius marmoreus*): 2RAD-

based Phylogeography Including an Isolated Population in Veracruz, Mexico

Kenneth Wang[1], Todd W. Pierson[2], and Joseph R. Mendelson III[3,4]

Running Head: Phylogeography of *Incilius marmoreus*

[1]H. Milton Stewart School of Industrial and Systems Engineering, Georgia Institute of

Technology, Atlanta, Georgia 30332, USA; Email: kennethwang@gatech.edu; ORCID: 0000-

0001-6954-1583.

[2]Department of Ecology, Evolution, and Organismal Biology, Kennesaw State University,

Kennesaw, Georgia 30144, USA; Email: tpierso3@kennesaw.edu; ORCID: 0000-0003-3818-

5146.

[3]School of Biological Sciences, Georgia Institute of Technology, Atlanta, Georgia 30332, USA.

[4]Department of Research, Zoo Atlanta, Atlanta, Georgia 30315, USA; Email:

jmendelson@zooatlanta.org; ORCID: 0000-0001-5620-377X.

Send reprint requests to JRM.

Inhabiting an extensive range along the Pacific Coast of Mexico and a smaller allopatric region

in the State of Veracruz, *Incilius marmoreus* has an unusual distribution amongst herpetofauna.

Günther (1901) considered the Pacific coastal toads *Bufo argillaceus* and *Bufo lateralis* to be

conspecific with *Incilius [Bufo] marmoreus,* for which the type locality is in Veracruz. Here, we

adopt an integrative approach to reevaluate the phylogeography and taxonomy of *I. marmoreus*

by gathering and analyzing morphological data and conducting phylogenetic and population

genetic analyses from genome-wide SNP data. Our results uphold the current taxonomy by

concurring with Günther (1901). Our phylogenetic and population genetic analyses suggest that

I. marmoreus from Veracruz are closely related to those from Oaxaca, and our coalescent

analyses recovered a north–south split along the Pacific Coast estimated to have occurred ~1.29

Mya followed by a shallow east–west split in the southern lineage that separates the Pacific

coastal populations and the allopatric population in Veracruz ~0.46 Mya. Present gene flow

throughout the western range is evident, and the species displays a marked amount of

morphological and genetic diversity throughout its range. The events that led to the formation of

the population isolated on the coast of Veracruz remain uncertain, but we hypothesize that

climatic and vegetation changes in the Late Pleistocene may have played a role.

Species are fundamental units of biology and systematics (Mayr, 1982; de Queiroz, 2007). Thus,

systematic studies often concern species delimitation—the determination of whether different

populations represent one or more distinct species (Liu et al., 2021). Despite general agreement

that species are independently evolving metapopulation lineages (de Queiroz, 2005, 2011),

investigators differ in their interpretation of which species criteria (e.g., reciprocal monophyly,

ecological differentiation, phenetic diagnosability, reproductive incompatibility) are most salient

to diagnose species. Consequentially, research in the field can sometimes yield variable,

unstable, or uncertain results due to theoretical disagreements over species concepts and

philosophy (Frost and Kluge, 1994; de Queiroz, 2007; Hillis, 2019), methodological concerns

such as the appropriateness of genetic loci used in phylogenetic analysis (Chan et al., 2022),

geographic inclusiveness of sampling with respect to hybridization zones and areas of sympatry

(Myers et al., 2020; Marshall et al., 2021), and discordance among and within morphological and

molecular datasets (e.g., Toews and Brelsford, 2012; Denton et al., 2014). Discordant patterns

may be the result of phenomena such as incomplete lineage sorting (Wang et al., 2018), poor

resolution from sampled loci (Whitney et al., 2009), or reticulate evolutionary patterns produced

by hybridization and introgression (Ruane et al., 2014; Johnson et al., 2015; Myers et al., 2020;

Pyron et al., 2020; Burbrink et al., 2021; Chambers et al., 2023).

Due to its complex geography and tremendous biodiversity, Mexico provides many

opportunities to study the biogeography and systematics of herpetofauna—of which there are

>1,000 species of non-avian reptiles (52% endemic to the country) and >400 species of

amphibians (70% endemic to the country; Suazo-Ortuño, 2023). Studies of the phylogeography

of these taxa shed light not only upon their focal species, but also upon the shared evolutionary

histories of other organisms in the region (e.g., Bryson et al., 2011; Ríos-Muñoz and Navarro-

Sigüenza, 2012; Suárez-Atilano et al., 2014). Broadly speaking, the Mexican Transition Zone is

the area where the Nearctic and Neotropical realms grade into each other (Halffter, 1987;

Morrone, 2020), and it includes five distinct mountainous features: the Trans-Mexican Volcanic

Belt, Sierra Madre Occidental, Sierra Madre Oriental, Sierra Madre del Sur, and the Chiapan–

Guatemalan Highlands. These mountain ranges and their associated highlands form the interior

Central Mexican Highlands, which is sandwiched by coastal lowlands. The formation of these

mountainous features was complex, with orogeny mostly occurring or finishing during the

Miocene (Mastretta-Yanes et al., 2015). For this reason, many phylogeographic studies attribute

recent vicariant speciation (e.g., in the Pleistocene) to climatic changes and older events to

geological activity (Bryson et al., 2012; Mastretta-Yanes et al., 2015; Schramm et al., 2021).

Nonetheless, the four magmatic episodes that gave rise to the relatively young Trans-Mexican

Volcanic Belt did not complete until ~3 Mya in the Plio-Pleistocene (Ferrari et al., 2012).

In southern Mexico, the Isthmus of Tehuantepec separates the Sierra Madre del Sur from

the Chiapan–Guatemalan Highlands, connecting the Veracruzan and Pacific Lowlands

biogeographic provinces across a contiguous, lowland ~200 km east–west zone (Morrone, 2017).

Some geological and biological evidence supports the existence of a Pliocene–Pleistocene trans-

isthmian seaway (Durham et al., 1955; Sullivan et al., 2000; Mulcahy et al., 2006; Daza et al.,

2010; Hardy et al., 2010), which may have further isolated highland species, although this

remains contentious (Durham et al., 1955; Barrier et al., 1998; Campbell, 1999; Mulcahy et al.,

2006). Regardless, the Isthmus of Tehuantepec is a well-recognized vicariant barrier between a

diverse variety of both highland taxa including mammals, birds, and herpetofauna (Duellman,

1960; Sullivan et al., 1997; Arellano et al., 2005; Weir et al., 2008; Castoe et al., 2009; Zamudio-

Beltrán et al., 2020). Although not as well-documented, it is also a biogeographic break between

lowland taxa found in the tropical moist forest on the northern (Gulf of Mexico) side and tropical

dry forest and thornscrub on the southern (Pacific) side. For example, sister species of *Coleonyx*

geckos are found on either side, with the abrupt transition from the northern wet tropical forest

habitat of the Gulf to the southern dry tropical forests of the Pacific acting as an ecological

barrier to gene flow (Butler et al., 2023).

Here, we focus on uncovering the phylogeography of a lowland species with allopatric

populations found on the Pacific and Gulf coasts (but not continuously across the intervening

lowlands in the Isthmus of Tehuantepec)—the Marbled Toad (*Incilius marmoreus*). Originally

described as *Bufo marmoreus* Wiegmann, 1833, we apply the genus name *Incilius* Cope, 1863

following recommendations suggested by morphology and molecular data (Frost et al., 2006;

Frost et al., 2009; Mendelson et al., 2011). *Incilius* spp. are commonly referred to as

"Mesoamerican Toads", although some species extend beyond the traditional boundaries of

Mesoamerica—south into the Ecuador (e.g., *I. coniferus*) and north into the southern United

States (e.g., *I. nebulifer* and *I. alvarius*). *Incilius marmoreus* is endemic to Mexico and is

distributed predominately in xeric to subhumid and deciduous habitat along the Pacific Coast,

from the southern tip of Sonora to near Nachig, Chiapas. Interestingly, the type specimens of *I.*

marmoreus originate from the city and state of Veracruz, which borders the Gulf of Mexico, and

are thus rather disjunct from the remainder of the distribution (Fig. 1A). These Veracruzan

records appear to be primarily associated with palm savannah and scrub habitat (Rzedowski,

1994). Records of *I. marmoreus* from northern Hidalgo (Lemos-Espinal and Dixon, 2016) were

determined to instead represent *I. nebulifer* (JRM unpubl. data), and a single record in Durango

(Lemos-Espinal et al., 2019) remains unconfirmed and is separated by a distance of ~220

kilometers from the populations on the Pacific. From our examination of museum specimens (see

Methods), this species does not exist in the Balsas Depression, but specimens of the

morphologically similar *I. perplexus* (Taylor, 1943) that occurs there are often confused with *I.*

marmoreus (see following paragraph). Thus, verified records of *I. marmoreus* are known from

two allopatric regions: a broad expanse of the Pacific Coast and a small area in Veracruz. While

broad distributions of animal species along the subhumid Pacific Coast of Mexico are common,

such a distribution with an allopatric population in coastal Veracruz is not (an example is noted

in the Discussion). *Incilius marmoreus* appears to be abundant throughout much of its range and

is currently considered a species of Least Concern (IUCN SSC Amphibian Specialist Group,

2020), but modeling suggests that it might face a loss of distribution in the next ~60 years from

climate change (Garcia et al., 2014).

Incilius marmoreus is sexually dimorphic; as males mature, they lose their juvenile

marbled pattern and develop a solid dorsal coloration that ranges from greenish yellow to

yellow-gold, whilst females develop a brown coloration and retain their dorsal pattern (van der

Heiden, 2022). Along the Pacific Coast, the morphologically similar *I. perplexus* is also sexually

dichromatic, but it can be distinguished from *I. marmoreus*. In females, the dorsal pattern of *I.*

perplexus is sprawled and rounded whereas *I. perplexus* bears more triangular blotches. In males,

it is more difficult to differentiate between species (e.g., from specimens that have lost their

patterning). However, although the parotoid glands of *I. marmoreus* can be either triangular or

rounded, the parotoid glands of *I. perplexus* are always rounded and generally "pill-shaped".

Along the Pacific Coast, *I. marmoreus* may also distinguished from sympatric *I. mazatlanensis*

(Taylor, 1940) by the absence or strong reduction of parietal crests, which are present and well-

developed in *I. mazatlanensis*. In Veracruz, *I. marmoreus* may be distinguished from sympatric *I.*

valliceps (Wiegmann, 1833) in the same manner.

According to Frost (2024), three taxa have been previously synonymized with *Incilius*

marmoreus Wiegmann, 1833. *Bufo argillaceus* Cope, 1868 and *Bufo lateralis* Werner, 1894

were described along the Pacific Coast, with the type localities in Colima and Oaxaca,

respectively. However, type specimens corresponding to these taxa were deemed to be

conspecific with *B. [Incilius] marmoreus* by Günther (1901). Until now, this recommendation to

consider the allopatric (and morphologically similar) *I. marmoreus* in the Pacific and Gulf

lowlands as conspecific has not received further attention. In addition, *Bufo eiteli* Ahl, 1927

(with the type locality in Veracruz) was later synonymized with *B. [Incilius] marmoreus* by

Gorham (1974). Because previous work on *Incilius* spp. has shown that many allopatric montane

populations represent distinct species (reviewed by Mendelson et al., 2011), reexamination of the

two allopatric populations of *I. marmoreus* is warranted. Here, we adopt an integrative approach

(e.g., Dayrat, 2005) involving both morphological comparison of museum specimens and

phylogenetic and population genetic analyses of genome-wide single nucleotide polymorphisms

(SNPs) to test divergence under the general lineage species concept (de Queiroz, 1998, 2007)

and evaluate alternative phylogeographic hypotheses. We also comment on differences in

interspecific relationships from our results vs. those from prior phylogenetic studies (Mendelson

et al., 2011).

MATERIALS AND METHODS

Morphological data collection and analysis

We obtained EtOH-preserved specimens of *I. marmoreus* from the following museums: AMNH,

CAS, KU, LACM, MCZ, MVZ, TCWC, UCM, UMMZ, UTA, and UTEP (museum codes per

Sabaj et al., 2023). These included specimens collected from Veracruz (and deposited in the

AMNH and UTA collections) and from across the remainder of the geographic range of the

species. After verifying identifications, we had a total of n = 171 specimens of *I. marmoreus*. Of

the 126 males, 29 were from Veracruz; of the 45 females, 7 were from Veracruz. We confirmed

sex through examination of nuptial excrescences on the first finger (present in males, absent in

females) and corroborated these determinations using differences in dorsal color patterns (van

der Heiden, 2022).

Following Duellman (1970) and Mendelson (1994), we collected the following

measurements, rounded to the nearest 0.1mm, via a digital calipers: SVL (snout–vent length; the

distance from the rostral tip to the posterior end of the ischium), TIB (tibial length), FT (foot

length), HL (head length), HW (head width), EYE (diameter of the bony orbit), EN (eye–nostril

length), TYMP (tympanum diameter, across the major axis if elliptical), PARL (longest parotoid

measurement parallel to the sagittal plane), and PARW (longest parotoid measurement parallel to

the coronal plane). We analyzed these data in three ways. First, we used Welch's t-tests ($\alpha = .05$)

to evaluate whether the average SVL differed between animals from Veracruz and those from the

Pacific Coast, separating males and females into two separate analyses. Second, we log-

transformed the measurements and used a series of linear models to test for regional differences

in the averages of other traits and for differences in how these traits scale with SVL; to do this,

we again separated males and females into different analyses and fit models using the function

"ln()" in R v4.3.1 (R Core Team, 2023). We evaluated statistical significance ($\alpha = .05$;

Bonferonni adjusted $\alpha = .0056$) of interaction terms using Type-III sums of squares with function

"Anova()" in the package "car" v3.1-2 (Fox and Weisberg, 2019). If the interaction term was not

significant, we refitted the models without the interaction and evaluated statistical significance of

the main effect (i.e., region) using Type-II sums of squares as described above. Third, we

conducted principal component analyses (PCA) through singular value decomposition (Wall et

al., 2003) using the "prcomp()" function in R v4.3.1 (R Core Team, 2023). We separated males

and females into different analyses, included all log-transformed morphological measurements,

and plotted the first two axes using the package "ggplot2" v3.5.1 (Wickham, 2016), with

separate ellipses denoting 95% confidence intervals surrounding points from Veracruz and those

from all other locations.

2RAD library preparation and sequencing

To generate genome-wide SNP data, we used a reduced-representation genome sequencing

method. These methods, which include Restriction site Associated DNA sequencing (RADseq)

and its many variants, sequence a subset of orthologous loci across genomes of many individuals

and have become popular tools to investigate phylogenetic and population genetic questions in

non-model organisms like amphibians and reptiles (e.g., Pante et al., 2015; McCartney-Melstad

et al., 2018; Liu et al., 2020; Scott et al., 2020). Here, we used 2RAD (Bayona-Vásquez et al.,

2019), which is similar to dual-digest RADseq (ddRAD) (Peterson et al., 2012) but offers several

advantages such as better utility with low-concentration DNA samples and lower startup costs.

We prepared 2RAD libraries for 16 *Incilius marmoreus* (including 2 from Veracruz), 5 *I.*

perplexus, 1 *I. canaliferus*, 5 *I. mazatlanensis*, 1 *I. valliceps*, and 1 *Rhinella horribilis*. We

extracted DNA using a Qiagen DNeasy Blood and Tissue kit and quantified extracts using a

Qubit 4.0 Fluorometer. We then prepared 2RAD libraries (Bayona-Vásquez et al., 2019) with 50

ng of input DNA and using the enzymes XbaI and EcoRI-HF. We conducted individual PCRs,

pooled PCR products by volume, and size-selected this pool for 450–550 bp on a Pippin Prep.

We then pooled these libraries with others from unrelated projected for paired-end 150-bp

(PE150) sequencing on an Illumina NovaSeq 6000.

2RAD data assembly and analysis

We assembled demultiplexed data in ipyrad v0.9.94 (Eaton and Overcast, 2020), with the

following parameters: reads trimmed to 130 bp, a clustering threshold of 0.85, a minimum read

depth of 6 reads per locus, and a minimum of 4 samples per locus. We then used this assembly in

the following phylogenomic and population genomic analyses.

First, we estimated a maximum-likelihood phylogeny from an alignment of all 2RAD

loci using the program RAxML v8.2.12 (Stamatakis, 2014) and through the analysis toolkit in

ipyrad v0.9.94 (Eaton and Overcast, 2020). We used a GTRGAMMA nucleotide substitution

model and inferred 100 rapid bootstrap replicates followed by a full maximum-likelihood search.

We rooted this phylogeny with the outgroup *R. horribilis*, and we plotted all bootstrap replicates

in a DensiTree-like (Bouckaert, 2010) visualization using the package "phangorn" v2.11.1

(Schliep, 2011) in R v4.3.1 (R Core Team, 2023). Second, we used the species-tree inference

program tetrad v.0.9.14, which is based upon the SVDQuartets algorithm (Chifman and

Kubatko,' 2014), to infer a species tree from SNPs. We used our full assembly, all quartets, and

100 bootstrap replicates, and we again rooted with the outgroup and plotted all replicates. Third,

we estimated a distance-based phylogenetic network using the Neighbor-Net algorithm (Bryant

and Moulton 2004) as implemented in the package "phangorn" (Schliep, 2011; Schliep et al.,

2017) in R v4.3.1 (R Core Team, 2023). For this analysis, we included only samples from *I.*

marmoreus, *I. perplexus, I. mazatlanensis,* and *I. canaliferus* and constructed a pairwise

Hamming distance matrix from one random SNP per locus.

Fourth, we conducted a principal components analysis (PCA) of *I. marmoreus* samples

using the analysis toolkit in ipyrad v0.9.94 (Eaton and Overcast, 2020). We first filtered our

assembly to: 1) include only ingroup samples (i.e., *I. marmoreus*); 2) include only SNPs present

in at least 80% of samples; 3) include only SNPs with a minor allele frequency (MAF) ≥ 0.2; and

4) impute missing data with *k*-means clustering and $k = 6$. We plotted the first two PC axes in R

v4.3.1 (R Core Team, 2023). Fifth, we used the Bayesian clustering program Structure v2.3.4

(Pritchard et al., 2000) to evaluate population genetic structure, and we likewise conducted this

analysis through the analysis toolkit in ipyrad v0.9.94 (Eaton and Overcast, 2020). We first

filtered our assembly to: 1) include only ingroup samples (i.e., *I. marmoreus*); 2) include only

SNPs present in at least 80% of samples; 3) include only SNPs with a MAF ≥ 0.05; and 4)

include only one SNP per locus. We then conducted 10 replicate runs of 300,000 Markov chain

Monte Carlo steps (with 30,000 burn-in steps) for $k = 1$–7. We removed replicates that did not

converge (i.e., by setting max_var_multiple = 100) and visually examined a plot of ΔK (Evanno

et al., 2005) to assess the "optimal" number of clusters. With the aid of the CLUMPP (Jakobsson

and Rosenberg, 2007), we then plotted the average of replicates for two strongly supported

values of *k* overlayed on a map.

Finally, we used the program Bayesian Phylogenetics & Phylogeography (BPP; Rannala

and Yang, 2003; Yang, 2015) through the analysis toolkit in ipyrad v0.9.94 (Eaton and Overcast,

2020) to estimate divergence times and effective population sizes under a multi-species

coalescent model (i.e., the A00 analysis), modeling our analyses after Butler et al. (2023). We

used only samples of *Incilius marmoreus*, and we conducted these analyses for two alternative

scenarios: 1) one in which we modeled coalescence between Veracruz samples and all others;

and 2) one in which we modeled coalescence between Veracruz and the remainder of southern

samples (see Results), then coalescence between these groups and the northern samples. For each

scenario, we assigned an inverse-gamma prior (3, 0.002) for the population size parameter θ and

an inverse-gamma prior (2, 0.0015) for τ_0, the divergence time of the root. We conducted two

independent runs for each scenario, with each consisting of 120,000 generations (with 20,000

discarded as burn-in). We then used the mutation rate of 7.76×10^{-10} mutations per site per year

(from the Tibetan frog *Nanorana parkeri;* Sun et al., 2015) and a generation time of two years

(Oliveira et al., 2017) to convert our parameters into absolute estimates of time and effective

population size; these reflect the best available estimates for these parameters, but we

acknowledge that there remains uncertainty about both. We visually assessed Markov chain

Monte Carlo (MCMC) mixing and convergence in Tracer v1.7.2 (Rambaut et al., 2018).

RESULTS

Morphological analyses

Male *Incilius marmoreus* from Veracruz (mean $\pm$ SD = 48.9 $\pm$ 2.9 mm) were smaller than those

from the Pacific Coast (57.0 $\pm$ 4.2 mm) ($t = -11.75$, df = 66.82, $P < 0.001$). Likewise, females

from Veracruz (52.6 $\pm$ 5.2 mm) were smaller than those from the Pacific Coast (61.5 $\pm$ 4.9 mm)

($t = -4.20$, df = 8.06, $P = 0.003$). In males, we found a significant interaction between SVL and

region (i.e., Veracruz vs. Pacific Coast) for HW (Table 1). For male toads from Veracruz, we

found that HW was narrower at small SVLs but increased at a greater rate with SVL. We also

found a significant interaction between SVL and region for EYE; smaller toads from Veracruz

had larger eyes, and eye size increased at a more gradual rate with SVL. However, neither of

these differences were significant after a Bonferroni correction for multiple comparisons (Table

1). We also found significant differences in the intercept (i.e., mean values) for EN and TYMP;

both these differences were significant even after correction for multiple comparisons (Table 1).

Males from Veracruz had proportionately larger EN and smaller TYMP. In females, we found a

significant interaction between SVL and region (i.e., Veracruz vs. Pacific Coast) for PARL

(Table 2). For female toads from Veracruz, we found that PARL was shorter at small SVL but

increased at a greater rate with SVL. However, this difference was not significant after correction

for multiple comparisons. We also found a significant difference in intercept (i.e., mean values)

for TIB (Table 2); females from Veracruz had smaller TIB than those from elsewhere. This

difference was not significant after correction for multiple comparisons (Table 2).

In both PCAs of morphological traits, we found that the first principal component (PC)

axis mostly separated animals from Veracruz from those from all other locations, but 95%

confidence ellipses overlapped considerably (Fig. 2). For both males and females, SVL had the

highest loading for this axis, with animals from Veracruz being smaller than those from

elsewhere (Tables 3, 4). Other measurements with high loadings on this first axis were TIB, HL,

HW, and EN. In the case of males, the second PC axis emphasized variation in size and shape of

the parotoid glands, with strong positive loadings on PARL and PARW. The third PC axis

assigned strong positive loadings on EYE and PARW. In females, the second PC axis mostly

explained eye-related measurements, with strong positive loadings on EYE and EN. The third

PC axis emphasized parotoid variation, with strong positive loadings on PARL and PARW.

Phylogenomic and population genomic analyses

We obtained 1.8–39.0 million PE150 reads per sample (median = 14.6 million), and our final

assembly included > 306,000 SNPs across > 75,000 loci. Our maximum-likelihood phylogenetic

analysis recovered *I. marmoreus* as sister to *I. perplexus,* and these two taxa together were then

sister to *I. canaliferus* (RAxML bootstrap support [b.s.] = 100). Together, these three taxa were

sister to *I. mazatlanensis* + *I. valliceps* (b.s. = 100; Fig. 3A). We recovered all these interspecific

relationships with high bootstrap support, and the topologies of both the species-tree inference

(Fig. 3B) and distance-based phylogenetic network (Fig. 3C) were concordant. Within *I.*

marmoreus, each of these analyses suggested a northern Pacific-Coast clade consisting of

samples from Sinaloa, Nayarit, and Jalisco (b.s. = 100) and a southern Pacific-Coast clade

consisting of samples from Guerrero, Oaxaca, and Veracruz (b.s. = 100); we recovered the

sample from Colima as sister to the southern clade in the maximum-likelihood and species-tree

analyses, and we recovered it as intermediate between the northern and southern clades in the

phylogenetic network (Fig. 3). In all three analyses, samples from Veracruz were nested within

samples from Oaxaca (Fig. 3).

The results of the PCA on the molecular data were consistent with our phylogenetic

analyses. The first PC axis explained 31.1% of variation and separated the northernmost samples

(i.e., from Sinaloa, Nayarit, and Jalisco) from the more southern samples, with the single sample

from Colima being somewhat intermediate, but closer to the southern samples (Fig. 1B). The

second PC axis explained 12.4% of variation and differentiated samples within the southern

clade, with those from Veracruz and Guerrero widely separated and those from Oaxaca at

intermediate positions (Fig. 1B). The Structure results provided corroborating evidence for these

same broad patterns. We found highest support for $k = 2$ and $k = 4$. At $k = 2$, the two clusters

corresponded with the northern and southern Pacific-Coast clades described above, and the

sample from Colima showed intermediate assignment values (Fig. 4A). At $k = 4$, the samples

from Sinaloa were placed in a third cluster (i.e., distinct from others in the northern clade), and

the fourth cluster had highest assignment probabilities in the samples from Veracruz, with those

from Guerrero and Oaxaca showing increasing assignment probabilities to this cluster moving

from west to east (Fig. 4B).

In our coalescent analyses in BPP, we found relatively concordant results across our two

modeled scenarios. In the first scenario, we estimated a coalescence time of 0.49 Mya for

Veracruz ($N_e = 0.1$ M) and all other populations ($N_e = 1.2$ M) (Fig. 5A). In the second scenario,

we estimated a coalescence time of 0.46 Mya for Veracruz ($N_e = 0.1$ M) and the remainder of

southern Pacific-Coast populations ($N_e = 0.5$ M) and a coalescence time of 1.29 Mya between

these groups ($N_e = 0.3$ M) and northern Pacific-Coast samples ($N_e = 0.4$ M) (Fig. 5B).

DISCUSSION

Our investigation revealed that *I. marmoreus* displays substantial morphological and

genetic variability throughout its range. Populations in Veracruz have smaller SVL and some

allometric differences from those along the Pacific Coast, but analyses of genome-wide SNP data

suggest that they are phylogenetically nested within populations from Oaxaca—the most

geographically proximate sampling sites from across the Isthmus of Tehuantepec. We argue that

these data are consistent with recognition of a single species, and below, and we further discuss

each of these results and their implications for the phylogeography and systematics of the toads.

From gross visual inspection, we did not observe any geographic cline in the

morphological proportions or color pattern along the Pacific Coast, nor could we identify any

diagnostic trait for distinguishing Pacific toads from those in Veracruz. However, our

morphometric analyses suggest that *I. marmoreus* from Veracruz have a smaller average SVL.

Additionally, we found some evidence that male *I. marmoreus* from Veracruz exhibit head

widths that scale more steeply in relation to SVL, have eye diameters that scale more gradually

with SVL, have proportionately longer eye–nostril measurements, and have proportionately

smaller tympana compared to their Pacific counterparts. We also found some evidence that

female *I. marmoreus* from Veracruz exhibit parotoid lengths that scale more steeply in relation to

SVL and have proportionately shorter tibiae.

We acknowledge that these morphological analyses have several limitations. First, they

include only adult specimens. Our statistical analyses implicitly assume linear relationships

between SVL and other morphological measurements, but some traits may scale otherwise.

Second, statistical power for the female comparisons (e.g., only seven females from Veracruz)

may have been impaired since our sampling was rather male-biased, as is often the case for toad

collections (Mendelson 1994). Perhaps most importantly, we note that all but two (UTA 54845,

54846) measured toads from Veracruz were drawn from a single museum series collected from

the type locality of the city of Veracruz, reducing our confidence in the generalizability of the

differences we measured.

Both the maximum-likelihood phylogeny and species tree we inferred from differed in

species-level topology from the summary hypothesis tree presented in Mendelson et al. (2011).

We found strong support for a sister relationship between *Incilius marmoreus* and *I. perplexus*,

with *I. canaliferus* being in-turn sister to *I. marmoreus* + *I. perplexus*; in contrast, Mendelson et

al. (2011) hypothesized a sister relationship between *I. canaliferus* and *I. perplexus*, with *I.*

marmoreus being sister to *I. canaliferus* + *I. perplexus*. Our topology is more concordant with

morphological evidence, for *I. perplexus and I. marmoreus* resemble each other in gross

morphology and pattern whilst *I. canaliferus* is readily distinguishable from them (and in fact, all

other *Incilius* spp.) by its unusually long parotoid glands.

Within *I. marmoreus,* our genome-wide SNP data were consistent with isolation-by-

distance along the Pacific Coast, with phylogenetic analyses demonstrating a northern and

southern clade with high support and PCA and Structure analyses corroborating this primary axis

of phylogeographic variation. Perhaps unexpectedly, we discovered that samples from Veracruz

were nested within samples from Oaxaca in phylogenetic analyses (but clustered separately in

the PCA); thus, we found evidence that toads from the southern end of their Pacific distribution

(Guerrero and Oaxaca) are more closely related to toads from Veracruz than they are to toads

from contiguous distribution farther north along the Pacific Coast (Jalisco, Nayarit, Colima). The

estimated coalescent timing ~0.46 or 0.49 Mya between toads in Veracruz and those nearest

them on the Pacific Coast is considerably later than the hypothesized ~2 Mya Pliocene–

Pleistocene marine introgression into the Isthmus of Tehuantepec (Maldonado-Koerdell, 1964;

Sullivan et al., 2000) and also rules out anthropogenic transportation (Ardelean et al., 2020;

Becerra-Valdivia and Higham, 2020).

Nonetheless, exactly what series of events may explain the current allopatric distribution

of *I. marmoreus* remains uncertain. With an extensive distribution along the western coast of

Mexico and a smaller allopatric population in Veracruz, Mexico, the geographic distribution of

Incilius marmoreus is rather unusual, but not unprecedented among herpetofauna. For example,

the crotaline viper *Agkistrodon bilineatus* covers a somewhat similar distribution, with those in

Veracruz having been described as *A. bilineatus lemosespinali* from a single specimen (Smith

and Chiszar, 2001). However, the provenance of that specimen and validity of that taxon have

been questioned (Campbell and Lamar, 2004; Bryson and Mendoza-Quijano, 2007; Porras et al.,

2013). Drawing from Gloyd and Conant (1990), Bryson and Mendoza-Quijano (2007)

hypothesized that a narrow band of seasonally dry forest—running from northern Guerrero to

Veracruz—might have acted as a dispersal corridor for *Agkistrodon*. We find this to be an

unlikely explanation for the biogeography of *I. marmoreus* for two reasons. First, unlike other

groups with similar distributions (e.g., *Ctenosaura* iguanas; Zarza et al., 2019), these toads

appear not to currently live in the interior lowlands (i.e., the Balsas Depression) nearest this

hypothetical corridor. Second, our data showed that *I. marmoreus* in Veracruz were most closely

related to those in southern Oaxaca—not those from Guerrero.

Instead, we suggest an alternative hypothesis. We propose that *I. marmoreus* diversified

throughout the Pacific Coast of Mexico and then dispersed eastward into Veracruz via the

Isthmus of Tehuantepec. Afterwards, relatively recent isthmian climactic and vegetative changes,

resulting in a wet Gulf forest and dry Pacific forest, may have ecologically isolated the Veracruz

population with the development of unsuitable wet-forest habitat on the northern side of the

isthmus. Today, the eastern terminus of the Trans-Mexican Volcanic Belt may prevent *I.*

marmoreus from expanding its distribution further north along the Gulf coast, limiting its

northern extent to near the town of Palma Sola, Veracruz. Indeed, this eastern terminus is a

biogeographic barrier between congeners *I. valliceps* and *I. nebulifer* (Mulcahy et al., 2006) and

has been implicated in the diversification of many other taxa, such as several fish species

(Mateos, 2005; Huidobro et al., 2006).

We recommend the continued recognition of *Incilius marmoreus* as a single species

exhibiting geographic variation. Consequently, we concur with Günther (1901) that the names

Bufo argillaceus and *Bufo lateralis* represent junior synonyms of *Incilius marmoreus*. We were

able to further corroborate this synonym by examining photographs of the syntypes or holotypes

of *B. marmoreus, B. argillaceus, and B. lateralis.* Furthermore, we were able to verify the later

synonymy of *B. eiteli* with *B. marmoreus* by Gorham (1974) in the same manner. Although the

populations in Veracruz remain disjunct and are apparently dwarfed, they share a very recent

evolutionary history with populations on the Pacific Coast and are not diagnosable by any

discrete morphological diagnostic characters. Thus, under the general lineage concept of species,

we do not recognize the *I. marmoreus* in Veracruz as representing an independently evolving

metapopulational lineage, and hence not as a distinct species (de Queiroz, 1998; de Queiroz,

2007). Our study is not the first to recognize populations of distinctly diminutive animals as

conspecific with "normal-sized" populations. For example, Hoza et al. (2023) used genomic data

to argue against the recognition of dwarf *Phrynosoma hernandesi* that is geographically isolated

in the San Luis Valley, Colorado, as a distinct species. Likewise, along the Pacific Coast, gene

flow, lack of morphological distinguishability, and lack of evidence of ecological divergence do

not support the evolutionary distinctiveness of the northern and southern metapopulations.

Finally, we note that the basic natural history of *I. marmoreus* remains nebulous. For

example, *Incilius marmoreus* has been reported to deposit eggs singly (Blair, 1972). However,

this behavior is highly unusual amongst bufonids and otherwise unknown within *Incilius* (but see

image by Campuzano León, 2016). Furthermore, single egg-laying behavior has been wrongly

attributed to other bufonids, such as *Anaxyrus debilis* (Girard, 1854) and *Anaxyrus retiformis*

(Sanders and Smith, 1951). In the case of these two species, eggs are, in truth, laid in short and

fragile strings that break easily, consequently appearing as if they have been laid singly

(Ferguson and Lowe, 1969; Altig and McDiarmid, 2015). We encourage future researchers to

describe and document additional aspects of this species' life history, behavior, and ecology.

DATA ACCESSIBILITY

All raw sequence reads are available through the NCBI SRA (PRJNA1173256). The assembled

2RAD dataset, morphological data, and code for all analyses are available through Zenodo

(10.5281/zenodo.14173692).

ACKNOWLEDGMENTS

We thank Greg Pauly for facilitating tissue and specimen loans, Ana Motta, Carol Spencer,

David Kizirian, Emily Braker, Greg Schneider, Lee Fitzgerald, Neftali Camacho, Stevie

Kennedy-Gold, and Vicky Zhuang for facilitating specimen loans, and Daniel G. Mulcahy, Frank

Tillack, Gassner Georg, Kevin de Queiroz, Mark-Oliver Rödel, and Schweiger Silke for

providing photos of the syntypes and holotypes associated with *Incilius marmoreus* and its junior

synonyms *I. argillaceus, I. lateralis, I. eiteli.*

LITERATURE CITED

Ahl, E. 1927 "1926". Ueber neue oder seltene Froschlurche aus dem Zoologischen Museum

Berlin. Sitzungsberichte der Gesellschaft Naturforschender Freunde zu Berlin 1926:111–

117.

Aiello-Lammens, M. E., R. A. Boria, A. Radosavljevic, and B. Vilela. 2015. spThin: an R

package for spatial thinning of species occurrence records for use in ecological niche

models. Ecography 38:541–545.

Altig, R. I., and R. W. McDiarmid. 2015. Handbook of Larval Amphibians of the United States

and Canada. Comstock Publishing Associates, Ithaca, New York.

Arellano, E., F. X. González-Cozátl, and D. S. Rogers. 2005. Molecular systematics of Middle

American Harvest Mice *Reithrodontomys* (Muridae), estimated from mitochondrial

cytochrome b gene Sequences. Molecular Phylogenetics and Evolution 37:529–540.

Barrier, É., L. Velasquillo, M. Chávez, and R. Gaulon. 1998. Neotectonic evolution of the

Isthmus of Tehuantepec (southeastern Mexico). Tectonophysics 287:77–96.

Bayona-Vásquez N. J., T. C. Glenn, T. J. Kieran, T. W. Pierson, S. L. Hoffberg, P. A. Scott,

K. E. Bentley, J. W. Finger, S. Louha, N. Troendle, P. Diaz-Jaimes, R. Mauricio,

and B. C. Faircloth. 2019. Adapterama III: Quadruple-indexed, double/triple-enzyme

RADseq libraries (2RAD/3RAD). PeerJ 7:e7724.

Becerra-Valdivia, L., and T. Higham. 2020. The timing and effect of the earliest human

arrivals in North America. Nature 584:93–97.

Blair, W.F. 1972. *Bufo* of North and Central America, p. 93–101. *In*: Evolution in the Genus

Bufo. W. F. Blair (ed.). University of Texas Press, Austin, Texas.

Bouckaert, R. R. 2010. DensiTree: making sense of sets of phylogenetic trees. Bioinformatics

26:1372–1373.

Bryson, Jr. R. W., and F. Mendoza-Quijano. 2007. Cantils of Hidalgo and Veracruz, Mexico,

with comments on the validity of *Agkistrodon bilineatus lemosespinali*. Journal of

Herpetology 41:536–539.

Bryson, Jr., R. W., R.W. Murphy, A. Lathrop, and D. Lazcano-Villareal. 2011. Evolutionary

drivers of phylogeographical diversity in the highlands of Mexico: a case study of the

Crotalus triseriatus species group of montane rattlesnakes. Journal of Biogeography

38:697–710.

Bryson, Jr., R. W., U.O. García-Vázquez, and B.R. Riddle. 2012. Diversification in the

Mexican horned lizard *Phrynosoma orbiculare* across a dynamic landscape. Molecular

Phylogenetics and Evolution 62:87–96.

Burbrink, F. T., G. M. Gehara, A. D. McKelvy, and E. A. Myers. 2021. Resolving spatial

complexities of hybridization in the context of the gray zone of speciation in North

American ratsnakes (*Pantherophis obsoletus* complex), Evolution 75:260–277

Butler, B. O., L. L. Smith, and O. Flores-Villela. 2023 Phylogeography and taxonomy of

Coleonyx elegans Gray 1845 (Squamata: Eublepharidae) in Mesoamerica: the Isthmus of

Tehuantepec as an environmental barrier. Molecular Phylogenetics and Evolution

178:107632

Campbell, J. A. 1999. Distribution patterns of amphibians in Middle America, p. 111–209. *In*:

Distribution Patterns of Amphibians: A Global Perspective. W.W. Duellman (ed.). John

Hopkins University Press, Baltimore, Maryland.

Campbell, J. A., and W. L. Lamar. 2004. Venomous reptiles of the Western Hemisphere.

Cornell University Press, Ithaca, New York.

Campuzano León, F. 2016. https://www.inaturalist.org/observations/12028134 (accessed 19

August 2024).

Castoe, T. A., J. M. Daza, E. N. Smith, M. M. Sasa, U. Kuch, J. A. Campbell, P. T.

Chippindale, and C. L. Parkinson. 2009. Comparative phylogeography of pitvipers

suggests a consensus of ancient Middle American highland biogeography. Journal of

Biogeography 36:88–103.

Chambers, E. A., T. L. Marshall, and D. M. Hillis. 2023. The importance of contact zones for

distinguishing interspecific from intraspecific geographic variation. Systematic Biology

72:357–371.

Chan, K. O., S. T. Hertwig, D. N. Neokleous, J. M. Flury, and R. M. Brown. 2022. Widely

used, short 16S rRNA mitochondrial gene fragments yield poor and erratic results in

phylogenetic estimation and species delimitation of amphibians. BMC Ecology and

Evolution 22:1–9.

Chifman, J., and L. Kubatko. 2014. Quartet inference from SNP data under the coalescent

model. Bioinformatics 30:3317–3324.

Dayrat, B. 2005. Towards integrative taxonomy. Biological Journal of the Linnean Society

85:407–417.

Daza, J.M., T. A. Castoe, and C. L. Parkinson. 2010. Using regional comparative

phylogeographic data from snake lineages to infer historical processes in Middle

America. Ecography 33:343–354.

de Queiroz, K. 1998. The general lineage concept of species, species criteria, and the process of

speciation: a conceptual unification and terminological recommendations, p. 57–76. *In*:

Endless Forms: Species and Speciation. D. J Howard, and S. H. Berlocher (eds.), Oxford

Academic, New York, New York.

de Queiroz, K. 2005. A unified concept of species and its consequences for the future of

taxonomy. Proceedings of the California Academy of Sciences 56:196–215.

de Queiroz, K. 2007. Species concepts and species delimitation. Systematic Biology 56:879–

886.

de Queiroz, K. 2011. Branches in the lines of descent: Charles Darwin and the evolution of the

species concept. Biological Journal of the Linnean Society 103:19–35.

Denton, R. D., L. J. Kenyon, K. R. Greenwald, and H. L. Gibbs. 2014. Evolutionary basis of

mitonuclear discordance between sister species of mole salamanders (*Ambystoma* sp.).

Molecular Ecology 23:2811–2824.

Duellman, W. E. 1960. A distributional study of the amphibians of the Isthmus of Tehuantepec,

Mexico. University of Kansas Publications, Museum of Natural History 13:19–72.

Duellman, W. E. 1970. The hylid frogs of Middle America. Monograph of the Museum of

Natural History, the University of Kansas 1:1–753.

Durham, J. W., A. R. V. Arellano, and J. H. Peck. 1955. Evidence for no Cenozoic Isthmus of

Tehuantepec seaways. Geological Society of America Bulletin 66:977–992.

Eaton, D.A., and I. Overcast. 2020. ipyrad: interactive assembly and analysis of RADseq

datasets. Bioinformatics 36:2592–2594.

Evanno, G., S. Regnaut, and J. Goudet. 2005. Detecting the number of clusters of individuals

using the software STRUCTURE: a simulation study. Molecular Ecology 14:2611–2620.

Ferguson, J. H., and C. H. Lowe, Jr. 1969. Evolutionary relationships in the *Bufo punctatus*

group. American Midland Naturalist 81:435–466.

Ferrari, L., T. Orozco-Esquivel, V. Manea, and M. Manea. 2012. The dynamic history of the

Trans-Mexican Volcanic Belt and the Mexico subduction zone. Tectonophysics 522:122–

149.

Fox, J., and S. Weisberg. 2019. An R Companion to Applied Regression. Third edition. Sage,

Thousand Oaks, California.

Frost, D. R. 2024. Amphibian Species of the World: an Online Reference. Version 6.2.

https://amphibiansoftheworld.amnh.org/index.php (accessed 16 August 2024). American

Museum of Natural History, New York, USA.

Frost, D. R., and A. G. Kluge. 1994. A consideration of epistemology in systematic biology,

with special reference to species. Cladistics 10:259–294.

Frost, D. R., J. R. Mendelson III, and J. Pramuk. 2009. Further notes on the nomenclature of

Middle American Toads (Bufonidae). Copeia 2009:418.

Frost, D. R., T. Grant, J. Faivovich, R. H. Bain, A. Haas, C. F. B. Haddad, R. O. de Sá, A.

Channing, M. Wilkinson, S. C. Donnellan, C. J. Raxworthy, J. A. Campbell, B. L.

Blotto, P. E. Moler . . . W. C. Wheeler. 2006. The amphibian tree of life. Bulletin of the

American Museum of Natural History 297:1–370.

Garcia, R. A., M. Cabeza, C. Rahbek, and M. B. Araújo. 2014. Multiple dimensions of

climate change and their implications for biodiversity. Science 344:1247579.

GBIF. 2024. GBIF Occurrence Download https://doi.org/10.15468/dl.5j329g (accessed 16

August 2024).

Girard, C. 1854. A list of North American bufonids, with diagnoses of new species.

Proceedings of the Academy of Natural Sciences of Philadelphia 7:86–88.

Gloyd, H. K., and R. Conant. 1990. Snakes of the *Agkistrodon* complex—a monographic

review. Contributions to Herpetology No. 6. Society for the Study of Amphibians and

Reptiles, Oxford, Ohio.

Gorham, S. W. 1974. Checklist of World Amphibians Up to January 1, 1970. New Brunswick

Museum, Saint John, Canada.

Günther, A. C. L. G. 1901. Reptilia and Batrachia, p. 237–252. *In*: Biologia Centrali

Americana. O. Salvin, and F. D. Godman (eds.). London, R. H. Porter and Dulau & Co.,

London, England.

Halffter, G. 1987. Biogeography of the montane entomofauna of Mexico and Central America.

Annual Review of Entomology 32:95–114.

Hardy D. K., F. X. González-Cózatl, E. Arellano, and D. S. Rogers. 2013. Molecular

phylogenetics and phylogeographic structure of Sumichrast's harvest mouse

(*Reithrodontomys sumichrasti*: Cricetidae) based on mitochondrial and nuclear DNA

sequences. Molecular Phylogenetics and Evolution 68:282–292.

Hillis, D. 2019. Species delimitation in herpetology. Journal of Herpetology 53:3–12.

Hoza, J., H. R. Davis, and A. D. Leaché. 2023. Genomic data do not support the species status

of the San Luis Valley Short-Horned Lizard (*Phrynosoma diminutum*). Ichthyology &

Herpetology 11:390–396.

Huidobro, L., J. J. Morrone, J. L. Villalobos, and F. Álvarez. 2006. Distributional patterns of

freshwater taxa (fishes, crustaceans and plants) from the Mexican Transition Zone.

Journal of Biogeography 33:731–741.

IUCN SSC Amphibian Specialist Group. 2020. *Incilius marmoreus*. IUCN Red List of

Threatened Species 2020.

Jakobsson, M., and N. A. Rosenberg. 2007. CLUMPP: a cluster matching and permutation

program for dealing with label switching and multimodality in analysis of population

structure. Bioinformatics 23:1801–1806.

Johnson, B. B., T. A. White, C. A. Phillips, and K. R. Zamudio. 2015. Asymmetric

introgression in a Spotted Salamander hybrid zone. Journal of Heredity 106:608–617

Lemos-Espinal, J. A., and J. R. Dixon. 2016. Anfibios y Reptiles de Hidalgo / Amphibians &

Reptiles of Hidalgo. CONABIO, Ciudad de México, Distrito Federal, México.

Lemos-Espinal, J. A., G. R. Smith, and R. Valdes-Lares. 2019. Amphibians and Reptiles of

Durango. ECO Herpetological Publishing & Distribution, Rodeo, New Mexico.

Liu, J-X., M-Y. Zhou, G-Q. Yang, Y-X. Zhang, P-F. Ma, C. Guo, M. S. Vorontsova, and D-

Z. Li. 2020. ddRAD analyses reveal a credible phylogenetic relationship of the four main

genera of *Bambusa-Dendrocalamus-Gigantochloa* complex (Poaceae: Bambusoideae).

Molecular Phylogenetics and Evolution 146:106758.

Liu, S., M. Hou, Y-H. Lwin, Q. Wang, and D. Rao. 2021. A new species of *Gonyosoma*

Wagler, 1828 (Serpentes, Colubridae), previously confused with *G. prasinum* (Blyth,

1854). Evolutionary Systematics 5:129–139.

Maldonado-Koerdell, M. 1964. Geohistory and paleogeography of Middle America, p. 3–32.

In: Handbook of Middle American Indians, vol. 1. R.C. West (ed.). University of Texas

Press, Austin, Texas.

Marshall T. L., E. A. Chambers, M. V. Matz, and D. M. Hillis. 2021. How mitonuclear

discordance and geographic variation have confounded species boundaries in a widely

studied snake. Molecular Phylogenetics and Evolution 162:107194.

Mastretta-Yanes, A., A. Moreno-Letelier, D. Piñero, T. H. Jorgensen, and B. C. Emerson.

2015. Biodiversity in the Mexican highlands and the interaction of geology, geography

and climate within the Trans-Mexican Volcanic Belt. Journal of Biogeography 42:1586–

1600.

Mateos, M. 2005. Comparative phylogeography of livebearing fishes in the genera *Poeciliopsis*

and *Poecilia* (Poeciliidae: Cyprinodontiformes) in central Mexico. Journal of

Biogeography 32:775–780.

Mayr, E. 1982. The Growth of Biological Thought: Diversity, Evolution, and Inheritance.

Belknap Press of Harvard University Press, Cambridge, Massachusetts.

McCartney-Melstad, E., M. Gidiş, and H. B. Shaffer. 2018. Population genomic data reveal

extreme geographic subdivision and novel conservation actions for the declining Foothill

Yellow-legged Frog. *Heredity* 121:112–125.

Mendelson, J. R., III. 1994. A new species of toad (Anura: Bufonidae) from the lowlands of

eastern Guatemala. Occasional Papers of the Museum of Natural History, University of

Kansas 166:1–21.

Mendelson, J. R., III, D. G. Mulcahy, T. S. Williams, and J. W. Sites. 2011. A phylogeny and

evolutionary natural history of Mesoamerican toads (Anura: Bufonidae: *Incilius*) based

on morphology, life history, and molecular data. Zootaxa 3138:1–34.

Morrone, J. J., T. Escalante, and G. Rodríguez-Tapia. 2017. Mexican biogeographic

provinces: map and shapefiles. Zootaxa 4277:277–279.

Morrone, J.J. 2020. Biogeographic regionalization of the Mexican Transition Zone, p. 103. *In*:

The Mexican Transition Zone. Springer, Cham, Switzerland.

Mulcahy, D. G., B. H. Morrill, and J.R. Mendelson, III. 2006, Historical biogeography of

lowland species of toads (*Bufo*) across the Trans-Mexican Neovolcanic Belt and the

Isthmus of Tehuantepec. Journal of Biogeography 33:1889–1904.

Myers, E. A., A. D. McKelvy, and F. T. Burbrink. 2020. Biogeographic barriers, Pleistocene
refugia, and climatic gradients in the southeastern Nearctic drive diversification in
cornsnakes (*Pantherophis guttatus* complex). Molecular Ecology 29:797–811.

Oliveira, B. F., V. A. São-Pedro, G. Santos-Barrera, C. Penone, and G. C. Costa. 2017.
AmphiBIO, a global database for amphibian ecological traits. Scientific Data 4:1–7.

Pante, E., J. Abdelkrim, A. Viricel, D. Gey, S. C. France, M. C. Boisselier, and S. Samadi.
2015. Use of RAD sequencing for delimiting species. Heredity 114:450–459.

Peterson, B. K., J. N. Weber, E. H. Kay, H. S. Fisher, and H. E. Hoekstra. 2012. Double
digest RADseq: an inexpensive method for de novo SNP discovery and genotyping in
model and non-model species. PLOS ONE 7:e37135.

Porras, L. W., L. D. Wilson, G. W. Schuett, and R. S. Reiserer. 2013. A taxonomic
reevaluation and conservation assessment of the Common Cantil, *Agkistrodon bilineatus*
(Squamata: Viperidae): a race against time. Amphibian and Reptile Conservation 7:48–
73.

Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of population structure using
multilocus genotype data. Genetics 155:945–959.

Pyron R. A., K. A. O'Connell, E. M. Lemmon, A. R. Lemmon, and D. A. Beamer. 2020.
Phylogenomic data reveal reticulation and incongruence among mitochondrial candidate
species in Dusky Salamanders (*Desmognathus*). Molecular Phylogenetics and Evolution
146:106751.

Rambaut, A., A. J. Drummond, D. Xie, G. Baele, and M. A. Suchard. 2018. Posterior
summarization in Bayesian phylogenetics Using Tracer 1.7. Systematic Biology 67:901–
904.

Rannala, B., and Z. Yang. 2003. Bayes estimation of species divergence times and ancestral

population sizes using DNA sequences from multiple loci. Genetics 164:1645–1656.

Ríos-Muñoz, C.A., and A.G Navarro-Sigüenza. 2012. Patterns of species richness and

biogeographic regionalization of the avifaunas of the seasonally dry tropical forest in

Mesoamerica. Studies on Neotropical Fauna and Environment 47:171–182.

Ruane, S., R. W. Bryson, R. A. Pyron, and F. T. Burbrink. 2014. Coalescent species

delimitation in Milksnakes (Genus *Lampropeltis*) and impacts on phylogenetic

comparative analyses. Systematic Biology 63:231–250.

Rzedowski, J. 1994. Vegetación de México. Limusa Noriega Editores, Ciudad de México,

Distrito Federal, México.

Sabaj, M. H. 2023. Codes for Natural History Collections in Ichthyology and Herpetology

(online supplement). Version 9.5 (10 Nov 2023). Electronically accessible at

https://asih.org, American Society of Ichthyologists and Herpetologists, Washington, DC.

Sanders, O., and H. M. Smith. 1951. Geographic variation in toads of the *debilis* group of *Bufo*.

Field and Laboratory 19:141–160.

Schliep, K. P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics 27:592–593.

Schliep, K., A. J. Potts, D. A. Morrison, and G. W. Grimm. 2017. Intertwining phylogenetic

trees and networks. Methods in Ecology and Evolution 8:1212–1220.

Schramm, F. D., A. Valdez-Mondragón, and L. Prendini. 2021. Volcanism and palaeoclimate

change drive diversification of the world's largest whip spider (*Amblypygi*). Molecular

Ecology 30:2872–2890.

Scott, P. A., L. J. Allison, K. J. Field, R. C. Averill-Murray, and H. B. Shaffer. 2020.

Individual heterozygosity predicts translocation success in threatened desert tortoises.

Science 370:1086–1089.

Smith, H. M., and D. Chiszar. 2001. A new subspecies of Cantil (*Agkistrodon bilineatus*) from

central Veracruz, Mexico (Reptilia: Serpentes). Bulletin of the Maryland Herpetological

Society 37:130–136.

Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics 30:1312–1313.

Suárez-Atilano, M., F. Burbrink, and E. Vázquez-Domínguez. 2014. Phylogeographical

structure within *Boa constrictor imperator* across the lowlands and mountains of Central

America and Mexico. Journal of Biogeography 41:2371–2384.

Suazo-Ortuño, I., A. Ramírez-Bautista, and J. Alvarado-Díaz. 2023. Amphibians and reptiles

of Mexico: diversity and conservation, p. 105–125. *In*: Mexican Fauna in the

Anthropocene. R. W. Jones, C. P. Ornelas-García, R. Pineda-López, and F. Álvarez.

(eds.). Springer, Cham, Switzerland.

Sullivan, J., E. Arellano, and D. S. Rogers. 2000. Comparative phylogeography of

Mesoamerican highland rodents: concerted versus independent response to past climatic

fluctuations. The American Naturalist 155:755–768.

Sullivan, J., J. A. Markert, and C. W. Kilpatrick. 1997. Phylogeography and molecular

systematics of the *Peromyscus aztecus* species group (Rodentia: Muridae) inferred using

parsimony and likelihood. Systematic Biology 46:426–40.

Sun, Y-B., Z-J. Xiong, X-Y. Xiang, S-P. Liu, W-W. Zhou, X-L. Tu, L. Zhong, L. Wang, D-

D. Wu, B-L. Zhang, C-L. Zhu, M-M. Yang, H-M. Chen, F. Li . . . Y-P. Zhang. 2015.

Whole-genome sequence of the Tibetan frog *Nanorana parkeri* and the comparative

evolution of tetrapod genomes. Proceedings of the National Academy of Sciences

112:1257–1262.

Taylor, E. H. 1940 "1939". Herpetological miscellany No. I. University of Kansas Science

Bulletin 26:489–571.

Toews, D. P. L., and A. Brelsford. 2012. The biogeography of mitochondrial and nuclear

discordance in animals. Molecular Ecology 21:3907–3930.

van der Heiden, A. 2022. *Incilius marmoreus* (Marbled Toad). Sexual dichromatism and

reproductive behavior. Herpetological Review 53:468–470.

Wall, M. E., A. Rechtsteiner, and L. M. Rocha. 2003. Singular value decomposition and

principal component analysis, p. 91–109. *In*: A Practical Approach to Microarray Data

Analysis. D. P. Berrar, W. Dubitzky, M. Granzow. (eds.). Springer, Boston,

Massachusetts.

Wang, K., J. A. Lenstra, L. Liu, Q. Hu, T. Ma, Q. Qiu, and J. Liu. 2018. Incomplete lineage

sorting rather than hybridization explains the inconsistent phylogeny of the wisent.

Communications Biology 1:169.

Weir, J. J., E. Bermingham, M. J. Miller, J. Klicka, and M. A. González. 2008.

Phylogeogoraphy of a morphologically diverse Neotropical montane species, the

Common Bush-Tanager (*Chlorospingus ophthalmicus*). Molecular Phylogenetics and

Evolution 47:650–64.

Whitney, K. D., J. R. Ahern, L. G. Campbell, L. P. Albert, and M. S. King. 2010. Patterns of

hybridization in plants. Perspectives in Plant Ecology, Evolution and Systematics

12:175–182.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York,

New York.

Yang, Z. 2015. The BPP program for species tree estimation and species delimitation. Current

Zoology 61:854–865.

Zamudio-Beltrán, L.E., Y. Licona-Vera, B. E. Hernández-Baños, J. Klicka, and J. F.

Ornelas. 2020. Phylogeography of the widespread white-eared hummingbird (*Hylocharis*

leucotis): pre-glacial expansion and genetic differentiation of populations separated by

the Isthmus of Tehuantepec. Biological Journal of the Linnean Society 130:247–267.

Zarza, E., V. H. Reynoso, C. M. Faria, and B. C. Emerson. 2019. Introgressive hybridization

in a Spiny-Tailed Iguana, *Ctenosaura pectinata*, and its implications for taxonomy and

conservation. PeerJ, 7:e6744.

Table 1. Results from a series of linear models for morphological measurements of male

Marbled Toads (*Incilius marmoreus*). We modeled each morphological variable as a function of

snout–vent length (SVL) and its interaction with region (i.e., Veracruz vs. Pacific Coast). We

used an analysis-of-variance (ANOVA) with Type III Sums of Squares to evaluate statistical

significance. If the interaction term was significant, we do not report the main effect. If the

interaction term was not significant, we refit the model without the interaction and used a Type-

II sums of squares to evaluate the main effect of region. Here, we indicate statistical significance

($\alpha = 0.05$) with an asterisk, and statistical significance after a Bonferroni correction (adjusted $\alpha =$

0.0056) with two asterisks. All values rounded to three decimal places. TIB = tibial length; FT =

foot length; HL = head length; HW = head width; EYE = eye diameter; EN = eye-nostril length;

TYMP = tympanum diameter; PARL = parotoid length; PARW = parotoid width.

	SS (Type III)	SS (Type II)	F-value	Pr(>F)
TIB slopes	0.002		1.285	0.259
TIB intercepts		0.001	0.523	0.471
FT slopes	0.004		0.403	0.527
FT intercepts		0.000	0.000	0.988
HL slopes	0.003		1.806	0.182
HL intercepts		0.000	0.007	0.935
HW slopes	0.013		4.590	0.034*
HW intercepts	—	—	—	—
EYE slopes	0.014		4.108	0.045*
EYE intercepts	—	—	—	—
EN slopes	0.005		1.936	0.167
EN intercepts		0.043	18.242	0.000**
TYMP slopes	0.023		2.124	0.148
TYMP intercepts		0.095	8.559	0.004**
PARL slopes	0.049		1.742	0.189
PARL intercepts		0.066	2.326	0.130
PARW slopes	0.000		0.000	0.991
PARW intercepts		0.005	0.208	0.650

Table 2. Results from a series of linear models for morphological measurements of female

Marbled Toads (*Incilius marmoreus*). We modeled each morphological variable as a function of

snout–vent length (SVL) and its interaction with region (i.e., Veracruz vs. Pacific Coast). We

used an analysis-of-variance (ANOVA) with Type III Sums of Squares to evaluate statistical

significance. If the interaction term was significant, we do not report the main effect. If the

interaction term was not significant, we refit the model without the interaction and used a Type II

Sums of Squares to evaluate the main effect of region. Here, we indicate statistical significance

($\alpha = 0.05$) with an asterisk, and statistical significance after a Bonferroni correction (adjusted $\alpha =$

0.0056) with two asterisks. All values rounded to three decimal places. Abbreviations are

provided in the text and Table 1 caption.

	SS (Type III)	SS (Type II)	F-value	Pr(>F)
TIB slopes	0.002		1.082	0.304
TIB intercepts		0.007	4.097	0.049*
FT slopes	0.018		3.896	0.056
FT intercepts		0.010	1.880	0.178
HL slopes	0.008		2.703	0.108
HL intercepts		0.000	0.017	0.897
HW slopes	0.010		4.070	0.050
HW intercepts		0.000	0.020	0.889
EYE slopes	0.000		0.001	0.981
EYE intercepts		0.015	3.880	0.055
EN slopes	0.003		0.956	0.334
EN intercepts		0.005	2.021	0.163
TYMP slopes	0.014		1.074	0.306
TYMP intercepts		0.030	2.252	0.141
PARL slopes	0.113		5.439	0.025*
PARL intercepts	—	—	—	—
PARW slopes	0.007		0.315	0.578
PARW intercepts		0.003	0.138	0.712

739 **Table 3.** Results from a principal components analysis (PCA) of morphological measurements

740 from male Marbled Toads (*Incilius marmoreus*). Here, we report results from all principal

741 components (PC) axes up to 80% cumulative variance explained. Abbreviations are provided in

742 the text and Table 1 caption.

	PC1	PC2	PC3
Proportion of Variance	0.5697	0.1493	0.0874
Cumulative Variance	0.5697	0.7190	0.8064
Eigenvalue	5.6974	1.4928	0.8741
SVL	-0.3973	-0.0113	-0.0953
TIB	-0.3724	0.0647	-0.2269
FT	-0.3231	0.0287	-0.1367
HL	-0.3782	-0.0956	-0.0276
HW	-0.3605	0.0136	-0.1008
EYE	-0.2475	-0.4432	0.5608
EN	-0.3354	-0.3412	0.3037
TYMP	-0.2999	0.1012	-0.4738
PARL	-0.2100	0.5434	0.2451
PARW	-0.1354	0.6059	0.4674

744 **Table 4.** Results from a principal components analysis (PCA) of morphological measurements

745 from female Marbled Toads (*Incilius marmoreus*). Here, we report results from all principal

746 components (PC) axes up to 80% cumulative variance explained. Abbreviations are provided in

747 the text and Table 1 caption.

	PC1	PC2	PC3
Proportion of Variance	0.5917	0.1219	0.0865
Cumulative Variance	0.5917	0.7135	0.8001
Eigenvalue	5.9168	1.2186	0.8652
SVL	0.3714	-0.0923	-0.1983
TIB	0.3613	-0.2457	-0.1049
FT	0.3266	-0.1941	0.0317
HL	0.3586	-0.0241	-0.2519
HW	0.3647	-0.0075	-0.2648
EYE	0.2125	0.6961	-0.0612
EN	0.3243	0.4577	-0.1033
TYMP	0.2490	-0.4209	0.0359
PARL	0.3042	-0.0782	0.4629
PARW	0.2416	0.1245	0.7650

Figure 1. (A) A map of Mexico showing the distribution of the Marbled Toad (*Incilius marmoreus*). Small white dots represent

research-grade and unobscured iNaturalist hosted occurrence points downloaded from GBIF; we used the R package spThin version

v0.2.0 (Aiello-Lammens et al., 2015) to spatially thin records at a threshold of 25 km. Larger solid symbols represent locations from

which we generated 2RAD sequencing data. Identifications were verified through dorsal patterning and crest morphology. Shading

indicates elevation. (B) The first two principal components (PC) axes from a principal components analysis (PCA) of 2RAD data.

Figure 2. (A) The first two principal components (PC) axes from a principal components analysis (PCA) of morphological data

collected from male Marbled Toads (*Incilius marmoreus*) from across their geographic distribution. (B) The first two PC axes from an

identical analysis of female *I. marmoreus*. In both, point colors and shapes reflect the collection location, and ellipses show 95%

confidence intervals around two groups: toads from Veracruz and toads from the Pacific Coast.

Figure 3. (A) DensiTree-like plots of RAxML rapid bootstrap replicates from a maximum-likelihood analysis of loci from 2RAD

data, showing relationships within the Marbled Toad (*Incilius marmoreus*) and between relatives; (B) a similar plot showing bootstrap

replicates from a tetrad species-tree analysis of SNPs from the same dataset; (C) a Neighbor-Net phylogenetic network inferred from

one random SNP per locus from the same dataset, but with a subset of taxa. In all, point colors and shapes reflect the collection

location within Mexico.

Figure 4. Genetic clustering results for the Marbled Toad (*Incilius marmoreus*), inferred using SNPs from 2RAD data and with the

program Structure. Plots show results from (A) $k = 2$ and (B) $k = 4$. In both plots, some pie charts represent averages of assignments

from more than one individual sample from the same collection site. Shading indicates elevation.

Figure 5. Results from BPP coalescent analyses with two alternative demographic scenarios: (A) one in which we estimated a date for the coalescence between Veracruz samples and all others; and (B) one in which we estimated a date for the coalescence of Veracruz and the remainder of southern samples (see Fig. 4) and a second date for the coalescence between these two and the northern samples. Within each figure, bar widths are proportional to estimated effective population sizes (N_e), and bar heights are proportional to estimated coalescence dates.

APPENDIX 1 [Specimens Examined]

Incilius marmoreus.— Mexico: Chiapas: 2.7 mi [4.3 km] W Chiapa de Corzo (AMNH 173339–40), 18 mi [29.0 km] N Arriaga

(AMNH 173341–42); 29.5 mi [47.5 km] NE Tapanatepec, Oaxaca (AMNH 173343–51); 21.6 mi [34.8 km] N Arriaga (AMNH

173352–54); 13.3 mi. [21.4 km] S. Tuxtla Gutierrez, along Chis. 28 (UCM 48536–41); 4.8 mi [7.7 km] S. Jct. Mexico 190 and road to

La Concordia, along road to La Concordia (UCM 48544–50); 10.6 mi [17.1 km] S. Jct. Mexico 190 and road to La Concordia, along

road to La Concordia (UCM 48551–56); 15.1 mi [24.3 km] S. Jct. Mexico 190 and road to La Concordia, along road to La Concordia

(UCM 48557); 0.4 mi [0.6 km] NNE (by road on Mexico Highway 35) of Arriaga (UTEP 4714); 4.3 mi [7.0 km] N of Arriaga (UTEP

5843); 1.0 mi [1.6 km] N of Arriaga (UTEP 5927).

Mexico: Colima: 7.5 mi [12.1 km] SW Colima (LACM 36957, 59–70).

Mexico: Guerrero: 3.5 mi [5.6 km] S San Andres de la Cruz (KU 86676, 86679, 86681–82, 86684, 86688–89); Hwy 200 btwn Pie de

la Ruesta turnoff and 15 mi [24 km] W (LACM 128416); .6 mi [1 km] NW Puerto Marquez above Hwy to airport (LACM 129899);

24 mi [39 km] (by Mex Hwy 200) WNW Coyuca de Benitez; 1.6 mi [2.6 km] by rd SE Alcholoa (LACM 129910–11); Acapulco

(MCZ 1436); Chilpancingo (MCZ 17756).

Mexico: Jalisco: 5 mi [8 km] E Melaque on Hwy 80 (KU 95440–42, 45–46); Village of Casimiro Castillo on Hwy 30 (LACM 87827);

6 mi [10 km] NE of Chamela, La Huerta (UTEP 5860–61).

Mexico: Michoacan: 23.1 mi [37.2 km] WNW Caleta de Campos (TCWC 55735–39); 1 mi [1.6 km] NE of La Placita 50 ft [15 m]

(UMMZ 104543, (1) specimen); NE of La Placita 75 ft [23 m] (UMMZ 104544, (1) specimen); 1 mi [1.6 km] NE of La Placita 75 ft

[23 m] (UMMZ 104547); 0.5 mi [0.8 km] SW of La Placita (UMMZ 104548, (2) specimens); Ostula 500 ft [152 m] (UMMZ 104550,

(5) specimens); near Pomaro along tributary to Rio Pasnori (UMMZ 104808, (15) specimens); near Rancho El Diezmo 4.5 mi [7.2

km] N of Coire 2920 ft [890 m] (UMMZ 104810, (2) specimens); Playa Azul (UMMZ 112790, (2) specimens).

Mexico: Nayarit: 21-25 mi [34-40 km] S. Acaponeta, 100' +/- Hwy 15 (LACM 25403); Compostela: 3 mi [5 km] SE of Sayulita

(UTEP 5855–56).

Mexico: Oaxaca: Pochutla 150 (KU 58322, 24–25, 28–29, 31); Hwy 185 between Tehuantepec & Salina Cruz (LACM 8172–81,

65054).

Mexico: Sinaloa: 1 mi [1.6 km] ENE San Lorenzo (KU 47926, 30, 32–33, 39–42).

Mexico: Veracruz: Vera Cruz, western part of city (AMNH 12282–83, 86–89, 91–92, 94–96, 12302–03, 10–11, 13–14, 16, 20, 23,

26–27, 29, 36, 38); [city of] Vera Cruz (AMNH 13390–91, 99, 13404, 14, 25, 28, 32); town of Pochote, 2.3 rd mi [3.7 km] NW jct

Hwy 140 at Tierra Colorada (UTA 54845–46).