**Diet and trophic niche partitioning of three aerial hawking bat species in Venezuela**

Yara Azofeifa Romero (1, 2, †)[ID], Jafet M Nassar (1) [ID], Jesús Mavárez (3, ‡)[ID]

(1) Laboratorio de Biología de Organismos, Centro de Ecología, Instituto Venezolano de Investigaciones Científicas, Carretera Panamericana km 11, Caracas 1204, Venezuela.

(2) Laboratorio de Sistemática, Genética y Evolución (LabSGE), Escuela de Ciencias Biológicas, Universidad Nacional. Heredia 40101, Costa Rica.

(3) Laboratoire d'Ecologie Alpine, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, F-38000 Grenoble, France.

Correspondence:

(†): LabSGE, Campus Omar Dengo, Calle 9, Avenidas 0 y 1, Costa Rica; yara.azofeifa.romero@una.ac.cr

(‡): LECA, Biologie Bâtiment D, 2233 rue de la Piscine, 38610 Gières, France; jesus.mavarez@univ-grenoble-alpes.fr

**ABSTRACT**

Since Neotropical insectivorous bats (IB) often share feeding sites and prey, their

coexistence likely depends on processes promoting divergence at trophic niche

dimensions. Here we conducted DNA metabarcoding analyses of diets and activity

times of three coexisting aerial hawking IB species — *Molossus molossus, Eptesicus

furinalis*, and *Myotis nigricans* — from rice fields in the Venezuelan Llanos, aiming to

understand their patterns of trophic niche partitioning. Insect prey were identified using

DNA sequences obtained from bat fecal samples and then compared against GenBank

and a local sequence database. Additionally, we recorded bat vocalizations over rice

fields simultaneously with the bat captures. We compared diversity, composition, and

breadth of the three IB diets, and employed statistical models to understand: (1) the

dietary and activity time overlap among IB species, and (2) the effect of IB species'

morphology and hunting strategies over prey's hardness and flight speed. We found

high prey richness for *M. molossus* (32 families), *M. nigricans* (19 families), and *E.

furinalis* (13 families). Bats exhibited differences in prey richness, proportions, types

consumed, and in their activity periods. For instance, as expected from its larger size

and fast flight, we found that *M. molossus* prefers larger, fast-flying, and medium-to-high

sclerotized preys such as Coleoptera and Lepidoptera, with hunting activities starting

and peaking significantly earlier than in the other two bat species. On the other hand,

also agreeing with its size and flight type, *M. nigricans* exhibits a significant preference

for small, slow-flying, and weekly sclerotized preys such as Diptera, although with

relatively high dietary and temporal overlaps with *E. furinalis.* Thus, in concordance with

niche theory predictions, the coexistence between these bats, particularly between *M.*

*molossus* and the other two species, seems enhanced by reduced competition driven by

divergence at trophic niche dimensions.

**Keywords:** agricultural lands, diets, DNA metabarcoding, insectivorous bats, trophic

niche, Venezuela

**INTRODUCTION**

Insectivorous bats (IB) are remarkably diverse predators with 20-45 species

coexisting in syntopy in tropical forests and crops in the Neotropics, Western Africa, and

Southeast Asia (Kingston et al. 2003; Williams-Guillén and Perfecto 2011; Reardon and

Schoeman 2017). The high local species richness of IB is likely a result of

ecomorphological differences allowing for a wide variety of hunting and echolocation

strategies among coexisting species, with which trophic niche differentiation is possible.

Indeed, coexisting IB usually differ in relation to wing morphology, body mass, skull

shape, and echolocation call design, all of which correlate with habitat use, foraging

strategies employed, and types of prey consumed (Marinello and Bernard 2014).

Foraging strategies and diets appear thus to be important axes of ecological

differentiation, contributing directly to resource partitioning among IB species.

Knowledge about foraging strategies in IB has been accumulating steadily in

recent decades, particularly with the advent of ultrasonic recording devices. Studies

based on acoustic surveys have indicated that sympatric aerial IB species partition food

habitats both horizontally and vertically, even at fine-grain spatial scales, in tropical and

temperate environments (Marques et al. 2016; Roemer et al. 2019; Beilke et al. 2021).

On the other hand, several studies have highlighted temporal differences in the onset

and/or maximum feeding activity periods among Neotropical IB species (Esbérard and

Bergallo 2010; Gomes et al. 2020). These spatiotemporal differences in foraging

strategies and patterns of activity time most likely led to dietary differences among IB

species, particularly in relation to size, hardness, position in the air column, and flight

mode of consumed prey (Segura-Trujillo et al. 2016, 2022). Regarding diets,

metabarcoding genetic analyses in IB from temperate regions have provided good

evidence supporting trophic flexibility, with different degrees of dietary overlap that can

only be explained through subtle spatiotemporal partitioning of food resources (Aizpurua

et al. 2018; Arrizabalaga-Escudero et al. 2019; Gordon et al. 2019; Andriollo et al.

2021). In contrast, molecular-based diet analyses in neotropical IB are still relatively

scarce, usually have low taxonomic resolution, and rarely make inter-specific

comparisons (but see Emrich et al. 2014, Salina-Ramos et al. 2015, Aguiar et al. 2021,

and Segura-Trujillo et al. 2022).

In this study, we performed a DNA metabarcoding analysis of diets of three aerial

hawking IB species from the Northwestern Llanos of Venezuela: *Molossus molossus*

(Molossidae), *Myotis nigricans*, and *Eptesicus furinalis* (Vespertilionidae). These bats

are frequently sympatric and can even inhabit the same roosts (Wilson and LaVal 1974).

They usually feed in the same areas, including relatively open spaces, forest gaps and

margins, and mosaics of agriculture and forestland in rural environments (Linares 1998).

However, the three species differ in various morphological and behavioral aspects that

greatly affect their feeding strategies. For example, *M. molossus* has larger body,

forearm, skull, and bite force than *M. nigricans*, while *E. furinalis* presents values that

are either similar to *M. nigricans* (body weight and skull length) or to *M. molossus*

(forearm length and bite force) (Linares 1998; Aguirre et al. 2002). Besides, differences

in wing shape and aspect ratio imply a fast and high flight in unobstructed spaces in *M.*

*molossus* (Marinello and Bernard 2014; Amador et al. 2020); but a comparatively slower

and more maneuverable flight in background-cluttered spaces in *M. nigricans* and *E.*

*furinalis* (Amador et al. 2020).

We analyzed diet composition and activity time of echolocation calls in the three

species to understand their patterns of trophic niche partitioning. We predicted that

given *M. molossus*' relatively larger size and distinctive flight type, its diet should both be

composed by a broader spectrum of prey and show relatively low overlap with the other

two species. Under the same argument, the diets of *E. furinalis* and *M. nigricans* should

exhibit some overlap given their smaller size and similar flight type. Furthermore, *M.*

*molossus*' relatively strong skull, fast flight, and long-term calls, should allow a diet with

important proportions of highly sclerotized and fast-flying insects. In contrast, these

types of insects should be relatively infrequent in the diets of *E. furinalis* and *M.*

*nigricans*, and be replaced by less sclerotized and slow-flying insects. Some predictions

can also be made in relation to bat activity time. For instance, if the activity time in these

bats is maximal during periods of high availability of preferred prey, *E. furinalis* and *M.*

*nigricans* should exhibit a significant degree of activity overlap over time given their

probably similar diets. For *M. molossus*, its overlap could be high if the availability of its

preferred prey is synchronous with that of the other two bats, and relatively low if it is

not.

**MATERIALS AND METHODS**

**Study area**

This study was conducted in the Northwestern Llanos, Portuguesa state,

Venezuela. Two rice farms located in this region were selected: Colonia Agrícola Turén,

Turén (hereafter "Turén"; 9.215005267° N, -69.122405063° W, 140 m asl.) and

Agropecuaria Durigua, Acarigua (hereafter "Acarigua"; 9.542531825° N, -

69.115494594° W, 180 m asl.) (Fig. 1).

**Sampling**

**Bats.** Bat captures followed the guidelines recommended by the American

Society of Mammalogists (Sikes et al. 2016). The two farms were visited six times

between November 2013 and September 2014 and bats were captured using mist nets

placed near their roosting sites. In Turén, the three bat species cohabitated the same

roosting site, while in Acarigua only *M. molossus* and *M. nigricans* were found in the

selected roosting site. However, *E. furinalis* certainly roosted in the forest remnants

nearby this farm, as it was recorded foraging in the rice fields. Bats were captured

between 20:00 and 21:30 h, when they returned to their roost after their first food intake.

Fecal pellets collected were placed under a small cotton ball at the bottom of a sterile 15

mL polypropylene centrifuge tube, and dried by filling the tube with silica gel type III

(Sigma-Aldrich H S7625-500 G). In total, fecal samples from 136 bats were collected (*M.*

*molossus*: 71, *M. nigricans*: 44, *E. furinalis*: 21).

**Insects**. Nocturnal insects were captured in the rice fields on the same dates and

hours of bat captures. The insects were identified and preserved in ethanol 70%. Of the

110.767 insects captured, we identified 11 orders, 52 families, and 98 morphospecies

(Moreno et al. 2020).

## DNA extractions

**Bat fecal samples**. DNA was extracted from ten fecal pellets of each bat using

the DNasy Blood & Tissue Kit (Qiagen, Valencia, CA, USA). Sterile 1.5 mL

microcentrifuge tubes containing fecal pellets and 1 mL lysis buffer solution were placed

on a shaker at 30 rpm in an incubator at 55-56°C for two hours. Afterward, samples

were centrifuged for 1 min at 1000 rpm, and 100 μL of supernatants were extracted and

deposited in new sterile 2 mL microcentrifuge tubes. The remaining protocol steps

followed the manufacturer's instructions. To detect cross-contamination, extractions

were performed in batches of 24 tubes, including two no-feces negative controls with

and without water. Extracted DNA was stored at -18°C until subsequent PCR analysis.

**Insects**. DNA was extracted from two legs of insects with body size > 10 mm and

from five whole bodies of smaller ones. Extractions were performed using the DNasy

Blood & Tissue Kit following the manufacturer's instructions.

## PCR amplification and sequencing

Two regions in the mitochondrial 16S rDNA unit were PCR-amplified for both

fecal and insect samples: "Inse01" with primers Inse01-F 5'-

RGACGAGAAGACCCTATARA-3' and Inse01-R 5'-ACGCTGTTATCCCTAARGTA-3',

and "Arth02" with primers Arth02-F 5'-GATAGAAACCAACCTGGYT-3' and LR-N-12945-

2 5'-GCGACCTCGATGTTGGATT-3' (Taberlet et al. 2018). All the primers had a

common adapter used for Illumina sequencing and an individual 8-bp barcode that

allowed the simultaneous sequencing of all samples.

DNA extracts were PCR-amplified in triplicate for fecal samples and once for

insects. PCR reactions (volume 20 µL) contained: 10 µL of AmpliTaq Gold Master Mix

(Applied Biosystems™, Foster City, USA), 0.2 µM of each primer, 5.84 µL of water, 0.04

µg of bovine serum albumin (Roche Diagnostic, Basel, Switzerland), and 2 µL of DNA

diluted 1:20. The PCR protocols were: 1x: 10 min at 95 °C, 45x: 30 s at 95 °C, 30 s at 52

°C (16S-1) or 50 ºC (16S-2), 60 s at 72 °C, 1x: 7 min at 72 °C. The PCR batches

included two extraction negatives, a PCR-negative (water), and a blank (empty well).

The PCR products were pooled per replicate, resulting in eight pools: six bat diets (three

per metabarcode) and two insects (one per metabarcode). Amplicon pools were purified

using the MinElute PCR Purification kit (Qiagen), quantified using a Qubit 2.0

Fluorometer (Life Technology Corporation), and sent to Fasteris SA (Geneva,

Switzerland) for library preparation and sequencing using the MetaFast protocol

(www.fasteris.com/metafast; Taberlet et al. 2018).

**Bioinformatic analyses**

The raw sequences were processed separately for each library using the

*OBITools* package (Boyer et al. 2016). Reads were assembled with a minimum quality

score of 40 and assigned to samples based on unique tag and primer combinations,

allowing for two mismatches with the primer. Sequences more abundant in extraction,

PCR and blank negative controls than in samples were considered as contaminants and

removed. Sequences with coverage <10 reads, or with lengths outside the expected

range (e.g., 76–168 bp for Arth02, Taberlet et al. 2018), or present in only one of the

three fecal sample replicates were also discarded. The remaining sequences were

clustered into prey MOTUs (*i.e.* Molecular Operational Taxonomic Unit) using the

sumaclust algorithm with a sequence similarity of 97% (Mercier et al. 2013). Taxonomic

assignment of MOTUs was achieved using the NCBI BLAST algorithm (Altschul et al.

1990), first against the 16S sequence database from local insects, and then against

GenBank for the sequences that could not be assigned in the previous way. The

following *ad hoc* assignment rules were applied in this study: (1) If there was a single

best match: species level for similarity ≥ 98%, genus level for 95% ≤ similarity < 98%,

family/sub-family level for 92% ≤ similarity < 95%, order level for similarity < 92%; (2) If

there were multiple best matches: assigned to the lowest taxonomic level subsuming all

best matches (e.g. assigned to the generic level if all best matches correspond to two or

more species in the same genus). Sequences that remained unassigned were classified

as "unknown". A table of diet items was created by removing rare occurrences (<

0.01%) and converting the counts into presence/absence values.

**Bat activity time**

We used two Echo Meter 3+ detectors placed on top of 2 m tall poles to record

bat calls as described by Azofeifa et al. (2019). Bat activity time was recorded

continuously for 240 min, beginning at sunset. Each sampling session consisted of two

consecutive nights, resulting in four bat recordings per rice farm. This two-night

recording scheme was repeated six times (three per climatic season), with a one-month

interval between two consecutive sampling sessions, resulting in 24 activity records per

farm.

**Data analyses**

We analyzed the diet composition of the three bat species using only insect

presence/absence data. Observed and expected richness of insect taxa in the bat diets

were calculated throughout the entire study period using the nonparametric estimator

Incidence-Based Bootstrap (specpool function, R package Vegan 2.6-4, Oksanen et al.

2022). Differences in diet composition were analyzed with PerMANOVA tests based on

Jaccard distance matrices using PRIMER v7 software (Clarke and Gorley 2015),

followed by pairwise comparisons among bat species using non-parametric pseudo-t

tests (akin to t-tests).

We conducted a comprehensive literature review to classify the feeding behavior

of adult insects into three main categories: phytophagous, saprophagous, and

zoophagous. Additionally, whenever possible, we assigned secondary categories (e.g.

anthophagous, detritivorous, hematophagous, etc). This review also searched for

biological characteristics of prey such as the degree of sclerotization (strong, moderate,

or weak), and flight behaviors (fast, intermediate, or slow).

We employed Generalized Linear Models (GLMs) with a Poisson distribution,

using the R package *MuMIn* 1.47.5 (Bartoń 2023), to assess the impact of bat factors

such as species, functional group, forearm length, skull length, or bite force on

characteristics of the prey consumed, using the prey count for each category as the

response variable. We determined the relative importance of each variable using

Akaike's Information Criterion (AIC) and the weight of evidence (wi). Statistical

significance was determined by $p < 0.05$, with confidence intervals excluding zero

indicating a significant impact for the predictor.

The trophic niche width, based on the prey proportion in the diet of each bat

species, was calculated using Levin's index (Levins 1968) with the niche.width function

in the R package *Spaa* 0.2.2 (Zhang 2016). The Levin's index was transformed into the

Best index, which standardizes it on a scale of 0 (*i.e.* specialized diet) to 1 (*i.e.*

diversified diet) (Krebs 1999). Diet overlap between species was evaluated using

Pianka's niche overlap index (Pianka 1973). We used the function niche_null_model in

the R package *EcoSimR* 0.1.0, to compare Pianka's index obtained with 100,000

replicates of null models (Gotelli et al. 2015). To illustrate the dietary overlap among the

three species, we conducted a Principal Component Analysis (PCA) using the Hill and

Smith method with the function dudi.hillsmith from the R package ade 4 1.7-22 (Dray et

al. 2023).

We also analyzed the overlap in activity time of the three bat species using a non-

parametric approach based on a kernel density estimated with the function density in the

R package *Stats* 4.3.2. We obtained estimates of the Weitzman Superposition

Coefficient Δ4 and its 95% confidence intervals using the *R* package *Overlapp* 0.3.9

(Meredith *et al*., 2024). We used the Watson and Watson-Wheeler tests available in R

package *Circular* 0.5-0 (Agostinelli and Lund 2023) to evaluate the goodness-of-fit of the

activity time data for individual species, and to compare the activity time data

distributions across pairs of bat species. Finally, we plotted the kernel density-time

curves with default smoothing parameters of the density function with the R package

*ggplot2* (Wickham 2016).

**RESULTS**

**Richness and composition of the bat diets**

A total of 65 insect taxa were identified in the feces of the three bats species, with

identifications ranging from order and family to, in some instances, the species level

(Table 1, Supplementary Data SD1). Diets included 50 MOTUs from 71 *Molossus*

*molossus*, 31 MOTUs from 44 *Myotis nigricans*, and 22 MOTUs from 21 *Eptesicus*

*furinalis*. Expected richness of insect taxa were 60.02 (SD = 2.89) for *M. molossus*,

36.17 (SD = 2.05) for *M. nigricans*, and 25.90 (SD = 1.89) for *E. furinalis*. Estimated

completeness of the diets were 83% for *M. molossus*, 86% for *M. nigricans*, and 85% for

*E. furinalis*. Statistical exploration of these MOTUs suggested differences in the dietary

composition of the bat species (Pseudo-$F_{2, 133}$ = 3.64, *P*-perm = 0.001). Specifically, the

pairwise comparisons *M. molossus* vs. *M. nigricans* (pseudo-$t$ = 2.39, *P*-perm = 0.001)

and *M. molossus* vs. *E. furinalis* (pseudo-$t$ = 1.70, *P*-perm = 0.003) were significantly

different.

Diet composition of the three bat species differed significantly at the order level

(Pseudo-$F_{2, 133}$ = 4.83, *P*-perm = 0.001). Ten insect orders were identified in the fecal

samples, with nine in *M. molossus*, seven in *M. nigricans*, and five *E. furinalis*. Highly

significant differences in composition of insect orders occurred between *M. molossus* vs.

*M. nigricans* (Pseudo-$t$ = 2.52, *P*-perm = 0.001) and *M. molossus* vs. *E. furinalis*

(Pseudo-$t$ = 2.31, *P*-perm = 0.003). For *M. molossus*, the preferred prey sequence was

Coleoptera, Lepidoptera, and Diptera. In contrast, *M. nigricans* primarily favored Diptera,

followed by Coleoptera and Hemiptera; whereas *E. furinalis* favored Coleoptera,

followed by Diptera and Hemiptera (Fig. 2A).

At the family level, differences were also notable among bat species (Pseudo-$F_{2,}$

$_{133}$ = 3.92, *P*-perm = 0.001). Forty insect families were identified in the fecal samples,

with 32 in *M. molossus*, 19 in *M. nigricans*, and 13 in *E. furinalis*. As above, the most

notable differences in composition of insect families were also observed between *M.*

*molossus* vs. *M. nigricans* (Pseudo-*t* = 2.39, *P*-perm = 0.001) and *M. molossus* vs. *E.*

*furinalis* (Pseudo-*t* = 1.85, *P*-perm = 0.002). For *M. molossus*, favored insect families,

from most to least preferred, were Tipulidae, Elateridae, Crambidae, Curculionidae,

Noctuidae, and Scirtidae. *Myotis nigricans*, on the other hand, primarily favored

Tipulidae, followed by Cicadelliadae, Heteroceridae, Crambidae, Ptilodactylinae, and

Culicidae, whereas *E. furinalis* preferred Tipulidae, followed by Cicadellidae, Carabidae,

Ptilodactylinae, Curculionidae, and Culicidae (Fig. 2B). Some prey were assigned to

genus or species level in orders Coleoptera (*Aeolus* sp., *Lissorhoptrus* sp.), Lepidoptera

(*Spodoptera frugiperda*, *Leucania* sp., *Condica sutor*, *Rupela albinella*, *Diatraea*

*saccharalis*, *Eoreuma loftini*), and Hemiptera (*Graminella* sp., *Tagosodes orizicolus*,

*Oebalus ypsilon-griseus*), most of which were ingested by *M. molossus* (Table 1).

No notable differences among bat species were detected as a function of prey

trophic groups. The three bat species primarily feed on phytophagous insects (39.9% of

diet in *M. molossus*, 31.1% in *E. furinalis*, 29.2% in *M. nigricans*, Fig. 2C).

Saprophagous prey represented 10.8% in *M. nigricans*, 8.2% in *E. furinalis*, and 5.2% in

*M. molossus*, while zoophagous prey accounted for just 8.2% in *E. furinalis*, 6.2% in *M.*

*nigricans*, and only 2.4% in *M. molossus*. Regarding the secondary functional

classification of the diet, *M. molossus* and *E. furinalis* exhibited four categories of

phytophagous insects (anthophagous, seed-feeding, phyllophagous, and xylophagous),

while *M. nigricans* displayed three categories (anthophagous, seed-feeding, and

phyllophagous). Additionally, four secondary categories within saprophagous insects

(detritivores and fungivores) and zoophagous insects (predators and hematophagous)

were detected in the diets of all three bat species.

**Prey types**

**Sclerotization**. The primary components in the diet of the two species with the

strongest bites, *M. molossus* and *E. furinalis*, were prey with a high degree of

sclerotization (Fig. 3). The results from the GLMs suggested that the lower bite force of

*M. nigricans* was negatively associated with highly sclerotized prey (wi = 0.63, *P* =

0.0114, CI = -0.92 to -0.12). Nonetheless, a notable similarity among the three species

was observed regarding the consumption of prey with intermediate sclerotization (Fig.

3). On the other hand, GLMs suggested that the lower bite force of *M. nigricans* was

positively associated with weakly sclerotized prey (wi = 0.29, *P* =0.001, CI = 0.27 to

0.98), although this type of prey was also present in the diets of the other species with

stronger bite force (see Fig. 3). Confidence intervals of these GLMs do not include zero,

indicating that bite force influences prey consumption with varying degrees of

sclerotization (Supplementary Data SD2).

**Flight speed**. Flight speed of prey appeared to be closely linked to bat hunting

ability in open areas. Specifically, fast-flying prey contribute largely to the diet of *M.*

*molossus*, moderately so to *M. nigricans*, and very little to *E. furinalis* (Fig. 4). The GLMs

indicated that the diets of *M. nigricans* and *E. furinalis* are negatively influenced by fast-

flying prey (wi = 0.38, p-value = 0.002, CI = -1.26 to -0.30). Bats with larger forearm, *M.*

*Molossus* and *E. furinalis*, showed a similar consumption of prey with intermediate flight

speeds. In fact, the GLMs indicated that the diet of the species with the smallest

forearm, *M. nigricans*, was negatively influenced by intermediate-flying prey (wi = 0.66,

*p*-value = 0.001, CI = -2.17 to -0.59). Slow-flying prey appeared to be more associated

with the edge-space foragers *E. furinalis* and *M. nigricans* (wi = 0.49, *p*-value = 0.00001,

CI = 0.34 to 1.02). However, slow-flying prey were also present in the diet of *M.*

*molossus*. Confidence intervals for these GLMs excluded zero, suggesting an impact of

bat forearm length and foraging strategy on the consumption of prey with diverse flight

behaviors (Supplementary Data SD2).

**Bat niche breadth**

321 The highest *Levin*'s index was found in *E. furinalis* ($B_{est}$ = 0.48), followed by *M.*

322 *nigricans* ($B_{est}$ = 0.29), and *M. molossus* ($B_{est}$ = 0.26). *Molossus molossus* and *M.*

323 *nigricans* had many low-proportion food items (75th percentile: values ≤ 1.54%), while *E.*

324 *furinalis* had a slightly more balanced diet although still with many low-proportion items

325 (75th percentile: values ≤ 1.64%).

**Bat trophic niche overlap**

The highest trophic niche overlap was observed in comparisons between *E.*

*furinalis* vs. *M. nigricans* (Pianka's index = 0.91), followed by *E. furinalis* vs. *M. molossus*

(Pianka's index = 0.76) and *M. molossus* vs. *M. nigricans* (Pianka's index = 0.72). For all

the pairwise comparisons, the probability of observing by chance a Pianka index equal

or higher than the one recorded was extremely low, indicating that the partial dietary

overlaps observed among species were higher than expected by chance alone (*P* <

0.001, n = 10,000 simulations). This result can be visualized in the Hill and Smith PCA,

which shows a clear lack of full overlap between the diet of *M. molossus* and the other

two species (Fig. 5). Furthermore, these analyses indicated the presence of 24 MOTUs

detected in single bat individuals, most of which were found only in *M. molossus*.

**Bat activity time**

*Molossus molossus* and *M. nigricans* represented 55.20% and 41.97%,

respectively, of the sound files obtained in both farms (n = 6007). In contrast, *Eptesicus*

*furinalis* had the lowest number of sound files during the entire study period (2.83%).

The distribution of activity over time is not uniform for any of the species (Watson test =

11.1 to 241.8, all *P* < 0.01). The activity time of *M. molossus* did not overlap much with

either *M. nigricans* (Δ4=0.36, CI bootstrap =0.34 and 0.38) or *E. furinalis* (Δ4=0.39, CI

bootstrap=0.34 and 0.44) (Fig. 6 A-B). A medium to high overlap in activity time between

*E. furinalis* and *M. nigricans* was observed throughout the study (Δ4=0.93, CI

bootstrap=0.88 and 0.97) (Fig. 6 C). Additional tests comparing the distributions of

activity over time in terms of orientation showed differences between *M. molossus* and

both *M. nigricans* and *E. furinalis* (Watson and Watson-Wheeler test = 248.1 to 2312.4,

all *P* = 2.2e-16). In contrast, the result of that same test in the comparison *M. nigricans*

vs. *E. furinalis* was not significant (Watson and Watson-Wheeler test = 3.1, *P* = 0.2)

(Supplementary Data SD3).

**DISCUSSION**

We investigated the diet and activity times of three bat species—*Molossus*

*molossus*, *Eptesicus furinalis*, and *Myotis nigricans*—in an agricultural setting,

employing metabarcoding of fecal samples and bioacoustic methods. These species

show some remarkable differences in the richness, taxonomic affinity, proportions, and

functional grouping of prey consumed, as well as in their activity times. As consequence,

these bats differ significantly in the breadth and degree of overlapping of their tropic

niche, all of which certainly resulting from morphological and behavioral attributes

related to bite force, size, and hunting abilities of these predators. Indeed, the diet and

activity differences observed suggest a division of food resources among these bats that

contribute to trophic niche partitioning, enhancing thereby their coexistence in close

sympatry.

**Dietary richness and composition**

The dietary taxonomic richness of insects in the studied species—*M. molossus*

(nine orders, 32 families), *M. nigricans* (seven orders, 19 families), and *E. furinalis* (five

orders, 13 families)— often surpassed richness values reported in other molecular diet

analyses of molossids and vespertilionids in agricultural areas. Notably, in our study, *M.*

*molossus* showed twice and 1.7 times the number of orders and families, respectively,

reported by Aguiar et al. (2021) for the same species in urban-agricultural environments

in Goiás, Brazil. Furthermore, molecular analyses of diets of other tropical molossid

species report family richness nearly three times lower than the value reported for *M.*

*molossus* in this study (Bohmann et al. 2011; Aguiar et al. 2021). However, Kolkert et al.

(2020) found 7.4 to 10.4 times more MOTUs richness in vespertilionid diets in

subtropical agricultural regions in New South Wales compared to our study. These

contrasting results suggest that bat dietary diversity is more variable than previously

understood, likely influenced by species-specific adaptations to climate, and agricultural

ecosystem characteristics (Hedrick and Dumont 2018).

In this study, differences in preference of insect orders among bat species were

conspicuous. For instance, in *M. molossus*, the three main types of prey were

Coleoptera (36.2%), Lepidoptera (22.5%) and Diptera (21.1%); whereas in *M. nigricans*

the ranking was Diptera (44.6%), Coleoptera (23.8%) and Hemiptera (13.1%), and in *E.*

*furinalis* was Coleoptera (39.3%), Diptera (32.8%) and Hemiptera (16.4%). The most

abundant insect orders captured in the study area were Coleoptera (61%), Hemiptera

(23%) and Diptera (15%) (Moreno et al. 2020). This suggests that these bats, especially

*M. molossus* and *E. furinalis*, are opportunistic predators whose diet compositions reflect

to a great extent the availability of potential insect prey. On the other hand, *M. nigricans*

exhibited a slightly more specialized behavior, with a marked preference for prey

(Diptera) that are not the most abundant in the study area. Thus, as observed in

generalist insectivorous birds, the structure of insect communities in agricultural crops

likely conditions the diets of aerial hawking bats (Durst et al. 2008; Aizpurua et al. 2018;

McClenaghan et al. 2019). However, the device and type of light used by Moreno et al.

(2020) to collect insects in our study area are probably biased towards capturing certain

taxa, so there is some uncertainty regarding the extent to which the bat diets reflect the

actual insect community structure.

Important sources of proteins and fats for bats include several phytophagous taxa

within Coleoptera, Lepidoptera, and Hemiptera. Notable among them are the elaterid

*Aeolus* sp. in the three bats species (Pi = 3.1% - 13.3%); the curculionid *Lissorhoptrus*

sp. in *M. molossus* (Pi = 5.6%) and *E. furinalis* (Pi = 4.9%); various moths in Crambidae

(*Rupela albinella* and *Diatraea saccharalis*) and Noctuidae (*Spodoptera frugiperda*,

*Condica sutor*, and *Leucania* sp.) in *M. molossus* ($\sum$Pi = 8.9%); and the cicadellid

*Graminella* sp. in *M. nigricans* (*Pi* = 10.8%) and *E. furinalis* (Pi = 8.2%). Notably, the

predominant prey type in the diets of these bats consists of phytophagous insects. This

indicates the high abundance of these insects, their nutritional value, and the efficiency

of converting them into bat body mass (García et al. 2012; Razeng and Watson 2015).

We found that Diptera represent an important fraction of the three bats diets (21.1

– 44.6%). For instance, the giant craneflies of the subfamily Limoniinae (Tipulidae)

accounted alone for 18.3% – 26.2% of the diets, more than any other food item

analyzed at the family/sub-family level. The identification of Diptera and other fragile

insects through morphological analysis of feces is not straightforward, which certainly

made difficult to understand their importance as food items, and to better ascertain the

drivers of food resource selection in non-molecular analyses of bat diets (Muñoz and

Molinari 2000; Whitaker and Philips 2012). Our research indicates that although Diptera

have lower nutritional value compared to Coleoptera or Lepidoptera, they remain

important food sources for bats due to their abundance and ease of capture in altered

environments (Liu et al. 2023), a result also observed in birds (Razeng and Watson

2015; Bellavance et al. 2018; McClenaghan et al. 2019). These vertebrates may use

Diptera either as substitutes of more nutritious but less available prey or as a

micronutrient supplement, which could be especially important in artificial wetlands.

**Types of prey captured**

We predicted that *M. molossus* should consume highly sclerotized prey, while for

*E. furinalis* and *M. nigricans* we suggested a higher consumption of weakly ones.

However, our findings partially align with these predictions, mainly due to *E. furinalis*.

Indeed, the best GLMs suggest that *M. molossus* and *E. furinalis* possess bite forces

suitable for consuming highly sclerotized prey, while *M. nigricans*' diet is driven by

weakly sclerotized prey due to its lower bite force. Previous studies have suggested that

bite force increases with skull length, which in turn correlates positively with the degree

of sclerotization of consumed prey (Senawi et al. 2015; Ramírez-Francel et al. 2021).

However, *E. furinalis* deviates from this relationship by having a skull length similar to *M.*

*nigricans,* yet with a bite force almost six times stronger (Aguirre et al. 2002). Although

the bite force of *E. furinalis* is slightly lower than that of *M. molossus*, it is sufficient to

overcome the mechanical limitations of processing highly sclerotized items, even with a

small skull. This aligns with Hedrick and Dumont (2018) finding that the dental rather

than the skull attributes are the primary focus of selective pressures related to food

processing, although such pattern seems more evident in phyllostomid than in non-

phyllostomid bat families.

The best GLM confirmed our prediction that the diets of *M. nigricans* and *E.*

*furinalis* should be negatively correlated with fast-flying prey, highlighting *M. molossus*'s

ability to hunt such prey through accelerated flight and long-range echolocation calls.

Similar to swallows and swifts, molossid bats have long, narrow wings, enabling agile

maneuvers during fast flights in open spaces at a lower cost compared to *M. nigricans*

and *E. furinalis*, which have short, wide wings (Norberg 1986; Voigt and Holderied 2012;

Matyjasiak et al. 2023). This same argument justifies the selection of the functional

group edge-space foragers by the GLM to explain the higher proportion of slow-flying

prey in *E. furinalis* and *M. nigricans.* The best GLM also indicated that forearm length

explains the significant similarities in intermediate-flying prey consumption between *M.*

*molossus* and *E. furinalis*. Forearm length, serving as a proxy for wing design (Thiagavel

et al. 2017), suggests that *E. furinalis*'s behavior of capturing slow-flying insects may

somewhat deviate from the typical edge-space foragers guild. In fact, *E. furinalis* has a

wing loading 1.2 times greater than *M. nigricans* (Amador et al. 2020), which could help

increase its flight speed, making it as efficient as *M. molossus* in capturing intermediate-

flying insects. However, all these arguments are part of a complex interplay of elements

that is barely understood, so they must be approached with caution.

**Trophic niche breadth**

As predicted, the diet of *M. molossus* had the highest taxonomic richness,

followed by *M. nigricans* and *E. furinalis*. However, contrary to expectations, the niche

breadth, as measured by the Best index, was nearly twice as big in *E. furinalis*

compared to the other bats. Thus, although *M. molossus* consumes a taxonomically

larger array of prey than *E. furinalis*, it is also less even in its consumption. These results

suggest that *M. Molossus* is perhaps better seen as a generalist species with great

trophic flexibility, a conclusion supported by our finding that 24% (17/71) *M. molossus*

individuals consumed prey detected only once, compared to 11% (5/44) and 10% (2/21)

in *M. nigricans* and *E. furinalis*, respectively. On a broader perspective, two types of

generalist species have been suggested to exist: somehow like *E. furinalis*, mostly

composed by individuals that consume a large array of prey equitably, and like *M.*

*Molossus*. and probably *M. nigricans*, composed by individuals that may consume

different kinds of prey (Bolnick et al. 2003). Although the latter category has been little

studied in bats, there is now some evidence supporting such pattern (Arrizabalaga-

Escudero et al. 2019; Magalhães de Oliveira et al. 2020).

**Trophic niche overlap**

As predicted, Pianka's index confirmed a significantly low dietary overlap between

*M. molossus* and the two vespertilionids, which, despite some differences in prey

contribution, are in turn highly overlapping. This observation aligns with findings by

Aguiar and Antonini (2008), who noted a high dietary overlap through morphological

analyses of fecal samples in *M. nigricans* and *E. furinalis* from the Brazilian Cerrado.

Other molecular analyses of bat diets have also found values of Pianka's index (0.60-

0.83) similar or slightly lower to those we obtained (0.72-0.91). These values have not

only resulted from comparisons between cryptic species (Razgour et al. 2011), but also

when inter-guild and even intra-guild comparisons are made (Salinas-Ramos et al.

2015; Gordon et al. 2019). In contrast, Segura-Trujillo et al. (2022) reported lower

dietary overlap among species from the same locality (Pianka's index ≤ 0.5) compared

to the same species from different localities (Pianka's index = ?), suggesting a

coexistence favored by reduced inter-specific competition via partitioning of trophic

niches. We believe that a similar process may enhance coexistence in the bats of this

study, although with higher overlapping of trophic niches, probably as a consequence of

the high productivity of rice fields in artificial wetlands, in which the elevated abundance

represents an almost unlimited food resource, leading to low competition and high

trophic niche overlap (Krüger et al. 2012).

**Bat activity time**

Consistent with our hypothesis, *M. molossus* exhibited low similarity in activity time

compared with *M. nigricans* and *E. furinalis*, which in turn had medium to high activity

time similarity among them. *Molossus molossus* is the first to appear in sound

recordings, followed closely by the near-simultaneous emergence of *M. nigricans* and *E.*

*furinalis*. Our observations and sound recordings, along with other studies indicating that

open-space bats forage before sunset (Esbérard and Bergallo 2010; Roemer et al.

2019), suggest that *M. molossus* is exploiting the insect community in a different

temporal window compared to the two vespertilionid species. The medium to high

overlapping times at activity peaks, occurring mainly between *M. nigricans* and *E.*

*furinalis*, suggest that competition between these bats might be at play, and that the

coexistence of the two species should be favored by divergence at trophic niche axes

other than diet composition and activity time.

**Concluding remarks**

The coexistence between *M. molossus*, *M. nigricans* and *E. furinalis* possibly

reflects differences in dietary preferences and the timing of hunting periods. *Molossus*

*molossus* shows a preference for relatively larger prey (e.g. Coleoptera and

Lepidoptera), that tend to fly fast and that exhibit a medium to high degree of

sclerotization, with both onset and peak of hunting activities occurring significantly

earlier than in the two other bats. Thus, divergence at these two trophic niche

dimensions should reduce inter-specific competition and enhance coexistence between

*M. molossus* and the vespertilionids, which agree with niche theory predictions (Finke

and Snyder, 2008). On the other hand, although *M. nigricans* exhibits a significant

preference for Diptera and other prey that tend to fly slowly and that are weekly

sclerotized, it shows relatively high dietary and temporal overlaps with *E. furinalis*. This

suggests that coexistence between these species is not favored by divergence at the

two trophic niche dimensions studied here but perhaps to the spatial structure of habitat

for hunting. In fact, these vespertilionids exhibit some aerodynamic differences that

make probable a differential use of space, both horizontal and vertical (Amador et al.

2020). Illuminating these aspects through integrative analyses will be key to unravel the

selective forces that contribute to the close coexistence and high diversification of bats

in tropical environments.

**References**

Agostinelli C., Lund U. 2023. Circular statistics. Version 0.5-0. [R package]. https://CRAN.R-project.org/package=circular

Aguiar LMS, Antonini Y. 2008. Diet of two sympatric insectivores bats (Chiroptera: Vespertilionidae) in the Cerrado of Central Brazil. *Revista Brasileira de Zoologia* 25(1):28–31. https://doi.org/10.1590/S0101-81752008000100005

Aguiar LMS, Bueno-Rocha ID, Oliveira G, Pires ES, Vasconcelos S, Nunes GL, Frizzas MR, Togni, PHB. 2021. Going out for dinner—The consumption of agriculture pests by bats in urban areas. *PLOS ONE* 16(10):e0258066. https://doi.org/10.1371/journal.pone.0258066

Aguirre LF, Herrel A, van Damme R, Matthysen E. 2002. Ecomorphological analysis of trophic niche partitioning in a tropical savannah bat community. *Proceedings of the Royal Society B: Biological Sciences* 269(1497):1271–1278. https://doi.org/10.1098/rspb.2002.2011

Aizpurua O, Budinski I, Georgiakakis P, Gopalakrishnan S, Ibañez C, Mata V, Rebelo H, Russo D, Szodoray-Parádi F, Zhelyazkova V, et al. 2018. Agriculture shapes the trophic niche of a bat preying on multiple pest arthropods across Europe: Evidence from DNA metabarcoding. *Molecular Ecology* 27(3):815–825. https://doi.org/10.1111/mec.14474

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *Journal of Molecular Biology* 215(3):403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Amador LI, Almeida FC, Giannini NP. 2020. Evolution of traditional aerodynamic variables in

bats (Mammalia: Chiroptera) within a comprehensive phylogenetic framework. *Journal of*

*Mammalian Evolution* 27:549–561. https://doi.org/10.1007/s10914-019-09475-8

Andriollo T, Michaux JR, Ruedi M. 2021. Food for everyone: Differential feeding habits of cryptic

bat species inferred from DNA metabarcoding. *Molecular Ecology* 30(18):4584–4600.

https://doi.org/10.1111/mec.16073

Arrizabalaga-Escudero A, Merckx T, García-Baquero G, Wahlberg N, Aizpurua O, Garin I, Goiti

U, Aihartza J. 2019. Trait-based functional dietary analysis provides a better insight into the

foraging ecology of bats. *Journal of Animal Ecology* 88(10):1587–1600.

https://doi.org/10.1111/1365-2656.13055

Azofeifa Y, Estrada-Villegas S, Mavárez J, Nassar JM. 2019. Activity of aerial insectivorous bats

in two rice fields in the Northwestern Llanos of Venezuela. *Acta Chiropterologica* 21(1):149–163.

https://doi.org/10.3161/15081109ACC2019.21.1.012

Bartoń K. 2023. MuMIn: Multi-Model Inference. Version 1.47.5. [R package]. https://CRAN.R-

project.org/package=MuMIn

Beilke EA, Blakey RV, O'Keefe JM. 2021. Bats partition activity in space and time in a large,

heterogeneous landscape. *Ecology and Evolution* 11(11):6513–6526.

https://doi.org/10.1002/ece3.7504

Bellavance V, Bélisle M, Savage J, Pelletier F, Garant D. 2018. Influence of agricultural

intensification on prey availability and nestling diet in tree swallows (*Tachycineta bicolor*).

*Canadian Journal of Zoology* 96(9):1053–1065. https://doi.org/10.1139/cjz-2017-0229

Bohmann K, Monadjem A, Lehmkuhl C, Rasmussen M, Zeale MRK, Clare E, Jones G,

Willerslev E, Gilbert MTP. 2011. Molecular diet analysis of two African free-tailed bats

(Molossidae) using high throughput sequencing. *PLoS ONE* 6(6):e21441.

https://doi.org/10.1371/journal.pone.0021441

Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML. 2003. The

ecology of individuals: Incidence and implications of individual specialization. *The American*

*Naturalist* 161(1):1–28. https://doi.org/10.1086/343878

Boyer F, Mercier C, Bonin A, Le Bras Y, Taberlet P, Coissac E. 2016. OBITOOLS: a Unix-

inspired software package for DNA metabarcoding. *Molecular Ecology Resources* 16(1):176–

182. https://doi.org/10.1111/1755-0998.12428

Clarke KR, Gorley RN. 2015. *PRIMER v7: User manual/tutorial*. Plymouth, UK: PRIMER-E Ltd.

Dray S, Dufour AB, Thioulouse J. 2023. Analysis of ecological data: Exploratory and euclidean

methods in environmental sciences. Version 1.7-22. [R package]. https://CRAN.R-

project.org/package=ade4

Durst SL, Theimer TC, Paxton EH, Sogge MK. 2008. Age, habitat, and yearly variation in the

diet of a generalist insectivore, the Southwestern Willow Flycatcher. *The Condor* 110(3):514–

525. https://doi.org/10.1525/cond.2008.8493

Emrich MA, Clare EL, Symondson WOC, Koenig SE, Fenton MB. 2014. Resource partitioning by

insectivorous bats in Jamaica. *Molecular Ecology* 23(15):3648–3656.

https://doi.org/10.1111/mec.12504

Esbérard CEL, Bergallo HG. 2010. Foraging activity of the free-tailed bat *Molossus molossus*

(Chiroptera; Molossidae) in southeastern Brazil. *Brazilian Journal of Biology* 70(4):1011–1014.

https://doi.org/10.1590/S1519-69842010000500013

Finke DL, Snyder WE. 2008. Niche partitioning increases resource exploitation by diverse

communities. *Science* 321(5895):1488–1490. https://doi.org/10.1126/science.1160854

García M, Farinós GP, Castañera P, Ortego F. 2012. Digestion, growth and reproductive

performance of the zoophytophagous rove beetle *Philonthus quisquiliarius* (Coleoptera:

Staphylinidae) fed on animal and plant based diets. *Journal of Insect Physiology 58*(10):1334–

1342. https://doi.org/10.1016/j.jinsphys.2012.07.007

Gomes DGE, Appel G, Barber JR. 2020. Time of night and moonlight structure vertical space

use by insectivorous bats in a Neotropical rainforest: An acoustic monitoring study. *PeerJ*

8:e10591. https://doi.org/10.7717/peerj.10591

Gordon R, Ivens S, Ammerman LK, Fenton MB, Littlefair JE, Ratcliffe JM, Clare EL. 2019.

Molecular diet analysis finds an insectivorous desert bat community dominated by resource

sharing despite diverse echolocation and foraging strategies. *Ecology and Evolution* 9(6):3117–

3129. https://doi.org/10.1002/ece3.4896

Gotelli NJ, Hart EM, Ellison AM. 2015. EcoSimR: Null model analysis for ecological data.

Version 0.1.0. [R package]. http://cran.nexr.com/web/packages/EcoSimR/index.html

Hedrick BP, Dumont ER. 2018. Putting the leaf-nosed bats in context: a geometric morphometric

analysis of three of the largest families of bats. *Journal of Mammalogy* 99(5):1042–1054.

https://doi.org/10.1093/jmammal/gyy101

Kingston T, Francis CM, Akbar Z, Kunz TH. 2003. Species richness in an insectivorous bat

assemblage from Malaysia. *Journal of Tropical Ecology* 19(1):67–79.

https://doi.org/10.1017/S0266467403003080

Krebs CJ. 1999. Ecological methodology (second edition). New York: Benjamin/Cummings.

Kolkert H, Andrew R, Smith R, Rader R, Reid N. 2020. Insectivorous bats selectively source

moths and eat mostly pest insects on dryland and irrigated cotton farms. *Ecology and Evolution*

10(1):371–388. https://doi.org/10.1002/ece3.5901

Krüger F, Harms I, Fichtner A, Wolz I, Sommer RS. 2012. High trophic similarity in the sympatric

North European trawling bat species *Myotis daubentonii* and *Myotis dasycneme*. *Acta*

*Chiropterologica* 14(2):347–356. http://dx.doi.org/10.3161/150811012X661666

Levins R. 1968. Evolution in changing environments: Some theoretical explorations. Princeton

University Press.

Linares O. 1998. Mamíferos de Venezuela. Sociedad Conservacionista Audubon de Venezuela,

Caracas, Venezuela.

Liu Y, Geng Y, Huang Z, Feng J, Jiang T. 2023. Pest suppression services and dietary niche

differentiation of bats in Chinese smallholder farming systems: implications for integrated pest

management. *Journal of Pest Science* 97:1587–1603. https://doi.org/10.1007/s10340-023-

01721-6

Magalhães de Oliveira HF, Camargo NF, Hemprich-Bennett DR., Rodríguez-Herrera B, Rossiter

SJ, Clare EL. 2020. Wing morphology predicts individual niche specialization in *Pteronotus*

*mesoamericanus* (Mammalia: Chiroptera). *PLOS ONE* 15(5):e0232601.

https://doi.org/10.1371/journal.pone.0232601

Marinello MM, Bernard E. 2014. Wing morphology of Neotropical bats: a quantitative and

qualitative analysis with implications for habitat use. *Canadian Journal of Zoology* 92(2):141–

147. https://doi.org/10.1139/cjz-2013-0127

Marques JT, Ramos Pereira MJ, Palmeirim JM. 2016. Patterns in the use of rainforest vertical

space by Neotropical aerial insectivorous bats: all the action is up in the canopy. *Ecography*

39(5):476–486. https://doi.org/10.1111/ecog.01453

Matyjasiak P, López-Calderón C, Ambrosini R, Balbontín J, Costanzo A, Kiat Y, Romano A,

Rubolini D. 2023. Wing morphology covaries with migration distance in a highly aerial

insectivorous songbird. *Current Zoology* 69(3):255-263. https://doi.org/10.1093/cz/zoac044

McClenaghan B, Nol E, Kerr KCR. 2019. DNA metabarcoding reveals the broad and flexible diet

of a declining aerial insectivore. *The Auk* 136(1):uky003. https://doi.org/10.1093/auk/uky003

Mercier C, Boyer F, Bonin A, Coissac E. 2013. SUMATRA and SUMACLUST: Fast and exact

comparison and clustering of sequences. Version 1.0.36. [Computer software].

http://metabarcoding.org/sumatra

Meredith M, Ridout M, Campbell LAD. 2024. overlap: Estimates of coefficient of overlapping for

animal activity patterns. Version 0.3.9. [R package]. https://CRAN.R-

project.org/package=overlap

Muñoz M, Molinari J. 2000. Insectos presa de los murciélagos molósidos Eumops perotis y

Molossus molossus. En: Péfaur J, editor. Ecología Latinoamericana. Actas del III Congreso

Latinoamericano; Universidad de Los Andes-Cons. Publ-CDCHT; Mérida (Venezuela); p. 223-

228.

Moreno A, Azofeifa Y, Domínguez C, Nassar JM. 2020. Ensamblaje de insectos con actividad

nocturna en dos arrozales de los Llanos Noroccidentales de Venezuela. *Anartia* 30:35–48.

https://produccioncientificaluz.org/index.php/anartia/article/view/34195

Norberg UM. 1986. Evolutionary convergence in foraging niche and flight morphology in

insectivorous aerial-hawking birds and bats. *Scandinavian Journal of Ornithology* 17(3):253–

260. https://doi.org/10.2307/3676835

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos

P, Stevens MHH, Szoecs E, et al. 2022. Vegan: Community Ecology Package. Version 2.6-4. [R

package]. https://CRAN.R-project.org/package=vegan

Pianka ER. 1973. The structure of lizard communities. Annual Review of *Ecology and*

*Systematics* 4:53–74. https://doi.org/10.1146/annurev.es.04.110173.000413

QGIS Development Team. 2023. QGIS Geographic Information System. Open-Source

Geospatial Foundation Project. Version 3.32. [Computer software]. https://www.qgis.org

Ramírez-Fráncel LA, García-Herrera LV, Losada-Prado S, Reinoso-Flórez G, Lim BK, Sánchez

F, Sánchez-Hernández A, Guevara G. 2021. Skull Morphology, bite force, and diet in

insectivorous bats from tropical dry forests in Colombia. *Biology* 10(10):1012.

https://doi.org/10.3390/biology10101012

Razgour O, Clare EL, Zeale MRK, Hanmer J, Schnell IB, Rasmussen M, Gilbert TP, Jones G.

2011. High-throughput sequencing offers insight into mechanisms of resource partitioning in

cryptic bat species. *Ecology and Evolution* 1(4):556–570. https://doi.org/10.1002/ece3.49

Razeng E, Watson DM. 2015. Nutritional composition of the preferred prey of insectivorous

birds: popularity reflects quality. Journal of Avian Biology 46(1):89–96.

https://doi.org/10.1111/jav.00475

Reardon S, Schoeman MC. 2017. Species richness, functional diversity and assemblage

structure of insectivorous bats along an elevational gradient in tropical West Africa. *Acta*

*Chiropterologica* 19(2):273–285. https://doi.org/10.3161/15081109ACC2017.19.2.005

Roemer C, Coulon A, Disca T, Bas Y. 2019. Bat sonar and wing morphology predict species

vertical niche. *The Journal of the Acoustical Society of America* 145(5):3242–3251.

https://doi.org/10.1121/1.5102166

Salinas-Ramos VB., Herrera Montalvo LG, León-Regagnon V, Arrizabalaga-Escudero A, Clare

EL. 2015. Dietary overlap and seasonality in three species of mormoopid bats from a tropical dry

forest. *Molecular Ecology* 24(20):5296–5307. https://doi.org/10.1111/mec.13386

Segura-Trujillo CA, Lidicker WZ, Álvarez-Castañeda ST. 2016. New perspectives on trophic

guilds of arthropodivorous bats in North and Central America. *Journal of Mammalogy 97*(2):644–

654. https://doi.org/10.1093/jmammal/gyv212

Segura-Trujillo CA, Álvarez-Castañeda ST, Castañeda-Rico S, Maldonado JE. 2022. Taxonomic

versus ecological prey traits among arthropodophagous bats: Implications for surveying trophic

partitioning patterns. *Journal of Mammalogy 103*(5):1071–1083.

https://doi.org/10.1093/jmammal/gyac027

Senawi J, Schmieder D, Siemers B, Kingston T. 2015. Beyond size – morphological predictors

of bite force in a diverse insectivorous bat assemblage from Malaysia. *Functional Ecology*

29(11):1411–1420. https://doi.org/10.1111/1365-2435.12447

Sikes RS, Animal Care and Use Committee of the American Society of Mammalogists. 2016.

2016 Guidelines of the American Society of Mammalogists for the use of wild mammals in

research and education. *Journal of Mammalogy* 97(3):663–688.

https://doi.org/10.1093/jmammal/gyw078

Taberlet P, Bonin A, Zinger L, Coissac É. 2018. Environmental DNA: For Biodiversity Research

and Monitoring. Oxford University Press.

Thiagavel J, Santana SE, Ratcliffe JM. 2017. Body size predicts echolocation call peak

frequency better than gape height in vespertilionid bats. *Scientific Reports* 7:828.

https://doi.org/10.1038/s41598-017-00959-2

Voigt CC, Holderied MW. 2012. High manoeuvring costs force narrow-winged molossid bats to

forage in open space. *Journal of Comparative Physiology B* 182:415–424.

https://doi.org/10.1007/s00360-011-0627-6

Whitaker Jr JO, Philips AF. (2012). Foods of the little free-tailed bats, *Molossus molossus*, from

Boca Chica Key, Monroe County, Florida. *Florida Scientist* 75(4):249–252.

https://www.jstor.org/stable/24321933

Wickham H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Version 3.4.4. [R package]. https://ggplot2.tidyverse.org

Williams-Guillén K, Perfecto I. (2011). Ensemble composition and activity levels of insectivorous

bats in response to management intensification in coffee agroforestry systems. PLoS ONE 6(1):

e16502. https://doi.org/10.1371/journal.pone.0016502

Wilson DE, LaVal RK. 1974. *Myotis nigricans*. *Mammalian Species* 39:1-3.

https://doi.org/10.2307/3503847

Zhang J. 2016. spaa: SPecies Association Analysis. Version 0.2.2. [R package].

https://CRAN.R-project.org/package=spaa

**Table 1**. Percentage of prey by taxon (Pi) in the diet of three species of aerial hawking bats at North-

western Venezuela's rice crops. Pi = ($\sum$Ni/N total) × 100, where $\sum$Ni is the total number of prey items for

taxon i found in the diet and N total is the total number of prey items across all taxa in the diet

| Taxa | *Molossus molossus* $N_{BATS} = 71$ $N_{PREYS} = 213$ | *Myotis nigricans* $N_{BATS} = 44$ $N_{PREYS} = 130$ | *Eptesicus furinalis* $N_{BATS} = 21$ $N_{PREYS} = 61$ |
|---|---|---|---|
| **Order Coleoptera** | | | |
| **Suborder Adephaga** | | | |
| **Carabidae** | | | |
| Unknown Carabidae | 2.35 | 0.00 | 6.56 |
| Cicindelinae | 0.00 | 0.00 | 1.64 |
| Dytiscidae | 0.47 | 0.00 | 0.00 |
| **Suborder Polyphaga** | | | |
| Unknown Polyphaga | 2.35 | 1.54 | 3.28 |
| Unknown Cucujiformia | 0.94 | 0.00 | 0.00 |
| Elateridae: Agrypninae (*Aeolus* sp) | 13.62 | 3.08 | 4.92 |
| Ptilodactylinae | 2.35 | 6.15 | 8.20 |
| **Curculionidae** | | | |
| Unknown Curculionidae | 0.47 | 0.00 | 1.64 |
| Scolytinae | 0.00 | 0.77 | 0.00 |
| Erirhininae*: Lissorhoptrus sp* | 5.63 | 1.54 | 4.92 |
| Scirtidae | 4.69 | 0.77 | 0.00 |
| Scarabaeidae: Dynastinae | 0.47 | 0.00 | 0.00 |
| Hydrophilidae | 0.94 | 1.54 | 0.00 |
| Heteroceridae | 0.00 | 6.92 | 4.92 |
| Tenebrionidae | 0.47 | 0.00 | 1.64 |
| Chrysomelidae | 0.94 | 0.00 | 1.64 |
| Throscidae | 0.00 | 0.77 | 0.00 |
| Zopheridae: Colydiinae | 0.47 | 0.77 | 0.00 |
| | | | |
| **Order Diptera** | | | |
| Unknown Diptera | 0.47 | 0.00 | 1.64 |
| **Suborder Nematocera** | | | |
| Unknown Nematocera | 0.00 | 1.54 | 0.00 |
| Sciaroidea | 0.00 | 1.54 | 1.64 |
| Tipuloidea: Limoniinae | 18.31 | 26.15 | 22.95 |
| Tipuloidea: Pediciidae | 0.47 | 0.00 | 0.00 |
| Culicoidea: Culicidae | 0.47 | 6.15 | 4.92 |
| Chironomoidea: Ceratopogonidae | 0.47 | 0.00 | 0.00 |
| Psychodoidea: Psychodinae | 0.00 | 3.08 | 0.00 |

**Suborder Brachycera**

| | | | |
|---|---|---|---|
| Unknown Muscomorpha | 0.00 | 1.54 | 0.00 |
| Calyptratae | 0.47 | 0.00 | 0.00 |
| Cyclorrhapha | 0.00 | 3.08 | 1.64 |
| Drosophilidae | 0.47 | 0.77 | 0.00 |
| Stratyiomidae | 0.00 | 0.77 | 0.00 |

**Order Lepidoptera**

| | | | |
|---|---|---|---|
| Unknown Lepidoptera | 0.47 | 0.00 | 0.00 |
| **Suborder Glossata** | | | |
| Rhopalocera | 2.35 | 0.00 | 0.00 |
| Obtectomera | 6.57 | 0.77 | 0.00 |
| **Crambidae** | | | |
| Unknown Crambidae | 2.35 | 3.85 | 0.00 |
| *Rupela albinella* | 4.23 | 1.54 | 0.00 |
| *Diatraea saccharalis* | 0.47 | 0.00 | 0.00 |
| *Eoreuma loftini* | 0.00 | 0.77 | 0.00 |
| **Noctuidae** | | | |
| Unknown Noctuidae | 0.94 | 0.00 | 0.00 |
| *Spodoptera frugiperda* | 3.29 | 0.00 | 0.00 |
| *Leucania* sp. | 0.47 | 0.00 | 0.00 |
| *Condica sutor* | 0.47 | 0.00 | 0.00 |
| Hesperiidae: Eudamidae | 0.00 | 0.00 | 1.64 |
| Geometridae | 0.00 | 0.77 | 0.00 |
| Saturnidae | 0.47 | 0.00 | 0.00 |
| Erebidae | 0.47 | 0.00 | 0.00 |

**Order Orthoptera**

| | | | |
|---|---|---|---|
| Unknown Orthoptera | 1.88 | 0.77 | 3.28 |
| **Suborder Ensifera** | | | |
| Gryllidae | 1.88 | 0.77 | 0.00 |

**Order Hemiptera**

| | | | |
|---|---|---|---|
| Unknown Hemiptera | 0.94 | 0.00 | 0.00 |
| **Suborder Heteroptera** | | | |
| Panheteroptera | 0.47 | 0.00 | 0.00 |
| Belostomatidae | 0.94 | 0.00 | 1.64 |
| Lygaeidae | 0.47 | 0.00 | 0.00 |
| Pentatomidae: *Oebalus ypsilongriseus* | 0.47 | 0.00 | 0.00 |
| **Suborder Auchenorrhyncha** | | | |
| **Cicadellidae** | | | |
| Unknown Cicadellidae | 1.41 | 2.31 | 3.28 |

|  | | | |
|---|---|---|---|
| *Graminella* sp | 1.41 | 10.77 | 8.20 |
| Cercopidae | 0.00 | 0.00 | 3.28 |
| Delphacidae: *Tagosodes orizicolus* | 0.94 | 0.00 | 0.00 |
|  |  |  |  |
| **Order Himenoptera**: Formicidae | 1.41 | 0.00 | 0.00 |
|  |  |  |  |
| **Order Blattodea** |  |  |  |
| Blaberidae | 0.47 | 0.00 | 0.00 |
| Rhinotermitidae | 0.47 | 1.54 | 0.00 |
| Kalotermitidae | 0.94 | 0.00 | 0.00 |
|  |  |  |  |
| **Order Neuroptera**: Chrysopinae | 0.47 | 0.00 | 0.00 |
|  |  |  |  |
| **Order Megaloptera**: Corydalidae | 0.00 | 2.31 | 0.00 |
|  |  |  |  |
| **Order Odonata**: Coenagrionidae | 0.47 | 0.00 | 0.00 |
|  |  |  |  |
| **Unknown Pterygota** | 5.16 | 5.38 | 6.56 |
| **Total (%)** | 100 | 100 | 100 |
| **Mean** | 1.54 | 1.54 | 1.54 |
| **Median** | 0.47 | 0.00 | 0.00 |
| **2sd** | 2.99 | 3.70 | 3.44 |
| **Min** | 0.00 | 0.00 | 0.00 |
| **Max** | 18.31 | 26.15 | 22.95 |
| **25th percentile** | 0.47 | 0.00 | 0.00 |
| **75th percentile** | 1.41 | 1.54 | 1.64 |

 Figure Legends

Fig. 1. Map of Acarigua and Turén rice farms in Northwestern Llanos, Portuguesa, Venezuela

(QGIS Development Team, 2023).

Fig. 2. Percentage of prey in the diet of three species of bats, classified by order, family and

trophic grpup throughout the study. For the classification of insects into trophic groups, only the

adult phase is considered. Unkn = Unknown, Phyt = Phytophagous, Zoop = Zoophagous, Sapr=

Saprophagous.

Fig. 3. Percentage of prey (%) strongly, medium, and weakly sclerotized on the diet of *M.*

*molossus*, *M. nigricans*, and *E. furinalis*.

Fig. 4. Percentage of prey (%) of fast, intermediate, and slow flying on the diet of *M. molossus*,

*M. nigricans*, and *E. furinalis*.

Fig. 5. Hill and Smith Principal Component Analysis (PCA hs) of dietary patterns in three bat

species. The analysis includes presence-absence data of the 65 MOTUs identified in this study.

Figure 6. Time-density curves of bat species pairs with overlap area in gray from both farms

(Acarigua and Turén) during the entire study period. Bootstrap estimates and 95% confidence

intervals for the overlap coefficient, Δ, are presented on the axis-x. The overlap coefficient Δ1 is

applied when the smaller sample size was less than 50, and Δ4 is used for sizes greater than

75.

Flight behavior of prey in the diet of three bat species

(A)

(B)

(B)