Diet and trophic niche partitioning of three aerial hawking bat species in the Venezuelan Llanos[1]

Yara Azofeifa Romero (1, 2, †) ⓘ, Jafet M Nassar (1) ⓘ, Jesús Mavárez (3, ‡) ⓘ

(1) Laboratorio de Biología de Organismos, Centro de Ecología, Instituto Venezolano de Investigaciones Científicas, Carretera Panamericana km 11, Caracas 1204, Venezuela.

(2) Laboratorio de Sistemática, Genética y Evolución (LabSGE), Escuela de Ciencias Biológicas, Universidad Nacional. Heredia 40101, Costa Rica.

(3) Laboratoire d'Ecologie Alpine, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, F-38000 Grenoble, France.

Correspondence:

(†): LabSGE, Campus Omar Dengo, Calle 9, Avenidas 0 y 1, Costa Rica; yara.azofeifa.romero@una.ac.cr

(‡): LECA, Biologie Bâtiment D, 2233 rue de la Piscine, 38610 Gières, France; jesus.mavarez@univ-grenoble-alpes.fr

[1]YA, JN, and JM formulated the idea. YA and JN developed the study's sampling design and sound recording methodology, YA and JM developed the genetic work. YA conducted fieldwork. JM generated sequencing data and conducted molecular analyses. YA performed the statistical analyses. YA and JM wrote the manuscript. JN provided feedback and additional information on earlier and final versions of the manuscript.

Diet and trophic niche partitioning of three aerial hawking bat species in the Venezuelan Llanos

ABSTRACT

Since Neotropical insectivorous bats (IB) share feeding sites and prey, their coexistence must depend on processes promoting divergence at trophic niches. We conducted diet and activity time analyses of three coexisting IB species in Venezuelan paddies, *Molossus molossus, Eptesicus furinalis*, and *Myotis nigricans*, to understand their trophic niche partitioning. We used statistical models to understand: (1) the dietary and activity time overlap among IB species, and (2) the relationship of IB species' morphology and hunting strategies with prey's hardness and flight speed. We found high prey richness for *M. molossus* (32 families), *M. nigricans* (19 families), and *E. furinalis* (13 families). Bats exhibited differences in prey richness, proportions, types consumed, as well as in their activity periods. As expected from its larger size and fast flight, *M. molossus* prefers larger, fast-flying, and medium-to-high sclerotized preys such as Coleoptera and Lepidoptera, with hunting activities starting and peaking significantly earlier than the other two bat species. On the other hand, also consistent with its size and flight type, *M. nigricans* exhibits a significant preference for small, slow-flying, and weekly sclerotized preys such as Diptera, although with relatively high dietary and temporal overlaps with *E. furinalis.* Thus, in concordance with niche theory predictions, the coexistence between these bats, particularly between *M. molossus* and the other two species, seems enhanced by reduced competition driven by divergence at trophic niche dimensions.

Keywords: coexistence, croplands, metabarcoding, insectivorous bats, activity time.

INTRODUCTION

Insectivorous bats (IB) are remarkably diverse predators with 20-45 species coexisting in syntopy in tropical forests and croplands in the Neotropics, Western Africa, and Southeast Asia (Kingston et al. 2003; Williams-Guillén and Perfecto 2011; Reardon and Schoeman 2017). The high local species richness of IB is likely a result of ecomorphological differences allowing for a wide variety of hunting and echolocation strategies among coexisting species, with which trophic niche differentiation is possible. Indeed, coexisting IB usually differ in relation to wing morphology, body mass, skull shape, and echolocation call design, all of which correlate with habitat use, foraging strategies employed, and types of prey consumed (Marinello and Bernard 2014). Thus, foraging strategies and diets appear to be important axes of ecological differentiation, contributing directly to resource partitioning among IB species.

Knowledge about foraging strategies in IB has been accumulating steadily in recent decades, particularly with the advent of ultrasonic recording devices. Studies based on acoustic surveys have indicated that sympatric aerial IB species partition food habitats both horizontally and vertically, even at fine-grain spatial scales, in tropical and temperate environments (Marques et al. 2016; Roemer et al. 2019; Beilke et al. 2021). On the other hand, several studies have highlighted temporal differences in the onset and/or maximum feeding activity periods among Neotropical IB species (Esbérard and Bergallo 2010; Gomes et al. 2020). These spatiotemporal differences in foraging strategies and patterns of activity time most likely led to dietary differences among IB species, particularly in relation to size, hardness, position in the air column, and flight mode of consumed prey (Segura-Trujillo et al. 2016, 2022). Regarding diet,

metabarcoding genetic analyses in IB from temperate regions have provided good

evidence supporting trophic flexibility, with different degrees of dietary overlap that can

only be explained through subtle spatiotemporal partitioning of food resources (Aizpurua

et al. 2018; Arrizabalaga-Escudero et al. 2019; Gordon et al. 2019; Andriollo et al.

2021). In contrast, molecular-based diet analyses in Neotropical IB are still relatively

scarce, usually have low taxonomic resolution, and rarely make inter-specific

comparisons (but see Emrich et al. 2014, Salina-Ramos et al. 2015, Aguiar et al. 2021,

and Segura-Trujillo et al. 2022).

In this study, we performed a DNA metabarcoding analysis of diets of three aerial

hawking IB species from the Northwestern Llanos of Venezuela: *Molossus molossus*

(Molossidae), *Myotis nigricans*, and *Eptesicus furinalis* (Vespertilionidae). These bats

are frequently sympatric and can even inhabit the same roosts (Wilson and LaVal 1974).

They usually feed in the same areas, including relatively open spaces, forest gaps and

margins, and mosaics of agriculture and forestland in rural environments (Linares 1998).

However, the three species differ in various morphological and behavioral aspects that

greatly affect their feeding strategies. For example, *M. molossus* has larger body,

forearm, skull, and bite force than *M. nigricans*, while *E. furinalis* presents measures that

are either similar to *M. nigricans* (body weight and skull length) or to *M. molossus*

(forearm length and bite force) (Linares 1998; Aguirre et al. 2002). Besides, differences

in wing shape and aspect ratio imply a fast and high flight in unobstructed spaces in *M.*

molossus (Marinello and Bernard 2014; Amador et al. 2020), but a comparatively slower

and more maneuverable flight in background-cluttered spaces in *M. nigricans* and *E.*

furinalis (Amador et al. 2020).

We analyzed diet composition and activity time of echolocation calls in the three

species to understand their patterns of trophic niche partitioning. We predicted that

given *M. molossus*' relatively larger size and distinctive flight type, its diet should both be

composed by a broader spectrum of prey and show relatively low overlap with the other

two species. Under the same argument, the diets of *E. furinalis* and *M. nigricans* should

exhibit some overlap given their smaller size and similar flight type. Furthermore, *M.*

molossus' relatively strong skull, fast flight, and long-term calls should allow a diet with

important proportions of highly sclerotized and fast-flying insects. In contrast, these

types of insects should be relatively infrequent in the diets of *E. furinalis* and *M.*

nigricans, and be replaced by less sclerotized and slow-flying insects. Some predictions

can also be made about to bat activity time. For instance, if the activity time in these

bats is maximal during periods of high availability of preferred prey, *E. furinalis* and *M.*

nigricans should exhibit significant activity overlap over time given their probably similar

diets. For *M. molossus*, its overlap could be high if the availability of its preferred prey is

synchronous with that of the other two bats, and relatively low if it is not.

MATERIALS AND METHODS

Study area

This study was conducted in the Northwestern Llanos, Portuguesa state,

Venezuela. Two rice farms located in this region were selected: Colonia Agrícola Turén,

Turén (hereafter "Turén"; 9.215005267° N, -69.122405063° W, 140 m asl.) and

Agropecuaria Durigua, Acarigua (hereafter "Acarigua"; 9.542531825° N, -

69.115494594° W, 180 m asl.) (Fig. 1).

Sampling

Bats. Bat captures followed the guidelines recommended by the American Society of Mammalogists (Sikes et al. 2016). The two farms were visited six times between November 2013 and September 2014 and bats were captured using mist nets placed near their roosting sites. In Turén, the three bat species cohabitated the same roosting site, while in Acarigua only *M. molossus* and *M. nigricans* were found in the selected roosting site. However, *E. furinalis* certainly roosted in the forest remnants nearby this farm, as it was recorded foraging in the rice fields. Bats were captured between 20:00 and 21:30 h, when they returned to their roost after their first food intake. Fecal pellets collected were placed under a small cotton ball at the bottom of a sterile 15 mL polypropylene centrifuge tube, and dried by filling the tube with silica gel type III (Sigma-Aldrich H S7625-500 G). In total, fecal samples from 136 bats were collected (*M. molossus*: 71, *M. nigricans*: 44, *E. furinalis*: 21).

Insects. Nocturnal insects were captured in the rice fields on the same dates and hours of bat captures. The insects were identified and preserved in ethanol 70%. Of the 110.767 insects captured, we identified 11 orders, 52 families, and 98 morphospecies (Moreno et al. 2020).

DNA extractions

Bat fecal samples. DNA was extracted from ten fecal pellets of each bat using the DNasy Blood & Tissue Kit (Qiagen, Valencia, CA, USA). Sterile 1.5 mL microcentrifuge tubes containing fecal pellets and 1 mL lysis buffer solution were placed on a shaker at 30 rpm in an incubator at 55-56°C for two hours. Afterward, samples

were centrifuged for 1 min at 1000 rpm, and 100 μL of supernatants were extracted and

deposited in new sterile 2 mL microcentrifuge tubes. The remaining protocol steps

followed the manufacturer's instructions. To detect cross-contamination, extractions

were performed in batches of 24 tubes, including two no-feces negative controls with

and without water. Extracted DNA was stored at -18°C until subsequent PCR analysis.

Insects. DNA was extracted from two legs of insects with body size > 10 mm and

from five whole bodies of smaller ones. Extractions were performed using the DNasy

Blood & Tissue Kit following the manufacturer's instructions.

PCR amplification and sequencing

Two regions in the mitochondrial 16S rDNA unit were PCR-amplified for both

fecal and insect samples: "Inse01" with primers Inse01-F 5'-

RGACGAGAAGACCCTATARA-3' and Inse01-R 5'-ACGCTGTTATCCCTAARGTA-3',

and "Arth02" with primers Arth02-F 5'-GATAGAAACCAACCTGGYT-3' and LR-N-12945-

2 5'-GCGACCTCGATGTTGGATT-3' (Taberlet et al. 2018). All the primers had a

common adapter used for Illumina sequencing and an individual 8-bp barcode that

allowed the simultaneous sequencing of all samples.

DNA extracts were PCR-amplified in triplicate for fecal samples and once for

insects. PCR reactions (volume 20 μL) contained: 10 μL of AmpliTaq Gold Master Mix

(Applied Biosystems™, Foster City, USA), 0.2 μM of each primer, 5.84 μL of water, 0.04

μg of bovine serum albumin (Roche Diagnostic, Basel, Switzerland), and 2 μL of DNA

diluted 1:20. The PCR protocols were: 1x: 10 min at 95 °C, 45x: 30 s at 95 °C, 30 s at 52

°C (16S-1) or 50 ºC (16S-2), 60 s at 72 °C, 1x: 7 min at 72 °C. The PCR batches

included two extraction negatives, a PCR-negative (water), and a blank (empty well).

The PCR products were pooled per replicate, resulting in eight pools: six bat diets (three

per metabarcode) and two insects (one per metabarcode). Amplicon pools were purified

using the MinElute PCR Purification kit (Qiagen), quantified using a Qubit 2.0

Fluorometer (Life Technology Corporation), and sent to Fasteris SA (Geneva,

Switzerland) for library preparation and sequencing using the MetaFast protocol

(www.fasteris.com/metafast; Taberlet et al. 2018).

Bioinformatic analyses

The raw sequences were processed separately for each library using the

OBITools package (Boyer et al. 2016). Reads were assembled with a minimum quality

score of 40 and assigned to samples based on unique tag and primer combinations,

allowing for two mismatches with the primer. Sequences more abundant in extraction,

PCR and blank negative controls than in samples were considered as contaminants and

removed. Sequences with coverage <10 reads, or with lengths outside the expected

range (e.g., 76–168 bp for Arth02, Taberlet et al. 2018), or present in only one of the

three fecal sample replicates were also discarded. The remaining sequences were

clustered into prey MOTUs (*i.e.* Molecular Operational Taxonomic Unit) using the

sumaclust algorithm with a sequence similarity of 97% (Mercier et al. 2013). Taxonomic

assignment of MOTUs was achieved using the NCBI BLAST algorithm (Altschul et al.

1990), first against the 16S sequence database from local insects, and then against

GenBank for the sequences that could not be assigned in the previous way. The

following *ad hoc* assignment rules were applied in this study: (1) If there was a single

best match: species level for similarity ≥ 98%, genus level for 95% ≤ similarity < 98%,

family/sub-family level for 92% ≤ similarity < 95%, order level for similarity < 92%; (2) If

there were multiple best matches: assigned to the lowest taxonomic level subsuming all

best matches (e.g. assigned to the generic level if all best matches correspond to two or

more species in the same genus). Sequences that remained unassigned were classified

as "unknown". A table of diet items was created by removing rare occurrences (<

0.01%) and converting the counts into presence/absence values.

Bat activity time

We used two Echo Meter 3+ detectors placed on top of 2 m tall poles to record

bat calls as described by Azofeifa et al. (2019). Bat activity time was recorded

continuously for 240 min, beginning at sunset. Each sampling session consisted of two

consecutive nights, resulting in four bat recordings per rice farm. This two-night

recording scheme was repeated six times (three per climatic season), with a one-month

interval between two consecutive sampling sessions, resulting in 24 activity records per

farm.

Data analyses

We analyzed the diet composition of the three bat species using only insect

presence/absence data. Observed and expected richness of insect taxa in the bat diets

were calculated throughout the entire study period using the nonparametric estimator

Incidence-Based Bootstrap (specpool function, R package Vegan 2.6-4, Oksanen et al.

2022). Differences in diet composition were analyzed with PerMANOVA tests based on

Jaccard distance matrices using PRIMER v7 software (Clarke and Gorley 2015),

followed by pairwise comparisons among bat species using non-parametric pseudo-t

tests (akin to t-tests).

We conducted a comprehensive literature review to classify the feeding behavior

of adult insects into three main categories: phytophagous, saprophagous, and

zoophagous. Additionally, whenever possible, we assigned secondary categories (e.g.

anthophagous, detritivorous, hematophagous, etc). This review also searched for

biological characteristics of prey such as the degree of sclerotization (strong, moderate,

or weak), and flight behaviors (fast, intermediate, or slow).

We employed Generalized Linear Models (GLMs) with a Poisson distribution,

using the R package *MuMIn* 1.47.5 (Bartoń 2023), to assess the impact of bat factors

such as species, functional group, forearm length, skull length, or bite force on

characteristics of the prey consumed, using the prey count for each category as the

response variable. We determined the relative importance of each variable using

Akaike's Information Criterion (AIC) and the weight of evidence (wi). Statistical

significance was determined by $p < 0.05$, with confidence intervals excluding zero

indicating a significant impact for the predictor.

The trophic niche width, based on the prey proportion in the diet of each bat

species, was calculated using Levin's index (Levins 1968) with the niche.width function

in the R package *Spaa* 0.2.2 (Zhang 2016). The Levin's index was transformed into the

Best index, which standardizes it on a scale of 0 (*i.e.* specialized diet) to 1 (*i.e.*

diversified diet) (Krebs 1999). Diet overlap between species was evaluated using

Pianka's niche overlap index (Pianka 1973). We used the function niche_null_model in

the R package *EcoSimR* 0.1.0, to compare Pianka's index obtained with 100,000

replicates of null models (Gotelli et al. 2015). To illustrate the dietary overlap among the

three species, we conducted a Principal Component Analysis (PCA) using the Hill and

Smith method with the function dudi.hillsmith from the R package ade 4 1.7-22 (Dray et

al. 2023).

We also analyzed the overlap in activity time of the three bat species using a non-

parametric approach based on a kernel density estimated with the function density in the

R package *Stats* 4.3.2. We obtained estimates of the Weitzman Superposition

Coefficient Δ4 and its 95% confidence intervals using the *R* package *Overlapp* 0.3.9

(Meredith *et al.*, 2024). We used the Watson and Watson-Wheeler tests available in R

package *Circular* 0.5-0 (Agostinelli and Lund 2023) to evaluate the goodness-of-fit of the

activity time data for individual species, and to compare the activity time data

distributions across pairs of bat species. Finally, we plotted the kernel density-time

curves with default smoothing parameters of the density function with the R package

ggplot2 (Wickham 2016).

RESULTS

Dietary richness and composition

A total of 65 insect taxa were identified in the feces of the three bat species, with

identifications ranging from order and family to, in some instances, the species level

(Online Resource 1 and 2). Diets included 50 MOTUs from 71 *Molossus molossus*, 31

MOTUs from 44 *Myotis nigricans*, and 22 MOTUs from 21 *Eptesicus furinalis*. Expected

richness of insect taxa were 60.02 (SD = 2.89) for *M. molossus*, 36.17 (SD = 2.05) for

M. nigricans, and 25.90 (SD = 1.89) for *E. furinalis*. Estimated completeness of the diets

were 83% for *M. molossus*, 86% for *M. nigricans*, and 85% for *E. furinalis*. Statistical

exploration of these MOTUs suggested differences in the dietary composition of the bat

species (Pseudo-$F_{2, 133}$ = 3.64, *P*-perm = 0.001). Specifically, the pairwise comparisons

M. molossus vs. *M. nigricans* (pseudo-*t* = 2.39, *P*-perm = 0.001) and *M. molossus* vs. *E.*

furinalis (pseudo-*t* = 1.70, *P*-perm = 0.003) were significantly different.

Diet composition of the three bat species differed significantly at the order level

(Pseudo-$F_{2, 133}$ = 4.83, *P*-perm = 0.001). Ten insect orders were identified in the fecal

samples, with nine in *M. molossus*, seven in *M. nigricans*, and five *E. furinalis*. Highly

significant differences in composition of insect orders occurred between *M. molossus* vs.

M. nigricans (Pseudo-*t* = 2.52, *P*-perm = 0.001) and *M. molossus* vs. *E. furinalis*

(Pseudo-*t* = 2.31, *P*-perm = 0.003). For *M. molossus*, the preferred prey sequence was

Coleoptera, Lepidoptera, and Diptera. In contrast, *M. nigricans* primarily favored Diptera,

followed by Coleoptera and Hemiptera; whereas *E. furinalis* favored Coleoptera,

followed by Diptera and Hemiptera (Fig. 2a).

At the family level, differences were also notable among bat species (Pseudo-$F_{2, 133}$

= 3.92, *P*-perm = 0.001). Forty insect families were identified in the fecal samples,

with 32 in *M. molossus*, 19 in *M. nigricans*, and 13 in *E. furinalis*. As above, the most

notable differences in composition of insect families were also observed between *M.*

molossus vs. *M. nigricans* (Pseudo-*t* = 2.39, *P*-perm = 0.001) and *M. molossus* vs. *E.*

furinalis (Pseudo-*t* = 1.85, *P*-perm = 0.002). For *M. molossus*, favored insect families,

from most to least preferred, were Tipulidae, Elateridae, Crambidae, Curculionidae,

Noctuidae, and Scirtidae. *Myotis nigricans*, on the other hand, primarily favored

Tipulidae, followed by Cicadelliadae, Heteroceridae, Crambidae, Ptilodactylinae, and

Culicidae; whereas *E. furinalis* preferred Tipulidae, followed by Cicadellidae, Carabidae,

Ptilodactylinae, Curculionidae, and Culicidae (Fig. 2b). Some prey were assigned to

genus or species level in orders Coleoptera (*Aeolus* sp., *Lissorhoptrus* sp.), Lepidoptera

(*Spodoptera frugiperda*, *Leucania* sp., *Condica sutor*, *Rupela albinella*, *Diatraea*

saccharalis, *Eoreuma loftini*), and Hemiptera (*Graminella* sp., *Tagosodes orizicolus*,

Oebalus ypsilon-griseus), most of which were ingested by *M. molossus* (Online

Resource 1).

No notable differences among bat species were detected as a function of prey

trophic groups. The three bat species primarily feed on phytophagous insects (39.9% of

diet in *M. molossus*, 31.1% in *E. furinalis*, 29.2% in *M. nigricans*, Fig. 2c). Saprophagous

prey represented 10.8% in *M. nigricans*, 8.2% in *E. furinalis*, and 5.2% in *M. molossus*;

while zoophagous prey accounted for just 8.2% in *E. furinalis*, 6.2% in *M. nigricans*, and

only 2.4% in *M. molossus*. Regarding the secondary functional classification of the diet,

M. molossus and *E. furinalis* exhibited four categories of phytophagous insects

(anthophagous, seed-feeding, phyllophagous, and xylophagous), while *M. nigricans*

displayed three categories (anthophagous, seed-feeding, and phyllophagous).

Additionally, four secondary categories within saprophagous insects (detritivores and

fungivores) and zoophagous insects (predators and hematophagous) were detected in

the diets of all three bat species (Online Resource 3).

Prey types

Sclerotization. The primary components in the diet of the two species with the

strongest bites, *M. molossus* and *E. furinalis*, were prey with a high degree of

sclerotization (Fig. 3a). The results from the GLMs suggested that the lower bite force of

M. nigricans was negatively associated with highly sclerotized prey (wi = 0.63, *P* =

0.0114, CI = -0.92 to -0.12). Nonetheless, a notable similarity among the three species

was observed regarding the consumption of prey with intermediate sclerotization (Fig.

3a). On the other hand, GLMs suggested that the lower bite force of *M. nigricans* was

positively associated with weakly sclerotized prey (wi = 0.29, *P* =0.001, CI = 0.27 to

0.98), although this type of prey was also present in the diets of the other species with

stronger bite force (Fig. 3a). Confidence intervals of these GLMs do not include zero,

indicating that bite force influences prey consumption with varying degrees of

sclerotization (Online Resource 4).

Flight speed. Flight speed of prey appeared to be closely linked to bat hunting

ability in open areas. Specifically, fast-flying prey contribute largely to the diet of *M.*

molossus, moderately so to *M. nigricans*, and very little to *E. furinalis* (Fig. 3b). The

GLMs indicated that the diets of edge-space foragers, *M. nigricans* and *E. furinalis,* are

negatively influenced by fast-flying prey (wi = 0.38, p-value = 0.002, CI = -1.26 to -0.30).

Bats with larger forearm, *M. Molossus* and *E. furinalis*, showed a similar consumption of

prey with intermediate flight speeds (Fig. 3b). In fact, the GLMs indicated that the diet of

the species with the smallest forearm, *M. nigricans*, was negatively influenced by

intermediate-flying prey (wi = 0.66, *p*-value = 0.001, CI = -2.17 to -0.59). Slow-flying prey

appeared to be more associated with the edge-space foragers *E. furinalis* and *M.*

nigricans (wi = 0.49, *p*-value = 0.00001, CI = 0.34 to 1.02). However, slow-flying prey

were also present in the diet of *M. molossus* (Fig. 3b). Confidence intervals for these

GLMs excluded zero, suggesting an impact of bat forearm length and foraging strategy

on the consumption of prey with diverse flight behaviors (Online Resource 4).

Bat niche breadth

The highest *Levin*'s index was found in *E. furinalis* (B_{est} = 0.48), followed by *M. nigricans* (B_{est} = 0.29), and *M. molossus* (B_{est} = 0.26). *Molossus molossus* and *M. nigricans* had many low-proportion food items (75th percentile: values ≤ 1.54%), while *E. furinalis* had a slightly more balanced diet although still with many low-proportion items (75th percentile: values ≤ 1.64%).

Bat trophic niche overlap

The highest trophic niche overlap was observed in comparisons between *E. furinalis* vs. *M. nigricans* (Pianka's index = 0.91), followed by *E. furinalis* vs. *M. molossus* (Pianka's index = 0.76) and *M. molossus* vs. *M. nigricans* (Pianka's index = 0.72). For all pairwise comparisons, the probability of observing by chance a Pianka index equal or higher than the one recorded was extremely low, indicating that the partial dietary overlaps observed among species were higher than expected by chance alone (P < 0.001, n = 10,000 simulations). This result can be visualized in the Hill and Smith PCA, which shows a clear lack of full overlap between the diet of *M. molossus* and the other two species (Fig. 4). Furthermore, these analyses indicated the presence of 24 MOTUs detected in single bat individuals, most of which were found only in *M. molossus*.

Bat activity time

Molossus molossus and *M. nigricans* represented 55.20% and 41.97%, respectively, of the sound files obtained in both farms (n = 6007). In contrast, *Eptesicus furinalis* had the lowest number of sound files during the entire study period (2.83%). The distribution of activity over time is not uniform for any of the species (Watson test =

11.1 to 241.8, all *P* < 0.01). The activity time of *M. molossus* did not overlap much with

either *M. nigricans* (Δ4=0.36, CI bootstrap =0.34 and 0.38) or *E. furinalis* (Δ4=0.39, CI

bootstrap=0.34 and 0.44). A medium to high overlap in activity time between *E. furinalis*

and *M. nigricans* was observed throughout the study (Δ4=0.93, CI bootstrap=0.88 and

0.97) (Fig. 5). Additional tests comparing the distributions of activity over time in terms of

orientation showed differences between *M. molossus* and both *M. nigricans* and *E.*

furinalis (Watson and Watson-Wheeler test = 248.1 to 2312.4, all *P* = 2.2e-16). In

contrast, the result of that same test in the comparison *M. nigricans* vs. *E. furinalis* was

not significant (Watson and Watson-Wheeler test = 3.1, *P* = 0.2).

DISCUSSION

We investigated the diet and activity times of three bat species—*Molossus*

molossus, *Eptesicus furinalis*, and *Myotis nigricans*—in an agricultural setting,

employing metabarcoding of fecal samples and bioacoustics methods. These species

show remarkable differences in richness, taxonomic affinity, proportions, and functional

grouping of prey consumed, as well as in their activity times. In consequence, these bats

differ significantly in the breadth and degree of overlapping of their tropic niche, all of

which resulting from morphological and behavioral attributes related to bite force, size,

and hunting abilities of these predators. Indeed, the diet and activity differences

observed suggest a division of food resources among these bats that contribute to

trophic niche partitioning, enhancing their coexistence in close sympatry.

Dietary richness and composition

The dietary taxonomic richness of insects in the studied species—*M. molossus*

(nine orders, 32 families), *M. nigricans* (seven orders, 19 families), and *E. furinalis* (five

orders, 13 families)— often surpassed richness values reported in other molecular diet

analyses of molossids and vespertilionids in agricultural areas. Notably, in our study, *M.*

molossus showed twice and 1.7 times the number of orders and families, respectively,

reported by Aguiar et al. (2021) for the same species in urban-agricultural environments

in Goiás, Brazil. Furthermore, molecular analyses of diets of other tropical molossid

species report family richness nearly three times lower than the value reported for *M.*

molossus in this study (Bohmann et al. 2011; Aguiar et al. 2021). However, Kolkert et al.

(2020) found 7.4 to 10.4 times more MOTUs richness in vespertilionid diets in

subtropical agricultural regions in New South Wales compared to our study. These

contrasting results suggest that bat dietary diversity is more variable than previously

understood, likely influenced by species-specific adaptations to climate, and agricultural

ecosystem characteristics (Hedrick and Dumont 2018).

In this study, differences in preference of insect orders among bat species were

conspicuous. For instance, in *M. molossus*, the three main types of prey were

Coleoptera (36.2%), Lepidoptera (22.5%) and Diptera (21.1%); whereas in *M. nigricans*

the ranking was Diptera (44.6%), Coleoptera (23.8%) and Hemiptera (13.1%), and in *E.*

furinalis was Coleoptera (39.3%), Diptera (32.8%) and Hemiptera (16.4%). The most

abundant insect orders captured in the study area were Coleoptera (61%), Hemiptera

(23%) and Diptera (15%) (Moreno et al. 2020). This suggests that these bats, especially

M. molossus and *E. furinalis*, are opportunistic predators whose diet compositions reflect

to a great extent the availability of potential insect prey. On the other hand, *M. nigricans*

exhibited a slightly more specialized behavior, with a marked preference for prey

(Diptera) that are not the most abundant in the study area. Thus, as observed in

generalist insectivorous birds, the structure of insect communities in agricultural lands

likely conditions the diets of aerial hawking bats (Durst et al. 2008; Aizpurua et al. 2018;

McClenaghan et al. 2019). However, the device and type of light used by Moreno et al.

(2020) to collect insects in our study area are probably biased towards capturing certain

taxa, so there is some uncertainty regarding the extent to which the bat diets reflect the

actual insect community structure.

Important sources of proteins and fats for bats include several phytophagous taxa

within Coleoptera, Lepidoptera, and Hemiptera. Notable among them are the elaterid

Aeolus sp. in the three bats species (Pi = 3.1% - 13.3%); the curculionid *Lissorhoptrus*

sp. in *M. molossus* (Pi = 5.6%) and *E. furinalis* (Pi = 4.9%); various moths in Crambidae

(*Rupela albinella* and *Diatraea saccharalis*) and Noctuidae (*Spodoptera frugiperda*,

Condica sutor, and *Leucania* sp.) in *M. molossus* ($\sum$Pi = 8.9%); and the cicadellid

Graminella sp. in *M. nigricans* (*Pi* = 10.8%) and *E. furinalis* (Pi = 8.2%). Notably, the

predominant prey type in the diets of these bats consists of phytophagous insects. This

indicates the high abundance of these insects, their nutritional value, and the efficiency

of converting them into bat body mass (García et al. 2012; Razeng and Watson 2015).

We found that Diptera represent an important fraction of the three bats' diets

(21.1 – 44.6%). For instance, the giant craneflies of the subfamily Limoniinae (Tipulidae)

accounted alone for 18.3% – 26.2% of the diets, more than any other food item

analyzed at the family/sub-family level. The identification of Diptera and other fragile

insects through morphological analysis of feces is not straightforward (Whitaker and

Philips 2012; Silva-Duarte et al. 2024), which certainly made it difficult to understand

their importance as food items, and to better ascertain the drivers of food resource

selection in non-molecular analyses of bat diets. Our research indicates that although

Diptera have lower nutritional value compared to Coleoptera or Lepidoptera, they remain

important food sources for bats due to their abundance and ease of capture in altered

environments (Liu et al. 2023), a result also observed in birds (Razeng and Watson

2015; Bellavance et al. 2018; McClenaghan et al. 2019). These vertebrates may use

Diptera either as substitutes for more nutritious but less available prey or as a

micronutrient supplement, which could be especially important in artificial wetlands.

Types of prey captured

We predicted that *M. molossus* should consume highly sclerotized prey, while for

E. furinalis and *M. nigricans* we suggested a higher consumption of weak ones*.*

However, our findings partially align with these predictions, mainly due to *E. furinalis*.

Indeed, the best GLMs suggest that *M. molossus* and *E. furinalis* possess bite forces

suitable for consuming highly sclerotized prey, while *M. nigricans*' diet is driven by

weakly sclerotized prey due to its lower bite force. Previous studies have suggested that

bite force increases with skull length, which in turn correlates positively with the degree

of sclerotization of consumed prey (Senawi et al. 2015; Ramírez-Francel et al. 2021).

However, *E. furinalis* deviates from this relationship by having a skull length similar to *M.*

nigricans, yet with a bite force almost six times stronger (Aguirre et al. 2002). Although

the bite force of *E. furinalis* is slightly lower than that of *M. molossus*, it is sufficient to

overcome the mechanical limitations of processing highly sclerotized items, even with a

small skull. This aligns with Hedrick and Dumont (2018) finding that the dental rather

than the skull attributes are the primary focus of selective pressures related to food

processing, although such a pattern seems more evident in phyllostomids than in non-

phyllostomid bat families.

The best GLM confirmed our prediction that the diets of *M. nigricans* and *E.*

furinalis should be negatively correlated with fast-flying prey, highlighting *M. molossus*'s

ability to hunt such type prey through accelerated flight and long-range echolocation

calls. Similar to swallows and swifts, molossid bats have long, narrow wings, enabling

agile maneuvers during fast flights in open spaces at a lower cost compared to *M.*

nigricans and *E. furinalis*, which have short, wide wings (Norberg 1986; Voigt and

Holderied 2012; Matyjasiak et al. 2023). This same argument justifies the selection of

the functional group edge-space foragers by the GLM to explain the higher proportion of

slow-flying prey in *E. furinalis* and *M. nigricans.* The best GLM also indicated that

forearm length explains the significant similarities in intermediate-flying prey

consumption between *M. molossus* and *E. furinalis*. Forearm length, serving as a proxy

for wing design (Thiagavel et al. 2017), suggests that *E. furinalis*' behavior of capturing

slow-flying insects may somewhat deviate from the typical edge-space foragers guild. In

fact, *E. furinalis* has a wing loading 1.2 times greater than *M. nigricans* (Amador et al.

2020), which could help increase its flight speed, making it as efficient as *M. molossus* in

capturing intermediate-flying insects. However, all these arguments are part of a

complex interplay of elements that is barely understood, so they must be approached

with caution.

Trophic niche breadth

As predicted, the diet of *M. molossus* had the highest taxonomic richness,

followed by *M. nigricans* and *E. furinalis*. However, contrary to expectations, the niche

breadth, as measured by the Best index, was nearly twice as big in *E. furinalis*

compared to the other bats. Thus, although *M. molossus* consumes a taxonomically

larger array of prey than *E. furinalis*, it is also less even in its consumption. These results

suggest that *M. Molossus* is perhaps better seen as a generalist species with great

trophic flexibility, a conclusion supported by our finding that 24% (17/71) *M. molossus*

individuals consumed prey detected only once, compared to 11% (5/44) and 10% (2/21)

in *M. nigricans* and *E. furinalis*, respectively. On a broader perspective, two types of

generalist species have been suggested to exist: somehow like *E. furinalis*, mostly

composed by individuals that consume a large array of prey equitably, and like *M.*

Molossus and probably *M. nigricans*, with individuals that may differ considerably in the

kinds of preys consumed (Bolnick et al. 2003). Although the latter category has been

little studied in bats, there is now some evidence supporting such pattern (Arrizabalaga-

Escudero et al. 2019; Magalhães de Oliveira et al. 2020).

Trophic niche overlap

As predicted, Pianka's index confirmed a significantly low dietary overlap between

M. molossus and the two vespertilionids, which, despite some differences in prey

contribution, are in turn highly overlapping. This observation aligns with findings by

Aguiar and Antonini (2008), who noted a high dietary overlap through morphological

analyses of fecal samples in *M. nigricans* and *E. furinalis* from the Brazilian Cerrado.

Other molecular analyses of bat diets have also found values of Pianka's index (0.60-

0.83) similar or slightly lower to those we obtained (0.72-0.91). These values have not

only resulted from comparisons between cryptic species (Razgour et al. 2011), but also

when inter-guild and even intra-guild comparisons are made (Salinas-Ramos et al.

2015; Gordon et al. 2019). In contrast, Segura-Trujillo et al. (2022) reported lower

dietary overlap among species from the same locality (Pianka's index ≤ 0.5) compared

to the same species from different localities, suggesting a coexistence favored by

reduced inter-specific competition via partitioning of trophic niches. We believe that a

similar process may enhance coexistence in the bats of this study, although with higher

overlapping of trophic niches, probably as a consequence of the high productivity of rice

fields in artificial wetlands, in which the elevated abundance represents an almost

unlimited food resource, leading to low competition and high trophic niche overlap

(Krüger et al. 2012).

Bat activity time

Consistent with our hypothesis, *M. molossus* exhibited low similarity in activity time

compared with *M. nigricans* and *E. furinalis*, which in turn had medium to high activity

time similarity among them. *Molossus molossus* is the first to appear in sound

recordings, followed closely by the near-simultaneous emergence of *M. nigricans* and *E.*

furinalis. Our observations and sound recordings, along with other studies indicating that

open-space bats forage before sunset (Esbérard and Bergallo 2010; Roemer et al.

2019), suggest that *M. molossus* is exploiting the insect community in a different

temporal window compared to the two vespertilionid species. The medium to high

overlapping times at activity peaks, occurring mainly between *M. nigricans* and *E.*

furinalis, suggests that competition between these bats might be at play, and that the

coexistence of the two species should be favored by divergence at trophic niche axes

other than diet composition and activity time.

Concluding remarks

The coexistence between *M. molossus*, *M. nigricans,* and *E. furinalis* in the

Venezuelan Llanos possibly reflects differences in dietary preferences and the timing of

hunting periods. *Molossus molossus* shows a preference for relatively larger prey (e.g.

Coleoptera and Lepidoptera), that tend to fly fast and that exhibit a medium to high

degree of sclerotization, with both onset and peak of hunting activities occurring

significantly earlier than in the two other bats. Thus, divergence at these two trophic

niche dimensions should reduce inter-specific competition and enhance coexistence

between *M. molossus* and the vespertilionids, which agree with niche theory predictions

(Finke and Snyder, 2008). On the other hand, although *M. nigricans* exhibits a

significant preference for Diptera and other prey that tend to fly slowly and that are

weekly sclerotized, it shows relatively high dietary and temporal overlaps with *E.*

furinalis. This suggests that coexistence between these species is not favored by

divergence at the two trophic niche dimensions studied here but perhaps to the spatial

structure of habitats used for hunting. In fact, these vespertilionids exhibit some

aerodynamic differences that make likely probable a differential use of space, both

horizontal and vertical (Amador et al. 2020). Illuminating these aspects through

integrative analyses will be key to unravel the selective forces that contribute to the

close coexistence and high diversification of bats in tropical environments.

Acknowledgements. Special thanks to José Antonio González-Carcacía (IVIC) for his

support during some fieldwork sessions. Claudia Domínguez (IVIC) and Abimel Moreno

(IVIC) provided assistance in identifying collected insects. Delphine Rioux (LECA) and

Eric Coissac (LECA) assisted with molecular and bioinformatics analyses.

Declarations

Funding. The Universidad Nacional in Costa Rica provided a scholarship to partially

cover the fieldwork for the first author. The Rufford Foundation (grant # 12366-1), Bat

Conservation International, and Idea Wild provided funding or equipment. The

Laboratoire d'Ecologie Alpine supported the molecular and bioinformatics analyses.

Conflicts of interest. The authors declares no conflicting or competing interests.

Ethics approval Bat captures followed the guidelines recommended by the American

Society of Mammalogists. Oficina de Diversidad Biológica (Ministerio del Poder Popular

para Ecosocialismo y Aguas) provided permits to collect bats and obtain their fecal

samples (Permit No. 1010-2013), to collect insects (Permit N°1011-2013), and to

conduct genetic analyses on these samples (Permit N° 0241-2013 and Permit N° 059-

2015).

Consent to participate. Not applicable.

Consent for publication. Not applicable.

Availability of data and material. The datasets used and/or analysed during the

current study are available from the corresponding author on reasonable request.

Author contribution statement. All authors contributed to the study's conception and

design. YA and JN contributed to fundraising for the purchase of equipment and other

materials, as well as material preparations. YA contributed to data collection and

statistical analyses. JM contributed to fundraising, genetic data acquisition and

analyses. The first drafts of the manuscript were written by YA and JM. All authors

contributed to the final version. All authors read and approved the final manuscript.

References

Agostinelli C, Lund U (2023) Circular statistics. Version 0.5-0. [R package]. https://CRAN.R-
project.org/package=circular

Aguiar LMS, Antonini Y (2008) Diet of two sympatric insectivores bats (Chiroptera:
Vespertilionidae) in the Cerrado of Central Brazil. Rev Bras Zool 25(1):28–31 doi:
10.1590/S0101-81752008000100005

Aguiar LMS, Bueno-Rocha ID, Oliveira G, Pires ES, Vasconcelos S, Nunes GL, Frizzas MR,
Togni, PHB (2021) Going out for dinner—The consumption of agriculture pests by bats in
urban areas. PLoS One 16(10):e0258066 doi: 10.1371/journal.pone.0258066

Aguirre LF, Herrel A, van Damme R, Matthysen E (2002) Ecomorphological analysis of trophic
niche partitioning in a tropical savannah bat community. Proc R Soc Lond B Biol Sci
269(1497):1271–1278 doi: 10.1098/rspb.2002.2011

Aizpurua O, Budinski I, Georgiakakis P, Gopalakrishnan S, Ibañez C, Mata V, Rebelo H, Russo
D, Szodoray-Parádi F, Zhelyazkova V, et al. (2018) Agriculture shapes the trophic niche
of a bat preying on multiple pest arthropods across Europe: Evidence from DNA
metabarcoding. Mol Ecol 27(3):815–825 doi: 10.1111/mec.14474

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J
Mol Biol 215(3):403–410 doi: 10.1016/S0022-2836(05)80360-2

Amador LI, Almeida FC, Giannini NP (2020) Evolution of traditional aerodynamic variables in
bats (Mammalia: Chiroptera) within a comprehensive phylogenetic framework. J Mamm
Evol 27:549–561 doi: 10.1007/s10914-019-09475-8

Andriollo T, Michaux JR, Ruedi M (2021) Food for everyone: Differential feeding habits of cryptic
bat species inferred from DNA metabarcoding. Mol Ecol 30(18):4584–4600 doi:
10.1111/mec.16073

Arrizabalaga-Escudero A, Merckx T, García-Baquero G, Wahlberg N, Aizpurua O, Garin I, Goiti
U, Aihartza J (2019) Trait-based functional dietary analysis provides a better insight into
the foraging ecology of bats. J Anim Ecol 88(10):1587–1600 doi: 10.1111/1365-
2656.13055

Azofeifa Y, Estrada-Villegas S, Mavárez J, Nassar JM (2019) Activity of aerial insectivorous bats
in two rice fields in the Northwestern Llanos of Venezuela. Acta Chiropt 21(1):149–163
doi: 10.3161/15081109ACC2019.21.1.012

Bartoń K (2023) MuMIn: Multi-Model Inference. Version 1.47.5. [R package]. https://CRAN.R-
project.org/package=MuMIn

Beilke EA, Blakey RV, O'Keefe JM (2021) Bats partition activity in space and time in a large,
heterogeneous landscape. Ecol Evol 11(11):6513–6526 doi: 10.1002/ece3.7504

Bellavance V, Bélisle M, Savage J, Pelletier F, Garant D (2018) Influence of agricultural intensification on prey availability and nestling diet in tree swallows (*Tachycineta bicolor*). Can J Zool 96(9):1053–1065 doi: 10.1139/cjz-2017-0229

Bohmann K, Monadjem A, Lehmkuhl C, Rasmussen M, Zeale MRK, Clare E, Jones G, Willerslev E, Gilbert MTP (2011) Molecular diet analysis of two African free-tailed bats (Molossidae) using high throughput sequencing. PLoS One 6(6):e21441 doi: 10.1371/journal.pone.0021441

Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML (2003) The ecology of individuals: Incidence and implications of individual specialization. Am Nat 161(1):1–28 doi: 10.1086/343878

Boyer F, Mercier C, Bonin A, Le Bras Y, Taberlet P, Coissac E (2016) OBITOOLS: a Unix-inspired software package for DNA metabarcoding. Mol Ecol Resour 16(1):176–182 doi: 10.1111/1755-0998.12428

Clarke KR, Gorley RN (2015) *PRIMER v7: User manual/tutorial*. PRIMER-E Ltd, Plymouth, UK.

Dray S, Dufour AB, Thioulouse J (2023) Analysis of ecological data: Exploratory and euclidean methods in environmental sciences. Version 1.7-22. [R package]. https://CRAN.R-project.org/package=ade4

Durst SL, Theimer TC, Paxton EH, Sogge MK (2008) Age, habitat, and yearly variation in the diet of a generalist insectivore, the Southwestern Willow Flycatcher. Condor 110(3):514–525 doi: 10.1525/cond.2008.8493

Emrich MA, Clare EL, Symondson WOC, Koenig SE, Fenton MB (2014) Resource partitioning by insectivorous bats in Jamaica. Mol Ecol 23(15):3648–3656 doi: 10.1111/mec.12504

Esbérard CEL, Bergallo HG (2010) Foraging activity of the free-tailed bat *Molossus molossus* (Chiroptera; Molossidae) in southeastern Brazil. Rev Bras Biol 70(4):1011–1014 doi: 10.1590/S1519-69842010000500013

Finke DL, Snyder WE (2008) Niche partitioning increases resource exploitation by diverse communities. Science 321(5895):1488–1490 doi: 10.1126/science.1160854

García M, Farinós GP, Castañera P, Ortego F (2012) Digestion, growth and reproductive performance of the zoophytophagous rove beetle *Philonthus quisquiliarius* (Coleoptera: Staphylinidae) fed on animal and plant based diets. J Insect Physiol *58*(10):1334–1342 doi: 10.1016/j.jinsphys.2012.07.007

Gomes DGE, Appel G, Barber JR (2020) Time of night and moonlight structure vertical space use by insectivorous bats in a Neotropical rainforest: An acoustic monitoring study. PeerJ 8:e10591 doi: 10.7717/peerj.10591

Gordon R, Ivens S, Ammerman LK, Fenton MB, Littlefair JE, Ratcliffe JM, Clare EL (2019) Molecular diet analysis finds an insectivorous desert bat community dominated by resource sharing despite diverse echolocation and foraging strategies. Ecol Evol 9(6):3117–3129 doi: 10.1002/ece3.4896

Gotelli NJ, Hart EM, Ellison AM (2015) EcoSimR: Null model analysis for ecological data. Version 0.1.0. [R package]. http://cran.nexr.com/web/packages/EcoSimR/index.html

Hedrick BP, Dumont ER (2018) Putting the leaf-nosed bats in context: a geometric morphometric analysis of three of the largest families of bats. J Mammal 99(5):1042–1054 doi: 10.1093/jmammal/gyy101

Kingston T, Francis CM, Akbar Z, Kunz TH (2003) Species richness in an insectivorous bat assemblage from Malaysia. J Trop Ecol 19(1):67–79 doi: 10.1017/S0266467403003080

Krebs CJ (1999) Ecological methodology, 2nd edn. Benjamin/Cummings, New York.

Kolkert H, Andrew R, Smith R, Rader R, Reid N (2020) Insectivorous bats selectively source moths and eat mostly pest insects on dryland and irrigated cotton farms. Ecol Evol 10(1):371–388 doi: 10.1002/ece3.5901

Krüger F, Harms I, Fichtner A, Wolz I, Sommer RS (2012) High trophic similarity in the sympatric North European trawling bat species *Myotis daubentonii* and *Myotis dasycneme*. Acta Chiropt 14(2):347–356 doi: 10.3161/150811012X661666

Levins R (1968) Evolution in changing environments: Some theoretical explorations. Princeton University Press, Princeton.

Linares O (1998) Mamíferos de Venezuela. Sociedad Conservacionista Audubon de Venezuela, Caracas, Venezuela.

Liu Y, Geng Y, Huang Z, Feng J, Jiang T (2023) Pest suppression services and dietary niche differentiation of bats in Chinese smallholder farming systems: implications for integrated pest management. J Pest Sci 97:1587–1603 doi: 10.1007/s10340-023-01721-6

Magalhães de Oliveira HF, Camargo NF, Hemprich-Bennett DR., Rodríguez-Herrera B, Rossiter SJ, Clare EL (2020) Wing morphology predicts individual niche specialization in *Pteronotus mesoamericanus* (Mammalia: Chiroptera). PLoS One 15(5):e0232601 doi: 10.1371/journal.pone.0232601

Marinello MM, Bernard E (2014) Wing morphology of Neotropical bats: a quantitative and qualitative analysis with implications for habitat use. Can J Zool 92(2):141–147 doi: 10.1139/cjz-2013-0127

Marques JT, Ramos Pereira MJ, Palmeirim JM (2016) Patterns in the use of rainforest vertical space by Neotropical aerial insectivorous bats: all the action is up in the canopy. Ecography 39(5):476–486 doi: 10.1111/ecog.01453

Matyjasiak P, López-Calderón C, Ambrosini R, Balbontín J, Costanzo A, Kiat Y, Romano A, Rubolini D (2023) Wing morphology covaries with migration distance in a highly aerial insectivorous songbird. Curr Zool 69(3):255-263 doi: 10.1093/cz/zoac044

McClenaghan B, Nol E, Kerr KCR (2019) DNA metabarcoding reveals the broad and flexible diet of a declining aerial insectivore. Auk 136(1):uky003 doi: 10.1093/auk/uky003

Mercier C, Boyer F, Bonin A, Coissac E (2013) SUMATRA and SUMACLUST: Fast and exact comparison and clustering of sequences. Version 1.0.36. [Computer software]. http://metabarcoding.org/sumatra

Meredith M, Ridout M, Campbell LAD (2024) overlap: Estimates of coefficient of overlapping for animal activity patterns. Version 0.3.9. [R package]. https://CRAN.R-project.org/package=overlap

Moreno A, Azofeifa Y, Domínguez C, Nassar JM (2020) Ensamblaje de insectos con actividad nocturna en dos arrozales de los Llanos Noroccidentales de Venezuela. *Anartia* 30:35–48. https://produccioncientificaluz.org/index.php/anartia/article/view/34195

Norberg UM (1986) Evolutionary convergence in foraging niche and flight morphology in insectivorous aerial-hawking birds and bats. Scand J Ornithol 17(3):253–260 doi: 10.2307/3676835

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Solymos
P, Stevens MHH, Szoecs E, et al. (2022) Vegan: Community Ecology Package. Version
2.6-4. [R package]. https://CRAN.R-project.org/package=vegan

Pianka ER (1973) The structure of lizard communities. Annu Rev Ecol Evol Syst 4:53–74 doi:
10.1146/annurev.es.04.110173.000413

QGIS Development Team (2023) QGIS Geographic Information System. Open-Source
Geospatial Foundation Project. Version 3.32. [Computer software]. https://www.qgis.org

Ramírez-Fráncel LA, García-Herrera LV, Losada-Prado S, Reinoso-Flórez G, Lim BK, Sánchez
F, Sánchez-Hernández A, Guevara G (2021) Skull Morphology, bite force, and diet in
insectivorous bats from tropical dry forests in Colombia. Biology 10(10):1012 doi:
10.3390/biology10101012

Razgour O, Clare EL, Zeale MRK, Hanmer J, Schnell IB, Rasmussen M, Gilbert TP, Jones G
(2011) High-throughput sequencing offers insight into mechanisms of resource
partitioning in cryptic bat species. Ecol Evol 1(4):556–570 doi: 10.1002/ece3.49

Razeng E, Watson DM (2015) Nutritional composition of the preferred prey of insectivorous
birds: popularity reflects quality. J Avian Biol 46(1):89–96 doi: 10.1111/jav.00475

Reardon S, Schoeman MC (2017) Species richness, functional diversity and assemblage
structure of insectivorous bats along an elevational gradient in tropical West Africa. Acta
Chiropt 19(2):273–285 doi: 10.3161/15081109ACC2017.19.2.005

Roemer C, Coulon A, Disca T, Bas Y (2019) Bat sonar and wing morphology predict species
vertical niche. J Acoust Soc Am 145(5):3242–3251 doi: 10.1121/1.5102166

Salinas-Ramos VB., Herrera Montalvo LG, León-Regagnon V, Arrizabalaga-Escudero A, Clare
EL (2015) Dietary overlap and seasonality in three species of mormoopid bats from a
tropical dry forest. Mol Ecol 24(20):5296–5307 doi: 10.1111/mec.13386

Segura-Trujillo CA, Lidicker WZ, Álvarez-Castañeda ST (2016) New perspectives on trophic
guilds of arthropodivorous bats in North and Central America. J Mammal *97*(2):644–654
doi: 10.1093/jmammal/gyv212

Segura-Trujillo CA, Álvarez-Castañeda ST, Castañeda-Rico S, Maldonado JE (2022) Taxonomic
versus ecological prey traits among arthropodophagous bats: Implications for surveying
trophic partitioning patterns. J Mammal *103*(5):1071–1083 doi:
10.1093/jmammal/gyac027

Senawi J, Schmieder D, Siemers B, Kingston T (2015) Beyond size – morphological predictors
of bite force in a diverse insectivorous bat assemblage from Malaysia. Funct Ecol
29(11):1411–1420 doi: 10.1111/1365-2435.12447

Silva-Duarte L, Herrera-Sarmiento MM., Serrano-Cardozo VH, Ramírez-Pinilla MP (2024)
Reproductive activity and diet of an urban population of Pallas' free-tailed bats *Molossus*
molossus (Molossidae). Acta Chiropt 25(2):237–250 doi:
10.3161/15081109ACC2023.25.2.004

Sikes RS, Animal Care and Use Committee of the American Society of Mammalogists (2016)
2016 Guidelines of the American Society of Mammalogists for the use of wild mammals
in research and education. J Mammal 97(3):663–688 doi: 10.1093/jmammal/gyw078

Taberlet P, Bonin A, Zinger L, Coissac É (2018) Environmental DNA: For Biodiversity Research
and Monitoring. Oxford University Press, Oxford.

Thiagavel J, Santana SE, Ratcliffe JM (2017) Body size predicts echolocation call peak
frequency better than gape height in vespertilionid bats. Sci Rep 7:828 doi:
10.1038/s41598-017-00959-2

Voigt CC, Holderied MW (2012) High manoeuvring costs force narrow-winged molossid bats to
forage in open space. J Comp Physiol B 182:415–424 doi: 10.1007/s00360-011-0627-6

Whitaker Jr JO, Philips AF (2012) Foods of the little free-tailed bats, *Molossus molossus*, from
Boca Chica Key, Monroe County, Florida. Fla Sci 75(4):249–252
https://www.jstor.org/stable/24321933

Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
Version 3.4.4. [R package]. https://ggplot2.tidyverse.org

Williams-Guillén K, Perfecto I (2011) Ensemble composition and activity levels of insectivorous
bats in response to management intensification in coffee agroforestry systems. PLoS
One 6(1): e16502 doi: 10.1371/journal.pone.0016502

Wilson DE, LaVal RK (1974) *Myotis nigricans*. Mamm Species 39:1-3 doi: 10.2307/3503847

Zhang J (2016) spaa: SPecies Association Analysis. Version 0.2.2. [R package].
https://CRAN.R-project.org/package=spaa

Figure Legends

Figure 1. Map of Acarigua and Turén rice farms in Northwestern Llanos, Portuguesa, Venezuela (QGIS Development Team, 2023).

Figure 2. Percentage of prey in the diet of three species of bats, classified by (a) order, (b) family, and (c) trophic group throughout the study. For the classification of insects into trophic groups, only the adult phase is considered. Unkn = Unknown, Phyt = Phytophagous, Zoop = Zoophagous, Sapr= Saprophagous.

Figure 3. Percentage of prey in the diet of three species of bats, classified by (a) degree of sclerotization, and (b) flight behavior.

Figure 4. Hill and Smith Principal Component Analysis (PCA hs) of dietary patterns in three bat species. The analysis includes presence-absence data of the 65 MOTUs identified in this study.

Figure 5. Time-density curves of bat species pairs with overlap area in gray from both farms during the entire study period. Bootstrap estimates and 95% confidence intervals for the overlap coefficient, Δ, are presented on the axis-x. The overlap coefficient $\Delta 4$ is used for sizes greater than 75.

Figure 1

Figure 2

Figure 3

a)

b)

Figure 4

Figure 5

a)

Time (Overlap Δ4 = 0.36, CI bootstrap = 0.34 to 0.38)

b)

Time (Overlap Δ4 = 0.39, CI bootstrap = 0.34 to 0.44)

c)

Time (Overlap Δ4 = 0.93, CI bootstrap = 0.88 to 0.97)