*Contact information of corresponding author:

Yara Azofeifa Romero

Escuela de Ciencias Biológicas, Universidad Nacional.

Campus Omar Dengo, Heredia, Costa Rica, 9th Street, Avenues 0 and 1. P.O. Box: 86-3000.

Email: yara.azofeifa.romero@una.ac.cr

Trophic flexibility in three bat species

Trophic flexibility favors coexistence of three aerial-hawking bat species in Venezuelan rice fields

Yara Azofeifa Romero*[1,2] https://orcid.org/0000-0003-2874-4207, Jafet M Nassar[1] https://orcid.org/0000-0001-9808-9362, Jesús Mavárez[3] https://orcid.org/0000-0002-2483-0178

[1]Laboratorio de Biología de Organismos, Centro de Ecología, Instituto Venezolano de Investigaciones Científicas, Carretera Panamericana km 11, Caracas 1204, Venezuela. E-mail: jafet.nassar@gmail.com

[2]Laboratorio de Sistemática, Genética y Evolución (LabSGE), Escuela de Ciencias Biológicas, Universidad Nacional. Heredia, 86-3000, Costa Rica. E-mail: yara.azofeifa.romero@una.ac.cr

[3]Laboratoire d'Ecologie Alpine, Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, F-38000 Grenoble, France. E-mail: jesus.mavarez@univ-grenoble-alpes.fr

Teaser Text

We simultaneously identified prey items in feces and measured the activity times of

three aerial-hawking bat species in Neotropical rice fields. We then explored the

relationships between these variables and the functional traits of both prey and

predators, confirming that coexistence might be facilitated by highly productive

environments.

ABSTRACT

Coexistence among Neotropical insectivorous bats (IB) that share roosts, foraging

areas, and prey likely relies on processes promoting trophic niche divergence. We

examined the diet and activity times of three coexisting IB species (*Molossus molossus,*

Neoeptesicus furinalis, and *Myotis nigricans*) in Venezuelan rice fields in Northwestern

Llanos to assess dietary and temporal overlap. Using published sources, we categorized

prey by hardness and flight speed, bats by bite force and functional group, and

examined the relationships among these variables. IB species showed differences in

prey composition, type, and activity timing. As expected from its fast flight and strong

bite, *M. molossus'* diet consisted primarily of fast-flying, highly sclerotized insects, with

activity times peaking significantly earlier than in the other two bat species. In contrast,

M. nigricans and *N. furinalis* had diets consisting primarily of slower-flying prey and

showed high temporal overlap in activity, although with different peak foraging times.

Notably, *N. furinalis*'s stronger bite may have enabled it to exploit more sclerotized prey

than *M. nigricans*, despite similar flight capabilities—indicating that prey hardness helps

reduce dietary overlap. These findings suggest that fine-scale trophic niche partitioning,

enhanced by the rich insect fauna of rice fields, facilitates the coexistence of these

ecomorphologically distinct IB species.

Keywords: activity time, artificial wetlands, croplands, insectivorous bats, metabarcoding.

Flexibilidad trófica favorece la coexistencia de tres especies de murciélagos cazadores aéreos en cultivos de arroz de Venezuela.

RESUMEN

La coexistencia entre murciélagos insectívoros (MI) neotropicales que comparten refugios, áreas de forrajeo y presas probablemente depende de procesos que promueven la divergencia del nicho trófico. Examinamos la dieta y los tiempos de actividad de tres especies coexistentes de MI (*Molossus molossus*, *Neoeptesicus furinalis* y *Myotis nigricans*) en arrozales de los Llanos Noroccidentales de Venezuela para evaluar superposición dietaria y temporal. Utilizando publicaciones, categorizamos a las presas según su dureza y velocidad de vuelo, a los murciélagos según su fuerza de mordida y grupo funcional, y examinamos las relaciones entre estas variables. Las especies de MI mostraron diferencias en la composición, tipo de presas y tiempo de actividad. Como se esperaba por su vuelo rápido y mordida fuerte, la dieta de *M. molossus* consistió principalmente en insectos de vuelo rápido y esclerotización alta, con tiempos de actividad que alcanzaron un máximo significativamente antes que en las otras dos especies de murciélagos. En contraste, *M. nigricans* and *N. furinalis* tuvieron dietas que consistieron principalmente en presas de vuelo lento que mostraron una alta superposición temporal en la actividad, aunque con distintos tiempos de forrajeo máximo. Cabe destacar que la mordida más fuerte de *N. furinalis* pudo haberle permitido explotar presas más esclerotizadas que *M. nigricans*, a pesar de sus capacidades de vuelo similares, lo que indica que la dureza de las presas ayuda a

reducir la superposición dietaria. Estos hallazgos sugieren que una partición fina del

nicho trófico, facilitada por la rica fauna de insectos en los cultivos de arroz, promueve

la coexistencia de estas especies de MI ecomorfológicamente distintas.

Palabras clave: cultivos, humedales artificiales, metabarcoding, murciélagos

insectívoros, tiempo de actividad.

Insectivorous bats (IB) are remarkably diverse predators with 20-45 species coexisting

in syntopy in tropical forests and croplands in the Neotropics, Western Africa, and

Southeast Asia (Kingston et al. 2003; Williams-Guillén and Perfecto 2011; Reardon and

Schoeman 2017). The high local species richness of IB is likely a result of

ecomorphological differences allowing for a wide variety of hunting and echolocation

strategies among coexisting species, making possible trophic niche differentiation.

Indeed, coexisting IB usually differ in wing morphology, body mass, skull shape, and

echolocation call design, all of which correlate with habitat use, foraging strategies, and

types of prey (Marinello and Bernard 2014). Accordingly, foraging site and diet represent

important dimensions of ecological differentiation that contribute to resource partitioning

among IB species.

Acoustic survey studies conducted in tropical and temperate environments have

shown that sympatric aerial IB species partition foraging sites both horizontally and

vertically, even at fine-grain spatial scales (Marques et al. 2016; Roemer et al. 2019;

Beilke et al. 2021). In addition to spatial segregation, these species also partition

resources along temporal dimensions, such as emergence time, timing of peak resource

use (e.g. prey, water, or roost), and activity duration (Adams and Thibault 2006; Holland

et al. 2011; Thomas and Jacobs 2013). However, how these temporal dimensions

contribute to reducing interspecific competition is still unclear, particularly when less

specialized or ecomorphologically distinct species coexist with dominant foragers

(Mayberry et al. 2020; Beilke et al. 2021). Although the mechanisms behind

spatiotemporal segregation remain understudied, such segregation promotes dietary

divergence among IB species.

Clear differences in the diets of IB are not always evident, which is why

increasingly precise dietary analyses are required to reveal underlying patterns of

resource partitioning among coexisting species. Recent genetic metabarcoding analyses

of diet in IB from temperate and tropical regions—conducted in diverse environments

such as deserts, protected areas, agricultural landscapes, and settlements near caves—

provide good evidence supporting trophic flexibility, with different degrees of dietary

overlap that can be explained by subtle spatiotemporal partitioning of food resources

(Bohmann et al. 2011; Salinas-Ramos et al. 2015; Gordon et al. 2019). On the other

hand, larger dietary overlaps can be seen among IB species from highly productive

periods or environments characterized by an elevated abundance of insects (Andriollo et

al. 2021; Srilopan et al. 2024). Such conditions are typical of agricultural landscapes,

where bats—capable of forming IB-rich ensembles—have become the focus of trophic

niche partitioning studies that highlight their role as controllers of insect pests (Liu et al.

2024; Srilopan et al. 2024).

Rice fields provide an ideal setting for examining trophic niche partitioning, as

their managed conditions create a humid, prey-rich environment that attracts

ecomorphologically distinct IB species (Liu et al. 2024). Building on previous research in

these rice fields (Azofeifa et al. 2019), this study provides unpublished dietary data on

syntopic IB species to explore whether their close coexistence is facilitated by their

differences in ecomorphological traits and activity times. With that aim, we performed

DNA metabarcoding analysis of diets and assessed activity times inferred from

echolocation calls, to examine trophic niche partitioning of three aerial-hawking IB

species in rice fields in the Northwestern Llanos of Venezuela: *Molossus molossus*

(Molossidae), *Myotis nigricans*, and *Neoeptesicus furinalis* (both Vespertilionidae).

These species are frequently sympatric, can inhabit the same roosts (Wilson and LaVal

1974) and forage in the same areas, including relatively open spaces, forest gaps and

edges, and mosaics of agriculture and forestland in rural environments (Linares 1998).

However, they differ in various morphological and behavioral traits that greatly affect

their feeding strategies (Freeman 1981; Norberg and Rayner 1987). In this regard, *M.*

molossus has a larger body, forearm, skull, and bite force than *M. nigricans*, while *N.*

furinalis has values that are intermediate (e.g., body weight) or similar to *M. molossus*

(e.g., forearm length and bite force) (Linares 1998; Aguirre et al. 2002). Additionally,

differences in wing shape and aspect ratio imply a fast and high flight in uncluttered

spaces in *M. molossus* (Marinello and Bernard 2014; Amador et al. 2020), but a

comparatively slower and more maneuverable flight in background-cluttered spaces in

M. nigricans and *N. furinalis* (Amador et al. 2020).

We predict that given the relatively larger size and distinctive flight type of *M.*

molossus, its diet should be composed of a broader spectrum of prey and show

relatively low overlap with the other two species. Under the same argument, the diets of

N. furinalis and *M. nigricans* should reflect the differences in their body sizes, but also

show some overlap given their similar flight types. Furthermore, the relatively strong

skull, fast flight, and long-term calls of *M. molossus* should allow for a significant

contribution of highly sclerotized and fast-flying insects in its diet. In contrast, these

types of insects should have a lower frequency in the diets of *N. furinalis* and *M.*

nigricans, being replaced by less sclerotized and slow-flying insects. Some predictions

can also be made about activity times. For instance, assuming that activity times in

these bats are maximal during periods of high availability of preferred prey, activity

overlaps are expected to be high if prey flights are synchronous and low if they are not.

According to this, *N. furinalis* and *M. nigricans* should exhibit significant activity overlap

given their probably similar diets. For *M. molossus*, the overlap with the activity of other

bats could be high if the maximum availability of preferred prey occurs at the same time

for all three species, and relatively low if the preferred prey of *M. molossus* are

asynchronous.

MATERIALS AND METHODS

Study area

This study was conducted in the Portuguesa state, part of the Northwestern Llanos of

Venezuela, where two rice farms were selected: Colonia Agrícola Turén, Turén

(hereafter "Turén"; 9.215005267, -69.122405063, 140 m asl.) and Agropecuaria

Durigua, Acarigua (hereafter "Acarigua"; 9.542531825, -69.115494594, 180 m asl.) (Fig.

1). The two farms, located 45.6 km apart, lie in a flat, rice-dominated landscape with

only 4–21% fragmented forest cover across 78.5 km² (Azofeifa et al. 2019).

Sampling

Bats. Bat captures followed the guidelines recommended by the American Society of

Mammalogists (Sikes et al. 2016). Between November 2013 and September 2014, each

farm was visited six times, with ca. one-month interval between visits. Bats were

captured using two 2.6 × 9.0 m mist nets (Avinet, Inc.), placed near roosting sites on two

consecutive nights per visit at each site, and subsequently identified using the keys by

Linares (1998) and Gardner (2007) (Supplementary Data SD1). In Turén, the three bat

species cohabited beneath a human-made tent constructed with dried palm leaves,

whereas in Acarigua, only *M. molossus* and *M. nigricans* were found roosting under the

zinc roof and within the concrete beams of a warehouse. *Neoeptesicus furinalis*

probably roosts in the forest remnants nearby this farm, as it was recorded foraging in

the rice fields. Bats were captured between 20:00 and 21:30 h, shortly after the main

peak of nightly foraging activity, as they returned to their roosts. Since the roosts were

located in front of the rice fields and we consistently observed bats flying toward them,

we assumed that their foraging activity was concentrated in these crops. The bats were

placed inside a clean cotton bag to collect feces. Subsequently, fecal pellets collected

were placed under a small cotton ball at the bottom of a sterile 15 mL polypropylene

centrifuge tube, and dried by filling the tube with silica gel type III (Sigma-Aldrich H

S7625-500 G). In total, fecal samples from 136 bats were collected: 71 *M. molossus*, 44

M. nigricans, 21 *N. furinalis*.

Insects. Nocturnal insects were captured using a white-light intersection trap placed on

the rice fields on the same dates and hours of bat captures. The purpose of these

collections was to create DNA reference libraries and to corroborate similarities among

insect communities, as suggested by the literature on agricultural crops. Insects were

identified by Moreno et al. (2020), preserved in 70% ethanol, and stored at room

temperature until genetic analysis.

DNA extractions

Bat fecal samples. DNA was extracted from the 10 largest fecal pellets of each bat

sample using the DNasy Blood & Tissue Kit (Qiagen, Valencia, CA, USA). Sterile 1.5 mL

microcentrifuge tubes containing fecal pellets and 1 mL lysis buffer solution were placed

on a shaker at 30 rpm in an incubator at 55-56°C for two hours. Afterward, samples

were centrifuged for 1 min at 1000 rpm, and 100 µL of supernatants were extracted and

deposited in new sterile 2 mL microcentrifuge tubes. The remaining protocol steps

followed the manufacturer's instructions. To detect cross-contamination, extractions

were performed in batches of 24 tubes, including two no-feces negative controls with

and without water. Extracted DNA was stored at -18°C until subsequent PCR analysis.

Insects. DNA was extracted from two legs of insects with body size > 10 mm and from

five whole bodies of smaller ones. Extractions were performed using the DNasy Blood &

Tissue Kit following the manufacturer's instructions.

PCR amplification and sequencing

Two regions in the mitochondrial 16S rDNA unit were PCR-amplified for both fecal and

insect samples: "Inse01" with primers Inse01-F 5'-RGACGAGAAGACCCTATARA-3'

and Inse01-R 5'-ACGCTGTTATCCCTAARGTA-3', and "Arth02" with primers Arth02-F

5'-GATAGAAACCAACCTGGYT-3' and LR-N-12945-2 5'-GCGACCTCGATGTTGGATT-

3' (Taberlet et al. 2018). All the primers had a common adapter used for Illumina

sequencing and an individual 8-bp barcode that allowed the simultaneous sequencing of

all samples.

DNA extracts were PCR-amplified in triplicate for fecal samples and once for

insects. PCR reactions (volume 20 µL) contained: 10 µL of AmpliTaq Gold Master Mix

(Applied Biosystems™, Foster City, USA), 0.2 µM of each primer, 5.84 µL of water, 0.04

µg of bovine serum albumin (Roche Diagnostic, Basel, Switzerland), and 2 µL of DNA

diluted 1:20. The PCR protocols were: 1x: 10 min at 95 °C, 45x: 30 s at 95 °C, 30 s at 52

°C (16S-1) or 50 °C (16S-2), 60 s at 72 °C, 1x: 7 min at 72 °C. The PCR batches

included two extraction negatives, a PCR-negative (water), and a blank (empty well).

The PCR products were pooled per replicate, resulting in eight pools: six bat diets (three

per metabarcode) and two insects (one per metabarcode). Amplicon pools were purified

using the MinElute PCR Purification kit (Qiagen), quantified using a Qubit 2.0

Fluorometer (Life Technology Corporation), and sent to Fasteris SA (Geneva,

Switzerland) for library preparation and sequencing using the MetaFast protocol

(www.fasteris.com/metafast; Taberlet et al. 2018).

Bioinformatic analyses

The raw sequences were processed separately for each library using the *OBITools*

package (Boyer et al. 2016). Reads were assembled with a minimum quality score of 40

and assigned to samples based on unique tag and primer combinations, allowing for two

mismatches with the primer. Sequences more abundant in extraction, PCR and blank

negative controls than in samples were considered as contaminants and removed.

Sequences with coverage <10 reads, or with lengths outside the expected range (e.g.,

76–168 bp for Arth02, Taberlet et al. 2018), or present in only one of the three fecal

sample replicates were also discarded. The remaining sequences were clustered into

prey MOTUs (*i.e.* Molecular Operational Taxonomic Unit) using the sumaclust algorithm

with a sequence similarity of 97% (Mercier et al. 2013). Taxonomic assignment of

MOTUs was achieved using the NCBI BLAST algorithm (Altschul et al. 1990), first

against the 16S sequence database from local insects, and then against GenBank for

the sequences that could not be assigned in the previous way. The following *ad hoc*

assignment rules were applied in this study: (1) if there was a single best match: species

level for similarity ≥ 98%, genus level for 95% ≤ similarity < 98%, family/sub-family level

for 92% ≤ similarity < 95%, order level for similarity < 92%; (2) if there were multiple best

matches: assigned to the lowest taxonomic level subsuming all best matches (e.g.,

assigned to the generic level if all best matches correspond to two or more species in

the same genus). Sequences that remained unassigned were classified as "unknown". A

table of diet items was created by removing rare occurrences (frequency < 0.01%) and

converting the counts into presence/absence values. We then calculated the percentage

of prey by taxon (P_i) in the diet of three IB species as $P_i = (\sum N_i/N \text{ total}) \times 100$, where

$\sum N_i$ is the total number of prey items for taxon i found in the diet and N total is the total

number of prey items across all taxa in the diet.

Bat activity time

We used two Echo Meter 3+ detectors placed on top of 2 m tall poles to record bat calls

as described by Azofeifa et al. (2019). Bat calls were recorded continuously for 240 min,

beginning at sunset. Each recording session occurred in the same two consecutive

nights of bat capture mentioned above, resulting in four bat recordings per rice farm and

session. This two-night recording scheme was therefore also repeated six times, with a

one-month interval between two consecutive sampling sessions, resulting in 24 activity

records per farm.

Data analyses

Prior to dietary analyses, we used data from Moreno et al. (2020) to compare the insect

communities between farms. We applied Hellinger transformations to the abundance

data, calculated Bray-Curtis distances, and performed a PerMANOVA using PRIMER v7

software (Clarke and Gorley 2015). As insect communities did not differ between farms

throughout the year (Pseudo-$F_{1,10}$ = 1.234, P-perm = 0.234), neither within the dry

(Pseudo-$F_{1,4}$ = 1.033, P-perm = 0.598) nor the rainy seasons (Pseudo-$F_{1,4}$ = 1.113, P-

perm = 0.398), we combined the data from both farms for subsequent dietary analyses.

All diversity and composition analyses were based on presence/absence

(incidence) data. Dietary diversity (Hill numbers q = 0, 1, 2) was quantified using iNEXT

function in the R package *iNEXT* 3.0.1. Sample-size and coverage-based

rarefaction/extrapolation curves were generated with ggiNEXT function. The percentage

of sampling completeness for richness (q = 0) was calculated as (1-Q1/U) x 100, where

Q1 is the number of prey singletons and U is the total number of prey observed (Hsieh

et al. 2016). Differences in diet composition were analyzed with PerMANOVA tests

based on Jaccard distance matrices using PRIMER v7 software (Clarke and Gorley

2015), followed by pairwise comparisons among bat species using non-parametric

pseudo-t tests (akin to t-tests).

We employed Generalized Linear Models (GLMs) with a Poisson distribution,

using the R package *MuMIn* 1.47.5 (Bartoń 2023), to assess the impact of three factors

—IB species, functional group (uncluttered space *vs*. background-cluttered space), and

bite force (high *vs*. low)— on prey characteristics such as the degree of sclerotization

(strong, intermediate, or soft) and flight behavior (fast, intermediate, or slow)

(Supplementary Data SD1 and SD2). The response variable in each model was the

count of prey items within each category. We determined the relative importance of each

predictor by comparing Akaike's Information Criterion (AIC), calculating Akaike weights

(wi), and examining the magnitude and direction of the estimated coefficients (β).

Statistical significance was determined by $P < 0.05$, with confidence intervals excluding

zero indicating a significant impact for the predictor.

Trophic niche breadth was estimated using Hill numbers of order q=1 and q=2

with the iNEXT function (Hsieh et al. 2016). Trophic niche overlap among the three bat

species was assessed as pairwise Hill-number-based similarity ($1-\beta_q$) for q=0, 1 and 2

using the hill_taxa_parti_pairwise function in the *hillR* package v. 0.5.3 (Li 2018).

Additionally, to visually illustrate dietary overlap, we conducted a PCA on the Hellinger-

transformed incidence matrix using the rda function in the *vegan* package v. 2.6-10,

which centers each prey variable to zero mean and applies a community-style PCA with

equal Euclidean weighting (Oksanen et al. 2025).

We also analyzed the overlap in activity time of the three bat species using a non-

parametric approach based on a kernel density estimated with the function density in the

R package *Stats* 4.3.2. We obtained estimates of the Weitzman Superposition

Coefficient Δ4 and its 95% confidence intervals using the *R* package *Overlapp* 0.3.9

(Meredith et al. 2024). We used the Watson (U^2 statistic) and Watson-Wheeler tests (W

statistic) available in R package *Circular* 0.5-0 (Agostinelli and Lund 2023) to evaluate

the goodness-of-fit of the activity time data for individual species, and to compare the

activity time data distributions across pairs of bat species. Finally, we plotted the kernel

density-time curves with default smoothing parameters of the density function with the R

package *ggplot2* (Wickham 2016).

RESULTS

Dietary richness and composition

A total of 65 MOTUs were identified in the feces of the three bat species. We assigned

93.4–94.8% of MOTUs to order level, 77.9–83.8% to family/sub-family level, and 17.7–

30.9% to species level (Supplementary Data SD3 and SD4). Diets included 50 MOTUs

from 71 *Molossus molossus*, 31 MOTUs from 44 *Myotis nigricans*, and 22 MOTUs from

21 *Neoeptesicus furinalis*. Estimated prey richness were 84.5 for *M. molossus*, 37.4 for

M. nigricans, and 30.6 for *N. furinalis*. Sampling completeness at the observed sample

sizes was approximately 88% for *M. molossus*, 91% for *M. nigricans*, and 85% for *N.*

furinalis (Supplementary Data SD5).

Diet composition of the three bat species differed significantly at the order level

(Pseudo-$F_{2, 133}$ = 4.83, P-perm = 0.001). Ten insect orders were identified in the fecal

samples, with nine in *M. molossus*, seven in *M. nigricans*, and five in *N. furinalis*.

Significant differences in the composition of insect orders occurred between *M.*

molossus vs. *M. nigricans* (Pseudo-t = 2.52, P-perm = 0.001) and *M. molossus* vs. *N.*

furinalis (Pseudo-t = 2.31, P-perm = 0.003). For *M. molossus*, the most frequently

detected prey were Coleoptera, followed by Lepidoptera and Diptera. In contrast, *M.*

nigricans had a higher incidence of Diptera, followed by Coleoptera and Hemiptera;

whereas *N. furinalis* showed the highest incidence of Coleoptera, followed by Diptera

and Hemiptera (Supplementary Data SD3; Fig. 2A).

At the family/sub-family level, differences were also notable among bat species

(Pseudo-$F_{2, 133}$ = 3.92, *P*-perm = 0.001). Forty insect families/sub-families were identified

in the fecal samples, with 32 in *M. molossus*, 19 in *M. nigricans*, and 13 in *N. furinalis*.

As above, the most notable differences in composition of insect families/sub-families

were observed between *M. molossus* vs. *M. nigricans* (Pseudo-*t* = 2.39, *P*-perm =

0.001) and *M. molossus* vs. *N. furinalis* (Pseudo-*t* = 1.85, *P*-perm = 0.002). Limoniinae

was the most frequently detected sub-family in the three bat species; however, *M.*

molossus showed a higher incidence of Agrypninae, *N. furinalis* of Carabidae, and both

M. nigricans and *N. furinalis* shared a greater incidence of Cicadellidae (Fig. 2B). All

these insect taxa were collected in the rice fields (Supplementary Data SD3).

Prey types

Sclerotization. The primary components in the diet of the two species with the

strongest bites, *M. molossus* and *N. furinalis*, were prey with a high degree of

sclerotization (Fig. 3A). Model selection supported bite force as the best predictor (wi =

0.70), with lower bite force in *M. nigricans* associated with a significantly lower incidence

of strongly sclerotized prey in its diet (β = –0.52, *P*-value = 0.011). Nonetheless, the

frequency of prey with intermediate sclerotization was similar across the three species

(Fig. 3A). On the other hand, model selection showed similar support for species and

bite force (wi = 0.51 and 0.49, respectively), with lower bite force in *M. nigricans*

significantly increasing the presence of less-sclerotized prey (β = 0.62, *P*-value = 0.001),

although this type of prey was also present in the diets of the other species with stronger

bite force (Fig. 3A) (Supplementary Data SD6).

Flight speed. Prey flight speed appeared to be closely associated with species that

forage in uncluttered spaces, such as *M. molossus*. Fast-flying prey made up a major

component of the diet of *M. molossus*, whereas in the other species their contribution

was lower but not negligible (Fig. 3B). Model selection moderately supported functional

group as the best predictor (wi = 0.38), with background-cluttered space foragers, *M.*

nigricans and N. furinalis, associated with a significantly lower incidence of fast flyers

prey in their diet (β = –0.78, *P*-value = 0.002). Model selection supported species as the

best predictor of intermediate-flight prey (wi = 0.89), with *M. nigricans* showing a

significantly lower presence of these prey in its diet (β = –1.19, *P*-value = 0.015). For this

same prey type, functional group also had a negative effect (β = –0.87, *P*-value =

0.002), indicating that background-cluttered space foragers overall were less likely to

capture intermediate flyers (Fig. 3B). On the other hand, model selection supported

again functional group as the best predictor of slow-flying prey consumption (wi = 0.58),

with background-cluttered space foragers showing a significantly higher incidence of

slow flyers prey in their diets (β = 0.68, *P*-value = 0.00001). However, slow-flying prey

were also present in the diet of *M. molossus* (Fig. 3B) (Supplementary Data SD6).

Trophic niche breadth and overlap of IB species

Trophic niche breadth was highest in *M. molossus* (q_1=23.3), followed by *M. nigricans*

(q_1=15.8) and *N. furinalis* (q_1=14.7). Pairwise niche overlap was highest between *N.*

furinalis and *M. nigricans* (q_1=0.73), and lower between *M. molossus* and either *N.*

furinalis (q_1=0.609) or *M. nigricans* (q_1=0.592) (Fig. 4; Supplementary Data SD5).

Bat activity time

Molossus molossus and *M. nigricans* represented 55.20% and 41.97%, respectively, of

the sound files obtained in both farms (n = 6007). In contrast, *N. furinalis* had the lowest

number of sound files during the entire study period (2.83%). The distribution of activity

over time is not uniform for any of the species (U^2 = 11.1 to 241.8, all *P* < 0.01). The

activity time of *M. molossus* did not overlap much with either *M. nigricans* (Δ4=0.36) or

N. furinalis (Δ4=0.39). A high overlap in activity time between *N. furinalis* and *M.*

nigricans was observed throughout the study (Δ4=0.93). However, peak activity times

differed among the three species: 6:45 p.m. for *M. molossus*; 7:19 p.m. and 8:35 pm for

N. furinalis; and 7:30 pm and 8:24 p.m. for *M. nigricans* (Fig. 5A-C). Additional tests

comparing the distributions of activity over time in terms of orientation showed

differences between *M. molossus* and both *M. nigricans* and *N. furinalis* (*W* = 248.1 to

2312.4, all *P* = 2.2e-16). In contrast, the same test comparing *M. nigricans* vs. *N.*

furinalis was not significant (*W* = 3.1, *P* = 0.2).

DISCUSSION

We investigated the diet and activity times of three species of aerial-hawking bats—

Molossus molossus, *Neoeptesicus furinalis*, and *Myotis nigricans*—in rice fields using

DNA metabarcoding of fecal samples and bioacoustics methods. These species

exhibited distinct degrees of differentiation in dietary taxonomic richness and

composition, biological attributes of prey consumed, and activity times. Consequently,

they differed in the breadth and degree of overlap of their trophic niches, all of which

resulted from traits related to bite force, hunting abilities, and intrinsic characteristics of

each IB species. In addition to the traits analyzed, the proximity of roosts to foraging

sites also influenced prey type by facilitating access to the insect community present in

the rice fields. The humid conditions and high prey richness in rice fields therefore

attractants of ecomorphologically distinct species, capable of exploiting these

anthropogenic wetlands in distinct ways. These adaptive differences promote the

coexistence of the three species, which appear to be structured by fine-scale trophic

niche partitioning.

Taxonomic richness and composition of prey

The diets of the IB species studied reflect the typical taxonomic richness and

composition of insect communities inhabiting rice fields (Acosta et al. 2017; Yunus et al

2022). These rice fields undergo cyclical disturbances—such as flooding and

harvesting—that promote temporal insect turnover, benefiting IB species that forage in

such dynamic habitats. This may explain the high prey richness observed in the diets of

the studied IB species—*M. molossus*, for example, showed nearly three times more

prey families than other molossid bats from less diverse systems such as sugarcane or

agro-urban environments (Bohmann et al. 2011; Aguiar et al. 2021). However, Kolkert et

al. (2020) reported much higher prey richness—7.4 to 10.4 times more—in vespertilionid

bat diets from subtropical agricultural regions with cotton crops in New South Wales.

This contrast likely reflects the broader range of habitat types present in that study,

including forest, floodplain woodland and semi-arid grassland. By comparison, the rice

fields in our study are embedded in a highly fragmented landscape, yet they appear to

provide sufficient prey to buffer IB bats from the typical effects of habitat fragmentation.

Several insect families/sub-families detected in the IB diets, such as Limoniinae,

Agrypninae, Curculionidae, Carabidae, Heteroceridae, Cicadellidae, and Crambidae,

exhibit a broad range of life histories, developing in soil, water, or vegetation. Many of

these insects engage in aerial activity during reproduction (Jackson and Campbell 1975;

Pathak and Khan 1994; Traugott et al. 2015), which bats specialized in open and edge

spaces are able to exploit. As expected, *M. molossus* showed the highest dietary

taxonomic richness, consistent with its superior hunting abilities in these environments

(Azofeifa et al. 2019). Moreover, edge-space species such as *M. nigricans* and *N.*

furinalis appear to benefit from foraging in rice fields rather than depending on the few

remaining forest patches, even if it means compromising prey richness in their diets.

This opportunistic behavior underscores the importance of rice fields as key foraging

habitats for IB in fragmented landscapes (Aizpurua et al. 2018).

Prey types, trophic breadth, and overlap

Prey type was associated with hunting abilities, bite force, and intrinsic characteristics of

each IB species. As predicted, *M. molossus*—adapted for fast, efficient flight in open

habitats like the studied rice fields (Voigt and Holderied, 2012)—had a diet primarily

composed of fast- and intermediate-flying insects (e.g., moths, beetles), while the two

vespertilionids showed a higher representation of slow-flying insects such as

leafhoppers and some dipterans. Tooth morphology, which ranges from compact in *M.*

nigricans and *N. furinalis* to more robust in *M. molossus* (Mies et al. 1996; Loureiro et al.

2018; Novaes et al. 2022), may reflect a gradient in the contribution of soft- to hard-

bodied prey (Freeman 1998), although, we did not detect this differentiation between *M.*

molossus and the vespertilionid species. Instead, *M. nigricans* diverged more clearly

from *N. furinalis* in the degree of prey sclerotization, likely due to its bite force being

almost six times lower than in the latter species (Aguirre et al. 2002). Bite force therefore

offered a more plausible explanation for the differences observed. However, although *M.*

molossus has a slightly larger skull and somewhat higher bite force than *N. furinalis*,

both species share a reduced third upper molar (Aguirre et al. 2002; Martínez-Coronel

and Hortelano-Moncada 2020; Mies et al. 1996), which may enhance grinding and

account for the similar representation of hard-bodied prey in their diets (Freeman 1998).

Therefore, even though skull structure and bite force probably represent the most

important drivers of hard-bodied prey processing in these bats (Giacomini et al. 2022),

some dental could still play a role on it.

Despite the rich insect community present in these artificial wetlands, we were

able to demonstrate that trophic niches are constrained by species-specific

ecomorphological limitations (Mayberry et al. 2020). For example, although soft-bodied

prey—mainly dipterans— are well represented in the diets of all three species, *M.*

nigricans appears to specialize in them, which is consistent with its slow flight and

relatively delicate skull. While this species can forage in open spaces by adjusting call

bandwidth and duration to improve prey detection (Siemers et al. 2001; Bader et al.

2015), its biomechanical constraints likely limit the range of prey it can efficiently exploit.

These results also support our prediction that *M. molossus* would include a broader

range of prey in its diet, highlighting its trophic flexibility. This is further evidenced by the

24% of individuals (17/71) consuming prey detected once, compared to 11% in M.

nigricans (5/44) and 10% in *N. furinalis* (2/21). Although singleton detections should be

interpreted with caution due to the limitations of DNA metabarcoding (Alberdi and Gilbert

2019), these observations align with the idea that some individuals in generalist species

may differ markedly in prey use (Bolnick et al. 2003), a trend that is receiving increasing

support in bats (Arrizabalaga-Escudero et al. 2019; Magalhães de Oliveira et al. 2020).

In general, our analyses support *M. molossus* as the dominant forager, with low

to moderate overlap in diet and activity time, which begins and peaks earlier than in the

other two species. This molossid began hunting shortly before sunset (but see Esbérard

and Bergallo 2010), coinciding with the rise in insect activity and the presence of some

diurnal raptors—an ecological compromise that slow-flying bats would not efficiently

cope with. The two vespertilionid species coexist with high overlap in both diet and

activity times, although their activity peaks differ— similar to observations of other IB

species reported in previous studies (Lambert et al. 2018; Beilke et al. 2021). These

observations align with Aguiar and Antonini's (2008) findings of high dietary overlap

between these species based on morphological fecal analysis from the Brazilian

Cerrado, and reinforce the idea that subtle ecological differences can facilitate

coexistence even when resource use broadly overlaps. We believe that coexistence is

possible despite higher trophic niche overlap, probably as a consequence of the high

productivity of rice fields in artificial wetlands, where elevated prey abundance

represents an almost unlimited food resource, leading to low competition and high

trophic niche overlap (Krüger et al. 2012).

Comparisons of niche overlap across studies should be made with caution due to

limitations in interpreting dietary data, particularly when using DNA metabarcoding. As

Alberdi and Gilbert (2019) pointed out, Hill numbers offer a more robust framework for

assessing diet diversity and overlap in metabarcoding studies. Direct comparisons using

Pianka's index are therefore not appropriate, as they may overestimate overlap.

Nevertheless, growing molecular evidence continues to reveal trophic flexibility in bats.

Reported values of Pianka's index have ranged from 0.50 to 0.83, in both cryptic

species (Razgour et al. 2011) and inter- or intra-guild comparisons (Salinas-Ramos et

al. 2015; Gordon et al. 2019; Segura-Trujillo et al. 2022).

Concluding remarks

The coexistence between *M. molossus*, *M. nigricans*, and *N. furinalis* in the rice fields

studied in the Venezuelan Llanos possibly reflects a combination of high insect

availability in these artificial wetlands and partial differences in diet and activity time.

Molossus molossus shows a higher incidence of fast-flying, medium-to-highly

sclerotized prey, and its activity times started and peaked significantly earlier than in the

other two bat species. Such divergence across these two trophic niche dimensions

should reduce inter-specific competition and enhance coexistence between *M. molossus*

and the vespertilionids, which agree with niche theory predictions (Finke and Snyder

2008). *Myotis nigricans* and *N. furinalis* are less specialized for foraging in open

environments, and their trophic niches are constrained by species-specific

ecomorphological limitations. Their diets were primarily composed of slow-flying prey

and showed high temporal overlap in activity, though with distinct peak times. A stronger

bite in *N. furinalis*, however, may have allowed for the inclusion of more strongly

sclerotized prey than in *M. nigricans*, suggesting that differences in prey hardness

reduce competition despite similar flight strategies. Although our analyses did not

account for seasonal or cyclical variation in rice cultivation, the non-migratory nature and

year-round presence of these species in the study area suggest that trophic interactions

may remain relatively stable over time. In this context, fine-scale trophic niche

partitioning—facilitated by the humid, prey-rich environment of rice fields—appears to

mediate the coexistence of IB species with marked ecomorphologically differences.

Acknowledgements. Special thanks to José Antonio González-Carcacía (IVIC) for his

support during some fieldwork sessions. Claudia Domínguez (IVIC) and Abimel Moreno

(IVIC) provided assistance in identifying collected insects. Delphine Rioux (LECA) and

Eric Coissac (LECA) assisted with molecular and bioinformatics analyses.

Supplementary Data

Supplementary Data SD1. — Taxonomic identification notes and ecomorphological traits

of the IB species

Supplementary Data SD2. — Prey hardness (order level) and flight type (family level) of

dietary items in three IB species.

Supplementary Data SD3. — Percentage of prey by taxon (Pi) in the diet of three

species of aerial hawking bats at North-western Venezuela's rice crops.

Supplementary Data SD4. —Sequences of insect MOTUs (Inse01_motus and

Arth02_motus) used for the assignment of MOTUs in bat feces

(bat_feces_Inse01.motus and bat_feces_Arth02_motus).

Supplementary Data SD5. — Estimated diversity, trophic niche breadth and dietary

overlap for three IB species.

518 Supplementary Data SD6— GLMs assessing the impact of bat factors (species,

519 functional group, and bite force) on prey characteristics.

References

Acosta LG, Jahnke SM, Redaelli LR, Pires PRS. 2017. Insect diversity in organic rice
fields under two management systems of levees vegetation. *Brazilian Journal of Biology*
77(4):731–744. https://doi.org/10.1590/1519-6984.19615

Adams RA, Thibault KM. 2006. Temporal resource partitioning by bats at water holes.
Journal of Zoology 270(3):466–472. https://doi.org/10.1111/j.1469-7998.2006.00152.x

Agostinelli C, Lund U. 2023. Circular statistics. Version 0.5-0. [R package].
https://CRAN.R-project.org/package=circular

Aguiar LMS, Antonini Y. 2008. Diet of two sympatric insectivores bats (Chiroptera:
Vespertilionidae) in the Cerrado of Central Brazil. *Revista Brasileira de Zoologia*
25(1):28–31. https://doi.org/10.1590/S0101-81752008000100005

Aguiar LMS, Bueno-Rocha ID, Oliveira G, Pires ES, Vasconcelos S, Nunes GL, Frizzas
MR, Togni, PHB. 2021. Going out for dinner—the consumption of agriculture pests by
bats in urban areas. *PLoS One* 16(10):e0258066.
https://doi.org/10.1371/journal.pone.0258066

Aguirre LF, Herrel A, van Damme R, Matthysen E. 2002. Ecomorphological analysis of
trophic niche partitioning in a tropical savannah bat community. *Proceedings of the*
Royal Society of London. Series B: Biological Sciences 269(1497):1271–1278.
https://doi.org/10.1098/rspb.2002.2011

Aizpurua O, Budinski I, Georgiakakis P, Gopalakrishnan S, Ibañez C, Mata V, Rebelo H,
Russo D, Szodoray-Parádi F, Zhelyazkova V, et al. 2018. Agriculture shapes the trophic
niche of a bat preying on multiple pest arthropods across Europe: evidence from DNA
metabarcoding. *Molecular ecology* 27(3):815–825. https://doi.org/10.1111/mec.14474

Alberdi A, Gilbert MTP. 2019. A guide to the application of Hill numbers to DNA-based
diversity analyses. *Molecular Ecology Resources* 19(4):804–817.
https://doi.org/10.1111/1755-0998.13014

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
search tool. *Journal of Molecular Biology* 215(3):403–410.
https://doi.org/10.1016/S0022-2836(05)80360-2

Amador LI, Almeida FC, Giannini NP. 2020. Evolution of traditional aerodynamic
variables in bats (Mammalia: Chiroptera) within a comprehensive phylogenetic
framework. *Journal of Mammalian Evolution* 27(3):549–561.
https://doi.org/10.1007/s10914-019-09475-8

Andriollo T, Michaux JR, Ruedi M. 2021. Food for everyone: differential feeding habits of
cryptic bat species inferred from DNA metabarcoding. *Molecular Ecology* 30(18):4584–
4600. https://doi.org/10.1111/mec.16073

Arrizabalaga-Escudero A, Merckx T, García-Baquero G, Wahlberg N, Aizpurua O, Garin
I, Goiti U, Aihartza J. 2019. Trait-based functional dietary analysis provides a better
insight into the foraging ecology of bats. *Journal of Animal Ecology* 88(10):1587–1600.
https://doi.org/10.1111/1365-2656.13055

Azofeifa Y, Estrada-Villegas S, Mavárez J, Nassar JM. 2019. Activity of aerial
insectivorous bats in two rice fields in the Northwestern Llanos of Venezuela. *Acta*
Chiropterologica 21(1):149–163. https://doi.org/10.3161/15081109ACC2019.21.1.012

Bader E, Jung K, Kalko EKV, Page RA, Rodriguez R, Sattler T. 2015. Mobility explains
the response of aerial insectivorous bats to anthropogenic habitat change in the
Neotropics. *Biological Conservation* 186: 97–106.
https://doi.org/10.1016/j.biocon.2015.02.028

Bartoń K. 2023. MuMIn: Multi-Model Inference. Version 1.47.5. [R package].
https://CRAN.R-project.org/package=MuMIn

Beilke EA, Blakey RV, O'Keefe JM. 2021. Bats partition activity in space and time in a
large, heterogeneous landscape. *Ecology and Evolution* 11(11):6513–6526.
https://doi.org/10.1002/ece3.7504

Bohmann K, Monadjem A, Lehmkuhl Noer C, Rasmussen M, Zeale MRK, Clare E,
Jones G, Willerslev E, Gilbert MTP. 2011. Molecular diet analysis of two African free-
tailed bats (Molossidae) using high throughput sequencing. *PLoS One* 6(6):e21441.
https://doi.org/10.1371/journal.pone.0021441

Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML. 2003.
The ecology of individuals: incidence and implications of individual specialization. *The*
American Naturalist 161(1):1–28. https://doi.org/10.1086/343878

Boyer F, Mercier C, Bonin A, Le Bras Y, Taberlet P, Coissac E. 2016. Obitools: a unix-
inspired software package for DNA metabarcoding. *Molecular Ecology Resources*
16(1):176–182. https://doi.org/10.1111/1755-0998.12428

Clarke KR, Gorley RN. 2015. *PRIMER v7: User manual/tutorial*. Plymouth (UK):
PRIMER-E Ltd.

Esbérard CEL, Bergallo HG. 2010. Foraging activity of the free-tailed bat *Molossus*
molossus (Chiroptera; Molossidae) in southeastern Brazil. *Brazilian Journal of Biology*
70(4):1011–1014. https://doi.org/10.1590/S1519-69842010000500013

Esri. 2024. ArGis Pro. Version 3.4.2. [Computer software]. Redlands (CA, USA):
Environmental System Research Institute, Inc. www.esri.com.

Finke DL, Snyder WE. 2008. Niche partitioning increases resource exploitation by
diverse communities. *Science* 321(5895):1488–1490.
https://doi.org/10.1126/science.1160854

Freeman PW. 1981. Correspondence of food habits and morphology in insectivorous
bats. *Journal of Mammalogy* 62(1):166–173. https://doi.org/10.2307/1380489

Freeman PW. 1998. Form, function, and evolution in skulls and teeth of bats. *Papers in*
Natural Resources 9.

Gardner AL, editor. 2007 [2008]. Mammals of South America. Vol. 1, Marsupials, xenarthrans, shrews, and bats. Chicago (IL, USA): University of Chicago Press.

Giacomini G, Herrel A, Chaverri G, Brown RP, Russo D, Scaravelli D, Meloro C. 2022. Functional correlates of skull shape in Chiroptera: feeding and echolocation adaptations. *Integrative Zoology* 17(3):430–442. https://doi.org/10.1111/1749-4877.12564

Gordon R, Ivens S, Ammerman LK, Fenton MB, Littlefair JE, Ratcliffe JM, Clare EL. 2019. Molecular diet analysis finds an insectivorous desert bat community dominated by resource sharing despite diverse echolocation and foraging strategies. *Ecology and Evolution* 9(6):3117–3129. https://doi.org/10.1002/ece3.4896

Holland RA, Meyer CFJ, Kalko EKV, Kays R, Wikelski M. 2011. Emergence time and foraging activity in Pallas' mastiff bat, *Molossus molossus* (Chiroptera: Molossidae) in relation to sunset/sunrise and phase of the moon. *Acta Chiropterologica* 13(2):399–404. https://doi.org/10.3161/150811011X624875

Hsieh TC, Ma KH, Chao A. 2016. iNEXT: an R package for rarefaction and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution* 7(12):1451–1456. https://doi.org/10.1111/2041-210X.12613

Jackson DM, Campbell RL. 1975. *Biology of the european crane fly, Tipula paludosa Meigen, in western Washington (Tipulidae: Diptera)*. Pullman (WA, USA): Washington State University, College of Agriculture Research Center. Technical Bulletin No. 81.

Kingston T, Francis CM, Akbar Z, Kunz TH. 2003. Species richness in an insectivorous bat assemblage from Malaysia. *Journal of Tropical Ecology* 19(1):67–79. https://doi.org/10.1017/S0266467403003080

Kolkert H, Andrew R, Smith R, Rader R, Reid N. 2020. Insectivorous bats selectively source moths and eat mostly pest insects on dryland and irrigated cotton farms. *Ecology and Evolution* 10(1):371–388. https://doi.org/10.1002/ece3.5901

Krüger F, Harms I, Fichtner A, Wolz I, Sommer RS. 2012. High trophic similarity in the sympatric north european trawling bat species *Myotis daubentonii* and *Myotis dasycneme*. *Acta Chiropterologica* 14(2):347–356. https://doi.org/10.3161/150811012X661666

Lambert CT, Hall LK, Larsen RT, Knight RN, McMillan BR. 2018. Temporal partitioning and the effects of climate change on two ecologically similar desert bats. *Journal of Mammalogy* 99(6):1486–1494. https://doi.org/10.1093/jmammal/gyy113

Li D. 2018. hillR: taxonomic, functional, and phylogenetic diversity and similarity through Hill Numbers. *Journal of Open Source Software* 3(31):1041. https://doi.org/10.21105/joss.01041

Linares O. 1998. *Mamíferos de Venezuela*. Caracas (VE): Sociedad Conservacionista Audubon de Venezuela

Liu Y, Geng Y, Huang Z, Feng J, Jiang T. 2024. Pest suppression services and dietary niche differentiation of bats in Chinese smallholder farming systems: implications for integrated pest management. *Journal of Pest Science* 97(3):1587–1603. https://doi.org/10.1007/s10340-023-01721-6

Loureiro LO, Lim BK, Engstrom MD. 2018. A new species of mastiff bat (Chiroptera,
Molossidae, *Molossus*) from Guyana and Ecuador. *Mammalian Biology* 90(1):10–21.
https://doi.org/10.1016/j.mambio.2018.01.008

Magalhães de Oliveira HF, Camargo NF, Hemprich-Bennett DR, Rodríguez-Herrera B,
Rossiter SJ, Clare EL. 2020. Wing morphology predicts individual niche specialization in
Pteronotus mesoamericanus (Mammalia: Chiroptera). *PLoS One* 15(5):e0232601.
https://doi.org/10.1371/journal.pone.0232601

Marinello MM, Bernard E. 2014. Wing morphology of Neotropical bats: a quantitative
and qualitative analysis with implications for habitat use. *Canadian Journal of Zoology*
92(2):141–147. https://doi.org/10.1139/cjz-2013-0127

Marques JT, Ramos Pereira MJ, Palmeirim JM. 2016. Patterns in the use of rainforest
vertical space by Neotropical aerial insectivorous bats: all the action is up in the canopy.
Ecography 39(5):476–486. https://doi.org/10.1111/ecog.01453

Martínez-Coronel M, Hortelano-Moncada Y. 2020. Nuevos registros de mamíferos en el
Centro de Oaxaca. *Revista Mexicana de Mastozoología (Nueva Época)* 10(2):45–52.
https://doi.org/10.22201/ie.20074484e.2020.10.2.314

Mayberry HW, McMillan MR, Chochinov AV, Hinds JC, Ratcliffe JM. 2020. Potential
foraging niche release in insectivorous bat species relatively unaffected by white-nose
syndrome? *Canadian Journal of Zoology* 98(10):667–680. https://doi.org/10.1139/cjz-
2019-0231

Mercier C, Boyer F, Bonin A, Coissac E. 2013. SUMATRA and SUMACLUST: Fast and
exact comparison and clustering of sequences. Version 1.0.36. [Computer software].
Grenoble (FR): LECA, Université Grenoble Alpes.
https://forge.metabarcoding.org/obitools/sumatra

Meredith M, Ridout M, Campbell LAD. 2024. overlap: estimates of coefficient of
overlapping for animal activity patterns. Version 0.3.9. [R package]. https://CRAN.R-
project.org/package=overlap

Mies R, Kurta A, King DG. 1996. *Eptesicus furinalis*. *Mammalian Species* (526):1–7.
https://doi.org/10.2307/3504316

Moreno A, Azofeifa Y, Domínguez C, Nassar JM. 2020. Ensamblaje de insectos con
actividad nocturna en dos arrozales de los Llanos Noroccidentales de Venezuela.
Anartia 30:35–48.
https://produccioncientificaluz.org/index.php/anartia/article/view/34195

Norberg UM, Rayner JMV. 1987. Ecological morphology and flight in bats (Mammalia;
Chiroptera): wing adaptations, flight performance, foraging strategy and echolocation.
Philosophical Transactions of the Royal Society of London B 316(1179):335–427.
https://doi.org/10.1098/rstb.1987.0030

Novaes RLM, Cláudio VC, Díaz MM, Wilson DE, Weksler M, Moratelli R. 2022.
Argentinean *Myotis* (Chiroptera, Vespertilionidae), including the description of a new
species from the Yungas. *Vertebrate Zoology* 72:1187–1216.
https://doi.org/10.3897/vz.72.e90958

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB,
Solymos P, Stevens MHH, Szoecs E, et al. 2025. Vegan: Community Ecology Package.
Version 2.6-10. [R package]. https://CRAN.R-project.org/package=vegan

Pathak MD, Khan ZR. 1994. *Insect pests of rice.* Manila (PH): International Rice
Research Institute.

Razgour O, Clare EL, Zeale MRK, Hanmer J, Schnell IB, Rasmussen M, Gilbert TP,
Jones G. 2011. High-throughput sequencing offers insight into mechanisms of resource
partitioning in cryptic bat species. *Ecology and Evolution* 1(4):556–570.
https://doi.org/10.1002/ece3.49

Reardon S, Schoeman MC. 2017. Species richness, functional diversity and
assemblage structure of insectivorous bats along an elevational gradient in tropical West
Africa. *Acta Chiropterologica* 19(2):273–285.
https://doi.org/10.3161/15081109ACC2017.19.2.005

Red MapBiomas, RAISG. 2023. Primera colección de mapas anuales de cobertura y
uso del suelo de Venezuela (1985–2022). Caracas (VE): Coordinación General de la
Red MapBiomas. [accessed 2025 Jun 7]. https://mapbiomas.org/coleccion-venezuela

Roemer C, Coulon A, Disca T, Bas Y. 2019. Bat sonar and wing morphology predict
species vertical niche. *The Journal of the Acoustical Society of America* 145(5):3242–
3251. https://doi.org/10.1121/1.5102166

Salinas-Ramos VB., Herrera Montalvo LG, León-Regagnon V, Arrizabalaga-Escudero
A, Clare EL. 2015. Dietary overlap and seasonality in three species of mormoopid bats
from a tropical dry forest. *Molecular Ecology* 24(20):5296–5307.
https://doi.org/10.1111/mec.13386

Segura-Trujillo CA, Álvarez-Castañeda ST, Castañeda-Rico S, Maldonado JE. 2022.
Taxonomic versus ecological prey traits among arthropodophagous bats: implications
for surveying trophic partitioning patterns. *Journal of Mammalogy 103*(5):1071–1083.
https://doi.org/10.1093/jmammal/gyac027

Siemers BM, Kalko EKV, Schnitzler HU. 2001. Echolocation behavior and signal
plasticity in the Neotropical bat *Myotis nigricans* (Schinz, 1821) (Vespertilionidae): a
convergent case with european species of *Pipistrellus*? *Behavioral Ecology and
Sociobiology* 50(4):317–328. https://doi.org/10.1007/s002650100379

Sikes RS, Animal Care and Use Committee of the American Society of Mammalogists.
2016. 2016 Guidelines of the American Society of Mammalogists for the use of wild
mammals in research and education. *Journal of Mammalogy* 97(3):663–688.
https://doi.org/10.1093/jmammal/gyw078

Srilopan S, Mata VA, Bumrungsri S. 2024. DNA metabarcoding reveals insights into diet
partitioning and pest suppression by open space bats in agricultural landscapes. *Acta
Chiropterologica* 26(2):171–187. https://doi.org/10.3161/15081109ACC2024.26.2.003

Taberlet P, Bonin A, Zinger L, Coissac É. 2018. *Environmental DNA: for biodiversity
research and monitoring.* Oxford (UK): Oxford University Press.

Thomas AJ, Jacobs DS. 2013. Factors influencing the emergence times of sympatric
insectivorous bat species. *Acta Chiropterologica* 15(1):121–132.
https://doi.org/10.3161/150811013X667920

Traugott M, Benefer CM, Blackshaw RP, van Herk WG, Vernon RS. 2015. Biology,
ecology, and control of elaterid beetles in agricultural land. *Annual Review of*
Entomology 60:313–334. https://doi.org/10.1146/annurev-ento-010814-021035

Voigt CC, Holderied MW. 2012. High manoeuvring costs force narrow-winged molossid
bats to forage in open space. *Journal of Comparative Physiology B* 182(3):415–424.
https://doi.org/10.1007/s00360-011-0627-6

Wickham H. 2016. *ggplot2: elegant graphics for data analysis*. Springer-Verlag New
York. Version 3.4.4. [R package]. https://ggplot2.tidyverse.org

Williams-Guillén K, Perfecto I. 2011. Ensemble composition and activity levels of
insectivorous bats in response to management intensification in coffee agroforestry
systems. *PLoS One* 6(1):e16502. https://doi.org/10.1371/journal.pone.0016502

Wilson DE, LaVal RK. 1974. *Myotis nigricans*. *Mammalian Species* 39:1-3.
https://doi.org/10.2307/3503847

Yunus M, Nasir B, Lasmini SA, Piri RL. 2022. Biodiversity and community structure of
arthropod in tropical rice fields under organic and conventional ecosystems. *Australian*
Journal of Crop Science 16(4):531–535. https://doi.org/10.21475/ajcs.22.16.04.p3551

Figure Legends

Fig. 1. Map of Acarigua and Turén rice farms in the Northwestern Llanos of Portuguesa,

Venezuela. The base map includes the 2014 land-cover layer from Red MapBiomas and Red

Amazónica de Información Socioambiental Georreferenciada (RAISG).

Fig. 2. Percentage of prey in the diet of three species of aerial-hawking bats, classified by (A)

order, and (B) family.

Fig. 3. Percentage of prey in the diet of three species of aerial-hawking bats, classified by (A)

degree of sclerotization, and (B) flight behavior.

Fig. 4. Hellinger-transformed PCA of dietary incidence (presence–absence of 65 MOTUs) in

three species of aerial-hawking bats. Pterygota was excluded from the analyses, reducing the

MOTU count by one (64 MOTUs).

Fig. 5. Time-density curves of bat species pairs with gray-shaded overlap areas, based on data

from both farms. Bootstrap estimates and 95% confidence intervals for the overlap coefficient, Δ,

are presented on the axis-x. The overlap coefficient $\Delta 4$ is used for sizes greater than 75.

Fig. 1. Map of Acarigua and Turén rice farms in the Northwestern Llanos of Portuguesa,

Venezuela. The base map includes the 2014 land-cover layer from Red MapBiomas and Red

Amazónica de Información Socioambiental Georreferenciada (RAISG).

Fig. 2. Percentage of prey in the diet of three species of aerial-hawking bats, classified by (A) order, and (B) family.

Fig. 3. Percentage of prey in the diet of three species of aerial-hawking bats, classified by (A)

degree of sclerotization, and (B) flight behavior.

780   Fig. 4. Hellinger-transformed PCA of dietary incidence (presence–absence of 65 MOTUs) in

781   three species of aerial-hawking bats. Pterygota was excluded from the analyses, reducing the

782   MOTU count by one (64 MOTUs).

785   Fig. 5. Time-density curves of bat species pairs with gray-shaded overlap areas, based on data

786   from both farms. Bootstrap estimates and 95% confidence intervals for the overlap coefficient, Δ,

787   are presented on the axis-x. The overlap coefficient Δ4 is used for sizes greater than 75.