1 **Two different and robustly modeled DNA binding modes of Competence Protein ComP**
2 **- systematic modeling with AlphaFold 3, RoseTTAFold2NA, Chai-1 and re-docking in**
3 **HADDOCK**
4

Stian Aleksander Helsem[1], Kristian Alfsnes[2], Stephan A. Frye[3], Alexander Hesselberg

Løvestad[1,4] and Ole Herman Ambur[1]*

[1]Department of Life Sciences and Health, Faculty of Health Sciences, OsloMet, Norway

[2] Division for Infection Control, Department of Bacteriology, Norwegian Institute of Public

Health, 0213 Oslo, Norway

[3] Division of Laboratory Medicine, Dept. Microbiology, Oslo University Hospital, Norway

[4]Department of Virology, Norwegian Institute of Public Health, Oslo, Norway

*Correspondence: olam@oslomet.no

Keywords: Competence protein; ComP; protein-DNA binding mode; minor pilin; DNA

Uptake Sequence; DUS; natural competence; transformation; interface; docking; AlphaFold

3; RoseTTAFold2NA; Chai-1; HADDOCK; PAE; pLDDT; DockQ; CPPM; Neisseriaceae;

Neisseria meningitidis; Neisseria cinerea; Neisseria mucosa; Neisseria subflava; Eikenella

corrodens; Kingella denitrificans; Bergeriella denitrificans

Abstract

The competence protein ComP is a Type IV minor pilin and the extracellular DNA binding protein involved in natural transformation in the human pathogens *Neisseria gonorrhoeae*, *Neisseria meningitidis, Eikenella corrodens* and related Neisseriaceae bacteria. Details of the DNA binding mode of ComP is enigmatic, and the 3D structure of the DNA :: protein complex remains unresolved. Here we characterize the ComP orthologs in a set of Neisseriaceae family members, model their common structural domains and their interaction with different preferred 12 base pair long DNA binding motifs, DNA Uptake Sequences (DUS) and scrambled versions of these. Through systematic in silico modeling using AlphaFold 3, RoseTTAFold2NA, and Chai-1 and model comparisons, we bring a new understanding of the interactions between DNA and ComP. We report six distinct binding modes of which two, here named Epsilon and Gamma, were robustly modeled across platforms and different ComPs. The characteristics and robustness of the predicted models and DNA binding modes from each tool are assessed and discussed. This work expands the knowledge on the ComP :: DUS interaction and guides further wet- and dry-lab systematic and experimental characterization of these complexes through which molecular and clinical interventions may be developed.

Introduction

The development of deep learning protein structure prediction algorithms is now allowing the study of protein function and interactions in a detail never seen before. AlphaFold3 (AF3) (Abramson et al., 2024), RoseTTAFold2NA (RF2NA) (Baek et al., 2024), RosETTAFold-All-Atom (Krishna et al., n.d.), NeuralPLexer (Qiao et al., 2024), DiffDock (Corso et al., 2022) and more have led the way to model protein interactions with other proteins, ligands and nucleic acids. Recently Chai-1 was released based with single model training and new functionality (Chai Discovery, 2024) and new implementations of AlphaFold 3 (AF3) are continuously being developed, with the release of HelixFold3 (Team, 2024) and Boltz-1 (Wohlwend et al., 2024) and the field is rapidly advancing. The underlying platform algorithms differ in fundamental ways and therefore also their emphasis on various confidence metrices. The Predicted Aligned Error (PAE) was introduced with AlphaFold and AF3 places a strong emphasis on PAE for providing information regarding confidence of predicted spatial relationships between residues and is used together with different Local Predicted Confidence Difference Test (LDDT) scores (Abramson et al., 2024). PAE is reported by default in AF3. RF2NA filters models based on a PAE < 10 threshold as acceptable but does not emphasize PAE-resolution below this this cut-off and like AF3, reports confidence from Local Distance Difference Test (lDDT) relative to available experimental structures and Predicted Local Distance Difference Test (pLDDT) as the model-intrinsic confidence score at the residue level (Baek et al., 2023). Chai-1 also applies per-residue and chain interface confidence metrics (LDDT and iLDDT) yet also integrates whole structure reliability from DockQ scoring (Chai discovery, 2024). In having an emphasis on PAE, AF3 also applies a Chain Pair PAE Minimum (CPPM) metric to assess the confidence

of the interaction between two chains, such as DNA and protein, within the complex and

propose the metric useful to distinguish binding from non-binding interactions (Alpha Fold

Server). Here, we explore the hypothesis that using fundamentally different platforms to

model the structure of an unresolved DNA :: Protein complex may educate our understanding

of their specific interaction and potentially provide a modeling consensus.

Natural transformation is the uptake and recombination of DNA and is an evolved

mechanism of many Gram positive and Gram negative bacteria (reviewed in Dubnau &

Blokesch, 2019). Competent bacteria able to use natural transformation to acquire DNA

display a range of different adaptations and competence states. Horizontal gene transfer

mechanisms facilitate the spread of advantageous traits including antibiotic resistance and

virulence genes and are therefore of clinical interest (Arnold et al., 2022).Transformation in

the Neisseriaceae family containing human pathogens *Neisseria gonorrhoeae*, *Neisseria*

meningitidis and *Eikenella corrodens* together with opportunistic and commensal relatives is

enhanced when short 10-12 nucleotide DNA binding motifs (e.g. ATGCCGT<u>CTG</u>AA in *N.*

meningitidis/ gonorrhoeae) are present in the transforming DNA (reviewed in Ambur et al.,

2016). These motifs are named DNA Uptake Sequences (DUS) and are employed as standard

practice for genetic manipulation of *Neisseria* species (Dillard & Chan 2024), for rapid

detection of *N. gonorrhoeae* using DUS-targeting gold nanoparticles (Carter et al., 2022) and

for molecular typing of *Kingella kingae* (Basmaci et al., 2014). DUS are genomically

enriched in permissive parts of their respective genomes (Treangen et al., 2008) and shown *in*

vivo to facilitate DNA mobility within species or between closely related genera (Frye et al.,

2013). Extracellular DNA containing multiple DUS has been shown trapped at the peripheral

edges of gonococcal colonies where it readily transforms (Bender et al, 2022). *In vitro*

acrylamide EMSA and surface plasmon resonance experiments using purified ComP, DUS-

DNA and competitive DNA have shown that ComP preferentially binds it cognate DUS with

high affinity and that this is an inherent property of ComP homologs (Cehovin et al., 2013;

Berry et al., 2013; 2016). Eight dialects of DUS have been described, all of which have a

conserved Watson strand CTG core essential for DUS-mediated transformation by means of

the competence protein ComP (Frye et al., 2013; Berry et al., 2013). In an evolutionary

perspective, DUS provides means to acquire homologous DNA for allelic re-shuffles (Ambur

et al., 2016).

ComP is a minor type IV pilin first identified as a positive effector of DUS-mediated

transformation in *N. gonorrhoeae* (Aas et al., 2002) and later identified as the DNA-binding

DUS receptor in *Neisseria subflava* and other *Neisseria sp.* (Cehovin et al., 2013). *In vivo*

overexpression of *N. subflava* ComP in a *N. meningitidis* model has been shown to increase

transformation also in a DUS-independent manner almost up to the level of DUS-specific

transformation (Berry et al., 2013). At least two different DNA binding modes of ComP may

therefore exist, one DUS-specific and one or more unspecific. Alternatively, DUS-specific

and unspecific DNA binding may share the same singular binding mode assuming only weak

DUS sequence binding biases with more relaxed interactions to parts of the DUS outside the

essential CTG core in accordance with wet-lab experiments (Berry et al., 2013; Frye et al.,

2013). ComP is integrated into the type IV pilus (T4P) protruding from the neisserial outer

membrane together with the major pilin PilE and other minor pilins (Berry et al., 2016). All

the minor pilins of T4P are structurally similar at the α-helical N-terminus required for pilin

multimer assembly (reviewed in Jacobsen et al., 2020). Relative to other minor pilins, ComP

has a unique, complex and exposed C-terminal globular domain where DNA can bind. The

high-resolution 3D structures of ComP (*N. meningitidis* and *N. subflava*) have been reported

and are based on crystallography of soluble ComP-maltose-binding-protein fusion proteins

and NMR of 6-His tagged ComP (Berry et al., 2016). ComP$_{N.men}$ and ComP$_{N.sub}$ are

structurally very similar and are characterized by an N-terminal α-helix, five antiparallel β-

strands, the α1-β1- and β1-β2-loops, a DD-region, and two disulfide bridges (Cys-Cys).

Intramolecular disulfide bridges are formed in exported proteins by thiol-disulfide

oxidoreductases (DsbAs) and wet-lab experiments have shown that *N. meningitidis* DsbA

knock-out mutants have reduced DNA uptake in a DUS-dependent manner (Sinha et. al,

2008). Whether this phenotype could directly link to ComP is unknown. The two ComP

disulfide bridges staples the DD-loop to the β1-strand and the β4-β5 strands together, were

found crucial for DUS-binding in electrophoretic mobility shift assays (Cehovin et al., 2013).

ComP has an electropositive stripe, and individual substitutions of three positively charged

amino acids (R78A, K94A, and K108A) were shown *in vivo* to affect binding and/or

transformation negatively (Cehovin et al., 2013). Several Neisseriaceae have been

demonstrated to have higher transformability and binding affinity for DUS containing DNA

compared to other sequences using transformation experiments and biophysical approaches

(Goodman & Scocca, 1988; Aas et al., 2002, Frye et al., 2013; Berry et al., 2013, 2016;

Cehovin et al., 2013). NMR spectra showed that the DD-region, the tip of the ß1-ß2 loop and

part of the α1-ß1 loop were involved in DUS binding and were modeled to establish contacts

with bases in successive grooves of the dsDNA (Berry et al., 2016). ComP has been found to

preferentially mediate transformation of dsDNA, although single-stranded Crick strand DUS

(TTCAGACGGCAT) shows some transforming ability in a *N. gonorrhoeae* model relative to

no-DUS and the Watson strand DUS (ATGCCGTCTGAA) (Duffin & Seifert, 2012).

Another study exploring uptake and transformation of fluorescently labelled ssDNA and

dsDNA found similarly that efficient DUS recognition was dependent on double-stranded

DNA whereas the rates of transport across the plasma membrane were comparable (Hepp et

al., 2016).

The ComP 3D structure with bound DUS/DNA remains unresolved. All major structural

work has focused on the ComP proteins of *N. meningitidis* and *N. subflava* (Berry et al.,

2013, 2016; Cehovin et al., 2013). Fewer efforts have been put into studying the ComP ::

DUS/DNA binding mode in silico. Berry et al. (2016) and Hughes-Games (2020) performed

docking analyses on *N. subflava* NJ9703$_{ComP}$ (here N.sub$_{nat}$) and *N. meningitidis* 8013$_{ComP}$

(here N.men$_{nat}$). Their molecular docking results both seem to show (our interpretation) that

the α1-β1 and β1-β2 loops interact with major groove (MG) and the DD-region with minor

groove (mG) of the DNA.

No systematic modeling of ComP with DUS/DNA has been performed using the now

available and fundamentally different deep learning-based platforms for structure prediction

of such complexes. AF3, RF2NA and Chai-1 represent the state-of-the-art of modeling

protein-nucleic acid (protein-NA) complexes and may hence provide means to better

understand the ComP :: DUS/DNA interaction. Using five of the DUS dialects from Frye et

al. (2013) as the basis for comparatively modeling ComP :: DUS/DNA interactions using

different modeling algorithms, this study brings a new understanding of the ComP ::

DUS/DNA complex and characterize two different and robust DNA binding consensuses.

Materials and methods

Data material

ComP orthologs were obtained using a series of BLASTp searches (Camacho et al., 2009),

using the structures with PDB accession 2NBA (Berry et al., 2013), 2M3K (Cehovin et al.,

2013) and 5HZ7 (Berry et al., 2013) as queries against the RefSeq non-redundant database in

the NCBI web server, specifying the Neisseriaceae clade as target. GenBank/RefSeq ComP

primary structure and accession numbers are listed in Table S1. Trimming of the primary

ComP structure removed 28 amino acids of the N-terminal part of the mature ComP$_{Nmen}$ and

the equivalent sections in the other six orthologous ComPs as they were trimmed in the

published 3D structures (Cehovin et al., 2013; Berry et al., 2013). ComP primary structures

from seven Neisseriaceae species were modeled with dsDNA containing their native 12 bp

DUS dialects in this study (Table S1). Native DUS dialects refer to the genomically most

abundant 12-mer DUS of each species as identified and described previously (Ambur et al.,

2007; Frye et al., 2013). The ComP :: DUS pairs were: *Eikenella corrodens* ATCC 23834

ComP :: AG-eikDUS (E. cor$_{nat}$), *Neisseria meningitidis* MC58 ComP :: AT-DUS (N. men$_{nat}$),

Kingella denitrificans ATCC 33394 ComP :: AA-king3DUS (K. den$_{nat}$), *Neisseria mucosa*

ATCC 25996 ComP :: AG-mucDUS (N. muc$_{nat}$), *Bergeriella denitrificans* ATCC 33394

ComP :: AG-DUS (B. den$_{nat}$), *Neisseria cinerea* ATCC 14685 ComP :: AT-DUS (N. cin$_{nat}$)

and *Neisseria subflava* NJ9703 ComP :: AG-DUS (N. sub$_{nat}$). These native ComP :: DUS

modeling complexes are described with the suffix "nat" appended to the species name (e.g.

N. sub$_{nat}$). Equivalently when using scrambled native DUS in ComP :: DNA modeling the

suffix "scr" was appended to the species name (e.g. N.sub$_{scr}$).

Modeling of ComP$_{nat}$s

To model ComP$_{nat}$s for the first time, seven different ComPs were modeled with their native

12 bp DUS (dsDNA) in three different deep-learning structure prediction platforms: AF3

(web server), RF2NA (local installation) and Chai-1 v. 0.2.0 (local installation). The

flowchart followed is presented in Fig 1.

Figure 1. Flowchart of the study showing the steps taken to model native $ComP_{nat}$ with Chai-

1, RF2NA and AF3 and re-docking of top-ranking RF2NA models in HADDOCK. AF3

modeling of $ComP_{scr}$ follow dashed lines on the right side. Boxes with inputs are white,

processes light gray and outputs dark grey. All data sources and parameters used are

described in the text.

A minimum of 20 native AF3 replicates (33 for $B.\ den_{nat}$ and 37 for $N.\ sub_{nat}$) and 100 native

runs with different seeds in RF2NA and Chai-1 were performed for each $ComP_{nat}$.

Customization of multiple sequence alignment (MSA) input to Chai-1 allowed for applying

also an MSA of DUS in addition to the ComP MSA, the details of the different MSA

schemes explored are listed in Table S2. These MSAs were generated using HHblits v. 3.0.0

(Steinegger, Meier, et al., 2019) to search the Big Fantastic Database (BFD) (Jumper et al.,

2021; Steinegger, Mirdita, et al., 2019; Steinegger & Söding, 2018) and Uniref30 (Mirdita et

al., 2022) databases for ComP MSAs, using the same HHblits parameters as in RF2NA. DUS

MSAs were created based on genomic DUS counts in the seven selected species (as in Frye et

al., 2013). 10 replicates were run for each MSA scheme. This generated a grand total of 3500

mmCIF/PDB models in Chai-1 (500 per $ComP_{nat}$), 850 models in AF3, with more than 100

models per $ComP_{nat,}$ and 700 models in RF2NA, with 100 models per $ComP_{nat}$. Criteria for

assigning successful models were ipTM > 0.6 in AF3 and Chai-1 and PAE < 10 in RF2NA,

following cut-off recommendations on the AF3 web server for ipTM and in Baek et al.

(2024) for PAE (Table 1).

Table 1. Model quality measures and confidence metrices used to assess predicted models

from Chai-1, AF3, RF2NA and HADDOCK.

Metrics	Score/test description	Range	Unit	Applied and reported	Relevant platform(s)
ipTM	Predicted interface template modeling score	0-1 Poor-excellent	-	≥ 0.6 considered successful	AF3 Chai-1
PAE	Predicted aligned error	Continuous	Å	< 10 considered successful in RF2NA	AF3 Chai-1 RF2NA
pLDDT	Predicted local distance difference test	0-100 Very poor-very high confidence	-	Significant differences between models	AF3 Chai-1 RF2NA
CPPM*	Chain pair PAE minimum	Continuous	Å	Significant differences between models	AF3
HADDOCK score	Weighted sum of various energy terms	Continuous	-	Normal mode	HADDOCK
DockQ**	Quality measure of similarity	Continuous	-	0-1 Poor-excellent	AF3, Chai-1 and RF2NA

* Used here to distinguish probable binders from non-binders in interactions between chains
in AF3

** Used here to measure consistency and mode convergence in predicted ComP$_{nat}$ models
within and between AF3, Chai-1 and RF2NA (Basu & Wallner, 2016; Mirabello & Wallner,
2024).

To investigate the confidence of AF3 to model the seven ComP$_{nat}$s in a DUS-specific manner

as measured by CPPM, AF3 modeling using also scrambled DUS sequences was carried out.

All seven ComPs were modeled with scrambled 12 bp versions of their native DUS dialects

in AF3, not allowing the scrambled DNA strings to contain the conserved trinucleotide 5'-

CTG-3' nor tetra-homomers (i.e. "GGGG") in the sequence, while containing the same

nucleotide content as the native DUS. Wilcoxon rank sum significance tests were then

performed to determine whether runs with native DUS yielded higher confidence metrics (see

below) than scrambled DUS runs. All 140 AF3 generated ComP$_{scr}$s models were assigned

binding mode as for ComP$_{nat}$s described below.

To evaluate the modeling reproducibility of the different binding modes while also assessing

the docking approaches used in previous studies (Berry et al., 2016; Hughes-Games, 2020),

re-docking experiments of RF2NA outputs were conducted in HADDOCK (Teixeira et al.,

2024). The top-ranking RF2NA ComP$_{nat}$s were re-docked. Following the suggestion by

(Esmaeeli et al., 2023), prior to molecular docking in HADDOCK, the first 10 all-atom non-

trivial normal modes with lowest frequency for both the ComPs and DUS were generated

using the R module bio3d (Grant et al., 2006, 2021; Skjærven et al., 2014; 2016). Selected

normal mode conformations were fed into HADDOCK as ensemble protein and DNA PDB

files and cross-docked (all against all), generating 400.000 rigid bodies per ComP$_{nat}$, of

which, after flexible and molecular dynamics refinement, the top 200 models (HADDOCK

score) were assigned binding mode. Because the input DNAs were highly distorted (due to

normal mode calculation), it was often challenging to discern DNA grooves (mG and MG)

and assign binding mode in the re-docking outputs. Binding modes for all HADDOCK

models were therefore very cautiously considered. This was also why the HADDOCK output

models were not used in the cross-platform DockQ consistency analyses. For N. sub$_{nat}$, all

solvent-accessible ComP residues that showed an NMR CSP shift below DNA concentration

of 20 mM in Berry et al., 2016 were specified as ambiguous restraints. For the other six

ComP$_{nat}$s, the same residues were used as ambiguous restraints only if conserved in the

ComP alignment (Fig S2).

All Chai-1 and AF3 models having ipTM $\geq$ 0.6 and RF2NA models having PAE < 10 (ipTM

is not an output in RF2NA) were deemed robust and inspected in 3D using PyMOL v. 3.0.0

and binding mode assigned according to the descriptions below. In ComP$_{nat/scr}$ models for

which ipTM never reached the 0.6 cut-off, all models were inspected in 3D and assigned

binding mode. Based on similarity across models in how central protein domains and DNA

grooves interact, general binding modes were defined (Table 2).

Table 2. ComP DNA binding modes. Modes are differentiated and assigned by how four

central ComP protein domains interact and enter the DNA minor- (mG) and major (MG)

grooves of DNA.

ComP domain/ Binding mode	α1-β1 loop	β1-β2 loop	DD-region	β4- and β5-sheets	Comment
γ (Gamma)	mG	mG	MG		
δ (Delta)	mG	mG	mG		
ε (Epsilon)	MG	MG	mG		
ζ (Zeta)	MG	MG	MG		
η (Eta)*				MG	N-terminus and DNA end close
θ (Theta)**	MG	MG	MG		DNA partly unwound
o (Other)	-	-	-	-	Unlike rest

254 * Only in HADDOCK for K. den$_{nat}$, B. den$_{nat}$ and N. men$_{nat}$. η part of o in Table 3.
255 **Only in Chai-1 for N. cin$_{nat}$
256

Representative examples of ComP$_{nat}$ complexes in the six different binding modes Gamma,

Delta, Epsilon, Zeta, Eta and Theta are shown in Fig S1 and representative PAE-plots for

each ComP$_{nat}$ assigned modes in Fig S4-S6. Since PAE plots were not generated by default in

either Chai-1 nor RF2NA, these were made using customized scripts.

Internal consistency and cross-platform convergence

To investigate intra-platform consistency of AF3, Chai-1 and RF2NA results separately, all

$ComP_{nat}$ against all $ComP_{nat}$ DockQ scores were computed for the top-ranking models across

replicate runs within the same platform. This was devised as a test of platform consistency for

repeatedly predicting the same binding mode. Additionally, the degree of agreement in the

results across AF3, Chai-1 and RF2NA was quantified by comparing the DockQ scores for

all robust models against all other robust models. To avoid giving weights to lower ranking

models, only AF3 and Chai-1 models having ipTM $\geq$ 0.6 were considered robust in the cross-

platform DockQ comparisons as advised (Abramson et al., 2024; Zhang & Skolnick, 2004),

except for N. men_{nat} and N. cin_{nat} in AF3 and N. muc_{nat} in Chai-1, for which ipTM $\geq$ 0.6 was

never reached and all models were included in these cases. Likewise, since ipTM is not

reported in RF2NA only those RF2NA models having mean interface PAE < 10 were

considered robust and included in the assessment as advised by Baek et al., 2024.

Results

The conserved ComP structure

Here, using AF3, the published structures of two ComP proteins ($ComP_{N.men}$ and $ComP_{N.sub}$)

(Berry et al., 2016; Cehovin et al., 2013) and five additional orthologous proteins from other

Neisseriaceae members (*B. denitrificans*, *E. corrodens*, *N. mucosa*, *N. cinerea* and *K.*

denitrificans) were aligned and compared (Fig 2).

Figure 2.   Structural overlay of ComPs from seven Neisseriaceae species. These are AF3

models aligned on the α-carbons of the model with the best ipTM score, Purple: $ComP_{K.den}$;

Green: $ComP_{N.sub}$. Turquoise: $ComP_{B.den}$. Yellow: $ComP_{N.muc}$. Pink: $ComP_{E.cor}$. Grey:

ComP$_{N.men}$. Blue: ComP$_{N.cin}$. The β-strands are numbered, and domains labelled. The three

representations are 90° right turns around the N-terminal α-helix aligned on the y-axis.

The structural alignment of the AF3 models revealed a high degree of similarity across all

seven ComPs and that all contained the same domains as previously described (Berry et al.,

2016; Cehovin et al., 2013). Identifiable and ubiquitous domains are the N-terminal α-helix,

five antiparallel β-strands, β-strand connecting loops, the DD-region, and two essential

disulfide bridges (connecting β1-DD-loop and β4-β5) (Berry et al., 2016). Minor differences

were seen across the ComPs, including prediction of a short, additional α-turn embedded in

the β1-β2-loop in ComPs of *B. denitrificans* and *E. corrodens* and a short α-turn located in

the DD-region in ComP of *N. mucosa*.

Minor structural variations in different domains for the same ComP were observed across

different replicate runs, even within the same platform. Particularly, parts of the DD-region

and the β1-β2 loop of ComP were modelled generally less consistently. In the modeled

structures in Fig 2, these flexible regions can be seen as imperfectly aligned domains

(separated protein strands). These are located in indel-variable regions of the investigated

ComPs, as shown in their aligned primary structure (Fig S2).

Due to strong overall structural predicted consistency across platforms, ComP structure

overlays from Chai-1 and RF2NA are not shown beyond their respective ComP$_{nat}$s in Figure

3 below, and their representative ComP$_{nat}$ PAE plots in the supporting data (Figs S4-S7). We

note however that despite overall structural consistency, RF2NA generally produced models

lacking the disulfide bridge connecting the β1-strand and the DD-region even though these

were present in all three experimentally predicted template structures commonly used in the

MSAs (2NBA, 5HZ7 and 2M3K). Furthermore, a smaller fraction of the RF2NA models,

also the disulfide bridge connecting β4- and β5-sheets were lacking which could affect the

integrity of the β-sheet. In contrast to all AF3 and Chai-1 models, RF2NA was therefore

found less prone to maintain modeled and functionally essential disulfide bridges in ComP.

Modeling and re-docking results

A series of iterated modelling experiments of all seven $ComP_{nat}s$ were conducted in all three

platforms, RF2NA, Chai-1 and AF3 to provide a grand total of 3620 $ComP_{nat}$ model

complexes. These $ComP_{nat}$ models are shown in Fig 3 as platform specific complex overlays

and their binding modes are detailed in Table 3 together with 1386 models from the re-

docking in HADDOCK.

Figure 3. Robust $ComP_{nat}$ models from RF2NA, Chai-1 and AF3 for all seven $ComP_{nat}$

complexes (A-G). **A**: $N.sub_{nat}$; **B**: $B.den_{nat}$; **C**: $K. den_{nat}$; **D**: $N. muc_{nat}$; **E**: $E.cor_{nat}$; **F**: $N. cin_{nat}$;

and **G**: $N. men_{nat}$. PAE and pLDDT score ranges of models included in the overlay are listed

next to each $ComP_{nat}$ overlay and in addition ipTM score ranges for AF3 and Chai-1 models.

Each overlay is aligned on the α-carbon of the models with the best ipTM score or lowest

PAE for RF2NA models. For each $ComP_{nat}$ modeled in AF3 and Chai-1, only the subset of

models with ipTM ≥ 0.6 are represented and when no models were predicted above the ipTM

threshold (Fig 3 F, G and D and Fig 6 A), all predicted models are represented and labeled

with *. For RF2NA, only models having PAE ≤10 are represented.

Tool	AF3					Chai-1					RF2NA*					HADDOCK				
Mode/ ComP$_{nat}$	☐	ζ	γ	δ	o	☐	ζ	γ	δ	o	☐	ζ	γ	δ	o	☐	ζ	γ	δ	o
N.sub$_{nat}$	258	-	-	-	-	428	-	-	-	-	-	-	1	45	-	26	-	159	-	15
B.den$_{nat}$	165	-	-	-	-	91	-	-	-	-	20	-	1	-	1	145	-	1	-	54
K.den$_{nat}$	146	-	-	-	-	445	-	-	-	-	25	69	4	-	-	200	-	-	-	-
N.muc$_{nat}$	105	-	-	-	-	*500	-	-	-	-	-	-	10	-	1	145	-	53	-	2
E.cor$_{nat}$	-	-	95	-	-	157	-	138	-	86	23	25	26	1	6	‡170	‡18	-	-	-
N.cin$_{nat}$	4	-	*96	-	-	-	-	-	-	†395	-	-	40	12	-	79	-	121	-	-

N.men$_{nat}$	*21	-	*76	*3	-	12	1	-	-	-	-	-	56	35	-	130	2	68	-	-
SUM	699	0	267	3	0	1633	1	138	-	481	68	94	136	93	7	895	20	402	-	69

*ipTM < 0.6 or ipTM not available (RF2NA)

† All in Theta mode

‡ For E. cor$_{nat}$ in HADDOCK, 200 docking solutions was not reached and 188 final docking solutions were found

Table 3. Total counts of ComP$_{nat}$ models from all three platforms AF3, Chai-1, RF2NA and re-docking in HADDOCK with assigned binding

modes. Greek letter symbols for modes are = Epsilon, **ζ** = Zeta, **γ**=Gamma, **δ**=Delta and **o**=other (here incl. Eta and Theta)

To investigate the modeling quality of each ComP$_{nat}$ prediction more closely, the PAE and

pLDDT distributions of each modeled complex by RF2NA, AF3 and Chai-1 are shown as

scatter plots per mode and platform in Fig 4A and subdivisions of the scatters within the

dominating Epsilon and Gamma modes per ComP$_{nat}$ in the B and C panels, respectively.

Figure 4.   **A**. PAE/pLDDT scatter plots or all binding modes of all ComP$_{nat}$s from AF3

(red), Chai-1 (blue) and RF2NA (green). **B**. Venn diagram of the resolved Epsilon scatters in

A per color coded ComP$_{nat}$ species. Modeling platforms are represented with symbols as

assigned in the right panel. **C**. Venn diagram of the resolved Gamma models in A per color

coded ComP$_{nat}$ species. Modeling platforms are represented with symbols as assigned in right

panel.

To further systematically explore AF3, Chai-1 and RF2NA mode convergence and quality of

each ComP$_{nat}$, all models with assigned modes from all platforms were paired and compared

directly using DockQ scores. The DockQ scores were categorized as higher or lower

according to the 0.23 (low), 0.49 (intermediate) and 0.8 (high) limits and the results are

shown as violin plots in Figure 5.

Figure 5.   Violin plots of DockQ scores for paired all against all model comparisons of

each ComP$_{nat}$ across platforms. The median and interquartile ranges are shown as horizontal

bars, and paired model comparisons are depicted as swarms inside the violins. Paired inter-

platform models which were assigned to two different binding modes are shown as

"Different". Horizontal red, yellow and green dashed lines represent DockQ of 0.23 (low),

0.49 (intermediate) and 0.8 (high), respectively. Red violins represent inter-platform model

pairs in Epsilon, blue Gamma, light grey Other, dark grey Delta and green Different.

General modeling performance

The distribution of AF3 models across modes covered the widest quality PAE/pLDDT score

ranges of the three platforms with PAE range 3-6 Å and pLDDT 78 (lowest)–93 (highest)

(Fig 4A). The distribution of Chai-1 models covered similar yet narrower ranges as AF3 with

PAE range 2.6 (lowest)–4.5 and pLDDT range 82-89. The RF2NA distributions showed a

systematically higher PAE range, 8-10, than AF3 and Chai-1 across modes, whilst the

pLDDT range 80-86 overlapped those of the other two platforms. The highly divergent PAE

distributions show how confidence from PAE is differentially emphasized in RF2NA relative

to AF3 and Chai-1. PAE, by not representing a universal metric were therefore very

cautiously considered. PAE may therefore be better suited for intra-platform confidence

scoring than between RF2NA and the other two platforms or in combination with other

metrics such as pLLDT and DockQ.

When assessing the internal model consistency of AF3, Chai-1 and RF2NA ComP$_{nat}$s using

DockQ scores, both AF3 and Chai-1 showed higher model consistency across replicate runs

of all ComP$_{nat}$s than RF2NA. AF3 showed a significantly higher median DockQ score

(0.841) compared to the other two platforms (Chai-1: 0.680; RF2NA: 0.617), as determined

by pairwise Wilcoxon rank-sum tests (Code Output S3). Also, Chai-1 showed a significantly

higher median DockQ score compared to RF2NA.

The DNA binding modes

The only two modes predicted by all three platforms and in the re-docking experiments in

HADDOCK were Epsilon (3295/5005) and Gamma (953/5005) (Table 3, Fig S1). Overall,

AF3, Chai-1 and HADDOCK predicted more Epsilon and RF2NA more Gamma. N.sub$_{nat}$,

B.den$_{nat}$, K.den$_{nat}$ and N.muc$_{nat}$ were more consistently predicted in Epsilon and E.cor$_{nat}$,

N.cin$_{nat}$ and N.men$_{nat}$ in Gamma. The Zeta mode (115/5005), which resembles Epsilon (Table

2), was not predicted by AF3 and mostly by RF2NA. The Delta mode (93/5005), which

resembles Gamma, was not predicted by Chai-1 and mostly by RF2NA. "Other" modes were

not predicted by AF3. The unique Theta mode (359/5005) was only predicted by Chai-1,

which permits entry of additional MSA DNA schemes. HADDOCK also found strong

general support for the Epsilon and Gamma modes and deviate from the three modeling

platforms most notably in assigning Gamma for N.sub$_{nat}$ (only 1 recorded in RF2NA), a

strong Epsilon bias for E. cor$_{nat}$ and a bias for Epsilon for N. men$_{nat}$.

All seven ComP$_{nat}$s were modeled in Epsilon in either AF3, Chai-1 or both. The AF3 and

Chai-1 models formed clusters of models with PAE range 2.5-6 Å and pLDDT range 78-93

(Fig 4B). The AF3 Epsilon models covered a wider range (strongest and weakest) than Chai-

1 Epsilon regarding both measures and included more ComP$_{nat}$s. Fig 4B shows an

overlapping region where models from four ComP$_{nat}$s (N.sub$_{nat}$, B.den$_{nat}$, K.den$_{nat}$, E.cor$_{nat}$)

converge on highly similar complexes modeled by either AF3 and Chai-1 or both (B.den$_{nat}$).

All RF2NA Epsilon models (E.cor$_{nat}$, B.den$_{nat}$, N.muc$_{nat}$ and K.den$_{nat}$) clustered together and

were skewed towards considerably higher PAE range 8.5-10 than AF3 and Chai-1 and

covered a pLDDT range of 82-85.

All seven ComP$_{nat}$ were modeled in Gamma. Gamma found its strongest models for E.cor$_{nat}$

modeled by both Chai-1 and AF3 (Fig 4B). As with Epsilon described above, the AF3

Gamma models covered a wider range than Chai-1 regarding both measures and reflected an

additional three ComP$_{nat}$ (N. cin$_{nat}$, N. men$_{nat}$ and a single N.muc$_{nat}$ model). RF2NA Gamma

models were clustered within narrower PAE and pLDDT ranges than AF3 and Chai-1 and

were again systematically skewed toward higher PAE.

Theta models were only found for N.cin$_{nat}$ in Chai-1 with PAE 2.5 and pLDDT 87. Theta

models formed an elongated cluster along both axes (Fig 4 A) in a similar manner as the

Chai-1 N.sub$_{nat}$ Epsilon models.

The Delta mode, which resembles Gamma (Table 2), was found in only three models in AF3

(N.men$_{nat}$), having PAE range 5-5.5 and comparatively low pLDDT range 78-82 and not in

Chai-1. This mode was however, modeled repeatedly in RF2NA with PAE range 8.3-10 and

pLDDT range 79-85 reflecting models for N.cin$_{nat}$, N.men$_{nat}$ and N.sub$_{nat}$.

Zeta was observed once in Chai-1 (N.men$_{nat}$) with PAE 3.5 and pLDDT 86 and repeatedly in

RF2NA (K.den$_{nat}$ and E.cor$_{nat}$) with PAE range 8.3-10 and pLDDT range 81-84. These robust

"Other" models which did not clearly fall into any of the other categorized binding modes

still had high pLDDT of 88 and low PAE of 3, while those from RF2NA were again

systematically skewed with PAE just below the cut-off, 9.5, and pLDDT 84.

All but one (B.den$_{nat}$) "Other" models in Chai-1 and RF2NA were E. cor$_{nat}$ models. The Chai-

1 models were again considerably better than those of RF2NA in terms of PAE/pLDDT.

Since no ComP$_{nat}$ modeled to the same distribution of modes across platforms, the main

results, confidence scores and cross-platform DockQ consistencies as presented in Figs 2-5

and Table 3 are described in further detail below for each ComP$_{nat}$. The statistics applied for

calculating significant differences in quality scores for each ComP$_{nat}$ are described in the

Methods section and results detailed in Code output S1-3.

Quality of the ComP$_{nat}$s

N.sub$_{nat}$: All models by AF3 (158/158) and Chai-1 (428/428) were in Epsilon, while RF2NA

models were in contrast assigned Delta (45/46) and Gamma (1/46). Overall, the Chai-1

models were found to be significantly better than AF3 models and both Chai-1's and AF3's

models were significantly better than RF2NA's in terms of PAE and pLDDT. In the

PAE/pLDDT scatter plot a smaller fraction of N.sub$_{nat}$ models from AF3 and Chai-

1overlapped each other in Epsilon and N.sub$_{nat}$ overlapped all other ComP$_{nat}$s except N.cin$_{nat}$

in Epsilon and none in Gamma (Figs 4B and C). N.sub$_{nat}$ models did however overlap in

terms of PAE/pLDDT with E.cor$_{nat}$, N.men$_{nat}$ and N.cin$_{nat}$ models in Delta (resolved scatter

not shown). A consistently flipped-out base and DNA strand deformation were observed in

the Chai-1 models (Fig 3A), showcasing the platform's unique ability to strongly alter the

DNA structure of these short templates, possibly to induce an optimized fit. Re-docking in

HADDOCK showed strong bias for Gamma in moderate agreement with the modeling results

(Table 3). HADDOCK predicted Gamma (159/200), Epsilon (25/200) and "Other" (15/200).

The high number of Delta in RF2NA and Gamma in HADDOCK could potentially be better

interpreted together in considering that these modes are minor variants of each other with

equal positioning of the β1-β2-loop. The cross-platform model DockQ consistency

comparisons for N.sub$_{nat}$ (Fig 5) showed that a sub-set of the Epsilon models were virtually

identical and reached high quality with excellent scores $\geq$ 0.8. The bulk of AF3 vs. Chai-1

model pairings, however, were in the medium quality range (DockQ $\approx$ 0.6), and all were in

Epsilon, showing high cross-platform convergence on Epsilon. Both these platform models

for N. sub$_{nat}$ compared to RF2NA models had most pairings in the acceptable quality range

and were consistently divergent on mode (i.e. "Different") reflecting the high number of

modeled Delta modes in RF2NA and Epsilon in the other two platforms.

B.den$_{nat}$: All models by AF3 (165/165) and Chai-1 (91/91) were Epsilon, as were the

majority (20/22) of the RF2NA models. In addition to Epsilon, RF2NA modeled a single

Gamma (1/22) and a single "Other" (1/22) model. The AF3 models were significantly better

in terms of PAE than those from Chai-1 whilst there was no significant difference in terms of

pLDDT. Both Chai-1 and AF3 B.den$_{nat}$ models were again significantly better than the

RF2NA models. In the PAE/pLDDT scatter plot B.den$_{nat}$ models overlapped with E.cor$_{nat}$,

N.sub$_{nat}$, K.den$_{nat}$ and N.muc$_{nat}$ models in Epsilon and with E.cor$_{nat}$, N.muc$_{nat}$, and N.cin$_{nat}$ in

Gamma (Fig 4B and C). The re-docking in HADDOCK corroborated the modeling

experiments in finding a strong bias for Epsilon (145/200), moderate number of Eta ("Other")

(54/200) and equivalently to RF2NA, only a single Gamma (1/200). The cross-platform

model DockQ consistency comparisons for B.den$_{nat}$ (Fig 5) showed that all AF3 vs. Chai-1

model pairings were in the medium DockQ quality range and convergently Epsilon, with

some pairings being near high quality (Fig 5). All AF3 vs. RF2NA pairings were in the

acceptable range, except for a residual of pairings that were in the lower medium range,

divergent as Different or convergently in the Epsilon binding mode. All Chai-1 vs. RF2NA

pairings were similarly also in the acceptable range with either Different or convergently

Epsilon modes.

K.den$_{nat}$: All models by AF3 (105/105) and Chai-1 (446/446) were Epsilon. AF3 models

were significantly better than Chai-1 in regard of pLDDT and insignificantly different for

PAE. RF2NA models were found in Zeta (69/98), Epsilon (25/98) and Gamma (4/98) with

significantly poorer PAE/pLDDT quality scores than AF3 and Chai-1. In the PAE/pLDDT

scatter plot a sub-fraction of AF3 and Chai-1 K.den$_{nat}$ Epsilon models overlapped in addition

to models of all other ComP$_{nat}$s except N.cin$_{nat}$ in Epsilon and all ComP$_{nat}$s in Gamma (Figs

4B and C). Re-docking in HADDOCK predicted unambiguously Epsilon (200/200) in full

agreement with AF3 and Chai-1 and moderately different from RF2NA. The high number of

Zeta in RF2NA and Epsilon in HADDOCK could potentially be better interpreted together in

considering in that these modes, like Gamma/Delta,are minor variants of each other with

identical positioning of the β1-β2-loop. The cross-platform model DockQ consistency

comparisons for K.den$_{nat}$ (Fig 5) showed that some model pairings reached high DockQ ≥ 0.7

for AF3 vs. Chai-1 comparisons, but most pairings were in the acceptable range and

convergently Epsilon. For the comparisons to RF2NA, most model pairings were in the

acceptable range, yet some were in the medium range. The distributions of divergently

Different and convergently Epsilon pairings were very similar.

N.muc$_{nat}$: All models by AF3 (105/105) were Epsilon. No Chai-1 models reached ipTM $\geq$

0.6, and all the 500 best scoring models were Epsilon (500/500). The relatively few

successful RF2NA models were Gamma (10/11) and "Other" (1/11). The AF3 N.muc$_{nat}$

models were significantly better than the Chai-1 models in terms of PAE and Chai-1 models

were significantly better than AF3 in terms of pLDDT. The models from both Chai-1 and

AF3 were significantly better than those from RF2NA in terms of PAE, but pLDDT was not

significantly different between Chai-1 and RF2NA. In the PAE/pLDDT scatter plot N.muc$_{nat}$

models overlapped with E.cor$_{nat}$, N.sub$_{nat}$, K.den$_{nat}$ and B.den$_{nat}$ models in Epsilon and all

ComP$_{nat}$ except N.sub$_{nat}$ in Gamma (Fig 4B and C). Re-docking in HADDOCK corroborated

findings from AF3 and RF2NA with a strong bias for Epsilon (145/200), yet both Gamma

(53/200) and "Other" (2/200) modes were re-docked in concordance with RF2NA. The cross-

platform model DockQ consistency comparisons for N.muc$_{nat}$ (Fig 5) showed that some

models reached high DockQ scores close to 0.8 for AF3 vs. Chai-1, with a broad range of

DockQ scores, both acceptable and medium and convergently Epsilon. The comparisons to

RF2NA were mostly in the acceptable range, represented by both divergent Different and

convergent Epsilon modes.

E.cor$_{nat}$: All models by AF3 (95/95) were uniquely Gamma. Chai-1 models were ambiguous

in terms of binding mode in that the models with lowest ipTM were either Gamma (138/381)

or "Other" (86/381), while the better models (157/381) were Epsilon. The RF2NA models

were also ambiguous in terms of binding mode, with Gamma (26/81), Epsilon (23/81), Zeta

(25/81), Delta (1/81) and Other (6/81) assignments making E.cor$_{nat}$ the ComP$_{nat}$ with most

modeled modes overall (see Fig 3E). The Chai-1 models were significantly better than both

AF3 and RF2NA and AF3 models were significantly better than RF2NA in terms of

PAE/pLDDT. In the PAE/pLDDT scatter plot $E.cor_{nat}$ models overlapped with $N.sub_{nat}$,

$B.den_{nat}$, $N.muc_{nat}$ and $K.den_{nat}$ models in Epsilon and with all $ComP_{nat}$ except $N.sub_{nat}$ in

Gamma (Fig 4B and C). The re-docking in HADDOCK showed in stark contrast to AF3 and

RF2NA no Gamma but a strong bias for Epsilon (170/188) and a few resembling Zeta

(18/188). $E.cor_{nat}$ was consequently the least coherently modeled $ComP_{nat}$ for either Epsilon

and Gamma across platforms and in re-docking. The cross-platform model DockQ

consistency comparisons for $E.cor_{nat}$ (Fig 5) showed that AF3 vs. Chai-1 paired with both

medium and acceptable DockQ scores, most often convergently Gamma with some divergent

and acceptable Different model pairings. In the AF3 vs. RF2NA comparisons (Fig 5) , most

pairings were in the acceptable range, convergently Gamma and divergently Different. The

bulk of Chai-1 vs. RF2NA model pairs were in the acceptable range and convergent on

Gamma, Epsilon and Other modes. Also, several divergently Different comparisons were

found.

$N.cin_{nat}$: The AF3 models never reached ipTM ≥ 0.6 and of the 100 best scoring models,

were Gamma (96/100) and few Epsilon (4/100). In Chai-1, a unique Theta mode was

observed for all models (395/395), a mode exclusively seen in Chai-1 for this $N.cin_{nat}$. The

RF2NA models were either Gamma (40/52) or the resembling Delta (12/52). Chai-1 models

were significantly better than AF3 and RF2NA in terms of PAE and pLDDT. AF3 models

were significantly better than RF2NA in terms of PAE while interestingly, RF2NA models

were significantly better than AF3 models in terms of pLDDT. In the PAE/pLDDT scatter

plot $N.cin_{nat}$ overlapped with $N.men_{nat}$ in Epsilon and all $ComP_{nat}$s except $N.sub_{nat}$ in Gamma

(Fig 4 B and C). In the unique Theta models, a complete strand separation in one end of the

DNA and one flipped out base were observed (Fig 3 F and Fig S1 C), again showcasing a

unique Chai-1 feature for altering DNA structure, possibly to induce an optimal fit. Re-

docking in HADDOCK showed like AF3 and RF2NA a bias for Gamma (121/200) with less

Epsilon (79/200). Ambiguities in both modeling and re-docking and the special strand-

splitting Theta mode predicted by Chai-1 suggests that N.cin$_{nat}$ has unique features in its

structure to significantly affect both modeling and re-docking of the AT-DUS template. The

cross-platform model DockQ consistency comparisons for N.cin$_{nat}$ (Fig 5) showed that AF3

vs. Chai-1 comparisons never reached acceptable DockQ, and all pairings were divergently

Different. Several AF3 vs. RF2NA pairings reached high-quality DockQ, albeit with most

pairings in the medium range and converged on Gamma. Several models were divergently in

Different modes. All Chai-1 vs. RF2NA comparisons were below the acceptable range and

consistently divergent with respect to mode.

N.men$_{nat}$: The AF3 models never reached ipTM $\geq$ 0.6, and the 100 best scoring models were

Gamma (76/100), Epsilon (21/100) and Delta (3/100). The relatively few Chai-1 models were

in Epsilon (12/13) and Zeta (1/13) and the RF2NA models were in Gamma (56/91) and Delta

(35/91). Chai-1 gave significantly better PAE and pLDDT scores than both AF3 and RF2NA.

As with Ncin$_{nat}$, the RF2NA models were significantly better than the AF3 in terms of

pLDDT, albeit having significantly weaker PAE. In the PAE/pLDDT scatter plot N.men$_{nat}$

models overlapped with N.sub$_{nat}$, K.den$_{nat}$ and N.cin$_{nat}$ models in Epsilon and K.den$_{nat}$,

N.cin$_{nat}$, E.cor$_{nat}$ and N.muc$_{nat}$ in Gamma (Fig 4 B and C). Re-docking in HADDOCK

showed in contrast to AF3 and RF2NA a bias for Epsilon (130/200), yet with several Gamma

(68/200) and a few Zeta (2/200). Few successful Chai-1 models may signify the platforms

difficulty in modeling this ComP$_{nat}$ complex. The cross-platform model DockQ consistency

comparisons for N.men$_{nat}$ (Fig 5) showed that most AF3 vs. Chai-1 pairings were in the

acceptable range, being Different or convergently Epsilon, with some Epsilon pairings

clustering in the medium range. AF3 vs. RF2NA convergently predicted Delta and Gamma in

the acceptable and medium ranges with some pairings close to the high-quality range. All

Chai-1 vs. RF2NA pairings were divergently Different and in the acceptable DockQ score

range.

562   AF3 modeling quality in matching native vs. scrambled DUS to ComP

In order to explore modeling quality and consistency in regard of DUS specificity, we

modeled $ComP_{nat}s$ and $ComP_{scr}$ and compared the resulting ipTM and Chain Pair PAE

minimum (CPPM) distributions for all models (Fig 6 A and B; Code output S2).

Figure 6.   **A:** Violin plots showing distributions of ipTM for native ("Native") and

scrambled DUS ("Scrambled") runs in AF3 for the docking set consisting of five DUS

dialects encompassing seven $ComP_{nat/scr}$ models. Box plots encompassing the median and

interquartile ranges are seen inside the violins. AA-king3DUS (K.den). AG-DUS (B.den +

N.sub). AG-eikDUS (E.cor). AG-mucDUS (N.muc). AT-DUS (N.men + N.cin). **B:** Same as

in A for CPPM. Refer to Code Output S2 for statistical significance tests for Native vs.

Scrambled for all DUS dialects.

Since the native DUS dialects were the same in the two pairs $N.men_{nat}/N.cin_{nat}$ (AT-DUS)

and $B.den_{nat}/N.sub_{nat}$ (AG-DUS), these were combined in the assessment. AF3 was chosen as

platform as it covered most $ComP_{nat}$ (6/7) with robust (ipTM $\geq$ 0.6) models. Wilcoxon rank

sum tests (Code output S3), using significance level α=0.05, showed that the ipTM

distributions (Fig 6 A) for native DUS models ("Native") and scrambled DUS models

("Scrambled") were significantly different in all comparisons except for AA-king3DUS.

Furthermore, ipTM in Native was significantly better (higher) than Scrambled for AG-DUS,

AG-eikDUS and AG-mucDUS. In contrast, for AT-DUS the ipTM distribution was

significantly lower (inferior) in Native than in Scrambled. This suggested a greater difficulty

for AF3 to generate high-quality models for Scrambled than Native for AG-DUS, AG-

eikDUS and AG-mucDUS, whereas AF3 modeled high quality models for both Scrambled

and Native for AA-king3DUS. For AT-DUS AF3 modeled better complexes with Scrambled

than with Native. The CPPM scores (Fig 5 B) reflected ipTM in that the distributions were

significantly different in all comparisons except AA-king3DUS. CPPM was significantly

better (lower) in Native than Scrambled for AG-DUS, AG-eikDUS and AG-mucDUS. For

AT-DUS, CPPM was significantly higher (inferior) in Native than Scrambled, corroborating

the results for ipTM and reflecting AF3's difficulty in predicting $ComP_{nat}$s with the AT-DUS

dialect and high variation in the models for these complexes. The CPPM results also

indicated that AF3 was not distinguishing AA-king3DUS from its scrambled versions. The

frequencies of assigned binding modes using scrambled DUS in AF3 are shown in Table 4,

showing that also these are mainly Epsilon and Gamma.

Complex	$ComP_{scr}$					$ComP_{nat}$				
Mode/ComP	⬚	ζ	γ	δ	o	⬚	ζ	γ	δ	o
N.sub	64	-	36	-		258	-	-	-	-
B.den	30	-	35	1	1	165	-	-	-	-
K.den	87	-	7	-	-	146	-	-	-	-

N.muc	16	-	18	-	1	105	-	-	-	-
E.cor	8	-	38	-	-	-	-	95	-	-
N.cin	4	-	29	-	5	4	-	-	-	-
N.men	4	-	22	-	2	-	-	-	-	-
SUM	213	0	195	1	9	678	1	95	0	0

Table 4. Binding mode distribution across all seven ComPs generated with either scrambled DUS (left) and native DUS (right) in AF3. Greek letter symbols for modes are ⬚ = Epsilon, ζ = Zeta, γ=Gamma, δ=Delta and **o**=other. Data for native DUS are as in Table 1 but with a strict ipTM ≥ 0.6 applied.

In considering the mode differences for each ComP going from native to scrambled DUS, as described in Table 4, we observe the following tendencies: Robust Gamma models appear in six ComP$_{scr}$ (N.sub$_{scr}$, Kden$_{scr}$, Bden$_{scr}$, N.muc$_{scr}$, N.cin$_{scr}$, N.men$_{scr}$) where there were no Gamma modes above cut-off for ComP$_{nat}$, and inversely Epsilon appear in E.cor$_{scr}$ where there were none in E.cor$_{nat}$. AF3 generated robust models for ComP$_{N.men}$ only when using scrambled DUS and below cut-off with native DUS. Representative ComP$_{scr}$ PAE-plots for each mode and ComP$_{scr}$ are shown in Fig S7.

Discussion

In this study, three different deep learning protein structure prediction platforms and one traditional molecular docking platform were used to model the interaction between ComP and DUS in native ComP$_{nat}$ complexes. The seven ComPs studied were chosen as they represented both a diverse set regarding primary structure from three different genera (*Neisseria/Bergeriella, Eikenella* and *Kingella*) (four genera if considering *Bergeriella* distinct from *Neisseria*) and had their respective genomes enriched with five different DUS dialects as previously described (Frye, 2016). The overall 3D structure of the ComP protein was found conserved in these investigated ComPs. The overall ComP domain organization and structure conservation were found highly similar and showed that all tested proteins likely were functionally conserved as DUS specific receptors as previously interpreted from ComP sequence alignments (Cehovin et al., 2013). Differences in amino acid composition of ComPs, as well as minor differences in structural domains in ComPs, e.g. the here predicted α-turns in AF3 ComP$_{B.den}$, ComP$_{E.cor}$, and ComP$_{N.muc}$, might together help understand general and specific characteristics of the ComP :: DUS interaction. It remains unclear why RF2NA generally did not predict the highly conserved disulfide bridge between β1-sheet and the DD-region. The disulfide bridged DD-region is of particular interest as it is expected to facilitate specific DUS-binding (Cehovin et al., 2013). Although the disulfide bridge between the β1-sheet and the DD-region is missing in RF2NA models, the relative positioning of the DD-loop in the overall ComP structure remains largely the same in the RF2NA models as in all other modeled structures and seems therefore not to have had significant impact on the RF2NA ComP$_{nat}$ models. It cannot be fully excluded however, that the propensity to model individual DNA binding modes by RF2NA could be influenced by more potential flexibility of the DD-loop lacking its disulfide bridge.

The ComP$_{nat}$ binding modes

By modeling ComP$_{nat}$s, six different binding modes and one heterogeneous "Other" category

were described. These categories were based on their primary and main interactions of two

consistently DNA-interacting loops in ComP, the α1-β1 loop and the β1-β2 loop.

Epsilon/Zeta/Theta were categorized by how these two loops interact through the DNA major

groove and Gamma/Delta through the minor groove (Table 2). The sub-divisions of Epsilon

and Gamma into Zeta/Theta and Delta, respectively, were based on how the DD-region

differentially interacted with DNA. Although ComP :: DUS docking by HADDOCK in our

hands frequently distorted the DNA structure making the grooves challenging to distinguish,

we have interpreted previously illustrated results from in silico ComP-DUS docking

experiments in Berry et al. (2016) and Hughes-Games (2020) to reflect the Epsilon mode, i.e.

main interactions through the major groove. Hence the Epsilon mode as such, has some

previous docking support.

Strong support for the Epsilon mode was also found here across platforms and for different

ComP$_{nat}$. All ComP$_{nat}$ were modeled in Epsilon by one or more platforms. Epsilon was the

most frequently modeled mode in AF3 and Chai-1 by a considerable margin, and this

therefore supports Epsilon as a very likely ComP$_{nat}$ complex. All platforms modeled Epsilon

for three or more ComP$_{nat}$ and the Epsilon models were particularly robust and consistent in

AF3 and Chai-1 for N.sub$_{nat}$, N. muc$_{nat}$, B. den$_{nat}$ and some of the K. den$_{nat}$ (Table 3; Fig3-5).

Fewer successful Epsilon models were found in E. cor$_{nat}$, N. cin$_{nat}$ and N. men$_{nat}$, for which

more Gamma was observed.

Zeta resembles Epsilon by having the α1-β1 and β1-β2 loops entering the major groove. Zeta

was not found in AF3. Zeta was always modeled together with Epsilon for the same ComP$_{nat}$

in the two platforms where Zeta was seen, Chai-1 and RF2NA, signifying the similarity of

the two modes. A single Zeta model was found for N. men$_{nat}$ in Chai-1 together with several

Epsilon. In RF2NA, Zeta and Epsilon were modeled similarly for K. den$_{nat}$ and E. cor$_{nat}$.

These observations support considering Zeta a DD-loop variant of Epsilon.

The Theta mode was uniquely identified in N.cin$_{nat}$ and only in Chai-1. Unlike all other

modes, Theta showed major conformational changes in the DNA, complete strand separation,

which potentially could allow better protein access to DNA bases.   Theta resembles

Epsilon/Zeta in that the major interactions were in the major groove, which was pried apart in

Theta. The consistently flipped-out base is the central thymine in the conserved 5'-CTG-3'

DUS motif, alluding to involvement in DUS-specific binding. The only other severe

alteration of the DNA structure was found in N. sub$_{nat}$ Epsilon Chai-1 models, also displaying

a flipped-out base. This fifth thymine base immediately precedes the conserved CTG core of

the AG-DUS and it is thus tempting to speculate that the modeled conformational change in

the DNA may instigate specific binding. The capability of Chai-1 to model substantial

conformational changes in DNA sets it apart from AF3 and RF2NA and may be due to

different emphasis on confidence metrics and that Chai-1 allowed also defining different

MSA schemes for the DNA. Further experimentation by e.g. molecular dynamics is required

to investigate these models in depth by manipulation of input parameters and MSA schemes.

Particularly strong support for Gamma was found in three ComP$_{nat}$, N. cin$_{nat}$, N. men$_{nat}$ and E.

cor$_{nat}$. Together all platforms modeled Gamma for at least one of these three ComP$_{nat}$ and all

ComP$_{nat}$s were modeled in Gamma in at least one platform. RF2NA generated Gamma

models for all ComP$_{nat}$s. Gamma was the most frequently modeled mode in RF2NA and

second most frequent mode in AF3 and Chai-1 after Epsilon.

Like Zeta/Theta can be considered variants of Epsilon, Delta resembles Gamma in having its

main interactions with the minor groove. When Delta was found for any ComP$_{nat}$, the same

platform also modeled Gamma, showing a correlation between the two modes as with

Epsilon/Zeta. These mode and sub-mode models may reflect a flexibility of the DD-region

which may be relevant, particularly in regard to the missing disulfide bridge in RF2NA. Delta

was modeled for $N.sub_{nat}$, $N.cin_{nat}$, $N.men_{nat}$ and $E.cor_{nat}$ by RF2NA and below ipTM cutoff in

AF3 ($N.men_{nat}$). $N.sub_{nat}$ was the only modeled complex in this RF2NA Delta subset with

also very robust Epsilon models in AF3 and Chai-1. In modeling predictions yielding Delta,

$N.cin_{nat}$, $N.men_{nat}$ and $E.cor_{nat}$ stand out as complexes imposing modeling challenges for all

platforms, although $E.cor_{nat}$ achieved moderately high ipTM scores in Gamma in AF3 and

Gamma/Epsilon in Chai-1. $E.cor_{nat}$ was also the only complex with a substantial fraction un-

categorized modes assigned Other.

The Eta model was never modeled but appeared in the re-docking experiments using

HADDOCK. Eta was therefore not considered a supported mode but draws attention to the

ability of re-docking to both reproduce other modeled complexes and predict new.

HADDOCK did robustly generate re-docks showing both Epsilon and Gamma modes in

support of these.

Other modes than the above were considered anomalies in modeling. As can be inspected in

Fig 3, the Other modes of $E.cor_{nat}$ in Chai-1 and a single model of $B.den_{nat}$ in RF2NA either

bind DNA to the opposite end of ComP or at greater distance and tilt relative to all other

models. If anything, these models again show the challenge in convergently modeling $E.cor_{nat}$

to any mode (modeled in all modes except Theta) and that the RF2NA outputs generally are

more mode-variable than AF3 and Chai-1.

Platform performances and consensus predictions

Generally, under our application of confidence scores and metrices, RF2NA predicted less robust models than AF3 and Chai-1. This observation aligns with those of Abramson et al. (2024) demonstrating RF2NA to be inferior to AF3 for protein-nucleic acid prediction on benchmark sets of the PDB database, having about 40% lower success rate than AF3. However, since the ComP :: DUS structure remains unresolved and may be unusual due to its unique extracellular confinement, it cannot be excluded that RF2NA models may be informative for identifying the DUS-specific mode. Abramson et al. (2024) does not describe if there was a perfect overlap in successful AF3 and RF2NA models or if RF2NA modelled certain complexes or types of complexes with greater success than AF3. However, the systematically high PAE values of successful RF2NA models contrasts the unity of nearly identical AF3 models with much lower PAE values and speaks at least for the robustness of the AF3 models and possibly the impropriety of emphasizing PAE metrics for RF2NA models. In support of the RF2NA Gamma models, it is notable that the pLDDT scores from RF2NA generally were relatively high (>82), like most of the models from AF3 and Chai-1 and that N.cin$_{nat}$ and N.men$_{nat}$ modeled with significant better pLDDT values than AF3 in Gamma. Although there is some variation within successful Chai-1 models, the consistent overlap with AF3 in terms of both Epsilon and Gamma is evident. Higher variation in the Chai-1 models compared to those from AF3 may also be explained by Chai-1's variable MSA input, whereas the AF3 web server does not allow for MSA customization (AF3 now released as open source). Thus, the cross-platform deviance may stem from the three platforms' unequal algorithmic modeling, score metric emphases and source inputs or simply that AF3 and Chai-1 models being of generally better quality than the those from RF2NA. However, all platforms and modes may prove informative when exploring DUS-specificity. Assuming that there is one and only one DUS-specific mode across all ComPs, it is paradoxical that

even in the cases where the models were of putatively high quality in both programs

(RF2NA: PAE < 10; AF3: ipTM $\geq$ 0.6), the modes were robustly either Epsilon or Gamma.

For instance, RF2NA may have provided insight into $ComP_{nat}s$ modeled with AT-DUS

($N.men_{nat}$ and $N.cin_{nat}$) with solely Gamma/Delta predictions, as both of these had models of

PAE <10, while AF3 always failed to predict $N.men_{nat}$ and $N.cin_{nat}$ models above its ipTM

threshold. At the same time, we never observed any RF2NA models of higher quality than

any AF3 or Chai-1 models when considering confidence from PAE, only pLDDT cases and

DockQ consistency, which is striking. RF2NA predicted generally more modes per $ComP_{nat}$

than AF3 and Chai-1. RF2NA therefore had challenges with converging to individual modes

and displayed greater modelling flexibility relative to AF3 and Chai-1. This may be due to

fundamentally different emphasis on different confidence metrices. Although more modes

were modelled per $ComP_{nat}$ in RF2NA, a converging trend towards Gamma (and Delta) was

observed in support of a robust loop-interaction prediction with the minor groove. Although it

is unclear to which degree RF2NA with its modelling flexibility can return "false positive"

$ComP_{nat}$ models of high confidence that fail to recover the true binding mode, Baek et al.

(2024) reported that 81% of models with PAE <10 correctly predicted the interface of the

complex, which is substantial.

In exploring the AF3 modeling quality of DUS against scrambled DUS using CPPM

confidence, $ComP_{N.cin}$ and $ComP_{N.men}$ were found as the only ComPs where scrambled DUS

models were significantly better than their native pairings of which most were below the

ipTM threshold (Table 3-4; Fig 5). Most of these scrambled models were in Gamma (Table

4). For AF3 to predict robust models for these two ComPs, a scrambled AT-DUS was

required, suggesting that individual bases in the native AT-DUS were preventive for AF3. In

contrast, $ComP_{B.den}/ComP_{N.sub}$, $ComP_{N.muc}$ and $ComP_{E.cor}$ AF3 models were significantly

better for native Epsilon or Gamma complexes than scrambled. These results suggest that

AF3 more accurately predicted the DUS-specific mode than any non-specific mode in these

ComPs although they were discrepantly Epsilon and Gamma and hence not parsimonious in

regard of mode. The ComP$_{scr}$ data show that AF3 predicted Gamma for N.men$_{scr}$ and N.cin$_{scr}$

in a seemingly DUS-independent manner and E.cor$_{nat}$ in a seemingly DUS-dependent manner

(Table 4). In contrast, AF3 robustly predicted Epsilon for B.den$_{nat}$, K.den$_{nat}$, N.sub$_{nat}$,

N.muc$_{nat}$ in a seemingly DUS-dependent manner and more Gamma in a seemingly DUS-

independent manner. Furthermore, ComP$_{K.den}$ did not significantly differentiate between its

native DUS AA-king3-DUS and the scrambled versions (Fig 5), questioning whether any of

the two modes could represent the ComP$_{K.den}$ DUS-specific mode. We have previously shown

that the DUS dialect phylogenetic distribution is consistent with the robust core genome

Neisseriaceae phylogeny (Frye, 2013). Finding here that the seven different ComP$_{nat}$s showed

propensities to model the two different modes in a manner inconsistent with the ComP

phylogeny remains puzzling. The strict conservation of the essential CTG core of DUS

suggests the molecular interactions and mechanism responsible for the ComP :: DUS binding

and consequential DNA uptake is one. Wet-lab experiments have shown that DNA

containing any single mutation in this CTG core completely and uniquely abrogates

transformation in both *N. meningitidis* and *N. subflava* (Berry et al., 2013; Frye et al., 2013)

whose ComPs here show propensities to model either Epsilon (*N. subflava*) or Gamma (*N.*

meningitidis) with their respective DUS in AF3 and other platforms. These *in vivo* results

support the notion that there is only one DUS-specific interaction and mechanism. The

modeling platforms used here do either inconsistently or not at all capture this expected

parsimony in their robustly predicted different modes. There could be different reasons for

this. Firstly, all predicted models are precisely that, predicted models. Although the here

applied platforms represent state-of-the-art tools for protein structure and protein-DNA

complex structure prediction, obtaining functional understanding is not immediate or straight

forward (Varadi et al., 2025). Issues with confidences of predicted disordered protein regions,

dynamic interactions, multi-chain complexes such as protein-DNA, training-set

representativeness and lack of benchmarking studies are different recently discussed

challenges (Varadi et al., 2025; Chakravarty et al., 2024; Ruff & Pappu, 2021). The ComPs

investigated here are to platform variable extents modeled based on pattern recognition,

learned protein energetics and possibly wet-lab resolved 3D structures inside user-

inextricable black-box algorithms. Basing our analysis on different such complex deep-

learning platform algorithms with diverging dependencies and potentially dynamic inputs, we

cannot exclude the possibility that the positioning of particularly flexible or disordered

regions in ComP are inaccurately modeled, such as the lack of biologically important

disulfide bridges in RF2NA. We also observe varying abilities to severely modulate the

structure of DNA, particularly seen in some Chai-1 models. A possible consequence of any

such platform biases could be that one mode is artificially favored in modeling not to

represent an expected ground truth. Hence, our search for cross-platform consensuses

applying fundamentally different modeling algorithms. As shown in Fig 2 the topologies of

the investigated ComPs are highly similar, yet three ComPs have modeled α-turns different

from the others. The β1-β2 loop have overlapping α-turns in $ComP_{Ecor}$, and $ComP_{Nmuc}$ and

$ComP_{Bden}$ have one in the DD-loop. We cannot find that these α-turn divergent topologies

could explain propensities for either Epsilon ($ComP_{Nmuc}$ and $ComP_{Bden}$) or Gamma

($ComP_{Ecor}$) in any systematic way and we have fair confidence in their predicted protein 3D

structures and overall structural coherence across platforms.

All our predicted models and modes are static snapshots which do not reflect any

spatiotemporal dynamics of the biological ComP :: DUS interaction. We cannot therefore

exclude the possibility that the DNA binding modes identified here may be limited to

showing the initial interactions between DUS and ComP, i.e. the first step towards making an

interaction which involves the essential CTG-core after sequential or simultaneous

conformational changes have been made in the protein, DNA, or both. We do note that it is

particularly the two AT-DUS binding ComPs and the AG-eikDUS binding ComP which are

more difficult to model in Epsilon than Gamma, particularly in AF3. The AT-DUS differs

from AG-DUS only in position -1 (A$\underline{T}$GCCGT$\underline{CTG}$AA vs. A$\underline{GG}$CCGT$\underline{CTG}$AA) five

nucleotide positions away from the CTG core which occupy DUS-positions 6-8. Hence the

singular -1 position away from the conserved core seems restrictive for making a robust

Epsilon mode in AT-DUS ComPs and should be investigated further (see below). The AG-

eikDUS (AGG$\underline{CTA}$C$\underline{CTG}$AA) dialect differs in 2-3 positions relative to the other dialects

and two of these differences are dialect-unique with T in position 3 and A in position 4 much

closer to the CTG-core than -1. For ComP$_{Nmen}$, Epsilon modes appear in AF3 only when

scrambled AT-DUS are used and relatively few Epsilon modes are found for ComP$_{Ncin}$ with

both templates (Table 4). ComP$_{Ecor}$ with AG-eikDUS and scrambled versions show similarly

that Epsilon appears only when using scrambled AG-eikDUS in AF3. Consequently, all three

ComPs harbour the propensity to model Epsilon *per se*, but are restricted by their native DUS

to do so. It is interesting that Epsilon modes remain scarce relative to Gamma in models of

all these three ComPs when using scrambled versions of DUS. This shows that these ComP's

retain an overall propensity for Gamma with different DNA templates even though their DUS

dialect-unique positions (-1 and 3+4) are not the same. The other four ComPs (ComP$_{Kden}$,

ComP$_{Bden}$, ComP$_{Nsub}$ and ComP$_{Nmuc}$) show an opposite tendency to loose their strict Epsilon

propensity when scrambled DUS versions are used and Gamma modes appear (Table 4).

These Epsilon to Gamma mode-shifts are particularly strong for ComP$_{Nmuc}$ and ComP$_{Bden}$

which show about equal numbers of Gamma and Epsilon modes when scrambled DNA

templates are used, and are phylogenetically closer to ComP$_{Nmen}$ and ComP$_{Ncin}$ than

ComP$_{Kden}$ and ComPE$_{cor}$. In contrast, ComP$_{Kden}$, retain much of its Epsilon propensity when

the scrambled AA-king3DUS was used resembling the disinclination to shift from Gamma to

Epsilon as discussed above. Since Epsilon and Gamma models are robustly predicted with

scrambled DUS in all investigated ComPs we find support for the notion that the models

predicted here could reflect initial contacts where DUS-specific interactions are yet to

completely unfold. Further systematic modeling studies of propensties to model different

modes using non-native DUS, artificial deconstructed DUS (incl. all 36 possible individual

transitions and transversions pr. DUS dialects) and DNA templates of different lengths and

internal DUS-positions could shed further light on these shifts in mode propensities and

better support Gamma or Epsilon as the one parsimonious DNA-binding mode responsible

for DUS-specificity. Any mode further supported by in silico approaches would require

confirming wet-lab assaying experiments to which we here provide direction.

Taken together, these data support that Gamma and Epsilon reflect the most robust and

inherent DNA-binding properties of ComPs irrespective of DUS-specificity. Experimental

data by Cehovin et al. (2013) showed that, in addition to DUS-specific transformation, non-

DUS transformation is also mediated by $ComP_{N.men}$. These *in vivo* results showed not only

the well-characterized wild-type bias for DUS-specific transformation, but also that non-DUS

transformation could reach the level of DUS-specific transformation when ComP was

overexpressed (Cehovin et al., 2013). We hypothesize that this DUS/non-DUS ambiguity

could be connected to the two distinct and robust binding modes identified here, Epsilon and

Gamma. Finally, DNA binding and DNA uptake are consecutive events in the transformation

process of which the modeling performed here considers the first step and initial interaction

between protein and DNA. Specific DNA-binding may require considerable conformational

changes in either or both ComP and DUS of which Gamma/Delta and Epsilon/Theta/Zeta,

could represent initial states. Molecular dynamics simulations may shed further light on

downstream conformational changes in binding partners in future in silico experiments and

ultimately resolve the specific molecular interactions responsible for DUS-specificity and

better explain the here documented divergent modeling performance of the different

platforms.

Summary of conclusions

Two distinctly different DNA binding modes of ComP, Epsilon and Gamma, were robustly

predicted by AF3, Chai-1 and RF2NA and HADDOCK. Epsilon and Gamma differ

characteristically in how two conserved ComP protein domains, the β1-β2-loop and the DD-

loop, interact with the minor and major grooves in the DNA. Although robustly modeled

across platforms, the modeling platforms showed considerable variation in their propensity to

model these two modes in the seven ComP :: DUS complexes explored here. RF2NA seemed

to place fundamentally different emphasis on PAE relative to AF3 and Chai-1 which skewed

the quality assessment in a systematic manner to the disadvantage of RF2NA models. In

contrast to PAE, pLDDT scores were considerably more overlapping and within high

confidence range (>82) across platforms in Epsilon and Gamma models and notably so in

Gamma. Robust support for Epsilon was found in N. sub$_{nat}$, B. den$_{nat}$, K. den$_{nat}$ and N. muc$_{nat}$,

and Gamma mainly found its support in E. cor$_{nat}$, N. cin$_{nat}$ and N. men$_{nat}$. RF2NA generally

predicted more Gamma whilst AF3, Chai-1 and re-docking in HADDOCK predicted more

Epsilon to show inherent differences between the three modeling platforms and relative to re-

docking. No parsimonious Epsilon vs. Gamma mode distribution could be identified to

elucidate if any of these modes could represent DUS-specific or DUS-independent binding.

AF3 predicted robustly both Epsilon and Gamma models of higher quality (CPPM) when

using native DUS relative to scrambled DUS and possible modeling limitations responsible

for these mode-incongruent results are discussed. Scrambling the DUS provided models

shifting their native propensities to bind either Epsilon or Gamma to the other mode, showing

inherent modeling capabilities of all ComPs to accommodate both modes irrespective of

DUS-specificity. Although six different binding modes were predicted here, we find only

strong consensus modeling support for all ComPs to bind DNA in Epsilon and Gamma.

Data availability

885 Scripts and data used are provided in the GitHub repository

886 https://github.com/stianale/ComP-DUS-modeling.

Author contributions

SAH: Data curation, formal analysis, investigation, methodology, implementation of

computer codes and supporting algorithms, data presentation, writing original draft and

review & editing. AHL; KA and SAF: Supervision of SH and scientific critical review,

writing – review & editing. OHA: Conceptualization, funding acquisition, project

administration, resources, supervision, visualization, writing – review & editing. A Ph.D.

grant was provided for SAH from Faculty of Health Sciences, OsloMet, Norway. The funders

had no role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript.

References

Aas, F. E., Wolfgang, M., Frye, S., Dunham, S., Løvold, C., & Koomey, M. (2002).
Competence for natural transformation in Neisseria gonorrhoeae: components of
DNA binding and uptake linked to type IV pilus expression. *Molecular
microbiology*, *46*(3), 749-760.

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,
Willmore, L., Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung, C.-
C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti,
E., … Jumper, J. M. (2024). Accurate structure prediction of biomolecular
interactions with AlphaFold 3. *Nature*, *630*(8016), 493–500.
https://doi.org/10.1038/s41586-024-07487-w

Ambur, O. H., Frye, S. A., & Tønjum, T. (2007). New functional identity for the DNA uptake
sequence in transformation and its presence in transcriptional terminators. *Journal of
bacteriology*, *189*(5), 2077-2085.

Ambur, O. H., Engelstädter, J., Johnsen, P. J., Miller, E. L., & Rozen, D. E. (2016). Steady at
the wheel: Conservative sex and the benefits of bacterial transformation.
*Philosophical Transactions of the Royal Society of London. Series B, Biological
Sciences*, *371*(1706). https://doi.org/10.1098/rstb.2015.0528

Arnold, B. J., Huang, I. T., & Hanage, W. P. (2022). Horizontal gene transfer and adaptive
evolution in bacteria. *Nature Reviews Microbiology*, *20*(4), 206-218.

Baek, M., McHugh, R., Anishchenko, I., Jiang, H., Baker, D., & DiMaio, F. (2024). Accurate
prediction of protein–nucleic acid complexes using RoseTTAFoldNA. *Nature
Methods*, *21*(1), 117–121. https://doi.org/10.1038/s41592-023-02086-5

Basmaci, R., Bidet, P., Yagupsky, P., Muñoz-Almagro, C., Balashova, N. V., Doit, C., & Bonacorsi, S. (2014). Major intercontinentally distributed sequence types of Kingella kingae and development of a rapid molecular typing tool. *Journal of Clinical Microbiology*, *52*(11), 3890-3897.

Basu, S., & Wallner, B. (2016). DockQ: A Quality Measure for Protein-Protein Docking Models. *PLOS ONE*, *11*(8), e0161879. https://doi.org/10.1371/journal.pone.0161879

Bender, N., Hennes, M., & Maier, B. (2022). Mobility of extracellular DNA within gonococcal colonies. *Biofilm*, *4*, 100078.

Berry, J.-L., Cehovin, A., McDowell, M. A., Lea, S. M., & Pelicic, V. (2013). Functional Analysis of the Interdependence between DNA Uptake Sequence and Its Cognate ComP Receptor during Natural Transformation in Neisseria Species. *PLoS Genetics*, *9*(12), e1004014. https://doi.org/10.1371/journal.pgen.1004014

Berry, J.-L., Xu, Y., Ward, P. N., Lea, S. M., Matthews, S. J., & Pelicic, V. (2016). A Comparative Structure/Function Analysis of Two Type IV Pilin DNA Receptors Defines a Novel Mode of DNA Binding. *Structure*, *24*(6), 926–934. https://doi.org/10.1016/j.str.2016.04.001

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1), 421. https://doi.org/10.1186/1471-2105-10-421

Carter, E., Davis, S. A., & Hill, D. J. (2022). Rapid detection of Neisseria gonorrhoeae genomic DNA using gold nanoprobes which target the gonococcal DNA uptake sequence. *Frontiers in Cellular and Infection Microbiology*, *12*, 920447.

Cehovin, A., Simpson, P. J., McDowell, M. A., Brown, D. R., Noschese, R., Pallett, M., Brady, J., Baldwin, G. S., Lea, S. M., Matthews, S. J., & Pelicic, V. (2013). Specific

DNA recognition mediated by a type IV pilin. *Proceedings of the National Academy of Sciences*, *110*(8), 3065–3070. https://doi.org/10.1073/pnas.1218832110

Chai Discovery. (2024). Chai-1: Decoding the molecular interactions of life. *bioRxiv*. https://doi.org/10.1101/2024.10.10.615955

Chakravarty, D., Schafer, J. W., Chen, E. A., Thole, J. F., Ronish, L. A., Lee, M., & Porter, L. L. (2024). AlphaFold predictions of fold-switched conformations are driven by structure memorization. *Nature communications*, *15*(1), 7296.

Corso, G., Stärk, H., Jing, B., Barzilay, R., & Jaakkola, T. (2022). *DiffDock: Diffusion Steps, Twists, and Turns for Molecular Docking*. https://doi.org/10.48550/arxiv.2210.01776

Dillard, J. P., & Chan, J. M. (2024). Genetic Manipulation of Neisseria gonorrhoeae and Commensal Neisseria Species. *Current Protocols*, *4*(9), e70000.

Dubnau, D., & Blokesch, M. (2019). Mechanisms of DNA Uptake by Naturally Competent Bacteria. *Annual Review of Genetics*, *53*, 217–237. https://doi.org/10.1146/annurev-genet-112618-043641

Duffin, P. M., & Seifert, H. S. (2012). Genetic transformation of Neisseria gonorrhoeae shows a strand preference. *FEMS Microbiology Letters*, *334*(1), 44–48. https://doi.org/10.1111/j.1574-6968.2012.02612.x

Esmaeeli, R., Bauzá, A., & Perez, A. (2023). Structural predictions of protein–DNA binding: MELD-DNA. *Nucleic Acids Research*, *51*(4), 1625–1636. https://doi.org/10.1093/nar/gkad013

Frye, S. A., Nilsen, M., Tønjum, T., & Ambur, O. H. (2013). Dialects of the DNA Uptake Sequence in Neisseriaceae. *PLOS Genetics*, *9*(4), e1003458. https://doi.org/10.1371/journal.pgen.1003458

Goodman, S. D., & Scocca, J. J. (1988). Identification and arrangement of the DNA sequence recognized in specific transformation of Neisseria gonorrhoeae. *Proceedings of the National Academy of Sciences*, *85*(18), 6982-6986.

Grant, B. J., Rodrigues, A. P. C., ElSawy, K. M., McCammon, J. A., & Caves, L. S. D. (2006). Bio3d: An R package for the comparative analysis of protein structures. *Bioinformatics*, *22*(21), 2695–2696. https://doi.org/10.1093/bioinformatics/btl461

Grant, B. J., Skjærven, L., & Yao, X. (2021). The Bɪᴏ3D packages for structural bioinformatics. *Protein Science*, *30*(1), 20–30. https://doi.org/10.1002/pro.3923

Hepp, C., Gangel, H., Henseler, K., Guenther, N., & Maier, B. (2016). Single-stranded DNA uptake during gonococcal transformation. *Journal of bacteriology*, *198*(18), 2515-2523.

Hughes-Games, A. (2020). *New Approaches to Characterising Mechanisms of Horizontal Gene Transfer in Neisseria gonorrhoeae Using Microscopy and Genome Sequence Analysis*. ProQuest Dissertations & Theses.

Jacobsen, T., Bardiaux, B., Francetic, O., Izadi-Pruneyre, N., & Nilges, M. (2020). Structure and function of minor pilins of type IV pili. *Medical microbiology and immunology*, *209*(3), 301-308.Teixeira, J. M. C. Honorato, R. V., Giulini, M., Bonvin, A., Alidoost, S., Reys, V., Jimenez, B., Schulte, D., van Noort, C., Verhoeven, S., Vreede, B., Schott, S. & Tsai, R. (2024). *haddocking/haddock3: V3.0.0-beta.5* (Version v3.0.0-beta.5) [Computer software]. Zenodo. https://doi.org/10.5281/ZENODO.10527751

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R.,

Adler, J., … Hassabis, D. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, *596*(7873), 583–589. https://doi.org/10.1038/s41586-021-03819-2

Krishna, R., Wang, J., Ahern, W., Sturmfels, P., Venkatesh, P., Kalvet, I., Lee, G. R., Morey-Burrows, F. S., Anishchenko, I., Humphreys, I. R., McHugh, R., Vafeados, D., Li, X., Sutherland, G. A., Hitchcock, A., Hunter, C. N., Kang, A., Brackenbrough, E., Bera, A. K., … Baker, D. (n.d.). Generalized biomolecular modeling and design with RoseTTAFold All-Atom. *Science*, *384*(6693), eadl2528. https://doi.org/10.1126/science.adl2528

Mirabello, C., & Wallner, B. (2024). *DockQ v2: Improved automatic quality measure for protein multimers, nucleic acids, and small molecules*. Bioinformatics. https://doi.org/10.1101/2024.05.28.596225

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., & Steinegger, M. (2022). ColabFold: Making protein folding accessible to all. *Nature Methods*, *19*(6), 679–682. https://doi.org/10.1038/s41592-022-01488-1

Qiao, Z., Nie, W., Vahdat, A., Miller, T. F., & Anandkumar, A. (2024). State-specific protein–ligand complex structure prediction with a multiscale deep generative model. *Nature Machine Intelligence*, *6*(2), 195–208. https://doi.org/10.1038/s42256-024-00792-z

Ruff, K. M., & Pappu, R. V. (2021). AlphaFold and implications for intrinsically disordered proteins. *Journal of molecular biology*, *433*(20), 167208.

Sinha, S., Ambur, O. H., Langford, P. R., Tønjum, T., & Kroll, J. S. (2008). Reduced DNA binding and uptake in the absence of DsbA1 and DsbA2 of Neisseria meningitidis due to inefficient folding of the outer-membrane secretin PilQ. *Microbiology*, *154*(1), 217-225.

Skjærven, L., Jariwala, S., Yao, X.-Q., & Grant, B. J. (2016). Online interactive analysis of protein structure ensembles with Bio3D-web. *Bioinformatics*, *32*(22), 3510–3512. https://doi.org/10.1093/bioinformatics/btw482

Skjærven, L., Yao, X.-Q., Scarabelli, G., & Grant, B. J. (2014). Integrating protein structural dynamics and evolutionary analysis with Bio3D. *BMC Bioinformatics*, *15*(1), 399. https://doi.org/10.1186/s12859-014-0399-6

Steinegger, M., Meier, M., Mirdita, M., Vöhringer, H., Haunsberger, S. J., & Söding, J. (2019). HH-suite3 for fast remote homology detection and deep protein annotation. *BMC Bioinformatics*, *20*(1), 473. https://doi.org/10.1186/s12859-019-3019-7

Steinegger, M., Mirdita, M., & Söding, J. (2019). Protein-level assembly increases protein sequence recovery from metagenomic samples manyfold. *Nature Methods*, *16*(7), 603–606. https://doi.org/10.1038/s41592-019-0437-4

Steinegger, M., & Söding, J. (2018). Clustering huge protein sequence sets in linear time. *Nature Communications*, *9*(1), 2542. https://doi.org/10.1038/s41467-018-04964-5

Team, P. (2024). Technical Report of HelixFold3 for Biomolecular Structure Prediction. *arXiv*. https://doi.org/10.48550/arXiv.2408.16975

Treangen, T. J., Ambur, O. H., Tonjum, T., & Rocha, E. P. C. (2008). The impact of the neisserial DNA uptake sequences on genome evolution and stability. *Genome Biology*, *9*(3), R60. https://doi.org/10.1186/gb-2008-9-3-r60

Varadi, M., Tsenkov, M., & Velankar, S. (2025). Challenges in bridging the gap between protein structure prediction and functional interpretation. *Proteins: Structure, Function, and Bioinformatics*, *93*(1), 400-410.

Wohlwend, J., Corso, G., Passaro, S., Reveiz, M., Leidal, K., Swiderski, W., Portnoi, T., Chinn, I., Silterra, J., Jaakkola, T., & Barzilay, R. (2024). Boltz-1: Democratizing

Biomolecular Interaction Modeling. bioRxiv.

https://doi.org/10.1101/2024.11.19.624167

Zhang, Y., & Skolnick, J. (2004). Scoring function for automated assessment of protein

structure template quality. *Proteins: Structure, Function, and Bioinformatics*, *57*(4),

702–710. https://doi.org/10.1002/prot.20264

Supporting information captions

Table S1   ComPs used for the different species and the paired DUS-dialect.

Table S2   The different MSA schemes used in Chai-1, listing the various settings.

Figure S1   Representative ComP-DUS structures showing six different binding modes
Epsilon, Gamma, Theta, Zeta, Delta and Eta modeled in AF3, Chai-1 and RF2NA.

Figure S2   Alignment of the ComPs, using default settings in MAFFT v. 7.505,
visualized in MVIEW v. 1.68. First 28 amino acids of the N. meningitidis mature ComP are
trimmed and the other ComPs accordingly.

Figure S3   Selected high-ranking HADDOCK models, representative of the most
abundant binding mode of each ComP$_{nat}$. A: N. muc$_{nat}$ (Epsilon). B: N. men$_{nat}$ (Epsilon). C:
K. den$_{nat}$ (Epsilon). D: E. cor$_{nat}$ (Epsilon). E: B. den$_{nat}$ (Epsilon). F: N. cin$_{nat}$ (Gamma). G: N.
sub$_{nat}$ (Gamma). AF3, Chai-1 and RF2NA DNA binding modes were used to guide the
assignment of grooves in the HADDOCK distorted DNAs.

Figure S4   Representative PAE plots of the top ranking native AF3 model for each
ComP$_{nat}$ and assigned mode.

Figure S5   Representative PAE plots of the top ranking native Chai-1 model for each
ComP$_{nat}$ and assigned mode.

Figure S6   Representative PAE plots of the top ranking native RF2NA model for each
ComP$_{nat}$ and assigned mode.

Figure S7   Representative PAE plots of the top ranking scrambled AF3 model for each
ComP$_{scr}$ and assigned mode.

923   **Code Output S1**   Wilcoxon rank-sum tests on PAE and pLDDT, comparing AF3, Chai-1

924   and RF2NA. Median PAE and pLDDT values are also shown.

925   **Code Output S2**   Wilcoxon rank-sum tests on ipTM and CPPM for each $ComP_{nat}$.

926   **Code Output S3**   Wilcoxon rank-sum test on DockQ scores for the internal platform

927   consistency check.

RF2NA	Chai-1	AlphaFold 3

A

N. sub_nat
8.34 ≤ PAE ≤ 9.98
79.69 ≤ plDDT ≤ 84.08

N. sub_nat
0.60 ≤ ipTM ≤ 0.84
2.87 ≤ PAE ≤ 4.78
81.4 ≤ plDDT ≤ 89.00

N. sub_nat
0.61 ≤ ipTM ≤ 0.79
3.36 ≤ PAE ≤ 4.73
83.00 ≤ plDDT ≤ 90.54

B

B. den_nat
9.76 ≤ PAE ≤ 9.99
83.06 ≤ plDDT ≤ 85.2

B. den_nat
0.60 ≤ ipTM ≤ 0.69
4.01 ≤ PAE ≤ 4.42
86.22 ≤ plDDT ≤ 87.

B. den_nat
0.70 ≤ ipTM ≤ 0.81
3.28 ≤ PAE ≤ 5.04
82.94 ≤ plDDT ≤ 90.34

C

K. den_nat
8.71 ≤ PAE ≤ 9.94
81.15 ≤ plDDT ≤ 84.33

K. den_nat
0.60 ≤ ipTM ≤ 0.78
3.20 ≤ PAE ≤ 3.86
83.34 ≤ plDDT ≤ 87.6

K. den_nat
0.65 ≤ ipTM ≤ 0.82
2.93 ≤ PAE ≤ 4.63
85.85 ≤ plDDT ≤ 92.64

D

N. muc_nat
9.28 ≤ PAE ≤ 9.99
81.98 ≤ plDDT ≤ 84.81

N. muc_nat
0.38 ≤ ipTM ≤ 0.51 *
4.16 ≤ PAE ≤ 5.23
81.45 ≤ plDDT ≤ 85.24

N. muc_nat
0.63 ≤ ipTM ≤ 0.78
3.62 ≤ PAE ≤ 4.37
81.20 ≤ plDDT ≤ 84.35

E

E. cor_nat
8.77 ≤ PAE ≤ 9.99
80.76 ≤ plDDT ≤ 85.06

E. cor_nat
0.60 ≤ ipTM ≤ 0.75
2.68 ≤ PAE ≤ 4.51
83.72 ≤ plDDT ≤ 89.67

E. cor_nat
0.65 ≤ ipTM ≤ 0.74
3.49 ≤ PAE ≤ 4.68
82.75 ≤ plDDT ≤ 87.65

F

N. cin_nat
8.99 ≤ PAE ≤ 9.99
81.49 ≤ plDDT ≤ 84.96

N. cin_nat
0.60 ≤ ipTM ≤ 0.77
2.70 ≤ PAE ≤ 4.38
81.50 ≤ plDDT ≤ 87.5s

N. cin_nat
0.34 ≤ ipTM ≤ 0.51 *
4.92 ≤ PAE ≤ 6.33
80.55 ≤ plDDT ≤ 83.66

G

N. men_nat
8.55 ≤ PAE ≤ 9.78
82.62 ≤ plDDT ≤ 85.25

N. men_nat
0.60 ≤ ipTM ≤ 0.63
3.26 ≤ PAE ≤ 3.37
86.08 ≤ plDDT ≤ 86.5o

N. men_nat
0.33 ≤ ipTM ≤ 0.46 *
4.92 ≤ PAE ≤ 6.33
76.74 ≤ plDDT ≤ 82.