Can AI modelling of protein structures distinguish between sensor
and helper NLR immune receptors?

AmirAli Toghani[1], Ryohei Terauchi[2,3], Sophien Kamoun[1*], Yu Sugihara[2*]

[1] The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, United Kingdom
[2] Iwate Biotechnology Research Center, Kitakami, Iwate, Japan
[3] Crop Evolution Laboratory, Kyoto University, Muko, Kyoto, Japan

*Corresponding authors: sophien.kamoun@tsl.ac.uk (S.K) and sugihara.yu.85s@kyoto-u.jp
(Y.S.)

ORCID IDs
AmirAli Toghani: 0000-0002-0644-699X
Ryohei Terauchi: 0000-0002-0095-4651
Sophien Kamoun: 0000-0002-0290-0315
Yu Sugihara: 0000-0001-6042-1091

ABSTRACT

NLR immune receptors can be functionally organized in genetically linked sensor-helper pairs.
However, methods to categorize paired NLRs remain limited, primarily relying on the presence of
non-canonical domains in some sensor NLRs. Here, we propose that the AI system AlphaFold 3
can classify paired NLR proteins into sensor or helper categories based on predicted structural
characteristics. Helper NLRs showed higher AlphaFold 3 confidence scores than sensors when
modelled in oligomeric configurations. Furthermore, funnel-shaped structures—essential for
activating immune responses—were reliably predicted in helpers but not in sensors. Applying this
method to uncharacterized NLR pairs from rice, we found that AlphaFold 3 can differentiate
between putative sensors and helpers even when both proteins lack non-canonical domain
annotations. These findings suggest that AlphaFold 3 offers a new approach to categorize NLRs
and enhances our understanding of the functional configurations in plant immune systems, even
in the absence of non-canonical domain annotations.

INTRODUCTION

NLR (nucleotide binding and leucine-rich repeat) proteins are intracellular immune receptors that occur across all kingdoms of life but are particularly highly diversified in plants (Barragan and Weigel, 2021). These proteins function as singletons, pairs, or networks (Adachi et al., 2019b; Contreras et al., 2023a). Singleton NLRs can detect pathogens and execute hypersensitive cell death and immune responses, while paired NLRs have subfunctionalized into sensor (pathogen detection) and helper (immune execution) NLRs that carry distinct biochemical activities. In monocots, paired NLRs typically emerge from distinct phylogenetic clades to function together and do not have a common phylogenetic origin, unlike NLR networks (Contreras et al., 2023a; Kourelis et al., 2021). Sensor and helper NLR pairs are often genetically clustered, and some sensors have non-canonical integrated domains (IDs) that function in pathogen sensing and are absent in helper NLRs (Białas et al., 2018; Marchal et al., 2022). However, the presence of IDs is currently the only effective way to distinguish sensor NLRs from helper NLRs *in silico*, and no other method has been established to distinguish sensors from helpers in paired NLRs.

In plants, NLRs that carry a coiled-coil (CC) domain at their N-termini are the most widespread class. Following pathogen recognition, CC-NLR proteins oligomerize into pentameric or hexameric pore-like complexes (Förderer et al., 2022; Liu et al., 2024; Madhuprakash et al., 2024; Wang et al., 2019; Zhao et al., 2022). These complexes, known as resistosomes, are a defining feature of NLRs that execute the immune response such as the helpers; they translocate to cellular membranes and trigger immune responses such as calcium influx and hypersensitive cell death (Contreras et al., 2022; Duggan et al., 2021; Ibrahim et al., 2024). The prevailing model is that the funnel-shaped structure of CC-NLR resistosomes inserts into membranes and is required for cell death and immune response (Adachi et al., 2019b; Förderer and Kourelis, 2023; Wang et al., 2019). However, this funnel-shaped structure is formed by the N-terminal α1 helix of NLR, which is a structurally dynamic region that is difficult to resolve using cryo-electron microscopy (Förderer et al., 2022; Liu et al., 2024; Madhuprakash et al., 2024; Zhao et al., 2022). About 20% of plant CC-NLRs have a conserved sequence motif, called MADA, at the N-terminal α1 helix, while a previous study indicates that MADA motifs have degenerated in some sensor NLRs of solanaceous plants (Adachi et al., 2019a).

AlphaFold revolutionized protein structural biology by accurately modelling protein structures from primary sequences (Jumper et al., 2021). Recently, AlphaFold 3 (AF3) was released, enabling the modelling of protein structures with oleic acids serving as proxy for cellular membranes (Abramson et al., 2024). This function is particularly useful for proteins that dynamically interact with cellular membranes, such as CC-NLRs, and has enabled high-confidence modelling of CC-NLR resistosomes (Madhuprakash et al., 2024; Ibrahim et al., 2024).

Here, we leveraged AF3 to explore the structural diversity of sensor and helper oligomers of a

curated set of CC-NLRs consisting of i) experimentally validated NLR pairs in rice (Pikm, Pii, and

Pia), ii) their orthologs (PIK5/6-NP, Pi5-3/1, and Pias), and iii) two previously cloned NLR pairs in

barley (RPG5/HvRGA1 and RGH2/3) as summarized in **Table S1**. Our analyses revealed that

helpers consistently exhibit higher AF3 confidence scores than sensors. Moreover, the funnel-

shaped structures, crucial for the immune response, were consistently observed in helpers but

not in sensors. We extended our investigations to uncharacterized NLR pairs of rice, and showed

that AF3 can distinguish between NLR pairs even when both proteins do not carry any

annotations for unconventional integrated domains. Based on these findings, we propose that

AF3 can be used to classify paired NLRs into functional categories, providing a novel approach

to understanding functional configurations of plant immune receptors.

RESULTS

Helper NLRs produce higher AlphaFold 3 pTM scores than their paired sensors

To benchmark the AF3 capacity to distinguish between sensor and helper NLRs, we curated

eight NLR pairs in which the sensor can be readily assigned because it carries an ID (**Fig. S1**;

Table S1). As in previous studies (Madhuprakash et al., 2024; Ibrahim et al., 2024), we used the

oligomerizing domains of the NLR proteins, from the N-terminus to the end of the NB-ARC

domain, and performed AF3 predictions of 5x and 6x stoichiometries with 50 oleic acids and

using three different seed values. We then compared the sensor and helper predicted template

modelling (pTM) scores (**Fig. 1a and b**; **Fig. S2–4**; **Table S2**). Helper NLRs consistently

exhibited higher pTM scores than sensor NLRs in both pentameric and hexameric

configurations, indicating that AF3 pTM scores can be used to distinguish between sensor and

helper NLRs.

Helper AF3 structures form funnel-shaped structures unlike sensors

In addition to the AF3 confidence scores, we also examined the AF3 structures for distinct

structural patterns between sensor and helper NLRs (**Fig. 1c**; **Fig. S5 and S6**). Helper NLRs

consistently formed funnel-shaped structures in contrast to sensor NLRs. This observation aligns

with previous studies demonstrating that sensor NLRs cannot execute the immune response on

their own and may have lost the capacity to oligomerise into resistosome-like structures (Adachi

et al., 2019b; Contreras et al., 2022). Sensors formed resistosome-like structures but with low

confidence scores (**Fig. 1c**; **Fig. S5 and S6**). These findings further indicate that AF3 can

distinguish between sensor and helper NLRs.

We then analyzed whether the presence of the MADA motif could classify sensors or helpers in

the curated NLR pairs using the MADA HMM model in Adachi et al., 2019a (**Table S3**). The

helpers of Pikm and PIK5/6-NP contained the MADA motif with HMM scores exceeding the cut-

off value of 10, while the other sensors and helpers did not. Based on these results, only Pikm

and PIK5/6-NP could be classified according to the presence of the MADA motif, whereas the

remaining NLR pairs could not be classified. These findings indicate that AF3 modelling-based

classification is more reliable than the sequence-based classification and highlight the utility of

AF3 for NLR classification.

AF3 can classify paired NLRs even when the putative sensors lack an integrated domain

annotation

Based on Stein et al., 2018, we extracted ten rice CC-NLR pairs that i) are genetically linked in

head-to-head orientations; ii) belong to distinct phylogenetic clades; and iii) carry a full N-terminal

CC domain (**Fig. 2a**; **Fig. S1; Table S4–S6**). In five of the ten pairs, one of the NLRs carries an

ID annotation and is presumed to be the sensor. The putative helpers (without ID annotation) had

higher AF3 pTM scores than sensors (with ID annotation) for these pairs (**Fig. 2b**; **Fig. S7; Table**

S7). Furthermore, putative helpers formed funnel-shaped structures, whereas putative sensors

did not (**Fig. 2c**; **Fig. S8 and S9**). These results further confirm the application of AF3 for

functional classification as noted with the eight previously characterized pairs.

Next, we examined the five remaining pairs whose NLRs do not carry any ID annotations. In

each pair, we averaged the pTM scores of pentameric and hexameric configurations and

assigned them as putative helpers or sensors based on higher or lower pTM scores, respectively

(**Fig. 2b**; **Fig. S1 and S7; Table S7**). Pair-13 is identical to the rice pair PIK5/6-NP, in which

PIK5-NP is known as a sensor with an integrated domain that is not annotated by InterProScan

(Białas et al., 2021; Kourelis et al., 2021). Nonetheless, AF3 successfully classified them into

helper or sensor based on the pTM scores even without the ID annotation. Among the other four

pairs, Pair-07 exhibited a consistent difference in the pTM scores between the putative sensor

and helper NLRs (**Fig. S7; Table S7**).

DISCUSSION

AlphaFold 3 can potentially distinguish between sensor and helper NLRs based on their

predicted structures and confidence scores. In AF3 predictions, helpers consistently exhibited

higher AF3 confidence scores than sensors (**Fig. 1**; **Fig. S2–4**). Moreover, the funnel-shaped

structures were consistently observed in helpers but not in sensors (**Fig. 1c and 2c**; **Fig. S5, S6,**

S8, and S9). These findings indicate that AF3 might be able to distinguish the functional

configurations of plant immune receptors. Our results highlight the utility of AF3 as a powerful

tool for distinguishing structural and functional differences in paired NLRs.

Why did sensor NLRs exhibit lower AF3 confidence scores than helper NLRs? Previous studies

proposed a model suggesting that paired NLRs evolved from singleton NLRs (Adachi et al.,

2019b). In this model, singleton NLRs, which can detect pathogens (sensor) and execute

hypersensitive cell death (helper), subfunctionalized into sensor or helper NLRs (Adachi et al.,

2019b; Contreras et al., 2023a). This subfunctionalization results in the loss of cell death activity

in sensor NLRs and the loss of pathogen perception activity in helper NLRs. Since the formation

of resistosomes with funnel-shaped structures is essential for induction of cell death and other

immune responses, the presence of stable resistosome and funnel-shaped structures in helper

NLRs, but not in sensor NLRs, supports the previously proposed model. Consequently, sensor

NLRs exhibit lower AF3 confidence scores than helper NLRs, reflecting their evolutionary

divergence and functional specialization.

Recently, several researchers have applied AlphaFold to plant-pathogen interactions (Contreras

et al., 2023b; Cruz et al., 2024; Ibrahim et al., 2024, 2023; Madhuprakash et al., 2024; Selvaraj

et al., 2024; Seong et al., 2024; Sugihara et al., 2023; Tamborski et al., 2023). These

advancements provide a novel approach to understanding functional configurations of immune

receptors beyond domain architectures, sequence motifs, and phylogenetic relationships.

METHODS

NLR annotation

NLRs were annotated using NLRtracker v1.0.3 (Kourelis et al., 2021) and InterProScan v5.67-

99.0 (Jones et al., 2014). Note that InterProScan did not annotate an integrated domain of PIK5-

NP as previously reported (Kourelis et al., 2021). Therefore, we manually replaced the domain

architecture of PIK5-NP from "CNL" to "CONL" as it contains the HMA domain between the NB-

ARC and LRR (Białas et al., 2021). The domain architectures were visualized using refplantnlR

(https://github.com/JKourelis/refplantnlR). The MADA motifs were analyzed using the HMM

model in Adachi et al., 2019a and HMMER v3.4 (http://hmmer.org) with the option "--max". The

NLRtracker and HMMER outputs are archived on Zenodo

(https://doi.org/10.5281/zenodo.13826775).

Curation of NLR sequences

The sequences of curated NLR pairs were derived from either RefPlantNLR (Kourelis et al.,

2021) or NCBI (**Table S1**). Regarding the NLR pairs described by Stein et al., 2018, the protein

sequences of *Oryza sativa* cv. Nipponbare (Oryza_sativa_vg_japonica.protein.fasta) were

downloaded from the URL (https://doi.org/10.7946/P2FC9Z). Based on Supplementary Data 6 in

Stein et al., 2018, the NLR pairs that i) are genetically linked in head-to-head orientations; ii)

belong to distinct phylogenetic clades were extracted (**Table S4**). Using DIAMOND BLASTP

v2.1.9 (Buchfink et al., 2021), the corresponding NLR sequences were identified from the NCBI

RefSeq annotation of rice cultivar Nipponbare genome (GCF_034140825.1). The best-hit

sequences were summarized in **Table S5** and confirmed to be genetically linked in a head-to-

head orientation. Based on NLRtracker outputs, the CC-NLR pairs, which carry a full N-terminal

CC domain, were retained (**Table S6**). Piar-05, Piar-13, and the sensor of Pair-14 correspond to

Pia, PIK5/6-NP, and Pi-ta, respectively (**Table S4 and S6**).

AF3 prediction

Based on NLRtracker outputs, the amino acid sequences from the N-terminus to the end of the

NB-ARC domain were extracted. Using the AlphaFold 3 webserver

(https://golgi.sandbox.google.com), the extracted sequences were modelled in both pentameric

and hexameric configurations with 50 oleic acids using three different seed values. The input

sequences and resulting models are archived on Zenodo

(https://doi.org/10.5281/zenodo.13826775).

Phylogenetic analysis

We built the tree of monocot paired NLRs with RefPlantNLR (Kourelis et al., 2021) and a set of

4,936 NLR proteins from the NLRtracker output of 13 RefSeq proteomes, including two dicot

species (*Arabidopsis thaliana* and *Solanum lycopersicum*) and 11 monocot species (*Zea mays*,

Triticum aestivum, *Setaria viridis*, *Phragmites australis*, *Phoenix dactylifera*, *Oryza sativa*, *Musa*

acuminata, *Lolium perenne*, *Hordeum vulgare* subsp. *vulgare*, *Brachypodium distachyon*, and

Asparagus officinalis) (**Data S1, S2, S3, and S4**) (Toghani and Kamoun, 2024). More details on

the phylogenetics analysis are available on GitHub

(https://github.com/amiralito/Paired_NLR_AF3).

FUNDING

The authors received funding from the sources listed below. The Gatsby Charitable Foundation

(A.T. and S.K.), Biotechnology and Biological Sciences Research Council (BBSRC) BB/P012574

(Plant Health ISP) (AT and SK), BBSRC BBS/E/J/000PR9795 (Plant Health ISP - Recognition)

(A.T. and S.K.), BBSRC BBS/E/J/000PR9796 (Plant Health ISP - Response) (A.T. and S.K.),

BBSRC BBS/E/J/000PR9797 (Plant Health ISP – Susceptibility) (A.T. and S.K.), BBSRC

BBS/E/J/000PR9798 (Plant Health ISP – Evolution) (A.T. and S.K.), European Research Council

(ERC) 743165 (S.K.), Engineering and Physical Sciences Research Council EP/Y032187/1

(S.K.), and Japan Society for the Promotion of Science (JSPS) 23K20042, 24H00010 (R.T.).

ACKNOWLEDGEMENTS

We thank Tolga Bozkurt (Imperial College London), Andres Posbeyikian (The Sainsbury

Laboratory), Raoul Frijters (Rijk Zwaan Breeding B.V.), and Kentato Fujita (Osaka University) for

valuable comments and scientific discussions.

CONFLICT OF INTERESTS

S.K. receives funding from industry on NLR biology and has co-founded a start-up company

(Resurrect Bio Ltd.) related to NLR biology. S.K. has filed patents on NLR biology.

DATA AVAILABILITY

The scripts and dataset used for the phylogenetic analysis are available on GitHub

(https://github.com/amiralito/Paired_NLR_AF3). The other datasets are archived on Zenodo

(https://doi.org/10.5281/zenodo.13826775).

REFERENCES

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard
AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C, O'Neill M, Reiman D,
Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O, Bridgland A,
Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB,
Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S,
Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A, Bapst
V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of
biomolecular interactions with AlphaFold 3. *Nature* **630**:493–500. doi:10.1038/s41586-
024-07487-w
Adachi H, Contreras MP, Harant A, Wu C, Derevnina L, Sakai T, Duggan C, Moratto E, Bozkurt
TO, Maqbool A, Win J, Kamoun S. 2019a. An N-terminal motif in NLR immune receptors
is functionally conserved across distantly related plant species. *eLife* **8**:e49956.
doi:10.7554/eLife.49956
Adachi H, Derevnina L, Kamoun S. 2019b. NLR singletons, pairs, and networks: evolution,
assembly, and regulation of the intracellular immunoreceptor circuitry of plants. *Curr Opin
Plant Biol* **50**:121–131. doi:10.1016/j.pbi.2019.04.007
Barragan AC, Weigel D. 2021. Plant NLR diversity: the known unknowns of pan-NLRomes. *Plant
Cell* **33**:814–831. doi:10.1093/plcell/koaa002
Białas A, Langner T, Harant A, Contreras MP, Stevenson CE, Lawson DM, Sklenar J, Kellner R,
Moscou MJ, Terauchi R, Banfield MJ, Kamoun S. 2021. Two NLR immune receptors

acquired high-affinity binding to a fungal effector through convergent evolution of their integrated domain. *eLife* **10**:e66961. doi:10.7554/eLife.66961

Białas A, Zess EK, De la Concepcion JC, Franceschetti M, Pennington HG, Yoshida K, Upson JL, Chanclud E, Wu C-H, Langner T, Maqbool A, Varden FA, Derevnina L, Belhaj K, Fujisaki K, Saitoh H, Terauchi R, Banfield MJ, Kamoun S. 2018. Lessons in Effector and NLR Biology of Plant-Microbe Systems. *Mol Plant-Microbe Interactions®* **31**:34–45. doi:10.1094/MPMI-08-17-0196-FI

Buchfink B, Reuter K, Drost H-G. 2021. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nat Methods* **18**:366–368. doi:10.1038/s41592-021-01101-x

Contreras MP, Lüdke D, Pai H, Toghani A, Kamoun S. 2023a. NLR receptors in plant immunity: making sense of the alphabet soup. *EMBO Rep* **24**:e57495. doi:10.15252/embr.202357495

Contreras MP, Pai H, Selvaraj M, Toghani A, Lawson DM, Tumtas Y, Duggan C, Yuen ELH, Stevenson CEM, Harant A, Maqbool A, Wu C-H, Bozkurt TO, Kamoun S, Derevnina L. 2023b. Resurrection of plant disease resistance proteins via helper NLR bioengineering. *Sci Adv* **9**:eadg3861. doi:10.1126/sciadv.adg3861

Contreras MP, Pai H, Tumtas Y, Duggan C, Yuen ELH, Cruces AV, Kourelis J, Ahn H-K, Lee K-T, Wu C-H, Bozkurt TO, Derevnina L, Kamoun S. 2022. Sensor NLR immune proteins activate oligomerization of their NRC helpers in response to plant pathogens. *EMBO J* **n/a**:e111519. doi:10.15252/embj.2022111519

Cruz DGDL, Zdrzałek R, Banfield MJ, Talbot NJ, Moscou MJ. 2024. Molecular mimicry of a pathogen virulence target by a plant immune receptor. doi:10.1101/2024.07.26.605320

Duggan C, Moratto E, Savage Z, Hamilton E, Adachi H, Wu C-H, Leary AY, Tumtas Y, Rothery SM, Maqbool A, Nohut S, Martin TR, Kamoun S, Bozkurt TO. 2021. Dynamic localization of a helper NLR at the plant–pathogen interface underpins pathogen recognition. *Proc Natl Acad Sci* **118**:e2104997118. doi:10.1073/pnas.2104997118

Förderer A, Kourelis J. 2023. NLR immune receptors: structure and function in plant disease resistance. *Biochem Soc Trans* **51**:1473–1483. doi:10.1042/BST20221087

Förderer A, Li E, Lawson AW, Deng Y, Sun Y, Logemann E, Zhang X, Wen J, Han Z, Chang J, Chen Y, Schulze-Lefert P, Chai J. 2022. A wheat resistosome defines common principles of immune receptor channels. *Nature* **610**:532–539. doi:10.1038/s41586-022-05231-w

Ibrahim T, Khandare V, Mirkin FG, Tumtas Y, Bubeck D, Bozkurt TO. 2023. AlphaFold2-multimer guided high-accuracy prediction of typical and atypical ATG8-binding motifs. *PLOS Biol* **21**:e3001962. doi:10.1371/journal.pbio.3001962

Ibrahim T, Yuen ELH, Wang H-Y, King FJ, Toghani A, Kourelis J, Vuolo C, Adamkova V, Castel B, Jones JD, Wu C-H, Kamoun S, Bozkurt TO. 2024. A helper NLR targets organellar membranes to trigger immunity. doi:10.1101/2024.09.19.613839

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong S-Y, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**:1236–1240. doi:10.1093/bioinformatics/btu031

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**:583–589. doi:10.1038/s41586-021-03819-2

Kourelis J, Sakai T, Adachi H, Kamoun S. 2021. RefPlantNLR is a comprehensive collection of experimentally validated plant disease resistance proteins from the NLR family. *PLOS Biol* **19**:e3001124. doi:10.1371/journal.pbio.3001124

Liu F, Yang Z, Wang C, You Z, Martin R, Qiao W, Huang J, Jacob P, Dangl JL, Carette JE, Luan S, Nogales E, Staskawicz BJ. 2024. Activation of the helper NRC4 immune receptor forms a hexameric resistosome. *Cell* **187**:4877-4889.e15. doi:10.1016/j.cell.2024.07.013

Madhuprakash J, Toghani A, Contreras MP, Posbeyikian A, Richardson J, Kourelis J, Bozkurt TO, Webster M, Kamoun S. 2024. A disease resistance protein triggers oligomerization of its NLR helper into a hexameric resistosome to mediate innate immunity.

Marchal C, Michalopoulou VA, Zou Z, Cevik V, Sarris PF. 2022. Show me your ID: NLR immune
receptors with integrated domains in plants. *Essays Biochem* EBC20210084.
doi:10.1042/EBC20210084
Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023. UCSF
ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**:e4792.
doi:10.1002/pro.4792
Selvaraj M, Toghani A, Pai H, Sugihara Y, Kourelis J, Yuen ELH, Ibrahim T, Zhao H, Xie R,
Maqbool A, Concepcion JCD la, Banfield MJ, Derevnina L, Petre B, Lawson DM, Bozkurt
TO, Wu C-H, Kamoun S, Contreras MP. 2024. Activation of plant immunity through
conversion of a helper NLR homodimer into a resistosome. *PLOS Biol* **22**:e3002868.
doi:10.1371/journal.pbio.3002868
Seong K, Wei W, Vega B, Dee A, Ramirez-Bernardino G, Kumar R, Parra L, Krasileva K. 2024.
Engineering the plant intracellular immune receptor Sr50 to restore recognition of the
AvrSr50 escape mutant. doi:10.1101/2024.08.07.607039
Stein JC, Yu Y, Copetti D, Zwickl DJ, Zhang L, Zhang C, Chougule K, Gao D, Iwata A,
Goicoechea JL, Wei S, Wang J, Liao Y, Wang M, Jacquemin J, Becker C, Kudrna D,
Zhang J, Londono CEM, Song X, Lee S, Sanchez P, Zuccolo A, Ammiraju JSS, Talag J,
Danowitz A, Rivera LF, Gschwend AR, Noutsos C, Wu C, Kao S, Zeng J, Wei F, Zhao Q,
Feng Q, El Baidouri M, Carpentier M-C, Lasserre E, Cooke R, Rosa Farias D da, da Maia
LC, dos Santos RS, Nyberg KG, McNally KL, Mauleon R, Alexandrov N, Schmutz J,
Flowers D, Fan C, Weigel D, Jena KK, Wicker T, Chen M, Han B, Henry R, Hsing YC,
Kurata N, de Oliveira AC, Panaud O, Jackson SA, Machado CA, Sanderson MJ, Long M,
Ware D, Wing RA. 2018. Genomes of 13 domesticated and wild rice relatives highlight
genetic conservation, turnover and innovation across the genus Oryza. *Nat Genet*
50:285–296. doi:10.1038/s41588-018-0040-0
Sugihara Y, Abe Y, Takagi H, Abe A, Shimizu M, Ito K, Kanzaki E, Oikawa K, Kourelis J, Langner
T, Win J, Białas A, Lüdke D, Contreras MP, Chuma I, Saitoh H, Kobayashi M, Zheng S,
Tosa Y, Banfield MJ, Kamoun S, Terauchi R, Fujisaki K. 2023. Disentangling the complex
gene interaction networks between rice and the blast fungus identifies a new pathogen
effector. *PLOS Biol* **21**:e3001945. doi:10.1371/journal.pbio.3001945
Tamborski J, Seong K, Liu F, Staskawicz BJ, Krasileva KV. 2023. Altering Specificity and
Autoactivity of Plant Immune Receptors Sr33 and Sr50 Via a Rational Engineering
Approach. *Mol Plant-Microbe Interactions®* **36**:434–446. doi:10.1094/MPMI-07-22-0154-
R
Toghani A, Kamoun S. 2024. Functional annotation of 180 RefSeq reference plant proteomes
reveals a dataset of 113,684 NLR proteins. doi:10.5281/zenodo.13627395
Wang Jizong, Hu M, Wang Jia, Qi J, Han Z, Wang G, Qi Y, Wang H-W, Zhou J-M, Chai J. 2019.
Reconstitution and structure of a plant NLR resistosome conferring immunity. *Science*
364:eaav5870. doi:10.1126/science.aav5870
Zhao Y-B, Liu M-X, Chen T-T, Ma X, Li Z-K, Zheng Z, Zheng S-R, Chen L, Li Y-Z, Tang L-R, Chen
Q, Wang P, Ouyang S. 2022. Pathogen effector AvrSr35 triggers Sr35 resistosome
assembly via a direct recognition mechanism. *Sci Adv* **8**:eabq5108.
doi:10.1126/sciadv.abq5108
372
373

374
Fig. 1. Helper NLRs produce higher AlphaFold 3 pTM scores than their paired sensors. a) Bar plot
comparing sensor and helper pTM scores in pentameric AlphaFold 3 predictions. The amino acid sequences of
the oligomerizing domains of the NLR proteins, from the N-terminus to the end of the NB-ARC domain, were
used for the prediction. The pentameric structures were modelled with 50 oleic acids using three different seed
values. b) Scatter plot comparing sensor and helper pTM scores in pentameric AlphaFold 3 predictions. The
resulting pTM scores were averaged across three seed values for each sensor and helper NLR. c) Pentameric
AlphaFold 3 structures of the four representative NLR pairs. The structures predicted using seed value 1 were
visualized using ChimeraX (Meng et al., 2023).

383
Fig. 2. AF3 can classify paired NLRs even when the putative sensors do not carry an integrated domain

annotation. a) Domain architectures of ten rice NLR pairs described by Stein et al., 2018. The domains were

annotated with NLRtracker (Kourelis et al., 2021) and visualized with refplantnlR

(https://github.com/JKourelis/refplantnlR). b) Bar plot comparing putative sensor and helper pTM scores in

pentameric AlphaFold 3 predictions. The amino acid sequences of the oligomerizing domains of the NLR

proteins, from the N-terminus to the end of the NB-ARC domain, were used for the prediction. The pentameric

structures were modelled with 50 oleic acids using three different seed values. Putative sensors and helpers

were assigned based on the average pTM scores of pentameric and hexameric structures, with a putative sensor

having the lower average score and a putative helper having the higher average score. c) Pentameric AlphaFold

3 structures of the four representative NLR pairs described by Stein et al., 2018. The structures predicted using

seed value 1 were visualized using ChimeraX (Meng et al., 2023).

Fig. S1. Phylogenetic tree of monocot paired NLRs with RefPlantNLR and 4,936 NLR proteins from 13

RefSeq proteomes, including two dicot species and 11 monocot species obtained from Toghani et al.,

2024. The tree was built using the NB-ARC domains of the eight known NLR pairs summarized in **Table S1**,

RefPlantNLR, and 4,936 NLR proteins from the NLRtracker output of 13 RefSeq proteomes, including two dicot

species (*Arabidopsis thaliana* and *Solanum lycopersicum*) and 11 monocot species (*Zea mays*, *Triticum*

aestivum, *Setaria viridis*, *Phragmites australis*, *Phoenix dactylifera*, *Oryza sativa*, *Musa acuminata*, *Lolium*

perenne, *Hordeum vulgare* subsp. *vulgare*, *Brachypodium distachyon*, and *Asparagus officinalis*) (**Data S1, S2,**

S3, and S4) (Toghani and Kamoun, 2024). The paired NLRs of rice cultivar Nipponbare (GCF_034140825.1),

described by Stein et al., 2018, are included among 13 RefSeq proteomes. The genetic linkage between NLR

pairs is indicated by a connection between linked nodes. Known sensor and helper NLR nodes are colored blue

and orange, respectively, while putative sensor and helper nodes are colored green and yellow. Numbers next to

the tree nodes indicate bootstrap values. The tree is rooted at the CC_R-NLR clade. AlphaFold 3 was not

performed for Pair-02, Pair-03, Pair-08, Pair-09, and Pair-10 because they do not carry a full N-terminal CC

domain.

411
412
Fig. S2. Correlations between pentameric and hexameric AlphaFold 3 confidence scores and between
pTM and ipTM scores for previously reported NLR pairs. a) Correlation between the average pentameric and
hexameric pTM scores. b) Correlation between the average pentameric and hexameric ipTM scores. c)
Correlation between the pentameric pTM and ipTM scores. d) Correlation between the hexameric pTM and ipTM
scores. The amino acid sequences of the oligomerizing domains of the NLR proteins, from the N-terminus to the
end of the NB-ARC domain, were used for the prediction. The pentameric and hexameric structures were
modelled with 50 oleic acids using three different seed values. The resulting pTM or ipTM scores were averaged
across three seed values for each NLR in Fig. S2a and S2b. The Pearson rank correlation coefficients (*r*) were
calculated with the SciPy library in Python.

422
423
Fig. S3. Comparisons of sensor and helper AlphaFold 3 scores in pentameric and hexameric
configurations for previously reported NLR pairs. a-d) pTM score comparisons for pentamers (a,b) and
hexamers (c,d). e-h) ipTM score comparisons for pentamers (e,f) and hexamers (g,h). The a, c, e, and g panels
show bar plots, and the b, d, f, and h panels display score differences (helper minus sensor). The amino acid
sequences of the oligomerizing domains of the NLR proteins, from the N-terminus to the end of the NB-ARC
domain, were used for the prediction. The pentameric and hexameric structures were modelled with 50 oleic
acids using three different seed values. Subtractions were performed for all possible pairs of sensor and helper
scores.

Fig. S4. Scatter plots comparing average sensor and helper scores in AlphaFold 3 predictions. a) Scatter

plot comparing average sensor and helper pTM scores in pentamer predictions. b) Scatter plot comparing

average sensor and helper pTM scores in hexamer predictions. c) Scatter plot comparing average sensor and

helper ipTM scores in pentamer predictions. d) Scatter plot comparing average sensor and helper ipTM scores in

hexamer predictions. The amino acid sequences of the oligomerizing domains of the NLR proteins, from the N-

terminus to the end of the NB-ARC domain, were used for the prediction. The pentameric and hexameric

structures were modelled with 50 oleic acids using three different seed values. The resulting pTM or ipTM scores

were averaged across three seed values for each sensor and helper NLR.

443
444
445 **Fig. S5. Pentameric AlphaFold 3 predictions for previously reported NLR pairs.** The amino acid sequences
446 of the oligomerizing domains of the NLR proteins, from the N-terminus to the end of the NB-ARC domain, were
447 used for the prediction. The pentameric structures were modelled with 50 oleic acids using seed value 1. The
448 predicted structures were visualized using ChimeraX (Meng et al., 2023).

449
450
451 **Fig. S6. Hexameric AlphaFold 3 predictions for previously reported NLR pairs**. The amino acid sequences
452 of the oligomerizing domains of the NLR proteins, from the N-terminus to the end of the NB-ARC domain, were
453 used for the prediction. The hexameric structures were modelled with 50 oleic acids using seed value 1. The
454 predicted structures were visualized using ChimeraX (Meng et al., 2023).

Fig. S7. Comparisons of putative sensor and helper AlphaFold 3 scores in pentameric and hexameric configurations for NLR pairs described by Stein et al., 2018. a-d) pTM score comparisons for pentamers (a,b) and hexamers (c,d). e-h) ipTM score comparisons for pentamers (e,f) and hexamers (g,h). The a, c, e, and g panels show bar plots, and the b, d, f, and h panels display score differences (putative helper minus putative sensor). The pentameric and hexameric structures were modelled with 50 oleic acids using three different seed values. Putative sensors and helpers were assigned based on the average pTM scores of pentameric and hexameric structures, with a putative sensor having the lower average score and a putative helper having the higher average score. Subtractions were performed for all possible pairs of putative sensor and helper scores.

Fig. S8. Pentameric AlphaFold 3 predictions for NLR pairs described by Stein et al., 2018. The amino acid sequences of the oligomerizing domains of the NLR proteins, from the N-terminus to the end of the NB-ARC domain, were used for the prediction. The pentameric structures were modelled with 50 oleic acids using seed value 1. The predicted structures were visualized using ChimeraX (Meng et al., 2023).

Fig. S9. Hexameric AlphaFold 3 predictions for NLR pairs described by Stein et al., 2018. The amino acid

sequences of the oligomerizing domains of the NLR proteins, from the N-terminus to the end of the NB-ARC

domain, were used for the prediction. The hexameric structures were modelled with 50 oleic acids using seed

value 1. The predicted structures were visualized with ChimeraX (Meng et al., 2023).

Table S1. List of previously reported NLR pairs. All sequences were derived from either RefPlantNLR
(Kourelis et al., 2021) or NCBI. Sensor and helper NLRs are defined based on the presence or absence of an
integrated domain, respectively. Domain architectures were annotated using NLRtracker (Kourelis et al., 2021).
For domain architecture, "C," "N," "L," and "O" represent the CC, NB-ARC, LRR, and other integrated domains,
respectively. Note that InterProScan did not annotate an integrated domain of PIK5-NP as previously reported
(Kourelis et al., 2021). Therefore, we manually replaced the domain architecture of PIK5-NP from "CNL" to
"CONL" as it contains the HMA domain between the NB-ARC and LRR (Białas et al., 2021).
485
486   **Table S2. Summary of AlphaFold 3 predictions for previously reported NLR pairs.**
487
488   **Table S3. HMM scores of MADA motifs in previously reported NLR pairs.** The presence or absence of the
489   MADA or MADA-like motif was analyzed using the HMM model in Adachi et al., 2019a. MADA, MADA-like, and
490   no MADA motifs were classified based on HMM scores (> 10, < 10, and NA, respectively).
491
Table S4. List of rice NLR pairs described by Stein et al., 2018. The NLR pairs were identified from the rice
cultivar Nipponbare genome annotation (Stein et al., 2018). Based on Supplementary Data 6 in Stein et al., 2018,
the NLR pairs that i) are genetically linked in head-to-head orientations; ii) belong to distinct phylogenetic clades
were extracted. Domain architectures were annotated using NLRtracker (Kourelis et al., 2021). Columns from
Supplementary Data 6 in Stein et al., 2018 and those added in this study are highlighted in yellow and green,
respectively.
498
499   **Table S5. BLASTP results using NLRs from Stein et al., 2018 as queries against those in the NCBI RefSeq**
500   **annotation as subjects.** DIAMOND BLASTP (Buchfink et al., 2021) was used to identify corresponding
501   sequences between the datasets. NLRs from Stein et al., 2018 and those in the NCBI RefSeq annotation of the
502   rice cultivar Nipponbare (GCF_034140825.1) served as the query and subject sets, respectively.
503
504   **Table S6. Summary of NLRtracker outputs for rice NLR pairs from the NCBI RefSeq annotation of**
505   **Nipponbare (GCF_034140825.1), corresponding to those described by Stein et al., 2018.**
506   Only NLR pairs with a full N-terminal CC domain in both NLRs were processed for AlphaFold 3 predictions.
507
508   **Table S7. Summary of AlphaFold 3 predictions for rice NLR pairs from the NCBI RefSeq annotation of**
509   **Nipponbare (GCF_034140825.1), corresponding to those described by Stein et al., 2018.**
510   Putative sensors and helpers were assigned based on the average pTM scores of pentameric and hexameric
511   structures, with a putative sensor having the lower average score and a putative helper having the higher
512   average score.
513
514   **Data S1. Taxonomic and genome metadata of species used for phylogenetic analysis.**
515
516   **Data S2. Sequence and metadata of full-length and NB-ARC domains of NLR sequences used for**
517   **phylogenetic analysis.**
518
519   **Data S3. Alignment of 5,509 NB-ARC sequences of NLRs used in Fig. S1.**
520
521   **Data S4. Phylogenetic tree of 5,509 NB-ARC sequences of NLRs used in Fig. S1.**
522