Horizontal transfer of a LINE-RTE retrotransposon among parasite, host, prey and environment.

Brann, T.[1]*, de Oliveira, F. S.[1,2], Protasio, A. V.[1,3]

Affiliations:

1. Department of Pathology, University of Cambridge, Cambridge CB1 2PQ, UK.
2. Departamento de Genética, Universidade Federal do Paraná, 81531-990, Brazil.
3. Christ's College, University of Cambridge, CB2 3BU, UK.

* corresponding author. tb708@cam.ac.uk

Abstract

Background Horizontal transfer of transposable elements is both impactful, owing to the subsequent transposition burst, and insightful, providing information on organisms' evolutionary history. In eukaryotes, horizontal gene transfer (HGT) often involves transposable elements (TEs), host-parasite relationships, aquatic environments or any of them combined. The flatworm *Schistosoma mansoni* is a human parasite with two free-living aquatic stages (intercalated between a definitive human host and intermediate snail host) and has a sizable TE content. We aimed to identify and characterise potential instances of HGT leveraging new genomic resources available.

Results Using the latest chromosome-scale genome assembly and available TE sequences we identify two putatively horizontally transferred elements, named Perere-3 and Sr3, in the *S. mansoni* genome. We demonstrate the presence of these TEs in the genomes of *Schistosoma* spp. intermediate hosts, most likely explained by HGT. Perere-3 / Sr3 were also found across a wide range of additional organisms not susceptible to schistosome infection, including turtles, fish and other molluscs.

Conclusions We propose that the patchy distribution of Perere-3/Sr3 across the phylogenetic tree is best explained by HGT. This phenomenon is likely linked to the parasitic nature of schistosomes, as several snail species sharing the elements are susceptible to infection. However, presence of Perere-3/Sr3 in species beyond this relationship may suggest wider ancestral *Schistosomatidae* host ranges and/or undescribed schistosomes.

Keywords *Schistosoma mansoni*, Transposable element, Horizontal gene transfer, host-parasite relationship, *Schistosomatidae*

Introduction

Horizontal gene transfer (HGT), unlike vertical genetic inheritance, refers to DNA acquisition via a non-parental genome. Although examples are found in all realms, HGT in eukaryotes occurs at a rate approximately 80 times less than in prokaryotes (Van Etten and Bhattacharya 2020). This is due in part to additional physical barriers present in eukaryotes, such as the nuclear envelope, compartmentalisation of the germline, where the transferred genes must be present to be propagated to future generations, and, in many metazoans, a protective skin or dermis (Kurland, et al. 2003).

Host-parasite relationships, which reduce or circumvent physical distances and barriers between organisms, are overrepresented in instances of eukaryotic HGT (Gilbert, et al. 2010). Parasites live for varied, but often extended, periods of time in direct contact or close proximity to their hosts, often feeding on host tissue. This intimate host-parasite relationship provides an excellent environment for exposure to the other organism's DNA. Another common medium for DNA "sharing" is aquatic environments. In them, many organisms such as algae, corals and fish, share and sample the same body of water, favouring HGT events (Abe, et al. 2020; Pérez-Etayo, et al. 2020).

Transposable Elements (TEs) are genetic entities capable of mobilisation within genomes. They are more frequently implicated in horizontal transfer events (referred to as "horizontal transfer of TEs" or HTT) than "traditional" host genes due to their inherent capacity to encode enzymes that mediate their excision and insertion into genomes. TEs are often present in many copies and can occupy large fractions of host

genomes, such as 52% in Humans (Tang, et al. 2018), and 78% in Antarctic Krill (Shao, et al. 2023).

TE mobilisation has consequences in all genomes, disrupting or altering host genes and providing a substrate for exaptation or structural rearrangements (Young, et al. 2020). To mitigate the potential disruptive effects of TEs, organisms have evolved TE suppression mechanisms, including heterochromatin formation of TE-derived DNA preventing transcription and/or post-transcriptional silencing that degrades nascent TE mRNAs (Slotkin and Martienssen 2007). Foreign TEs acquired by HTT initially do not encounter any inherent resistance, resulting in a "transposition burst": a rapid increase in TE copy number in the genome. Such a burst can change genome size, interrupting and shuffling genes and skewing GC content. For example, the horizontally transferred LINE-RTE BovB constitutes 18% of the cow (*Bos taurus*) genome and is also present in other mammals such as the platypus (*Ornithorhynchus anatinus*) and multiple lineages of snake (Walsh, et al. 2013; Puinongpo, et al. 2020). BovB dissemination was likely mediated by parasitic ticks with a broader range of ancestral hosts than currently observed (Ivancevic, et al. 2013). Examples such as this offer valuable insights into the evolutionary relationships and natural history of implicated species.

With approximately 40% of its genome derived from TEs, *Schistosoma mansoni*, a human parasitic flatworm and the best characterised species of the *Schistosomatidae* family, is considered TE-rich (International Helminth Genome Consortium 2019). These parasites have complex life cycles alternating between asexual reproduction in snails and sexual reproduction in vertebrate hosts (such as humans) with two short-lived aquatic stages: miracidia and cercariae (Nelwan 2019). Adult schistosomes can survive in their vertebrate hosts for many years or decades, the oldest recorded survival of individual worms being 37 years (Chabasse, et al.

1985). Parasite persistence in the snail host is variable but lifecycle completion

requires at least 5 weeks in laboratory conditions (Kapp, et al. 2003).

The theoretical basis of *S. mansoni* involvement in HTT seems plausible given

that the parasite has multiple aquatic life stages, extended periods of host association

and TEs constitute at least one third of its genome (International Helminth Genome

Consortium 2019), characteristics known to promote HTT (Gilbert, et al. 2010; Abe, et

al. 2020; Pérez-Etayo, et al. 2020). In fact, an HTT event from salmonid to schistosome

was proposed based on analysis of an EST library and subsequent amplification of

salmonid-specific transposon-like sequences in *Schistosoma* DNA / cDNA (Melamed,

et al. 2004). This work was later disproved on the basis that the level of salmon DNA

contamination in the EST library by salmon was much higher than expected,

potentially due to salmon sperm DNA being used as a DNA carrier during DNA

isolation (Grunau and Boissier 2010; Wijayawardena, et al. 2015). Other claims of HTT

in *Schistosoma* include transfers of MHC sequences from mice or rabbits (Imase, et

al. 2001; Zhao, et al. 2009), DNA transposons from fish and frogs (Leaver 2001),

receptor sequences from mammals (Hu, et al. 2003) and albumin from mice (Williams,

et al. 2006; DeMarco, et al. 2007) have equally been contested based on DNA

contamination, inadequate sampling of taxa, sequence conservation or incomplete /

fragmented genome assemblies (Wijayawardena, et al. 2015). More recently, the

transfer of a functional *cki* homolog in parasitic flatworms (including *S. mansoni*) from

an unidentified metazoan has been proposed (Wendt and Collins III 2024), perhaps

representing one of the few (as yet) undisputed cases of HGT.

Recently, a chromosomal-level genome assembly for *S. mansoni*

(Buddenborg, et al. 2021) and high quality genomes of Schistosoma snail hosts (Nong,

et al. 2022; Young, et al. 2022; Bu, et al. 2023) have been made available, opening

the opportunity for a comprehensive investigation into horizontal transfer in a medically

relevant organism with a high potential for gene transfer. Here, we leverage these

genomic resources to identify and characterise instances of HTT between *S. mansoni*

and its hosts and wider metazoans implicated in this phenomenon.

Methods

For a preliminary assessment of horizontal transfer in selected genomes (Table 1), we

used a library of available and curated *S. mansoni* TEs ([Supplementary Table 1](#),

[Supplementary File 1](#)) as a query in a BLASTN search (v2.12.0) (Camacho, et al.

2009) with predefined "relaxed" parameters that improve the identification of

contiguous TE sequences in species where they may have diverged and/or be highly

fragmented in relation to the query sequences used (Galbraith, et al. 2020). These

parameters are `-reward 3 -penalty -4 -xdrop_ungap 80 -xdrop_gap

130 -xdrop_gap_final 150 -word_size 14 -dust yes -gapopen 30 -

gapextend 6 -outfmt 6 -evalue 1e-5'` and define what we, from now on

call, a "relaxed BLASTN". In addition to published and available TEs (Drew and

Brindley 1997; Drew, et al. 1999; Laha, et al. 2002; Arkhipova, et al. 2003; Copeland,

et al. 2003; DeMarco, et al. 2004; Feschotte 2004; Laha, et al. 2004; Copeland, et al.

2005; DeMarco, et al. 2005; Laha, et al. 2005; DeMarco, et al. 2006; Jacinto, et al.

2011), we manually curated Perere-3 and Sr3 consensuses ([Supplementary File 2](#))

using the previously published sequences (DeMarco, et al. 2005; Laha, et al. 2005;

Venancio, et al. 2010) and published methodology (Goubert, et al. 2022) with the

S. mansoni genome (Buddenborg et al*.* 2021, PRJEA36577, WormBase ParaSite

release WBPS19).

Table 1 - Genome accessions used in preliminary identification of horizontally transferred elements. Organisms were selected for their relationship, either evolutionarily or parasitically to *Schistosoma mansoni*, as described in "Notes".

Organism	Assembly	Notes
Schistosoma mansoni	GCA_000237925.5	Parasite
Trichobilharzia regenti	GCA_944472135.2	Other *Schistosomatidae*
Heterobilharzia americana	GCA_944470555.2	
Fasciola hepatica	GCA_948099385.1	
Clonorchis sinensis	GCA_003604175.2	Other trematodes
Bulinus truncatus	GCA_021962125.1	Intermediate host
Biomphalaria straminea	GCA_021533235.1	
Lymnaea stagnalis	GCA_900036025.1	
Homo sapiens	GCF_000001405.40	Definitive

We used the compiled TE library with manually curated Perere-3 and Sr3 to annotate the *S. mansoni* genome (Howe, et al. 2017; Buddenborg, et al. 2021) available in WormBase ParaSite release WBPS19 (PRJEA36577), using RepeatMasker v4.1.236 (Smit, et al. 2013) with default parameters except for `-no_is -gff -s -a`. To investigate conserved protein domains in consensus sequences of Perere-3 and Sr3, we retrieved open reading frames with EMBOSS' `getorf -minsize 500` (Rice, et al. 2000) from their DNA sequences and submitted them to InterProScan5 (Jones, et al. 2014). Percentage identities of consensus and relevant domains (L1-EN - cd09076 and RT-Pol - PS50878) were generated using Jalview's Pairwise Alignment (Waterhouse, et al. 2009). When evaluating Perere-3/Sr3 genomic insertions, these were considered "Complete" when conserved protein domains "Exo_endo_phos" and "RVT_1" were both present in at least 99% of their full length, as identified with Pfam (Mistry, et al. 2021).

To study the relationship between Perere-3 and Sr3 insertions within the *S. mansoni* genome, we retrieved copies with a minimum length of 3,200bp and used them to construct a distance tree with iqtree (v2.2.3) (Minh, et al. 2020) with default parameters except for `--seqtype DNA -T AUTO --alrt 1000`. The resulting

tree was visualised in ITOL (v5) (Letunic and Bork 2021) and Perere-3 and Sr3

coloured to highlight segregation. We calculated relative TE quantities directly from

our output of RepeatMasker (see above) and represented these using the

`treemapify` (v2.5.5) and `ggplot2` (v3.4.3) (Wickham 2011) packages,

implemented in R Studio (Build 554) (RStudio Team 2020). To investigate substitution

rates of these elements with respect to their consensus sequences, we collected

Kimura values (with divCpGMod) from RepeatMasker's alignment file generated using

the `-a` option and represented them in a histogram.

To further assess TE activity, we used the *terminal branch length* (TBL) method,

previously used for analysis of zebrafish TEs (Chang, et al. 2022). Fragments greater

than 500bp were aligned with `mafft --auto` (Katoh, et al. 2005), trimmed with

TrimAl (Capella-Gutiérrez, et al. 2009) with the `-gt` parameter set to 0.01. Trees

were then constructed per TE using `FastTree -nt -gamma` (Price, et al. 2010)

and TBLs of the element fragment tree were then extracted using

`termlength.py`.

Table 2 - Genome accessions used in construction of *Schistosoma* phylogeny. All genomes used
are available at WormBase Parasite release 19. *Trichobilharzia regenti* was used as an outgroup for
analysis.

Organism	Assembly	Accession
Plakobranchus ocellatus	PoB_v1	GCA_019648995.1
Schistosoma mansoni	v10	GCA_000237925.5
Schistosoma japonicum	ASM636876v1	GCA_025215515.1
Schistosoma turkestanicum	tdSchTur1.1	GCA_944470395.2
Schistosoma rodhaini	tdSchRodh2.1	GCA_944470435.2
Schistosoma spindale	tdSchSpin1.1	GCA_946903255.1
Schistosoma margrebowiei	tdSchMarg1.1	GCA_944470205.2
Schistosoma mattheei	tdSchMatt1.1	GCA_944470405.2
Schistosoma intercalatum	tdSchInte2.1	GCA_944470385.2
Schistosoma haematobium	tdSchHaem2.1	GCA_944470465.2
Schistosoma guineensis	tdSchGuin1.1	GCA_944470375.2
Schistosoma curassoni	tdSchCurr1.1	GCA_944474815.3
Schistosoma bovis	tdSchBovi2.1	GCA_944470445.2
Trichobilharzia regenti	tdTriRege1.1	GCA_944472135.2

Table 3 - Genome accessions used in construction of molluscan phylogeny. All genomes used are available on NCBI Genome List. *Plakobranchus ocellatus* was used as an outgroup for analysis.

Organism	Assembly	Accession
Plakobranchus ocellatus	PoB_v1	GCA_019648995.1
Achatina immaculata	ASM976088v1	GCA_009760885.1
Arion vulgaris	ASM2079622v1	GCA_020796225.1
Cepaea nemoralis	Cnem_1.0	GCA_014155875.1
Candidula unifasciata	ASM976088v1	GCA_905116865.2
Physella (Physa) acuta	ASM2847654v2	GCA_028476545.2
Lymnaea stagnalis	v1.0	GCA_900036025.1
Radix auricularia	ASM207201v1	GCA_002072015.1
Ampullaceana balthica	Rbalt_genome_v1	GCA_944989445.1
Bulinus truncatus	Btru.v1	GCA_021962125.1
Anisus vortex	AniVort1.1	GCA_949126835.1
Biomphalaria straminea	Bstr_hk_v1	GCA_021533235.1
Biomphalaria glabrata	BioGlab47.1	GCF_947242115.1
Biomphalaria pfeifferi	UNM_Bpfe_1.0	GCA_030265305.1

To provide a framework for our proposed TE transfer among *Schistosoma* and selected species, we constructed a species tree using sequences obtained with reciprologs (https://github.com/glarue/reciprologs, with default parameters), an approach that identifies "reciprocal best hits" *i.e.* amino acid sequences that are pairwise best hits across all species sampled. These reciprologs genes were then aligned per-gene using 'mafft --auto' (Katoh, et al. 2005) and subsequently concatenated to generate a single alignment per species of all reciprocal best hits. Species level phylogeny was then constructed with iqtree (Minh, et al. 2020) with parameters '--seqtype AA -T AUTO --alrt 1000 -B 1000'. For the input into reciprologs, we obtained trematode amino acid sequences from WormBase ParaSite (release WBPS19) and filtered them to include just the longest isoform of every gene. For the snail genomes, for which amino acid sequences were not readily available, BUSCO (Simão, et al. 2015) was used (with default parameters and the 'mollusca_obd10' database) to identify putative conserved amino acid sequences derived from the mollusc genomes. The resulting amino acid sequences were then used as inputs for reciprologs. Further to our identification of Perere-3/Sr3 sequences

in non-*S. mansoni* genomes, we proceeded to manually curate (Goubert, et al. 2022)

their consensus sequences in these genomes (Table 2 and Table 3). We introduced

a variation on the first step of the manual curation (see protocol by Goubert et al.

2022), replacing the stringent BLASTN search with our "relaxed BLASTN" approach

to account for greater sequence differences between query (*S. mansoni* Perere-3 and

Sr3) and subject.

To investigate the relationship between consensus sequences of Perere-3 and

Sr3-like elements from Schistosoma and snail genomes, we calculated their pairwise

nucleotide distances (including that of Sr2 as outgroup) using EMBOSS' `distmat`

and the Jukes-Cantor substitution model (Rice, et al. 2000). These relationships are

represented using the `pheatmap` package (Kolde and Kolde 2015) to generate a

hierarchical clustering tree.

To assess any potential transfer of Perere-3 and Sr3 across metazoa, we

constructed a dendrogram of combined novel Perere-3 / Sr3-like sequences from

Schistosoma and molluscan organisms (Table 2 and 3) and LINE_RTE sequences

from RepBase (Bao, et al. 2015). Sequences were aligned with `mafft -auto` and

trimmed with TrimAl (Capella-Gutiérrez, et al. 2009) with `-gt` set to 0.25. This

dendrogram was generated with iqtree using default parameters except for `-bb

1000 -alrt 1000`. To identify the extent of transfer of Perere-3 and Sr3 across

metazoa, we scanned 3446 high-quality genome assemblies (Scaffold N50 > 1Mbp,

accessed from GenBank using the esearch function, August 2023) (Benson, et al.

2018) using our "relaxed BLASTN" approach. The species list was filtered for entries

with available speciation data (n=2271) at timetree.org (Kumar, et al. 2022). We

manually annotated the tree to indicate "taxa by phyla" for all non-chordates and "taxa

by class" for chordates. Some clades representing a single or small numbers of

species were not annotated to improve clarity of figure.

To explore the species implicated in the metazoan transfer, we highlighted

species with either the highest "Max Length Hit", "Percentage Identity" or "Number of

hits over 1,000bp". We also included hits from notable taxa - the only bird, mammal

and a coral (highest of the seven corals identified) and *Schistosoma japonicum* (the

only *Schistosoma* organism with speciation data). To explore horizontal transfer

across these metazoan species we first manually curated Perere-3 / Sr3-like TEs for

all organisms in Table 5 and generated both a species phylogeny and TE consensus

tree for comparison. Species phylogeny for organisms in Table 5 was generated with

timetree.org (Kumar, et al. 2022) and TE consensus tree constructed with iqtree and

`-bb 1000 -wbt -alrt 1000` parameters after alignment with `mafft --auto`.

All processing, analysis and visualisation scripts used in this publication are

available on github at: https://github.com/tbrann99/Schisto_HTT/

Results

Finding *Schistosoma* TE-like sequences in the parasite's

intermediate hosts

We explored the possibility of HTT in *Schistosoma mansoni* using a relaxed BLASTN

search of known *S. mansoni* TEs against other Schistosomatidae (*Trichobilharzia*

regenti, *Heterobilharzia americana*) and trematodes (*Fasciola hepatica, Clonorchis*

sinensis) as well as intermediate (*Bulinus truncatus, Biomphalaria straminea,*

Lymnaea stagnalis) or definitive hosts (*Homo sapiens*). Using this approach, we found

significant similarity to two LINE-RTEs known as Perere-3 and Sr3 in the three genomes corresponding to Schistosoma intermediate hosts, *i.e.* molluscs (Figure 1). Prospective TEs are present at near-full-length of the *S. mansoni* consensus sequence used as query in the search (91.3% and 90.8% in *Bu. truncatus,* 98.3% and 97.9% in *Bi. straminea*, 91.3% and 90.8% in *L. stagnalis* for Perere-3 and Sr3 respectively), suggesting similarity is not simply due to the presence of highly conserved TE domains but instead represents a full-length match. Additionally, in these organisms, many near-full-length (more than 90% full length, approximately 2,900bp for Perere-3 and Sr3) copies were identified, indicating transposition of these elements. In contrast, few "Other TEs" were identified above the e-value filter in intermediate snail hosts. We did not detect significant hits of Perere-3/Sr3 in the other trematodes closely related to *Schistosoma*, namely *C. sinensis* or *F. hepatica*. As these trematodes are evolutionarily closer to *S. mansoni* than the molluscan intermediate host, the presence of Perere-3 and Sr3 is unlikely to be a product of vertical inheritance. This first observation led us to hypothesise that a HTT event between molluscs and their parasites could have occurred, either directly or indirectly.

Organism	Perere-3 Max % FL	Perere-3 n(>90% FL)	Sr3 Max % FL	Sr3 n(>90% FL)	Other TE Name	Other TE Max % FL	Other TE n(>90% FL)
Schistosoma mansoni	100.0	1081	100.0	1084	RTESj-like	100.0	617
Trichobilharzia regenti	21.5	0	42.4	0	Cercyon	100.0	226
Heterobilharzia americana	32.4	0	29.6	0	Cercyon	100.0	6
Fasciola hepatica	0.0	0	0.0	0	Colo	88.9	0
Clonorchis sinensis	0.0	0	2.0	0	Colo	86.8	0
Bulinus truncatus	91.3	143	90.8	171	n/a	0.0	0
Biomphalaria straminaea	98.3	72	97.9	75	Saci2	8.7	0
Lymnaea stagnalis	91.3	45	90.7	65	Columbina	6.3	0
Homo sapiens	0.0	0	0.0	0	n/a	0.0	0

Figure 1 - The genome of the intermediate host of *Schistosoma mansoni* contains *S. mansoni-like* TEs. Select organisms with varying relationships to *S. mansoni* were scanned with a relaxed BLASTN to identify the presence of *S. mansoni* TEs in these genomes. "Max % FL" represents the highest percentage of the *S. mansoni* TE consensus sequence that was mapped to the genome. "n(90% FL) was calculated as the number of copies identified over 90% of the full length of the respective elements. The highest non-Perere-3 or non-Sr3 element ("Other TE") is also shown to highlight potential differences between Perere-3 / Sr3-derived results and other elements. Simplified phylogeny (left) was extracted from timetree.org (Kumar, et al. 2022), representative genomes used for each group highlighted with *.

Perere-3 and Sr3 in the *Schistosoma mansoni* genome

We characterised Perere-3 and Sr3 in *S. mansoni*. In the first instance, we took the sequences of Perere-3 and Sr3 previously described by Venancio et al. (2010) and used the most recent version of the *S. mansoni* assembly to update their consensus sequences using a manual curation approach that better represent the range of TE insertions found in the genome. The resulting consensus sequences for Perere-3 and Sr3 are 3,207 and 3,209 base pairs respectively ([Supplementary File 2](#)). These elements have high sequence similarity: at the DNA level, they share a nucleotide percentage identity of 79.81%. Both elements encode full-length L1-endonuclease (L1-EN) and the reverse transcriptase (RT Pol), for which their amino acid percentage identities are 83.81% and 82.59% related to one another respectively (Figure 2A).

Despite their close sequence similarity, these TEs are considered to transpose independently from one another. To confirm this, we retrieved from the genome all Perere-3 and Sr3 copies with a minimum length of 3,200bp and built a sequence similarity dendrogram (Figure 2B). Our results show that sequences arising from Perere-3 or Sr3 TE transposition events segregate into two distinct clusters. Furthermore, the relative distance between these two LINE-RTEs and others from the same organism suggests that Perere-3 and Sr3 may have relatively recently shared a common TE ancestor (Figure 2B). Differential transposition activity has resulted in these elements having different proportions of genomic coverage: Perere-3 and Sr3 occupy 2.1% and 4.0% of the genome respectively (Figure 2C). Further evidence for their independent activity can be seen in their profile of accumulated mutations represented in the differential kimura distributions (Figure 2D). Kimura distances represent the evolutionary distance of a given insertion from the consensus sequence from which relative age may be inferred. Our data show that Sr3 has a distinct bimodal

distribution of kimura values, with a peak at both 12 and 22, in comparison to Perere-3 which is unimodal with a single peak corresponding to a kimura value of 12. This indicates differential bursts of transposition between these elements and pointing to independent transposition.

To further understand the dynamics of Perere-3 and Sr3 in *S. mansoni*, particularly given their potential involvement in horizontal transfer events, we aimed to assess ongoing transposition activity of these elements. Based on their completeness, these elements are capable of transposing: of the 43,800 Perere-3/Sr3 annotated fragments, 432 (approximately 1%) have at least 99% of the length of the protein domains required for TE transposition, namely the endonuclease and reverse transcriptase domains. However, final proof of ongoing or very recent transposition would be given by presence of identical TE insertions. To this end, we use a TBL approach which provides an estimation of the relative age of all genomic TE copies (Chang, et al. 2022), with low TBLs indicating more recent transpositions. We demonstrate that Sr3 and Perere-3 have 173 and 124 copies with a TBL of 0 (Figure 2E), indicating many identical pair-wise matches, and providing evidence of recent enough transposition events that did not have time to accumulate mutations.

Figure 2 – Perere-3 and Sr3 are similar LINE-RTEs with different transposition histories within the *S. mansoni* genome. A) Protein domain topology of Perere-3 and Sr3, with percentage identity (DNA) of the entire element (in black) and the L1-EN and RT Pol protein domains (in red). **B)** Tree of full-length (>2,900bp) TE insertions in the *S. mansoni* genome with Sr2 LINE-RTE as outgroup. **C)** Genomic coverage of TEs of the *S. mansoni* genome. Total coverage with TE library of 24.7%. **D)** Kimura profile of Perere-3 and Sr3 in the *S. mansoni* genome, divided into "Incomplete" and "Complete" elements, where complete indicates at least 99% of both the L1-EN and RT Pol domains. **E)** *Terminal branch length* (TBL) analysis of all *S. mansoni* TE copies, assessing copy number against median terminal branch length. The size of the datapoint indicates the number of instances where the TBL was 0. Red dashed line represents the mean terminal branch length of all elements.

Evolutionary Spread and Activity of Perere-3 and Sr3 Across Schistosoma and Molluscan Hosts

In the first section, we shown that Perere-3 and Sr3 are found extensively across the *Schistosoma* genus but not in full length nor in high numbers in *Trichobilharzia*, a member of the *Schistosomatidae* family, or wider Trematoda (Figure 1). We broadened our search beyond the *S. mansoni* genome to evaluate the presence and extent of transposition across the *Schistosoma* genus and to more accurately

determine the evolutionary timing of a potential transfer event. We searched for Perere-3/Sr3 sequences in 11 additional *Schistosoma* genomes and found that all contained full copies of the TEs (Table 4).

Table 4 - Full length copies of Perere-3 and Sr3 are found across *Schistosoma spp.* Perere-3 and Sr3 copies identified using the 'relaxed BLASTN' and manually curated *S. mansoni* Perere-3 and Sr3 consensus sequences. All genomes used are available on WormBase ParaSite, with accessions listed in Table 2.

Species	Perere-3			Sr3		
	% Genome	n(>2,900bp)	Max Full Length	% Genome	n(>2,900bp)	Max Full Length
S. japonicum	3.81	977	3239	3.88	982	3242
S. turkestanicum	5.98	1156	3225	5.36	1135	3241
S. rodhaini	5.31	1563	3241	5.81	1574	3241
S. mansoni	4.91	1073	3241	5.43	1079	3244
S. spindale	5.5	1482	3237	6.04	1486	3252
S. margrebowiei	4.8	891	3257	5.35	903	3257
S. mattheei	5.1	1277	3247	5.65	1284	3287
S. intercalatum	5.13	1253	3256	5.69	1257	3256
S. haematobium	4.81	1511	3253	5.32	1514	3269
S. guineensis	4.94	1130	3253	5.49	1134	3268
S. curassoni	4.96	1168	3259	5.51	1177	3266
S. bovis	4.93	1115	3249	5.46	1122	3249

Our analysis of TBL across the *Schistosoma* genus (Figure 3A) suggests that these elements have been active throughout speciation. Whilst similar numbers of near full length (>2,900bp) elements are observed across *Schistosoma*, Perere-3/Sr3 from *S. japonicum* and *S. turkestanicum* have higher TBLs medians and fewer near 0 TBLs, indicating that a higher proportion of transposition events in these organisms are older than in those species most recently diverged (Figure 3A). We found a significant increase in terminal branch length between *S. japonicum* and *S. turkestanicum* and the newer diverged schistosomes (Kruskal-Wallis and Dunn's test, *p<0.001*). The widespread presence and activity of these elements across *Schistosoma* indicate that the horizontal transfer event likely occurred in a common ancestor of the genus.

Figure 3 - *S. mansoni* Perere-3 and Sr3-like sequences are found across *Schistosoma sp.* and molluscs, including snails susceptible to infection. A) *Schistosoma* species tree and terminal branch length (TBL) distribution of Perere-3 and Sr3-like insertions aligned to said tree. Species tree was constructed from reciprocal orthologs across all species. *Schistosoma* phylogeny was constructed with all species available in WormBase ParaSite. *S. mattheei*, *S. intercalatum*, *S. haematobium*, *S. guineensis*, *S. curassoni* and *S. bovis* were collapsed into a single clade representing the *haematobium* group (*), in line with previous phylogenetic assessment of *Schistosoma (Leger and Webster 2017)*. *Trichobilharzia regenti* was used to root the tree, though no full-length Perere-3 / Sr3-like sequences were identified in this organism (Figure 1). TBL values plotted correspond to the *S. haematobium* genome. Diamonds indicate the mean. Testing for terminal branch length differences

between species with a Kruskal-Wallis test and Dunn's post-hoc analysis demonstrated significance ($p<0.001$) across all pairwise-species comparisons except *S. spindale* and *S. rodhaini* ($p>0.05$). Number of elements greater than 2,900bp is shown to the right of the violin plot. **B)** Rooted species tree of Panpulmonata, including Planorbidae snails, the intermediate hosts of *Schistosoma spp.* and related Hygrophila snails. *Plakobranchus ocellatus* was used as an outgroup. Molluscan genes were obtained with BUSCO (see Methods). Leaves are colour-coded to indicate presence of elements over 2,900bp (green), over 1,000bp (orange) or shorter (red). **C)** Heatmap representing distance matrix (DNA) of manually curated TE consensus sequences from *Schistosoma* and Panpulmonate snails alongside hierarchical clustering tree. Snails are labelled Planorbidae (green), Hygrophila (yellow) or panpulmonate (pink). *S. mansoni* LINE-RTE Sr2 was added as an outgroup, as indicated by the '*' in the tree.

Having found extended presence of Perere-3 and Sr3 in a wider range of *Schistosoma spp*, we wanted to test whether these TEs are also present in the respective intermediate hosts (class: Planorbidae) and other molluscs across Panpulmonata. We use the 'relaxed BLASTN' approach with *S. mansoni* Perere-3 and Sr3 as queries on the genomes of selected Planorbidae and Panpulmonata and build a corresponding species tree to represent species relatedness (Figure 3B). We found significant BLASTN hits corresponding to more than 90% of the full length of *S. mansoni* Perere-3 and Sr3 consensus sequences (green in Figure 3B, Supplementary Table 2) in 8 molluscan species. For instance, *Am. balthica* contains 324 copies over 2,900bp in length. Two of the three species of *Biomphalaria* (namely *B. glabrata* and *B. pfeifferi*) have a marked reduction in number of copies as well as their maximum length, suggesting TEs in these species are no longer active and the remaining identifiable copies have accumulated many mutations.

To explore the relationship between *Schistosoma* and molluscan Perere-3 and Sr3-like sequences, we manually curated these TEs from the 12 molluscan genomes

for which significant BLASTN hits were found. To find out how closely related these

sequences are to each other, we built a nucleotide distance matrix (Figure 3C) of the

Schistosoma and molluscan TEs consensus sequences. A hierarchical clustering

approach was used to identify intersequence relationships and we found that

Schistosoma Perere-3 and Sr3-like sequences cluster in the centre of the heatmap

within the curated molluscan TEs (Grey, Figure 3C). The high similarity observed

between *Schistosoma* and molluscan Perere-3 and Sr3-like elements and positioning

of the *Schistosoma* clade within a continuum of molluscan TE sequences, further adds

to the evidence of a horizontal transfer event. Additionally, we found Perere-3 / Sr3 in

Hygrophila snails, which are not Planorbidae and are not known to be susceptible to

Schistosoma infection.

Assessment of Perere-3 and Sr3 across metazoa

To assess the relationship between *Schistosoma*/molluscan Perere-3 and Sr3-like

sequences and the wider LINE-RTE superfamily of TEs, we extracted consensus

sequences of LINE-RTEs from the RepBase database (Bao, et al. 2015) and used

these to construct a sequence similarity dendrogram. Our analysis included two well-

known horizontal TE transfers of LINE-RTEs namely BovB (Ivancevic, et al. 2013) and

AviRTE (Suh, et al. 2016). We show that Perere-3 and Sr3 sequences from

Schistosoma and gastropoda cluster tightly together and, at the same time, separately

from the known TE transfers, indicating a distinct transfer event from those previously

described (Figure 4A). Several LINE-RTEs from RepBase cluster around our curated

Sr3 and Perere-3 sequences, such as those from *Crassostrea gigas* (Pacific Oyster),

Aplysia californica (Californian Sea Hare, a sea slug), *Saccoglossus kowalevskii*

(Acorn Worm, a free-living, hemichordate), *Danio rerio* (Zebrafish) and *Chrysemys picta bellii* (Western Painted Turtle).

We therefore hypothesised that Perere-3 and Sr3 may exist outside *Schistosoma* and molluscs and, as RepBase is limited to user-submitted, previously curated TEs, we extended our search to all available, high quality metazoan genomes. This was done with the relaxed BLASTN approach using *S. mansoni* Perere-3 and Sr3 as queries against all metazoan genomes (contig N50 > 1Mbp) available in GenBank (n=3,446, Supplementary Table 3). An organism was determined to be a "weak" candidate for HTT if it returned a blast hit >1,500bp, while "strong" candidate required at least one larger than 2,000bp and at least five hits >1,000bp. Stringency around the number of hits identified helps to differentiate between contamination in the genome assembly resulting in a single hit and *bona fide* hits, which would have many copies. Using this criteria we found, out of the 3,446 genomes tested, 117 organisms to be "weak" candidates for Perere-3/Sr3 HTT while 256 of these were "strong" candidates. To visualise the presence/absence of Perere-3/Sr3-like sequences in these species and their respective relationships, we used timetree.org and filtered for genomes for which speciation data was available in this database (n=2,271). From this subset of genomes, 207 and 91 were "weak" and 'strong' hits respectively.

We found Perere-3/Sr3-like sequences discontinuously across the metazoan tree (Figure 4B), including reptiles, such as the previously mentioned *Chrysemys picta bellii*, amphibians, marine invertebrates, molluscs, cnidaria, a single bird and a single mammal and more extensively across fish. Such a diverse number of taxa from which these hits derived suggests that Perere-3 and Sr3 have been extensively transferred across metazoans. Given the similarity of Perere-3 and Sr3 (Figure 2A) and the expected nucleotide distances of such TEs, highly similar results were seen for

Perere-3 and Sr3 when searched against metazoan genomes (Supplementary Figure

2).

Of the 298 organisms identified as containing Perere-3 / Sr3-like sequences,

we highlight several species (Table 5) which were selected either due to having large

numbers of hits over 1,000bp, highest single largest hits or being notable taxonomic

clades. To investigate TEs and species relatedness, we curated Perere-3 and Sr3-like

sequences in these species. We first evaluated their sequence similarity and presence

of conserved protein domains and found that, despite a large evolutionary distance

across taxa (Figure 5A), all but two species of curated TEs encode full length L1-end

and RT domains and shared the overall TE domain topology (Figure 5B,

Supplementary Figure 3). We then compared a dendrogram of curated consensus

sequences to the corresponding species tree. We show extensive phylogenetic

incongruence between the species and TE trees (Figure 5). For example, despite fish

and *S. mansoni* diverging more than 600 million years ago (Figure 5A), curated

Perere-3 and Sr3-like sequences from fish are closely related to their *S. mansoni*

counterparts, contradicting what would be expected by vertical inheritance. Such

phylogenetic incongruence between species for sequences sharing such levels of

similarity suggests extensive horizontal transfer of Perere-3 and Sr3, beyond

Schistosoma and mollusca and across metazoa.

Figure 4 - Perere-3 and Sr3 are found across metazoa. A) Dendrogram of consensus sequences of
LINE-RTEs found in RepBase plus *Schistosoma* and mollusc Perere-3 / Sr3-like sequences,
highlighting known horizontally transferred groups BovB (Ivancevic, et al. 2013) and AviRTE (Suh, et
al. 2016). * Indicates additional BovB sequence that did not cluster within other BovB sequences. **B)**
Presence of Perere-3 / Sr3-like sequences across metazoan genomes available at Timetree.org
(Kumar, et al. 2022). Relevant hits were classified as either weak (light brown: at least one hit > 1,500bp)
or strong (dark brown: at least one hit > 2,000bp and at least five hits > 1,000bp). Perere-3 and Sr3
were used as a query, results shown for the TE generating higher results, TEs were plotted separately
in Supplementary Figure 2.

Table 5. Select metazoan species identified from Figure 4. Species were chosen based on the highest maximum length, highest percentage identities and highest n(>1,000bp) found. In addition to these, notable taxa identified were also included (coral, mammal and bird).

Taxa		Organism	Alternative Name	Max Length	% ID	n(>1,000bp)
Schistosoma

Schistosoma japonicum	-	3239	74.1	5349
Tortoise / Turtle		*Chelonoidis abingdonii*	Pinta Island Tortoise	3199	66.2	78
		Trachemys dorbigni	D'Orbignys slider	3195	65.6	104
		Cuora mccordi	McCord's box turtle	3189	65.9	692
Fish		*Scleropages formosus*	Asian arowana	3192	65.3	62
		Sinocyclocheilus grahami	Golden-line barbel	3036	72.1	4
		Culter alburnus	Topmouth culter	3042	72.0	7
		Megalobrama amblycephala	Wuchang bream	3037	71.7	2
Mollusc		*Cerithideopsis pliculosa*	Plicate horn shell	3039	68.1	2019
		Conus ventricosus	Mediterranean cone	3172	66.6	800
Coral		*Desmophyllum pertusum*	Lophelia pertusa	3028	68.9	55
Mammal		*Gracilinanus agilis*	Agile gracile opossum	1619	60.4	28
Bird		*Pseudopodoces humilis*	Grount tit	2582	64.6	1

Figure 5. **Juxtaposition of TE and species trees highlights the phylogenetic incongruence between them and is evidence in support of the horizontal transfer of Sr3 and Perere-3 A)** Speciation tree generated with timetree.org showing divergence times of selected organisms. **B)** Tree of manually curated TE consensus sequences of Perere-3 and Sr3 from the same organisms in A. Two distinct elements were curated for *Conus ventricosus* and *Cerithideopsis pliculosa*. *Culter alburnus* and *G. agilis* curated elements were 5' truncated. Ultrafast bootstrap values lower than 98 are shown. TE domains of curated elements are represented with boxes to the right of the panel; L1-EN (red), RT_Pol (green) and DUF-6451. Protein domains of all consensus sequences were mapped using the methodology for Figure 2A.

Discussion

Discussion

Our preliminary investigation using BLASTN on the genomes of trematodes and

Schistosoma hosts (namely human and Planorbidae snails) with *S. mansoni* TEs,

revealed that two elements, Perere-3 and Sr3, are well represented in the genomes

of *Schistosoma spp.* intermediate hosts (*Lymnaea stagnalis*, *Bu. truncatus*,

Bi. straminea), while no other element shares the same degree of similarity among

this group. Perere-3 and Sr3 were absent from the definitive human host and from the

non-*Schistosoma* trematode genomes sampled, suggesting either loss in the

trematode lineages sampled or a later gain in *Schistosoma*. Owing to the parasitic

relationship and prolonged contact between *Schistosoma* and their intermediate host,

we hypothesised that Perere-3 and Sr3 could represent examples of horizontally

transferred TEs. Previously documented horizontal transfer events involving

S. mansoni, e.g. with Salmonids (Melamed, et al. 2004), Humans (Yu, et al. 2008) and

Sturgeon (Hale, et al. 2010) have all subsequently been refuted (Wijayawardena, et

al. 2015). As these rare transfer events can provide insights into the evolutionary

history of organisms and their relatedness (Ivancevic, et al. 2013; Daubin and Szöllősi

2016) we aimed to further characterise and confirm our initial observation of HTT of

Perere-3 / Sr3 and examine evidence for the direction of transfer.

Our first focus was on assessing Perere-3 and Sr3 within the *S mansoni*

genome. Perere-3 and Sr3 are two LINE-RTEs with high sequence similarity that exist

at high frequency in the *S. mansoni* genome. With a percentage identity of 79.81%

across their full-length sequence, these elements fall just outside the "80:80:80 rule"

(Wicker, et al. 2007) which justifies their classification as separate TE families (Figure

2A). We demonstrate that the diversity of near-full length insertions of Perere-3 and

Sr3 in the *S. mansoni* genome segregated into two distinct clusters (Figure 2B). We

propose that these elements were at one point a single element that underwent

extensive transposition and divergence.

Within the *S. mansoni* genome, Perere-3 and Sr3 represent the second and

fourth most abundant TEs, respectively (Figure 2C). Together, these elements

comprise 6% of the *S. mansoni* genome and account for approximately one-fifth of the

organism's TE content, indicating extensive past transposition. In addition to their

considerable abundance, we show that these elements are capable of transposition,

as indicated by 432 insertions having full-length (Figure 2D) conserved L1-EN and RT

protein domains, the two enzymes needed for active transposition. In fact, these TEs

are likely actively transposing, as evidenced by the 430 copies with terminal branch

lengths of 0 (Figure 2E), *i.e.* not even one nucleotide difference between the pairs of

insertions considered. While by no means a confirmation of horizontal transfer,

historical transposition bursts and periods of ongoing activity suggest lack or reduced

capability of sequence-specific resistance by the host genome, making these TEs

good candidates of HTT (Gilbert and Feschotte 2018). Perere-3 and Sr3's

transposition activity and high abundance in *S. mansoni* has contributed to its

increased genome size, providing a large source of homologous substrate for

recombination, which may have contributed in turn to the expansion of gene families

in the TE-rich subtelomere of *S. mansoni* (Brann, et al. 2024).

We further investigated the presence or absence of Perere-3 and Sr3 across

Schistosoma and panpulmonata lineages by manually curating consensus sequences

of these elements in selected species. We were able to identify and curate Perere-3

and Sr3-like elements from representative *Schistsosoma* lineages having constructed

a phylogeny for all genome assemblies available at the time of writing. We show that

the most recently diverged Schistosoma species (*S. haematobium, S. spindale,*

S. mansoni and *S. rodhaini*) have a larger proportion of active elements (reflected in

the lower TBL median) than the more basal species *S. japonicum* and

S. turkestanicum (Figure 3A). We cannot directly attribute speciation of *Schistosoma*

to the horizontal acquisition of an element in the last common ancestor but such an

event is likely to have had a large impact on the genus' evolution, interrupting genes

and / or genic regions or promoting structural rearrangements. Venancio et al. (2010)

hypothesised the transposition of Perere-3 and Sr3 contributed to the speciation of

African schistosomes. To confirm this, 'speciation genes' associated with *Schistosoma*

diversification would first need to be identified (Nosil and Schluter 2011) and the

subsequent contribution of Perere-3 and Sr3 to these be investigated.

Our analysis identified significant hits of Perere-3 and Sr3-like sequences in

Planorbidae genomes, a family of snails susceptible to *Schistosoma* infection. The

large evolutionary distance that separates these two animal groups, approximately

600 million years (Kumar, et al. 2022), suggests a horizontal transfer event has

occurred. To identify the direction of transfer, we extended our relaxed BLASTN to

include other, related snails from Hygrophila and the wider Panpulmonata order and

demonstrated the extended presence of these elements (Figure 3B). For instance,

Bu. truncatus, susceptible to *S. japonicum* infection, and *Bi. straminea*, susceptible to

Schistosoma spp. infection, have 142 (~ 1Mbp or 0.25% of the genome) and 72 copies

above 90% of the full length of the consensus sequences of *S. mansoni* Perere-3 and

Sr3. Given the multiple occurrences and completeness of copies found across

molluscan genomes, such an observation is unlikely to result from DNA contamination

during genome assembly, as has been proposed for previous putative *S. mansoni*

HGT observations (Wijayawardena, et al. 2015). Interestingly, *Bi. glabrata* and *B.*

pfeifferi had only degraded copies, despite their close relationship to *Bi. straminea*

(Figure 3B), suggesting that these elements have stopped transposition sometime

post-divergence from *Bi. straminea*.

To find further evidence in support of HTT and discern potential evolutionary

relationships between TE–derived sequences in *Schistosoma* and Papulmonate, we

calculated the pairwise nucleotide distance of manually curated, consensus

sequences of Perere-3 and Sr3-like from species in these groups (Figure 3C). We

show that *Schistosoma* Perere-3 and Sr3-like sequences cluster within the

comparatively more diverse molluscan sequences. This observation suggests a

potential route for the horizontal transfer of Perere-3 and Sr3 from Panpulomnate to

Schistosoma presumably via the parasitic relationship of an ancestral *Schistosoma*

panpulmonate host-parasite pair. The two groups of molluscan sequences formed by

the "insertion" of the *Schistosoma* cluster (Figure 3C) only loosely correspond to either

the constructed phylogenetic tree or panpulmonates' susceptibility to schistosome

infection. This discordance could be explained by as yet undescribed parasitic

relationships between snail and host, or relationships that cannot occur due to a lack

of geographic overlap rather than biological incompatibility. From the sequence and

species relationships investigated here, we propose that Perere-3 and Sr3 LINE-

RTEs, which have extensively transposed in *Schistosoma spp.*, could have been

acquired via horizontal transfer from an ancestral snail host.

To investigate if these TEs could have been further transferred in more distantly

related species, we expanded our search through a collection of metazoan genomes.

In this approach we queried the genomes of 2,271 organisms (for which speciation

data was available at timetree.org) for sequences with high similarity to *S. mansoni*

Perere-3 or Sr3. We identified Perere-3/Sr3-like sequences across 298 organisms,

204 of which were classified as "strong" hits (more than five hits larger than 1,000bp

and one more than 2,000bp), including reptiles, gastropods and fish (Figure 4B). A

weak hit was identified from one mammal (*Gracilanus agilis*) and bird (*Pseudopodoces*

humilis). These weak hits are likely contamination of the genome assembly, as the

multiple large copies were not identified, indicating transposition had not occurred in

these genomes. Other species had strong hits, indicative of not only presence of a

single TE but TE activity. For example, the painted turtle *Chrysemys picta bellii*, which

organisms from the superfamily *Schistosomatoidae* do infect turtles, and trematode

infection has been described (Johnson, et al. 1998). Fish host a remarkable number

of digenean parasites (Bullard and Overstreet 2008) and the horizontal transfer of

genetic material between predator and prey has been documented via parasite

intermediaries (Kambayashi, et al. 2022). Fish, such as *Oreochromis* identified here

have been trialled in the biological control of schistosomiasis via predation of snail

populations (Slootweg, et al. 1994) and will also eat free-living aquatic stages of the

trematode life cycle (Orlofske, et al. 2015). It is possible that the feeding of fish, on

either Perere-3 / Sr3-containing snails and schistosomes may provide an avenue by

which HGT has occurred. Many of the identified snails are hosts for multiple parasites

with complex life cycles, for example, *Lymnaea stagnalis* is "a cosmopolitan vector of

trematodes infecting diverse vertebrates" (Gilbert, et al. 2010). It is possible that these

panpulmonates act as a distribution centre for horizontally transferred material across

different host-parasite combinations and life cycles that facilitate the extensive transfer

of Perere-3 and Sr3 across metazoan organisms in and around aquatic environments.

Water-filtering organisms such as sponges and corals are known for their ability to

accurately reflect the metagenome of a given environment (Cai, et al. 2022). In aquatic

environments where Perere-3 and Sr3 are actively expanding within and transferring

between organisms, it seems likely that this phenomenon would extend to organisms

sampling such an environment (Figure 6.)

Figure 6. Proposed network of organisms implicated in the transfer of Perere-3 and Sr3. Organisms not to scale, arrows not representative of direction of Perere-3 and Sr3 transfer but of network interactions. Image generated using biorender.com.

The phylogenetic distribution of Perere-3/Sr3-containing species is patchy and

cannot be explained by vertical inheritance alone (unless many simultaneous TE

losses occurred in many genomes). This observation suggests that the horizontal

transfer of Perere-3 and Sr3 extends well beyond the host-parasite relationships

between Schistosoma and molluscs. A comparison of the speciation tree and the TE

phylogeny generated for these organisms (Figure 5) demonstrates this phylogenetic

incongruence.

Conclusion

We were able to leverage available genomic resources for *S. mansoni* and the high-quality genomes of other organisms to identify Perere-3 and Sr3-like sequences and characterise the extent of horizontal transfer of these LINE-RTEs in a way not previously possible. We show evidence in support of horizontal transfer of Perere-3 / Sr3 from panpulmonate snails to *Schistosoma* parasites, which, in light of further evidence showing the spread of these sequences further through the phylogenetic tree, appears as the tip of the iceberg of a much broader transfer dynamic. Owing to the rebuking of previous *S. mansoni* transfer events, we believe this represents one of the few robust horizontal transfer examples available in *S. mansoni*, due to the number of organisms implicated, quality of genomes sampled, and breadth of organisms involved.

Acknowledgments:

Dr James Galbraith, University of Exeter, for discussions around early observations.

Dr Alex Suh, University of Bonn, for help and ideas around early drafts of analysis.

Kristian Sorensen, for contributing to literature analysis, manuscript review and discussions on horizontal transfer context.

Caroline Walker, University of Cambridge, for illustrative feedback of proposed network of organisms.

Author Contributions:

Study was conceived by TB. Methods and analysis were implemented by TB and AVP. Manual curation of Perere-3 and Sr3 was completed by AVP and manual curation of elements in *Schistosoma* (excluding *S. mansoni*) and snails was completed by FdSO. The manuscript was drafted by TB and AVP. All authors edited and approved the final version of the manuscript.

Funding:

T.B. funded by the Department of Pathology, University of Cambridge.

F. S. dO financed by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES, Finance Code 001) and the Institutional Internationalization Program (CAPES/PRINT) from Universidade Federal do Paraná (UFPR).

A.V.P. funded by Christ's College Cambridge (Research Fellowship in Animal Parasitology).

Supplementary Information

	Supplementary Table 1. Source of _Schistosoma mansoni_ transposable element consensus sequences used in analysis. Reported refers to the relevant citation in which the TE was identified / described and accessions of respective elements provided where consensus sequences were submitted.
	Supplementary Table 2. Relaxed BLASTN output of Perere-3 and Sr3 on molluscan genomes used in analysis. Results reported for Perere-3 and Sr3 separately. '% Genome' refers to the percentage of the genome occupied by the respective element, 'n(>2,900bp)' the number of hits above 2,900bp and 'Max Full Length' the length of the largest hit identified.
	Supplementary Table 3. Assembly information and relaxed BLASTN output for all genomes selected for analysis. All metazoan genome assemblies with an N50 greater than 1,000,000bp were identified and searched for the presence of Perere-3 and Sr3 using the 'Relaxed BLASTN'. Blast results for all species are shown for Perere-3 and Sr3 respectively, highlighting the percentage identity (%id),length and Evalue for the best hit, as identified by bit score. Additionally, the number of hits over 1,000bp for the given element was also highlighted (n>1,000bp). For visualisation, a species phylogeny was constructed using timetree; species with an appropriate entry are highlighted in "Present in TimeTree" column.
	Supplementary Figure 1 – E-Value distribution of metazoan 'relaxed' BLASTN hits demonstrates hits unlikely to arise by chance alone . Hits were classified as those greater than 1,000bp, of these approximately 80% had an E-Value of 0. These were segmented into those >1,000bp (red), or >2,000bp (blue), as the definition of a "strong" hit required at least one hit greater than 2,000bp. Non-0 E-values were plotted by magnitude (i.e. 100 refers to a value of e-100). Black dashed line indicates mean of non-0 E-values.
	Supplementary Figure 2 - Separate Perere-3 (A) and Sr3 (B) blast results across high-quality metazoan genomes. Presence of Perere-3 / Sr3-like sequences across metazoan genomes available at Timetree.org.

	Relevant hits were classified as either weak (light brown: at least one hit > 1,500bp) or strong (dark brown: at least one hit > 2,000bp and at least five hits > 1,000bp).
Supplementary Figure 3	**Supplementary Figure 3 – Reverse transcriptase and transposable element consensus distance trees provide similar topology.** Trees of manually curated TE consensus sequences of Perere-3 and Sr3 from organisms of Table 4. Two distinct elements were curated for Conus ventricosus and Cerithideopsis pliculosa. Culter alburnus and G. agilis curated elements were 5' truncated. Reverse transcriptase domain tree generated using InterProScan5 to identify relevant protein domains. Both trees generated using `iqtree -bb 1000 -wbt -alrt 1000` after alignment with mafft.
Supplementary File 1	**Supplementary File 1. *Schistosoma mansoni* transposable element library used for analysis.** Compiled from sources listed in Supplementary Table 1.
Supplementary File 2	**Supplementary File 2. Manually curated Perere-3 and Sr3 sequences from *Schistosoma mansoni*.** Sequences curated using methodology outlined by Goubert et al. 2022.

664   All processing, analysis and visualisation scripts used in this publication are available

665   on GitHub at: https://github.com/tbrann99/Schisto_HTT/

References

Abe K, Nomura N, Suzuki S. 2020. Biofilms: hot spots of horizontal gene transfer (HGT) in aquatic environments, with a focus on a new HGT mechanism. FEMS microbiology ecology 96:fiaa031.

Arkhipova IR, Pyatkov KI, Meselson M, Evgen'ev MB. 2003. Retroelements containing introns in diverse invertebrate taxa. Nature genetics 33:123-124.

Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. Mobile DNA 6:1-6.

Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Ostell J, Pruitt KD, Sayers EW. 2018. GenBank. Nucleic acids research 46:D41-D47.

Brann T, Beltramini A, Chaparro C, Berriman M, Doyle S, Protasio A. 2024. Subtelomeric plasticity contributes to gene family expansion in the human parasitic flatworm Schistosoma mansoni. BMC genomics 25:217.

Bu L, Lu L, Laidemitt MR, Zhang S-M, Mutuku M, Mkoji G, Steinauer M, Loker ES. 2023. A genome sequence for Biomphalaria pfeifferi, the major vector snail for the human-infecting parasite Schistosoma mansoni. PLoS neglected tropical diseases 17:e0011208.

Buddenborg SK, Tracey A, Berger DJ, Lu Z, Doyle SR, Fu B, Yang F, Reid AJ, Rodgers FH, Rinaldi G. 2021. Assembled chromosomes of the blood fluke Schistosoma mansoni provide insight into the evolution of its ZW sex-determination system. bioRxiv:2021.2008. 2013.456314.

Bullard SA, Overstreet RM. 2008. Digeneans as enemies of fishes. In. Fish Diseases (2 Vols.): CRC Press. p. 831-911.

Cai W, Harper LR, Neave EF, Shum P, Craggs J, Arias MB, Riesgo A, Mariani S. 2022. Environmental DNA persistence and fish detection in captive sponges. Molecular Ecology Resources 22:2956-2966.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. BMC bioinformatics 10:1-9.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972-1973.

Chabasse D, Bertrand G, Leroux J, Gauthey N, Hocquet P. 1985. Developmental bilharziasis caused by Schistosoma mansoni discovered 37 years after infestation. Bulletin de la Societe de pathologie exotique et de ses filiales 78:643-647.

Chang N-C, Rovira Q, Wells J, Feschotte C, Vaquerizas JM. 2022. Zebrafish transposable elements show extensive diversification in age, genomic distribution, and developmental expression. Genome research 32:1408-1423.

Copeland CS, Brindley PJ, Heyers O, Michael SF, Johnston DA, Williams DL, Ivens AC, Kalinna BH. 2003. Boudicca, a retrovirus-like long terminal repeat retrotransposon from the genome of the human blood fluke Schistosoma mansoni. Journal of Virology 77:6153-6166.

Copeland CS, Mann VH, Morales ME, Kalinna BH, Brindley PJ. 2005. The Sinbad
retrotransposon from the genome of the human blood fluke, Schistosoma mansoni, and the
distribution of related Pao-like elements. BMC Evolutionary Biology 5:1-18.

Daubin V, Szöllősi GJ. 2016. Horizontal gene transfer and the history of life. Cold Spring
Harbor perspectives in biology 8:a018036.

DeMarco R, Kowaltowski AT, Machado AA, Soares MB, Gargioni C, Kawano T, Rodrigues
V, Madeira AM, Wilson RA, Menck CF. 2004. Saci-1,-2, and-3 and Perere, four novel
retrotransposons with high transcriptional activities from the human parasite Schistosoma
mansoni. Journal of Virology 78:2967-2978.

DeMarco R, Machado AA, Bisson-Filho AW, Verjovski-Almeida S. 2005. Identification of
18 new transcribed retrotransposons in Schistosoma mansoni. Biochemical and biophysical
research communications 333:230-240.

DeMarco R, Mathieson W, Dillon GP, Wilson RA. 2007. Schistosome albumin is of host,
not parasite, origin. International journal for parasitology 37:1201-1208.

DeMarco R, Venancio TM, Verjovski-Almeida S. 2006. SmTRC1, a novel Schistosoma
mansoni DNA transposon, discloses new families of animal and fungi transposons belonging to
the CACTA superfamily. BMC Evolutionary Biology 6:1-13.

Drew AC, Brindley PJ. 1997. A retrotransposon of the non-long terminal repeat class from
the human blood fluke Schistosoma mansoni. Similarities to the chicken-repeat-1-like elements of
vertebrates. Molecular biology and evolution 14:602-610.

Drew AC, Minchella DJ, King LT, Rollinson D, Brindley PJ. 1999. SR2 elements, non-long
terminal repeat retrotransposons of the RTE-1 lineage from the human blood fluke Schistosoma
mansoni. Molecular biology and evolution 16:1256-1269.

Feschotte C. 2004. Merlin, a new superfamily of DNA transposons identified in diverse
animal genomes and related to bacterial IS 1016 insertion sequences. Molecular biology and
evolution 21:1769-1780.

Galbraith JD, Ludington AJ, Suh A, Sanders KL, Adelson DL. 2020. New environment, new
invaders—repeated horizontal transfer of LINEs to sea snakes. Genome Biology and Evolution
12:2370-2383.

Gilbert C, Feschotte C. 2018. Horizontal acquisition of transposable elements and viral
sequences: patterns and consequences. Current opinion in genetics & development 49:15-24.

Gilbert C, Schaack S, Pace II JK, Brindley PJ, Feschotte C. 2010. A role for host–parasite
interactions in the horizontal transfer of transposons across phyla. Nature 464:1347-1350.

Goubert C, Craig RJ, Bilat AF, Peona V, Vogan AA, Protasio AV. 2022. A beginner's guide
to manual curation of transposable elements. Mobile DNA 13:7.

Grunau C, Boissier J. 2010. No evidence for lateral gene transfer between salmonids and
schistosomes. Nature genetics 42:918-919.

Hale MC, Jackson JR, DeWoody JA. 2010. Discovery and evaluation of candidate sex-
determining genes and xenobiotics in the gonads of lake sturgeon (Acipenser fulvescens).
Genetica 138:745-756.

Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. 2017. WormBase ParaSite− a
comprehensive resource for helminth genomics. Molecular and Biochemical Parasitology 215:2-
10.

Hu W, Yan Q, Shen D-K, Liu F, Zhu Z-D, Song H-D, Xu X-R, Wang Z-J, Rong Y-P, Zeng
L-C. 2003. Evolutionary and biomedical implications of a Schistosoma japonicum complementary
DNA resource. Nature genetics 35:139-147.

Imase A, Kobayashi K, Ohmae H, Matsuda H, Iwamura Y. 2001. Horizontal and vertical
transmission of mouse class I MHC sequence in Schistosoma mansoni. Parasitology 123:163-168.
International Helminth Genome Consortium. 2019. Comparative genomics of the major parasitic
worms. Nature genetics 51:163-174.

Ivancevic AM, Walsh AM, Kortschak RD, Adelson DL. 2013. Jumping the fine LINE
between species: horizontal transfer of transposable elements in animals catalyses genome
evolution. Bioessays 35:1071-1082.

Jacinto DS, Muniz HDS, Venancio TM, Wilson RA, Verjovski-Almeida S, Demarco R. 2011.
Curupira-1 and Curupira-2, two novel Mutator-like DNA transposons from the genomes of human
parasites Schistosoma mansoni and Schistosoma japonicum. Parasitology 138:1124-1133.

Johnson CA, Griffith JW, Tenorio P, Hytrek S, Lang CM. 1998. Fatal trematodiasis in
research turtles. Comparative Medicine 48:340-343.

Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell
A, Nuka G. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics
30:1236-1240.

Kambayashi C, Kakehashi R, Sato Y, Mizuno H, Tanabe H, Rakotoarison A, Künzel S,
Furuno N, Ohshima K, Kumazawa Y. 2022. Geography-dependent horizontal gene transfer from
vertebrate predators to their prey. Molecular biology and evolution 39:msac052.

Kapp K, Coustau C, Wippersteg V, Jourdane J, Kunz W, Grevelding CG. 2003.
Transplantation of in vitro-generated Schistosoma mansoni mother sporocysts into Biomphalaria
glabrata. Parasitology research 91:482-485.

Katoh K, Kuma K-i, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of
multiple sequence alignment. Nucleic acids research 33:511-518.

Kolde R, Kolde M. 2015. Package 'pheatmap'. r package, Vol. 1. 790. In.

Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G, Hedges
SB. 2022. TimeTree 5: an expanded resource for species divergence times. Molecular biology and
evolution 39:msac174.

Kurland CG, Canback B, Berg OG. 2003. Horizontal gene transfer: a critical view.
Proceedings of the National Academy of Sciences 100:9658-9662.

Laha T, Brindley PJ, Verity CK, McManus DP, Loukas A. 2002. pido, a non-long terminal
repeat retrotransposon of the chicken repeat 1 family from the genome of the Oriental blood fluke,
Schistosoma japonicum. Gene 284:149-159.

Laha T, Kewgrai N, Loukas A, Brindley PJ. 2005. Characterization of SR3 reveals
abundance of non-LTR retrotransposons of the RTE clade in the genome of the human blood fluke,
Schistosoma mansoni. BMC genomics 6:1-14.

Laha T, Loukas A, Smyth DJ, Copeland CS, Brindley PJ. 2004. The fugitive LTR
retrotransposon from the genome of the human blood fluke, Schistosoma mansoni. International
journal for parasitology 34:1365-1375.

Leaver MJ. 2001. A family of Tc1-like transposons from the genomes of fishes and frogs:
evidence for horizontal transmission. Gene 271:203-214.

Leger E, Webster JP. 2017. Hybridizations within the genus Schistosoma: implications for
evolution, epidemiology and control. Parasitology 144:65-80.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. Nucleic acids research 49:W293-W296.

Melamed P, Chong KL, Johansen MV. 2004. Evidence for lateral gene transfer from
salmonids to two Schistosome species. Nature genetics 36:786-787.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,
Lanfear R. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic inference in the
genomic era. Molecular biology and evolution 37:1530-1534.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer EL, Tosatto
SC, Paladin L, Raj S, Richardson LJ. 2021. Pfam: The protein families database in 2021. Nucleic
acids research 49:D412-D419.

Nelwan ML. 2019. Schistosomiasis: life cycle, diagnosis, and control. Current Therapeutic
Research 91:5-9.

Nong W, Yu Y, Aase-Remedios ME, Xie Y, So WL, Li Y, Wong CF, Baril T, Law ST, Lai
SY. 2022. Genome of the ramshorn snail Biomphalaria straminea—an obligate intermediate host
of schistosomiasis. GigaScience 11:giac012.

Nosil P, Schluter D. 2011. The genes underlying the process of speciation. Trends in
ecology & evolution 26:160-167.

Orlofske SA, Jadin RC, Johnson PT. 2015. It'sa predator–eat–parasite world: how
characteristics of predator, parasite and environment affect consumption. Oecologia 178:537-547.

Pérez-Etayo L, González D, Vitas AI. 2020. The aquatic ecosystem, a good environment
for the horizontal transfer of antimicrobial resistance and virulence-associated factors among
extended spectrum β-lactamases producing E. coli. Microorganisms 8:568.

817 Price MN, Dehal PS, Arkin AP. 2010. FastTree 2–approximately maximum-likelihood trees
818 for large alignments. PloS one 5:e9490.

819 Puinongpo W, Singchat W, Petpradub S, Kraichak E, Nunome M, Laopichienpong N,
820 Thongchum R, Intarasorn T, Sillapaprayoon S, Indananda C. 2020. Existence of Bov-B LINE
821 retrotransposons in snake lineages reveals recent multiple horizontal gene transfers with copy
822 number variation. Genes 11:1241.

823 Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European molecular biology open
824 software suite. Trends in Genetics 16:276-277.

825 RStudio Team. 2020. Integrated development for R. RStudio, Pbc.

826 Shao C, Sun S, Liu K, Wang J, Li S, Liu Q, Deagle BE, Seim I, Biscontin A, Wang Q. 2023.
827 The enormous repetitive Antarctic krill genome reveals environmental adaptations and population
828 insights. Cell 186:1279-1294. e1219.

829 Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
830 assessing genome assembly and annotation completeness with single-copy orthologs.
831 Bioinformatics 31:3210-3212.

832 Slootweg R, Malek E, McCullough F. 1994. The biological control of snail intermediate
833 hosts of schistosomiasis by fish. Reviews in Fish Biology and Fisheries 4:67-90.

834 Slotkin RK, Martienssen R. 2007. Transposable elements and the epigenetic regulation of
835 the genome. Nature reviews genetics 8:272-285.

836 Smit A, Hubley R, Green P. 2013. RepeatMasker, http://www.repeatmasker.org/.

837 Suh A, Witt CC, Menger J, Sadanandan KR, Podsiadlowski L, Gerth M, Weigert A, McGuire
838 JA, Mudge J, Edwards SV. 2016. Ancient horizontal transfers of retrotransposons between birds
839 and ancestors of human pathogenic nematodes. Nature communications 7:11396.

840 Tang W, Mun S, Joshi A, Han K, Liang P. 2018. Mobile elements contribute to the
841 uniqueness of human genome with 15,000 human-specific insertions and 14 Mbp sequence
842 increase. DNA Research 25:521-533.

843 Van Etten J, Bhattacharya D. 2020. Horizontal gene transfer in eukaryotes: not if, but how
844 much? Trends in Genetics 36:915-925.

845 Venancio TM, Wilson RA, Verjovski-Almeida S, DeMarco R. 2010. Bursts of transposition
846 from non-long terminal repeat retrotransposon families of the RTE clade in Schistosoma mansoni.
847 International journal for parasitology 40:743-749.

848 Walsh AM, Kortschak RD, Gardner MG, Bertozzi T, Adelson DL. 2013. Widespread
849 horizontal transfer of retrotransposons. Proceedings of the National Academy of Sciences
850 110:1012-1016.

851 Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. 2009. Jalview Version 2—
852 a multiple sequence alignment editor and analysis workbench. Bioinformatics 25:1189-1191.

Wendt G, Collins III JJ. 2024. Horizontal gene transfer of a functional cki homolog in the
human pathogen Schistosoma mansoni. bioRxiv.

Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy P,
Morgante M, Panaud O. 2007. A unified classification system for eukaryotic transposable
elements. Nature reviews genetics 8:973-982.

Wickham H. 2011. ggplot2. Wiley interdisciplinary reviews: computational statistics 3:180-
185.

Wijayawardena BK, Minchella DJ, DeWoody JA. 2015. Horizontal gene transfer in
schistosomes: a critical assessment. Molecular and Biochemical Parasitology 201:57-65.

Williams DL, Sayed AA, Ray D, McArthur AG. 2006. Schistosoma mansoni albumin, a
major defense against oxidative damage, was acquired by lateral gene transfer from a mammalian
host.

Young E, Abid HZ, Kwok P-Y, Riethman H, Xiao M. 2020. Comprehensive analysis of
human subtelomeres by whole genome mapping. PLoS genetics 16:e1008347.

Young ND, Stroehlein AJ, Wang T, Korhonen PK, Mentink-Kane M, Stothard JR, Rollinson
D, Gasser RB. 2022. Nuclear genome of Bulinus truncatus, an intermediate host of the
carcinogenic human blood fluke Schistosoma haematobium. Nature communications 13:977.

Yu F, Li Y, Liu L, Li Y. 2008. Comparative genomics of human-like Schistosoma japonicum
genes indicates a putative mechanism for host–parasite relationship. Genomics 91:152-157.

Zhao G, Li J, Zou F, Liu W, Mo X, Lin R, Yuan Z, Weng Y, Song H, Zhu X. 2009.
Heterogeneity of class I and class II MHC sequences in Schistosoma japonicum from different
endemic regions in mainland China. Parasitology research 106:201-206.

875