# Evidence for non-optimal codon choice in highly transcribed sex-biased genes of *Drosophila melanogaster*

**Authors:** Carrie A. Whittle[1], Cassandra G. Extavour[1,2]

[1]Department of Organismic and Evolutionary Biology, Harvard University, 16 Divinity Avenue, Cambridge MA 02138, USA

[2]Department of Molecular and Cellular Biology, Harvard University, 16 Divinity Avenue, Cambridge MA 02138, USA

Corresponding author: Cassandra G. Extavour

Email: extavour@oeb.harvard.edu

Phone: (617)-496-1935

Fax: (617)-496-9507

**ORCIDs**

C.A. Whittle 0000-0002-9331-0520

C.G. Extavour 0000-0003-2922-5855

**Author Contributions** CAW and CGE devised the study, conducted analyses and wrote the manuscript.

**Competing Interest Statement:** None

**Classification:** Research Report, Genetics

**Keywords:** Non-optimal codons, sex-bias, protein folding, tRNAs, *Drosophila*

**This PDF file includes:**

Main Text
Figures 1 to 3

## Abstract

Biases in synonymous codon use occur across multicellular organisms. Optimal codons, defined as those most commonly used in highly transcribed genes, are thought to arise from selection for cost-efficient translation, which would favor codons with abundant matching tRNAs. Such presumed selection is described as optimal codon choice. Non-optimal codons, defined as those least commonly used in highly transcribed genes, may in principle also play important roles, including regulating protein folding or mRNA translation. However, the dynamics of non-optimal codon use remain understudied. Here we rigorously examine non-optimal codon use, using sex-biased genes expressed in the gonads of *Drosophila melanogaster* as a case study. Using this model, we show that as genes evolve sex-biased expression, they tend to evolve a preference for non-optimal codon use, and that this preference is stronger for testis-biased genes than for ovary-biased genes. Further, from thorough analyses of each of 18 degenerate amino acids, we found evidence that the use of non-optimal codons is not random. Instead, specific non-optimal codons are favored over their synonymous non-optimal codons, an effect again especially pronounced in testis-biased genes. Moreover, we found that non-optimal codon use is positively linked to elevated disorder of the encoded proteins, a pattern most consequential in testis-biased genes. Remarkably, all 18 degenerate amino acids were associated with higher disorder when they were encoded by the identified primary non-optimal codon, than when encoded by its sister optimal codon. We hypothesize that selection may have acted to promote non-optimal codon choice to regulate translation. We discuss the putative roles of tRNA gene copy numbers, pleiotropy, and sex-biased expression in the evolution of this level of gene regulation.

## Introduction

The genetic code contains 18 amino acids encoded by synonymous codons (1-5). Optimal codons are defined as those codons most frequently used for amino acids in abundant transcripts, and are thought to promote cost efficient or more accurate translation, a phenomenon often described as optimal codon choice (1, 6). To date, optimal codons have been identified in unicellular (1, 7, 8) and some multicellular organisms, including insects such as flies, beetles, crickets and mosquitoes (3, 7, 9-16). The optimization of codons in highly expressed genes is thought to have resulted from adaptation of the cDNA to favor use of the most common tRNAs in the cells, often measured as the number of matching tRNA genes in the genome, which gives rise to a high ratio of tRNA supply to codon demand (8, 17-22). However, a number of studies have found that some optimal codons have no tRNAs (wobble codons) in certain taxa, a scenario that may slow translation (11, 12, 23). While studies in multicellular taxa have largely focused on optimal codons (1, 6, 16, 19, 24-29), non-optimal codons, defined as those least commonly used in highly expressed genes, may also have translational roles (4, 11, 12, 30-33). When non-optimal codons have few or no matching tRNA genes, some data suggest that their roles could include targeted slowing of translation for co-translational protein folding or to prevent ribosome jamming (4, 30, 33-37). In addition, under a somewhat paradoxical scenario, some non-optimal codons have been found to have abundant tRNAs, and yet are not used as optimal codons. In these cases, this high tRNA abundance to codon use ratio could lead to rapid and preferential translation of specific mRNAs, namely, those mRNAs with unusually high use of the non-optimal codons (11, 12, 38, 39). At present however, studies on non-optimal codon use and its potential molecular consequences remain sparse. Key outstanding questions include whether some non-optimal codons may be favored over others, that is, whether non-optimal codons may exhibit non-random use, and if so, whether favored non-optimal codons may be linked to specific properties of the proteins whose transcripts deploy them.

An attractive model system to study codon use, particularly non-optimal codon use, is sex-biased genes expressed in the gonads of multicellular organisms. Because codon use in such genes is especially apt to affect reproductive success and fitness, highly expressed gonad-biased genes are expected to be subject to heritable translation-related selection on codon use. Indeed, pioneering studies in the insect *D. melanogaster* and in certain plants have shown that optimal codon use differs between sexes, and is lower in male- than in female-biased reproductive genes (13, 27, 40). Thus, gonadal genes exhibit sufficient selection differences in codon use to be detectable on a gene-wide level (e.g., Fop, or frequency of optimal codons (41) (13, 27, 40)), and thus provide an informative system to fill gaps in our knowledge on the

dynamics of non-optimal codon use. *D. melanogaster* as a study system is especially valuable given its vast genomic and gene expression resources available for study (42-44).

Some previous observations suggest that the dynamics of non-optimal codon use in genes may be dependent on protein-folding properties (4, 34, 45). For example, studies of the circadian clock proteins FQR (in *Neurospora*) and PER in *D. melanogaster* have found that experimentally changing non-optimal codons to optimal codons in disordered regions impaired protein form and function, a pattern not found upon optimization of codons in structurally ordered regions (α-helices) (30, 31, 34). Thus, non-optional codons in those two genes may contribute to proper protein structure, particularly for disordered regions (30, 34). Further, changes in a single synonymous codon, including non-optimal codons, have been linked to altered protein folding and disease phenotypes (31, 36, 46-48). Thus, the extent and accuracy of protein folding may be a factor linked to a gene's overall non-optimal codon use (30), or potentially to the use of specific non-optimal codons. One approach to quantify the contribution to protein folding of individual amino acid sites (and thus, on a codon-by-codon level) and also on a gene-wide basis, is to assess the relative solvent accessibility (RSA). The RSA is an index that quantifies the exposed surface area of an amino acid within the protein structure (that reflects secondary and tertiary structures), relative to its maximum possible surface area in Angstroms (49, 50). Low RSA values typically indicate deeply buried hydrophobic amino acids or protein regions, while exposed residues at the protein surface are typically hydrophilic with relatively high RSA values. RSA analyses have proven effective in revealing links between protein folding and the rate of protein sequence evolution, including slow protein evolution in highly ordered (low RSA) regions (51-56). In this regard, it is reasonable to infer that RSA may also provide an effective means to study protein folding properties with respect to optimal and non-optimal codon use.

Here, from the study of highly expressed sex-biased genes in *D. melanogaster*, we provide evidence of non-random non-optimal codon use, that is, selection for the use of non-optimal codons. Specifically, we report a stepwise increase in non-optimal codon use from highly transcribed sexually unbiased, to ovary-biased, to testis-biased genes. We show that certain non-optimal codons are markedly preferred over other sister non-optimal codons, especially in the testis-biased genes. In addition, genes with the highest frequency of non-optimal codon use (>60% of amino acids) show significantly elevated gene wide RSA, and this pattern is most prevalent in testis-biased genes. We further reveal that for a single amino acid, the most strongly favored non-optimal codon per amino acid is consistently linked to elevated RSA as compared to its sister optimal codon, a pattern observed for 18 of 18 degenerate amino acids in

the genetic code. The pronounced accumulation of non-optimal codons in testis-biased genes includes both codons with few and with plentiful matching tRNA gene copy numbers. This suggests that selection for non-optimal codons in testis-biased genes could enhance at least two potential functions, namely slowed translation and targeted up-translation (4, 11, 35)) We hypothesize that the accumulation of favored non-optimal codons in sex-biased genes, and especially in male-biased genes, may be mediated by very low cross-tissue gene pleiotropy (57-59), and by weak or absent potential for sexual conflict (60, 61).

## Results and Discussion

### *A Priori* Classifications of Organismal Optimal and Non-optimal codons in *D. melanogaster*

Prior to analyses of sex-biased gene sets and codon use, we first aimed to define the organism-wide optimal and non-optimal codon status for *D. melanogaster* for each of the 59 codons encoding the 18 degenerate amino acids. For this, we compared the relative synonymous codon usage (RSCU) (62) of a set of highly expressed (ribosomal protein genes) versus all genes in the genome, and defined this value as the $\Delta RSCU_{Ribosome-All}$ (see Materials and Methods (63)). All codon classifications obtained with this method are provided in Table S1. We identified optimal codons as universally G3 and C3 codons, as previously suggested for *D. melanogaster* (3, 13, 28, 64). This pattern of codon use is not explainable by mutational biases, but rather likely arose from selection pressures (3, 28, 64). Herein, for our study, for each of the 18 degenerate amino acids we explicitly defined the (primary) optimal codon (11, 12) as that with the largest positive $\Delta RSCU_{Ribosome-All}$ value compared with its sister codons (Table S1). All other codons for a given amino acid (a total of 41) were classed as non-optimal codons. These typically had a negative or near-zero $\Delta RSCU$ value (and an RSCU value 1.5-fold to six-fold smaller than that of the optimal codon). We then focused on further analysis of these non-optimal codons, which have been minimally considered or excluded in codon use studies to date (3, 13, 28, 64)(see Supplementary Text File S1 for details).

We classified all optimal and non-optimal codons based on the number of exact matching tRNA genes in the *D. melanogaster* genome (Table S1, GtRNA database (65)), because tRNA abundances, often well reflected by gene copy numbers, underlie purported translational functions (8, 11, 17-20, 23, 35)). 12 of the 18 optimal codons had the most abundant perfectly matching tRNA genes among its sister codons (e.g., GCC for Gly with 14 tRNAs; denoted as $Opt_{high-tRNAs}$ status; high/low tRNAs were demarcated by

the median of 5 tRNA genes). The other six optimal codons had no matching tRNAs, and were classed as

wobble codons (Opt$_{wobble}$; e.g. Pro-CCC). In turn, 26 of the 41 non-optimal codons had low tRNA numbers

(Non-opt$_{low-tRNAs}$, status; e.g., ATA for Ile) and the remaining 15 had abundant tRNA genes (Non-opt$_{high-}$

$_{tRNAs}$ status; e.g., ACT for Thr) (Table S1). Hereafter, we focus on the study of sex-biased genes and their

patterns of optimal and non-optimal and optimal codon use (Table S1). When pertinent, we also consider

statuses of their tRNA gene abundances.

**Identification and Characterization of the Sex-Biased Gene Sets**

We identified highly expressed testis-biased (N=916), ovary-biased (N=258) and unbiased

(N=605) genes in *D. melanogaster* (Fig. S1A) using strict criteria of at least five-fold-bias and ≥100

RPKM in at one sex versus the other (testis versus ovaries (mated)) based on transcriptome data from

modEncode (Table S2, (42, 44) see Materials and Methods). For stringency, we also compared gene

expression in the male accessory glands and virgin ovaries (Table S2). We found that 86.9% of the

identified testis-biased genes were also male-biased in accessory glands (using a cutoff of ≥ 5 fold sex-

bias), and that 77.1% of the identified ovary-biased genes were also female-biased in virgin ovaries,

indicating that strong male-female transcriptional differences were also present in those alternate

reproductive tissue types. The predicted functions of these genes based on GO analysis showed testis-

biased genes were strongly linked to testicular roles such as spermatogenesis, and ovary-biased genes

were associated with female reproductive process roles such as the chorion and oogenesis. In contrast,

unbiased genes were preferentially involved in the ribosome and other housekeeping functions (Table S3;

chromosomal locations of genes are discussed in Supplementary Text File S1). In terms of characteristics

of these genes with sex-biased expression, the CDS lengths were shorter for testis-biased (median=273

codons) and unbiased genes (median=269 codons) than for the ovary biased genes (median=498 codons;

MWU tests P<0.05) (Fig. S1B). We then determined the *tau* value, an index of gene expression specificity

with values from 0 to 1 (59), across 59 tissue stages in *D. melanogaster* (Table S2). *tau* exhibited a

stepwise decline from testis-biased genes (median=0.95) to ovary-biased genes (median=0.83) to unbiased

genes (median=0.74) (ranked ANOVA and Dunn's P<0.05 for each paired contrast; median genome-wide

*tau*=0.85, Fig. S1C). Thus, the testis-biased genes exhibited higher specificity, and lower cross-tissue

pleiotropy (58, 66, 67), than ovary-biased and unbiased genes.

We assessed the relationships between tRNA gene numbers, amino acid use, and amino acid size

complexity (S/C) scores, which are thought to reflect biosynthetic costs of amino acids (provided in Table

S4 (68). The tRNA gene copy number per amino acid (pooled across all codons) was strongly positively

correlated to the amino acid use (percentage of each amino acid relative to all concatenated amino acids

per gene set) for all genes (testis-biased genes: Spearman's R=0.91, P<$10^{-7}$; ovary-biased genes: R=0.85,

P<$10^{-7}$; unbiased genes: R=0.83, P<$10^{-7}$; Fig. S2A-C). Moreover, the S/C score per amino acid (68) was

inversely correlated to the organismal tRNA gene numbers (R=-0.59,P<$10^{-7}$, Fig. S2D). In sum, the

patterns suggest that all three gene sets have a history of selection on codon use (8, 11, 18), further

affirming their utility as a model to study dynamics of codon use in this taxon.

By examining the codon adaptation index (CAI), an index that reflects the gene-wide codon

frequencies relative to a highly expressed reference gene set (here, ribosomal genes) (62, 63), we found a

stepwise decrease in CAI from unbiased, to ovary-biased, to testis-biased genes (ranked ANOVA and

Dunn's P<0.05, Fig. S3A; note that CAI was highly correlated to Fop across the genome, Spearman's

R=0.967, P<$10^{-7}$). In other words, as genes evolved increased gonad-biased expression, they also evolved

increases non-optimal codon use, with the highest non-optimal codon use in testis-biased genes (see

Supplementary Text File S1). Further, we found that the expression level (two classes, ≥200 RPKM, and

≥100-200 RPKM) and CDS length (short and long CDS lengths, with long defined as ≥272 codons; cutoff

at the 33$^{rd}$ percentile of all gene lengths) were linked to CAI within each sex-biased gene set (Fig. S3B).

The differences in CAI between gene sets (Fig. S3A) were robust to both transcription level and length

cutoffs (Fig. S3A,B; see also Supplementary Text File S1). The high non-optimal codon use in testis-

biased genes is not due to enhanced genetic interference, as those processes would be expected to

ameliorate non-optimal codon use in longer rather than shorter genes, a pattern that was absent from our

data (Fig. S3) (13, 69-71). The testis-biased genes herein were short (Fig. S1B), and even within the testis-

biased gene set (Fig. S3B), shorter genes (<272 codons) had greater non-optimal codon use (lower CAI)

than longer genes for both expression classes ((MWU-tests P<0.05, Fig. S3B; see Supplementary Text

File S1). Thus, other factors must explain the stepwise increase in non-optimal codon use from unbiased,

to ovary-biased to testis-biased genes.

**Non-optimal Codon Use is Non-random in *D. melanogaster* Sex-biased Genes**

To further characterize codon use in unbiased and sex-biased genes, we assessed codon use on a

codon-by-codon basis for each gene set described above. We determined the ΔRSCU between testis-

biased genes versus ovary-biased genes (ΔRSCU$_{\text{Testis-Ovary}}$), and between testis-biased genes versus

unbiased genes (ΔRSCU$_{\text{Testis-Unbiased}}$), for each of the 59 codons with synonymous codons (*cf.* (27)). In

these contrasts, a positive $\Delta$RSCU value indicates higher use of a codon in the testis-biased genes, and a negative value denotes a reduction in its use within testis-biased genes (and thus elevated use in the respective ovary or unbiased genes, see Materials and Methods). The $\Delta$RSCU$_{\text{Testis-Ovary}}$ and $\Delta$RSCU$_{\text{Testis-Unbiased}}$ values per codon are shown in Table S5. Remarkably, we found that testis-biased genes reduced their use of optimal codons, and this pattern was universal to all of the degenerate amino acids. Specifically, 16 of the 18 optimal codons in *D. melanogaster* had a statistically significant negative $\Delta$RSCU$_{\text{Testis-Ovary}}$ (P<0.05, only exceptions were for Ser and Thr P>0.05), and 18 of 18 optimal codons had a negative value for $\Delta$RSCU$_{\text{Testis-Unbiased}}$. The magnitude of negative $\Delta$RSCU values was consistently larger for the $\Delta$RSCU$_{\text{Testis-Unbiased}}$ contrast than for the $\Delta$RSCU$_{\text{Testis-Ovary}}$ contrast, consistent with a stepwise decline in the use of optimal codons from unbiased genes, to ovary-biased genes to testis-biased genes (Table S5, Fig. S3). This suggests the possibility that there are weaker translational selection pressures for optimal codons in both types of sex-biased genes, and especially in the testis-biased genes. However, another significant hypothesis worthy of consideration here is that the patterns may reflect a non-random displacement of optimal codons by the favored use of non-optimal codons in testis-biased genes. In other words, it could be the case that testis-biased genes reduce their use of optimal codons not arbitrarily, but rather in favor of preferential use of non-optimal codons.

To test this hypothesis, we asked whether there was any pattern to the non-optimal codons that were used in testis-biased genes. We found that for the nine amino acids encoded by two synonymous codons (Asn, Asp, Cys, Gln, Glu, His. Lys, Phe, Tyr), the universal reduction in use of the optimal codon in testis-biased genes was compensated by increased use of the sister non-optimal codon (as indicated by positive $\Delta$RSCU$_{\text{Testis-Unbiased}}$ and $-\Delta$RSCU$_{\text{Testis-Ovary}}$ values, Table S5), consistent with a hypothesis of preferential use of the non-optimal codons in testis-biased genes. Nonetheless, the two-fold degenerate amino acids cannot be used to fully differentiate whether there may also be codon choice among the non-optimal codons. Thus, we examined the nine amino acids that were encoded by three, four or six synonymous codons (Ala, Arg, Gly, Ile, Leu, Pro, Ser, Thr, Val). If the low use of optimal codons in testis-biased genes were solely due to weaker selection for optimal codons, then one would expect a similar degree of enhanced use of all the synonymous non-optimal codons per amino acid in testis-biased genes (under a neutral model). In contrast, however, we found that the elevated use of non-optimal codons in testis-biased genes was non-arbitrary (Fig. 1A, Table S5). As an example, for the four-fold degenerate amino acid Thr, a neutral model would predict similar and increased use of the three non-optimal codons (ACG, ACA, ACT) under reduced use of the optimal codon (ACC). However, we found that only ACT

(Non-opt$_{high\text{-}tRNAs}$) had elevated use in testis-biased genes versus ovary-biased genes ($\Delta$RSCU$_{Testis\text{-}Ovary}$

=+0.166; t-test P<0.05. Fig. 1B), while the other two non-optimal codons showed no elevation in use. This

is not consistent with a neutral model of non-optimal codon use, and instead suggests a profound

preference for one specific non-optimal codon over the others (Fig. 1B). The same preference for ACT

was also observed using the $\Delta$RSCU$_{Testis\text{-}Unbiased}$ comparison ($\Delta$RSCU=+0.201, Table S5), a codon that

was detected even though the ovary-biased and unbiased genes were from fully non-overlapping gene

sets, indicating that the preference for this non-optimal codon in the testis-biased genes is robust to the

two contrasts. Another example is Ile (three-fold degenerate) where the reduced frequency of the optimal

codon ATC in testis-biased versus unbiased genes contrast was accompanied by a strong preferential use

of just one of the two non-optimal codons (ATA; $\Delta$RSCU$_{Testis\text{-}Ovary}$ =+0.114; 9.5 fold higher than that of

its synonymous non-optimal codon ATT; Table S5), and the same non-optimal codon preference was also

identified using the $\Delta$RSCU$_{Testis\text{-}Unbiased}$ contrast (value was +0.266. t-test P<0.05, Fig. 1C, Table S5).

While the same primary non-optimal codons (largest positive $\Delta$RSCU) tended to occur in both the

$\Delta$RSCU$_{Testis\text{-}Ovary}$ and the $\Delta$RSCU$_{Testis\text{-}Unbiased}$ contrasts, the latter were more often statistically significant

(Fig. 1, Table S5). This likely reflects not only the larger $\Delta$RSCU values in the latter contrasts, but also

the larger N values and thus power for the unbiased gene set (N=605 for unbiased, and 258 for ovary-

biased). Overall, the pattern of the highly favored use of a specific non-optimal codon over others per

synonymous codon family in testis-biased genes was also observed for Gly (GGA, Fig. 1D), Ala (GCA),

Leu (TTG), Ser (AGT) and Val (GTA as primary non-optimal codon, GTT as secondary), Pro (CCT,

particularly $\Delta$RSCU$_{Testis\text{-}Ovary}$ contrast) and Arg (AGG as primary, AGA as a close secondary) (Table S5,

Fig. 1A). All primary non-optimal codons ended in A3, T3 or G3 (Table S5), thus excluding a role of

mutational biases for a particular nucleotide at the 3$^{rd}$ "silent" position, and suggesting instead that

selective processes have shaped the use of those codons. Together, these patterns show that testis-biased

genes show little use of use optimal codons, but instead preferentially use specific non-optimal codons,

rather than randomly choosing among the available non-optimal codons.

Notably, the primary non-optimal codon per amino acid (Table S5, Fig. 1A) in testis-biased genes

was not predicted by tRNA status. Specifically, for Arg, Leu and Ser, the primary non-optimal codon had

low tRNA abundance (Non-opt$_{low\text{-}tRNAs}$ status), while others such as Gly, Pro and Thr favored the use of

codons with highly abundant tRNAs (Non-opt$_{high\text{-}tRNAs}$ status (Table S5)). We speculate that this pattern

suggests that selection of non-optimal codons with few tRNAs (Arg, Leu and Ser) may typically be

involved in slowing translation for protein folding or to limit ribosome jamming, while those with

abundant tRNAs (Gly, Pro and Thr) may often be involved in ensuring up-translation of mRNAs carrying those codons. These predictions are based on presumed supply-demand ratios of tRNAs:codons (11, 30, 32, 35, 72). Thus, the primary non-optimal codons identified here may regulate gene function via specific translational roles.

We determined the percentage of genes in each gene set under study that exhibited extreme use of the primary non-optimal codons (defined as RSCU values $\geq 1.5$). As shown in Fig. S4A, a larger percentage of testis-biased genes exhibited extreme use of the primary non-optimal codons than did the comparable ovary-biased and/or unbiased genes (Chi-square P<0.05; shown as examples are Thr-ACT, Gly-GGA, Ile-ATA and Val-GTA; see Supplementary Text File S1 for parallel data for optimal codons, Fig. S4B). It is worth noting that the percent of genes with extreme use of optimal codons with plentiful tRNAs differed from those optimal codons with wobble status (Fig. S4B; in all gene sets, P<0.05). This observation is important as it allows us to infer that these two classes of optimal codons may not be equivalent in functions (see Supplementary Text File S1), and thus in future studies should not be treated as equal optimal codons, nor studied without including tRNA gene data.

**Amino Acids Using Non-optimal Codons Have High Surface Exposure and Disorder**

To determine whether these patterns of non-optimal codon use were linked to specific biophysical properties of the encoded proteins, we first assessed the solvent accessibility RSA value for every amino acid (and its associated codon) in the testis-biased genes (N=299,817 amino acids), ovary-biased genes (155,282 amino acids) and unbiased genes (218,580 amino acids). We found that RSA value was strongly correlated to the disorder probability scores (from NetsurfP (73)) in all three datasets (Spearman's R $\geq 0.68$ for each of three datasets, P<$10^{-7}$). Thus, higher RSA reflects both surface exposure and intrinsic disorder (74-76). In terms of major secondary structures of proteins ($\alpha$-helices, $\beta$-strands and random coils (73)), a majority of amino acids in the proteins of all three gene sets were located within random coils (60.6% for testis-biased, 57.2% of unbiased, and 70.8% for ovary-biased genes, Chi-square tests P<0.05). $\alpha$-helices were the next most common structures in all three gene sets (31.1, 22.4 and 31.5%, respectively), and $\beta$-strands the least common (11.2, 8.3 and 6.8%, Fig. S5A). The lowest RSA values occurred for amino acids located in $\beta$-strands, while these values were intermediate for $\alpha$-helices and highest for random coils for all three gene sets (Fig. S5B, median RSA values across all residues for testis-biased genes were 0.13, 0.34 and 0.53, for ovary-biased genes were 0.11, 0.33 and 0.60, and for unbiased genes were 0.11, 0.30 and 0.57 for $\beta$-strands, $\alpha$-helices and random coils respectively; MWU-tests P<0.05 for

all contrasts per gene set). These patterns are consistent with disordered regions having the greatest surface exposure.

For each gene under study, we then determined the percent of amino acids encoded by non-optimal codons, as a gene-wide direct measure of non-optimal codon use (denoted as Percent-Non-opt). The genes were placed into bins of low ($\leq$50%), moderate (>50 to $\leq$60%) and high Percent-Non-opt codon use (>60%; note that we excluded Met and Typ from this analysis, discussed in Supplementary Text File S1). We found the average RSA per gene increased from the lowest Percent-Non-opt bin (median RSA=0.40), to the moderate (median=0.43) to high (median=0.47) bins (MWU-tests each P<0.05) in testis-biased genes (Fig. 2A). In other words, the elevated non-optimal codon use in these testis-biased genes was linked to greater disorder in their encoded proteins, whereas genes using more optimal codons tended to be more well-folded (Fig. 2A). For ovary-biased genes and unbiased genes, no statistically significant differences were observed in RSA between the highest Percent-Non-opt bin and the moderate and low bins (P>0.05, Fig. 2B,C), likely because the highest non-optimal codon class was so uncommon in those gene sets for testing (N$\leq$23 genes in each case). Nonetheless, the moderate Percent-Non-Opt bins had elevated average RSA per gene as compared to the low bins for both the ovary-biased and the unbiased genes (MWU-tests P<0.05). Overall, elevated non-optimal codon use is associated with greater protein surface exposure in testis-biased, ovary-biased and unbiased genes, with the most striking stepwise correlation observed for the testis-biased genes (Fig. 2A-C).

Importantly, of the 255 genes (out of 1,779 highly expressed genes under study; Fig. S1A) within the highest Percent-Non-opt class (>60% non-optimal codons, Fig. 2D), 83% were testis-biased (N=212), while only 9.0% were ovary-biased (N=23) and 7.8% were unbiased (N=20; Fig. 2D). Thus, the highest degree of non-optimal codon use, and elevated protein RSA and disorder, are core and novel features of a major fraction of the testis-biased genes, but these features are not prevalent among ovary-biased or unbiased genes. Even when taking into account that there were more testis-biased genes than ovary-biased or unbiased genes, we find that 23.1% (212 of 916) of the testis-biased gene set belongs to the highest class bin of non-optimal codon use, versus only 8.9% (23 of 258) and 3.3% (20 of 605) of the ovary-biased and unbiased gene sets respectively. In terms of genes in the moderate Percent-Non-opt bin (Fig. 2E), testis-biased genes were most common in absolute numbers (N=325; that 35.4% of all testis-biased genes), with less than a third as many ovary-biased genes (N=99 of 258, 38.3% of the ovary genes), and even fewer unbiased genes (N=78; 12.9% of unbiased genes). In the lowest Percent-Non-opt bin (with the highest optimal codon use), unbiased genes were most strongly represented with 509 genes (84.1% of the

unbiased gene set), while 389 were testis-biased (42.5% of testis-biased gene set), and 136 were ovary-

biased genes (52.7% of the ovary-biased gene set; Fig. 2F). Overall, it is evident that a major portion of

testis-biased genes were linked to the highest non-optimal codon use bin (Percent-Non-opt>60%), a group

that also had very high protein-wide RSA (Fig. 2A-C). These RSA patterns add another layer of support

(in addition to the results in Fig. 1) to the conclusion that the high non-optimal codon use in *D.*

*melanogaster* genes is non-random, given that it is also negatively associated with the degree of protein

folding in all gene sets under study (Fig. 2A-C), and that that the highest degree of non-optimal codon use

(and thus high RSA) is especially common in testis-biased genes (Fig. 2A,D).

In Figs. 2G-I, we provide examples of the predicted structures of a testis-biased gene from the

highest Percent-Non-opt bin, an ovary-biased gene from the moderate Percent-Non-opt gene bin, and an

unbiased gene from the low Percent-Non-opt bin. The testis-biased sperm-related gene protein Mst57Da

(FBgn0259139) encodes 73.0% of its amino acids with non-optimal codons, and is predicted to be highly

unstructured (Fig.2G) with an average CDS-wide RSA of 0.72±0.09. The ovary-biased gene protein Jabba

(FBgn0259682), that is involved in early embryogenesis (42), has 55.1% of its amino acids encoded by

non-optimal codons, and shows more helical structures and a lower RSA value (0.50 ±0.01; Fig. 2H) than

the sperm protein (Fig. 2G). The unbiased gene encoding ribosomal protein RpL3 has 73.2% optimal

codon use an average RSA of 0.36 ±0.01 (Fig. 2I), and high levels of helical and beta-sheets illustrating

its extensive degree of folding (NetsurfP predictions for all three proteins are in Fig. S6). In sum, these

patterns demonstrate the marked positive associations in non-optimal codon use and RSA.

Functional clustering shows that the 212 testis-biased genes in the highest Percent-Non-opt bin

(>60% non-optimal codons, Fig. 2D) are enriched in genes involved in sperm-related functions (Table

S6). Examples include genes encoding accessory gland proteins Acp54A1*,* Acp26Aa and Acp36DE*,* the

testis-enriched gene *RpL10Aa,* the seminal fluid protein Sfp87B*,* and spermatogenesis genes *soti*, *Mst89B*,

M*st57Da*, and other Mst genes (Table S7; gene functions from DAVID and FlyBase (42, 77)). Thus, these

testis-biased genes with the highest percentage of non-optimal codons were highly specialized to testicular

functions, including spermatogenesis.

***tau suggests a mechanism to accumulate non-optimal codons***

Analyses of gene expression specificity across 59 tissues/stages in *D. melanogaster* (Table S2),

revealed that testis-biased genes had exceptionally elevated *tau* values, with a median near the maximum

possible value of 1 for genes in the high, moderate and low Percent-Non-opt bins (median values≥0.96,

Fig. 2D-F; see Materials and Methods). Thus, the expression of these genes is highly specific to the testis.

In turn, ovary-biased genes, had intermediate *tau* values (median between 0.83 to 0.87) among low,

moderate and high Percent-Non-opt bins, while unbiased genes had the lowest *tau* (0.71 to 0.77) (Fig. 2D-

F). Assuming expression breadth reflects a gene's cross-tissue pleiotropy (58, 66, 67), then testis-biased

genes comprise the least pleiotropic gene set under study. The low pleiotropy of these testis-biased genes

may allow them to be under less severe functional constraint than those involved in a wide range of tissues

or pathways (57, 66, 78, 79), possibly allowing them the flexibility to evolve the observed high non-

optimal codon use (Figs. 1, 2). Moreover, sex-biased gene expression, including sex-specific expression

(79.0% of testis-biased genes were sex-specific herein, Fig. S1 legend), may act as a means to limit sexual

conflict (60, 61, 80). Thus, we predict that non-optimal codon use in the testis-biased genes may typically

be of little or no fitness consequence to the ovaries or other tissues,  facilitating accumulation of non-

optimal codons in testis genes to regulate gene expression for specialized testicular functions.

**All 18 Degenerate Amino Acids have Higher RSA Values When Encoded by the Primary Non-**

**optimal Codon than by the Optimal Codon**

Finally, we assessed the RSA value for each individual amino acid when encoded by the primary

non-optimal codon (Table S5) versus by the optimal codon (Tale S1) in the testis-biased genes. Our

analyses of all amino acids encoded by testis-biased genes showed that for all 18 degenerate amino acids,

the primary non-optimal codon was linked to a higher average RSA than the primary optimal codon (Fig.

3; sign test P<0.001; averages and standard errors are in Table S8). Thus, the elevated RSA of non-optimal

codons included every degenerate amino acid, regardless of its maximum solvent accessibility and its

polarity (which vary among amino acids (50)). When encoded by a non-optimal codon, residues were

consistently located in less deeply folded protein regions, than when that same amino acid was encoded

by its optimal codon. This result includes the primary non-optimal codons Thr-ACT, Ile-ATA, and Gly-

GGA (Fig. 2B-D) and spans all two-fold, three-fold, four-fold and six-fold degenerate amino acids (Table

S5). Thus, the favored use of the primary non-optimal codons versus the optimal codon in more disordered

regions was universal across all 18 degenerate amino acids.

In addition, the primary non-optimal codons, which all exhibited elevated RSA compared to

optimal codons, included those with few and those with plentiful matching tRNA genes (i.e. with Non-

opt$_{low-tRNAs}$ and Non-opt$_{high-tRNAs}$ statuses (Fig. 3, Table 1, Table S5; optimal codons included both Opt$_{high-}$

$_{tRNAs}$ and Opt$_{wobble}$ statuses). The rarity of exact matching tRNAs for Non-opt$_{low-tRNAs}$ codons (Table 1,

Table S5, Fig. 3) may make these types of codons poised to slow translation, and allow proper co-

translational protein folding (32, 35), including in disordered regions (30). However, the high ratio of

tRNA genes to codon use for Non-opt$_{high\text{-}tRNAs}$ codons suggests that this class of codons likely has different

functions, possibly in the rapid translation of mRNA segments containing the codons, or the up-translation

of any mRNAs with high use of these (otherwise) rare codons within a cellular mRNA pool (11, 12, 39).

Thus, despite potentially having distinct functions in slowing translation (Non-opt$_{low\text{-}tRNAs}$) or in targeted

enhanced translation of mRNAs (Non-opt$_{high\text{-}tRNAs}$)) (4, 11, 30-32, 35, 72), the two types of non-optimal

codons (Table 1), are each consistently favored over optimal codons in less-structured protein segments

(Fig. 3). Based on these findings, we speculate that the combined use of primary Non-opt$_{low\text{-}tRNAs}$ and Non-

opt$_{high\text{-}tRNAs}$ codons (Fig. 3) may respectively act to slow down and speed up local rates of translation

within specific types of mRNAs (for instance, in highly transcribed mRNAs that have ameliorated use

these otherwise rare codons, such as testis and sperm genes, Table S5, Fig. 3), particularly within the

disordered regions, thereby providing a means to control translational pacing and potentially allowing

proper protein conformation and functionality.

**Conclusions and Future Directions**

The collective evidence herein supports a model of selection for non-optimal codon use. Non-

optimal codons are non-arbitrarily used in sex-related genes, and certain non-optimal codons are favored

over others, a pattern particularly prevalent in testis-biased genes, and with likely functions in regulating

translation.

Future studies should assess the prevalence and evidence for selection for non-optimal codon use

in other multicellular organisms, including a system whereby non-optimal codons are more common in

the ovaries (15) or in other specific organ systems. In addition, investigation of the placement of all four

categories of codons studied herein, namely Opt$_{high\text{-}tRNAs}$, Opt$_{wobble}$, Non-opt$_{low\text{-}tRNAs}$ and Non-opt$_{high\text{-}tRNAs}$,

with respect to the starting ramp of translation, versus middle and terminal CDS regions, and especially

with respect to the RSA in disordered regions, will help reveal whether these codons may cause localized

pacing (fast/slow) of translation (35, 81, 82) in *D. melanogaster* and other multicellular models.

Experimental studies to quantify translation rates and protein folding of testis genes with extreme use of

Non-opt$_{high\text{-}tRNAs}$ codons (Fig. S4, Table S5; cf. (30, 83)) may further reveal their putative roles in the rapid

translation of specific mRNAs in a cellular mRNA pool or of mRNA regions (11, 39), while investigations

of those genes with extreme use of Non-opt$_{low\text{-}tRNAs}$ codons (Fig. S4) may inform on their function in the

slowed translation and precise protein conformation in testis-involved genes, as inferred for circadian genes (4, 30, 31). Additional data on the tRNA populations in the testes and ovaries in this species as the technology improves (38) (discussed in Supplementary Text File S1), will also help determine whether or how there is variation in tRNA populations among the gonads, beyond that predicted by tRNA gene frequency (Fig. 2). Research should also evaluate whether younger testis-biased genes have greater flexibility to evolve a set of preferred non-optimal codons with functional roles (*cf.*(83)). Finally, *in vitro* or *in vivo* studies that replace non-optimal codons with optimal codons (30, 31, 83), and vice-versa, to evaluate their effects on protein structure and functions for testis-biased and ovary-biased genes studied here, may provide further insights into the potential specific protein biophysical roles of the non-optimal codons in this species. As part of such analyses, an evaluation of altered Non-opt$_{low-tRNAs}$ and Non-opt$_{high-tRNAs}$ statuses within disordered protein regions may reveal their individual roles on the pacing of translation, protein conformation and protein functionality.

**Materials and Methods**

**Genomic Data and Expression Analyses**

We downloaded the genome of *D. melanogaster* (13,986 protein coding genes) version 6.50 (42). For our analysis, we extracted the longest CDS per gene (11, 12, 84). To identify sex-biased genes, we used the modEncode database that contains expression data (42, 44) for a wide range of *D. melanogaster* tissue types and developmental stages (N=59 tissues/stages studied herein, Table S2 (42-44)), and that we also used to measure *tau* (59) across all tissues. We identified highly expressed sex-biased genes as those with ≥100RPKM and ≥5 fold sex-bias in one sex versus the other in the contrast of 4-day mated testes versus 4-day mated ovaries, and unbiased genes as those with <5 fold bias (and ≥100RPKM in at least one gonad). For additional rigor, we also identified sex-biased genes using male accessory glands versus virgin ovaries (Table S2). Further details on identification of sex-biased genes and calculations of expression specificity using *tau* ((59), described previously in (85) using the data in Table S2, are provided in Supplementary Text File S1.

**Codon Use and ΔRSCU**

The organism-wide RSCU values (Table S1) for *D. melanogaster* were obtained from the Codon Statistics Database (CSD) (63). From these, we calculated the $\Delta RSCU_{Ribosome-All}$ =RSCU$_{Ribosomal\ genes}$ -

RSCU$_{\text{All genes}}$ (using the concatenated genes per dataset), which is a more conservative version of the

approach using the highest and least expressed genes (25-28, 86). We identified the largest positive

ΔRSCU$_{\text{Ribosome-All}}$ values as the optimal codon (and each was also identified as optimal at the CSD, with

P-values<0.05). All other codons per amino acid were classed as non-optimal, given their large disparity

in use from the optimal codon (see Supplementary Text File S1 section "Genome Datasets and Gonadal

Expression Analyses"). The tRNA gene counts in the *D. melanogaster* genome were determined using the

GtRNA database ((http://gtrnadb.ucsc.edu) (65) which utilizes tRNAscan-SE to extract tRNAs from the

genome (87, 88). CAI values that reflect adaption similarity of codon use to the highly expressed

ribosomal gene set, and the Fop values, were obtained from the CSD (63).

For the study of differences codon use among sex-biased genes, we determined herein the RSCU

values and the codon raw counts per codon for every testis-biased, ovary-biased and unbiased gene using

Codon W (89). For each of the studied 59 codons encoding a degenerate amino acid, the RSCU values

were used to calculate ΔRSCU$_{\text{Testis-Ovary}}$=average RSCU$_{\text{Testis-biased genes}}$ – average RSCU$_{\text{Ovary-biased genes}}$ and

the ΔRSCU$_{\text{Testis-Unbiased}}$=average RSCU$_{\text{Testis-biased genes}}$ – average RSCU$_{\text{Unbiased genes}}$. The RSCU values per

codon were compared across all genes per contrast using t-tests to obtain P-values (27). The codon counts

per gene under study (for each testis-biased, ovary-biased and unbiased gene) were used to calculate the

Percent-Non-opt codon use per gene (relative to all optimal and non-optimal codons per gene), based on

the non-optimal and optimal codon identities that were defined in Table S1 (see also Supplementary Text

File S1).

**475    Protein Folding, Relative Solvent Accessibility and Codon Use in the Sex-biased Genes**

To determine the RSA per amino acid we used NetsurfP v3.0  set to default parameters (73). This

program is a machine-learning based tool proven effective for determining RSA and other structural data

of proteins (73, 90). NetsurfP v3.0 has been widely used to characterize protein folding and structures (73,

90), and its RSA calculations have been shown to be equivalent to those predicted by other top predictor

tools such as SPOT-1D-Single (73). We also found strong correlations using protein structures from the

Protein Data Base and RSA from GETAREA (49) (data not shown). We determined the type of secondary

structure that each amino acid was located within, namely a helix, string (β-string/sheet) or a random coil,

and the disorder scores per amino acid, using NetsurfP 3.0 (73).

Protein folding predictions for the visualization of structures in Fig. 2G-I were determined using

Alphafold2 (91), available for the *D. melanogaster* proteins (Uniprot proteome Identity UP000000803,

taxon identity= 7227) from the AlphaFold2 database ([https://alphafold.ebi.ac.uk](https://alphafold.ebi.ac.uk)) (92). Visualization was
conducted using Geneious Prime (93).
488
489 **GO Functions**
490 Gene ontology and functional analysis was performed using DAVID (77) .
491

## Acknowledgements

493 The authors thank members of the Extavour lab for valuable discussions. C. Extavour is an
494 Investigator of the Howard Hughes Medical Institute, which supported this work.

# References

1. T. Ikemura, Correlation between the abundance of *Escherichia coli* transfer RNAs and the occurrence of the respective codons in its protein genes: a proposal for a synonymous codon choice that is optimal for the *E. coli* translational system. *J. Mol. Biol.* **151**, 389-409 (1981).
2. P. M. Sharp, T. M. Tuohy, K. R. Mosurski, Codon usage in yeast: cluster analysis clearly differentiates highly and lowly expressed genes. *Nucleic acids research* **14**, 5125-5143 (1986).
3. D. C. Shields, P. M. Sharp, D. G. Higgins, F. Wright, "Silent" sites in Drosophila genes are not neutral: evidence of selection among synonymous codons. *Mol. Biol. Evol.* **5**, 704-716 (1988).
4. Y. Liu, Q. Yang, F. Zhao, Synonymous but Not Silent: The Codon Usage Code for Gene Expression and Protein Folding. *Annu. Rev. Biochem.* **90**, 375-401 (2021).
5. J. B. Plotkin, G. Kudla, Synonymous but not the same: the causes and consequences of codon bias. *Nature Reviews Genetics* **12**, 32-42 (2011).
6. R. Hershberg, D. A. Petrov, General rules for optimal codon choice. *PLoS genetics* **5**, e1000556 (2009).
7. T. Ikemura, Codon usage and tRNA content in unicellular and multicellular organisms. *Mol. Biol. Evol.* **2**, 13-34 (1985).
8. H. Akashi, Translational selection and yeast proteome evolution. *Genetics* **164**, 1291-1303 (2003).
9. N. Galtier *et al.*, Codon Usage Bias in Animals: Disentangling the Effects of Natural Selection, Effective Population Size, and GC-Biased Gene Conversion. *Mol Biol Evol* **35**, 1092-1103 (2018).
10. E. N. Moriyama, J. R. Powell, Codon usage bias and tRNA abundance in *Drosophila*. *J. Mol. Evol.* **45**, 514-523 (1997).
11. C. A. Whittle, A. Kulkarni, N. Chung, C. G. Extavour, Adaptation of codon and amino acid use for translational functions in highly expressed cricket genes. *BMC Genomics* **22**, 234 (2021).
12. C. A. Whittle, A. Kulkarni, C. G. Extavour, Evidence of multifaceted functions of codon usage in translation within the model beetle *Tribolium castaneum*. *DNA Res.* **26**, 473-484 (2019).
13. T. M. Hambuch, J. Parsch, Patterns of synonymous codon usage in Drosophila melanogaster genes with sex-biased expression. *Genetics* **170**, 1691-1700 (2005).
14. A. Williford, J. P. Demuth, Gene expression levels are correlated with synonymous codon usage, amino acid composition, and gene architecture in the red flour beetle, *Tribolium castaneum*. *Mol. Biol. Evol.* **29**, 3755-3766 (2012).
15. C. A. Whittle, C. G. Extavour, Rapid Evolution of Ovarian-Biased Genes in the Yellow Fever Mosquito (Aedes aegypti). *Genetics* **206**, 2119-2137 (2017).
16. S. K. Behura, D. W. Severson, Coadaptation of isoacceptor tRNA genes and codon usage bias for translation efficiency in *Aedes aegypti* and *Anopheles gambiae*. *Insect Mol. Biol.* **20**, 177-187 (2011).
17. H. Gingold, O. Dahan, Y. Pilpel, Dynamic changes in translational efficiency are deduced from codon usage of the transcriptome. *Nucleic Acids Res.* **40**, 10053-10063 (2012).
18. M. Z. Du *et al.*, Co-adaption of tRNA gene copy number and amino acid usage influences translation rates in three life domains. *DNA Res.* **24**, 623-633 (2017).
19. L. Duret, tRNA gene number and codon usage in the *C. elegans* genome are co-adapted for optimal translation of highly expressed genes. *Trends Genet.* **16**, 287-289 (2000).
20. R. Percudani, A. Pavesi, S. Ottonello, Transfer RNA gene redundancy and translational selection in *Saccharomyces cerevisiae*. *J. Mol. Biol.* **268**, 322-330 (1997).

540    21.    M. Zhou *et al.*, Non-optimal codon usage affects expression, structure and function of clock
541           protein FRQ. *Nature* **495**, 111-115 (2013).
542    22.    M. dos Reis, R. Savva, L. Wernisch, Solving the riddle of codon usage preferences: a test for
543           translational selection. *Nucleic Acids Res.* **32**, 5036-5044 (2004).
544    23.    M. Stadler, A. Fire, Wobble base-pairing slows in vivo translation elongation in metazoans. *RNA*
545           **17**, 2063-2073 (2011).
546    24.    H. Akashi, Gene expression and molecular evolution. *Curr. Op. Genet. Dev.* **11**, 660-666 (2001).
547    25.    A. D. Cutter, J. D. Wasmuth, M. L. Blaxter, The evolution of biased codon and amino acid usage
548           in nematode genomes. *Mol. Biol. Evol.* **23**, 2303-2315 (2006).
549    26.    P. K. Ingvarsson, Molecular evolution of synonymous codon usage in Populus. *BMC Evol. Biol.*
550           **8**, 307 (2008).
551    27.    C. A. Whittle, M. R. Malik, J. E. Krochko, Gender-specific selection on codon usage in plant
552           genomes. *BMC Genomics* **8**, 169-179 (2007).
553    28.    L. Duret, D. Mouchiroud, Expression pattern and, surprisingly, gene length shape codon usage in
554           Caenorhabditis, Drosophila, and Arabidopsis. *Proc. Natl. Acad. Sci. USA* **96**, 4482-4487 (1999).
555    29.    B. Wang *et al.*, Optimal codon identities in bacteria: implications from the conflicting results of
556           two different methods. *PLoS ONE* **6**, e22714 (2011).
557    30.    M. Zhou, T. Wang, J. Fu, G. Xiao, Y. Liu, Nonoptimal codon usage influences protein structure
558           in intrinsically disordered regions. *Mol. Microbiol.* **97**, 974-987 (2015).
559    31.    J. Fu *et al.*, Codon usage affects the structure and function of the Drosophila circadian clock
560           protein PERIOD. *Genes Dev* **30**, 1761-1775 (2016).
561    32.    C. E. Brule, E. J. Grayhack, Synonymous Codons: Choose Wisely for Expression. *Trends Genet.*
562           **33**, 283-297 (2017).
563    33.    T. Tuller *et al.*, An evolutionarily conserved mechanism for controlling the efficiency of protein
564           translation. *Cell* **141**, 344-354 (2010).
565    34.    C. H. Yu *et al.*, Codon Usage Influences the Local Rate of Translation Elongation to Regulate
566           Co-translational Protein Folding. *Mol Cell* **59**, 744-754 (2015).
567    35.    K. C. Stein, J. Frydman, The stop-and-go traffic regulating protein biogenesis: How translation
568           kinetics controls proteostasis. *J. Biol. Chem.* **294**, 2076-2084 (2019).
569    36.    D. P. Letzring, K. M. Dean, E. J. Grayhack, Control of translation efficiency in yeast by codon-
570           anticodon interactions. *RNA* **16**, 2516-2528 (2010).
571    37.    S. Pechmann, J. Frydman, Evolutionary conservation of codon optimality reveals hidden
572           signatures of cotranslational folding. *Nat Struct Mol Biol* **20**, 237-243 (2013).
573    38.    J. C. Guimaraes *et al.*, A rare codon-based translational program of cell proliferation. *Genome*
574           *Biol.* **21**, 44 (2020).
575    39.    M. Torrent, G. Chalancon, N. S. de Groot, A. Wuster, M. Madan Babu, Cells alter their tRNA
576           abundance to selectively regulate protein synthesis during stress conditions. *Sci Signal* **11**, DOI:
577           10.1126/scisignal.aat6409 (2018).
578    40.    Z. Zhang, T. M. Hambuch, J. Parsch, Molecular evolution of sex-biased genes in Drosophila.
579           *Mol. Biol. Evol.* **21**, 2130-2139 (2004).
580    41.    S. Bahiri-Elitzur, T. Tuller, Codon-based indices for modeling gene expression and transcript
581           evolution. *Comput Struct Biotechnol J* **19**, 2646-2663 (2021).
582    42.    L. S. Gramates *et al.*, FlyBase: a guided tour of highlighted features. *Genetics* **220** (2022).
583    43.    J. J. Li, H. Huang, P. J. Bickel, S. E. Brenner, Comparison of *D. melanogaster* and *C. elegans*
584           developmental stages, tissues, and cells by modENCODE RNA-seq data. *Genome Res.* **24**, 1086-
585           1101 (2014).

44.  B. R. Graveley *et al.*, The developmental transcriptome of *Drosophila melanogaster*. *Nature*
**471**, 473-479 (2011).
45.  Y. M. Zalucki, M. P. Jennings, Experimental confirmation of a key role for non-optimal codons
in protein export. *Biochem Biophys Res Commun* **355**, 143-148 (2007).
46.  C. Kimchi-Sarfaty *et al.*, A "silent" polymorphism in the MDR1 gene changes substrate
specificity. *Science* **315**, 525-528 (2007).
47.  S. Kirchner *et al.*, Alteration of protein function by a silent polymorphism linked to tRNA
abundance. *PLoS Biol.* **15**, e2000779 (2017).
48.  H. Goodarzi *et al.*, Modulated Expression of Specific tRNAs Drives Gene Expression and
Cancer Progression. *Cell* **165**, 1416-1427 (2016).
49.  R. Fraczkiewicz, Braun, W., Exact and efficient analytical calculation of the accessible surface
areas and their gradients for macromolecules. *J. Comput. Chem.* **19**, 319-333 (1998).
50.  M. Z. Tien, A. G. Meyer, D. K. Sydykova, S. J. Spielman, C. O. Wilke, Maximum allowed
solvent accessibilites of residues in proteins. *PLoS One* **8**, e80635 (2013).
51.  M. P. Scherrer, A. G. Meyer, C. O. Wilke, Modeling coding-sequence evolution within the
context of residue solvent accessibility. *BMC Evol. Biol.* **12**, 179 (2012).
52.  G. C. Conant, P. F. Stadler, Solvent exposure imparts similar selective pressures across a range
of yeast proteins. *Mol Biol Evol* **26**, 1155-1161 (2009).
53.  E. A. Franzosa, Y. Xia, Structural determinants of protein evolution are context-sensitive at the
residue level. *Mol Biol Evol* **26**, 2387-2395 (2009).
54.  A. F. Moutinho, F. F. Trancoso, J. Y. Dutheil, The Impact of Protein Architecture on Adaptive
Evolution. *Mol Biol Evol* **36**, 2013-2028 (2019).
55.  J. Echave, S. J. Spielman, C. O. Wilke, Causes of evolutionary rate variation among protein sites.
*Nat. Rev. Genet.* **17**, 109-121 (2016).
56.  R. Bricout, D. S. Weil, D., A. C. Genovesio, H.R., Evolution is not uniform along protein
sequences. *Molecular Biology and Evolution* **40**, msad042 (2023).
57.  S. P. Otto, Two steps forward, one step back: the pleiotropic effects of favoured alleles. *Proc.*
*Biol. Sci.* **271**, 705-714 (2004).
58.  J. E. Mank, H. Ellegren, Are sex-biased genes more dispensable? *Biol. Lett.* **5**, 409-412 (2009).
59.  I. Yanai *et al.*, Genome-wide midrange transcription profiles reveal expression level
relationships in human tissue specification. *Bioinformatics* **21**, 650-659 (2005).
60.  H. Ellegren, J. Parsch, The evolution of sex-biased genes and sex-biased gene expression. *Nat.*
*Rev. Genet.* **8**, 689-698 (2007).
61.  H. W. S. Wong, L. Holman, Pleiotropic fitness effects across sexes and ages in the Drosophila
genome and transcriptome. *Evolution* 10.1093/evolut/qpad163 (2023).
62.  P. M. Sharp, W. H. Li, The codon Adaptation Index--a measure of directional synonymous
codon usage bias, and its potential applications. *Nucleic Acids Res.* **15**, 1281-1295 (1987).
63.  K. Subramanian, B. Payne, F. Feyertag, D. Alvarez-Ponce, The Codon Statistics Database: A
Database of Codon Usage Bias. *Mol Biol Evol* **39** (2022).
64.  A. Heger, C. P. Ponting, Variable strength of translational selection among 12 Drosophila
species. *Genetics* **177**, 1337-1348 (2007).
65.  P. P. Chan, T. M. Lowe, GtRNAdb 2.0: an expanded database of transfer RNA genes identified
in complete and draft genomes. *Nucleic Acids Res.* **44**, D184-189 (2016).
66.  R. P. Meisel, Towards a more nuanced understanding of the relationship between sex-biased
gene expression and rates of protein-coding sequence evolution. *Mol. Biol. Evol.* **28**, 1893-1900
(2011).

67.  R. Assis, Q. Zhou, D. Bachtrog, Sex-biased transcriptome evolution in Drosophila. *Genome Biol.*
*Evol.* **4**, 1189-1200 (2012).
68.  M. J. Dufton, Genetic code synonym quotas and amino acid complexity: cutting the cost of
proteins? *J. Theor. Biol.* **187**, 165-173 (1997).
69.  L. Loewe, B. Charlesworth, Background selection in single genes may explain patterns of codon
bias. *Genetics* **175**, 1381-1393 (2007).
70.  A. J. Betancourt, D. C. Presgraves, Linkage limits the power of natural selection in Drosophila.
*Proc. Natl. Acad. Sci. USA* **99**, 13616-13620 (2002).
71.  J. M. Comeron, M. Kreitman, M. Aguade, Natural selection on synonymous sites is correlated
with gene length and recombination in Drosophila. *Genetics* **151**, 239-249 (1999).
72.  C. A. Brackley, M. C. Romano, M. Thiel, The dynamics of supply and demand in mRNA
translation. *PLoS Comput. Biol.* **7**, e1002203 (2011).
73.  M. H. Hoie *et al.*, NetSurfP-3.0: accurate and fast prediction of protein structural features by
protein language models and deep learning. *Nucleic Acids Res.* **50**, W510-W515 (2022).
74.  D. Piovesan, A. M. Monzon, S. C. E. Tosatto, Intrinsic protein disorder and conditional folding
in AlphaFoldDB. *Protein Sci.* **31**, e4466 (2022).
75.  M. Akdel *et al.*, A structural biology community assessment of AlphaFold2 applications. *Nat.*
*Struct. Mol. Biol.* **29**, 1056-1067 (2022).
76.  D. Ilzhofer, M. Heinzinger, B. Rost, SETH predicts nuances of residue disorder from protein
embeddings. *Front Bioinform* **2**, 1019597 (2022).
77.  B. T. Sherman *et al.*, DAVID: a web server for functional enrichment analysis and functional
annotation of gene lists (2021 update). *Nucleic Acids Res.* **50**, W216-W221 (2022).
78.  R. Dean, J. E. Mank, Tissue Specificity and Sex-Specific Regulatory Variation Permit the
Evolution of Sex-Biased Gene Expression. *Am. Nat.* **188**, E74-E84 (2016).
79.  J. E. Mank, L. Hultin-Rosenberg, M. Zwahlen, H. Ellegren, Pleiotropic constraint hampers the
resolution of sexual antagonism in vertebrate gene expression. *Am. Nat.* **171**, 35-43 (2008).
80.  F. C. Ingleby, I. Flis, E. H. Morrow, Sex-biased gene expression and sexual conflict throughout
development. *Cold Spring Harb Perspect Biol* **7**, a017632 (2014).
81.  T. Quax, N. Claassens, Soll D, J. van der Ooost, Codon Bias as a Means to Fine-Tune Gene
Expression. *Mol. Cell* **59**, 149-161 (2015).
82.  J. B. Miller, L. R. Brase, P. G. Ridge, ExtRamp: a novel algorithm for extracting the ramp
sequence based on the tRNA adaptation index or relative codon adaptiveness. *Nucleic Acids Res.*
**47**, 1123-1131 (2019).
83.  S. R. Allen *et al.*, Distinct responses to rare codons in select *Drosophila* tissues. *Elife* **11** (2022).
84.  T. Sandmann, M. C. Vogg, S. Owlarn, M. Boutros, K. Bartscherer, The head-regeneration
transcriptome of the planarian Schmidtea mediterranea. *Genome Biol.* **12**, R76 (2011).
85.  C. A. Whittle, C. G. Extavour, Revealing the links between gene protein sequence evolution and
the rapid divergence of ovariole numbers in *Drosophila*. *Biorxiv*  (2023).
86.  B. Wang *et al.*, Optimal codon identities in bacteria: implications from the conflicting results of
two different methods. *PLoS One* **6**, e22714 (2011).
87.  T. M. Lowe, P. P. Chan, tRNAscan-SE On-line: integrating search and context for analysis of
transfer RNA genes. *Nucleic Acids Res.* **44**, W54-57 (2016).
88.  P. P. Chan, T. M. Lowe, tRNAscan-SE: Searching for tRNA Genes in Genomic Sequences.
*Methods Mol. Biol.* **1962**, 1-14 (2019).
89.  J. F. Peden (1999) Analysis of codon usage.  (University of Nottingham).

90. M. S. Klausen *et al.*, NetSurfP-2.0: Improved prediction of protein structural features by
integrated deep learning. *Proteins* **87**, 520-527 (2019).
91. T. Juniper *et al.*, Use of beta-D-glucan in diagnosis of suspected *Pneumocystis jirovecii*
pneumonia in adults with HIV infection. *Int J STD AIDS* **32**, 1074-1077 (2021).
92. M. Varadi *et al.*, AlphaFold Protein Structure Database: massively expanding the structural
coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res.* **50**, D439-
D444 (2022).
93. M. Kearse *et al.*, Geneious Basic: an integrated and extendable desktop software platform for the
organization and analysis of sequence data. *Bioinformatics* **28**, 1647-1649 (2012).

Fig. 1. A) The values for $\Delta RSCU_{Testis-Ovary}$ (left panel) and $\Delta RSCU_{Testis-Unbiased}$ (right panel) per codon for amino acids with three or more synonymous codons (P-values in Table S5). The amino acid-codon and the codon status (left) with respect to tRNAs (right) are each indicated. Close-up examples of the $\Delta RSCU$ for B) all four codons for Thr; C) all three codons for Ile; and D) all four codons for Gly (* indicates P<0.05 in B-D). In panel A, $\Delta RSCU_{Testis-Unbiased}$ for Leu-CTG $Opt_{high-tRNAs}$ =0.818 and Arg-CGC $Opt_{wobble}$=1.11 and were outside the shown limit of X-axis limit.

Fig. 2. A-C: The relationships between high non-optimal codon use (>60% of amino acids), moderate non-optimal use (50 to ≤60%) and low optimal codon use (≤50%) and the relative solvent accessibility for: A) testis-biased genes; B) ovary-biased genes; and C) unbiased genes in *D. melanogaster*. Pie charts of the number genes with D) high non-optimal codon use; E) moderate non-optimal; and F) high optimal codon use, also showing the median values of *tau* across genes. Examples of AlphaFold2 predicted protein structures of a typical: G) testis-biased gene with high non-optimal codons and RSA; H) ovary-biased genes wth moderate non-optimal codons and RSA; and I) an unibiased gene with low non-optimal codons and RSA. In A-C, different letters below bars indicate a statistically signficant difference (MWU-test P<0.05)

Fig. 3.  The average RSA of testis-biased genes when encoded by the primary non-optimal codon (defined in Table S5) versus the optimal codon (Table S1) for all 18 degenerate amino acids. Results are from all amino acids and codons in testis-biased genes (N= N=299,817 amino acids). In total, 18 of 18 degenerate amino acids had higher average RSA when encoded by the non-optimal codon, and sign test across all amino acids had P<0.001. The standard errors were very small (all <0.005) and are provided in Table S8.