Discovery of allodiploid hybrid species *Aspergillus latus* in Japan

Saho Shibata[1], Momotaka Uchida[1], Sayaka Ban[1], Katsuhiko Kamei[1], Akira Watanabe[1], Takashi

Yaguchi[1], Vit Hubka[2,3], Hiroki Takahashi[1,4,5]

[1]Medical Mycology Research Center, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba 260-8673,

Japan

[2]Laboratory of Fungal Genetics and Metabolism, Institute of Microbiology, Czech Academy of

Sciences, Vídeňská 1083, 142 00 Prague 4, Czech Republic

[3]Department of Botany, Faculty of Science, Charles University, Benátská 2, 128 00 Prague 2,

Czech Republic

[4]Molecular Chirality Research Center, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba 263-

8522, Japan.

[5]Plant Molecular Science Center, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba 260-8675,

Japan.

Address for correspondence: Hiroki Takahashi

E-mail: hiroki.takahashi@chiba-u.jp

ORCID

S. Shibata: 0000-0003-2336-6948

M. Uchida: 0000-0001-5642-1435

S. Ban: 0000-0003-3442-308X

K. Kamei: 0000-0002-6067-9049

A. Watanabe: 0000-0002-3057-2937

T. Yaguchi: 0000-0002-9697-8470

V. Hubka: 0000-0003-4583-6496

H. Takahashi: 0000-0001-5627-1035

Abstract

Allodiploid hybrid species, *Aspergillus latus*, belonging to section *Nidulantes*, is a hybrid of *A.*

spinulosporus and unknown species closely related to *A. quadrilineatus* and *A. subltus*. This

hybrid has often been misidentified as the species in section *Nidulantes*, such as *A. nidulans*, *A.*

spinulosporus, *A. sublatus*, or other cryptic species. *A. latus* has not been reported in Japan as

well as Asia so far. In this study, we screened 23 clinical strains identified as *A. spinulosporus*

isolated in Japan from 2012 to 2023 and found seven *A. latus* strains. To characterize the *A. latus*

strains, we conducted comprehensive phenotyping including morphological observation, whole

genome sequences and phylogenetic analysis based on calmodulin (*CaM*) gene. In addition, we

conducted antifungal susceptibility testing for *A. latus* strains. As a result, the morphological

characters of *A. latus* were more similar to those of *A. spinulosporus* compared to *A. sublatus*.

However, the ascospore of *A. latus* differed from that of *A. spinulosporus*. *A. latus*

phylogenetically clustered with *A. spinulosporus* and *A. sublatus*. Furthermore, *A. latus* strains

showed reduced susceptibility to caspofungin and amphotericin B compared to *A. spinulosporus*,

while they were susceptible to azoles. Our results suggest that *A. latus* has been a causative

pathogen of aspergillosis in Japan since 2013.

Keywords: *Aspergillus latus, Aspergillus spinulosporus*, *Aspergillus sublatus*, section *Nidulantes*,

aspergillosis

Introduction

Aspergillus species are the causal agents for aspergillosis, mainly affecting immunocompromised individuals [1, 2]. Although *A. fumigatus* is a main causative agent, other species such as *A. nidulans*, *A. flavus*, *A. niger*, and *A. terreus* also contribute to the disease [3–6]. *Aspergillus nidulans*, a member of the section *Nidulantes* poses a serious threat to immunocompromised patients, especially those with chronic granulomatous disease [3]. Additionally, eleven other species within the section *Nidulantes* have been isolated from patients with invasive aspergillosis [7]. Recently, Steenwyk et al. (2020) through screening of *A. nidulans* strains identified that *A. latus* is an allodiploid hybrid species between *A. spinulosporus* and unknown species closely related to *A. quadrilineatus*, and exhibits higher virulence and minimum inhibitory concentrations to some antifungals than *A. nidulans* [8]. This highlights the importance of accurate species identification to ensure effective treatment, as misidentification could compromise patient care.

Steenwyk et al. (2020 & 2024) reported 30 *A. latus* strains from Belgium, Brazil, France, Germany, Netherlands and Portugal identified as *A. nidulans*, *A. spinulosporus*, or other cryptic species, suggesting that *A. latus* infections are likely underreported [9]. Prospective survey in single center in Japan reported that six *A. nidulans* strains were identified among 107 *Aspergillus* strains [11]. To date, *A. latus* has not been reported in Asia including Japan. However, *A. nidulans*

and some cryptic species were isolated from patients in Japan, namely *A. latus* could had also

been isolated [10, 11].

In this study, we identified seven clinical *A. latus* strains isolated in Japan by screening 23 *A.*

spinulosporus strains which were collected between 2012 and 2023. To characterize these seven

A. latus strains isolated in Japan, we performed comprehensive phenotyping including

morphological observations, whole genome sequences and phylogenetic analysis based on

calmodulin (*CaM*) gene. In addition, we measured the antifungal susceptibility of these strains to

six antifungals. Here, we present the first report of *A. latus* being isolated in Japan. Our findings

indicate that *A. latus* has been the causative agent of aspergillosis in Japan since 2013.

Material and methods

Strains

A. latus, *A. sublatus* and *A. spinulosporus* strains used in this study have been stored and

maintained at the Medical Mycology Research Center, Chiba University (IFM strains) in Japan.

A total of 23 strains of *A. latus* and *A. spinulosporus* are clinical isolate in Japan, and *A. sublatus*

IFM 42029 (= IFM 4553 = CBS 140630) is a type strain collected from soil in Brazil.

Genomic DNA extraction

Fungi were incubated on Potato Dextrose Agar (Thermo Fischer Scientific, MA, USA) at

30 °C for 7 days to obtain fully mature conidia. The conidia were inoculated in Potato Dextrose

Broth (Thermo Fischer Scientific) at 37 °C at 180 rpm for 16 h. Mycelia were washed with H_2O,

and genomic DNA was purified with phenol-chloroform extraction and Monarch® Genomic DNA

Purification Kit (New England Biolabs, MA, USA) [12, 13]. The DNA concentration and quality

were then measured by Qubit Fluorometer (Thermo Fischer Scientific). The extracted DNA were

used for cloning of *CaM* gene and whole-genome sequencing.

Whole genome sequencing and assembly

Whole-genome 150 bp paired-end (PE) sequencing of four *A. latus* strains, IFM 65030, IFM

65233, IFM 65239 and IFM 66778 was performed using Illumina NovaSeq 6000 (Illumina, CA,

USA) by Novogene Biotech Co. (Beijing, China). The raw genomic reads were trimmed using

fastp v.0.20.1 [14]. The mitochondrial genomes were assembled using GetOrganelle v.1.6.4 [15].

To filter the mitochondrial reads, the reads were aligned with the mitochondrial genome using

BWA-MEM v.0.7.17-r1188 [16]. The mapped reads were filtered using SAMtools v.1.10 [17] and

SeqKit v.0.10.1 [18]. The nuclear genomes were assembled using VelvetOptimiser v.2.2.6 [19].

The completeness of draft genomes was evaluated by BUSCO v. 5.2.1 [20] with the database

eurotiales_odb10. blastn v.2.15.0+ [21] was used for extracting the sequences of *CaM* of *A. latus*

strains by serving the sequence of *A. nidulans* FGSC A4 as query sequence. The genome sizes of

A. latus strains were estimated using Jellyfish [22] and GenomeScope with 21 k-mers [23].

DNA Cloning and Sequencing

The cloning of two *CaM* genes of three strains, IFM 61956, IFM 63852 and IFM 64630, were

performed. PCR amplification was performed by Quick Taq® HS DyeMix (TOYOBO, Tokyo,

Japan) in a ProFlex™ PCR System (Thermo Fisher Scientific) with a program consisting of an

initial denaturing step at 94 °C for 2 min: 35 cycles of denaturation at 94 °C for 30 sec, annealing

at 50 °C for 30 sec, and extension at 68 °C for 1 min. The amplicons of *CaM* were obtained by

PCR with primer pairs CF1L_SacI (5′-AAAGAGCTCGCCGACTCTTTGACYGARGAR-3′)

and CF4_SacI (5′-AAAGAGCTCTTTYTGCATCATRAGYTGGAC-3′). These primers contain

the sequence of CF1 and CF4 primers, respectively, from a previous study [24, 25], with the SacI

restriction site at the 5′ end. The PCR amplicons were purified by gel extraction with QIAquick

Gel Extraction Kit (QIAGEN, Hilden, Germany). The objective gene fragments were cloned into

pMK-dGFP vector [26], and introduced into competent cells of *Escherichia coli* DH5α. FastGene

Plasmid Mini Kit (Nippon Genetics, Tokyo, Japan) was used to extract the plasmid DNA. The

plasmid (ca. 2.0 µL) was used as the templates for the sequencing reaction with a BigDye

Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher Scientific) along with PtrpC-Rev (5′-

AAATGCTCCTTCAATATCATCTTCTGTCGA-3′)   and   M13R   (5′-

CAGGAAACAGCTATGAC-3′) primers [27], following the manufacturer's instructions. Ethanol

precipitation was used to remove excess fluorescent dye-terminators from cycle sequencing

reactions prior to analysis on an automated sequencer. Sequence was performed with an ABI 3100

Genetic Analyzer (Thermo Fisher Scientific).

Phylogenetic analysis

The sequences of *CaM* ranging from 438 bp to 824 bp in length were aligned with MAFFT

v.7.508 [28, 29]. Aligned sequences were trimmed using trimAl v.1.4.rev15 [30], resulting in 686

bp in length. The maximum likelihood tree of *CaM* was constructed using the multithreaded

version of RAxML v.8.2.12 [31], the GTRCAT model, and 1,000 bootstrap replicates. The

phylogenetic tree was visualized using the FigTree (free download available at

http://tree.bio.ed.ac.uk/software/figtree/). A total of 112 taxa were included in our tree

(Supplementary Table S1).

Morphological observations

A. latus IFM 65329 was used for microscopic and macroscopic observation. The colony

character of the other six strains of *A. latus*, *A. sublatus* IFM 42029^{T} (= IFM 4553^{T}) and *A.*

spinulosporus IFM 66771 were observed. The observations of macroscopic characters were

curried on the agar media, i.e., Czapek yeast autolysate agar (CYA, Czapeck Dox Agar, Duchefa

Biochemie, Haarlem, Netherlands and Difco™ Yeast Extract, Thermo Fischer Scientific), oatmeal

agar (OA, Sigma Aldrich, MO, USA) and malt extract agar (MEA, Thermo Fischer Scientific)

with trace elements (0.1 g $ZnSO_4 \cdot 7H_2O$ and 0.5 g $CuSO_4 \cdot 5H_2O$ in 100 mL distilled water) at 25

or 37 °C for 7 days. Microscopic observations of conidia and conidiophores were made from 1

week old colonies on MEA. Ascomata and ascospores were observed from 8-week-old colonies

on OA. A light microscope (Nikon ECLIPSE Ni, Tokyo, Japan and Swift Optical Instruments S7-

TP520, TX, USA) was used to examine morphological characters including the size and shape of

ascomata, ascospores, conidiophores and conidia. These were mounted in a drop of water on glass

slides. The slide preparations were examined and photographed using a digital camera (Nikon

DS-Fi3, Japan). Approximately 30 structures were randomly chosen, and their length and width

were measured using 'ImageJ' software (free download available at http://rsbweb.nih.gov/ij/).

Antifungal susceptibility testing

Minimum inhibitory concentration (MIC) tests for amphotericin B (AMPH-B), itraconazole

(ITCZ), voriconazole (VRCZ), miconazole (MCZ), micafungin (MCFG), and caspofungin

(CPFG), were performed using the dried plate for antifungal susceptibility testing of yeasts

method (Eiken Chemicals, Tokyo, Japan). The method follows the Clinical and Laboratory

Standards Institute M38-E2 with slight modifications [32–34].

Results

Screening and identification of *A. latus*

To identify the presence of allodiploid hybrid strains of *A. latus*, we sequenced the *CaM* genes

of 23 *A. spinulosporus* strains. Sequences with double peaks were observed in seven strains. The

sequences of two copies of *CaM* were determined in all seven strains by either cloning or whole-

genome sequencing (Table 1). These strains were isolated from the patients between 2013 and

2022. The ages of the patients ranged from 13 to 79 years. The patients were diagnosed with

invasive pulmonary aspergillosis (IFM 63852 and IFM 66778) and chronic pulmonary

aspergillosis (IFM 65233 and IFM 65329), the other patients were not diagnosed with

aspergillosis (IFM 61956, IFM 64360, and IFM 65030). However, the patients were diagnosed

with other diseases such as leukemia. Therefore, we could not conclude that *A. latus* is a primary

cause of death.

Draft genomes for four *A. latus* strains (IFM 65030, IFM 65233, IFM 65239 and IFM 66778)

were generated, with assembled genome sizes ranging from 62.1 Mb to 63.2 Mb, aligning with

the typical genome sizes of *A. latus* (previous study showed average genome size 69.09 ± 5.68

Mb [8]) (Supplementary Table S2).

Phylogenetic analysis

We conducted a phylogenetic analysis to characterize the position of seven Japanese *A. latus*

strains. One copy of *CaM* gene sequence clustered with the sequences of *A. spinulosporus*, while

the other copy clustered with *A. sublatus* (Fig. 1). The *CaM* sequence of *A. latus* differed from

that of *A. sublatus* in length by 1 bp, that of *A. quadrilineatus* in length by 6 bp, but the *CaM*

sequence of *A. latus* harbors no specific substitution compared to *A. sublatus*, *A. quadrilineatus*

and *A. spinulosporus.*

Morphological observations

The colony morphologies of *A. latus* varied among strains (Fig. 2, Supplementary Fig. 1). On

MEA, the growth was slower compared to that of *A. sublatus* and *A. spinulosporus* (Fig. 2B,

Supplementary Fig. 1B, E, H, K, N, Q, T, W). In terms of colony appearance, *A. latus* was more

similar to *A. spinulosporus* than to *A. sublatus*, as its colonies were covered with green conidia

on OA and sometimes on CYA (Table 2 and Supplementary Fig. 1C, F, I, L, O, R, X). The

microscopic of *A. latus* also closely resembled those of the other two species but ascospores of *A.*

spinulosporus are echinulated (Table 2) [35, 36].

Antifungal susceptibility testing

Antifungal susceptibility testing was conducted on seven *A. latus*, one *A. sublatus*, and three

A. spinulosporus strains (Table 3). Reduced susceptibility to CPFG was observed in all tested

strains, geometric mean MICs of 7.2 for *A. latus*, 16 for *A. sublatus*, and 5.0 for *A. spinulosporus*.

Reduced susceptibility to AMPH-B was noted in *A. latus* and *A. sublatus*, while *A. spinulosporus*

strains displayed low MICs, with geometric mean MICs of 4.4, 4.0, and 0.7, respectively.

Aspergillus latus strains displayed low MICs to MCFG, whereas MICs of *A. sublatus*, and *A.*

spinulosporus were elevated. Susceptibility to VRCZ and MCZ varied: reduced susceptibility to

VRCZ was observed in *A. latus* IFM 66778 strain (MIC 2.0), and reduced susceptibility to MCZ

was noted in *A. latus* strains IFM 61956 and IFM 65329 (MIC 2.0), as well as *A. sublatus*. All

tested strains were susceptible to ITCZ.

Discussion

Here, we report the allodiploid hybrid *A. latus* strains isolated in Japan. Although 30 strains

have been found worldwide [8, 9], *A. latus* has not been reported in Japan as well as Asia. By

screening 23 *A. spinulosporus* strains, we found seven clinical *A. latus* strains (30.4%). *A. latus*

IFM 61956 was isolated in 2013, suggesting that this species might be more prevalent in the

country. The pathogenicity of *A. latus* was shown to be comparable to that of *A. spinulosporus*

and *A. nidulans* [8, 9]. In this study, the pathogenicity of *A. latus* was unclear, because *A. latus*

strains could be colonized in most cases (Table 1). However, ongoing monitoring is necessary, as

changes in pathogenicity or resistance profiles could impact treatment strategies in the future.

We conducted a phylogenetic analysis of *A. latus* (Fig. 1) with results indicating that it is

derived from a hybridization between *A. spinulosporus* and a species related to *A. sublatus* and *A.*

quadrilineatus. Morphological characters of seven *A. latus* strains, such as colony color and

growth on MEA, were most stable among strains (Fig. 2B, Table 2, and Supplementary Fig. 1B,

E, H, K, N, Q). The differences between *A. latus*, *A. spinulosporus* and *A. sublatus*, however, were

slight, and given the variability observed in previous studies, species identification relying on

these traits may be challenging (Supplementary Fig. 1) [9].

In terms of antifungal susceptibility, strains of *A. latus* found in Japan demonstrated increased

susceptibility to ITCZ compared to overseas strains [8], while showing reduced susceptibility to

CPFG, aligning with the previous research [8, 9] (Table 3). It has been reported that *A.*

spinulosporus exhibits reduced susceptibility to CPFG compared to *A. nidulans* and other

clinically relevant species in section *Nidulantes* [8, 37]. In this study, we confirmed that *A.*

spinulosporus and *A. sublatus* showed reduced susceptibility to CPFG, suggesting that this

characteristic in *A. latus* may be derived from these species. In addition, *A. latus* strains exhibited

AMPH-B resistance. Although high AMPH-B MIC values have been reported in other *Aspergillus*

species, especially *A. nidulans* and related species [8, 37, 38], the mechanisms underlying the

resistance remain poorly understood. Increased catalase production has been implicated in

AMPH-B resistance in *A. terreus* [39], suggesting that *A. latus* may produce high level of catalase.

In conclusion, our study provides significant insights into the ecology and clinical relevance

of *A. latus*, a newly recognized hybrid pathogen and cryptic species of *A. nidulans*. We

documented the first occurrence of this pathogen in Japan, demonstrating its prevalence and

potential role as a causative agent of aspergillosis since 2013. The comprehensive phenotyping,

including morphological and genetic analysis, underscored the diagnostic challenges due to its

similarities with *A. spinulosporus* and *A. sublatus*. Moreover, antifungal susceptibility testing

revealed reduced susceptibility to certain antifungal agents, highlighting the need for accurate

identification to inform effective treatment strategies. Our findings add to the growing body of

knowledge about the distribution and pathogenic characteristics of *A. latus* and emphasize the

importance of continued surveillance and accurate diagnostics in clinical mycology.

Acknowledgements

245  We thank Dr Takayuki Arazoe (Tokyo University of Science) for providing us with the pMK-

246 dGFP plasmid. We thank Junko Ito and Emi Shirai for technical assistance with the experiments.

247 We are also grateful to Machiko Zen for assistance with the data analysis.

249 **References**

250 1. Latgé JP. *Aspergillus fumigatus* and aspergillosis. Clin Microbiol Rev. 1999;12:310–50.

251  https://doi.org/10.1128/CMR.12.2.310

252 2. Sugui JA, Kwon-Chung KJ, Juvvadi PR, Latgé J-P, Steinbach WJ. *Aspergillus fumigatus* and

253  related species. Cold Spring Harb Perspect Med. 2014;5:a019786.

254  https://doi.org/10.1101/cshperspect.a019786

255 3. Bastos RW, Valero C, Silva LP, Schoen T, Drott M, Brauer V, et al. Functional

256  Characterization of Clinical Isolates of the Opportunistic Fungal Pathogen *Aspergillus*

257  *nidulans*. mSphere. 2020;5:e00153-20. https://doi.org/10.1128/mSphere.00153-20

258 4. Paulussen C, Hallsworth JE, Álvarez-Pérez S, Nierman WC, Hamill PG, Blain D, et al.

259  Ecology of aspergillosis: insights into the pathogenic potency of *Aspergillus fumigatus* and

260  some other *Aspergillus* species. Microb Biotechnol. 2017;10:296–322.

261  https://doi.org/10.1111/1751-7915.12367

5.   Zakaria A, Osman M, Dabboussi F, Rafei R, Mallat H, Papon N, et al. Recent trends in the

epidemiology, diagnosis, treatment, and mechanisms of resistance in clinical *Aspergillus*

species: A general review with a special focus on the Middle Eastern and North African

region. J Infect Public Health. 2020;13:1–10. https://doi.org/10.1016/j.jiph.2019.08.007

6.   Hashimoto A, Hagiwara D, Watanabe A, Yahiro M, Yikelamu A, Yaguchi T, et al. Drug

Sensitivity and Resistance Mechanism in *Aspergillus* Section *Nigri* Strains from Japan.

Antimicrob Agents Chemother. 2017;61:e02583-16. https://doi.org/10.1128/AAC.02583-16

7.   Arastehfar A, Carvalho A, Houbraken J, Lombardi L, Garcia-Rubio R, Jenks JD, et al.

Aspergillus fumigatus and aspergillosis: From basics to clinics. Stud Mycol.

2021;100:100115. https://doi.org/10.1016/j.simyco.2021.100115

8.   Steenwyk JL, Lind AL, Ries LNA, Dos Reis TF, Silva LP, Almeida F, et al. Pathogenic

Allodiploid Hybrids of *Aspergillus* Fungi. Curr Biol. 2020;30:2495-2507.e7.

https://doi.org/10.1016/j.cub.2020.04.071

9.   Steenwyk JL, Knowles S, Bastos RW, Balamurugan C, Rinker D, Mead ME, et al.

Evolutionary origin and population diversity of a cryptic hybrid pathogen. Nat Commun.

2024;15:8412. https://doi.org/10.1038/s41467-024-52639-1

10.   Tashiro T. Pulmonary Aspergillosis and clinical significance of *Aspergillus* isolation from

respiratory samples. J Jap Soc Clin Microbiol. 2009;19:67–75.

11. Toyotome T, Saito S, Koshizaki Y, Komatsu R, Matsuzawa T, Yaguchi T. Prospective survey

of *Aspergillus* species isolated from clinical specimens and their antifungal susceptibility: A

five-year single-center study in Japan. J Infect Chemother. 2020;26:321–3.

https://doi.org/10.1016/j.jiac.2019.09.002

12. Bian C, Kusuya Y, Sklenář F, D'hooge E, Yaguchi T, Ban S, et al. Reducing the number of

accepted species in *Aspergillus* series *Nigri*. Stud Mycol. 2022;102:95–132.

https://doi.org/10.3114/sim.2022.102.03

13. Kusuya Y, Bian C, Hagiwara D, Ban S, Takahashi H. A novel Zn2-Cys6 transcription factor

clcA contributes to copper homeostasis in *Aspergillus fumigatus*. Curr Genet. 2022;68:605–

17. https://doi.org/10.1007/s00294-022-01250-2

14. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018;34:i884–90. https://doi.org/10.1093/bioinformatics/bty560

15. Jin J-J, Yu W-B, Yang J-B, Song Y, dePamphilis CW, Yi T-S, et al. GetOrganelle: a fast and

versatile toolkit for accurate de novo assembly of organelle genomes. Genome Biol.

2020;21:241. https://doi.org/10.1186/s13059-020-02154-5

16. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25:1754–60. https://doi.org/10.1093/bioinformatics/btp324

17. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of

SAMtools   and   BCFtools.   Gigascience.   2021;10:giab008.

https://doi.org/10.1093/gigascience/giab008

18. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File

Manipulation. PLoS One. 2016;11:e0163962. https://doi.org/10.1371/journal.pone.0163962

19. Zerbino DR, Birney E. Velvet: algorithms for de novo short read assembly using de Bruijn

graphs. Genome Res. 2008;18:821–9. https://doi.org/10.1101/gr.074492.107

20. Seppey M, Manni M, Zdobnov EM. BUSCO: Assessing Genome Assembly and Annotation

Completeness. Methods Mol Biol. 2019;1962:227–45. https://doi.org/10.1007/978-1-4939-

9173-0_14

21. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture   and   applications.   BMC   Bioinformatics.   2009;10:421.

https://doi.org/10.1186/1471-2105-10-421

22. Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of

occurrences   of   k-mers.   Bioinformatics.   2011;27:764–70.

https://doi.org/10.1093/bioinformatics/btr011

23. Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, et al.

GenomeScope: fast reference-free genome profiling from short reads. Bioinformatics.

2017;33:2202–4. https://doi.org/10.1093/bioinformatics/btx153

24. Peterson SW. Multilocus DNA sequence analysis shows that *Penicillium biourgeianum* is a

distinct species closely related to *P. brevicompactum* and *P. olsonii*. Mycol Res.

2004;108:434–40. https://doi.org/10.1017/S0953756204009761

25. Peterson SW, Vega FE, Posada F, Nagai C. *Penicillium coffeae*, a new endophytic species

isolated from a coffee plant and its phylogenetic relationship to *P. fellutanum*, *P. thiersii* and

P. brocae based on parsimony analysis of multilocus DNA sequences. Mycologia.

2005;97:659-66. https://doi.org/10.3852/mycologia.97.3.659

26. Arazoe T, Ohsato S, Arie T, Yoneyama K, Kuwata S. Construction of a system for exploring

mitotic homologous recombination in the genome of *Pyricularia oryzae*. J Gen Plant Pathol.

2013;79:422–30. https://doi.org/10.1007/s10327-013-0474-2

27. Huey B, Hall J. Hypervariable DNA fingerprinting in *Escherichia coli*: minisatellite probe

from bacteriophage M13. J Bacteriol. 1989;171:2528–32.

https://doi.org/10.1128/jb.171.5.2528-2532.1989

28. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. Nucleic Acids Res. 2002;30:3059–66.

https://doi.org/10.1093/nar/gkf436

29. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol Biol Evol. 2013;30:772–80.

https://doi.org/10.1093/molbev/mst010

30. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment

trimming in large-scale phylogenetic analyses. Bioinformatics. 2009;25:1972–3.

https://doi.org/10.1093/bioinformatics/btp348

31. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics. 2014;30:1312–3.

https://doi.org/10.1093/bioinformatics/btu033

32. Kikuchi K, Watanabe A, Ito J, Oku Y, Wuren T, Taguchi H, et al. Antifungal susceptibility of

Aspergillus fumigatus clinical isolates collected from various areas in Japan. J Infect

Chemother. 2014;20:336–8. https://doi.org/10.1016/j.jiac.2014.01.003

33. CLSI. Reference method for broth dilution antifungal susceptibility testing of filamentous

fungi. In: CLSI standard M38, 3rd edn. Clinical and Laboratory Standards Institute, Wayne,

PA; 2017.

34. He X, Kusuya Y, Hagiwara D, Toyotome T, Arai T, Bian C, et al. Genomic diversity of the

pathogenic fungus *Aspergillus fumigatus* in Japan reveals the complex genomic basis of

azole resistance. Commun Biol. 2024;7:274. https://doi.org/10.1038/s42003-024-05902-6

35. Chen AJ, Frisvad JC, Sun BD, Varga J, Kocsubé S, Dijksterhuis J, et al. *Aspergillus* section

Nidulantes (formerly *Emericella*): Polyphasic taxonomy, chemistry and biology. Stud Mycol.

2016;84:1–118. https://doi.org/10.1016/j.simyco.2016.10.001

36. Horie Y. New or interesting *Emericella* from herbal drugs. Trans Mycol Soc Japan.

1979;20:481–91.

37. Tavakoli M, Rivero-Menendez O, Abastabar M, Hedayati MT, Sabino R, Siopi M, et al.

Genetic diversity and antifungal susceptibility patterns of *Aspergillus nidulans* complex

obtained from clinical and environmental sources. Mycoses. 2020;63:78–88.

https://doi.org/10.1111/myc.13019

38. Van Der Linden JWM, Warris A, Verweij PE. *Aspergillus* species intrinsically resistant to

antifungal agents. Med Mycol. 2011;49 Suppl 1:S82-89.

https://doi.org/10.3109/13693786.2010.499916

39. Blum G, Perkhofer S, Haas H, Schrettl M, Würzner R, Dierich MP, et al. Potential basis for

amphotericin B resistance in *Aspergillus terreus*. Antimicrob Agents Chemother.

2008;52:1553–5. https://doi.org/10.1128/AAC.01280-07

Statements & Declarations

Funding

This work was partly supported by JSPS KAKENHI Grant No. 24K19262.

Competing Interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

Saho Shibata, Vit Hubka, and Hiroki Takahashi contributed to the study conception and design.

Material preparation, data collection and analysis were performed by Saho Shibata, Momotaka

Uchida, Sayaka Ban, Kamei Katsuhiko, Akira Watanabe, Vit Hubka, and Hiroki Takahashi. The

draft of the manuscript was written by Saho Shibata, Momotaka Uchida, Takashi Yaguchi, Vit

Hubka and Hiroki Takahashi, and all authors commented on the manuscript. All authors read and

approved the final manuscript.

Ethical approval

No ethical approval is required

383 **Table 1.** Clinical information of *Aspergillus latus* strains examined in this study

| Strain No. | Species name | | Isolation year | Disease record | Age | Outcome |
	this study	previous identified				
IFM 61956	*A. latus*	*A. spinulosporus*	2013	colon cancer, pneumonia	N/A	dead
IFM 63852	*A. latus*	*A. spinulosporus*	2016	interstitial lung disease, invasive pulmonary aspergillosis	79	dead
IFM 64360	*A. latus*	*A. spinulosporus*	2016	ulcerative colitis, pneumonia	45	dead
IFM 65030	*A. latus*	*A. spinulosporus*	2017	bronchiolitis	62	alive
IFM 65233	*A. latus*	*A. spinulosporus*	2018	liver cirrhosis, chronic pulmonary aspergillosis	66	alive
IFM 65329	*A. latus*	*A. spinulosporus*	2015	pulmonary non-tuberculous mycobacteriosis, chronic pulmonary aspergillosis	68	alive
IFM 66778	*A. latus*	*A. spinulosporus*	2020	acute myeloid leukemia, invasive pulmonary aspergillosis	13	alive

386   **Table 2.** Morphological comparison of *Aspergillus latus*, *A. spinulosporus* and *A. sublatus*.

| | | A. latus | A. spinulosporus | A. sublatus |
		This study	This study & [35]	This study & [36]
Colony	CYA	Obverse: yellow green, olive green, pale brown edge	Obverse: pale brown, yellow edge, grey, green conidia forming all over	Obverse: reddish brown, grey, pale brown edge, reddish conidia forming in the center
		Reverse: dark brown, yellowish brown, green conidia formed in the center	Reverse: pale brow, brown,	Reverse: pale brown, orange, brown edge
	MEA	Obverse: pale yellow, irregular edge with white mycelia	Obverse: concentrically growth, pale-yellow	Obverse: concentrically growth, pale-yellow, white edge, olive-green conidia formed in the center
		Reverse: pale-yellow, soluble yellowish pigments present and exudates absent	Reverse: pale yellow, soluble pigments and exudates absent	Reverse: pale yellow, no soluble pigments and exudates
	OA	Obverse: pale brown, green, partially reddish, covered in green conidia, the center pigmented yellow to brown	Obverse: pale brown with olive green conidia and sometime wrinkled	Obverse: brown with pale brown edge
		Reverse: soluble orange or brown pigments present and exudates absent	Reverse: orange to brown, brown soluble pigments present and exudates absent	Reverse: orange to brown, brown soluble pigments present and exudates absent

Ascoma ta	character	cleistothecial, superficial, black, globose, subglobose, surrounded by Hülle cells	cleistothecial, superficial, dark brown, globose, subglobose, surrounded by Hülle cells	cleistothecial, superficial, red to purple, globose, subglobose
Hülle cells	size (μm)	8.7–20 (diameter)	14–30 (diameter)	12.5–30× 10–28
	character	hyaline, globose to ovoid	hyaline, globose to ovoid	hyaline, globose to ovoid
Ascospo res	size (μm)	3.5–10 (pleated equatorial crests: 0.6–2.3)	3.5–4.5 × 3.0–4.5 (pleated equatorial crests: 0.8–1.0)	4.0–4.5 × 3.0–3.5 (pleated equatorial crests: 1.0–1.5)
	character	purple red, reddish brown, in surface view globose, subglobose, spore bodies smooth, incompletely reticulate, ribbed, globose, subglobose, in side view lenticular, with two pleated equatorial crests	orange,reddish brown, in surface view globose, subglobose, spore bodies echinulate, globose, subglobose, in side view lenticular	purple red, reddish brown, globose, subglobose, spore bodies smooth, in side view lenticular
Conidio phores	size (μm)	50–179 × 2.0–3.6	70–120 × 5.0–6.0	200× 4.0–6.0
	character	smooth stipes, pale brown	smooth stipes, yellowish brown	pale grey, grayish yellow
Vesicles	size (μm)	4.8–9.0 (wide)	9.0–11 (wide)	9.0–12.5 (wide)
	character	pale brown, subglobose to subclavate	yellowish brown, subclavate, fertile over the upper half	hemispherical, flask-shaped
Metulae	size (μm)	5.1–8.3 × 2.0–4.0	6.0–8.0 × 3.0–4.0	4.0–6.0 × 3.0–4.0
	character	hyaline to pale green	pale brown to pale green	hyaline to pale yellowish brown
Phialide s	size (μm)	5.2–8.4 × 2.0–4.0	6.0–8.5 × 2.0–3.0	6.0–8.0 × 2.0–3.0
	character	hyaline, flask-shaped	hyaline to pale green, flask-shaped	hyaline to pale yellowish brown

| Conidia | size (μm) | 2.7–4.8 | 3.0–4.0 | 2.5–3.5(4.0) |
| | character | echinulate, globose to subglobose, green in mass | echinulate, globose to subglobose, green in mass | echinulate, globose to subglobose, green to pale olive green |

388   **Table 3.** MIC values for antifungal drugs

Species name	Strain No.	MIC (µg/mL)					
		MCFG	CPFG	AMPH-B	ITCZ	VRCZ	MCZ
Aspergillus latus	IFM 61956	0.06	16	4	0.5	0.12	2
	IFM 63852	0.12	16	4	0.25	0.12	1
	IFM 64360	0.06	16	4	0.25	0.25	1
	IFM 65030	0.06	4	4	0.25	0.06	1
	IFM 65233	0.06	2	4	0.5	0.06	1
	IFM 65329	0.03	8	4	0.25	0.12	2
	IFM 66778	0.06	4	8	0.25	2	0.5
A. sublatus	IFM 42029$^{\text{T}}$	>16	16	4	0.25	0.12	2
A. spinulosporus	IFM 66771	0.15>	2	1	0.25	0.25	0.5
	IFM 61449	>16	8	0.5	0.25	0.12	0.5
	IFM 64750	>16	8	1	0.25	0.12	1

MCFG = micafungin, CPFG = caspofungin, AMPH-B = amphotericin B, ITCZ = itraconazole, VRCZ = voriconazole,

MCZ = miconazole

 Figure legends

Fig. 1. Phylogenetic tree constructed using maximum likelihood (ML) analysis based on the

models selected with the GTRCAT for *CaM* (Bootstrap value: > 70%). *A. aurantiobrunneus* and

A. fructus were used as outgroup. Numbers after taxa are isolate, specimen or strain numbers. The

scale bar represents the number of nucleotide substitutions per site. The sequences obtained in

this study are shown in bold. T = type sequence.

Fig. 2. *A. latus* IFM 65329 colonies: obverse CYA 37 °C (A), MEA 37 °C (B), OA 25 °C (C),

reverse: CYA 37 °C (D), MEA 37 °C (E), OA 25 °C, 7d (F), conidiophores (G–J), conidia (K),

Hülle cells (L), ascomata (M), ascospores (N, O) Scale bars: G–L, N, O 10 μm.

Shibata et al. Figure 1