1 *De novo* assembled databases enable species-specific protein-based

2 stable isotope probing of microbiomes without prior knowledge of

3 the community composition

4 Simon Klaes[1,2*], Christian White[1,3], Lisa Alvarez-Cohen[3], Lorenz Adrian[1,2], Chang Ding[1]

5 [1]Molecular Environmental Biotechnology, Helmholtz Centre for Environmental Research – UFZ,
6 Leipzig, Germany
7 [2]Geobiotechnology, Technische Universität Berlin, Berlin, Germany
8 [3]Civil and Environmental Engineering, University of California, Berkeley, California, United States
9
10
11 Running title: denovo-SIP
12
13 *To whom correspondence should be addressed: Simon Klaes (simon.klaes(at)ufz.de)

15 Keywords: metaproteomics, *de novo* peptide sequencing, peptide databases, microbiome analysis,
16 microbial communities, mass spectrometry, database assembly, isotopic labelling, microbial activity
17 profiling, database quality optimization

Abstract

Background: Protein-based stable isotope probing (Protein-SIP) is a powerful approach that can directly link individual taxa to activity and substrate assimilation, elucidating metabolic pathways and trophic relationships within microbiomes. In Protein-SIP, peptides and corresponding taxa are identified by database matches. Thus, database quality is crucial for accurate Protein-SIP analyses. For samples with unknown community composition, Protein-SIP usually employs broad-spectrum or metagenome-derived databases. However, broad-spectrum databases require advanced post-processing strategies and metagenome-derived databases can be costly to acquire. Further, both can inflate database size, negatively impacting peptide identification.

Results: Here, an approach in which we use *de novo* assembled databases for Protein-SIP on microbiomes with unknown community compositions is introduced. This approach enables databases to be directly generated from the mass spectrometry raw data using *de novo* peptide sequencing. We then use the mass spectrometric data from labeled cultures to quantify isotope incorporation into specific peptides. We benchmark our approach against the canonical approach in which a sample-matching database is generated by DNA sequencing on three different datasets: 1) a proteome analysis from a defined microbial community containing ^{13}C-labeled *E. coli* cells, 2) time-course data of an anammox-dominated continuous reactor after feeding with ^{13}C-labeled bicarbonate, and 3) a model of the human distal gut simulating a high-protein and high-fiber diet cultivated in either ^2H_2O or $H_2^{18}O$. Our results show that *de novo* database assembly is applicable to different isotopes and detects similar amounts of labeled peptides compared to sample-matching databases. Furthermore, we show that peptide-centric Protein-SIP allows species-specific resolution, enabling the assessment of activity related to individual biological processes. Finally, we provide access to our modular Python pipeline to assist the *de novo* assembly of databases and subsequent peptide-centric Protein-SIP data analysis (https://git.ufz.de/meb/denovo-sip).

Conclusions: *De novo* assembled databases enable species-specific Protein-SIP of microbiomes without prior knowledge of the community composition.

Introduction

Microbiomes are microbial communities in a defined habitat with distinct physio-chemical properties [1]. As microbiomes are ubiquitous and crucial for the functioning and health of myriad ecosystems, microbiome studies are essential for understanding fundamental ecological processes and advancing knowledge of human health and the environment [2–4]. Microbiomes are frequently studied via omics approaches [5], which can provide insights into their state and dynamics [6]. However, the complexity of microbiomes often hinders attempts to link ecosystem functions to individual taxa. To establish direct links between substrates, metabolites, functions, and taxa, omics approaches can be combined with stable isotope probing (SIP) [7].

In SIP experiments, microbial communities are exposed to a substrate highly enriched in a stable isotope. Upon assimilation, microorganisms can incorporate the stable isotopes into biomolecules, such as DNA [8, 9], RNA [9, 10], phospholipid-derived fatty acids [11, 12], and proteins [13, 14]. Analyzing the relative amount of labeled biomolecules and their degree of isotope incorporation can establish a direct link of individual taxa to general (for ^{2}H or ^{18}O labeled water) or substrate-specific (for ^{13}C, ^{15}N, or ^{34}S labeled substrate) activity [8-19]. By establishing this link, SIP can identify key microorganisms [20, 21], elucidate metabolic pathways [22, 23], and unravel complex food webs in microbiomes [9, 24]. The choice of the investigated biomolecule affects the sensitivity and taxonomic resolution of SIP analyses [as reviewed in 7]. DNA and RNA-based SIP provide high taxonomic resolution but rely on the separation of heavy and light fractions via ultracentrifugation leading to low sensitivity and inaccuracy in isotope quantification. Fatty acid and protein-based SIP (Protein-SIP) employ mass spectrometry to accurately quantify isotope incorporation. While fatty acids only provide limited taxonomic information, proteins can characterize communities down to the strain level. Thus, Protein-SIP offers the advantage of accurately quantifying isotope incorporation while preserving high taxonomic resolution.

Protein-SIP is based on bottom-up metaproteomics. In brief, proteins are extracted from the sample and digested into peptides that are then analyzed by high-resolution tandem mass spectrometry (MS/MS). Obtained spectra are mapped back to peptides and relative isotope abundances (RIA) are calculated from shifts in precursor ion peaks [25–27]. In principle, spectra can be mapped to peptides by database or spectral library searches as well as *de novo* peptide sequencing. To date, only database searches are used for Protein-SIP. In database searches, acquired mass spectra are compared to theoretical spectra of peptides in a constructed target database to obtain peptide-spectrum matches (PSMs) that are then validated by a target-decoy approach. While database-dependent approaches are widely accepted for proteomics and Protein-SIP, they are heavily reliant on the quality of the database leading to several constraints. One key constraint is that database creation ideally requires prior knowledge of the community composition. For defined cultures, such as pure strains, sample-matching databases are readily available in public repositories. However, for samples of unknown composition, e.g., environmental samples, databases usually must be derived from metagenome sequencing, which can be a time-consuming and costly step. In addition, metagenome-derived databases often cover only a fraction of the protein-encoding genes in complex microbiomes, so that

uncovered proteins cannot be identified [28–30]. Alternatively, broad-spectrum databases such as
NCBI non-redundant proteins (nr) or Uniprot can be used [13, 31–34]. However, broad-spectrum
databases increase computational time substantially, require advanced filtering strategies, and often
lead to lower taxonomic resolution [35–38]. Subsets of broad-spectrum databases focusing on
taxonomic marker proteins were successfully applied to Protein-SIP, reducing computational time, but
also limiting the taxonomic resolution [39]. Moreover, metagenome-derived and broad-spectrum
databases contain sequences of proteins that are not expressed in the sample, inflating database size.
Inflated databases can negatively impact peptide identification during target-decoy validation
strategies [36, 40]. Furthermore, database-dependent search approaches are inherently biased
towards the database content making it impossible to detect proteins missing in the database [37].

De novo peptide sequencing addresses the above-mentioned limitations of database-dependent
searches by directly inferring peptides from MS/MS spectra without using references. Thereby, *de*
novo peptide sequencing does not provide a link of peptides to proteins like database-dependent
approaches, making it a peptide-centric approach, in which taxonomic and functional information of
peptides can be inferred by text matching (Unipept [41]) or sequence similarity (BLASTP [42]). To date,
de novo sequencing has been used to characterize microbial community compositions at the family
level [43] and to assess database suitability [44]. Until recently *de novo* sequencing identified many
fewer peptides than database-dependent approaches [45]. However, current advances in artificial
intelligence and the increased amount of publicly available training data have massively improved *de*
novo sequencing algorithms, challenging database-dependent approaches [46, 47].

Despite the advances in *de novo* sequencing, it remains unclear whether *de novo* algorithms can
reliably identify peptides in partially labeled samples, where isotope peaks may cause signal
interference. Moreover, *de novo* sequencing alone cannot identify labeled peptides or quantify isotope
incorporation, necessitating their integration with SIP tools. To address this knowledge gap, we
propose the creation of sample-matching databases derived from *de novo* sequencing of unlabeled
peptides, enabling Protein-SIP analysis without prior knowledge of community compositions
(Figure 1).

Figure 1: Proposed workflow for protein-based stable isotope probing (Protein-SIP) using *de novo* assembled peptide databases.

Here, we investigate whether *de novo* algorithms can accurately identify peptides in partially labeled samples and assess whether these identified peptides can be used to assemble sample-matching databases for Protein-SIP analysis using ^{2}H, ^{13}C, and ^{18}O labeled data. Additionally, we benchmark databases assembled with different state-of-the-art *de novo* sequencing tools with each other and with sample-matching reference databases.

Results

De novo sequencing of peptides in partially labeled samples

To investigate (1) if *de novo* algorithms can identify target peptides in partially labeled samples and (2) how the RIA affects *de novo* peptide sequencing, we analyzed previously published proteomic data (PRIDE accession number: PXD04141) of standard *E. coli* cultures cultivated with a defined ^{13}C RIA of 1.07–99% [27]. Specifically, we performed *de novo* peptide sequencing using the state-of-the-art engines PepNet and Casanovo and compared the number of *de novo* identified target and non-target peptides across the reported quality scores with a conventional database search using MS-GF+ and MetaProSIP (Figure 2).

Figure 2: Comparison of the non-redundant number and quality score of peptides identified by *de novo* sequencing in standard *Escherichia coli* K12 cultures across biological triplicates cultivated with a defined ¹³C relative isotope abundance (RIA) of 1.07–99%. *De novo* peptide sequencing was performed using Casanovo and PepNet. Negative quality scores are plotted as zero. Solid lines represent identified peptides matching the proteome of *E. coli* K12 + universal contaminants. Dotted lines indicate identified peptides not matching the proteome of *E. coli* K12 + universal contaminants. Dashed lines are only plotted in the bottom graphs and represent the number of identifications using the sample-matching reference database with MS-GF+ and MetaProSIP.

Casanovo and PepNet successfully identified 6,956–15,824 unique target peptides in samples containing a ¹³C RIA of 1.07–5% ¹³C. An increase in the ¹³C RIA resulted in a reduced number of peptide

identifications. In samples with a ^{13}C RIA of 25–99%, both *de novo* algorithms as well as conventional database search failed to detect a relevant number of target peptides. However, this limitation is not critical, as SIP tools usually require an unlabeled reference for detecting labeled peptides. These unlabeled references can then be used for de novo sequencing to create the peptide database.

Casanovo outperformed PepNet by identifying more target peptides and fewer non-target peptides. Moreover, applying higher quality score thresholds led to a pronounced decrease in non-target peptide identifications but only a marginal decrease in target peptide identifications. This indicates effective separation of true and false-positive identifications by Casanovo at high quality scores. Compared to conventional database searches, Casanovo identified 103.1–157.7% of the number of unique target peptides.

Benchmarking *de novo* assembled databases with a defined microbial community

To test (1) if *de novo* algorithms can identify peptides from partially labeled samples, and (2) whether identified peptides can be an adequate database for identifying labeled peptides, we analyzed previously published proteomic data (PRIDE accession number: PXD024174) from a defined microbial community containing unlabeled and ^{13}C-labeled *E. coli* cells in a 1:1 ratio [48]. The community consisted of 21 bacterial species belonging to 15 families, 1 eukaryote, 1 archaeon and 5 bacterial viruses (Supplementary Table 1 in [49]). The labeled *E. coli* cells were cultivated with 10% fully labeled (100% ^{13}C) and 90% unlabeled (1.1% ^{13}C) glucose. We performed *de novo* peptide sequencing with PepNet and Casanovo. Subsequently, peptides were filtered applying different defined score thresholds to assemble databases of 14,700 to 875,366 peptides (Supplementary Table S1) which were then used to identify labeled peptides with a MetaProSIP workflow (Figure 3).

Figure 3: Comparison of *de novo* assembled databases for identifying unlabeled and [13]C-labeled peptides in a mock microbial community spiked with labeled *Escherichia coli* K12 cells obtained from [48] using a MetaProSIP workflow. *De novo* databases were assembled using Casanovo or PepNet and identifications were filtered with different thresholds for the quality score. The color represents whether the identified peptide is present in the proteome of *E. coli* K12 + universal contaminants (navy circle), of the residual mock community (purple rectangles), or in none of them (grey crosses). Dashed lines represent the number of identifications using the sample-matching reference database. Only peptides detected in at least two biological replicates are plotted.

We identified more peptides from the defined microbial community and fewer unrelated peptides using Casanovo-derived databases compared to those from PepNet. Casanovo identified 14,700 peptides with a score greater than 0.99, and a database built from these peptides (CS>0.99) yielded the highest number of labeled *E. coli* K12 peptides among the *de novo* assembled databases. Specifically, using the CS>0.99 database in our MetaProSIP workflow, we identified 8,496 peptides, including 505 labeled *E. coli* K12 peptides, just 20 fewer than with the reference database. Thus, the CS>0.99 database was used for further analyses. Using this database, we investigated the level of

taxonomic resolution that could be identified for the labeled organisms within the unlabeled residual community. For this purpose, we inferred the taxonomy of all identified peptides with a lowest-common-ancestor approach using Unipept and plotted the peptides with their assigned family against the calculated ^{13}C RIA (Figure 4A).

Figure 4: Analysis of ^{13}C relative isotope incorporation (RIA) into peptides of different taxa identified in a mock microbial community with spike-in of ^{13}C-labeled *E. coli* cells. RIA values are shown for peptides detected in at least two replicates and for taxa for which at least three peptides were detected. Dots indicate peptides assigned to all detected families (**A**) or *Enterobacteriaceae* spp. (**B**). 'n' denotes the number of peptides identified for a taxon across all replicates. Solid lines indicate medians. Dotted lines indicate means. Statistically significant differences in RIA between species are marked with '**' based on Student's t-test for the means of two independent samples with p < 0.01. **C**: Comparison of drop-off curves from *in silico* digestion of the *E.coli* K12 proteome and labeled peptides detected in at least two replicates of the mock microbial community with spike-in of ^{13}C-labeled *E. coli* cells. **D**: Number of best hits to *Salmonella enterica* and *E. coli* when querying by BLASTP the 31 labeled peptides not assigned to any lowest common ancestor by Unipept and detected in at least two replicates of the mock microbial community with spike-in of ^{13}C-labeled *E. coli* cells. Sequence similarity search was performed using BLASTP against the nr database. Organisms with only one hit and hits with e-value ≥ 0.001 were removed before plotting.

Our MetaProSIP workflow identified 11,497 non-redundant peptides out of which 8,449 peptides were consistently detected in at least two replicates. From these 8,449 peptides, 7,897 were assigned to a lowest common ancestor using Unipept. Additionally, 450 of the 8,449 peptides were labeled in at least two sample out of which 419 were assigned to the lowest common ancestor via Unipept.

Overall, we identified 13 out of 15 bacterial families, as well as the eukaryotic and archaeal family in the mock community. We did not detect any bacterial viruses. However, we consistently detected 6 peptides that were assigned to the fungal family *Trichomeriaceae* that was not expected in the mock community. It is unclear whether these peptides are false-positives or originate from a contamination. Most of the identified peptides cluster around a RIA of 1.1%, corresponding to the natural ^{13}C-abundance indicative of unlabeled peptides. Only for *Enterobacteriaceae* we observed a second cluster of peptides centered between a RIA of approximately 7% and 11%. *Salmonella enterica* and *E. coli* were the only members of *Enterobacteriaceae* identified at higher taxonomic resolution, whereby peptides assigned to *E. coli* exhibited a median RIA of 8.5%, which is significantly higher (p < 0.001) than that found for peptides from *S. enterica,* indicating that only *E. coli* peptides were labeled (Figure 4B).

To test for hidden labeled populations, we applied three approaches. First, we compared the number of labeled peptides assigned to *E coli* or superordinate taxa of its lineage with the number of labeled peptides assigned to other taxa. Our results showed that most labeled peptides were assigned to E. coli or superordinate taxa of its lineage, indicating a high population share of E. coli in the labeled subpopulation of the sample (Supplementary Table S2). Secondly, we compared the theoretical drop-off rate of *E. coli* K12 peptides with the observed drop-off rate of detected labeled peptides. Drop-off rates describe the decline in taxon-specific peptide counts from higher to lower taxonomic levels [43]. By comparing observed and theoretical drop-off rates, it is possible to assess whether peptide counts at a given taxonomic level can account for counts at higher levels [43]. Consequently, drop-off rates can reveal hidden subpopulations not captured by the taxonomic database. Here, the observed drop-off rate deviated only slightly from the theoretical one, indicating the absence of hidden side populations (Figure 4C). Thirdly, we performed a sequence similarity search for the 31 labeled peptides

that Unipept could not assign to any lowest common ancestor (Figure 4D). For 11of those peptides,
BLASTP found highly similar sequences with an e-value < 0.001. After removing organisms with only
one hit, all highly similar sequences were related to *E. coli* or its close relative *S. enterica* [50].
Consequently, all performed tests indicate the absence of hidden labeled subpopulations in
accordance with the setup of the defined microbial community.

Case study 1: Investigating trophic relations in a continuous flow reactor for the cultivation of
anammox bacteria by tracing ^{13}C incorporation into peptides

To test the power and applicability of our approach on real samples, we analyzed proteomics data
from a continuous flow anammox reactor enriched with '*Candidatus* Kuenenia stuttgartiensis' strain
CSTR1 at three different time points after changing the carbon source from non-labeled to ^{13}C-labeled
bicarbonate while leaving nitrogen sources unchanged.

Again, it was possible to detect and subsequently quantify the RIA of peptides using *de novo* assembled
databases. The assembled databases had a size of 5,313 to 303,595 non-redundant peptides
(Supplementary Table S1). For samples acquired on day 21 (5.25 × hydraulic retention time), it was
necessary to mix labeled peptides with unlabeled peptides from day 0 to enable *de novo* peptide
sequencing and subsequent RIA calculation by MetaProSIP (Supplementary Table S3). Across all
samples, we detected more labeled '*Ca.* Kuenenia stuttgartiensis' strain CSTR1 peptides and less non-
CSTR1 labeled peptides with the Casanovo-derived databases compared to the PepNet-derived ones
(Supplementary Figure S1). Similarly, we compared different quality score thresholds for filtering *de*
novo identified peptides before creating the database for SIP analysis (Figure 5).

Figure 5: Comparison of *de novo* assembled databases for identifying unlabeled and [13]C-labeled peptides in an

enrichment of '*Candidatus* Kuenenia stuttgartiensis' strain CSTR1 from a continuous flow reactor fed with [13]C-

bicarbonate using a MetaProSIP workflow. *De novo* databases were assembled using Casanovo and filtering

identifications with different thresholds for the quality score. Navy circles denote the number of identified

peptides present in the proteome of '*Ca.* Kuenenia stuttgartiensis' + contaminants. Grey crosses denote the

number of identified peptides absent in the proteome of '*Ca.* Kuenenia stuttgartiensis' + contaminants. Dashed

lines represent the number of identifications using the sample-matching reference database derived from

metagenome sequencing [51]. Only peptides detected in at least two biological replicates are plotted.

By applying a score threshold of 0.99 (CS>0.99), we identified 1372 labeled '*Ca.* Kuenenia

stuttgartiensis' strain CSTR1 peptides, surpassing all other thresholds investigated. Additionally, the *de*

novo assembled database demonstrated comparable performance to a genome-based one yielding

1459 labeled '*Ca.* Kuenenia stuttgartiensis' strain CSTR1 peptides. To investigate the [13]C incorporation

into taxa over time, we used the CS>0.99 database for the MetaProSIP workflow. Then, we analyzed
the RIA and taxonomy of detected peptides (Figure 6).

Figure 6: Distribution of the ^{13}C relative isotope abundance (RIA) in detected peptides across taxa identified in samples from a continuous flow reactor enriching '*Candidatus* Kuenenia stuttgartiensis' strain CSTR1 at different time points after switching to medium containing ^{13}C-bicarbonate. Dots indicate peptides assigned to detected phyla (**A**) or species (**B**). RIA values are shown for peptides detected in at least two replicates at each time point. Taxa are shown for which at least three peptides were detected. 'n' denotes the number of peptides assigned to a taxon across all replicates and time points. Solid lines indicate medians. Dotted lines indicate means. Statistically significant differences in RIA between time points are indicated with '***' based on Student's t-test for the means of two independent samples with $p < 0.001$. **C**: Comparison of drop-off curves from *in silico* digestion of the '*Candidatus* Kuenenia stuttgartiensis' strain CSTR1 proteome and labeled peptides detected in at least two samples of the continuous flow reactor enriching '*Ca*. Kuenenia stuttgartiensis' strain CSTR1. **D**: Top-hit species distribution after sequence similarity search of the 59 labeled peptides not assigned to any lowest common ancestor by Unipept but detected in at least two samples of the continuous flow reactor enriching '*Candidatus* Kuenenia stuttgartiensis' strain CSTR1. The sequence similarity search was performed with BLASTP against the nr database. Organisms with only one hit and hits with e-value ≥ 0.001 were removed before plotting.

The MetaProSIP workflow identified 4,440 non-redundant peptides out of which 1,947 peptides were consistently detected in at least two replicates of each time point. From these 1,947 peptides, 1,846 were successfully assigned to a lowest common ancestor using Unipept. Additionally, 1,232 of the 1,947 peptides were labeled in at least two sample out of which 1,173 were assigned to lowest common ancestor via Unipept.

At the phylum level, most detected peptides were assigned to *Plancomycetota*, consistent with their high abundance in the reactor (Figure 6A). Some peptides were assigned to the phylum *Chordata*, indicating human contamination, and a few were linked to *Pseudomonadota*, previously described as a subpopulation of the reactor [52]. At the species level, we only detected '*Ca.* Kuenenia stuttgartiensis' belonging to the phylum *Planctomycetota,* suggesting high enrichment of the culture (Figure 6B). Substantial incorporation of ^{13}C has been detected only for peptides assigned to '*Ca.* Kuenenia stuttgartiensis' or superordinate taxa of its lineage, indicating its activity and assimilation of ^{13}C-bicarbonate (Supplementary Table S4). Specifically, we observed a significant ($p < 0.001$) increase of ^{13}C RIA in '*Ca.* Kuenenia stuttgartiensis' peptides at each investigated time point, resulting in a median RIA of 95.7% after 5.25 × hydraulic retention time demonstrating almost complete substitution of ^{12}C with ^{13}C.

To investigate for labeled populations not covered by Unipept, we again compared observed and theoretical drop-off curves of peptides (Figure 6C). The drop-off curve of detected labeled peptides closely resembled the theoretical one, with a maximum deviation of only 4.9% (Supplementary Table S4). This suggests that all detected labeled peptides were derived from '*Ca.* Kuenenia stuttgartiensis'.

Furthermore, we searched for sequences similar to the 59 labeled peptides that Unipept could not assign to any taxon (Figure 6D). For 15 of those peptides, BLASTP identified highly similar sequences with e-value < 0.001. After removing organisms with only one hit, most of the highly similar sequences belonged to '*Ca.* Kuenenia stuttgartiensis' or superordinate taxa of its lineage. Three hits with high sequence similarity were affiliated with '*Candidatus* Jettenia sp.' and '*Candidatus* Jettenia sp. AMX1',

respectively, which are anammox bacteria. In brief, our results indicate that '*Ca.* Kuenenia
stuttgartiensis' also dominates the labeled population in the reactor.

Case study 2: Investigating species-specific and function-specific activity between two different
simulated diets in a human distal gut microbiome model by tracing ^{2}H and ^{18}O incorporation
into peptides

To test our strategy for comparative purposes and also for stable isotopes other than ^{13}C, we analyzed
previously published raw data (PRIDE accession number: PXD024291) from a model community of the
human gut microbiome ('Robogut community') cultivated for 12 h in a protein-rich or fiber-rich
medium simulating different diets [18, 48]. The medium contained unlabeled water (for controls) or
water in which 25% was substituted with either ^{2}H$_2$O or H$_2$^{18}O. First, we compared different *de novo*
assembled databases for identifying unlabeled and labeled peptides expected to be present in the
sample based on the published reference database (Figure 7).

Figure 7: Comparison of *de novo* assembled databases for identifying unlabeled and $^{18}O/^{2}H$-labeled peptides in a model of the human gut microbiome (Robogut community) cultivated with $H_2^{18}O$ or $^{2}H_2O$, respectively, and obtained from [18] using a MetaProSIP workflow. *De novo* databases were assembled using Casanovo after filtering all identifications with defined thresholds for the quality score. Navy circles denote the number of identified peptides present in the sample-matching reference proteome + contaminants. Grey crosses denote the number of identified peptides absent in the sample-matching reference proteome + contaminants. Dashed lines represent the number of identifications that we generated using the published sample-matching reference database. Only peptides detected in at least two biological replicates are plotted.

The *de novo* assembled databases contained 48,715 to 715,031 non-redundant peptides (Supplementary Table S1). We identified more Robogut peptides with databases created using Casanovo compared to those created using PepNet, for both investigated stable isotopes ^{18}O and ^{2}H (Supplementary Figure S2). The database for which peptides were *de novo* sequenced by Casanovo and filtered with a score threshold of 0.99 (CS>0.99) led to the most identifications of labeled Robogut

peptides among all investigated databases. Furthermore, we identified more labeled peptides in samples cultivated with ^{18}O than with ^{2}H and more unlabeled peptides in the ^{2}H samples.

Next, we analyzed the RIA and taxonomy of peptides identified under different dietary conditions using the CS>0.99 database. Our MetaProSIP workflow identified 35,115 non-redundant peptides, out of which 8,099 and 8,999 peptides were detected in at least two replicates of both media compositions in presence of ^{18}O and ^{2}H, respectively. From these 8,099 and 8,999 peptides, 7,669 and 8,430 were assigned to a lowest common ancestor using Unipept, respectively.

Overall, the incorporation of ^{18}O was substantially higher than the incorporation of ^{2}H. Moreover, the identified peptides generally had significantly higher RIA (p < 0.01) in protein-rich compared to fiber-rich conditions (Supplementary Figure S3).

Figure 8: Distribution of the ^{18}O and ^{2}H relative isotope abundance (RIA) in detected peptides across species identified in a model of the human gut microbiome cultivated in either a high fiber or high protein media in the presence of either $H_2{}^{18}O$ or $^{2}H_2O$. RIA values are shown for peptides detected in at least two replicates of both media compositions. Species detected in the presence of $H_2{}^{18}O$ and $^{2}H_2O$ are plotted for which at least three peptides were detected. 'n' denotes the number of peptides assigned to a specific taxon across triplicates. Statistically significant differences in RIA between media are indicated with '*', '**', and '***' based on Student's t-test for the means of two independent samples with p < 0.05, p < 0.01, and p < 0.001, respectively. To improve readability, outliers were removed from the raw graph (Supplementary Figure S4).

After inferring the taxonomy of peptides via Unipept, we identified 22 species present in all tested setups (Figure 8). Notably, we detected *Copranaerobaculum intestinale* and *Hominenteromicrobium mulieris*, which were recently described [53, 54] but not reported by previous analyses of this dataset due to their absence in the originally used database [18, 48]. In ^{18}O incubated samples, more taxa were found to differ significantly in their RIA between the two simulated diets than ^{2}H incubated samples.

Specifically, we detected significantly higher RIA under high protein conditions compared to high fiber
conditions for peptides assigned to *Akkermansia muciniphilia*, *Bacteroides ovatus*, *Bacteroides*
uniformis, *Clostridium hylemonae*, *Clostridium symbiosum*, and *Copraanaerobaculum intestinale*, as
well as significantly lower RIA for *Hominenteromicrobium mulieris* and *Hungatella hathewayi*.

To assess changes in activity related to individual biological processes, we assigned identified peptides
and their RIA to biological processes via gene ontology (GO) numbers using Unipept. For an overview,
GO numbers were first merged into GO slims of the prokaryote subset (Figure 9).

Figure 9: Distribution of the ^{18}O and ^{2}H relative isotope abundance (RIA) in detected peptides across general
biological processes of prokaryotes (goslim_prokaryote) identified in a model of the human distal gut
microbiome cultivated in either a high fiber or high protein medium in the presence of either $H_2^{18}O$ or $^{2}H_2O$. RIA
values are shown for peptides detected in at least two replicates of both media and for biological processes
detected in presence of $H_2^{18}O$ and $^{2}H_2O$ for which at least three peptides were detected (10 to 3457 peptides per
strip). Statistically significant differences in RIA between media are indicated with '*', '**', and '***' based on
Student's t-test for the means of two independent samples with p < 0.05, p < 0.01, and p < 0.001, respectively.
To facilitate readability, outliers were removed from the raw graph (Supplementary Figure S5).

For most biological processes peptides exhibited similar RIA values between the diets, while they
exhibited significantly different RIAs for other processes. Specifically, peptides assigned to
carbohydrate derivative metabolic processes, protein folding, response to stimulus, and translation
exhibited significantly higher ^{2}H and ^{18}O RIAs in the high protein medium. Peptides assigned to cell
adhesion, cell wall organization or biogenesis, photosynthesis, protein catabolic processes, protein

maturation, protein-containing complex assembly, and small molecule biosynthetic processes only
exhibited significantly higher ^{18}O RIA in the protein-rich medium. Peptides assigned to carbohydrate
metabolic processes, cytoskeleton organization, generation of precursor metabolites and energy,
nitrogen cycle metabolic processes, nucleobase-containing small molecule metabolic processes, and
protein modification processes exhibited significantly higher ^{2}H RIA in the protein-rich medium.

To further investigate biological processes related to dietary differences, we specifically analyzed all
detected GO child terms of the parental GO slims of amino acid metabolic processes (GO:0006520),
carbohydrate derivative metabolic processes (GO:1901135), and carbohydrate metabolic processes
(GO:0005975). The identified peptides were assigned to a total of 160 biological processes related to
the above-mentioned GO slims, out of which 28 biological processes exhibited significant differences
($p < 0.05$) in ^{2}H or ^{18}O RIA between both media compositions (for specific processes see Supplementary
Table S5). Differences in RIA related to amino acid or carbohydrate metabolism were isotope-specific.
In the protein-rich medium, significantly higher ^{18}O and ^{2}H RIA were observed in peptides associated
with L-histidine catabolism to glutamate and formate, as well as carbohydrate metabolism. The
protein-rich medium also showed significantly elevated ^{18}O RIA in peptides linked to
acetylglucosamine, arabinose, and fucose catabolisms, as well as fructose-6-phosphate metabolism,
while ^{2}H RIA was significantly higher in peptides involved in aminoacylation, biosynthesis of arginine,
UTP, and GMP, as well as aspartate, glucose, and fructose-1,6-bisphosphate metabolism. Conversely,
the fiber-rich medium exhibited significantly higher ^{18}O RIA in peptides related to amino acid
biosynthesis (particularly valine and isoleucine), and higher ^{2}H RIA in peptides associated with cellulose
catabolism, as well as arabinose and nucleoside metabolisms.

Discussion

Our results show that *de novo* peptide sequencing can generate sample-matching databases for ^{2}H,

^{13}C, and ^{18}O Protein-SIP analyses without prior knowledge of community compositions. In our work,

databases derived from Casanovo allowed for the identification of consistently more target peptides

than PepNet-derived databases. Filtering *de novo* identified peptides to those with a quality score

> 0.99 before assembling the database (CS>0.99) enabled the identification of more labeled target

peptides than using a non-filtered database, supporting the conclusion that inflated database size can

challenge database search algorithms, as reported previously [36, 40]. By using the CS>0.99 databases,

we identified 72.6–103.6% of all target peptides and 94.0–100.5% of labeled target peptides compared

to the reference databases generated from (meta)genome information. These results indicate that the

CS>0.99 databases perform comparably to reference databases. Similarly, casanovo-derived databases

were recently reported to outperform the broad-spectrum database Swiss-Prot [55].

Based on the analyzed data and the obtained results, we propose a modular Denovo-SIP pipeline

(Figure 1). In the proposed Denovo-SIP pipeline, peptides are first *de novo* sequenced using Casanovo

[46] to create a peptide sequence database for subsequent database-dependent search. The database-

dependent search is carried out by MS-GF+ [56] and MetaProSIP [26] to validate peptides with a target-

decoy approach and infer isotope incorporation. Unipept [41] and BLASTP [42] can then retrieve taxa

and functions of peptides. Python scripts from the Denovo-SIP pipeline allow the integration of isotope

incorporation data with functional and taxonomic annotation for subsequent analysis. The modular

structure of the Denovo-SIP pipeline allows flexibility, enabling the integration of different engines and

the replacement or supplementation of modules with alternative tools to meet specific needs.. For

example, Sage could increase the speed of database search [57]; Sipros 4 could improve the accuracy

and sensitivity in detecting labeled peptides [27]; Calis-p could enhance the analysis of low amounts

of isotope incorporation [48]; and MetaX could further analyze peptide-associated taxa and functions

[58]. Furthermore, *de novo* peptide sequencing algorithms are developing fast: Engines such as π-

HelixNovo [59], PowerNovo [60], and NovoB [61], recently reported to outperform Casanovo on some

datasets, could be used in the Denovo-SIP pipeline. Moreover, rescoring algorithms could be employed

to increase sensitivity and precision of *de novo* peptide sequencing [62] as well as database searches

[63].

We first tested the Denovo-SIP pipeline by analyzing proteomic data from a defined microbial

community containing ^{13}C-labeled *E. coli* cells. Using the pipeline, we successfully identified the isotope

labeling in *E. coli* and showed that the residual community was unlabeled. We were unable to detect

the unlabeled families *Alteromonadaceae* and *Roseobacteriaceaee*, which may have been caused by

the raw data quality since even with the reference database, only 0 to 2 peptides of those families

could be isotopically quantified in most replicates (Supplementary Table S5 in [48]). Previous analyses

also suggest that raw data quality is a major bottleneck of *de novo* sequencing [45, 64]. We also did

not detect bacterial viruses in the mock community, despite their protein abundance of 0.08–0.14%

(Supplementary Table 3 in [49]). However, these bacterial viruses were also not reliably identified by

metaproteomics using a metagenome-derived database [49]. Similarly, another bottom-up proteomics

study reported difficulties in detecting viruses having a low relative abundance within a complex
microbiome [65].

By using the Denovo-SIP pipeline, we could also confirm that the anammox bacterium '*Candidatus*
Kuenenia stuttgartiensis' strain CSTR1 used bicarbonate as its sole carbon source. Furthermore, we did
not detect any other organisms in the reactor assimilating bicarbonate. This experiment demonstrated
that Denovo-SIP can detect highly enriched peptides with RIA of more than 95% as well as differentially
labeled peptides between time points. Therefore, Denovo-SIP can be applied to time-series studies.

We also employed the Denovo-SIP workflow to analyze previously published data from the human
distal gut model microbiome cultivated in a fiber-rich or protein-rich medium containing either ^{2}H- or
^{18}O-labeled water. This approach is used to detect general activity as deuterium and oxygen from water
is always incorporated when a microorganism is active. Through this analysis, we showed that changes
in activity due to dietary compositions are species- and process-specific in agreement with previous
analyses [18, 48]. Previous work reported that a fiber-rich medium increases activity, thus RIA in
peptides, related to amino acid biosynthesis [18]. Our work specifically suggests that the biosynthesis
of valine and isoleucine, as well as the catabolism of histidine and the metabolism of carbohydrates
were affected by the medium composition.

In conclusion, our results show that *de novo* assembled databases perform comparably to sample-
matching reference databases in detecting labeled peptides. The main advantages of *de novo*
assembled databases are the saved time and costs associated with metagenome sequencing and the
ability to detect foreign peptides, which are typically absent in reference databases. Such foreign
peptides could originate from contamination [66], mutation [67], alternative splicing [68], or
nonspecific cleavage [69]. Moreover, *de novo* assembled databases allow the detection of peptides
from organisms not expected in the sample, as shown here for *C. intestinale* and *H. mulieris* in the
human distal gut model.

De novo assembled databases also have certain limitations. First, *de novo* sequencing was only
successful on peptides with RIA of up to 5%. Usually such low or unlabeled peptides are present in the
beginning or in controls of experiments allowing database creation for subsequent analyses of labeled
samples. This limitation is not specific to *de novo* sequencing. To date, all major SIP tools (SIPPER [25],
MetaProSIP [26], and Sipros [27]) require unlabeled reference samples to identify labeled peptides.
Secondly, *de novo* peptide sequencing is computationally intensive and often requires high-
performance computing graphics processing units (GPUs) to achieve inference speeds comparable to
database searches. Thirdly, *de novo* assembled databases contain amino acid sequences derived from
mass spectra. Amino acid sequences are less variable than nucleic acid sequences. Hence, *de novo*
assembled databases inherently result in lower taxonomic resolution than databases derived from
nucleic acid sequences, such as metagenome sequencing. Furthermore, taxonomy inference via a
lowest common ancestor approach requires the detection of taxon-specific peptides. The number of
taxon-specific peptides decreases with increasing taxonomic resolution [70]. Moreover, the
abundance of taxon-specific peptides varies between taxa [38]. Thus, the overall sensitivity and
taxonomic resolution of lowest common ancestor approaches is dependent on the specific taxa

present. Still, we demonstrated that *de novo* assembled databases can achieve species-specific resolution. Another limitation is that although identifying peptides by *de novo* sequencing does not require their presence in databases, interpreting the identified peptides by inferring proteins, taxa, or functions still requires a reference database. We propose using Unipept for taxa/function inference. However, the current Unipept version 5.1.5 can only infer information from perfect matches of tryptic peptides. Recently, Unipept update 6.0 has been announced to allow the analysis of non-tryptic peptides enhancing the synergy with *de novo* sequencing [71]. To get information about peptides that do not perfectly match, taxonomic drop-off rates and BLAST can be integrated into our automated workflow. In doing so, Denovo-SIP can assess the potential of unreferenced organisms. Estimating false-discovery rates of *de novo*-identified peptides is challenging [72], and currently no tools for *de novo* sequencing of labeled peptides exist. Therefore, our approach of assembling databases through *de novo* peptide sequencing – followed by database searches to verify these peptides and calculate isotope incorporation rates – shows promise. Denovo-SIP could advance the application of Protein-SIP across various microbiological research fields, particularly when screening of many different samples is necessary.

Methods

Cultivation and protein extraction

Proteomics data of the anammox community were obtained with samples taken from a continuous stirred-tank reactor cultivating a mixed culture containing predominantly the anammox bacterium 'Candidatus Kuenenia stuttgartiensis' strain CSTR1 [52]. The reactor had an effective volume of 500 mL and was fed with two medium solutions containing 60 mM ammonium and 60 mM nitrite (concentration after mixing with ratio 1:1), respectively. Sodium bicarbonate (10 mM, concentration after mixing) was the only carbon source provided, provided with the nitrite medium. For a detailed medium recipe, please refer to [52]. The reactor was running stably with non-labeled bicarbonate at 30 °C, constant stirring, and 4-day hydraulic retention time. The cell density was around 2.5×10^8 cells mL^{-1}. At day 0, the nitrite medium containing non-labeled sodium bicarbonate was replaced with a medium containing ^{13}C-labeled sodium bicarbonate, while all other medium components and reactor operational conditions remained the same. Three samples (as triplicates) were taken from the reactor on day 0 (non-labeled), day 4 (1 × hydraulic retention time), and day 21 (5.25 × hydraulic retention time). Samples of 1.5 mL were centrifuged at 10,000 × g, 10 °C for 10 min, and the pellets were frozen at -20 °C until proteomic analysis was done.

Sample preparation and mass spectrometry

Saved pellets from reactor liquid were resuspended in 50 µL of 100 mM ammonium bicarbonate buffer, and went for a 3× freeze/thaw cycle to disrupt the cells (-80 °C followed by 1 min gentle shaking at 40 °C). Two microliter of 40 ng $µL^{-1}$ bovine serum albumin was added to each sample as internal standard. Then, an equal volume of 10% (w/v) sodium deoxycholate (DOC) was added to each sample, before 12 mM dithiothreitol (DTT) and 40 mM 2-iodoacetamide (IAA) (both were final concentrations) were added sequentially to reduce and alkylate the disulfide bond (incubation conditions: 30 min, 37 °C on a rotary shaker at 400 rpm for DTT; and 45 min, room temperature, 400 rpm, in the dark for IAA). Afterwards, 100 mM ammonium bicarbonate was added to dilute DOC to 1% (w/v). Sequencing-grade trypsin (6.3 µL, 0.1 µg $µL^{-1}$, Promega, Madison, USA) was then added to digest the sample at 37 °C, 400 rpm overnight. After trypsin digestion, neat formic acid was added to a final concentration of 2% (v/v), samples were vortexed, and two rounds of centrifugation at 16,000 × g for 10 min were performed to completely remove precipitated DOC. Peptides in the supernatant were desalted using 100 µL C_{18} tips (Pierce, Thermo Fisher Scientific, Massachusetts, USA) following the manufacturer's instruction. Eluted peptides were vacuum dried and resuspended in 50 µL of 0.1% (v/v) formic acid. Peptide concentrations were determined based on the UV absorbance at 205 nm measured on a DS-11 micro-volume spectrophotometer (DeNovix, Delaware, USA). Peptides from day 4 (1 × hydraulic retention time) and day 21 (5.25 × hydraulic retention time) were mixed with peptides from day 0 in a 1:1 ratio before 3 µL were injected for measurement on nano-LC-MS/MS (Orbitrap) as described previously [51] except that the MS2 resolution was lowered from 60,000 to 30,000. The proteomics

data were deposited at ProteomeXchange via the PRIDE partner repository under the accession
number PXD057215 (reviewer token: ejwbd9cUNtVw).

Generation of *de novo* assembled peptide databases

Mass spectrometric raw files (Supplementary Table S6) were converted into mgf format after peak-
picking, removing zero values in spectra, encoding in 64-bit precision, and zlib compression using
msConvert (ProteoWizard v3.0.23304) [73]. Mass spectra were translated to peptide sequence lists by
de novo sequencing using PepNet (v0.3.2) [47] and Casanovo (v3.5.0) [46] with the pre-trained models.
Peptide sequence lists were filtered by a threshold of the quality score (default: >0.99) reported by the
de novo engines. For each dataset, filtered peptide sequence lists were merged and converted to fasta
format using our DeNovo-SIP pipeline (https://git.ufz.de/meb/denovo-sip).

Identification of isotopically labeled peptides

Mass spectrometric raw files (Supplementary Table S6) were converted into mzML format after peak-
picking and encoding in 32-bit precision using msConvert (ProteoWizard v3.0.19046). [73] Next, mzML
files were analyzed using the *de novo* assembled peptide databases and for comparison also with the
sample-matching protein databases appended with universal contaminants [66] by a customized
MetaProSIP workflow [74] based on OpenMS v2.8 [75] on the Galaxy [76] instance of the Helmholtz
Centre for Environmental Research – UFZ. In brief, peptides were detected and validated by MS-GF+
[56] using a target-decoy approach, and peptide relative isotope abundances (RIA), as well as labeling
ratios (LR), were calculated by MetaProSIP [26]. The precursor mass tolerance was set to ± 10 ppm (for
the mock community with spike-ins of ^{13}C-labeled *Escherichia coli* [48] and the model of the human
gut microbiome [18]) or ± 5 ppm (for the '*Candidatus* Kuenenia stuttgartiensis' enrichment). We
allowed a peptide length of 6–100 amino acids, no (for peptide databases) or semi-tryptic (for protein
databases) cleavage and up to two of the following modifications: variable oxidation of methionine,
deamidation of asparagine or glutamine, N-terminal acetylation, carbamylation, or ammonia-loss and
fixed C-terminal carbamidomethylation. The weight merge window was set to ± 1.5 (for ^{2}H) and ± 5.0
(for ^{18}O and ^{13}C) in MetaProSIP. Identifications not assigned to a feature during pattern detection and
purely natural peptides were included in the report. The peptide-centric output of MetaProSIP was
used for further processing. Peptides for which MetaProSIP reported at least one non-natural weight
were considered as labeled. The taxonomy and gene ontology (GO) [77] terms of peptides were
inferred based on the lowest common ancestor approach by Unipept Desktop v2.0.2 [78] using UniProt
2024.03 [79]. Advanced missed cleavage handling and indistinguishability of leucine and isoleucine
were enabled. GO terms of biological processes were mapped to GO slims of the prokaryote subset
maintained by the GO Consortium [80] (goslim_prokaryote) using GOATOOLS [81]. Python scripts from
our Denovo-SIP pipeline (https://git.ufz.de/meb/denovo-sip) merged results from Unipept and
MetaProSIP for downstream analysis.

Sequence similarity search of unassigned peptides

Labeled peptides that could not be assigned to a lowest common ancestor by Unipept were searched for similar sequences using BLASTP 2.16.0+ [42] with the nr database (accession date: 2024/11/01). Hits with e-value ≥ 0.001 were removed before plotting top-hit species distributions using Blast2GO 6.0.3 [82].

Taxonomy drop-off rates

The number of taxon-specific peptides decreases as taxonomic resolution increases, resulting in a 'drop-off' from higher to lower taxonomic levels. Comparing observed and theoretical drop-off rates within a lineage allows assessing whether the peptide count at a given taxonomic level (e.g., species level) can explain the peptide count at higher taxonomic levels (e.g., phylum level) [43]. By this, drop-off rates can reveal 'hidden' organisms not captured by a taxonomic database (here: Unipept). In this manuscript, we use the term 'superordinate taxa' to refer to taxonomic ranks above a given taxon. For example, the superordinate taxa of *Escherichia coli* (species) include *Escherichia* (genus), *Enterobacteriaceae* (family), *Enterobacterales* (order), *Gammaproteobacteria* (class), *Pseudomonadota* (phylum), and *Bacteria* (domain).

Labeled peptides were tested for hidden organisms comparing observed with theoretical taxonomy drop-off rates of labeled taxa. To calculate theoretical drop-off rates, we extracted expressed proteins from the proteome of *E. coli* K12 and '*Ca.* K. stuttgartiensis' strain CSTR1 based on previous studies (Table S7 in [51] and Run4_Ecoli268_R3a_0_13C_400ng.pdResult in PXD023693 [83] from [48]). Expressed proteins were *in-silico* digested into tryptic peptides with a length of 7–15 amino acids. The detectability of peptides was assessed by DeepMSpeptide [84] and detectable peptides were processed by Unipept as explained above. We accounted for false-positive Unipept annotations by employing a sequence randomization strategy as described previously [43].

Availability of data and materials

The raw mass spectrometric proteomics data generated during this study have been deposited to the ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE partner repository [83] with the dataset identifier PXD057215 (reviewer token: ejwbd9cUNtVw). Raw mass spectrometry proteomics data reanalyzed during this study is also available via the PRIDE partner repository (Supplementary Table S6). Data obtained from our reanalyses has been deposited to Zenodo (Supplementary Table S6). The Denovo-SIP pipeline's Python scripts are available via GitLab (https://git.ufz.de/meb/denovo-sip). Any other relevant data will be made available upon request.

References

1. Berg G, Rybakova D, Fischer D, Cernava T, Vergès MCC, Charles T, et al. Microbiome definition re-visited: old concepts and new challenges. Microbiome. 2020;8:103.

2. Layeghifard M, Hwang DM, Guttman DS. Disentangling interactions in the microbiome: a network perspective. Trends Microbiol. 2017;25:217–28.

3. Lynch S V, Pedersen O. The human intestinal microbiome in health and disease. N Engl J Med. 2016;375:2369–79.

4. Falkowski PG, Fenchel T, Delong EF. The microbial engines that drive earth's biogeochemical cycles. Science. 2008;320:1034–9.

5. Knight R, Vrbanac A, Taylor BC, Aksenov A, Callewaert C, Debelius J, et al. Best practices for analysing microbiomes. Nat Rev Microbiol. 2018;16:410–22.

6. Zhang X, Li L, Butcher J, Stintzi A, Figeys D. Advancing functional and translational microbiome research using meta-omics approaches. Microbiome. 2019;7:154.

7. Jehmlich N, Vogt C, Lünsmann V, Richnow HH, von Bergen M. Protein-SIP in environmental studies. Curr Opin Biotechnol. 2016;41:26–33.

8. Radajewski S, Webster G, Reay DS, Morris SA, Ineson P, Nedwell DB, et al. Identification of active methylotroph populations in an acidic forest soil by stable-isotope probing. Microbiology. 2002;148:2331–42.

9. Lueders T, Wagner B, Claus P, Friedrich MW. Stable isotope probing of rRNA and DNA reveals a dynamic methylotroph community and trophic interactions with fungi and protozoa in oxic rice field soil. Environ Microbiol. 2004;6:60–72.

10. Manefield M, Whiteley AS, Griffiths RI, Bailey MJ. RNA stable isotope probing, a novel means of linking microbial community function to phylogeny. Appl Environ Microbiol. 2002;68:5367–73.

11. Treonis AM, Ostle NJ, Stott AW, Primrose R, Grayston SJ, Ineson P. Identification of groups of metabolically-active rhizosphere microorganisms by stable isotope probing of PLFAs. Soil Biol Biochem. 2004;36:533–7.

12. Boschker HTS, De Graaf W, Köster M, Meyer-Reil LA, Cappenberg TE. Bacterial populations and processes involved in acetate and propionate consumption in anoxic brackish sediment. FEMS Microbiol Ecol. 2001;35:97–103.

13. Jehmlich N, Schmidt F, Von Bergen M, Richnow HH, Vogt C. Protein-based stable isotope probing (Protein-SIP) reveals active species within anoxic mixed cultures. ISME J. 2008;2:1122–33.

14. Jehmlich N, Schmidt F, Hartwich M, von Bergen M, Richnow H-H, Vogt C. Incorporation of carbon and nitrogen atoms into proteins measured by protein-based stable isotope probing (Protein-SIP). Rapid Commun Mass Spectrom. 2008;22:2889–97.

15. Justice NB, Li Z, Wang Y, Spaudling SE, Mosier AC, Hettich RL, et al. ^{15}N- and ^{2}H proteomic stable isotope probing links nitrogen flow to archaeal heterotrophic activity. Environ Microbiol.

2014;16:3224–37.

16. Berry D, Mader E, Lee TK, Woebken D, Wang Y, Zhu D, et al. Tracking heavy water (D_2O) incorporation for identifying and sorting active microbial cells. Proc Natl Acad Sci U S A. 2015;112:E194–203.

17. Taubert M, Stöckel S, Geesink P, Girnus S, Jehmlich N, von Bergen M, et al. Tracking active groundwater microbes with D_2O labelling to understand their ecosystem function. Environ Microbiol. 2018;20:369–84.

18. Starke R, Oliphant K, Jehmlich N, Schäpe SS, Sachsenberg T, Kohlbacher O, et al. Tracing incorporation of heavy water into proteins for species-specific metabolic activity in complex communities. J Proteomics. 2020;222:103791.

19. Pfeffer M, Maurer M, Köllensperger G, Hann S, Graf AB, Mattanovich D. Modeling and measuring intracellular fluxes of secreted recombinant protein in *Pichia pastoris* with a novel ^{34}S labeling procedure. Microb Cell Fact. 2011;10:47.

20. Kümmel S, Herbst FA, Bahr A, Duarte M, Pieper DH, Jehmlich N, et al. Anaerobic naphthalene degradation by sulfate-reducing *Desulfobacteraceae* from various anoxic aquifers. FEMS Microbiol Ecol. 2015;91:fiv006.

21. Ouyang WY, Su JQ, Richnow HH, Adrian L. Identification of dominant sulfamethoxazole-degraders in pig farm-impacted soil by DNA and protein stable isotope probing. Environ Int. 2019;126 February:118–26.

22. Nowak KM, Miltner A, Gehre M, Schäffer A, Kästner M. Formation and fate of bound residues from microbial biomass during 2, 4-D degradation in soil. Environ Sci Technol. 2011;45:999–1006.

23. Grob C, Taubert M, Howat AM, Burns OJ, Dixon JL, Richnow HH, et al. Combining metagenomics with metaproteomics and stable isotope probing reveals metabolic pathways used by a naturally occurring marine methylotroph. Environ Microbiol. 2015;17:4007–18.

24. Bastida F, Jechalke S, Bombach P, Franchini AG, Seifert J, von Bergen M, et al. Assimilation of benzene carbon through multiple trophic levels traced by different stable isotope probing methodologies. FEMS Microbiol Ecol. 2011;77:357–69.

25. Slysz GW, Steinke L, Ward DM, Klatt CG, Clauss TRW, Purvine SO, et al. Automated data extraction from in situ protein-stable isotope probing studies. J Proteome Res. 2014;13:1200–10.

26. Sachsenberg T, Herbst FA, Taubert M, Kermer R, Jehmlich N, Von Bergen M, et al. MetaProSIP: automated inference of stable isotope incorporation rates in proteins for functional metaproteomics. J Proteome Res. 2015;14:619–27.

27. Xiong Y, Mueller RS, Feng S, Guo X, Pan C. Proteomic stable isotope probing with an upgraded Sipros algorithm for improved identification and quantification of isotopically labeled proteins. Microbiome. 2024;12:148.

28. Zhang X, Ning Z, Mayne J, Moore JI, Li J, Butcher J, et al. MetaPro-IQ: a universal metaproteomic approach to studying human and mouse gut microbiota. Microbiome. 2016;4:1–12.

29. Mise K, Iwasaki W. Unexpected absence of ribosomal protein genes from metagenome-assembled genomes. ISME Commun. 2022;2:118.

30. Rocha U, Kasmanas JC, Toscan R, Sanches DS, Magnusdottir S, Saraiva JP. Simulation of 69 microbial communities indicates sequencing depth and false positives are major drivers of bias in prokaryotic metagenome-assembled genome recovery. PLOS Comput Biol. 2024;20:e1012530.

31. Jechalke S, Franchini AG, Bastida F, Bombach P, Rosell M, Seifert J, et al. Analysis of structure, function, and activity of a benzene-degrading microbial community. FEMS Microbiol Ecol. 2013;85:14–26.

32. Starke R, Kermer R, Ullmann-Zeunert L, Baldwin IT, Seifert J, Bastida F, et al. Bacteria dominate the short-term assimilation of plant-derived N in soil. Soil Biol Biochem. 2016;96:30–8.

33. Herbst FA, Bahr A, Duarte M, Pieper DH, Richnow HH, von Bergen M, et al. Elucidation of in situ
polycyclic aromatic hydrocarbon degradation by functional metaproteomics (protein-SIP). Proteomics.
2013;13:2910–20.

34. Bozinovski D, Herrmann S, Richnow HH, Von Bergen M, Seifert J, Vogt C. Functional analysis of an
anaerobic *m*-xylene-degrading enrichment culture using protein-based stable isotope probing. FEMS
Microbiol Ecol. 2012;81:134–44.

35. Blakeley-Ruiz JA, Kleiner M. Considerations for constructing a protein sequence database for
metaproteomics. Computational and Structural Biotechnology Journal. 2022;20:937–52.

36. Kumar D, Yadav AK, Dash D. Choosing an optimal database for protein identification from tandem
mass spectrometry data. In: Keerthikumar S, Mathivanan S, editors. Proteome Bioinformatics. New
York, NY: Springer New York; 2017. p. 17–29.

37. Knudsen GM, Chalkley RJ. The effect of using an inappropriate protein database for proteomic data
analysis. PLoS One. 2011;6:e20873-.

38. Tanca A, Palomba A, Deligios M, Cubeddu T, Fraumene C, Biosa G, et al. Evaluating the impact of
different sequence databases on metaproteome analysis: insights from a lab-assembled microbial
mixture. PLoS One. 2013;8:e82981.

39. Klaes S, Madan S, Deobald D, Cooper M, Adrian L. Revealing taxonomy, activity, and substrate
assimilation in mixed bacterial communities by GroEL-proteotyping-based stable isotope probing.
iScience. 2024;27:111249.

40. Muth T, Kolmeder CA, Salojärvi J, Keskitalo S, Varjosalo M, Verdam FJ, et al. Navigating through
metaproteomics data: a logbook of database searching. Proteomics. 2015;15:3439–53.

41. Mesuere B, Devreese B, Debyser G, Aerts M, Vandamme P, Dawyndt P. Unipept: Tryptic peptide-
based biodiversity analysis of metaproteome samples. J Proteome Res. 2012;11:5773–80.

42. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST:
a new generation of protein database search programs. Nucleic Acids Res. 1997;25:3389–402.

43. Kleikamp HBC, Pronk M, Tugui C, Guedes da Silva L, Abbas B, Lin YM, et al. Database-independent
de novo metaproteomics of complex microbial communities. Cell Syst. 2021;12:375-383.e5.

44. Johnson RS, Searle BC, Nunn BL, Gilmore JM, Phillips M, Amemiya CT, et al. Assessing protein
sequence database suitability using de novo sequencing. Mol Cell Proteomics. 2020;19:198–208.

45. Muth T, Renard BY. Evaluating de novo sequencing in proteomics: already an accurate alternative
to database-driven peptide identification? Brief Bioinform. 2018;19:954–70.

46. Yilmaz M, Fondrie WE, Bittremieux W, Melendez CF, Nelson R, Ananth V, et al. Sequence-to-
sequence translation from mass spectra to peptides with a transformer model. Nat Commun.
2024;15:6427.

47. Liu K, Ye Y, Li S, Tang H. Accurate de novo peptide sequencing using fully convolutional neural
networks. Nat Commun. 2023;14:7974.

48. Kleiner M, Kouris A, Violette M, D'Angelo G, Liu Y, Korenek A, et al. Ultra-sensitive isotope probing
to quantify activity and substrate assimilation in microbiomes. Microbiome. 2023;11:24.

49. Kleiner M, Thorson E, Sharp CE, Dong X, Liu D, Li C, et al. Assessing species biomass contributions
in microbial communities via metaproteomics. Nat Commun. 2017;8:1588.

50. Masao F, Kenichi K, Ryuji K. Phylogenetic analysis of *Salmonella*, *Shigella*, and *Escherichia coli*
strains on the basis of the *gyrB* gene sequence. J Clin Microbiol. 2002;40:2779–85.

51. Ding C, Adrian L. Comparative genomics in '*Candidatus* Kuenenia stuttgartiensis' reveal high
genomic plasticity in the overall genome structure, CRISPR loci and surface proteins. BMC Genomics.
2020;21:851.

52. Ding C, Enyi FO, Adrian L. Anaerobic ammonium oxidation (Anammox) with planktonic cells in a
redox-stable semicontinuous stirred-tank reactor. Environ Sci Technol. 2018;52:5671–81.

53. Feng J-H, Chang Y-X, Li L-H, Li M-M, Huang Z-Y, Han N, et al. *Copranaerobaculum intestinale* gen. nov., sp. nov., a novel anaerobic bacterium isolated from human faeces. Int J Syst Evol Microbiol. 2022;72.

54. Afrizal A, Hitch TCA, Viehof A, Treichel N, Riedel T, Abt B, et al. Anaerobic single-cell dispensing facilitates the cultivation of human gut bacteria. Environ Microbiol. 2022;24:3861–81.

55. Chiang Y, Collins MJ. Orthrus: an AI-powered, cloud-ready, and open-source hybrid approach for metaproteomics. bioRxiv. 2024;:2024.11.15.623814.

56. Kim S, Pevzner PA. MS-GF+ makes progress towards a universal database search tool for proteomics. Nat Commun. 2014;5:5277.

57. Lazear MR. Sage: an open-source tool for fast proteomics searching and quantification at scale. J Proteome Res. 2023;22:3652–9.

58. Wu Q, Ning Z, Zhang A, Zhang X, Sun Z, Figeys D. MetaX: a peptide centric metaproteomic data analysis platform using Operational Taxa-Functions (OTF). bioRxiv. 2024;:2024.04.19.590315.

59. Yang T, Ling T, Sun B, Liang Z, Xu F, Huang X, et al. Introducing π-HelixNovo for practical large-scale de novo peptide sequencing. Brief Bioinform. 2024;25.

60. Petrovskiy D V, Nikolsky KS, Kulikova LI, Rudnev VR, Butkova T V, Malsagova KA, et al. PowerNovo: de novo peptide sequencing via tandem mass spectrometry using an ensemble of transformer and BERT models. Sci Rep. 2024;14:15000.

61. Lee S, Kim H. Bidirectional de novo peptide sequencing using a transformer model. PLoS Comput Biol. 2024;20:e1011892.

62. Klaproth-Andrade D, Hingerl J, Bruns Y, Smith NH, Träuble J, Wilhelm M, et al. Deep learning-driven fragment ion series classification enables highly precise and sensitive de novo peptide sequencing. Nat Commun. 2024;15:151.

63. Ananth V, Sanders J, Yilmaz M, Wen B, Oh S, Noble WS. A learned score function improves the power of mass spectrometry database search. Bioinformatics. 2024;40 Supplement_1:i410–7.

64. McDonnell K, Howley E, Abram F. The impact of noise and missing fragmentation cleavages on de novo peptide identification algorithms. Comput Struct Biotechnol J. 2022;20:1402–12.

65. Lozano C, Pible O, Eschlimann M, Giraud M, Debroas S, Gaillard J-C, et al. Universal identification of pathogenic viruses by liquid chromatography coupled to tandem mass spectrometry proteotyping. Mol Cell Proteomics. 2024;:100822.

66. Frankenfield AM, Ni J, Ahmed M, Hao L. Protein contaminants matter: building universal protein contaminant libraries for DDA and DIA proteomics. J Proteome Res. 2022;21:2104–13.

67. Sinitcyn P, Richards AL, Weatheritt RJ, Brademan DR, Marx H, Shishkova E, et al. Global detection of human variants and isoforms by deep proteome sequencing. Nat Biotechnol. 2023;41:1776–86.

68. Faridi P, Li C, Ramarathinam SH, Vivian JP, Illing PT, Mifsud NA, et al. A subset of HLA-I peptides are not genomically templated: evidence for cis- and trans-spliced peptide ligands. Sci Immunol. 2018;3:eaar3947.

69. Gattiker A, Bienvenut W V., Bairoch A, Gasteiger E. FindPept, a tool to identify unmatched masses in peptide mass fingerprinting protein identification. Proteomics. 2002;2:1435–44.

70. Klaes S, Madan S, Deobald D, Cooper M, Adrian L. GroEL-proteotyping of bacterial communities using tandem mass spectrometry. International Journal of Molecular Sciences. 2023;24:15692.

71. Vande Moortele T, Devlaminck B, Van de Vyver S, Van Den Bossche T, Martens L, Dawyndt P, et al. Unipept 6.0: expanding metaproteomics analysis with support for missed cleavages, semi-tryptic and non-tryptic peptides. bioRxiv. 2024;:2024.09.26.615136.

72. O'Bryon I, Jenson SC, Merkley ED. Flying blind, or just flying under the radar? The underappreciated power of de novo methods of mass spectrometric peptide identification. Protein Sci. 2020;29:1864–78.

73. Kessner D, Chambers M, Burke R, Agus D, Mallick P. ProteoWizard: open source software for rapid
proteomics tools development. Bioinformatics. 2008;24:2534–6.

74. Jehmlich N, von Bergen M. Protein-based stable isotope probing (Protein-SIP): applications for
studying aromatic hydrocarbon degradation in microbial communities. In: Anaerobic utilization of
hydrocarbons, oils, and lipids. Cham: Springer International Publishing; 2020. p. 277–84.

75. Röst HL, Sachsenberg T, Aiche S, Bielow C, Weisser H, Aicheler F, et al. OpenMS: a flexible open-
source software platform for mass spectrometry data analysis. Nat Methods. 2016;13:741–8.

76. The Galaxy Community. The Galaxy platform for accessible, reproducible and collaborative
biomedical analyses: 2022 update. Nucleic Acids Res. 2022;50:W345–51.

77. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the
unification of biology. Nat Genet. 2000;25:25–9.

78. Verschaffelt P, Tanca A, Abbondio M, Van Den Bossche T, Moortele T Vande, Dawyndt P, et al.
Unipept Desktop 2.0: construction of targeted reference protein databases for metaproteogenomics
analyses. J Proteome Res. 2023;22:2620–8.

79. The UniProt Consortium. UniProt: the universal protein knowledgebase in 2023. Nucleic Acids Res.
2023;51:D523–31.

80. Aleksander SA, Balhoff J, Carbon S, Cherry JM, Drabkin HJ, Ebert D, et al. The gene ontology
knowledgebase in 2023. Genetics. 2023;224:iyad031.

81. Klopfenstein D V., Zhang L, Pedersen BS, Ramírez F, Vesztrocy AW, Naldi A, et al. GOATOOLS: a
Python library for gene ontology analyses. Sci Rep. 2018;8:10872.

82. Götz S, García-Gómez JM, Terol J, Williams TD, Nagaraj SH, Nueda MJ, et al. High-throughput
functional annotation and data mining with the Blast2GO suite. Nucleic Acids Res. 2008;36:3420–35.

83. Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S, et al. The
PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. Nucleic
Acids Res. 2022;50:D543–52.

84. Serrano G, Guruceaga E, Segura V. DeepMSPeptide: peptide detectability prediction using deep
learning. Bioinformatics. 2020;36:1279–80.

Acknowledgements

Proteomics raw data published in this study was acquired at UFZ at the Centre for Chemical Microscopy (ProVIS), which is supported by the European Regional Development Fund (EFRE) and the Helmholtz Association. *De novo* sequencing was performed at the High-Performance Computing (HPC) Cluster EVE, a joint effort of both the Helmholtz Centre for Environmental Research – UFZ (http://www.ufz.de/) and the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (http://www.idiv-biodiversity.de/). We would like to thank Thomas Schnicke, Ben Langenberg, Guido Schramm, Toni Harzendorf, Tom Strempel, Lisa Schurack, and Christian Krause for maintaining EVE as well as Benjamin Scheer for technical assistance in protein mass spectrometry. We acknowledge the assistance of generative AI (ChatGPT and Claude) in code development and troubleshooting, with all code subsequently reviewed and verified by the authors. Figure 1 was created in BioRender (https://BioRender.com/o64a634). S.K. was funded by the German Research Foundation (DFG) grant number GRK 2032/2. C.W. received funding from the Erasmus+ ICM (KA171) program of the European Commission. Open Access funding was enabled by Project DEAL.

Author contributions

Conceptualization: S.K.; Data curation: S.K. and C.D.; Formal analysis: S.K. and C.D.; Funding acquisition: L.A.-C. and L.A.; Investigation: S.K. and C.W.; Methodology: S.K. and C.D.; Project administration: S.K. and C.D.; Resources: C.W., L.A.-C., and L.A..; Software: S.K. and C.D.; Supervision: C.D.; Validation: S.K. and C.D.; Visualization: S.K.; Writing – original draft: S.K.; Writing – review & editing: S.K., C.W., L.A.-C., L.A. and C.D.

Competing financial interests

The authors declare that they have no competing financial interest.

Supplementary Material

815   Supplementary Table S1: Sizes of *de novo* assembled databases used in this study.

Non-redundant *de novo* identified peptides						
Dataset identifier	PXD024174		AMX		PXD024291	
Score threshold	Casanovo	PepNet	Casanovo	PepNet	Casanovo	PepNet
0	290,680	875,366	256,559	303,595	512,680	715,031
0.2	149,626	258,054	82,754	33,867	250,323	225,748
0.4	148,847	175,840	74,423	23,321	247,553	170,283
0.6	139,704	119,472	54,101	16,636	232,755	128,653
0.8	96,835	80,093	29,304	12,001	180,762	97,990
0.85	76,673	71,180	22,380	10,960	153,699	90,692
0.9	53,981	61,634	15,785	9,819	120,752	82,718
0.95	31,873	50,450	9,934	8,473	84,176	72,797
0.99	14,700	35,375	5,313	6,605	48,715	58,476

817   Supplementary Table S2: Drop-off data from *Escherichia coli*.

Labeled peptides	Total submissions	Root	Superkingdom	Phylum	Class	Order	Family	Genus	Species
in silico	3,499	3,499	1,344	1,123	1,071	779	659	44	29
main	450	419	150	120	112	71	53	3	3
others			0	1	7	9	9	9	7
random_main			8	3	1	0	0	0	0
random_others			5	7	8	10	9	9	7
main-random_main			142	117	111	71	53	3	3
others-random_others			-5	-6	-1	-1	0	0	0
purity [%]			103.6	105.4	100.9	101.4	100.0	100.0	100.0
main-random_main normalised			100.0	82.4	78.2	50.0	37.3	2.1	2.1
in silico normalised			100.0	83.6	79.7	58.0	49.0	3.3	2.2
drop off delta [%]				1.4	1.9	15.9	31.4	55.0	2.1

820   Supplementary Table S3: Comparison of the number of detected target peptides in unmixed and mixed

821   samples of case study 1.

x hydraulic retention time		0	1		5.25	
			unmixed	mixed	unmixed	mixed
Peptides identified by MetaProSIP with genome-based database		5,830	3,088	4,449	130	3,329
Target peptides identified by *de novo* sequencing	Casanovo	6,695	3,628	5,120	322	3,713
	PepNet	4,336	2,035	3,171	97	2,131

823   Supplementary Table S4: Drop-off data from '*Candidatus* Kuenenia stuttgartiensis'.

Labeled peptides	Total submissions	Root	Domain	Phylum	Class	Order	Family	Genus	Species
in silico	3,500	3,500	2,544	2,326	2,312	2,312	2,258	2,081	2,081
main	1,232	1,173	847	768	765	765	745	635	635
others		0	0	1	1	1	1	1	1
random_main		314	33	0	0	0	0	0	0
random_others		0	24	43	38	37	37	38	38
main-random_main		859	814	768	765	765	745	635	635
others-random_others		0	-24	-42	-37	-36	-36	-37	-37
purity [%]		100.0	103.0	105.8	105.1	104.9	105.1	106.2	106.2
main-random_main normalised			100.0	94.3	94.0	94.0	91.5	78.0	78.0
in silico normalised			100.0	91.4	90.9	90.9	88.8	81.8	81.8
drop off delta [%]			0.0	3.1	3.3	3.3	3.0	4.9	4.9

825   Supplementary Table S5. Relative isotope abundance of peptides assigned to gene ontology (GO) terms

826   of biological processes related to subterms of amino acid metabolic processes (GO:0006520),

827   carbohydrate derivative metabolic processes (GO:1901135), and carbohydrate metabolic processes

828   (GO:0005975). P-values were calculated by Student's t-test for the means of two independent samples.

 Supplementary Table S6: Overview of analyzed datasets

Dataset	Identifier	Files analyzed by MetaProSIP workflow	Analysis deposited to
Standard *Escherichia coli* K12 cultures cultivated with 1.07–99% ^{13}C [27]	PXD041414	none	https://zenodo.org/records/14030621?token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6ImE3YmQzOTE1LTkwMmUtNDUxOS04M2I1LTc3NjQyMTBkNGFjNSIsImRhdGEiOnt9LCJyYW5kb20iOiIyM2NkYjRmMmEyZDAxZWFmZjAxMzcwNzViNDExZGZiZCJ9.P7_Y1CciVFvJLeMB_8oFEVkOAbEMy5W5fE4Lz_9KoKuc8p8-mqYh3Epw2HkK2SwmwuFKwHNcUii7VXFxEZCG9A
Mock community with spike-in of ^{13}C-labeled *Escherichia coli* [48]	PXD024174	Run1_MockU2_EcoliR1_10_2000ng.raw Run1_MockU2_EcoliR2_10_2000ng.raw Run1_MockU2_EcoliR3_10_2000ng.raw	https://zenodo.org/records/14030389?token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6IjU3NDVlYWRkLTE3ZmYtNDVhOC05OTU1LWQ2ZTQxYWIyNTMzOSIsImRhdGEiOnt}LCJyYW5kb20iOiI3ZGQ3NjAxZDE4MjYwYzZkMjhjOTlhNzJjMTBkMGU3ZiJ9.dkyBgYoi3K4yZOJZCKCCy0FS1EFw5HsiedYpqDnKQXMebDMwRhQppR5cnVqelvFEknwAddDKNlaurQRBc2VwVA
'Candidatus Kuenenia stuttgartiensis' strain CSTR1 reactor	PXD057215	01_6_27_12C_01.raw 02_6_27_12C_02.raw 03_6_27_12C_03.raw 19_7_01_mix_01.raw 20_7_01_mix_02.raw 21_7_01_mix_03.raw 22_7_18_mix_01.raw 23_7_18_mix_02.raw 24_7_18_mix_03.raw	https://zenodo.org/records/14030687?token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6IjdmMGRiMGQ1LWJkMGMtNDBjOC1hOGFiLTZmODM2ZGU0ODE1YyIsImRhdGEiOnt9LCJyYW5kb20iOiI4Y2M3ZGViMWZjZGRiMjRhMWY2NDIzOGYzNzExNWYwNiJ9.NULxsUIzIhoxfapIhgyv-OaY5TPKPsXHpDbQeaEh98Dn84KhFK4nrInCFpLe0x5jZmgMJ9lxzHxYOmTc13YW3w
Model of the human distal gut microbiome [18, 48]	PXD024291	2017-02-07_RS_Robo_9.raw 2017-02-07_RS_Robo_10.raw 2017-02-07_RS_Robo_11.raw 2017-02-07_RS_Robo_12.raw 2017-02-07_RS_Robo_13.raw 2017-02-07_RS_Robo_14.raw 2017-02-07_RS_Robo_15.raw 2017-02-07_RS_Robo_16.raw 2017-02-07_RS_Robo_17.raw 2017-02-07_RS_Robo_18.raw 2017-02-07_RS_Robo_19.raw 2017-02-07_RS_Robo_20.raw	https://zenodo.org/records/14030664?token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6ImZlNWU0MDM4LTRkMjEtNDYxNC1hN2NkLWYzZDY1YjYwMDZiZSIsImRhdGEiOnt9LCJyYW5kb20iOiIyZmY4ZjgwODY5NDY5ZGRkMmI3OTNjZTRlNjdmOTlkYiJ9.CupCCNWTKUZ8ekCcyxHXkmuMNedLw75j0-LJm5ljf3cs24WbrQI38qbFXCy4m7CmadsWMuHIG17_lCvqWzRVxw

Supplementary Figure S1: Comparing *de novo* assembled databases for identifying unlabeled and [13]C-labeled

peptides with a MetaProSIP workflow in a '*Candidatus* Kuenenia stuttgartiensis' strain CSTR1 enrichment from a

continuous stir tank reactor fed with [13]C-bicarbonate. *De novo* databases were assembled using PepNet and

filtering identifications with different thresholds for the quality score. Navy circles denote the number of

identified peptides present in the proteome of '*Ca.* Kuenenia stuttgartiensis' + contaminants. Grey crosses

denote the number of identified peptides absent in the proteome of '*Ca.* Kuenenia stuttgartiensis' +

contaminants. Dashed lines represent the number of identifications using the sample-matching reference

database derived from metagenome sequencing [51]. Only peptides detected in at least two biological replicates

are plotted.

Supplementary Figure S2: Comparing *de novo* assembled databases for identifying unlabeled and $^{2}H/^{18}O$-labeled
peptides with a MetaProSIP workflow in a model of the human gut microbiome (Robogut community) cultivated
with $^{2}H_2O$ or $H_2^{18}O$, respectively, and obtained from [18]. *De novo* databases were assembled using PepNet and
filtering identifications with different thresholds for the quality score. Navy circles denote the number of
identified peptides present in the sample-matching reference proteome + contaminants. Grey crosses denote
the number of identified peptides absent in the sample-matching reference proteome + contaminants. Dashed
lines represent the number of identifications that we generated using the published sample-matching reference
database. Only peptides detected in at least two biological replicates are plotted.

Supplementary Figure S3: Distribution of the ^{18}O and ^{2}H relative isotope abundance (RIA) in detected peptides
identified in a model of the human gut microbiome cultivated in either a high fiber or high protein medium in
the presence of either $H_2^{18}O$ or $^{2}H_2O$. RIA values are shown for peptides detected in at least two replicates of
both media. 'n' denotes the number of peptides identified across triplicates. Statistically significant differences
in RIA between media are indicated with '***' based on Student's t-test for the means of two independent
samples with p < 0.001.

Supplementary Figure S4: Figure 8 without removal of outliers.

Supplementary Figure S5: Figure 9 without removal of outliers.