The presence of a marker associated with Pacific oyster resistance to OsHV-1

does not affect susceptibility to *Vibrio aestuarianus*

Liam B. Surry[1†], Ben J. G. Sutherland[1†], Spencer L. Lunda[2], Andrew H. Loudon[1,3], Konstantin Divilov[4],

Chris J. Langdon[5], Timothy J. Green[1]

[1] Vancouver Island University, Center for Shellfish Research, Nanaimo, BC, Canada V9R 5S5

[2] Oregon State University, Department of Microbiology, 226 Nash Hall, Corvallis, OR 97331

[3] Vancouver Island University, Department of Biology, Nanaimo, BC, Canada V9R 5S5

[4] Center for Quantitative Life Sciences, Oregon State University, Corvallis, OR 97331

[5] Department of Fisheries, Wildlife, and Conservation Sciences, Coastal Oregon Marine Experiment

Station, Oregon State University, Hatfield Marine Science Center, Newport, OR 97365, US

[†] these authors contributed equally to this work

*Author for correspondence: LBS and BJGS

Current Addresses:

Faculty of Science and Technology, Vancouver Island University, Nanaimo, BC, Canada V9R 5S5

liambsurry@gmail.com

Ben.Sutherland@viu.ca

Running Title: Chr8 marker and *V. aestuarianus*

Abstract

The Pacific oyster, *Crassostrea* (*Magallana*) *gigas*, is an important species in aquaculture globally, but its production is threatened by pathogens including ostreid herpesvirus 1 (OsHV-1) and *Vibrio aestuarianus*. A genetic marker associated with resistance to OsHV-1 infection has been identified on chromosome 8 (Chr8) of the Pacific oyster genome. Marker-assisted selective breeding has used the Chr8 marker to produce oyster families with increased survivorship and resistance to OsHV-1 infection. The potential effect of the Chr8 marker on susceptibility to other pathogens remains largely unknown, but was recently associated with increased resistance of spat to *Vibrio coralliilyticus*. Here we assess the effect of the presence and allelic dosage of the Chr8 marker on the susceptibility of selectively bred juvenile Pacific oysters to *V. aestuarianus* infection. Sixteen families were produced with various Chr8 marker genotypes, and a *V. aestuarianus* disease challenge was conducted. Challenged oysters were individually genotyped at the Chr8 marker, and *Vibrio* susceptibility was evaluated among genotypes within and between families. Prior to the challenge trial, the Chr8 marker did not occur at the expected Mendelian ratios in families with heterozygous parents; in multiple families the homozygous alternate genotype occurred at a lower frequency than was expected. Mortality rates in the *Vibrio* challenge differed between families, ranging from 47.9% to 85.4%, but no association was observed between the Chr8 marker and survival to *V. aestuarianus* exposure. Therefore, we observed no pleiotropic effect (positive or negative) of the Chr8 marker on survivorship to *V. aestuarianus* at the evaluated life stage. Reduced-representation sequencing was used to genotype the challenged oysters and a genome-wide association study for *V. aestuarianus* survivorship was performed, but no significant associations were found, suggesting polygenic architecture for the trait.

Keywords: genome-wide association study; OsHV-1; ostreid herpesvirus 1; Pacific oyster; *Vibrio aestuarianus*; virus.

Introduction

The Pacific oyster *Crassostrea* (*Magallana*) *gigas* plays a prominent role in the aquaculture industry, accounting for more than 98% of all oyster production worldwide, with over five million metric tons produced annually (Azéma et al., 2017; Botta et al., 2020; Laurence et al., 2009; Okon et al., 2023). Native to the northwest Pacific Ocean, *C. gigas* has been introduced globally for aquaculture purposes because of its high growth rate and adaptability to various environments (Zhai et al., 2021). Aquaculture aims to supply aquatic-sourced proteins to meet human dietary requirements for the rapidly growing global population (Yang et al., 2022). However, the industry faces substantial challenges owing to pathogenic infections and outbreaks caused by viruses and bacteria (Yang et al., 2022). Since 2008, there has been a global increase in oyster mass mortality events, which has substantially decreased *C. gigas* production in Europe and Australia (Barbosa-Solomieu et al., 2015; Fuhrmann et al., 2019). For example, Pacific oyster production in New South Wales (Australia) decreased by 48% within three years of the first OsHV-1 outbreak. Decreases in production can cause significant increases in prices. In France, the price of oysters increased from € 4.93/kg in 2001 to € 7.78/kg in 2012, when the surviving spat from a 2008 mortality event reached market size (Fuhrmann et al., 2019).

Oysters do not possess adaptive immunity, and so resistance to pathogens cannot be enhanced through vaccination, but rather by methods such as immunostimulation, diet, and genetic selection (Divilov et al., 2021; Robinson & Green, 2020; Yang et al., 2022). Furthermore, oysters are typically cultivated in open environments within straits and bays, with complete contact with flow-through seawater, making it impossible to fully control diseases through the administration of disinfectants, antibiotics, or similar treatments. Quarantining is not feasible, and therefore, large-scale pathogen outbreaks can occur. Genetic improvement through selective breeding is considered to be the most practical approach to enhance disease tolerance and resistance (Green et al., 2023; Zhai et al., 2021).

Ostreid herpesvirus 1 (OsHV-1) is a pathogen that can cause large-scale mortality events in Pacific oysters. In 2008, a highly virulent form of OsHV-1 termed OsHV-1 microvariant (µVar), was detected in France (Segarra et al., 2010). Outbreaks of novel OsHV-1 µVars have subsequently occurred in Italy, the United Kingdom, Australia, New Zealand and the United States (de Lorgeril et al., 2018; Jenkins et al., 2013; Keeling et al., 2014; Burge et al., 2021; Divilov et al., 2023b). OsHV-1 affects oyster immunity and allows opportunistic bacteria, such as *Vibrio* species, to infect hosts (Siboni et al., 2024). OsHV-1 can infect all oyster life stages, especially larval and juvenile stages, resulting in up to 100% mortality (de Lorgeril et al., 2018). Therefore, substantial efforts have been made to limit the spread of OsHV-1. In 2018, a novel OsHV-1 µVar was detected in San Diego Bay, California, USA, posing a significant risk to oyster aquaculture on the west coast of North America (Burge et al., 2021; Thompson et al., 2024). Recent research

using high-throughput sequencing indicated that this strain is different from the µVar detected in France as

it has a 6 bp deletion in the microsatellite region instead of a 12 bp deletion, along with other differences

(Pelletier, 2023). Given the threat posed by OsHV-1, substantial efforts have been made to implement

strategies to mitigate the spread of the virus.

Extensive efforts in selective breeding programs have improved the OsHV-1 survivability of Pacific

oysters in the USA and Australia as a potential mitigation strategy against OsHV-1 related mortality

(Divilov et al., 2023; Kube et al., 2018). A quantitative trait locus (QTL) on chromosome 8 (Chr8) was

identified as associated with an upregulated interferon regulatory factor (IRF) antiviral gene (i.e., *irf2*), and

this is thought to have increased resistance to OsHV-1 infections (Divilov et al., 2021). Oysters with the

alternate allele of this locus (hereafter referred to as the Chr8 marker) may be genetically primed to defend

against viral infections. The Chr8 marker has been used in marker-assisted selection using a rhAmp

genotyping assay (Beltz et al., 2018; Divilov et al., 2023), a fluorescence-based PCR method used to detect

specific alleles at a locus. Selectively bred families carrying the Chr8 marker were produced by the

Molluscan Broodstock Program (MBP) of Oregon State University. However, selective breeding may lead

to fitness trade-offs (Divilov et al., 2024). For example, increased OsHV-1 resilience in offspring of oysters

primed with an OsHV-1 viral mimic has been reported to co-occur with higher *Vibrio* infection levels in

offspring (Robinson & Green, 2020). OsHV-1 resilience has also been correlated with decreased larval size

and altered microbiome composition. Recently, families produced by the MBP via marker-assisted selection

to produce either heterozygous or homozygous reference genotypes at the Chr8 marker were shown to have

increased survivorship in heterozygous larvae and spat to *V. coralliilyticus* (Lunda et al., 2026).

Collectively, this raises the question as to whether oysters bred for increased survival to OsHV-1 may have

increased or decreased susceptibility to other bacterial pathogens, such as *Vibrio aestuarianus*.

Bacteria from the genus *Vibrio* are prevalent in marine environments, and several pathogenic strains

threaten vertebrates and invertebrates (Wendling et al., 2014). *Vibrio* species can occupy a variety of niches,

and include forms that are free-living, mutualistic, opportunistic, or pathogenic. Some *Vibrio* spp. can result

in vibriosis, a disease that can cause mass oyster mortalities. Common species of *Vibrio* that are pathogenic

to oysters include *V. splendidus*, *V. anguillarum*, *V. tubiashii* and *V. aestuarianus* (Wendling et al., 2014). In

Europe, *V. aestuarianus* has caused recurrent mortalities since 2001, first in France and then in other

European countries, including Italy, Ireland, and Spain (Mesnil et al., 2023). In recent years, the frequency

of mortality events in France has increased from 30% in 2011 to 77% in 2013, when it has emerged as the

primary pathogen identified during summer mortality events (Coyle et al., 2023). In British Columbia (BC),

Canada, *V. aestuarianus* is one of the most common pathogenic *Vibrio* species and is emerging as a serious

threat to oyster aquaculture of spat and adult *C. gigas* (Cowan et al., 2024). Considering the potential for

the application of marker-assisted selection using the Chr8 marker on the west coast of North America and

given that the effects of the Chr8 marker OsHV-1 survivorship allele on *V. aestuarianus* infection remain

unknown, further investigation is required. This is particularly relevant considering the recent findings of

an increase in resistance to oysters carrying the alternate allele of the Chr8 marker against *V. coralliilyticus*

(Lunda et al., 2026). Additionally, although loci associated with increased survival to *V. alginolyticus* in

Pacific oysters have been identified (Yang et al., 2022), the identification of loci associated with increased

survival to *V. aestuarianus* would also be beneficial for selective breeding purposes.

In the present study, we evaluated the effect of allele presence and allele dosage of the Chr8 marker

on the susceptibility to *V. aestuarianus* infection in a laboratory challenge. Given the functional differences

in immunological responses to viruses and bacteria (Green et al., 2016), we hypothesized that there may be

immunological trade-offs between resistance to viral and bacterial pathogens that may impact the resistance

of oysters with the OsHV-1 survivorship allele to a bacterial infection. However, we may also see an

increase in resistance, as was observed by Lunda et al. (2026) in infection trials with *V. coralliilyticus*. In

either case, it will be important to characterize the response of oysters carrying the Chr8 marker alternate

allele to *V. aestuarianus* infection. Furthermore, using the phenotypic data and samples from the laboratory

trial, we performed double digest RAD-sequencing (Peterson et al., 2012) to conduct a genome-wide

association study (GWAS) for *V. aestuarianus* survivorship.

Methods

Samples and crosses

Pacific oyster families with various genotypic states of the Chr8 marker (i.e. position 9,719,736 on Chr8;

Divilov et al., 2023) were generated by Oregon State University's Molluscan Broodstock Program (MBP;

Oregon, USA). Spat from 16 families were provided to Vancouver Island University (VIU) to be used in

experimental exposures to *V. aestuarianus*. Oysters (18 weeks old; > 4 mm long and < 6 mm wide) were

transported to the Deep Bay Marine Field Station of VIU (Bowser, British Columbia (BC), Canada), held

overnight in 12.5 °C flow-through seawater tanks, then transported to the Center for Shellfish Research

(CSR) of VIU (Nanaimo, BC). Seawater and algae (*Chaetoceros muelleri*) samples were collected from

Deep Bay Marine Field Station and transported to the VIU campus (Nanaimo).

Four of the 16 families had parents that were both heterozygous for the Chr8 marker (i.e., families

F114-F117). The other 12 families had at least one parent with a homozygous reference genotype at the

Chr8 marker locus (i.e., F101-F103, F105-F113; Table S1). These 12 families were the same families that

were tested for resistance or susceptibility to *V. coralliilyticus* by Lunda et al. (2026). No families were

produced by crosses with parents that were both homozygous for the alternate allele (i.e., the allele

associated with increased survivorship to OsHV-1). Additionally, one family from the VIU oyster breeding

program (Oyster Family Run 6, family 10; OFR6-10) was also included in the *Vibrio* challenge. OFR6-10

has not been selected for OsHV-1 resistance.

Disease challenge with Vibrio aestuarianus

In western Canada, *V. aestuarianus* can cause farm mortality in juvenile and adult oysters during periods of

elevated seawater temperatures (Cowan et al., 2024). Current laboratory methods for disease challenges in

shellfish include injection, co-habitation, and bath treatments. Injecting oysters in the adductor muscle with

a known amount of *V. aestuarianus* has been shown to provide reliable results (Green et al., 2016), but this

method by-passes natural barriers of immunity (De Decker et al., 2011). Co-habitation potentially mimics

a natural infection in the field (Dégremont et al., 2020, 2021) but this method can be unreliable in the timing

and magnitude of mortality compared to injection or bath exposures (Nordmo et al., 1997). Therefore, here

we conducted a bath exposure following procedures outlined by (Mackenzie et al., 2022) with a minor

modification to include one oyster per well in the tissue culture plate to ensure that oysters are exposed to

the same concentration of *V. aestuarianus*. It is worth noting here that factors known to affect disease risk

to *V. aestuarianus* include oyster age, size, and exposure dose (Azema et al., 2015; Travers et al., 2017).

Forty-eight *C. gigas* from each of the 16 MBP families and the VIU family (total n = 816) were

exposed to *V. aestuarianus*. An additional mixture of samples (n = 48) from multiple families was used as

a non-exposed control group, treated in the same way as the exposed oysters but without exposure to *Vibrio*.

For the *V. aestuarianus* exposure, *C. gigas* were placed in 12 well plates, where each well contained 3 mL

seawater and one oyster, and then plates were acclimatized overnight in a temperature-controlled room at

16 °C.

A *Vibrio* suspension was prepared by inoculating *V. aestuarianus* (strain: BS032 2018) in two flasks

containing TSB + 2% NaCl overnight in an orbital shaker (23 °C and 200 rpm). The cultures were then

removed from the incubator, transferred into Falcon tubes, and centrifuged (5000 x g) at ambient

temperature for 5 min. The supernatant was discarded and the cell pellets were resuspended in seawater.

The optical density of the resuspended cells was measured at 600 nm using an BioSpectrometer and

µCuvette® G1.0 (Eppendorf). A standard curve was prepared to estimate the amount of *V. aestuarianus* in

the preparation. The concentration of the *V. aestuarianus* inoculum was also estimated by serial dilution

and plating on tryptone soy agar with 2% NaCl, incubation at 23 °C for 48 h, followed by enumeration of

colonies.

The exposure trial was initiated by adding 300 µL of the *V. aestuarianus* suspension to each well

on each experimental plate (final concentration = 2.45×10^7 CFU per ml). Sterile seawater was added to

the control plate in place of the *V. aestuarianus* suspension. The temperature of the room was then increased

to 23 °C for five days to simulate a marine heatwave event, and the temperature was reduced to 16 °C on

the sixth day of exposure (Green et al., 2019). After *Vibrio* inoculation, 100 µL of algae (*C. muelleri*) was

added to each well of all plates to encourage oyster feeding and uptake of *V. aestuarianus*. Morbidity was

measured daily for six days, after which time the disease challenge ended. Oysters were considered

moribund or deceased when their shell valves were gaping and failed to close after gently touching the

oysters with a sterile toothpick. Oysters were placed into 25 mL Falcon tubes containing 100% ethanol,

where whole individuals were pooled in storage tubes labeled with day of death and family. After six days,

surviving oysters were preserved and stored in 100% ethanol. The ethanol-preserved oysters were stored at

room temperature until DNA extraction was conducted. Mortality data were analyzed using IBM SPSS

Statistics v28.0.1.0 (142) (IBM Corp., Armonk, NY) and R (R Core Team, 2026). Mortality distributions

were assessed using a Kaplan-Meier survival analysis. A log-rank test was conducted to determine the

differences in survival distributions among families, and a pairwise log-rank test was performed to

determine which families had different survival distributions.

Genomic DNA extraction

DNA extractions were performed on oysters from families F114-F117 (i.e., those with both parents

heterozygous for the Chr8 marker) to extract total genomic DNA using a Monarch® Genomic DNA

Purification Kit (#T3010L) according to manufacturer instructions for extractions from tissue. Briefly,

oyster spat were removed from their shells using sterilized forceps, rinsed with deionized water to remove

residual ethanol, then placed in a digestive solution containing lysis buffer and proteinase K (NEB).

Samples were homogenized using a sterile pestle and digested at 56 ºC for 2 h, after which an RNase

incubation was performed. Genomic DNA was purified through an extraction column as per manufacturer's

guidelines and eluted in 35 µL molecular-grade water held at 60 ºC. DNA quantification was performed via

absorbance and Qubit fluorimetry using a BioSpectrometer and µCuvette® G1.0 (Eppendorf). Samples

were checked for purity using A260/A280 and A260/A230 ratios on with the BioSpectrometer, and for

quality using 1% agarose gel electrophoresis (see Supplemental Information). Extracted genomic DNA was

then stored at -20 ºC.

Chr8 marker OsHV-1 resistance allele dosage

The rhAmp assay (Design ID: CD.GT.ZKJS3200.1; Integrated DNA Technologies) targeting a SNP with

G/A (REF/ALT) alleles at position 9,719,736 bp on Chr8 (Divilov et al., 2023) was used to genotype the

OsHV-1 resistance locus (i.e., Chr8 marker) in all individuals from the *V. aestuarianus* challenge trial in

families F114-F117. The assay was conducted on a CFX96 Touch Real-Time PCR Detection System (BIO-

RAD) using the manufacturer's instructions for a 5 µL reaction volume in a 96-well plate (Table S2).

Samples were normalized to 5 ng/µl in molecular-grade water and 10 ng (2 µL) of each sample added to

each well, where each sample was run in duplicate. A negative control well was added to each plate, where
instead of sample DNA it contained only molecular-grade water. The assay was previously confirmed to
work on gBlocks as per manufacturer's recommendations, and samples with known genotypes were used
as positive controls in this study. The fluorophore dyes FAM and VIC were used for genotyping the
reference (REF) and alternate (ALT) alleles, respectively, and to call genotypes as homozygous reference,
heterozygous, or homozygous alternate.

Results of the rhAmp assay were exported from the instrument in text format and processed in R
v4.3.2 using a custom script (see *Data Availability*). As some likely false positives were observed, where
one dye was detected at a much later Cq value than the other detected dye, a correction was applied in R to
convert the suspected false-positive heterozygote to a homozygote. Specifically, Cq differences between
FAM and VIC fluorophores were calculated per well, and any cases where the absolute value of the Cq
difference between the two fluorophores was greater than four, the call was determined to be a false positive
for the later Cq fluorophore, which was then set to zero, making the call a homozygote for the earlier
fluorophore (Figure S1 and Figure S2). Using the corrected data, genotypes were assigned based on the
retained fluorophore calls: FAM only = homozygous reference; VIC only = homozygous alternate; both
FAM and VIC = heterozygote; no fluorophore call = missing. Technical reproducibility was evaluated at
the genotype level (Table S3), and the final call for technical replicates was based on the majority call, as
inconsistent calls were re-run. Genotyping results were plotted in R (see *Data Availability*).

Statistical analysis of the Chr8 marker genotypes was carried out using IBM SPSS Statistics
v28.0.1.0 (142) and R. The effect of the Chr8 marker genotype on mortality in the *Vibrio* challenge was
determined using a chi-square test of homogeneity (or a Fisher's exact test if the chi-square assumptions
could not be met). To use the chi-square test, samples had to meet the following criteria: one dependent
variable measured at the dichotomous level (mortality: dead vs. alive or time until death), one independent
variable with three or more categorical groups (genotype: *AA, Aa, aa*), and independence of observations.
Finally, the sample had to meet a minimal sample size requirement (minimal sample size for calculating
expected frequencies required an expected count ≥ 5). If samples did not meet the sample size criterion, a
Fisher's exact test was performed. Baseline genotype frequencies in each family of F114-F117 prior to the
Vibrio challenge were also assessed using this method.

Genotyping by ddRADseq
Samples from families F114-F117, their respective parents, and OFR6-10 (VIU) were used for double-
digest restriction site-associated DNA sequencing (ddRADseq) at the Institut de Biologie Intégrative et des
Systems (IBIS; Quebec, Canada) using *Nsi*I/ *Msp*I cutting enzymes, as previously described (Sutherland et
al., 2020). Samples were pooled to sequence 32 oysters per chip on an Ion Torrent Proton using a

ThermoFisher Ion PI Chip Kit v3 chip, as per manufacturer's instructions. Base calling was performed

using Torrent Suite 5.10.1 (ThermoFisher). Multiplexed FASTQ files with sample identifying barcodes

present were exported using the FileExporter plugin.

Single-end read processing and genotyping was conducted using the repository *stacks_workflow*

(Normandeau, 2013/2025) and Stacks v.2.3e (Rochette et al., 2019), accompanied by custom analysis

scripts (see *Data Availability*). FASTQ file quality was assessed before and after trimming using FastQC

v0.11.8 with results consolidation by MultiQC (Andrews, 2010; Ewels et al., 2016). Universal adapters

were eliminated and reads longer than 50 bp were retained using cutadapt v1.18 run in parallel (Martin,

2011; Tange, 2015). Quality trimming was performed on all reads to retain Phred >10 in sliding windows

of size 15% of the read length, followed by read truncation to 80 bp and sample demultiplexing using the

process_radtags module of Stacks2 (Catchen et al., 2011; Rochette et al., 2019), with two enzymes selected

(i.e., *Nsi*I and *Msp*I) in parallel.

Demultiplexed and trimmed reads were aligned to a chromosome-level genome assembly for

Pacific oyster (i.e., GCF_902806645.1_cgigas_uk_roslin_v1_genomic.fna) (Peñaloza et al., 2021) using

bwa mem v0.7.12-r1039 (Li et al., 2009; Peñaloza et al., 2021). SAMtools v1.9 was used to remove

unmapped reads and secondary or low-quality alignments (Li et al., 2009). The remaining reads following

quality control were genotyped using Stacks2. The number of reads and alignments per sample, as well as

alignment rates were calculated using SAMtools and custom scripts.

RAD-loci were identified with Stacks2 using the reference-based approach through the *gstacks*

module with default settings. Genotype data were filtered using the *populations* module of Stacks2,

requiring that a locus was genotyped in at least 70% of the individuals per population in all six populations

(i.e., F114-F117, parents, and OFR6.10), and with a minimum global minor allele frequency (MAF) of 0.01.

Samples with fewer than 1 M reads were identified and removed from the analysis, as previously described

(Sutherland et al., 2020). Samples were genotyped again using *gstacks* as described above, and outputs

were used for downstream analysis. Single SNP per RAD-tag data were exported using the *populations*

module as a VCF file. The VCF file was imported into R using vcfR (Knaus & Grünwald, 2017) and

converted to genind format for use with adegenet functions (Jombart & Ahmed, 2011). Functions from the

simple_pop_stats repository were employed for analysis (see *Data availability*). The genotyping rate per

sample was determined and individuals with less than 30% missing data were retained. Minor allele

frequency (MAF) was re-calculated per locus, and loci with MAF < 0.01 were excluded. H_{OBS} and Hardy–

Weinberg (HW) equilibrium filters were not applied due to anticipated family effects. A principal

components analysis (PCA) based on genotypes was conducted using the *glPca* function of adegenet.

GWAS for survivorship to V. aestuarianus *challenge*

A GWAS for survivorship in the *V. aestuarianus* challenge was conducted using GEMMA v0.98.5 (Zhou
& Stephens, 2012) using the single-SNP per locus VCF file, as described above, converted to GEMMA
input format in R (see *Data Availability*). NCBI chromosome identifiers were converted to linkage groups
using the Pacific oyster assembly *Cgigas_uk_roslin_v1,* and linkage groups were converted to
chromosomes following Peñaloza et al. (2021). A GWAS analysis was conducted using unimputed
genotypes, as well as using genotypes that were imputed per family using the *impute_mean* function of the
missMethods R package (Rockel, 2022). Only families F114-F117 were used for the GWAS (excluding the
VIU family OFR6-10). The GEMMA analyses were conducted on families F114-F117 together by using
GEMMA to calculate kinship coefficients and using either a binary phenotype (i.e., dead or alive) or a
phenotype of days-to-death, where survivors of the trial were given the value of day 7 (i.e., one day after
the trial concluded). A Bonferroni-corrected *p*-value of 0.05 was used as the genome-wide significance
threshold. The R package *fastman* (Paria et al., 2022) was used to generate Manhattan plots from the GWAS
analysis.

Results

Baseline genotype frequencies of Chr8 marker
Prior to the disease challenge, the Chr8 marker genotype frequencies for families F114-F117 differed from
expected Mendelian proportions considering all heterozygous parents (i.e., 1:2:1 for REF/REF, REF/ALT,
and ALT/ALT, respectively; Figure 1). Considering averages across the four families, heterozygotes were
most frequently observed (57%), followed by the homozygous reference (25%), and the homozygous
alternates were the rarest (18%). When inspecting individual families (Figure 1), a skew was observed for
ALT/ALT genotypes where family F115 had significantly more than expected (35%), and F114 and F117
had significantly fewer (11% and 7%, respectively; χ^2 test of homogeneity p<0.05). No other genotypes
were significantly different among the families (Table S4).

V. aestuarianus *disease challenge of all families*
No mortality was observed in the first two days of the trial. Considering all the exposed, selectively bred
families together (i.e., excluding the VIU breeding program OFR6-10), 37.8% of oysters died on day 3, and
67.2% died by day 6 (n = 516) (Figure S3). The remaining 252 (32.8%) survived the disease challenge. The
multiple-family unexposed control group exhibited 100% survival.
The per-family cumulative mortality was on average (± s.d.) 67.2 ± 9.5% (Figure 2). Family F106
exhibited the highest mortality (85.4%), and F116 the lowest (47.9%). Rates of mortality also differed
among families (p<0.001; Figure 3); F106 died more rapidly than F103, F113, F114, F115, and F116
(pairwise comparison p < 0.003; Table S5; Bonferroni correction significance threshold p = 0.003). F106

offspring were all heterozygous for the Chr8 marker, and F103 and F113 were all homozygous reference

(Table S1), but when grouping families by parental genotypes at the Chr8 marker, there was no consistent

cumulative mortality trend (Figure 2) and substantial inter-family variation was observed.

Effects of Chr8 marker allele presence and dosage on V. aestuarianus *challenge survival*

Families F114-F117 had offspring with all genotypic states of the Chr8 marker, and therefore allowed the

investigation of the effects of the presence and dosage of the OsHV-1 resistance allele on mortality (days

to death, Figure 4; survival, Figure 5). No significant associations were observed between offspring

genotype and mortality for F114-F117 considering days to death, nor dead or alive phenotype (p>0.05).

The analysis was conducted for each family individually to avoid potential confounding interfamily effects.

ddRAD-seq genotyping of Vibrio-exposed families

Oysters from families F114-F117 and OFR6-10 (VIU family) from the *Vibrio* challenge were genotyped by

ddRADseq (avg. oysters per family = 36.4). The parents of MBP families were also sequenced (Table S6).

On average, 92.8 M single-end reads were produced per chip from each of the six chips sequenced (n =

523.1 M reads total). Demultiplexing resulted in an average of 91.8% reads retained. On average (± s.d.),

$65.0 \pm 2.2\%$ of reads aligned against the chromosome-level reference genome (Peñaloza et al., 2021) and

340,291,536 alignments were used for genotyping with *gstacks*. In total, 277.6 M primary alignments

(81.6%) were retained (10.9% of alignments were discarded owing to insufficient MAPQ and 7.5% due to

excessive soft clipping). On average, 1.79 M alignments were obtained per sample (range: 0.265-4.681 M

records per sample). The removal of samples with fewer than 1 M reads resulted in the removal of 18 of

the 190 samples, retaining 172 samples (Table S6). Notably, 15 of these 18 removed samples were oysters

that had died during the trial, suggesting that these samples may have had substantial DNA degradation

occurring between the mortality event and sampling, which has also seen in other disease challenge trials

in Pacific oysters (Sutherland et al., 2025).

After low-yield samples were removed, genotyping used 330.6 M alignments (269.7 M primary

alignments) to build 393,153 RAD loci with an average (± s.d.) effective coverage of $56.9 \pm 22.6x$. Low

genotyping rate loci were removed (see Methods), as were those with MAF < 0.01, resulting in the retention

of 10,202 RAD loci comprising 889,291 genomic sites, of which 25,801 contained variants (percent

polymorphic = 2.9%). The exported VCF file with a single SNP per RAD locus retained resulted in 8,252

variants for downstream analysis. Low genotyping rate samples (missing ≥ 30% of loci) were removed,

retaining 168 individuals (Figure S4; Table S6), including 32, 36, 35, 33, and 25 individuals from F114,

F115, F116, F117, and OFR6-10, respectively, as well as seven of the eight parents of families F114-F117.

Reapplying the MAF filter resulted in the retention of 8,193 SNPs (4,218 with MAF > 0.1). Clustering

samples based on genotypes by PCA showed close clustering of samples within the four MBP families,

with the parents of these families interspersed among the offspring and the VIU family OFR6-10 grouping

together but apart from the MBP families on PC1 (Figure S5). One individual labeled as F116 clustered

within the F115 family, suggesting a sample labelling or handling error (Figure S5).

GWAS for V. aestuarianus *survival*

Selectively bred oysters (families F114-F117) were used in GWAS analyses considering survival to the *V.*

aestuarianus challenge as a binary response (dead or alive) or days-to-death. This included 49 oysters that

survived and 87 that died (total n = 136). Considering only loci on chromosomes (n = 7,870 SNPs), without

imputation 3,656 SNPs were analyzed by GEMMA with a MAF > 0.05 filter applied. The percent variation

explained (PVE) ± standard error (S.E.) was 19.2 ± 14.7% for survival as a binary trait and 6.3 ± 9.8% for

days post-exposure survival. There were no significantly associated loci to survivorship to the *Vibrio*

challenge considering survival as dead or alive, or as days-to-death (Figure 6). Similarly, no significant

associations were identified using the day-to-death phenotype (Figure S6A). Using mean imputed

genotypes, 5,478 of the 7,870 chromosomal SNPs were analyzed by GEMMA (MAF > 0.05), resulting in

similar PVE values (PVE ± S.E. for dead/alive variable = 18.1 ± 14.8% and for days-to-death = 5.7 ± 9.6%).

The GWAS with imputed genotypes also did not identify any loci significantly associated with the

phenotype survival or days-to-death (Figure S6B and Figure S6C).

Discussion

Host genetic and epigenetic factors are known to influence disease dynamics (Krickeberg et al., 2012;

Magwire et al., 2012). In this study, we assessed whether the presence of a marker associated with increased

survival and resistance to a virus (OsHV-1) affects susceptibility to a bacterial pathogen (*V. aestuarianus*).

Our results indicate that the presence and allelic dosage of the OsHV-1 resistance allele of the Chr8 marker

(Divilov et al., 2023) in selectively bred families of Pacific oysters does not affect the susceptibility of the

juvenile oysters (spat) to *V. aestuarianus* infection. This contrasts with recent findings that the marker in

many of these same families is associated with increased survivorship to *V. coralliilyticus* infection (Lunda

et al., 2026). The cause of the different responses to the two bacteria species is not known, since both *V.*

coralliilyticus and *V. aestuarianus* would potentially be present in the environment in which the field trials

for survivorship occurred, and therefore the selective advantage of the surviving oysters could have

captured increased resistance to the bacteria in addition to the endemic OsHV-1, as discussed for *V.*

coralliilyticus by Lunda et al. (2026).

Juvenile Pacific oysters are susceptible to *V. aestuarianus* in British Columbia (BC), Canada

(Cowan et al., 2024). The present results and the results of Lunda et al. (2026) provide important insights

for selective breeding for disease resistance in Pacific oysters. The present results suggest a lack of
pleiotropy in the Chr8 marker for *V. aestuarianus* resistance and an absence of genetic correlation between
survivorship to OsHV-1 and *V. aestuarianus*, at least in regard to the survivorship mechanism associated
with the Chr8 marker. The results of Lunda et al. (2026) indicate that there should be no negative
consequence of using the Chr8 marker for marker-assisted selection on the resistance of the selected oysters
for *V. coralliilyticus*, and in contrast should improve their resilience to that bacterial pathogen. Given the
independence observed between the Chr8 protective allele and *V. aestuarianus* resistance, identifying loci
involved in resistance to *V. aestuarianus* may enable marker-assisted selection of oysters that are resistant
to both OsHV-1 (and *V. coralliilyticus*; Lunda et al., 2026) and *V. aestuarianus*. However, in the present
study we were not able to identify any major effect loci for resistance to *V. aestuarianus* and rather identify
that resistance may be controlled by polygenic architecture.
395
Effects of the OsHV-1 resistance allele on survival to Vibrio aestuarianus *infection*
The presence of the Chr8 marker allele associated with increased resistance to OsHV-1 had no effect on the
survival to *V. aestuarianus* infection. This finding is supported by another study that reported the absence
of genetic correlations between OsHV-1 and *V. aestuarianus* resistance (Azéma et al., 2017). Azéma et al.
(2017) analyzed the genetic parameters for resistance to OsHV-1 and *V. aestuarianus* at three different life
stages: spat (3 to 6 months old); juvenile (11 to 15 months old); and adult (25 to 26 months old). They
reported no observable genetic correlation between survival to viral or bacterial infection. However, the life
stage of the oyster significantly affected susceptibility to both pathogens; susceptibility to OsHV-1
decreased with age and growth, whereas susceptibility to *V. aestuarianus* increased with age and growth.
The different effects of age on susceptibility to OsHV-1 and *Vibrio* may reflect the different immune
responses of oyster hosts. Further evidence of the independence of the resistance mechanisms was provided
by Green et al. (2016), who found different transcriptome responses to OsHV-1 and *V. aestuarianus*
infections. Thus, the immune genes involved in the defence against OsHV-1 and *Vibrio aestuarianus*
infections appear to be distinct.
The absence of a correlation between the Chr8 marker and *V. aestuarianus* susceptibility suggests
that the mechanisms of resistance to these pathogens are independent. The Chr8 marker allele associated
with increased survivorship is thought to upregulate *interferon regulatory factor 2* (*irf2*), an antiviral gene
that can provide oysters with constitutively upregulated antiviral immunity (Divilov et al., 2023). With
constitutively overexpressed *irf2*, overexpression of antiviral genes could be expected even in the absence
of the virus. Therefore, the lack of increased susceptibility to *V. aestuarianus* is notable, as it suggests that
the upregulated antiviral immunity does not impact (negatively or positively) the oyster's ability to survive
the bacterial infection. However, the finding of increased resistance to *V. coralliilyticus* (Lunda et al., 2026)

suggests that this is not a universal conclusion, and that the mechanism of resistance does provide resistance

to a bacteria pathogen (i.e., *V. coralliilyticus*). The genetic mechanisms underlying *Vibrio* resistance are not

well known. However, a recent study using 52 Pacific oyster families found 18 SNPs in three QTL

associated with resistance to *V. alginolyticus* in *C. gigas*, and the authors suggested that *Vibrio* resistance is

polygenic (Yang et al., 2022). Most of the associated genes are related to pathogen recognition and immune

defence responses (Yang et al., 2022). Collectively, genes that induce the expression of immune-related

genes are potential targets for selective breeding to increase *Vibrio* resilience.

Selective breeding for one trait can lead to fitness trade-offs by negatively affecting a second trait

and consequently impacting overall fitness (Divilov et al., 2024; Dwivedi et al., 2021). Our understanding

of the compromises in selection for disease resistance is limited, particularly regarding the effects on other

traits or resistance to other pathogens. Trade-offs can arise when beneficial traits are linked to detrimental

ones, thereby affecting fitness outcomes (Gallardo-Hidalgo et al., 2021). For example, although selection

for increased growth rate has also improved larval resilience to ocean acidification, fast-growing oysters

are more susceptible to marine heatwaves and *V. aestuarianus* infections later in life; mortality has been

found to increase by 76% per gram of body weight gained (Khtikian, 2021; Nordio et al., 2021). However,

selection for OsHV-1 resistance in the MBP breeding population has not been observed to impact growth

(Divilov et al., 2024). Selection for disease resistance is complex, as resistance is expected to be influenced

by multifactorial biological functions, including reproduction and immunity (Bai & Plastow, 2022; Wang

et al., 2021).

We expected that a trade-off could be observed in the present study because up-regulation of the

immune system can increase the energy requirements of the host (Lochmiller & Deerenberg, 2000;

Robinson & Green, 2020). This can reduce the overall reproductive success, growth rates, and survival to

events such as pathogen outbreaks (Lochmiller & Deerenberg, 2000; Robinson & Green, 2020).

Additionally, in some cases, over-reactive immune responses to pathogens can be more damaging than a

weak immune response (Green et al., 2016), as discussed in relation to immune tolerance (Råberg et al.,

2007). The expression of extremely potent antibacterial or antiviral effectors can result in a toxic cellular

environment that can be detrimental to the organism (Green et al., 2016), and strong immune effectors and

responses can inflict damage on host tissues. In the case of infectious diseases, a fitness trade-off might

occur if the heightened immune response leads to greater collateral damage (Råberg et al., 2007).

Furthermore, other environmental stress factors may impact immune responses due to energetic constraints

(Sutherland et al., 2014). Here, we did not measure other factors that may represent energetic trade-offs for

increased viral resistance (e.g., growth rate and response to other stressors), but these factors could be

integrated into future studies to evaluate the impact of the Chr8 marker on *Vibrio* resistance. Notably, before

the *V. aestuarianus* exposure occurred, there were baseline differences in the genotype frequencies of the

Chr8 marker, specifically in the proportion of the homozygous alternate genotype. Therefore, other

unknown factors may cause distortions in the frequency of this marker, which could affect future breeding

practices around the Chr8 marker.

Although we did not observe a trade-off, selection for disease resistance may also indirectly

increase resistance to other stresses such as marine heat waves (MHW). MHWs increase water temperature,

resulting in high mortality rates (Cowan et al., 2024). Vibriosis and *V. aestuarianus* proliferation are further

linked to seawater temperature, with increased temperature increasing host susceptibility by weakening the

immune system and increasing the transmission and proliferation of pathogenic *Vibrio* species (Cowan et

al., 2024; Wendling et al., 2014). MHWs are a consequence of climate change, which is driven by increased

anthropogenic emissions of CO_2. The global average MHW frequency and duration increased by 34% and

17%, respectively, resulting in a 54% increase in annual marine heatwave days globally (Oliver et al., 2018).

Bolstering resilience to MHWs may be achieved by producing oysters with increased resistance to viral and

bacterial pathogens. This can mitigate the detrimental effects of these temperature spikes on oyster

populations. This dual-pathogen resistance strategy could serve as a multifaceted approach to bolster oyster

health and ensure the sustainability of oyster aquaculture in the presence of various environmental stressors.

Family effects and GWAS for V. aestuarianus *resistance*

In this study, we also analyzed family differences in survival following *V. aestuarianus* infection, both in

cumulative mortality and mortality rates (days to death). Although different families have known OsHV-1

genotypes, they also would have other genetic differences that could affect resistance to *V. aestuarianus*.

As the Chr8 marker did not explain the variance observed across families in *Vibrio* susceptibility, other

factors may be involved.

In the GWAS analysis for survivorship to *V. aestuarianus*, we did not identify any loci significantly

associated with mortality or mortality rate in response to infection. The applied sample size may have been

too small to detect a more subtle genetic association, such as that which would occur with a polygenic trait.

Resistance to *Vibrio* infections has previously been reported to exhibit a polygenic architecture (Yang et al.,

2022), and therefore may require a significantly larger sample size to identify significant associations. It is

also possible that inspecting other populations of Pacific oysters could identify major effect loci for *Vibrio*

survivorship that are not present in the MBP families investigated here. Furthermore, differences in

resistance across families could involve other determining factors, such as epigenetic differences impacting

immune responses to the bacterial infection. For example, a recent GWAS for survival to Pacific oyster

mortality syndrome indicated that epigenetic factors play a larger role than genetic factors for the syndrome

(Gawra et al., 2023). Future studies may benefit from including a larger number of individuals to account

for the potential polygenic nature of *Vibrio* resistance mechanisms, but also may need to consider that *Vibrio*

resistance as a trait could be best addressed by other selective breeding approaches, including genomic
selection (Boudry et al., 2021; Meuwissen et al., 2016), rather than marker-assisted selection, in part
because of the challenge in identifying major effect loci.

Conclusion

The genotype and allelic dosage of the Chr8 marker allele associated with survivorship to OsHV-1 did not
affect susceptibility to *V. aestuarianus* infection in selectively bred Pacific oyster spat evaluated here. This
suggests a lack of pleiotropic effect of the Chr8 marker on *V. aestuarianus* resistance, as well as a lack of
genetic correlation between these traits. Given the recent findings that the OsHV-1 resistance allele also
provides protection to *V. coralliilyticus*, this may suggest different immune pathways for defense against
these two *Vibrio* species. The GWAS analysis did not find any loci significantly associated with *Vibrio*
survivorship, suggesting a polygenic architecture in these families for this trait. Nonetheless, significant
family differences in *Vibrio* susceptibility were observed here, even within the MBP breeding program
families that have been selectively bred for at least six generations (de Melo et al., 2016) and therefore are
expected to be more genetically homogenous than outbred oysters. These observations emphasize the
potential for further investigation into selective breeding for resistance to *V. aestuarianus*, which may
benefit from genomic selection rather than marker-assisted selection, unless major effect loci can be
identified.

Supplementary Materials

506 Supplementary tables and figures are provided in Supplementary Results.
507 **Additional File S1.** Read counts and alignments for all samples retained in the analysis.

Data Availability

Raw sequencing data have been uploaded to SRA under BioProject PRJNA1145972 with BioSamples
SAMN43080749 - SAMN43080938.
RADseq genotypes and supporting files: 10.6084/m9.figshare.26524321
rhAmp assay raw data plate files: 10.6084/m9.figshare.26515438
Sample and family IDs for mapping families: 10.6084/m9.figshare.28599923

This analysis was supported by the following repositories:
Manuscript code and README: https://github.com/bensutherland/ms_cgig_chr8_vibrio
simple_pop_stats repository: https://github.com/bensutherland/simple_pop_stats

Funding

521   Funding for this study was provided by the Pacific States Marine Fisheries Commission (PSMFC) project

522   "Preparing for Future Challenges – Threats from Ocean Acidification, *Vibrio coralliilyticus* and OsHV-1

523   µvar to West Coast Oyster Farmers" (Award No. NA18NMF4720007).

Conflicts of Interest

526   BJGS is affiliated with Sutherland Bioinformatics. The other authors declare no conflicts of interest.

Acknowledgements

529   Thanks to the members of the Centre for Shellfish Research of Vancouver Island University for laboratory

530   assistance, to Antonia Barela, Gary Fleener, and William Schoeneck for assistance in the hatchery for the

531   production of juvenile oysters, and to the Deep Bay Marine Field Station staff for storing and acclimating

532   the juvenile oysters. Thanks to Brian Boyle at IBIS for ddRADseq library preparations and sequencing.

533   Thanks to three anonymous reviewers for comments on an earlier version of this manuscript.

Author Contributions

536   BS, LS, TG, and CL conceived of the study. LS, SL, TG conducted laboratory experiments. BS, LS, and

537   KD contributed to software development and analysis. BS curated the data. LS and BS drafted the

538   original manuscript. TG and BS provided laboratory and staff supervision. TG and CL acquired funding

539   for the work and administered the project. All authors (LS, BS, SL, AL, KD, CL, TG) supported result

540   interpretation and contributed to manuscript review and editing.

References

Andrews, S. (2010). *Babraham Bioinformatics—FastQC A Quality Control tool for High Throughput*
Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Azéma, P., Lamy, J.-B., Boudry, P., Renault, T., Travers, M.-A., & Dégremont, L. (2017). Genetic
parameters of resistance to *Vibrio aestuarianus*, and OsHV-1 infections in the Pacific oyster,
Crassostrea gigas, at three different life stages. *Genetics, Selection, Evolution : GSE*, *49*, 23.
https://doi.org/10.1186/s12711-017-0297-2

Azema, P., Travers, M.-A., Lamy, J.-B., & Degremont, L. (2015). Genetic parameters for OsHV-1 and
Vibrio aestuarianus resistance in *Crassostrea gigas*: First results using controlled challenges.
Journal of Shellfish Research, *34*(2), 606–607.

Bai, X., & Plastow, G. S. (2022). Breeding for disease resilience: Opportunities to manage polymicrobial
challenge and improve commercial performance in the pig industry. *CABI Agriculture and*
Bioscience, *3*(1), 6. https://doi.org/10.1186/s43170-022-00073-y

Barbosa-Solomieu, V., Renault, T., & Travers, M.-A. (2015). Mass mortality in bivalves and the intricate
case of the Pacific oyster, *Crassostrea gigas*. *Journal of Invertebrate Pathology*, *131*, 2–10.
https://doi.org/10.1016/j.jip.2015.07.011

Beltz, K., Tsang, D., Wang, J., Rose, S., Bao, Y., Wang, Y., Larkin, K., Rupp, S., Schrepfer, D., Datta, K.,
Gunderson, K., Sailor, C., Hansen, S., Dobosy, J., Lewis, L., Menezes, A., Walder, J., Behlke, M.,
& Chen, C. (2018). A high-performing and cost-effective SNP genotyping method using rhPCR
and universal reporters. *Advances in Bioscience and Biotechnology*, *09*, 497–512.
https://doi.org/10.4236/abb.2018.99034

Botta, R., Asche, F., Borsum, J. S., & Camp, E. V. (2020). A review of global oyster aquaculture
production and consumption. *Marine Policy*, *117*, 103952.
https://doi.org/10.1016/j.marpol.2020.103952

Boudry, P., Allal, F., Aslam, M. L., Bargelloni, L., Bean, T. P., Brard-Fudulea, S., Brieuc, M. S. O.,
Calboli, F. C. F., Gilbey, J., Haffray, P., Lamy, J.-B., Morvezen, R., Purcell, C., Prodöhl, P. A.,
Vandeputte, M., Waldbieser, G. C., Sonesson, A. K., & Houston, R. D. (2021). Current status and
potential of genomic selection to improve selective breeding in the main aquaculture species of
International Council for the Exploration of the Sea (ICES) member countries. *Aquaculture*
Reports, *20*, 100700. https://doi.org/10.1016/j.aqrep.2021.100700

Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., & Postlethwait, J. H. (2011). Stacks: Building and
genotyping loci *de novo* from short-read sequences. *G3 Genes|Genomes|Genetics*, *1*(3), 171–182.
https://doi.org/10.1534/g3.111.000240

Cowan, M. W., Pearce, C. M., Green, T. J., Finston, T., Meyer, G. R., McAmmond, B., Van Hamme, J. D.,
Bottos, E. M., Marshall, R., Evans, W., Sutherland, T. F., & de la Bastide, P. Y. (2024).
Abundance of *Vibrio aestuarianus*, water temperature, and stocking density are associated with
summer mortality of Pacific oysters in suspended culture. *Aquaculture International*.
https://doi.org/10.1007/s10499-024-01415-5

Coyle, N. M., O'Toole, C., Thomas, J. C. L., Ryder, D., Feil, E. J., Geary, M., Bean, T. P., Joseph, A. W.,
Waine, A., Cheslett, D., & Verner-Jeffreys, D. W. (2023). *Vibrio aestuarianus* clade A and clade B
isolates are associated with Pacific oyster (*Magallana gigas*) disease outbreaks across Ireland.
Microbial Genomics, *9*(8), mgen001078. https://doi.org/10.1099/mgen.0.001078

De Decker, S., Normand, J., Saulnier, D., Pernet, F., Castagnet, S., & Boudry, P. (2011). Responses of
diploid and triploid Pacific oysters *Crassostrea gigas* to *Vibrio* infection in relation to their
reproductive status. *Journal of Invertebrate Pathology*, *106*(2), 179–191.
https://doi.org/10.1016/j.jip.2010.09.003

de Lorgeril, J., Lucasson, A., Petton, B., Toulza, E., Montagnani, C., Clerissi, C., Vidal-Dupiol, J.,
Chaparro, C., Galinier, R., Escoubas, J.-M., Haffner, P., Dégremont, L., Charrière, G. M., Lafont,
M., Delort, A., Vergnes, A., Chiarello, M., Faury, N., Rubio, T., … Mitta, G. (2018). Immune-

suppression by OsHV-1 viral infection causes fatal bacteraemia in Pacific oysters. *Nature*
Communications, *9*(1), Article 1. https://doi.org/10.1038/s41467-018-06659-3

de Melo, C. M. R., Durland, E., & Langdon, C. (2016). Improvements in desirable traits of the Pacific
oyster, *Crassostrea gigas*, as a result of five generations of selection on the West Coast, USA.
Aquaculture, *460*, 105–115. https://doi.org/10.1016/j.aquaculture.2016.04.017

Dégremont, L., Azéma, P., Maurouard, E., & Travers, M.-A. (2020). Enhancing resistance to *Vibrio*
aestuarianus in *Crassostrea gigas* by selection. *Aquaculture*, *526*, 735429.
https://doi.org/10.1016/j.aquaculture.2020.735429

Dégremont, L., Morga, B., Maurouard, E., & Travers, M.-A. (2021). Susceptibility variation to the main
pathogens of *Crassostrea gigas* at the larval, spat and juvenile stages using unselected and
selected oysters to OsHV-1 and/or *V. aestuarianus*. *Journal of Invertebrate Pathology*, *183*,
107601. https://doi.org/10.1016/j.jip.2021.107601

Divilov, K., Merz, N., Schoolfield, B., Green, T. J., & Langdon, C. (2023). Marker-assisted selection in a
Pacific oyster population for an antiviral QTL conferring increased survival to OsHV-1 mortality
events in Tomales Bay. *Aquaculture*, *567*, 739291.
https://doi.org/10.1016/j.aquaculture.2023.739291

Divilov, K., Schoolfield, B., Mancilla Cortez, D., Wang, X., Fleener, G. B., Jin, L., Dumbauld, B. R., &
Langdon, C. (2021). Genetic improvement of survival in Pacific oysters to the Tomales Bay strain
of OsHV-1 over two cycles of selection. *Aquaculture*, *543*, 737020.
https://doi.org/10.1016/j.aquaculture.2021.737020

Divilov, K., Schoolfield, B., Merz, N., & Langdon, C. (2024). No trade-off between growth and OsHV-1
tolerance in Pacific oysters (*Crassostrea gigas*) farmed on the West Coast, USA. *Animal*
Genetics, *55*(1), 158–162. https://doi.org/10.1111/age.13375

Dwivedi, S. L., Reynolds, M. P., & Ortiz, R. (2021). Mitigating tradeoffs in plant breeding. *iScience*,
24(9), 102965. https://doi.org/10.1016/j.isci.2021.102965

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048.
https://doi.org/10.1093/bioinformatics/btw354

Fuhrmann, M., Castinel, A., Cheslett, D., Furones Nozal, D., & Whittington, R. j. (2019). *The impacts of*
ostreid herpesvirus 1 microvariants on Pacific oyster aquaculture in the Northern and Southern
Hemispheres since 2008. *38(2)*, 491. https://doi.org/10.20506/rst.38.2.3000

Gallardo-Hidalgo, J., Barría, A., Yoshida, G. M., & Yáñéz, J. M. (2021). Genetics of growth and survival
under chronic heat stress and trade-offs with growth- and robustness-related traits in rainbow
trout. *Aquaculture*, *531*, 735685. https://doi.org/10.1016/j.aquaculture.2020.735685

Gawra, J., Valdivieso, A., Roux, F., Laporte, M., de Lorgeril, J., Gueguen, Y., Saccas, M., Escoubas, J.-
M., Montagnani, C., Destoumieux-Garzón, D., Lagarde, F., Leroy, M. A., Haffner, P., Petton, B.,
Cosseau, C., Morga, B., Dégremont, L., Mitta, G., Grunau, C., & Vidal-Dupiol, J. (2023).
Epigenetic variations are more substantial than genetic variations in rapid adaptation of oyster to
Pacific oyster mortality syndrome. *Science Advances*, *9*(36), eadh8990.
https://doi.org/10.1126/sciadv.adh8990

Government of Canada, F. and O. C. (2018, December 4). *Herpes-Type Virus Disease of Oysters*.
Fisheries and Oceans Canada. https://www.dfo-mpo.gc.ca/science/aah-saa/diseases-
maladies/htvdoy-eng.html

Green, K. M., Spalding, A. K., Ward, M., Levine, A., Wolters, E. A., Hamilton, S. L., & Rice, L. (2023).
Oregon shellfish farmers: Perceptions of stressors, adaptive strategies, and policy linkages. *Ocean*
& Coastal Management, *234*, 106475. https://doi.org/10.1016/j.ocecoaman.2022.106475

Green, T. J., Siboni, N., King, W. L., Labbate, M., Seymour, J. R., & Raftos, D. (2019). Simulated marine
heat wave alters abundance and structure of Vibrio populations associated with the Pacific oyster
resulting in a mass mortality event. *Microbial Ecology*, *77*(3), 736–747.
https://doi.org/10.1007/s00248-018-1242-9

Green, T. J., Vergnes, A., Montagnani, C., & de Lorgeril, J. (2016). Distinct immune responses of juvenile and adult oysters (*Crassostrea gigas*) to viral and bacterial infections. *Veterinary Research*, *47*(1), 72. https://doi.org/10.1186/s13567-016-0356-7

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: New tools for the analysis of genome-wide SNP data. *Bioinformatics*, *27*(21), 3070–3071. https://doi.org/10.1093/bioinformatics/btr521

Khtikian, N. (2021). *Investigating the role of Vibrio aestuarianus in summer mortality of farmed Crassostrea gigas in Baynes Sound, British Columbia*. University of Victoria Libraries. http://hdl.handle.net/1828/13534

Knaus, B. J., & Grünwald, N. J. (2017). vcfr: A package to manipulate and visualize variant call format data in R. *Molecular Ecology Resources*, *17*(1), 44–53. https://doi.org/10.1111/1755-0998.12549

Krickeberg, K., Pham, V. T., & Pham, T. M. H. (2012). The Idea of Epidemiology. In K. Krickeberg, V. T. Pham, & T. M. H. Pham (Eds.), *Epidemiology: Key to Prevention* (pp. 1–10). Springer. https://doi.org/10.1007/978-1-4614-1205-2_1

Kube, P., Dove, M., Cunningham, M., Kirkland, P., Gu, X., Hick, P., O'Connor, W., & Elliott, N. (2018). *Genetic Selection for Resistance to Pacific Oyster Mortality Syndrome*. CSIRO Agriculture and Food, NSW Department of Primary Industries, Australian Seafood Industries, Seafood CRC, and FRDC.

Laurence, M., Rose-marie, L., & Philippe, G. (2009). *Alien Species Alert: Crassostrea gigas (Pacific oyster)*.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Lochmiller, R. L., & Deerenberg, C. (2000). Trade-offs in evolutionary immunology: Just what is the cost of immunity? *Oikos*, *88*(1), 87–98. https://doi.org/10.1034/j.1600-0706.2000.880110.x

Lunda, S. L., Divilov, K., Green, T. J., Sutherland, B.J.G., Wang, X., Ling, J., Jennings, J., Mandour, M., Mueller, R.S., & Langdon, C. (2026). Selection for a single nucleotide polymorphism on chromosome 8 of the genome of *Crassostrea gigas* improves resistance to both ostreid herpesvirus 1 and *Vibrio coralliilyticus*. *Aquaculture*, https://doi.org/10.1016/j.aquaculture.2025.743597

Mackenzie, C. L., Pearce, C. M., Leduc, S., Roth, D., Kellogg, C. T. E., Clemente-Carvalho, R. B. G., & Green, T. J. (2022). Impacts of seawater pH buffering on the larval microbiome and carry-over effects on later-life disease susceptibility in Pacific oysters. *Applied and Environmental Microbiology*, *88*(22), e01654-22. https://doi.org/10.1128/aem.01654-22

Magwire, M. M., Fabian, D. K., Schweyen, H., Cao, C., Longdon, B., Bayer, F., & Jiggins, F. M. (2012). Genome-wide association studies reveal a simple genetic basis of resistance to naturally coevolving viruses in *Drosophila melanogaster*. *PLoS Genetics*, *8*(11), e1003057. https://doi.org/10.1371/journal.pgen.1003057

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal*, *17*(1), Article 1. https://doi.org/10.14806/ej.17.1.200

Mesnil, A., Jacquot, M., Garcia, C., Tourbiez, D., Canier, L., Bidois, A., Dégremont, L., Cheslett, D., Geary, M., Vetri, A., Roque, A., Furones, D., Garden, A., Orozova, P., Arzul, I., Sicard, M., Charrière, G. M., Destoumieux-Garzón, D., & Travers, M.-A. (2023). Emergence and clonal expansion of *Vibrio aestuarianus* lineages pathogenic for oysters in Europe. *Molecular Ecology*, *32*(11), 2869–2883. https://doi.org/10.1111/mec.16910

Meuwissen, T., Hayes, B., & Goddard, M. (2016). Genomic selection: A paradigm shift in animal breeding. *Animal Frontiers*, *6*(1), 6–14. https://doi.org/10.2527/af.2016-0002

Nordio, D., Khtikian, N., Andrews, S., Bertotto, D., Leask, K., & Green, T. (2021). Adaption potential of *Crassostrea gigas* to ocean acidification and disease caused by *Vibrio harveyi*. *ICES Journal of Marine Science*, *78*(1), 360–367. https://doi.org/10.1093/icesjms/fsaa080

Normandeau, E. (2023). *STACKS Workflow2* [Python]. https://github.com/enormandeau/stacks_workflow
(Original work published 2013)
Okon, E. M., Birikorang, H. N., Munir, M. B., Kari, Z. A., Téllez-Isaías, G., Khalifa, N. E., Abdelnour, S.
A., Eissa, M. E. H., Al-Farga, A., Dighiesh, H. S., & Eissa, E.-S. H. (2023). A global analysis of
climate change and the impacts on oyster diseases. *Sustainability*, *15*(17), Article 17.
https://doi.org/10.3390/su151712775
Oliver, E. C. J., Donat, M. G., Burrows, M. T., Moore, P. J., Smale, D. A., Alexander, L. V., Benthuysen,
J. A., Feng, M., Sen Gupta, A., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E.,
Scannell, H. A., Straub, S. C., & Wernberg, T. (2018). Longer and more frequent marine
heatwaves over the past century. *Nature Communications*, *9*(1), 1324.
https://doi.org/10.1038/s41467-018-03732-9
Paria, S. S., Rahman, S. R., & Adhikari, K. (2022). *fastman: A fast algorithm for visualizing GWAS*
results using Manhattan and Q-Q plots. https://doi.org/10.1101/2022.04.19.488738
Pelletier, C. (2023). *Diversité et évolution de l'Ostreid Herpesvirus type 1* [Phdthesis, Université de La
Rochelle]. https://theses.hal.science/tel-04549017
Peñaloza, C., Gutierrez, A. P., Eöry, L., Wang, S., Guo, X., Archibald, A. L., Bean, T. P., & Houston, R. D.
(2021). A chromosome-level genome assembly for the Pacific oyster *Crassostrea gigas*.
GigaScience, *10*(3), giab020. https://doi.org/10.1093/gigascience/giab020
Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest RADseq:
an inexpensive method for *De Novo* SNP discovery and genotyping in model and non-model
species. *PLoS ONE*, *7*(5), e37135. https://doi.org/10.1371/journal.pone.0037135
R Core Team. (2026). *R: A Language and Environment for Statistical Computing* [Computer software].
https://www.r-project.org/
Råberg, L., Sim, D., & Read, A. F. (2007). Disentangling genetic variation for resistance and tolerance to
infectious diseases in animals. *Science*, *318*(5851), 812–814.
https://doi.org/10.1126/science.1148526
Robinson, A. N., & Green, T. J. (2020). Fitness costs associated with maternal immune priming in the
oyster. *Fish & Shellfish Immunology*, *103*, 32–36. https://doi.org/10.1016/j.fsi.2020.04.047
Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods for paired-
end sequencing improve RADseq-based population genomics. *Molecular Ecology*, *28*(21), 4737–
4754. https://doi.org/10.1111/mec.15253
Rockel, T. (2022). *missMethods: Methods for Missing Data* (Version 0.4.0) [Computer software].
https://cran.r-project.org/web/packages/missMethods/index.html
Segarra, A., Pépin, J. F., Arzul, I., Morga, B., Faury, N., & Renault, T. (2010). Detection and description
of a particular Ostreid herpesvirus 1 genotype associated with massive mortality outbreaks of
Pacific oysters, *Crassostrea gigas*, in France in 2008. *Virus Research*, *153*(1), 92–99.
https://doi.org/10.1016/j.virusres.2010.07.011
Siboni, N., King, W. L., Williams, N. L. R., Scanes, E., Giardina, M., Green, T. J., Ostrowski, M.,
O'Connor, W., Dove, M., Labbate, M., & Seymour, J. R. (2024). Increased abundance of
potentially pathogenic Vibrio and a marine heatwave co-occur with a Pacific oyster summer
mortality event. *Aquaculture*, *583*, 740618. https://doi.org/10.1016/j.aquaculture.2024.740618
Sutherland, B. J. G., Divilov, K., Thompson, N. F., Delomas, T. A., Lunda, S. L., Langdon, C. J., &
Green, T. J. (2026). Pedigree-based genome-wide imputation using a low-density amplicon panel
for the highly polymorphic Pacific oyster *Crassostrea* (*Magallana*) *gigas*. *Aquaculture*, 612(1),
743096.
Sutherland, B. J. G., Hanson, K. C., Jantzen, J. R., Koop, B. F., & Smith, C. T. (2014). Divergent
immunity and energetic programs in the gills of migratory and resident *Oncorhynchus mykiss*.
Molecular Ecology, *23*(8), 1952–1964. https://doi.org/10.1111/mec.12713
Sutherland, B. J. G., Rycroft, C., Ferchaud, A.-L., Saunders, R., Li, L., Liu, S., Chan, A. M., Otto, S. P.,
Suttle, C. A., & Miller, K. M. (2020). Relative genomic impacts of translocation history, hatchery

practices, and farm selection in Pacific oyster *Crassostrea gigas* throughout the Northern
Hemisphere. *Evolutionary Applications*, *13*(6), 1380–1399. https://doi.org/10.1111/eva.12965

Tange, O. (2015). *GNU Parallel 20150322 ('Hellwig')* [Computer software]. Zenodo.
https://doi.org/10.5281/zenodo.16303

Thompson, N. F., Agnew, M. V., Calla, B., & Burge, C. A. (2024). Assessing selection potential for
Pacific oyster (*Crassostrea gigas*) to a North American OsHV-1 µvar: Comparing two
experimental assay methods. *Aquaculture*, *590*, 741076.
https://doi.org/10.1016/j.aquaculture.2024.741076

Travers, M.-A., Tourbiez, D., Parizadeh, L., Haffner, P., Kozic-Djellouli, A., Aboubaker, M., Koken, M.,
Dégremont, L., & Lupo, C. (2017). Several strains, one disease: Experimental investigation of
Vibrio aestuarianus infection parameters in the Pacific oyster, *Crassostrea gigas*. *Veterinary*
Research, *48*(1), 32. https://doi.org/10.1186/s13567-017-0438-1

Wang, D., Loor, A., Bels, L. D., Stappen, G. V., den Broeck, W. V., & Nevejan, N. (2021). Dynamic
immune response to vibriosis in Pacific oyster *Crassostrea gigas* larvae during the infection
process as supported by accurate positioning of GFP-Tagged Vibrio strains. *Microorganisms*,
9(7), 1523. https://doi.org/10.3390/microorganisms9071523

Wendling, C. C., Batista, F. M., & Wegner, K. M. (2014). Persistence, seasonal dynamics and pathogenic
potential of Vibrio communities from Pacific oyster hemolymph. *PLOS ONE*, *9*(4), e94256.
https://doi.org/10.1371/journal.pone.0094256

Yang, B., Zhai, S., Zhang, F., Wang, H., Ren, L., Li, Y., Li, Q., & Liu, S. (2022). Genome-wide
association study toward efficient selection breeding of resistance to *Vibrio alginolyticus* in
Pacific oyster, *Crassostrea gigas*. *Aquaculture*, *548*, 737592.
https://doi.org/10.1016/j.aquaculture.2021.737592

Zhai, S., Yang, B., Zhang, F., Li, Q., & Liu, S. (2021). Estimation of genetic parameters for resistance to
Vibrio alginolyticus infection in the Pacific oyster (*Crassostrea gigas*). *Aquaculture*, *538*, 736545.
https://doi.org/10.1016/j.aquaculture.2021.736545

Zhou, X., & Stephens, M. (2012). Genome-wide efficient mixed-model analysis for association studies.
Nature Genetics, *44*(7), 821–824. https://doi.org/10.1038/ng.2310

TABLES AND FIGURES

Figure 1. Chr8 marker genotype proportions in juveniles from families F114-F117 prior to the *Vibrio*
exposure trial. Arrows indicate a genotype with a significant difference (p < 0.05) between families, where
more homozygous alternate genotypes were observed in F115 relative to F114 and F117 (χ^2=15.183, p-
value = 0.019).

Figure 2. Proportions of oysters that died (light grey) or survived (dark grey) in each family in the *V. aestuarianus* disease challenge. Families are grouped by the parental genotypes at the Chr8 marker, as indicated at the top of the plot. The unexposed control group exhibited 100% survival.

Figure 3. Kaplan-Meier survival curves for Pacific oyster families after exposure to *V. aestuarianus*. Survival distributions for the families showed significant differences (χ^2=39.070, p<0.001), and pairwise differences are described in Results. Parental genotypes for the Chr8 marker per family are given in Figure 2.

Figure 4. The effects of the Chr8 marker allelic dosage on mortality as quantified as days to death in the *V. aestuarianus* trial in families F114-F117 (panels A-D, respectively; i.e., those with both parents heterozygous for the Chr8 marker). No significant association was found (p>0.05) (F114 χ^2=5.370, p>0.05; F115 χ^2=14.172, p>0.05; F116 χ^2=11.280, p>0.05; F117 χ^2=4.429, p>0.05).

Figure 5. The effects of the Chr8 marker allelic dosage on mortality as quantified by the frequency of survival (light grey) or mortality (black) in the *V. aestuarianus* trial in families F114-F117 (panels A-D, respectively, i.e., those with both parents heterozygous for the Chr8 marker). No significant association was found (Chi^2 test p > 0.05).

Figure 6. Manhattan plot showing associations of each SNP from the unimputed RAD-seq dataset for
survivorship (mortality or survival) in the *V. aestuarianus* trial for families F114-F117. The red horizontal
line indicates the genome-wide Bonferroni-corrected p-value significance threshold of 0.05. No significant
associations were observed.