1 **Elevational and Oceanic Barriers Shape the Distribution, Dispersal and Diversity of Aotearoa's**
2 **Kapokapowai (*Uropetala*) Dragonflies**
3

Authors: Ethan R. Tolman*[1,2], Christopher D. Beatty*[2,3], Aaron Goodman*[2,4], Priscilla Ramchand*[2,5],
Tara McAllister[6], Kelly Reyes[7], Akifa Zahara[7], Kassandra Taveras[7], Vincent Wade[7], John Abbott[8], Seth
Bybee[9], Robert Guralnick[11], Kathleen M. Harding[12], Manpreet K. Kohli[2,13], Paul B. Frandsen[10], Jessica
Ware[†2], Anton Suvorov[†1]
*Equal contributing authors to the manuscript
†Senior authors
[1]Department of Biological Sciences, Virginia Tech, Blacksburg, Virginia
[2]Division of Invertebrate Zoology, American Museum of Natural History, New York, New York
[3]Program for Conservation Genomics, Department of Biology, Stanford University, Stanford, California
[4]The Graduate Center, The City University of New York, New York, New York
[5] Brooklyn College, The City University of New York, Brooklyn, New York
[6]Te Wānanga o Aotearoa, New Zealand
[7]Science Research Mentorship Program, American Museum of Natural History, New York, New York
[8]Alabama Museum of Natural History and Department of Research and Collections, The University of
Alabama, Tuscaloosa, Alabama
[9]Department of Biology, Brigham Young University, Provo, Utah
[10]Department of Plant and Wildlife Sciences, Brigham Young University, Provo, Utah
[11]Florida Museum of Natural History, University of Florida, Gainesville, Florida
[12]Redwood City, California, USA
[13]Department of Natural Sciences, Baruch College, New York, New York

Acknowledgements
We would like to thank Thomas Beatty and Mahealani Kaloi for their support of this project. We would
also like to acknowledge Dr. Milen Marinov for his generosity in providing Odonata specimens, as well
as the Department of Entomology at the Smithsonian Institution's National Museum of Natural History,
Florida State Collection of Arthropods, and American Museum of Natural History for providing
specimens used in this analysis.
Funding
This work was funded by the Graduate Student Research Award from the Society of Systematic
Biologists, the Tow Undergraduate and Graduate International Research Stipends from The City
University of New York Brooklyn College, the Maxwell/Hanrahan Award from the American Museum of
Natural History, and by the National Science Foundation (NSF) Awards [Bybee: 2002432; Ware:
2002473; Guralnick: 2002457; Abbott: 2002489: Genealogy of Odonata (GEODE): Dispersal and color
as drivers of 300 million years of global dragonfly evolution].

Data availability
Filtered VCF files with and without *P. raptor*, and log and xml files from the BEAST2 runs have been
uploaded to figshare (10.6084/m9.figshare.27896910). The Raw reads have been uploaded to the
sequencing read archive (bioproject PRJNA1190089).
Conflict of Interests Statement
The authors declare no conflicts of interest.
44

Abstract

Mountains and islands provide an opportunity for studying the biogeography of diversification and population fragmentation. Aotearoa (New Zealand) is an excellent location to investigate both phenomena due to alpine emergence and oceanic separation. While it would be expected that separation across oceanic and elevation gradients are major barriers to gene flow in animals, including aquatic insects, such hypotheses have not been thoroughly tested in these taxa. By integrating population genomic from sub-genomic Anchored-Hybrid Enrichment sequencing, ecological niche modeling, and morphological analyses from scanning-electron microscopy, we show that tectonic uplift and oceanic vicariance are implicated in speciation and population structure in Kapokapowai (*Uropetala*) dragonflies. Although Te Moana o Raukawa (Cook Strait), is likely responsible for some of the genetic structure observed, speciation has not yet occurred in populations separated by the strait. We find that the altitudinal gradient across Kā Tiritiri-o-te-Moana (the Southern Alps) is not impervious but it significantly restricts gene flow between aforementioned species. Our data support the hypothesis of an active colonization of Kā Tiritiri-o-te-Moana by the ancestral population of Kapokapowai, followed by a recolonization of the lowlands. These findings provide key foundations for the study of lineages endemic to Aotearoa.

Keywords

Anisoptera, Epiprocta, biogeography, population genomics, Petaluridae, Islands, Odonata

Introduction

Islands are widely recognized for their unique role in shaping biodiversity (Barreto et al., 2021; Conway & Olsen, 2019; Gleditsch et al., 2022; Wilson, 2009). The Theory of Island Biogeography by MacArthur and Wilson (1963) summarized several of these phenomena, and this theory was later extended to mountaintop habitats (Wyckhuys et al., 2022). The islands of Aotearoa (New Zealand; see table 1 for nomenclature) have often been considered as a natural laboratory to study both island and mountain biogeography (Shepherd et al., 2024; Thomas et al., 2023; Wallis & Buckley, 2024; Waters & Craw, 2006). This is due to the opening and closing of Te Moana o Raukawa (Cook Strait) which separates Te Ika-a-Māui and Te Waipounamu (also known as the North and South Islands respectively) and last opened 500,000 years ago (Lewis et al., 1994), and the uplift of Kā Tiritiri-o-te-Moana (the Southern Alps), a relatively young mountain range on Te Waipounamu (South Island; Table 1).

Despite its young geologic age, Te Moana o Raukawa has been proposed as a driver of allopatric speciation and population divergence in several taxa including amphipods (Stevens & Hogg, 2004), Weka (terrestrial birds) and their ectoparasites (Trewick et al., 2017), and skink lizards (O'Neill et al., 2008). The tectonic uplift of the Kā Tiritiri-o-te-Moana mountain range likely opened niches and acted as a driver of diversification in flora (Thomas et al., 2023) and fauna (Buckley et al., 2024) of Te Waipounamu. Most lineages of alpine flora arrived in the later Tertiary period, and the environmental upheaval in the Pliocene and Pleistocene drove diversity in these lineages (Thomas et al., 2023). Most alpine invertebrate fauna are relatively young, with ages corresponding to the increased uplift of the Kā Tiritiri-o-te-Moana range 5 million years ago (Ma), and the development of a permanent snow line just over 1 Ma (Buckley et al., 2024). Similar to the flora, it is broadly understood that most invertebrate fauna evolved from lineages already on the island as niches opened with the increased elevation (Buckley et al., 2024), although it has been hypothesized that invertebrates have ridden tectonic uplift to higher altitudes (Heads, 2017).

Kapokapowai as an Important Study System in Island Biogeography

Aquatic insects have been used to better understand the past climate and geology of Aotearoa, and are
useful models in the study of evolution and ecology (Córdoba-Aguilar et al., 2022), but they have been
relatively underrepresented in biogeographic and broader biodiversity work (Leschen et al., 2023) and
knowledge about biogeography remains limited for many taxa. One such group of aquatic
macroinvertebrates are the dragonflies first referred to as Kapokapowai (among other names) by Māori
(Grant, 2014), and later, named *Uropetala* by scientists from colonial Great Britain and Belgium (Selys
1858).
Kapokapowai are a particularly promising system for studying the impact of tectonic uplift and
oceanic barriers in speciation as a member of the so-called "relict" (Ware et al., 2014) dragonfly lineage
Petaluridae, a member of the previously diverse superfamily Petalurida. Petaluridae originated in the mid
Jurassic, as sister to the species rich Gomphidae family, but comprising only 11 currently described
species (see Tolman et al. (2024) and Cairns et al. (2025) for discussion of the status of one species of the
genus *Petalura*) it is among the most species poor lineages of the order Odonata (Tolman et al., 2024).
Despite inhabiting fen habitats which should readily promote fossilization, there is only one crown fossil
Petaluridae, which is dated to the mid Cretaceous and closely resembles the extant species found at the
fossil locality (Tolman et al., 2024). Thus, there is no compelling paleontological evidence that the
Petaluridae were previously diverse. Speciation in this family appears to largely be driven by vicariance,
as diversification in Petaluridae is strongly correlated with continental drift, and the expansion and
contraction of inland seas and land bridges (Ware et al., 2014; Tolman et al., 2024).
Kapokapowai likely diverged from the Australian petalurids during continental split between
Zealandia and Gondwana, and persisted through the Oligocene drowning of ~18 Ma (Cooper and Cooper
1997; Ware et al., 2014; Tolman et al., 2024). Within Kapokapowai there are two currently described
species, *Uropetala carovei* (Selys, 1843) which is thought to inhabit lower elevations of both Te Ika-a-
Māui and Te Waipounamu (North and South Islands, respectively), and *Uropetala chiltoni* (Tillyard,
1921), which inhabits higher elevations of Te Waipounamu (South Island). Kapokapowai is the only
lineage within Petaluridae where speciation is thought to have occured within the past 5 Ma (Tolman et
al., 2024), thus the implications of geographical changes on species and population diversity can be
studied in finer detail using this species complex than with other lineages of Petaluridae. The study of
Kapokapowai can provide important hypotheses for the study of other "relict" lineages on island systems,
defined by McCulloch and Waters (2019) to refer to phylogenetically "old lineages" found on
geologically "young islands." Island relicts have been considered to be a high priority for conservation
biology (Grandcolas et al. 2014) and for the broader study of extinction and speciation more generally
(Grandcolas et al. 2014).
121
Māori Knowledge of Kapokapowai
In systems such as Kapokapowai, Indigenous knowledge is an important source of ecological and
evolutionary information (Jessen et al. 2022). According to Māori cosmology, dragonflies and other
insects are considered the children of Tāne Māhuta (the god of forests, birds and insects). Dragonflies
appear in whakaairo (traditional carvings which often adorn the walls of Māori meeting houses) and
waiata (songs) of Māori (Indigenous Peoples of Aotearoa). Stories about dragonflies also feature in
pūrakau (legends) belonging to specific iwi (tribes) and hapū (subtribes). All insects, including
dragonflies, were created before humans in Māori genealogical histories and are therefore tuakana
(roughly translated as an elder family member) and are respected accordingly. Kapokapowai or Kapowai
are names commonly-used for the Petaluridae of Aotearoa, which loosely translates to water snatchers,

132	though there are other recorded names including Tītīwaiora, Kihihiwaru and Uruururoroa (Grant, 2014).
133	Thus, these historical and cultural accounts provide valuable information about indirect as well as direct
134	spatiotemporal distribution records of Kapokapowai. As this work was conducted with the express
135	permission of the iwi who own the lands of Aotearoa, we refer to these dragonflies by the Māori name,
136	Kapokapowai, in our work here. We also note that the name Kapokapowai was used to describe these
137	dragonflies in the literature by Taylor (1855), italicizing "*Kapokapowai*" as to explicitly refer to a genus,
138	three years prior to the first use of *Uropetala* by Selys (Grant, 2014; Taylor, 1855, 1870), and
139	Kapokapowai has continued to be used to refer to dragonflies in this genus in scientific literature
140	(Durrant, 2024; Fong, 2024; E. R. Tolman et al., 2024). However, we are not making any change to the
141	binary taxonomic nomenclature of this genus here. In referring to Kapokapowai by their Māori name in
142	this work, we designate *U. chiltoni* as Kapokapowai Montaine (KM), and *U. carovei* from either island as
143	Kapokapowai North (KN) and Kapokapowai South (KS; Table 1), representing the three putative species
144	in the Kapokapowai species complex.
145	
146	*The Kapokapowai Species Complex*
147	Debate about whether this complex should appropriately be considered to have two species has been
148	ongoing for decades (Andress, 1998; L. S. Wolfe, 1952; Winstanley & Rowe, 1980). According to
149	Tillyard's original description, KM is characterized by large pale blotches on its labrum, which are not
150	present in KN and KS, and by black femur leg segments, which are brown in KN and KS. However,
151	variation in the splotches of the labrum have been used to suggest that hybrid populations between KM
152	and KS can be found at the north end of Te Waipounamu (Winstanley & Rowe, 1980). Behavioral work
153	suggests that female KM lay all their eggs in one plant, and in repeating this egg laying process females
154	tend to badly damage their wings; such damage has not been seen in KN or KS (W. J. Winstanley, 1981).
155	This behavioral work should be replicated to confirm that this is a species–wide behavior. It has also been
156	suggested that KN and KS may be different species, as KN on Te Ika-a-Māui may have broader superior
157	appendages and larger adult body size than their counterparts on Te Waipounamu (L. S. Wolfe, 1952).
158	Two different karyotypes have been identified for this genus, but it is unclear if this is due to laboratory
159	error (Kuznetsova & Golub, 2020); no other attempts have yet been made to determine the number of
160	species in this genus using molecular data. Based on known morphological variation and behavioral data
161	(L. S. Wolfe, 1952; Winstanley & Rowe, 1980), the dragonflies known as Kapokapowai are hypothesized
162	here to comprise between one and three species, the diversification of which was shaped by both Te
163	Moana o Raukawa (Cook Strait) and Kā Tiritiri-o-te-Moana (Southern Alps).
164	Here, we use population genomics, alongside scanning electron microscopy images of secondary
165	male genitalia, and environmental niche models to determine the number of species and the population
166	dynamics within Kapokapowai and to study how Kā Tiritiri-o-te-Moana and Te Moana o Raukawa may
167	have driven speciation in this genus, and consider how our findings may inform evolution on Aotearoa
168	more generally
169	
170	**Materials and Methods**
171	***Population genomics***
172	*Data acquisition*
173	We utilized molecular data from a total of 37 dragonflies, 18 KM, 7 KN, 5 KS, and 7 *Phenes raptor*
174	which were used as an outgroup (Supplemental table 1). We generated a 93 locus, 20kb (on average)
175	nucleotide Anchored-Hybrid Enrichment (AHE) read library (Tolman et al., 2024) for 16 KM, and all

other sequenced individuals, following the protocol used by Goodman et al (2023) and Tolman et al.
(2024). Briefly, we extracted DNA from one leg of each specimen, which was removed using sterilized
forceps, in the case of nymphs and adults. The entire thorax was used to extract DNA from exuvia. We
incubated the tissue in proteinase K for one week at 55C, and then followed the manufacturers protocol
for the remainder of the DNA extraction using the Zymo DNA Micro-prep kit (Hilden Germany). We
sent the DNA extract to RAPID Genomics (Gainesville, Florida) for library preparation and sequencing.
Additionally, we incorporated in the current study previously generated 1000 locus (which contained the
93 targeted loci for the 35 individuals sequenced here) AHE read libraries for 2 KM individuals, which
were used in (Tolman et al., 2024) to estimate phylogenetic relationships of all species of Petaluridae.
185
Population Structure Analysis
We mapped the 37 sequenced AHE read libraries to a previously compiled soft-masked whole genome
assembly (NCBI GenBank assembly GCA_036926305.1) of an individual KS (Tolman et al., 2023) with
BWA2 (0.7.17-r1188; H. Li & Durbin, 2009) using default parameters, and calculated mapping statistics,
average depth and converted the output to a sorted BAM file with samtools v1.16.1 (Danecek et al.,
2021). We called variants with bcftools v1.15.1 (Danecek et al., 2021), using the options: `bcftools`
`mpileup -a AD,DP,SP -Ou -f reference.fasta Input_bam | bcftools call -`
`f GQ,GP -mO z -o output`. We then filtered the resulting VCF file for a minimum quality score of
30, and a minimum and maximum depth of 10 and 350 respectively to capture the region targeted with
hybrid probes and the adjacent flanking region that is also sequenced and often more variable (Goodman
et al., 2023).
To visualize population structure we performed a PCA analysis on a covariance matrix generated
from the called SNPs with VCFtools v0.1.16 (Purcell et al., 2007) without pruning for linkage-
disequilibrium (LD). To test if the outliers in the PCA could be a result of missing data, we identified the
missingness of all individuals in the unpruned VCF file which excluded *Phenes*, and compared the
missingness of KM with positive and negative PC1 scores. To model the ancestry of each sampled
individual, we generated admixture plots for each putative species from admixture models assuming 2-7
ancestral populations with ADMIXTURE v1.3.0 (Skotte et al., 2013) and compared model fit using a
cross-validation approach. The admixture graph under the best fitting models and the PCA plots were
generated in R v2024.04.2+764 using mapmixture v1.1.0 library (Jenkins, 2024). Additionally, we
calculated the F_{ST} between and within each of the three putative species both using VCFtools v0.1.16
(Purcell et al., 2007). To test for a correlation between genetic and geographic distance among our
sampled individuals, we generated pairwise genetic distance matrices between sampled individuals with
VCFtools v0.1.16 (Danecek et al., 2021), and then used the Mantel test in the vegan package in R
v2024.04.2+764 (Oksanen et al., 2024), with 1,000 permutations to determine the significance of the
correlation between geographic and genetic distance.
Genetic linkage can bias the recovery of population structure, as can inappropriately pruning sites
(Bercovich et al. 2025). To determine the impact of genetic linkage on this dataset, and test the
appropriateness of considering SNPs without LD pruning we pruned linked sites with PLINK v1.9.0-b.8
(Purcell et al., 2007) using the flags `--indep-pairwise 10 5 0` and re-ran the PCA analysis
using the resulting VCF file.
217
Species delimitation

To delimit species based on our SNP data, we used the SPEEDEMON v.1.1.1 (Douglas & Bouckaert,
2022) pipeline in BEAST2 v2.7.7 (Bouckaert et al., 2014), using all 395 biallelic recovered SNPs after
filtering.t,. We used snapper v1.1.4 (Stoltz et al., 2021) to run the SPEEDEMON pipeline on SNPs as
opposed to alignments, and BICEPS v1.1.2 (Bouckaert, 2022) to implement a Yule Skyline Collapse
Model to test for species delimitations, implemented with a chain length of 10^6 draws. We designated
KM as a reference species, with KN and KS as putative species. We used the mean pairwise percent
genetic divergence between individuals of KM (~.54), as calculated from the VCF file using the pandas
(McKinney, 2010) library for data manipulations and numpy (Harris et al., 2020) for numerical
computations, as the prior mean on the theta prior (Douglas & Bouckaert, 2022). We used an alpha of 2.0,
and beta of 0.26942 to achieve this mean in the Gamma distribution of the theta prior (see Stoltz et al.,
2021 for a full description of these options), and followed the developer's suggestions for all other
parameters in the BEAST2 run (Stoltz et al., 2021) . We independently ran the analysis two times to test
for model convergence. We computed the posterior mean and upper credible (95%) limit of the Gelman-
Rubin-Potential-Scale-Reduction Factor (PSRF) of each parameter, using the coda v0.19.4.1 library
(Plummer et al. 2006), in r4.4.1(R Core Team 2021), to test for adquate model convergence.
We also calculated the Effective Sample Size (ESS) for both runs in TRACER v1.7.2 (Rambaut et al.,
2018) = to test for adequate mixing (at ESS >150) and model convergence (Rambaut et al., 2018).
236
Divergence time estimation
We utilized SNAPPER v1.1.4 (Stoltz et al., 2021) implemented in BEAST2 v2.7.7 (Bouckaert et al.,
2014) to determine when the three putative species diverged, and to further explore which populations are
most closely related. We used a different SNP set for this analysis, which included seven sampled *Phenes*
raptor to use as an outgroup. The sequencing and variant call pipeline used to include these samples was
identical to the protocol listed earlier. We used PLINK v1.9.0-b.8 (Purcell et al., 2007) with the flags `--`
`indep-pairwise 10 5 0.2` to prune linked sites from the VCF containing both *Phenes* and
Kapokapowai. We used this pruned input file, with uniform age constraints of 113-135 Ma for the root
age and of 0.1-15 Ma for the age of Kapokapowai (as previously identified by Tolman et al. 2024), to
generate the snapper input file with a MCMC chain length of 10^7. SNAPPER can only implement a
strict-clock model, a reasonable assumption for recently diverged populations and species, and (Stoltz et
al., 2021), and we did not adjust the default tree model employed by SNAPPER. We calculated ESS for
both runs in TRACER v1.7.2 (Rambaut et al., 2018) to test for adequate mixing, and computed the
posterior mean and upper confidence (95%) limit of the PSRF of each parameter, using the coda v0.19.4.1
library (Plummer et al. 2006), in r4.4.1(R Core Team 2021), to test for adequate model convergence.
252
Tests of gene flow
As the resulting tree and species delimitation analyses supported KN and KS as sister, we tested for
evidence of introgression between KM and KN and KM and KS with an ABBA-BABA test from Dsuite
v0.5 r57, which is optimized to work on lineages with shorter evolutionary time scales. We used the
unpruned VCF file which included *Phenes raptor* as input (Malinsky et al., 2021), and used *P. raptor* as
an outgroup (E. R. Tolman et al., 2024).
259
Morphological analysis
To identify potential morphological synapomorphies of the putative species of Kapokapowai we
evaluated the male secondary genitalia. We specifically looked at the vesica spermalis, or sperm pump.

This structure is located on the ventral side of the second abdominal segment in male Odonata, to which
sperm is transferred prior to copulation, and may provide informative morphological characters for
species determination (Kennedy, 1922; May, 1997; Ware, 2008). The vesica spermalis of Odonata is
composed of four segments that are variable between species in their relative size and shape. In
Petaluridae, a barb on the second segment near the junction with the third segment and the horns on the
lateral side of the third segment have been used to support intrageneric difference (E. Tolman, 2024). We
imaged the vesica spermalis (later referred to as penes) from two males from each putative species to try
and identify more definitive diagnostic features for species delimitation. We followed the protocol for
scanning electron microscopy outlined by Tolman et al. (2024). Briefly, penes were mounted laterally and
dorsally on aluminum stubs and coated with gold palladium using a Cressington 108E sputter coater and
specimens were then imaged using an S-7400 Hitachi scanning electron microscope.
274
Environmental niche modeling
Occurrence Records
We acquired occurrence records of adult KN, KS, and KM from the Global Biodiversity Information
Facility (GBIF)(available in Supplemental data). We selected occurrences from GBIF possessing
preserved museum samples and research grade observations, which include verified latitude and longitude
coordinates, a photograph of the sighting, observation date, and at least ⅔ agreement on species
identification by the community. We filtered occurrences by removing sightings from erroneous localities
(middle of the ocean, locations of large museums). We divided occurrences for KN and KS between Te
Ika-a-Māui and Te Waipounamu (North and South Islands) to be used as individual models
(Supplemental Table 3).
285
Environmental Data
We acquired averaged sets of environmental predictor variables for modeling consisting of purely
bioclimatic variables. All analyses were conducted using the statistical programming language R v. 4.1.2.
We acquired environmental rasters at 2.5 arc-second resolution (~5km at the equator) from the CHELSA
climate database v2.1 (Karger et al., 2017, 2023) . We downloaded the 'Anthropocene' bioclimatic
dataset (1979 – 2013) consisting of 19 bioclimatic variables which follow Worldclim v2 (Fick &
Hijmans, 2017). We omitted bioclimatic variables 8 and 9 after visual inspection of interpolation
discontinuities (Booth, 2022). Although such spatial artifacts were minor within our initial study extent,
we observed major breaks of climate smoothing when we extrapolated our model to a larger spatial extent
within the last glacial maximum and interglacial periods.
296
Model Construction
We spatially thinned occurrences to the resolution of our environmental variables to reduce the effects of
sampling bias and artificial clustering (Veloz, 2009). We thinned occurrences by 5km to match the spatial
resolution of our environmental data. We chose study extents for occurrences of KN and KS found in Te
Ika-a-Māui and Te Waipounamu, defined as polygons around all localities between both islands
respectively; we chose the same study extent for occurrences of KM found in Te Ika-a-Māui. We chose
these study extents to include potential suitable habitat, while excluding large areas outside the species'
dispersal limitations (Peterson & Lieberman, 2012). Within this extent, we randomly sampled 50,000
background points for modeling and extracted their environmental values. We used these values to
calculate correlations between variables using the 'vifcor' and 'vifstep' functions in the usdm package

(Naimi, 2023) and filtered out variables with correlation coefficients higher than 0.7 and a Variance
Inflation Factor (VIF) threshold of 10, both standard cutoffs in such analyses (Goodman et al., 2024).
To model the distribution of putative Kapokapowai species, we used the presence-background
algorithm MaxEnt v3.4.4 (Phillips et al., 2017), using the R Package ENMeval 2.0.0 (Kass et al., 2021)
for model building, parameterization, evaluation with different complexity settings, and reporting of
results. We partitioned our data using the 'checkerboard2' strategy. To prevent our model from
extrapolating beyond the bounds of training data, we clamped our models by omitting ranges of
environmental data which fall outside of the training data. All final models were fitted to the full dataset.
We tuned model complexity to find optimal settings using different feature classes, including linear (L),
quadratic (Q), and hinge (H) as well as regularization multipliers 1 through 5 (higher numbers penalize
complexity more) (Radosavljevic & Anderson, 2014; Warren & Seifert, 2011).
We assessed our model using averages of threshold-dependent (omission rate) and
threshold-independent (Area Under the Receiver Operating Characteristic or AUC) discrimination
performance metrics calculated on withheld validation data (Warren & Seifert, 2011). The 10th-percentile
omission rate sets a threshold as the lowest suitability value for occurrences after removing the lowest
10% suitability values (Kass et al., 2021; Radosavljevic & Anderson, 2014). Validation AUC is a
measure of discrimination accuracy that can be used to make relative comparisons between ENMs with
different settings fit on the same data (Lobo et al., 2008; Radosavljevic & Anderson, 2014). Finally, we
checked the results against the Akaike Information Criterion with correction for small sample sizes
(AICc; calculated on the full dataset) (Warren & Seifert, 2011). To investigate model behavior, we
examined marginal response to suitability. Marginal response curves show the modeled relationship of
each variable individually with the occurrence data when all other variables are held constant (Phillips et
al., 2017).
We made habitat suitability predictions for KM, and KN and KS from both islands using our
'Anthropocene' environmental predictor variables. We projected our modern-day models to a new study
extent to encapsulate the potential habitat of Kapokapowai within the historic past, defined as a polygon
around both Islands of Aotearoa. We generated thresholdless predictions by converting raw predictions to
a scale of 0-1 to approximate probability of occurrence using the 'cloglog' transformation (Phillips et al.
2017). We also generated a threshold prediction, calculated from the 10-percentile omission rate from our
model evaluation. To estimate habitat suitability of Kapokapowai within the geologic past, we generated
predictions using environmental predictor variables from the PaleoClim dataset (Brown et al., 2018). We
generated predictions within three distinct time periods: Mid-Holocene (8.3 – 4.2ka), Last Glacial
Maximum (21ka), and the Last Interglacial (130ka).
340
Ordination Analysis
To determine niche differentiation between putative species, we compared niche overlap in occurrences
of KN and KS from Te Ika-a-Māui and Te Waipounamu using an ordination framework by first reducing
dimensionality within the datasets via a Principal Component analysis (PCA). Using the 'espace_pca'
function in the *Wallace* v2.0.5 and *ade4* package v1.7 in R (Dray & Dufour, 2007; Kass et al., 2018)), we
generated a Principal Components Analysis (PCA) using environmental variables from Kapokapowai
occurrences and plotted with correlation loadings to infer the degree of influence particular environmental
variables possess in the distribution of the 50,000 background points within niche space; we chose
bioclimatic variables which were uncorrelated and shared among both Kapokapowai MaxEnt models.
Using the 'espace_occDens' function in the package *ecospat* v3.2 (Di Cola et al., 2017, p. 2) an

occurrence density grid was estimated for both the environmental values at each occurrence point and
background extent points using a kernel density estimation approach. Niche overlap between occurrence
density grids of environmental values at each occurrence point and background points were compared
using Schoener's D (Schoener, 1968) 'espace_nicheOv' function. Finally, using the
'ecospat.plot.overlap.test' function, we conducted a niche similarity test in which Kapokapowai niches
from both islands are randomly shifted in the background extent and permuted 1000 times, and the niche
overlap between the two are recalculated. A p-value <0.05 indicates that the niches from both time
periods are significantly similar with each other.
359
Ancestral State Reconstruction of Elevation Ranges in Petaluridae
Having considered the niche space and population structure of low and high elevation Kapokapowai, then
determined whether inhabiting mountains (KM) or lowlands (KN and KS) is the ancestral state for
Kapokapowai–an important consideration in how these dragonflies are impacted by geography–by
conducting an ancestral state reconstruction of the maximum elevation range for all of Petaluridae. We
downloaded all occurrences of Petaluridae from the Global Biodiversity Information Facility (GBIF;
GBIF.org, 2025) and determined the elevation of each occurence point using the elevatr v0.99.0 R
package (Hollister et al. 2023). We determined the 90th percentile of elevation for each species of
Petaluridae to minimize the impact of outlier observations, and classified the species as occupying high
(>900 M), medium (400-900 M) or low (<400 M) elevations based upon this statistic. To model ancestral
states we fit three variants of the Markov k-state model: Equal-Rates (ER), a single transition rate
between all pairs of states, Symmetric-Rates (SR), distinct forward and backwards rates that are equal in
reciprocal pairs; All-Rates-Differed (ARD), allowing each transition a unique rate. We fit each model
with phytools v2.4-4 (Revell 2012) with estimated stationary frequencies. We also fit a custom model
with equal rates between low and medium elevation, medium and high elevation, and a different rate
between low and high elevation. We then compared model fit for the four models using AIC and used
model averaging from ANCR to integrate node-state estimates across models according to their AIC
rates. We used the dated species tree for Petaluridae (including the two currently described species of
Kapokapowai) from Tolman et al. (2024) and added *Petalura pulcherrima* as sister to *P. ingentissima* at
14 Ma (within the confidence interval identified by Ware et al. (2014)). We tested the model with and
without the fossil *Argentinopetala archangelskyi* which has been identified as sister to *P. raptor* (Tolman
et al., 2024). To determine the elevation of this fossil at ~115 Ma we converted the modern coordinates
into the paleo-coordinates at 115 Ma using the palaeorate function in palaeoverse v1.4 (Jones et al. 2023)
with grid rotation under the MERDITH2021 plate model. We used raster v3.6.31 (Hijmans, 2025) to load
the 1° resolution Early Cretaceous Digital Elevation Model (DEM; Scotese and Wright 2018) and extract
the paleo-elevation at the rotated point. As all states were equally probable at each internal node with and
without *A. archangelskyi*, we reconducted searches with the averaged model using stochastic character
mapping. To infer the frequency of changes between character states, we simulated 1000 discrete
character histories under the selected averaged model, with empirical root-state frequencies set using the
fitzjohn option in phytools v2.4-4 (Revell 2012). All aforementioned ancestral state reconstruction
analyses were conducted in R v4.4.1 (R Core Team 2021).
391
Results
Population genomics
Population Structure Analysis

Mean sequencing depth varied from 0.02x to 0.19x, and missingness (calculated from the LD pruned
VCF file) ranged from 0.01 to 0.27 (Supplemental Table 1). After filtering for mapping quality (> q30)
and depth (10-350), we retained 365 SNPs for the initial population structure analysis. The weighted F_{ST}
values were estimated at 0.23 between KM and KS, 0.29 between KM and KN and 0.11 between KN and
KS (Table 2). We then separated the data into subsets as follows by their distributions along PC1 (Fig
1A): KM (all individuals), KM (negative PC axes), KN (Te Ika-a-Māui) and KS (Te Waipounamu). The
F_{ST} values differed when only a subset of KM was used (Supplemental Table 3). There was no evidence
that geographic separation was correlated with pairwise genetic distance within Kapokapowai (Pearson's
correlation coefficient $r = 0.02462$, p-value = 0.322).
The centroid of KM was distal from the centroids of KN and KS across PC1, while the centroid
of KN and KS differed along PC2 (Fig. 1A). Despite the differing centroids across PC1 and PC2, there
was overlap in the space occupied between all three putative species (Fig. 1A). The missingness of KM
individuals with a positive PC1 ranged from 0.02 to 0.17, while the missingness of KM individuals with a
negative PC1 value ranged from 0.07 to 0.16 (Supplemental Table 3). 118 SNPs were retained after LD
pruning the VCF, although the resulting population structure visualized through PCA (Supplemental Fig.
1) largely reflected the structure generated from pruned SNPs (Fig. 1A), with a slightly reduced distance
between KN and KS.
Because of the low cross validation-error indicating comparable fit to other tested models
(Supplemental table 5), and the three clusters identified in our PCA analysis, we provide the three
ancestral populations model in our main figure and discussion (Fig. 1B), but also provide assumptions of
K = 2, and K = 4 (Supplemental fig. 2). Ancestral population 1 was only present in KM, while ancestral
populations 2 and 3 were present in all three putative species (Fig. 1B).

417
418   *Species delimitation analysis*
419   All parameters in the SPEEDEMON analysis showed adequate mixing (ESS > 400), and model
420   convergence (PSRF = 1.0 with 95% CI <1.01 for all parameters). The species were recovered as currently
421   described, specifically KM as *U. chiltoni* and KN and KS as *U. carovei*, with very high (99%) posterior
422   support.

423
Divergence time estimation
Including the outgroup *Phenes raptor* we recovered 727 SNPs and retained 204 of these SNPs after LD
pruning. The SPEEDEMON runs showed adequate mixing (ESS>600 for posterior, likelihood, prior,
lambda and tree height; ESS >150 for clock rate) and strong evidence of convergence (PSRF=1.0 with
95% CI <1.06 for all parameters). In the dated tree Kapokapowai and *Phenes* shared a Most Recent
Common Ancestor (MRCA) between 113.0 and 133.7 Ma, KM shared a MRCA with KN and KS
between 2.6 and 15.0 Ma, while KN and KS diverged between 0.1 and 3.2 Ma (Fig. 1E; Supplemental
Fig. 2).

432
433   *Tests of gene flow*
434   The ABBA-BABA test, implemented in D-suite using all 727 SNPs identifie devidence of gene flow
435   between KM and KS (p < .01; Table 2)). It is important to note that this test cannot determine the
436   directionality of gene flow.

437
438

Figure 1: Population Genomics of Kapokapowai

Fig. 1: (A) PCA of Kapokapowai from SNPs, colored by putative species: Kapokapowai Montane (Blue),

KN (Green) and KS (Teal). Each dot represents one individual with the distance from each point to the

centroid of each putative species shown with a line. (B) Admixture analysis of Kapokapowai assuming

three ancestral populations, with the ancestral populations 1-3 colored as blue, green, and teal respectively

per each sampled individual (x-axis). Pie charts are mapped according to their proximate locality, and

represent the proportion of each ancestral population in the gene pool of each locality. Ancestral

population 1 is only found in KM. (C) Adult Female Kapokapowai North from Te Ika-a-Māui. (D)

Nymphs of Kapokapowai Montane. (E) Dated tree displaying the divergence between the three putative

species and the outgroup *Phenes raptor*. Notable geographic events in Aotearoa are annotated on the tree,

including the Oligocene drowning when Aotearoa was reduced to 18% of its current size (Cooper and

Cooper 1997), the reemergence of taxa following the drowning, the uplift of Kā Tiritiri o te Moana, and

the Pleistocene land bridge events between Te Ika-a-Māui and Te Waipounamu.

Morphological analysis

Figure 2: Male Secondary Genitalia of the Three Hypothesized Kapokapowai Species

Scanning electron microscopy. Images of male penes from six sampled Kapokapowai (two from each

putative species). Divergent features include (1) curves at the end of the horns only found in

Kapokapowai Montane, (2) the barb near the joint of the second and third segments, and (3) a knob found

near the barb from one individual from Kapokapowai North.

The "horns" of KM (see figure 2, feature 1) are downcurved at the tips in a way not observed in any of

the three KN or KS individuals (Fig. 3). The angle and curvature of the barb (figure 2, feature 2), and the

presence of a "knob" near it (figure 2, feature 3) differs between individuals, but does not appear to be

fixed between any populations. One individual of KN had a knob near the barb.

Ecological Niche Modeling

Optimal model settings varied considerably for Kapokapowai based on 10-percentile omission rate and

mean validation AUC metrics. Based on the results of the collinearity analysis, we used the following six

predictor variables to build all models: Isothermality (bio03), temperature seasonality (bio04), maximum

temperature of the warmest month (bio05), precipitation seasonality (bio15), precipitation of the warmest

quarter (bio18), and precipitation of the coldest quarter (bio19). Models varied in complexity with

expressing either hinge-loss, or linear-quadratic feature classes with regularization multipliers of 3-4

(higher complexity penalty). Mean validation AUC was adequate for KS and KM (0.75 and 0.86

respectively), while 10% omission rate was low relative to its expected value of 0.1 (0.11 and 0.09

respectively) (Supplemental Table 2). Furthermore, optimal models for KS and KM possessed low

amounts of non-zero lambdas (coefficients) and AICc values suggesting low amounts of model

overinflation (Supplemental Table 2). Optimal models for KN performed poorly, possessing an

unacceptably low AUC value (0.44) and high 10% omission rate (0.16) (Supplemental Table 2).

Optimal models retained 4 – 6 predictor variables after regularization, and relied mainly on

contributions from Isothermality (bio03), temperature seasonality (bio04), maximum temperature of the

warmest month (bio05), and precipitation of the wettest and driest quarters (bio18 and bio19 respectively)
(See Supplemental Fig. 3). Response curves of climate variables revealed differing relationships with
suitability among Kapokapowai models (Supplemental Fig. 1). Response curves for KN expressed jagged
optimal ranges, with habitat suitability not following a smooth or continuous gadient. Temperature
Seasonality (bio03) and precipitation of the coldest quarter expressed positive linear/hinge relationships
with suitability among all three Kapokapowai models, while precipitation of the wettest quarter (bio18)
expressed negative linear/hinge relationships with suitability for KN and KS; precipitation of the wettest
quarter possessed no relationship in KM. Isothermality (bio03), and maximum temperature of the
warmest month (bio05) expressed opposite relationships with suitability within Kapokapowai.
Isothermality (bio03) expressed positive linear or linear/quadratic relationships with suitability within
KM and KN, but a negative linear relationship with KS. Maximum temperature of the warmest month
(bio05) expressed negative linear or quadratic/linear relationships with suitability within KM and KS but
a positive linear relationship within Kapokaowai South (Supplemental Fig. 1).
Predictions using our 'Anthropocene' environmental variables for KS within Te Waipounamu
revealed areas of high suitability along the coast (Westland, Nelson and Marlborough provinces) (Fig
3D). Suitability was highest for KM across the Southern Alps (interior Nelson and Marlborough
provinces) (Fig. 3A). Predictions for Kapokaowai North within Te Ika-a-Māui revealed areas of high
suitability within the interior (Wellington and Hawke's Bay, along the Tararura, Ruahine, and
Kaimanawa Mountain Ranges), as well as the Northern portion of the island within Auckland (Fig 3G).
Threshold predictions reflect our 'cloglog' predictions (See Supplemental Fig. 3).
Predictions generated from our paleoclimate variables demonstrated variability in suitability over
time for Kapokapowai compared to our modern-day 'Anthropocene' variables. For KM, suitability
expands within the mid-Holocene (8.3 – 4.2ka), encompassing most of Te Waipounamu except for the
coastal margins (Fig 3H.). Projection to the Te Ika-a-Māui within the mid-Holocene indicates suitable
habitat within the island's center (Te Ika-a-Māui Volcanic Plateau, Tararua Range, and Raukumara
Range) (Fig 3H.). Within the LGM, suitability contracts within Te Waipounamu, with suitable habitat
only being located within a small portion of the Kā Tiritiri-o-te-Moana (Southern Alps) and Kaikōura
Mountain Ranges (Fig 3I.). Within Te Ika-a-Māui, suitability expands, encompassing most of the western
and southern mountain ranges. Although a land bridge connected both Islands during the LGM (Te
Moana o Raukawa), suitability was low (Fig 3I.) while our threshold predictions suggest suitable habitat
across the strait (See Supplemental Fig. 4).
For KS within Te Waipounamu, suitability contracts within the mid-Holocene, being restricted to
a small portion of the coast (northern Westland) (Fig 3E.). When projected to Te Ika-a-Māui, suitability is
high within the Northern Cape and southwestern cape (Taranaki Province), and Auckland (Fig 3E).
Within the LGM, suitability expands encompassing the northern coastal regions of Te Waipounamu
(Westland and Nelson), as well as the Northern Cape, Taranaki Province, and Auckland within Te Ika-a-
Māui (Fig 3F.). Suitability across the Te Moana o Raukawa is intermediate, with a few pockets of high
suitability (Fig 3F.) while our threshold predictions suggest suitable habitat across the strait (See
Supplemental Fig. 4).
For KN within Te Ika-a-Māui, suitability contracts within the mid-Holocene, with only a few
pockets of high suitability within the southern coasts (Tararua Ranges), and Auckland (Fig. 3B.). When
projected to Te Waipounamu, suitability is highest along the coastal ranges of southern Westland and
Otago. Within the LGM, suitability expands to include most of the southern Mountain Ranges within Te
Ika-a-Māui, as well as Kaimanawa Mountain Ranges within the middle of the Island (Fig 3C). Within Te

Waipounamu, suitability slightly expands southward into the coastal ranges of Otago (Fig 3D). Suitability
within the Te Moana o Raukawa remains intermediate with a small pocket of high suitability off the coast
of Taranaki (Fig 3D.) while our threshold predictions suggest suitable habitat across the strait (See
Supplemental Fig. 4).
531
Ordination Analysis
Environmental variables for KN and KSwithin Te Ika-a-Māui and Te Waipounamu expressed an
intermediate degree of overlap in ordination analysis, but were statistically more similar than random
(Fig. 4). Within our PCA analysis, the first principal component explained ~34% of the variance with
PC2 explaining ~32%. Isothermality (bio03), temperature seasonality (bio04), and precipitation of the
coldest quarter (bio19) loaded strongly on PC1, while precipitation of the warmest and coldest quarters
(bio18 and bio19) loaded strongly on PC2 (Table 4). Environmental niche overlap for Kapokapowai was
intermediate between Te Ika-a-Māui and Te Waipounamu (Schoener's D = 0.24), with niche similarity
tests showing observed overlaps being higher than 95% of simulated overlaps ($P = 0.01$), supporting
niche similarity between both islands (Fig. 4).
542
Figure 3: Niche Models of Kapokapowai Populations. MaxEnt predictions for Kapokapowai projected to
the modern day using our 'Anthropocene variables', mid-Holocene (Northgrippian: 4.2 – 8.326ka)
generated from the PaleoClim dataset (Brown et al. 2018), and Last Glacial Maximum (LGM)(21ka) and
Last Interglacial (LIG)(130ka) generated from the CHELSA database (Karger et al. 2017). Predictions
were derived from the optimal model using the criterion of mean valuation AUC (AUC_{val}) values being
the highest, and 10% omission rate being the lowest. Predictions were transformed using the 'cloglog'
function in Maxent v3.4.4 (Phillips et al., 2017), in which raw values are converted to a range of 0 - 1 to
approximate a probability of occurrence. Brighter colors (yellow, green, blue) indicate areas of higher
suitability (higher probability of occurrence), while darker colors (violet) indicate areas of lower
suitability (lower probability of occurrence).
553
Figure 4: *Niche Overlap of Kapokapowai North and Kapokapowai South.* Ordination plot (PCA), niche
overlap, correlation circle, and niche similarity tests showing niche differences between Kapokapowai
North and South from Te Ika-a-Māui (North) and Te Waipounamu (South). Ordination plot represents
principal component points of occurrence and background environmental values using type 2 scaling
(Mahalanobis distance). PCA points for Te Waipounamu are in blue, while Te Ika-a-Māui is red. Solid
contour lines in the niche overlap illustrate full range (100%) of climate space ('fundamental niche')
while dashed lines indicate 50% confidence intervals. Contour lines of Te Waipounamu are in orange,
and Te Ika-a-Māui is blue. Shading shows the density of species occurrences per grid cell of kernel
density analysis ('realized niche'), and violet pixels show niche stability (climate conditions occupied in
both time periods). Orange shading indicates climate conditions only occupied by Te Waipounamu, while
blue indicates climate conditions only occupied by Te Ika-a-Māui. Correlation circle indicates climactic
variable loadings on the PCA space. Length and direction of arrows indicates influence and distribution of
variables within PCA space. Bioclimatic variables which follow Worldclim v2 (Fick and Hijmans 2017).
Histograms represent observed (red bar) and randomly simulated overlaps of niche similarity. The
observed overlap is higher than 95% ($p < 0.05$) for both analyses supporting the niche similarity across
both islands.
570

Ancestral State Reconstruction
Figure 5: *Ancestral State Reconstruction of the Maximum Elevation inhabited by Petaluridae.* Maximum
likelihood (ML) reconstructions and stochastic character mapping (SM) are shown both with (ML A, SM
B) and without (ML C, SM D) the fossil taxa *Argentinopetala archangelskyi*, which is considered to be
sister to *P. raptor* based upon wing characteristics (Tolman et al., 2024). In all models, all three ancestral
states (high, medium, and low maximum elevation) are equally likely as the ancestral state at each node,
including at the root.
578
The ER model was strongly supported by AIC (Supplemental Table 6) both with and without *A.*
archangelskyi, indicating gains and losses of different elevation habitats are equally probable across the
tree. All nodes were equivalent with near equal support for each elevational maxim, including the
ancestral state of Petaluridae, in both maximum likelihood models and using stochastic character
mapping, both with and without *A. archangelskyi* (Fig. 5). Across 1000 stochastic maps under the
averaged model with *A. archangeslskyi* included in the tree, we inferred on average 235 instances of
medium to high elevation and high to medium elevation with transition and 236 medium to low and low
to medium transitions, meaning each of these transitions are expected to occur once every 2.9 million
years for a given branch of Petaluridae on average. We inferred 49 high to low and low to high elevation
transitions each, averaging one of each transition every 14.5 million years for a given branch of
Petaluridae on average. The expected frequency of transitions per million years for a given branch of
Petaluridae did not change with *A. archangelskyi* removed.
591
Discussion
Historical Island Geography Underlies a Species Complex
As hypothesized, our results demonstrate that uplift and oceanic barriers have shaped the evolution of
Kapokapowai. The divergence time analysis suggested that the MRCA (2.6-15 Ma) of all sampled
Kapokapowai could very well have coincided with the accelerated tectonic uplift which occurred ~5Ma,
or the more gradual uplift from 10-5Ma (Fig. 1E). Intriguingly, a strong bottleneck in the ancestral
lineage of KS was identified between 2.5-10Ma from a reference genome (Tolman et al., 2023) using the
Pairwise Sequential Coalescent (PSMC; Liu & Hansen, 2017). This bottleneck could plausibly be
reflective of speciation between KM and KN+KS. KM has higher genetic diversity than either KN or KS
(Fig. 1A,B) supporting the hypothesis that this bottleneck was caused by a founder effect similar to many
other systems (Anopheles gambiae 1000 Genomes Consortium et al., 2017; Auton et al., 2015; B. Li et
al., 2024).
The genetic differentiation between KN and KS is much smaller than the differentiation between
either population and KM (Table 2). Although Te Moana o Raukawa (Cook Strait) is currently a barrier to
the dispersal of Kapokapowai, our species distribution models suggest that this has not always been the
case (Fig. 4, Supplemental Fig. 4). Perhaps Te Moana o Raukawa has not been a barrier to the dispersal
of Kapokapowai for a sufficient period of time for a strong genetic structure to evolve. On the other hand,
the peaks of Kā Tiritiri-o-te-Moana have not contained suitable habitat for KN or KS in any of the tested
time periods, and have likely served as a longer barrier to the dispersal of these populations, shaping the
genetic structure currently observed in Kapokapowai. The physical boundary of the mountains is further
compounded by the fact that the alpine habitat has shortened the flight season of KM, compared to KN
and KS (L. S. Wolfe, 1952).

The data we have here support a hypothesis in which all sampled Kapokapowai are descendents of a population that colonized alpine habitats (the most supported pattern in the fauna of Aotearoa (Buckley et al., 2024), although there is speculation that some lineages may have ridden tectonic uplift to higher elevations (Heads, 2017)), while KN and KS descend from populations that later moved into lowland habitats. Our ancestral state reconstruction analysis of Petaluridae would support the former, as across 1000 simulations, transitions from high elevation to low and medium elevational maxims occurred on average every 2.9 million years on each branch. It is entirely plausible that such flexibility in adjusting to different elevations is a common trait in lineages that have persisted on Zealandia for tens of millions of years.This hypothesis of recolonization of the lowlands is not without precedent, as plant lineages have also been hypothesized to have colonized mountains, and then recolonized the lowlands of Aoteora (Thomas et al., 2023). A demographic model for KM from a reference genome, and revisiting the specimens sampled here with genome-wide resequencing data could be used to generate a demographic model for all populations of Kapokapowai, and a historical demographic model of dispersal and gene flow between populations using the site-frequency-spectrum (Noskova et al., 2023), both of which could be used to robustly test this hypothesis.

Gene Flow in the Presence of Geographic Barriers and Implications for Species Delimitation

Evidence of Current gene flow

Despite the aforementioned geographic barriers, the PCA (Fig. 1A), admixture (Fig. 1B), and introgression analyses all suggest there is some level of interbreeding and movement between populations. Neither barrier seems to be entirely restricting gene flow, as evidenced by lower fixation (F_{ST} < 0.30) than might be expected given > 5 million years of separation between KM and KN and KS (Fig. 1E), and evidence of introgression between KS and KM.

One individual KM notably inherited 75% of its ancestry from ancestral population one, which is unique to KM, and 25% of its ancestry from ancestral population three (Fig. 1B). Thus, this individual is likely the offspring of an F1-hybrid back-crossed with an individual with 100% of its ancestry from population one, suggesting the genetic populations can breed with one another, and gene flow may still occur in the present.

Implications for species delimitation

Given the dynamics of gene flow, current species delimitations in this complex are difficult to make. Given the genetic differentiation between KM and KN and KS (F_{ST} >.23), our identification of a diagnostic characteristic of the curvature on the horns of KM (Fig. 2), and the strong posterior support (>99%) of the currently described species (*U. carovei* comprised of KN and KS and *U. chiltoni* comprised of KM) in our species delimitation analysis, we suggest that current species delimitations remain unchanged. Although KN and KS do show some genetic differentiation, they do not show evidence of niche divergence, and we do not believe that they merit a new species description at this point in time.

Generating Species Distribution Models for Island Species

Although the utilization of GBIF occurrences in conjunction with genomic analyses provide the opportunity for highly robust niche models which have the potential to match population admixture or structuring among populations of odonates, a consequence is the underfitting of model performance due to the abundance, range, and density of the species itself. ENM results for KN revealed low AUC values,

which is a continuous metric that provides discriminative ability in false positives and false negatives
(Phillips et al. 2017). Since KN is found throughout the entirety of Te Ika-a-Māui, our models were
unable to discern gradients of high and low suitability, resulting in poor model output. This pattern is also
observed in our AICc value (Supplementary Table S2). Our thinned dataset consisted of 239 occurrences,
sufficient to generate a robust model; the widespread distribution throughout Te Ika-a-Māui contributed
to high AICc scores.
ENMs in the context of island biogeography is an area of ongoing study, as model extrapolation
becomes poor for species endemic to small or singular islands (Sutherland et al. 2021, Goodman et al.
2024), limiting the incorporation of novel climatic conditions that could improve model fit. Furthermore
AUC values may be skewed if only a few climatic variables drive the model, leading to oversimplified
relationships with suitability (Phillips et al. 2017), as evidenced by the number of nonzero lambda
coefficients within our KN model (2)(See Supplementary Table S2).
Interestingly, our 10-percentile omission rate metrics were more robust (0.16). The 10-percentile
omission rate metric excludes the lowest 10 percent of raw occurrence values, thus creating a threshold
within suitability. Our lower omission rates suggest minimal outliers in our dataset, suggesting that our
predictions for KN on Te Ika-a-Māui are not overly generalist.
An aspect to consider in paleo-ecological modeling (PaleoENM) is the use of multiple lines of
evidence to assess model fidelity. Incorporating supplemental analyses such as response curves and
ordination analyses provide clarity to model over- or under-extrapolation, which can be interpreted as
either model error or accuracy. Response curves demonstrate if the full environmental range of the
species has been sampled for each species or population, while ordination analyses allow
multidimensional visualization of the niche outside of a spatial context. In the case of our data, response
curves, albeit jagged, map the full relationship of each climatic variable within the model, while our
ordination analysis demonstrates statistical differences among the KN and KS populations. Overall,
although genetic divergence was low among KN and KS populations, the ecological and distributional
differences among both populations are the result in differences in climatic conditions, and not a result of
poor model performance because of inappropriate apriori grouping of populations based on their presence
on Te Ika-a-Māui or Te Waipounamu.
686
Conclusion
To look at species diversity and the potential for speciation in a "relict" group such as the petaltail
dragonflies offers a unique perspective in evolutionary biology, as well as island ecology. Analyses
suggest that this family is extremely old, even for an order as ancient as Odonata, with some extant
species persisting as independent evolutionary lineages since the Cretaceous era (Kohli et al., 2021;
Suvorov et al., 2022; E. R. Tolman et al., 2024). With their reliance on fen habitats, Petaluridae are also
somewhat limited in their range, compared to more generalist dragonflies that use ponds or rivers to
complete their life cycle. Despite this, our results demonstrate that speciation in Petaluridae is associated
with shifts to habitats at different elevations, in both Kapokapowai and other groups of petaltails, which
may be associated with their long persistence. While species of Kapokapowai have potentially been
separated for significant periods of time, either across islands or in different habitats within an island, it
may be that their adaptations to these specialized fen habitats has limited the genetic variability that
would lead to speciation. This may help to explain why this group, despite its age, continues to be the
smallest family within Anisoptera.
701

References

Andress, R. (1998). *Description of the larva of Petalura ingentissima Tillyard, 1907 (Anisoptera: Petaluridae)*. http://archive.org/details/odonatologica-27-353-359

Anopheles gambiae 1000 Genomes Consortium, Data analysis group, Partner working group, Sample collections—Angola:, Burkina Faso:, Cameroon:, Gabon:, Guinea:, Guinea-Bissau:, Kenya:, Uganda:, Crosses:, Sequencing and data production, Web application development, & Project coordination. (2017). Genetic diversity of the African malaria vector Anopheles gambiae. *Nature*, *552*(7683), 96–100. https://doi.org/10.1038/nature24995

Auton, A., Abecasis, G. R., Altshuler, D. M., Durbin, R. M., Abecasis, G. R., Bentley, D. R., Chakravarti, A., Clark, A. G., Donnelly, P., Eichler, E. E., Flicek, P., Gabriel, S. B., Gibbs, R. A., Green, E. D., Hurles, M. E., Knoppers, B. M., Korbel, J. O., Lander, E. S., Lee, C., … National Eye Institute, N. (2015). A global reference for human genetic variation. *Nature*, *526*(7571), 68–74. https://doi.org/10.1038/nature15393

Barreto, E., Rangel, T. F., Pellissier, L., & Graham, C. H. (2021). Area, isolation and climate explain the diversity of mammals on islands worldwide. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1965), 20211879. https://doi.org/10.1098/rspb.2021.1879

Bercovich, U., M. S. Rasmussen, Z. Li, C. Wiuf, and A. Albrechtsen. 2025. Measuring linkage disequilibrium and improvement of pruning and clumping in structured populations. Genetics 229:iyaf009.

Booth, T. H. (2022). Checking bioclimatic variables that combine temperature and precipitation data before their use in species distribution models. *Austral Ecology*, *47*(7), 1506–1514. https://doi.org/10.1111/aec.13234

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M. A., Rambaut, A., & Drummond, A. J. (2014). BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. PLOS Computational Biology, 10(4), e1003537. https://doi.org/10.1371/journal.pcbi.1003537

Bouckaert, R. R. (2022). An Efficient Coalescent Epoch Model for Bayesian Phylogenetic Inference. *Systematic Biology*, *71*(6), 1549–1560. https://doi.org/10.1093/sysbio/syac015

Brown, J. L., Hill, D. J., Dolan, A. M., Carnaval, A. C., & Haywood, A. M. (2018). PaleoClim, high spatial resolution paleoclimate surfaces for global land areas. *Scientific Data*, *5*(1), Article 1. https://doi.org/10.1038/sdata.2018.254

Buckley, T. R., Hoare, R. J. B., & Leschen, R. A. B. (2024). Key questions on the evolution and biogeography of New Zealand alpine insects. *Journal of the Royal Society of New Zealand*, *54*(1), 30–54. https://doi.org/10.1080/03036758.2022.2130367

Cairns, J., I. R. C. Baird, F. Johnson, L. Noble, and W. Glamore. 2025. Detecting□: Novel eDNA Method and Water Balance Modelling Insights Enable Improved Catchment Management and Conservation Outcomes. Ecohydrology 18:e70013.

Conway, M., & Olsen, B. J. (2019). Contrasting drivers of diversification rates on islands and continents across three passerine families. *Proceedings. Biological Sciences*, *286*(1915), 20191757. https://doi.org/10.1098/rspb.2019.1757

Córdoba-Aguilar, A., Beatty, C. D., & Bried, J. T. (2022). Introduction to Dragonflies and Damselflies, Second Edition. In A. Cordoba-Aguilar, C. Beatty, & J. Bried (Eds.), Dragonflies and Damselflies: Model Organisms for Ecological and Evolutionary Research (p. 0). Oxford University Press. https://doi.org/10.1093/oso/9780192898623.003.0001

Cooper, A., and R. A. Cooper. 1997. The Oligocene bottleneck and New Zealand biota: genetic record of a past environmental crisis. Proceedings of the Royal Society of London. Series B: Biological Sciences 261:293–302. Royal Society.

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

Di Cola, V., Broennimann, O., Petitpierre, B., Breiner, F. T., D'Amen, M., Randin, C., Engler, R., Pottier, J., Pio, D., Dubuis, A., Pellissier, L., Mateo, R. G., Hordijk, W., Salamin, N., & Guisan, A. (2017). ecospat: An R package to support spatial analyses and modeling of species niches and distributions. *Ecography*, *40*(6), 774–787. https://doi.org/10.1111/ecog.02671

Douglas, J., & Bouckaert, R. (2022). Quantitatively defining species boundaries with more efficiency and more biological realism. *Communications Biology*, *5*(1), 1–10. https://doi.org/10.1038/s42003-022-03723-z

Dray, S., & Dufour, A.-B. (2007). The ade4 Package: Implementing the Duality Diagram for Ecologists. *Journal of Statistical Software*, *22*, 1–20. https://doi.org/10.18637/jss.v022.i04

Durrant, R. (2024). *Invertebrate Justice: Extending The Boundaries of Non-Speciesist Green Criminology*. Springer Nature.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, *37*(12), 4302–4315. https://doi.org/10.1002/joc.5086

Fong, H. M. (2024). *A revision of the genus Mecodema Blanchard 1853 (Coleoptera: Carabidae) species groups ducale and laterale from the South Island, Aotearoa New Zealand* [Thesis, ResearchSpace@Auckland]. https://researchspace.auckland.ac.nz/handle/2292/70309

GBIF.org (1 May 2025) GBIF Occurrence Download https://doi.org/10.15468/dl.t648we

Gleditsch, J., Behm, J., Ellers, J., Jesse, W., & Helmus, M. (2022). *Contemporizing island biogeography theory with anthropogenic drivers of species richness.* https://doi.org/10.22541/au.163955172.21711050/v3

Goodman, A., Tolman, E., Uche-Dike, R., Abbott, J., Breinholt, J. W., Bybee, S., Frandsen, P. B., Gosnell, J. S., Guralnick, R., Kalkman, V. J., Kohli, M., Lontchi, J. F., Lupiyaningdyah, P., Newton, L., & Ware, J. L. (2023). Assessment of targeted enrichment locus capture across time and museums using odonate specimens. Insect Systematics and Diversity, 7(3), 5. https://doi.org/10.1093/isd/ixad011

Goodman, A. M., Beatty, C. D., Büsse, S., Ubukata, H., Miyazaki, T., Blair, M. E., & Ware, J. L. (2024). Paleoecological niche modeling of Epiophlebia (Epiophlebioptera: Epiophlebiidae) reveals continuous distribution during the Last Glacial Maximum. *International Journal of Odonatology*, *27*, 60–76. https://doi.org/10.48156/1388.2024.1917262

Grandcolas, P., R. Nattier, and S. Trewick. 2014. Relict species: a relict concept? Trends in Ecology & Evolution 29:655–663.

Grant, E. A. (2014). *Bronze as a non-customary intervention in the interpretation of insects from the natural world of Māori: An exhibition report presented in partial fulfilment of the requirements for the degree of Doctor of Philosophy in Fine Arts at Massey University, Palmerston North, Aotearoa, New Zealand* [Massey University]. http://hdl.handle.net/10179/6512

Harris, C. R., Millman, K. J., Van Der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Van Kerkwijk, M. H., Brett, M., Haldane, A., Del Río, J. F., Wiebe, M., Peterson, P., … Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2

Heads, M. J. (2017). *Biogeography and evolution in New Zealand*. CRC press, Taylor & Francis group.

Hijmans R (2025). *raster: Geographic Data Analysis and Modeling*. R package version 3.6-32, https://github.com/rspatial/raster.

Hollister, J., T. Shah, J. Nowosad, A. L. Robitaille, M. W. Beck, and M. Johnson. 2023. elevatr: Access Elevation Data from Various APIs.

Jenkins, T. L. (2024). mapmixture: An R package and web app for spatial visualisation of admixture and population structure. *Molecular Ecology Resources*, e13943. https://doi.org/10.1111/1755-0998.13943

801   Jessen, T. D., N. C. Ban, N. X. Claxton, and C. T. Darimont. 2022. Contributions of Indigenous
802   Knowledge to ecological and evolutionary understanding. Frontiers in Ecology and the
803   Environment 20:93–101.
804   Jones, L. A., W. Gearty, B. J. Allen, K. Eichenseer, C. D. Dean, S. Galván, M. Kouvari, P. L.
805   Godoy, C. S. C. Nicholl, L. Buffan, E. M. Dillon, J. T. Flannery-Sutherland, and A. A. Chiarenza.
806   2023. palaeoverse: A community-driven R package to support palaeobiological analysis. Methods
807   in Ecology and Evolution 14:2205–2215

808 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E.,
809   Linder, H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's land surface
810   areas. *Scientific Data*, *4*(1), Article 1. https://doi.org/10.1038/sdata.2017.122

811 Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H., & Zimmermann, N. E. (2023). CHELSA-
812   TraCE21k – high-resolution (1 km) downscaled transient temperature and precipitation
813   data since the Last Glacial Maximum. *Climate of the Past*, *19*(2), 439–456.
814   https://doi.org/10.5194/cp-19-439-2023

815 Kass, J. M., Muscarella, R., Galante, P. J., Bohl, C. L., Pinilla-Buitrago, G. E., Boria, R. A., Soley-
816   Guardia, M., & Anderson, R. P. (2021). ENMeval 2.0: Redesigned for customizable and
817   reproducible modeling of species' niches and distributions. *Methods in Ecology and Evolution*,
818   *12*(9), 1602–1608. https://doi.org/10.1111/2041-210X.13628

819 Kass, J. M., Vilela, B., Aiello-Lammens, M. E., Muscarella, R., Merow, C., & Anderson, R. P. (2018).
820   Wallace: A flexible platform for reproducible modeling of species niches and distributions built
821   for community expansion. *Methods in Ecology and Evolution*, *9*(4), 1151–1156.
822   https://doi.org/10.1111/2041-210X.12945

823 Kennedy, C. H. (1922). The morphology of the penis in the genus Libellula (Odonata). *Entomology*
824   *Newsletter*, *33*, 33–40.

825 Kohli, M., Letsch, H., Greve, C., Béthoux, O., Deregnaucourt, I., Liu, S., Zhou, X., Donath, A., Mayer,
826   C., Podsiadlowski, L., Gunkel, S., Machida, R., Niehuis, O., Rust, J., Wappler, T., Yu, X., Misof,
827   B., & Ware, J. (2021). Evolutionary history and divergence times of Odonata (dragonflies and
828   damselflies) revealed through transcriptomics. *iScience*, *24*(11), 103324.
829   https://doi.org/10.1016/j.isci.2021.103324

830 Kuznetsova, V. G., & Golub, N. V. (2020). A checklist of chromosome numbers and a review of
831   karyotype variation in Odonata of the world. *Comparative Cytogenetics*, *14*(4), 501–540.
832   https://doi.org/10.3897/CompCytogen.v14i4.57062

833 L. S. Wolfe. (1952). A study of the genus Uropetala Selys (Order Odonata) from New Zealand.
834   *Transactions of the Royal Society of New Zealand*, *80*, 245–275.

835 Leschen, R., Collier, K., Death, R., Harding, J., & Smith, B. (2023). Systematics expertise and taxonomic
836   status of New Zealand's freshwater insects. *New Zealand Science Review*, *73*(3–4), 92–95.
837   https://doi.org/10.26686/nzsr.v73i3-4.8534

838 Lewis, K. B., Carter, L., & Davey, F. J. (1994). The opening of Cook Strait: Interglacial tidal scour and
839   aligning basins at a subduction to transform plate edge. *Marine Geology*, *116*(3), 293–312.
840   https://doi.org/10.1016/0025-3227(94)90047-7

841 Li, B., Duan, Y., Du, Z., Wang, X., Liu, S., Feng, Z., Tian, L., Song, F., Yang, H., Cai, W., Lin, Z., & Li,
842   H. (2024). Natural selection and genetic diversity maintenance in a parasitic wasp during
843   continuous biological control application. *Nature Communications*, *15*(1), 1379.
844   https://doi.org/10.1038/s41467-024-45631-2

845 Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform.
846   *Bioinformatics*, *25*(14), 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

847 Liu, S., & Hansen, M. M. (2017). PSMC (pairwise sequentially Markovian coalescent) analysis of RAD
848   (restriction site associated DNA) sequencing data. *Molecular Ecology Resources*, *17*(4), 631–641.
849   https://doi.org/10.1111/1755-0998.12606

Lobo, J. M., Jiménez-Valverde, A., & Real, R. (2008). AUC: A misleading measure of the performance of predictive distribution models. *Global Ecology and Biogeography*, *17*(2), 145–151. https://doi.org/10.1111/j.1466-8238.2007.00358.x

Malinsky, M., Matschiner, M., & Svardal, H. (2021). Dsuite—Fast D-statistics and related admixture evidence from VCF files. *Molecular Ecology Resources*, *21*(2), 584–595. https://doi.org/10.1111/1755-0998.13265

May, M. L. (1997). Reconsideration of the status of the genera Phyllomacromia and Macromia (Anisoptera: Corduliidae). *Odonatologica*, *26*(4), 405–414.

McCulloch, G. A., and J. M. Waters. 2019. Phylogenetic divergence of island biotas: Molecular dates, extinction, and "relict" lineages. Molecular Ecology 28:4354–4362.

McKinney, W. (2010). *Data Structures for Statistical Computing in Python*. 56–61. https://doi.org/10.25080/Majora-92bf1922-00a

Naimi, B. (2023). *usdm: Uncertainty Analysis for Species Distribution Models* (Version 2.1-6) [Computer software]. https://cran.r-project.org/web/packages/usdm/index.html

Noskova, E., Abramov, N., Iliutkin, S., Sidorin, A., Dobrynin, P., & Ulyantsev, V. (2023). *GADMA2: More efficient and flexible demographic inference from genetic data* (p. 2022.06.14.496083). bioRxiv. https://doi.org/10.1101/2022.06.14.496083

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., … Weedon, J. (2024). *vegan: Community Ecology Package* (Version 2.6-6.1) [Computer software]. https://cran.r-project.org/web/packages/vegan/index.html

O'Neill, S. B., Chapple, D. G., Daugherty, C. H., & Ritchie, P. A. (2008). Phylogeography of two New Zealand lizards: McCann's skink (*Oligosoma maccanni*) and the brown skink (*O. zelandicum*). *Molecular Phylogenetics and Evolution*, *48*(3), 1168–1177. https://doi.org/10.1016/j.ympev.2008.05.008

Peterson, A. T., & Lieberman, B. S. (2012). Species' Geographic Distributions Through Time: Playing Catch-up with Changing Climates. *Evolution: Education and Outreach*, *5*(4), 569–581. https://doi.org/10.1007/s12052-012-0385-2

Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E., & Blair, M. E. (2017). Opening the black box: An open-source release of Maxent. *Ecography*, *40*(7), 887–893. https://doi.org/10.1111/ecog.03049

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *American Journal of Human Genetics*, *81*(3), 559–575.

Radosavljevic, A., & Anderson, R. P. (2014). Making better Maxent models of species distributions: Complexity, overfitting and evaluation. *Journal of Biogeography*, *41*(4), 629–643. https://doi.org/10.1111/jbi.12227

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior Summarization in Bayesian Phylogenetics Using Tracer 1.7. *Systematic Biology*, *67*(5), 901–904. https://doi.org/10.1093/sysbio/syy032

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other things). Methods in Ecology and Evolution 3:217–223.

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Schoener, T. W. (1968). The Anolis Lizards of Bimini: Resource Partitioning in a Complex Fauna. *Ecology*, *49*(4), 704–726. https://doi.org/10.2307/1935534

Shepherd, L., Simon, C., Langton-Myers, S., & Morgan-Richards, M. (2024). Insights into Aotearoa New Zealand's biogeographic history provided by the study of natural hybrid zones. *Journal of the Royal Society of New Zealand*, *54*(1), 55–74. https://doi.org/10.1080/03036758.2022.2061020

Skotte, L., Korneliussen, T. S., & Albrechtsen, A. (2013). Estimating individual admixture proportions from next generation sequencing data. *Genetics*, *195*(3), 693–702. https://doi.org/10.1534/genetics.113.154138

Scotese, C. R., and N. M. Wright. 2018. PALEOMAP Paleodigital Elevation Models (PaleoDEMS) for the Phanerozoic. Zenodo.

Stevens, M. I., & Hogg, I. D. (2004). Population genetic structure of New Zealand's endemic corophiid amphipods: Evidence for allopatric speciation. *Biological Journal of the Linnean Society*, *81*(1), 119–133. https://doi.org/10.1111/j.1095-8312.2004.00270.x

Stoltz, M., Baeumer, B., Bouckaert, R., Fox, C., Hiscott, G., & Bryant, D. (2021). Bayesian Inference of Species Trees using Diffusion Models. *Systematic Biology*, *70*(1), 145–161. https://doi.org/10.1093/sysbio/syaa051

Sutherland, L. N., G. S. Powell, and S. M. Bybee. 2021. Validating species distribution models to illuminate coastal fireflies in the South Pacific (Coleoptera: Lampyridae). Sci. Rep. 11: 17397.

Suvorov, A., Scornavacca, C., Fujimoto, M. S., Bodily, P., Clement, M., Crandall, K. A., Whiting, M. F., Schrider, D. R., & Bybee, S. M. (2022). Deep Ancestral Introgression Shapes Evolutionary History of Dragonflies and Damselflies. *Systematic Biology*, *71*(3), 526–546. https://doi.org/10.1093/sysbio/syab063

Taylor, R. (1855). *Te Ika a Maui, or, New Zealand and its inhabitants: With the geology, natural history, productions, and climate of the country, &c* (Vol. 1855, pp. 1–554). https://doi.org/10.5962/bhl.title.161768

Taylor, R. (with Smithsonian Libraries). (1870). *Te Ika a Māui, or, New Zealand and its inhabitants: Illustrating the origin, manners, customs, mythology, religion, rites, songs, proverbs, fables and language of the Māori and Polynesian races in general□; together with the geology, natural history, productions, and climate of the country*. London□: William Macintosh□; Wanganui, N.Z.□: H. Ireson Jones. http://archive.org/details/teikamauiornewze00tayl

Thomas, A., Meudt, H. M., Larcombe, M. J., Igea, J., Lee, W. G., Antonelli, A., & Tanentzap, A. J. (2023). Multiple origins of mountain biodiversity in New Zealand's largest plant radiation. *Journal of Biogeography*, *50*(5), 947–960. https://doi.org/10.1111/jbi.14589

Tolman, E. R., Beatty, C. D., Frandsen, P. B., Bush, J., Bruchim, O. R., Driever, E. S., Harding, K. M., Jordan, D., Kohli, M. K., Park, J., Park, S., Reyes, K., Rosari, M., Ryu, J. L., Wade, V., & Ware, J. L. (2023). *Newly Sequenced Genomes Reveal Patterns of Gene Family Expansion in select Dragonflies (Odonata: Anisoptera)* (p. 2023.12.11.569651). bioRxiv. https://doi.org/10.1101/2023.12.11.569651

Tolman, E. R., Beatty, C. D., Kohli, M. K., Abbott, J., Bybee, S. M., Frandsen, P. B., Stephen Gosnell, J., Guralnick, R., Kalkman, V. J., Newton, L. G., Suvorov, A., & Ware, J. L. (2024). A molecular phylogeny of the Petaluridae (Odonata: Anisoptera): A 160-Million-Year-Old story of drift and extinction. *Molecular Phylogenetics and Evolution*, *200*, 108185. https://doi.org/10.1016/j.ympev.2024.108185

Trewick, S. A., Pilkington, S., Shepherd, L. D., Gibb, G. C., & Morgan-Richards, M. (2017). Closing the gap: Avian lineage splits at a young, narrow seaway imply a protracted history of mixed population response. *Molecular Ecology*, *26*(20), 5752–5772. https://doi.org/10.1111/mec.14323

Veloz, S. D. (2009). Spatially autocorrelated sampling falsely inflates measures of accuracy for presence-only niche models. *Journal of Biogeography*, *36*(12), 2290–2299. https://doi.org/10.1111/j.1365-2699.2009.02174.x

W. J. Winstanley. (1981). AN EMERGENCE STUDY ON UROPETALA CAROVEI CAROVEI (ODONATA: PETALURIDAE) NEAR WELLINGTON, NEW ZEALAND, WITH NOTES ON THE BEHAVIOUR OF THE SUBSPECIES. *Tuatara*, *25*(1).

Wallis, G. P., & Buckley, T. R. (2024). Evolutionary biogeography of Aotearoa. *Journal of the Royal Society of New Zealand*, *54*(1), 1–7. https://doi.org/10.1080/03036758.2023.2260539

Ware, J. L. (2008). *MOLECULAR AND MORPHOLOGICAL SYSTEMATICS OF LIBELLULOIDEA (ODONATA: ANISOPTERA) AND DICTYOPTERA* [PhD Thesis, Rutgers University]. https://rucore.libraries.rutgers.edu/rutgers-lib/24985/PDF/1

Ware, J. L., Beatty, C. D., Herrera, M. S., Valley, S., Johnson, J., Kerst, C., May, M. L., & Theischinger, G. (2014). The petaltail dragonflies (Odonata: Petaluridae): Mesozoic habitat specialists that survive to the modern day. *Journal of Biogeography*, *41*(7), 1291–1300. https://doi.org/10.1111/jbi.12273

Warren, D. L., & Seifert, S. N. (2011). Ecological niche modeling in Maxent: The importance of model complexity and the performance of model selection criteria. *Ecological Applications*, *21*(2), 335–342. https://doi.org/10.1890/10-1171.1

Waters, J. M., & Craw, D. (2006). Goodbye Gondwana? New Zealand Biogeography, Geology, and the Problem of Circularity. *Systematic Biology*, *55*(2), 351–356. https://doi.org/10.1080/10635150600681659

Wilson, E. O. (2009). Island Biogeography in the 1960s: Theory and Experiment. In *The Theory of Island Biogeography Revisited* (pp. 1–12). Princeton University Press. https://doi.org/10.1515/9781400831920.1

Winstanley, W. J., & Rowe, R. J. (1980). The larval habitat of Uropetala carovei carovei (Odonata: Petaluridae) in the North Island of New Zealand, and the geographical limits of the subspecies. *New Zealand Journal of Zoology*, *7*(1), 127–134. https://doi.org/10.1080/03014223.1980.10423769

Wyckhuys, K. A. G., Sanchez Garcia, F. J., Santos, A. M. C., Canal, N. A., Furlong, M. J., Melo, M. C., Gc, Y. D., & Pozsgai, G. (2022). Island and Mountain Ecosystems as Testbeds for Biological Control in the Anthropocene. *Frontiers in Ecology and Evolution*, *10*. https://doi.org/10.3389/fevo.2022.912628

Table 1: Reference table of colonial and Māori names of localities and taxa.

Geographic localities	
<u>Māori name</u>	<u>Colonial name</u>
Aotearoa	New Zealand
Te Moana o Raukawa	Cook Strait
Te Ika-a-Māui	North Island
Te Waipounamu	South Island
Kā Tiritiri-o-te-Moana	Southern Alps
Population Names	
Māori names used in this work	Scientific name and locality
Kapokapowai North (KN)	*Uropetala carovei*, Te Ika-a-Māui

Kapokapowai South (KS)	*Uropetala carovei*, Te Waipounamu (below 1.2 km)
Kapokapowai Montane (KM)	*Uropetala chiltoni*, Kā Tiritiri-o-te-Moana

975
976

Table 2: F_{st} values between putative species of Kapokapowai, using 365 called SNPs. Kapokapowai North (KN), Kapokapowai Montaine (KM), and Kapokapowai South (KS) are abbreviated. The results show stronger segmentation between KM and KN and KS.

Pairwise comparison	Mean F_{ST}	Weighted F_{ST}
KM-KS	0.20	0.23
KM-KN	0.27	0.29
KN-KS	0.05	0.11

977
978

Table 3: gene flow between Populations of Kapokapowai

P1	P2	P3	Dstatistic	Z-score	p-value	f4-ratio	BBAA	ABBA	BABA
KN	KS	KM	0.15	3.74	< 0.01	0.08	25.89	28.52	20.86

Table 3: ABBA-BABA test output generated with Dsuite provides evidence of gene flow between KN and KS, as variants shared by KS and KM are significantly more common than variants shared by KM and KN (p < 0.01).

979
980

Table 4: PCA loadings, importance of variables, percent overlap, percent of unique niche overlap from both islands, environmental similarity (Schoener's D), and results of similarity tests from our EcoSpat analyses.

Species	PC1	PC2	Overlap (%)	Unique North Island (%)	Unique South Island (%)	Schoener's D	Similarity test (p)
Kapokapowai (North and South)	**34.9**	**32.6**	**93**	**6.9**	**13.2**	**24.7**	**0.01**
Environmental	**PC1**	**PC2**	**PC3**				

Variable			
bio 3	-0.51	-0.27	-0.38
bio 4	-0.51	-0.17	-0.48
bio 5	0.11	0.47	-0.54
bio 15	0.24	0.49	-0.37
bio 18	0.38	-0.56	-0.20
bio 19	0.51	-0.35	-0.40

A

Axis 2 (9.3 %)

Axis 1 (56.0 %)

South

Montane

North

B

Anc1 Anc2 Anc3

Latitude

Longitude

Montane

North

South

Admixture proportions

C

D

E

North

South

Montane

Phenes raptor (Outgroup)

Time (Ma)

1. Oligocene "drowning" (~23 Ma)
2. Early Miocene re-emergence (23-16 Ma)
3. Gradual uplift of Kā Tiritiri o te Moana (10-5 Ma)
4. Accelerated uplift of of Kā Tiritiri o te Moana (5 Ma-present)
5. Pleistocene land bridge events (2.6 Ma-10Ka

Kapokapowai Montane **Kapokapowai Montane** **Kapokapowai North** **Kapokapowai North** **Kapokapowai South** **Kapokapowai South**

Predicted Suitability

- <= 0.2
- 0.2 - 0.4
- 0.4 - 0.6
- 0.6 - 0.8
- > 0.8