Coordination of flux-related leaf and root traits impacts forest community assembly

Zeqing Ma[1*], Mengcheng Duan[1,5], Lawren Sack[2], Chengen Ma[1], Le Li[1], M. Luke McCormack[3], Mingzhen Lu[4], Dali Guo[1]

[1]Qianyanzhou Ecological Research Station, Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China
[2]Department of Ecology and Evolutionary Biology, University of California Los Angeles; Los Angeles, California, 90095, USA
[3]The Center for Tree Science, The Morton Arboretum, Lisle, Illinois, 60532, USA
[4]Department of Environmental Studies, New York University, New York City, New York 10003, USA
[5]University of Chinese Academy of Sciences, Beijing 100049, China

*Corresponding authors
Zeqing Ma, mazeqing@gmail.com, ORCID: https://orcid.org/0000-0003-1197-8066;

E-mail addresses (ORCID):
Z.M.: mazq@igsnrr.ac.cn (0000-0003-1197-8066)
L.S.: lawrensack@ucla.edu (0000-0002-7009-7202)
L.M.: mingzhen.lu@nyu.edu (0000-0002-8707-8745)
M.L.M.: lmccormack@mortonarb.org (0000-0002-8300-5215)

Running title: Linking leaf and root traits to community structure
Competing Interest Statement: The authors declare no competing interests.
Classification: Biological Sciences (Ecology)

Keywords: Above- and belowground, plant functional traits, leaf vein density, ecological strategy, hydraulics, root, woody angiosperms, community composition.

Abstract

Understanding the functions and coordination of plant traits is critical for predicting how diverse species respond to climate change. According to hydraulic and economic theories, leaves and roots—key organs for resource acquisition—are expected to function in coordination, such that species with faster resource utilization would possess leaf and root traits that facilitate rapid carbon, nutrient, and water uptake and fluxes. However, there has been limited evidence supporting leaf-root trait coordination and a role for that coordination on community structure. Here, we measured 13 leaf and root functional traits for 101 woody species from six tropical and subtropical forests, and assessed coordination and its association with community dominance. Hydraulic traits, such as leaf vein density and root vessel density, were coordinated between organs and showed compensation trade-offs between traits within organs, such as, leaf vein density and diameter. Economic traits relating to composition, such as nitrogen concentration, were coordinated between organs, whereas economic structural traits were decoupled, such as leaf mass per area and specific root length. Overall, hydraulic traits and economics traits were partially independent. The coordination of flux-related leaf and root traits was associated with ectomycorrhizal symbiosis and with dominance within the community. These findings indicate how trait organization within and across organs contributes to optimal whole plant function, with implications for performance in natural communities.

Main text

Leaves and roots are the organs through which plants acquire light energy, carbon, water, and nutrients, the fuel and material for the growth of individuals and ecosystems. As the two key resource-acquiring organs, their coordination for resource uptake has been predicted in hydraulic and economic theories (Tyree & Zimmermann 2002; Reich 2014; Weemstra *et al.* 2023). However, evidence of the functional coordination of leaves and roots based on associations between their traits has been scarce and controversial (Bergmann *et al.* 2017; Carmona *et al.* 2021; Weigelt *et al.* 2021). Thus, extensive research has demonstrated tissue nitrogen (N) correlations between roots and leaves (Tjoelker *et al.* 2005; Westoby & Wright 2006; Geng *et al.* 2014; Weigelt *et al.* 2023). However, controversies persist regarding how leaf and root structural and hydraulic traits may be coordinated (Withington *et al.* 2006; Valverde-Barrantes *et al.* 2017; McCulloh *et al.* 2019; Weigelt *et al.* 2021; Verslues *et al.* 2022).

Theories for the coordinated function of leaves and roots assume that both organs are acted on by the same selective forces during evolution (Wright *et al.* 2004; Boyce 2005; Valverde-Barrantes *et al.* 2017; Weigelt *et al.* 2023). In that case, leaf and root traits might be unified along a single overall dimension of plant design (Freschet *et al.* 2010; Weigelt *et al.* 2023). The plant economic spectrum (PES) predicts that species with rapid resource uptake should achieve this with acquisitive economic traits in both leaves and roots, resulting in correlations between key traits including leaf mass per area (LMA) and specific root length (SRL) (Freschet *et al.* 2010; Reich 2014; Bergmann *et al.* 2020; Weigelt *et al.* 2021). Plant hydraulic theory also predicts that maximum flux rates would be coordinated across organs, as increasing transport through one organ alone would increase bottlenecks in other organs (Tyree & Zimmermann 2002; Noblin *et al.* 2008; Wolfe *et al.* 2023). These theories of unified selection to optimize leaf and root function would predict positive coordination of leaf and root hydraulic traits such as leaf vein density (i.e., vein length per unit area, VLA) and root vessel density ($VesDens_{root}$) (Fig. 1a), and potential "compensatory" trade-offs among functional traits within organs due to selection and/or allocation or design conflicts(Agrawal *et al.* 2010; Fox 2011; Garland *et al.* 2022), for example, VLA *vs.* leaf vein diameter ($VeinDiam_{leaf}$) (Feild & Brodribb 2013).

An alternative hypothesis is that roots and leaf traits are decoupled in response to distinct constraints and selective pressures (Weemstra *et al.* 2016; Carmona *et al.* 2021; Vleminckx *et al.* 2021). Even under the hypothesis of coordinated leaf and root function at whole-plant scale, their trait decoupling could arise if, independently of their structures (i.e., their quality), the amounts of leaves and roots (i.e., their quantity) vary widely. For example, a root system constructed of roots with relatively slow uptake rates can still achieve very high total system uptake if there are many individual roots. If so, the goal of understanding whole-plant organization may be considerably more complex, and depend, in part, on constraints on allocation to above and below ground parts (Freschet *et al.* 2015; Vleminckx *et al.* 2021; Weemstra *et al.* 2023).

Figure 1 | The coordination of root and leaf traits across scales from tissues within organs, to individual species performance to community assembly. a, At the organ scale, plant leaves are designed for efficient light capture and gas/water exchange, while plant roots and their associated mycorrhizal fungi support nutrient and water uptake. Water and nutrients enter through roots and are transported to plant leaves via the vascular system that coordinates whole-plant function. We hypothesized that the vascular hydraulic flux-related traits are coordinated between roots and leaves across species. Within organs, certain traits may show compensation trade-offs, such as vein density (VLA) and vein diameter (Veindiam_{leaf})(Feild & Brodribb 2013). b, Yet, each species will show a unique trait coordination, which would influence its performance in its community. For example, species of the family Magnoliaceae generally have thick roots with wide root vessels, and thick leaf minor veins, relative to other species, corresponding to a low abundance value in their home community. c, We hypothesized that across species, the opitmzied coordination between leaves and roots of vascular flux-related traits, and of economics-related traits, would be linked with mycorrhizal association and with their relative dominance within forest communities.

Root-leaf relations are further complicated by trade-offs and interactions between roots and symbiotic fungi (Kong *et al.* 2019; Bergmann *et al.* 2020; Weigelt *et al.* 2021). Almost all trees form symbioses with arbuscular mycorrhizal (AM) or ectomycorrhizal (EcM) fungi to acquire nutrients and water against the background of competition with numerous soil microbes (Brundrett 2002; Liu *et al.* 2015; Chen *et al.* 2016; Meng *et al.* 2023). The great variety in the identities and intensities of mycorrhizal associations raises the fundamental question of how mycorrhizal fungi affect whole plant water and nutrient acquisition and transport efficiency, and thus, species competitiveness and community assembly (Fig.1b). Notably, community ecology has historically emphasized niche differentiation according to leaf or stem traits, with much less focus on below-ground traits (Laughlin 2014; Laughlin *et al.* 2021; Da *et al.* 2023; Werden *et al.* 2023). A holistic understanding of whole-plant coordination and community organization will improve the prediction of species dynamics with different below and above -ground strategies, especially as species and ecosystems turn over under climate change.

Here, we focused on the coordination of root and leaf traits, association with the two main mycorrhizal types, and their scaled influence on the dominance of individual

species within their respective communities. We first tested the generality of the plant economic spectrum, in which, across species, the leaf and root economic traits (LMA, leaf nitrogen (LeafN) vs. SRL, root nitrogen (RootN)) would show coordinated variation for rapid metabolism and growth, on the one hand, or slower, more conservative growth. Further, we hypothesized that across species, the vascular hydraulic transport-related traits would be coordinated between roots and leaves, with some traits showing compensatory trade-offs within organs. We especially focused on traits related to vascular hydraulic transport in roots and leaves, that is, the ratio of root stele to diameter (Stele:Diam), root vessel diameter and density, and leaf vein diameter and density. Despite species differing dramatically in their root and leaf design within a similar forest – each species might have its unique coordination (Fig.1b) resulting in performance differences in its environment and community. We thus hypothesized that the optimized flux-related traits and mycorrhizal associations of individual species would scale up to an association with their relative dominance in forest community structure (Fig. 1c). Such a mechanism would assume a trait-by-competitiveness pattern(Bennett *et al.* 2016), in which a plant species optimizes coordination of different resource acquisition traits that can enhance its dominance in the community.

We measured 13 leaf and root traits for species with two contrasting mycorrhizal associations across 101 diverse woody angiosperms (59 genera within 31 families) from six subtropical and tropical forests (Table 1). This dataset is unique in that we: (i) applied consistent methods to measure traits, differing from database meta-analyses that used various sources and measurement methods; (ii) included hydraulic traits, such as root anatomy, often ignored in meta-analyses; (iii) focused on the first-order roots (the most distal absorptive fine roots), which are most comparable in function to leaves; and (iv) calculated an "importance value" for each species as a quantitative indicator for evaluating the species dominance in their natural community.

Table 1. Summary of the 13 key leaf and root traits and species importance value for woody species in six (sub)-tropical forest

Trait (abbreviation)	Functions	Units	N	Mean	SE	Max	Min	CV(%)	Blomberg's K	Pagel's λ
Root traits										
Root diameter (RD)	economic & flux-related	μm	101	338	2.0	1010	72.6	58.6	**0.42**	**0.98**
Specific root length (SRL)	economic & flux-related	m g^{-1}	101	57.7	0.4	170	4.4	68.4	**0.18**	**0.87**
Root tissue density (RTD)	economic structural	g cm^{-3}	99	0.4	0.0	0.7	0.1	36.1	**0.12**	**0.51**
Root carbon concentration (RootC)	economic composition	mg g^{-1}	96	467	0.5	590	323	9.8	0.05	<0.01
Root nitrogen concentration (RootN)	economic composition	mg g^{-1}	101	18.2	0.1	40.7	5.4	37.7	0.11	**0.77**
Root C:N ratios	economic composition		96	27.2	0.1	77.8	9.8	44.1	**0.10**	**0.71**
Root vessel diameter (VesDiam$_{root}$)	flux-related	μm	101	0.3	0.03	0.5	0.1	22.9	**0.11**	**0.58**
Root vessel density (VesDens$_{root}$)	flux-related	mm^{-2}	96	4509	32	18684	1095	67.6	0.04	**0.36**
Root stele:diameter ratio (Stele:Diam)	flux-related		101	8.7	0.0	17.5	2.7	39.7	**0.06**	**0.42**
Leaf traits										
Leaf dry mass per area (LMA)	economic structural	g m^{-2}	101	78.2	0.3	159	32.5	33.9	0.05	0.38
Leaf nitrogen concentration (LeafN)	economic composition	mg g^{-1}	101	19.5	0.1	44.6	5.6	32.5	0.04	**0.42**
Leaf vein density (VLA)	flux-related	mm mm^{-2}	82	6.1	0.02	13.0	2.3	31.5	0.04	**0.37**
Leaf minor vein diameter (VeinDiam$_{leaf}$)	flux-related	μm	96	9.3	0.03	27.6	5.0	30.3	0.03	0.09
Community structure										
Importance value (IV)		%	95	1.4	0.02	10.8	0.0	135.8		
Relative basal area (RBA)		%	68	1.8	0.06	24.8	0.0	222.5		

N, number of species analyzed. SE, standard error. CV (%), the coefficient of variance. Blomberg's *K* and Pagel's *λ* value: bold values indicate traits have significant phylogenetic signal ($p < 0.05$), i.e., show phylogenetic influence.

183
Results and Discussion
Separation of flux-related and economic trait axes in roots and leaves
Within the (sub)-tropical woody plant trait space, we identified two main independent
dimensions of variation (Fig.2a; Extended Data Table. 4,5,6). Six root and leaf
hydraulic traits (including SRL) were strongly loaded in the first axis, which
explained 40.5% of the total variation, whereas economic traits, including leaf
nitrogen (LeafN) and root nitrogen (RootN) and leaf dry mass per unit area (LMA),
were loaded in the second axis (21.6%), indicating the decoupling of flux traits and
economics traits.
193
Between-organ coordination (and within-organ compensation) of flux-related
traits in leaves and roots
Across species, the vein length per leaf area (VLA, i.e., leaf vein density) was
positively related to root stele: diam ratio (Fig. 2b; $r = 0.47$, $P < 0.001$) and specific
root length (SRL) (Fig. 2c; $r = 0.30$, $P < 0.01$), indicating coordinated vascularization
in roots and leaves. These results support the expectation that rapid leaf transpiration
is coupled with high capacity for root water delivery to maintain plant transport by
optimizing traits that would minimize resistance to water flow in these organs that
represent bottlenecks within the plant (Tyree & Sperry 1989; Steudle & Peterson
1998; Sack & Holbrook 2006; Verslues *et al.* 2022). Thus, root-leaf hydraulic flux
coordination likely resulted from maintaining a continuous water potential gradient
due to the continuity of the xylem.
206
Across species, leaf vein diameter (VeinDiam$_{leaf}$) was positively related to the root
vessel diameter (VesDiam$_{root}$) (Fig. 2d; $r = 0.39$, $P<0.001$), indicating the xylem cell
size coordination within whole-plant design. This cell size coordination also supports
the conservation of hydraulic architecture during evolution (McCulloh *et al.* 2019;
Koçillari *et al.* 2021) compared to root and leaf morphology. Indeed, we found that
size-related root and leaf hydraulic traits showed a narrow range of variation,
suggesting significant phylogenetic conservatism in conduits diameter (Table 1). Still,
small increases in conduit diameter would disproportionately increase hydraulic
conductivity, as the flow rate in capillaries is proportional to the fourth power of the
radius as described by the Hagen–Poiseuille law (Tyree & Ewers 1991). The water
entering plants is determined by two root parts—the stele and cortex (Steudle &
Peterson 1998; Robinson *et al.* 2003). Generally, the water potential in the cortex
would be close to that of soil water while stele water potential would be closer to leaf
water potential (Robinson *et al.* 2003). The coordination of leaf and root hydraulic
traits would thus contribute to a conservative gradient in water potential from the root
stele to the leaf vein due to the anatomical continuity of the xylem. Within the
hydraulic dimension, the absorptive roots serve as the water inlet and leaf stomata act
as a water outlet.
225
We found evidence for compensation trade-offs within organs that are associated with
whole-plant hydraulic coordination across organs (Fig.2b-c, inset). Thus, root vessel
diameter (VesDiam$_{root}$) was negatively related to SRL, and leaf minor vein diameter
(VeinDiam$_{leaf}$) was negatively related to vein density (VLA) (Fig.2b-c, inset;
Extended Data Fig 1). Smaller root vessel diameter and smaller leaf veins may be an
adaptation to enable the development of a greater density of conduits (i.e., VLA,
SRL), potentially at a reduced cost in space and materials (Sack *et al.* 2012). For

example, the leaf vein diameter is typically negatively related to vein density (Feild & Brodribb 2013). This organization may help maximize the conduit network transport distribution efficiency with respect to construction costs. Notably, the size of conduits and the diameter of fine roots and veins are not the only determinants of transport capacity. Indeed, i) conduit number rather than diameter is the basis of differences in transport in these fine structures (Koçillari *et al.* 2021); ii) thicker veins and roots come at the cost of their length density (i.e., VLA, SRL) (Fig. 2; Extended Data Fig. 2), which is more important for determining their transport efficiency, and iii) at the terminal ends of the collecting or distribution system, most transport is radial, not axial. Therefore, for both leaf minor veins and absorptive fine root the exchange surface area would tend to be more important than the pipe diameters inside.

The coordinated evolution of root and leaf traits was supported by our phylogenetic analyses (Boyce *et al.* 2009). Most species in Magnoliaceae and Lauraceae tended to have thicker roots and wider root vessels, distinct from species of other more recently diverged families (Table.1, Extended Data Fig. 3, 7) (Kong *et al.* 2014; Ma *et al.* 2018). As atmospheric CO_2 has decreased since the late Cretaceous, stomata density and vein density significantly increased to improve leaf carbon fixation (Raven 2002; Boyce *et al.* 2009; Comas *et al.* 2012). Given the increased water consumption, plants would also need to adapt to increase water acquisition (Boyce 2005; Boyce *et al.* 2009; Scoffoni *et al.* 2017). Thus, thinner root diameter and cortex would enable greater specific root length and root length density to optimize water acquisition (Rieger 1999; Hernandez *et al.* 2010; Masumoto *et al.* 2022). The selective force for increased CO_2 uptake would drive the evolution of the hydraulic network from leaves to roots for maximized water flow (Fig. 2), leading to hydraulic coordination to a higher degree as the whole plant functions. As a result of these evolutionary processes, the small but dense leaf vein and root conduits enhanced the adaptive capacity to utilize pulsed resource supplies and helped plants forage for water and nutrients (Comas *et al.* 2012).

Figure 2 | Root-leaf trait coordination and compensation across 101 subtropical/tropical forest species. a, Principal component analysis highlights independent axes of vascular-related traits (PC1) and economic traits (PC2) across 101 woody species in six (sub)-tropical forests. These two orthogonal axes suggest that tradeoffs operate independently along the hydraulic and economic dimensions. PC1 indicates significant correlations among root diameter (RD), Stele:Diam ratio, root vessel diameter (VesDiam_root), specific root length (SRL), vein length per unit leaf area (VLA), and minor leaf vein diameter (VeinDiam_leaf). PC2 consists of the "economics traits", showing strong correlations between root nitrogen (RootN) and leaf nitrogen concentration (LeafN), and leaf dry mass per area (LMA). We acquired importance values (relative coverage, relative abundance, relative dominance) for 95 species in the different native communities. Each species' importance value is mapped onto a size scale; bigger circles associated with higher importance value. The color gradient in **a** indicates regions of highest (green) to lowest (white) occurrence probability of species in the trait space, with contour lines indicating 0.5, 0.95, and 0.99 quantiles. **b**, Stele:Diam ratio was positively related to VLA. **c**, SRL (proxy of root conduits density) of first-order roots is positively related to VLA (proxy for leaf conduits density) at the organ level. Both measures of conduit density, SRL and VLA, are key determinants of transport capacity. As the rules of the branching systems, thicker veins and root vessels come at the cost of their length density (inset panels, compensation in VLA *vs.* VeinDiam_leaf, SRL *vs.* VesDiam_root), which is important for determining their transport efficiency. **d,** At the terminal ends of the distribution

282 system, there is a positive coordination between VeinDiam$_{leaf}$ and VesDiam$_{root}$. The density of conduits
283 (**b, c**) more than the size of conduits (**d**) is the basis of differences in transport in root or leaf, despite
284 the long-standing expectation that the size of conduits is the leading determinant of transport capacity.
285 Economic composition traits show between organ coordination (RootN *vs.* LeafN; LeafN *vs.* 1/LMA;
286 RootN *vs.* 1/LMA) (**e, f**), but economic structural traits show decoupling between organs (**g**).

Coordination and decoupling of economic traits in roots and leaves

Leaves and roots showed coordination in economic composition traits, i.e., in leaf
(LeafN) and root nitrogen concentrations (RootN); LeafN *vs.* 1/LMA; RootN *vs.*
1/LMA (Fig. 2e, f). A leaf with higher nitrogen concentration and photosynthetic rates
would match a root with a higher nitrogen and potentially higher respiration rate and
nitrogen uptake rate (Volder *et al.* 2005), according to stoichiometric conservatism in
active metabolic tissues (Sterner & Elser 2002). In addition, some nitrogen-rich
proteins act as signal material or enzymes, enhancing coordination (Reich *et al.*
2008). Moderate positive coordination in nitrogen between leaf and root seems to be a
general pattern globally (Reich *et al.* 2008; Wang *et al.* 2022), with variation
primarily associated with soil substrate and temperature (Reich & Oleksyn 2004). The
positive correlation in nitrogen suggests similar metabolism in the leaf and absorptive
roots (Extended Data Fig.1, Fig. 2e), and integration of whole-plant functioning.

Notably, root economic structural traits — specific root length (SRL) and root
diameter (RD) were strongly loaded at the first axis, while leaf mass per area (LMA)
was strongly loaded at the second axis (Fig 2a; Extended Data Table 6), indicating
that the economic structural dimension is not strongly coordinated between leaves and
roots (Kramer-Walter *et al.* 2016; Weemstra *et al.* 2016; Han *et al.* 2023). Overall,
'fast' leaves with low LMA were not necessarily associated with high SRL for
quickly acquiring soil resources (Liu *et al.* 2010; Holdaway *et al.* 2011; Fort *et al.*
2013; Poorter & Ryser 2015; Weigelt *et al.* 2021). One of the main reasons for this
decoupling is that leaves and roots can adjust biomass allocation to maintain the
functional balance (Poorter *et al.* 2012; Freschet *et al.* 2015; Freschet *et al.* 2021).

Further, structurally, the two organs have fundamentally distinct designs (Vogelmann
et al. 1996; Brodribb *et al.* 2010), with leaves generally deploying a two-layer laminar
design (Fig. 1a), where the upper mesophyll cells (palisade and spongy) layer
primarily capturing the light and radiant energy, and the lower epidermis and stomata
maximizing the CO_2 uptake and minimizing water loss (Brodribb *et al.* 2007; Sack &
Scoffoni 2013). These two layers may vary independently (Li *et al.*, 2015), leading to
the same LMA but different functional modules (Li *et al.*, 2017). By contrast, roots
follow a multi-layer cylinder design (Robinson *et al.* 2003; Guo *et al.* 2008), and the
cortex and stele tightly co-vary (Kong *et al.* 2014). Despite the general decoupling of
the leaf and root structure owing to their unique design requirement, we infer that the
physical structure of the leaf layer containing flux-related parts might coordinate with
root anatomical traits owing to their hydraulic connectivity.

Distinct resource types may also facilitate decoupling of economics modules. Roots
are designed differently from leaves to acquire physical mass resources rather than
light energy and air flux. Water and nutrients require physical connections to soil, and
all mass must cross the epidermis, cortex, and endodermis before entering the xylem
(Fig. 1a) (Steudle & Peterson 1998; Barlow 2002). Leaves have diverse shapes,
functioning as semi-closed systems protected by various tissues (Fig. 1a), and
demonstrate multi-functionality, finely tuning their behavior to manage water loss,

light capture, and carbon fixation (Westoby *et al.* 2002; Laughlin 2014; Díaz *et al.*
2016). The diverse shapes and complex functions of leaves make them difficult to
compare to the simple cylindrical shape of roots. Overall, root and leaf morphological
traits are independently adapted and compete either belowground for heterogeneous
soil resources or aboveground for homogeneous CO_2, respectively (Boyce 2005;
Freschet *et al.* 2015).

Overall, we found that flux-related traits were decoupled from economics traits.
Specifically, hydraulics traits display between-organ coordination and within-organ
compensation tradeoffs while economic composition traits indicate between-organ
coordination but economic structural traits suggest decoupling between organs (Fig.
2).

Mycorrhizal association mediates root-leaf functional coordination.
We investigated the degree that the coordination patterns between leaf and root traits
might differ with mycorrhizal type. Thus, we tested these patterns separately for the
AM- and EcM- plant species (Extended Data Table 7), which are thought to support
distinct strategies for resource acquisition belowground and nutrient cycling (Phillips
et al. 2013).

The correlation among certain hydraulic traits (Stele:Diam *vs.* VLA) and among
economic composition flux traits (RootN *vs.* LeafN) were significant across both
mycorrhizal types. However, for given trait associations, the patterns were found
between AM- and EcM species (Extended Data Fig. 2; Table 7). It is likely that, in
these cases of pairwise relationships, the correlation between traits were mediated by
mycorrhizal type (Extended Data Table 8). Thus, species with thick roots can employ
mycorrhizal hyphae to achieve high absorptive surface area equivalent to that of
species with much thinner roots (Eissenstat *et al.* 2015; Liu *et al.* 2015), essentially
achieving functional equivalence (e.g., total absorptive surface) albeit with quite
distinct root morphological traits (McCormack & Iversen 2019). This influence of
mycorrhizal association can potentially complicate the root-leaf trait coordination,
which might help explain the variations we observed (Extended Data Fig. 2).

Flux-related root-leaf traits influence species dominance within a community.
The discovery of root-leaf coordination in water flux-related traits raises the question
of whether such coordination might impact species dominance within a community
setting (Fig. 1; Fig. 3; Extended Data Fig. 4). Further, given the close relationship
between roots and their mycorrhizal partners, it will be important to also consider how
these relationships contribute to tree species dominance. The leading hydraulic flux-
related dimension (PC1, mainly driven by SRL, Stele:Diam ratio, $VesDiam_{root}$ and
VLA) was associated with tree species importance value (Fig. 3a; $r = 0.32$, $P < 0.01$)
and relative basal area (Extended Data Fig. 4a) within diverse forest communities.
This association suggests that species with enhanced hydraulic capacity may exploit
larger resource hypervolumes, contributing to their ecological dominance. The
individual traits SRL, Stele: Diam ratio, and VLA are also closely associated species
importance value (Fig. 3c-e), whereas economic traits show little association
(Extended Data Fig. 5).

Figure 3 | Flux-related root-leaf traits link to the species dominance in the communities.
Different species with multiple water-use and light-capture strategies among different mycorrhizal
types each occupy a unique position within their respective communities, ranging from low to
high dominance indicated by their importance value (IV). **a,** we found a correlation between a
species position within PC1 and their importance value ($R^2 = 0.09$, $P < 0.01$). PC1 is the linear
combination of a range of important hydraulic traits, with low values of PC1 denoting species with
low root vessel diameter, low leaf vein diameter, low root diameter and high SRL (proxy of root
conduits density), high vein leaf per leaf area (proxy of leaf conduits density), high Stele: Diam
(Fig.2a). **b,** we found no correlation between PC2 and species importance value. PC2 is a linear
combination of leaf mass per unit area (LMA), leaf nitrogen concentration (LeafN), and root
nitrogen concentration (RootN). **c, d,** conduit number (i.e., VLA, SRL), indicated that the exchange
surface area, rather than diameter, is the basis of differences in transport in these fine structures,
significantly influencing the performance of species in the communities. **e,** Stele:Diam ratio is
important for determining their transport efficiency, constrained by the phylogenetic history, and
is correlated with the importance value of species in the communities. Yellow and green represent
the mycorrhizal fungi type; the EcM species tend to competitively dominate in seasonal forests.
The x and y-axis were log scaled.

Based on a linear mixed effects model, we found that the tree mycorrhizal identity
had a great influence on the species importance value in a given community (Fig. 4),
stronger than plant functional types such as growth form (tree vs shrub), leaf habit
(evergreen vs deciduous) across 95 species (Extended Data Table 8, 9). We observed
that EcM species had a higher importance value, on average, than AM species in these
seasonal forests ($P < 0.001$, linear mixed-effects model). These EcM host species
typically have higher leaf vein density and greater specific root length, also associated
with their dominance in these (sub)-tropical forests. These findings suggest that
symbioses-root-leaf coordination scales up as an influence on the evolutionary
assembly of plant communities and plant–plant interactions.

There are at least three mechanisms for the advantages of EcM host species with thin
roots and greater leaf vein density in the community. First, EcM trees with thin roots
are typically more tolerant of water stress than AM trees in seasonal forests (Brzostek
et al. 2014). Thin roots are relatively easily coated with EcM (Comas *et al.* 2014;

Kong *et al.* 2019), increasing water foraging space with hydrophilic hyphae (Agerer 2006). Thin roots also may work as circuit breakers of the plant hydraulic system (McCulloh *et al.* 2019; Cuneo *et al.* 2021), and can quickly sense signals of soil drought (Bais *et al.* 2006), showing strong structural plasticity and flexibility to adapt to seasonal water supplies. As a result, EcM roots possibly can adjust better to pulsed water and wet-dry season shifts (Liese *et al.* 2019). Second, EcM may form a network of mycelium and some may decompose organic matter directly to obtain nutrients, enabling these associations to gain a nutritional advantage compared to their AM neighbors (Lindahl & Tunlid 2015). Third, the Hartig net of EcM protects the host roots from various pathogens and predators: this effect is especially important when species are in the early stage (i.e., seedlings) (Connell & Lowman 1989; Peay *et al.* 2010; Chen *et al.* 2019). As to their leaf design, EcM species in these forests typically have higher leaf vein density, higher hydraulic conductivity, and shorter vein spacing compared with AM species, which can make water transfer to mesophyll cells more localized, effectively supporting photosynthesis (Sack & Holbrook 2006; Brodribb *et al.* 2010; Feild & Brodribb 2013).

Figure 4 | The influence of key root and leaf traits on the species' importance value (IV). We performed linear mixed-effects models, with all 9 root-leaf traits and plant functional type (mycorrhizal associations, leaf habit and growth form) as fixed effects and species phylogeny and sampling sites as random effects. **a**, **b**, Optimal model results for predictors of species IV using the complete and imputed datasets, respectively. Root diameter (RD), root vessel diameter (VesDiam$_{root}$), and mycorrhizal type (MYT) were significantly correlated with the IV of species. We also interpreted the key factors of variation in the species IV in the community by variance partitioning, based on the complete (**c**) and imputed datasets (**d**). Root traits play a significant role in community assembly, and the relative importance of VesDiam$_{root}$, a flux-related trait, is four times greater than Leaf N concentration (LeafN). SRL: specific root length; VLA: leaf vein density.

Overall, our results identify impacts of flux-related root-leaf traits and mycorrhizal association on plant performance and forest community assembly (Fig. 3, 4, Extended Data Fig. 6). Selective pressures in these seasonal forests may favor hydraulic designs that efficiently transport water, enable flexible nutrient foraging, and minimize construction costs. We speculate that the innovation of EcM association plus higher

specific root length and leaf vein density might enhance the root resistance to
herbivores (Chen *et al.* 2019)and plants' adaptive ability to variable soil water supply,
in turn, these plant species had higher survivorship during the long history of
disturbances in these seasonal forests (Brzostek *et al.* 2014). Our trait-based evidence
points to probable explanations for why EcM species and families (e.g.,
Dipterocarpaceae, Fagaceae, Myrtaceae) have the ability to become dominant species
at (sub)-tropical latitude despite the exceptionally high diversity of AM tree species
coexisting in these communities (McGuire *et al.* 2010; Peay *et al.* 2010; Lu & Hedin
2019).

By linking whole-plant traits organization with their influence on community
dynamics, our study extends the current knowledge boundary of trait-based ecology.
Instead of a "single economic dimension" design (Reich 2014), a plant may be
constructed by different resource-capturing modules with multiple dimensions. Yet,
all these dimensions are integrated by the whole-plant level vascular system to form a
coherent above-belowground resource supply strategy. Further, our results point to the
ability to use easier-to-measure leaf hydraulic traits to infer or predict root traits that
are much harder to measure, as a foundation for fully parameterizing the hydraulic
architecture of plants, a mission that is of high importance in the modeling ecosystems
under climate change.

Methods Summary

We collected leaf and root samples from 101 diverse woody angiosperms (59 genera
within 31 families) from six (sub)-tropical forests, covering key clades of common
species in southern China (Table 1; Extended Data Table 1, Fig. 6). We focused on
mature plants and sampled at least three per species. Root samples were collected
following the procedure described in Guo *et al.* 2008. We focused on the first-order
roots, as they can be considered functionally comparable to leaves regarding resource
acquisition. Leaf samples were collected from the upper part of the tree canopy,
with > 40 mature fully expanded sun-exposed leaves collected from each individual
trees.

We measured 13 key functional traits of leaves and absorptive roots associated with
plant hydraulics, chemistry, and morphology (Table 1). The Methods provide more
details of trait measurement*s*. We constructed a plant phylogeny using rbcL and matK
sequences of each species (Extended Data Table 2). The phylogenetic tree was
constructed using maximum likelihood and Bayesian approaches, with divergence
time estimated by BEAST 1.7.1 (Drummond *et al.* 2012).

The importance value (IV) is an integrated measure of a species' density, occurrence
frequency, and basal area, offering a comprehensive assessment of its potential niche
hyperspace and competitiveness within a community (Curtis & McIntosh 1951;
Whittaker 1972). We obtained IV data for 95 of the 101 species examined, sourcing
data from two primary sources. For 82 species, IV data was obtained from survey data
collected in natural forest communities managed by the China Ecosystem Research
Network (CERN), publicly available for approval through their website
(http://www.cnern.org.cn). The remaining 13 species' IV data were derived from
published literature describing the community structure of nearby forests (Zeng *et al.*
1999; Fang 2005; Xu *et al.* 2015; Wen *et al.* 2018). The importance value for each

species across different communities was calculated as the sum of relative density,
relative frequency, and relative basal area (RBA), divided by three (Mueller-Dombois
& Ellenberg 1974). The absence of IV for six species is likely due to their low
abundance within these forest communities or the limited spatial scale of the
community census. Further, we obtained the relative basal area (RBA), another
indicator of species' dominance, for 68 out of the 101 species from these data sources.
Both IV and RBA were used to assess the influence of root-leaf traits on species' roles
in the community.
507
508 We performed trait imputation using Multivariate Imputation with Chained Equations
509 (MICE) based on ecological and phylogenetic relationships between species to obtain
510 complete trait information. The imputed datasets were compared with the complete
511 dataset to evaluate the reliability of the imputation procedure (Extend Data Fig. 3,
512 Table 4, 5, 6).
513

Statistical analyses
All statistical analyses were performed using the R software (version 4.2.0.). Mean,
minimum, maximum, and coefficient of variation (CV) were estimated for each trait
across 101 species. Blomberg's K and Pagel's λ tests were conducted to assess the
phylogenetic signal of each trait. Phylogenetically independent contrasts (PICs) were
calculated to examine the effect of phylogeny on interspecific comparisons. Principal
component analysis (PCA) was performed on transformed and standardized root-leaf
traits of the complete (95 species) and imputed (101 species) datasets. Varimax
rotation was applied to enhance interpretability. Phylogenetic PCA was also
employed to reduce evolutionary noise and reveal functional coordination between
root and leaf traits. Dissimilarities (Euclidean distance) among leaf habits, growth
forms, mycorrhizal types were calculated using the permutational multivariate
analysis of variance (PERMANOVA). The relationships between leaf and root traits
were analyzed using standardized major axis regressions for all species and separately
for AM and EcM species. A linear mixed model with restricted maximum likelihood
was fitted to examine the impact of root-leaf traits and plant functional groups on
species competitiveness. Species IV was used as the response variable, with nine key
root-leaf traits, mycorrhizal types (AM vs. EcM), growth form (Tree vs. shrub), leaf
habit (deciduous vs. evergreen) included as explanatory variables. Before fitting the
model, we streamlined the fixed effects by selecting significant explanatory variables
with a percent increase in mean squared error (%IncMSE) greater than zero using a
random forest analysis (Liaw & Wiener 2001). We then used the dredge function in
the MuMIn package (Barton 2009) for the model selection procedure.
537

Materials and Methods

Sampling approach. We collected leaf and root samples from the six tropical-subtropical forests of South China. The sites covered a latitudinal range from 18°40´ N to 24°32´ N, with mean annual temperature ranging from 10.7 to 22.4 °C, and mean annual precipitation ranging from 1539 to 2651 mm (Extended Data Table 1). We sampled 101 angiosperm woody species (73 trees and 28 shrubs) from 70 genera and 33 families, covering key clades of common species in southern China, such as magnoliids, fabids, and lamiids (Extended Data Fig. 1b). All these species are native and dominant in the sub-canopy and canopy layers.

We sampled at least three mature individuals for each species to derive the mean species trait value. We collected root samples following the procedure described in Guo *et al.*, (2008). We first removed the surface soil (0–20 cm) at the base of the sample trees to expose the main lateral roots. We selected and cut these root branches with intact terminal branch orders (> 5 g of fresh first-order roots to ensure accuracy of measurements). Subsamples of the roots from each tree were gently washed in deionized water to remove soil adhering to the roots. These samples were immediately put into plastic tubes filled with formalin-aceto-alcohol (FAA) solution (90 ml of 50% ethanol, 5 ml of 100% glacial acetic acid, and 5 ml of 37% methanol) for later anatomical measurements. The remaining samples were transported in plastic bags in a cooler to the laboratory and frozen at -20°C until subsequent morphological and chemical analyses.

We collected leaf samples from the upper part of the tree canopy by tree climbing, with > 40 mature fully expanded sun-exposed leaves collected from each individual trees. Once collected, more than five leaves per individual were immediately put into a buffered FAA fixation solution to analyze anatomical and venation traits. The remaining 15-25 leaf samples were used to measure morphological and chemical traits.

Trait measurement. We measured fourteen key functional traits of leaves and absorptive roots (first-order root) that are thought to play an important role in resource acquisition and transportation in woody plants (Fig. 1; Table 1). We especially focused on the traits of first-order roots because these roots, which are the most short-lived and metabolically active, can be considered functionally comparable to leaves as resource acquisition organs (Pregitzer *et al.* 2002; Guo *et al.* 2008; Xia *et al.* 2010; McCormack *et al.* 2015). For anatomical traits, we randomly chose 20 first-order root segments fixed in FAA solution in the field from each species. The root segments were immersed in a sequence of alcohol solutions for dehydration before being embedded in paraffin (Guo *et al.* 2008). We sliced the roots into eight-μm-thick cross-sections. The cross-sections were then stained with safranine and fast green, with the cortex staining green and the stele staining red, and photographed using a Leica DFC450 camera (Nussloch, Germany) mounted on a Leica DM 2500 microscope. To ensure the maximum quality of the slices, we initially selected well over 20 root cross-sections and randomly chose 20 from all successful segments for our anatomical trait measurements. The anatomical traits (root diameter (RD), mean vessel diameter (VesDiam$_{root}$), stele diameter, and root vessel density (VesDens$_{root}$)) were measured using ImageJ (NIH Image, Bethesda, MD, USA). The ratio of stele diameter to root diameter (Stele:Diam) was calculated to indicate the proportion of resource transportation within a root.

More than five intact root branches for each species were dissected for morphological
measurement as described in Pregitzer et al. (2002). The length of the first-order roots
was measured using an electronic digital caliper. Specific root length (SRL) was
calculated as the root length divided by its dry mass. The oven-dried (60°C, 48 h) root
subsamples were ground to fine powder using a SPEX 8000-D mixer mill (SPEX,
Edison, NJ, USA), and their N concentrations (RootN) were measured using an
elemental analyzer (Vario Microcube; Elementar, Hanau, Germany).
594
For leaf venation measurements, paradermal sections were prepared according to the
general protocols described by Carins Murphy *et al.* (2012); Brodribb *et al.* (2013),
with modifications depending on the species. The adaxial epidermis and palisade
mesophyll were carefully removed using a sharp razor blade to expose the minor
veins. Sections were then placed in bleach (50 g L^{-1} sodium hypochlorite and 13 g L^{-1}
sodium hydroxide) for several hours to several days, depending on the species, until
clear. For some species that resisted clearing in bleach alone, sections were first
placed in 5% KOH. After clearing, sections were carefully rinsed to remove bleach
and stained in 1% toluidine blue. For each section, ten fields of view were
photographed. Using ImageJ, leaf minor vein diameter (VeinDiam$_{leaf}$, μm) was
measured, and leaf vein density (VLA, mm mm^{-2}) was calculated as the total vein
length per unit leaf area. For each species, at least four leaves from different
individuals were used, and 30 fields of view (each field with an area of 1250×937.5
μm) were selected between the midrib and the margin.
609
For leaf morphological and chemical traits, leaf surface area was scanned in the field
site immediately after sampling using a portable scanner (Canon LiDE 110, Tokyo,
Japan) and then measured using ImageJ software (NIH Image, Bethesda, MD, USA).
The leaf samples were oven-dried to a constant mass at 60 °C for 48 hours. The dried
mass of leaf samples was estimated with a precision of 0.1 mg. Leaf dry mass per unit
area (LMA, g m^{-2}) was calculated as the ratio of leaf dry mass to projected leaf area.
The oven-dried leaf samples were ground to fine powder. Leaf carbon and nitrogen
concentration (LeafN) was determined using an elemental analyzer (Vario EL III,
Elementar, Hanau, Germany). We classified these traits as associated with plant
hydraulics, chemistry, and morphology at the leaf and root (Table 1).
620
The construction of plant phylogeny. Plant genomic DNA of each species was
extracted from silica gel-dried leaves, which were collected at the same time as the
roots were sampled. We determined each species' rbcL and matK sequences
(chloroplast gene fragments) (Extended Data Table 2). After model selection using
jModelTest v2.1.1 (Posada 2008), the phylogenetic tree was constructed using
maximum likelihood (ML) and Bayesian approaches. In the phylogenetic analyses,
the tree branch length was proportional to the difference in divergence time between
neighbor clades. Divergence time was estimated by BEAST1.7.1 (Drummond *et al.*
2012) with eight fossil calibration nodes (Extended Data Table 3).
630
Evaluation of species competitiveness. Importance value (IV) is a widely used
measure of a species' competitiveness within a natural community. We obtained IV
data for 95 out of 101 species from nearby forest communities. To determine the IV
for each species within its respective community, we collected community survey
data from six long-term forest plots between 2015 and 2018. These plots are part of
the Chinese Ecosystem Research Network (CERN) ecological stations (see

Acknowledgements) and are subject to comprehensive forest inventories every five
years. Our plant tissue sampling sites were located near these plots.
639
During the community survey, each forest plot was divided into multiple 10×10-
meter survey units. In each survey plots, woody plants with a diameter at breast height
(DBH) $\geq$ 1 cm were marked and identified, and their DBH was recorded. The data
includes the basal area of each species, the number of stems for each species, the
frequency of occurrence (the proportion of survey plots in which the species appears
out of the total number of survey plots). The IV of each species was calculated as
follows:

647
$$IV(x) = \frac{1}{3} \times \left(\frac{Basal\ area\ (x)}{Basal\ area\ (All)} + \frac{Numbers\ (x)}{Numbers\ (All)} + \frac{Frequency\ (x)}{Frequency\ (All)} \right)$$

648 Where *x* is a particular species on a plot, and *All* is the sum of all species on each plot.
649
650 Relative basal area (RBA) is another indicator of species' dominance. We obtained
651 RBA data for 68 out of the 101 species from the same data sources. In assessing the
652 impact of root-leaf traits on plant dominance, both IV and RBA serve as comparative
653 benchmarks, helping to identify key root and leaf traits that influence community
654 assembly.
655
Plant functional group. Plant growth form (tree vs shrub), leaf habit (deciduous vs
evergreen) and light demand (shade-tolerance vs sun preference) were provided in
community census data. The identification of mycorrhizal types in all woody plants
relies on the anatomical examination of absorptive roots. Plants characterized by a
dense mycelial sheath encasing the root surface were categorized as ectomycorrhizal
(EcM) species, while those with cortical cells penetrated by arbuscules were
designated as arbuscular mycorrhizal (AM) species.
663
Trait imputation. Our dataset encompassed nine root-leaf traits of 101 woody
species in Chinese tropical-subtropical forests, with only 2.7% missing values.
Notably, the data for all traits were nearly complete, with the sole exception being
VLA, which had 19 missing values due to insufficient material available for
measurement (Table 1). 82 species have complete trait measurements for the key nine
traits forming the 'complete dataset'. We employed Multivariate Imputation with
Chained Equations (MICE) to obtain complete trait information, leveraging ecological
and phylogenetic relationships between species. This method is chosen for its superior
accuracy and reduced bias compared to single imputation methods (Cooke *et al.*
2019). The first ten phylogenetic eigenvectors were incorporated into the matrix to be
imputed. To capture the uncertainty in the imputation process, 20 trait datasets were
imputed and then averaged ('imputed dataset'). These imputed datasets are based on
the same input trait data but differ in their estimations for the missing data. Before the
imputation process, all traits were log10-transformed. The reliability of the imputation
procedure was evaluated by comparing statistical results for complete and imputed
datasets.
680
Statistical analysis. All statistical analyses were performed using the R software
(4.2.0.). We estimated the mean value, minimum, maximum, and coefficient of
variation (CV) for each trait across all 101 species (Table 1). We employed
Blomberg's K and Pagel's λ test assuming a Brownian evolution model to test each

trait's phylogenetic signal. Phylogenetically independent contrasts (PICs) were
subsequently calculated for all the functional traits to examine the effect of phylogeny
on interspecific comparisons. Both descriptive statistics and phylogenetic analyses
were performed using the original dataset, without incorporating imputed trait values
derived from phylogenetic relationships.
690
A principal component analysis (PCA) was performed to examine the coordination of
root-leaf traits using the transformed and standardized traits of the 'complete dataset'
(82 species) and 'imputed dataset' (101 species), respectively. Two independent
functional trait dimensions were identified via PCA. Subsequently, we applied a
varimax rotation to the selected components to improve the results' interpretability.
The phylogenetic PCA was also used to reduce the evolutionary noise, revealing the
possible functional coordination between root and leaf traits under an evolutionary
context.
699
For exploring the impact of different groups of species on functional trait variation,
we grouped species according to their life habit (Deciduous vs. Evergreen), growth
form (Tree vs. Shrub), mycorrhizal types (AM vs. EcM). The dissimilarities
(Euclidean distance) among these groups of species were calculated along two
independent trait variation axes using the permutational multivariate analysis of
variance (PERMANOVA) (vegan package).
706
The relationships between leaf and root traits were analyzed by standardized major
axis regressions (Warton *et al.* 2006). We first conducted this analysis across all
species and then separately for AM and EcM species. In contrast to simple linear
regression, we used SMA regressions because they do not assume a unidirectional
effect of one parameter over the other (i.e., SMA minimizes the areas of the triangles
formed by the observations and the line).
713
We fitted a linear mixed model with restricted maximum likelihood using species IV
as the response variable to determine how root-leaf traits and associated plant
functional groups affect community structure. As explanatory variables, we included
nine key root-leaf traits, mycorrhizal types, growth form and leaf habit. To reflect our
dataset's large-scale spatial and phylogenetic structure, we treated the sampling sites
and species phylogeny as crossed random effects. Before model fitting, we simplified
the fixed effects by screening out the important explanatory variables with a percent
increase in mean squared error (%IncMSE) larger than zero using a random forest.
We then standardized the response variable and all fixed effects to allow for a direct
comparison of estimates. We fitted this simplified model with the R lme4qtl package
using the lme4qtl function and diagnosed the normality and homogeneity of residuals
(Ziyatdinov *et al.* 2018). To obtain the best-fitting models, we created a set of models
with all possible combinations of the initial variables and sorted them according to the
Akaike Information Criterion (AIC) fitted with Maximum Likelihood. We performed
a model-averaging procedure based on the AICc (ΔAICc < 2) to determine parameter
coefficients for the best final set of predictors of IV. This model selection procedure
was performed using the dredge function in the MuMIn package. We ran linear
regression between IV and root-leaf traits to test the effect of individual traits on IV.
732
733

734
Data availability
All data will be deposited at Figshare upon publication of this paper.

Acknowledgments
We thank Shenglei Fu, Zhongliang Huang, Yuhong Liu, Yiping Zhang, Handong
Wen, Xiaobao Deng and Mingxian Lin for their assistance in field sampling, Deliang
Kong for assistance in data collection. We thank the following field research stations
and government agencies for their kind support: Dinghushan and Heshan Station for
Subtropical Forest Studies of South China Botany Garden, Chinese Academy of
Sciences; Xianhu Botanic Park of Shenzhen; Ailaoshan Station for Subtropical Forest
Studies and Xishuangbanna Station for Tropical Forest Studies of Xishuangbanna
Tropical Botanical Garden, Chinese Academy of Sciences; Jianfengling Nature
Reserve Management Bureau. This study was funded by the Natural Science
Foundation of China (NSFC grant nos. 31325006 and 31530011).

Author Contributions
ZM and DG developed the conceptual approach, compiled the data, and drafted the
manuscript; LS, ZM, and ML developed the analytical approach, and all authors
contributed to revisions. Conceptualization: ZM, DG, LS, ML; Methodology: ZM,
DL, LS, ML; Investigation: CM, LL, DM; Visualization: ZM, ML; Funding
acquisition: ZM, DG; Project administration: ZM, LS; Supervision: LS, ZM, DG;
Writing – original draft: ZM; Writing – review & editing: ZM, LS, ML.

References

Agerer, R. (2006). Fungal relationships and structural identity of their ectomycorrhizae. *Mycological Progress*, 5, 67-107.

Agrawal, A., Conner, J. & Rasmann, S. (2010). Tradeoffs and negative correlations in evolutionary ecology. In: *Evolution after darwin: the first 150 years*. Sinauer Associates, pp. 243-268.

Bais, H.P., Weir, T.L., Perry, L.G., Gilroy, S. & Vivanco, J.M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annual Review of Plant Biology*, 57, 233-266.

Barlow, P.W. (2002). Cellular patterning in root meristems: its origins and significance. In: *Plant roots: the hidden half* (eds. Waisel, Y, Eshel, A & Kafkafi, U). Marcel Dekker New York, NY, USA, pp. 49-82.

Barton, K. (2009). MuMIn : multi-model inference, R package version 0.12.0. Retrieved from http://r-forge.r-project.org/projects/mumin/ 2009.

Bennett, J.A., Riibak, K., Tamme, R., Lewis, R.J. & Pärtel, M. (2016). The reciprocal relationship between competition and intraspecific trait variation. *Journal of Ecology*, 104, 1410-1420.

Bergmann, J., Ryo, M., Prati, D., Hempel, S. & Rillig, M.C. (2017). Root traits are more than analogues of leaf traits: the case for diaspore mass. *New Phytologist*, 216, 1130-1139.

Bergmann, J., Weigelt, A., van der Plas, F., Laughlin, D.C., Kuyper, T.W., Guerrero-Ramirez, N. *et al.* (2020). The fungal collaboration gradient dominates the root economics space in plants. *Science Advance*, 6, eaba3756.

Boyce, C.K. (2005). The Evolutionary History of Roots and Leaves. In: *Vascular Transport in Plants* (eds. Holbrook, NM & Zwieniecki, MA). Academic Press Burlington, pp. 479-499.

Boyce, C.K., Brodribb, T.J., Feild, T.S. & Zwieniecki, M. (2009). Angiosperm leaf vein evolution was physiologically and environmentally transformative. *Proceedings of the Royal Society B: Biological Sciences*, 276, 1771 - 1776.

Brodribb, T.J., Feild, T.S. & Jordan, G.J. (2007). Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Physiology*, 144, 1890-1898.

Brodribb, T.J., Feild, T.S. & Sack, L. (2010). Viewing leaf structure and evolution from a hydraulic perspective. *Functional Plant Biology*, 37, 488-498.

Brodribb, T.J., Jordan, G.J. & Carpenter, R.J. (2013). Unified changes in cell size permit coordinated leaf evolution. *New Phytologist*, 199, 559-570.

Brundrett, M.C. (2002). Coevolution of roots and mycorrhizas of land plants. *New Phytologist*, 154, 275-304.

Brzostek, E.R., Dragoni, D., Schmid, H.P., Rahman, A.F., Sims, D., Wayson, C.A. *et al.* (2014). Chronic water stress reduces tree growth and the carbon sink of deciduous hardwood forests. *Global Change Biology*, 20, 2531-2539.

Carins Murphy, M.R., Jordan, G.J. & Brodribb, T.J. (2012). Differential leaf expansion can enable hydraulic acclimation to sun and shade. *Plant, Cell & Environment*, 35, 1407-1418.

Carmona, C.P., Bueno, C.G., Toussaint, A., Träger, S., Díaz, S., Moora, M. *et al.* (2021). Fine-root traits in the global spectrum of plant form and function. *Nature*, 597, 683-687.

Chen, L., Swenson, N.G., Ji, N., Mi, X., Ren, H., Guo, L. *et al.* (2019). Differential soil fungus accumulation and density dependence of trees in a subtropical forest. *Science*, 366, 124-128.

Chen, W.L., Koide, R.T., Adams, T.S., DeForest, J.L., Cheng, L. & Eissenstat, D.M.

(2016). Root morphology and mycorrhizal symbioses together shape nutrient
foraging strategies of temperate trees. *Proceedings of the National Academy of
Sciences*, 113, 8741-8746.
Comas, L.H., Callahan, H.S. & Midford, P.E. (2014). Patterns in root traits of woody
species hosting arbuscular and ectomycorrhizas: implications for the evolution
of belowground strategies. *Ecology and Evolution*, 4, 2979-2990.
Comas, L.H., Mueller, K.E., Taylor, L., Midford, P.E., Callahan, H.S. & Beerlingz, D.J.
(2012). Evolutionary patterns and biogeochemical significance of angiosperm
root traits. *International Journal of Plant Sciences*, 173, 584-595.
Connell, J.H. & Lowman, M.D. (1989). Low-diversity tropical rain forests - some
possible mechanisms for their existence. *The American Naturalist*, 134, 88-119.
Cooke, R.S.C., Eigenbrod, F. & Bates, A.E. (2019). Projected losses of global mammal
and bird ecological strategies. *Nature Communications*, 10, 2279.
Cuneo, I.F., Barrios-Masias, F., Knipfer, T., Uretsky, J., Reyes, C., Lenain, P. *et al.*
(2021). Differences in grapevine rootstock sensitivity and recovery from
drought are linked to fine root cortical lacunae and root tip function. *New
Phytologist*, 229, 272-283.
Curtis, J.T. & McIntosh, R.P. (1951). An Upland Forest Continuum in the Prairie-Forest
Border Region of Wisconsin. *Ecology*, 32, 476-496.
Da, R., Fan, C., Zhang, C., Zhao, X. & von Gadow, K. (2023). Are absorptive root traits
good predictors of ecosystem functioning? A test in a natural temperate forest.
New Phytologist, 239, 75-86.
Díaz, S., Kattge, J., Cornelissen, J.H., Wright, I.J., Lavorel, S., Dray, S. *et al.* (2016).
The global spectrum of plant form and function. *Nature*, 529, 167-171.
Drummond, A.J., Suchard, M.A., Xie, D. & Rambaut, A. (2012). Bayesian
phylogenetics with BEAUti and the BEAST 1.7. *Molecular biology and
evolution*, 29, 1969-1973.
Eissenstat, D.M., Kucharski, J.M., Zadworny, M., Adams, T.S. & Koide, R.T. (2015).
Linking root traits to nutrient foraging in arbuscular mycorrhizal trees in a
temperate forest. *New Phytologist*, 208, 114-124.
Fang, Y. (2005). Species composition and diversity of evergreen broad-leaved forest of
Castanopsis carlesii and C. eyrei in Wuyishan National Nature Reserve, Fujian,
China. *Biodiversity Science*, 13, 148-155.
Feild, T.S. & Brodribb, T.J. (2013). Hydraulic tuning of vein cell microstructure in the
evolution of angiosperm venation networks. *New Phytologist*, 199, 720-726.
Fort, F., Jouany, C. & Cruz, P. (2013). Root and leaf functional trait relations in Poaceae
species: implications of differing resource-acquisition strategies. *Journal of
Plant Ecology*, 6, 211-219.
Fox, J. (2011). Why expect trade-offs in ecology and evolution? In: *Oikos blog*.
Freschet, G.T., Cornelissen, J.H.C., van Logtestijn, R.S.P. & Aerts, R. (2010). Evidence
of the 'plant economics spectrum' in a subarctic flora. *Journal of Ecology*, 98,
362-373.
Freschet, G.T., Pagès, L., Iversen, C.M., Comas, L.H., Rewald, B., Roumet, C. *et al.*
(2021). A starting guide to root ecology: strengthening ecological concepts and
standardising root classification, sampling, processing and trait measurements.
New Phytologist, 232, 973-1122.
Freschet, G.T., Swart, E.M. & Cornelissen, J.H. (2015). Integrated plant phenotypic
responses to contrasting above- and below-ground resources: key roles of
specific leaf area and root mass fraction. *New Phytologist*, 206, 1247-1260.
Garland, T., Downs, C.J. & Ives, A.R. (2022). Trade-offs (and constraints) in

organismal biology. *Physiological and biochemical zoology*, 95, 82-112.

Geng, Y., Wang, L., Jin, D., Liu, H. & He, J.S. (2014). Alpine climate alters the
relationships between leaf and root morphological traits but not chemical traits.
Oecologia, 175, 445-455.

Guo, D.L., Xia, M.X., Wei, X., Chang, W.J., Liu, Y. & Wang, Z.Q. (2008). Anatomical
traits associated with absorption and mycorrhizal colonization are linked to root
branch order in twenty-three Chinese temperate tree species. *New Phytologist*,
180, 673-683.

Han, M.G., Chen, Y., Sun, L.J., Yu, M., Li, R., Li, S.F. *et al.* (2023). Linking rhizosphere
soil microbial activity and plant resource acquisition strategy. *Journal of
Ecology*, 111, 875-888.

Hernandez, E.I., Vilagrosa, A., Pausas, J.G. & Bellot, J. (2010). Morphological traits
and water use strategies in seedlings of Mediterranean coexisting species. *Plant
Ecology*, 207, 233-244.

Holdaway, R.J., Richardson, S.J., Dickie, I.A., Peltzer, D.A. & Coomes, D.A.J.J.o.E.
(2011). Species‐ and community‐level patterns in fine root traits along a
120000‐year soil chronosequence in temperate rain forest. *Journal of Ecology*,
99, 954-963.

Koçillari, L., Olson, M.E., Suweis, S., Rocha, R.P., Lovison, A., Cardin, F. *et al.* (2021).
The Widened Pipe Model of plant hydraulic evolution. *Proceedings of the
National Academy of Sciences*, 118, e2100314118.

Kong, D.L., Ma, C.E., Zhang, Q., Li, L., Chen, X.Y., Zeng, H. *et al.* (2014). Leading
dimensions in absorptive root trait variation across 96 subtropical forest species.
New Phytologist, 203, 863-872.

Kong, D.L., Wang, J.J., Wu, H.F., Valverde-Barrantes, O.J., Wang, R.L., Zeng, H. *et al.*
(2019). Nonlinearity of root trait relationships and the root economics spectrum.
Nature Communications, 10, 2203.

Kramer-Walter, K.R., Bellingham, P.J., Millar, T.R., Smissen, R.D., Richardson, S.J. &
Laughlin, D.C. (2016). Root traits are multidimensional: specific root length is
independent from root tissue density and the plant economic spectrum. *Journal
of Ecology*, 104, 1299-1310.

Laughlin, D.C. (2014). The intrinsic dimensionality of plant traits and its relevance to
community assembly. *Journal of Ecology*, 102, 186-193.

Laughlin, D.C., Mommer, L., Sabatini, F.M., Bruelheide, H., Kuyper, T.W.,
McCormack, M.L. *et al.* (2021). Root traits explain plant species distributions
along climatic gradients yet challenge the nature of ecological trade-offs. *Nature
Ecology Evolution*, 5, 1123-1134.

Liaw, A. & Wiener, M. (2001). Classification and Regression by RandomForest. *R
News*, 23.

Liese, R., Leuschner, C. & Meier, I.C. (2019). The effect of drought and season on root
life span in temperate arbuscular mycorrhizal and ectomycorrhizal tree species.
Journal of Ecology, 107, 2226-2239.

Lindahl, B.D. & Tunlid, A. (2015). Ectomycorrhizal fungi - potential organic matter
decomposers, yet not saprotrophs. *New Phytologist*, 205, 1443-1447.

Liu, B.T., Li, H.B., Zhu, B., Koide, R.T., Eissenstat, D.M. & Guo, D.L. (2015).
Complementarity in nutrient foraging strategies of absorptive fine roots and
arbuscular mycorrhizal fungi across 14 coexisting subtropical tree species. *New
Phytologist*, 208, 125-136.

Liu, G.F., Freschet, G.T., Pan, X., Cornelissen, J.H.C., Li, Y. & Dong, M. (2010).
Coordinated variation in leaf and root traits across multiple spatial scales in

Chinese semi-arid and arid ecosystems. *New Phytologist*, 188, 543-553.

Lu, M. & Hedin, L.O. (2019). Global plant-symbiont organization and emergence of biogeochemical cycles resolved by evolution-based trait modelling. *Nature Ecology Evolution*, 3, 239-250.

Ma, Z.Q., Guo, D.L., Xu, X.L., Lu, M.Z., Bardgett, R.D., Eissenstat, D.M. *et al.* (2018). Evolutionary history resolves global organization of root functional traits. *Nature*, 555, 94-97.

Masumoto, T., Ito, T., Akatsuki, M. & Makita, N. (2022). Fine root hydraulic conductivity relates to root functional traits in four coniferous species. *Rhizosphere*, 21, 100489.

McCormack, M.L., Dickie, I.A., Eissenstat, D.M., Fahey, T.J., Fernandez, C.W., Guo, D. *et al.* (2015). Redefining fine roots improves understanding of below-ground contributions to terrestrial biosphere processes. *New Phytologist*, 207, 505-518.

McCormack, M.L. & Iversen, C.M. (2019). Physical and functional constraints on viable belowground acquisition strategies. *Frontiers in Plant Science*, 10, 1215.

McCulloh, K.A., Domec, J.-C., Johnson, D.M., Smith, D.D. & Meinzer, F.C. (2019). A dynamic yet vulnerable pipeline: Integration and coordination of hydraulic traits across whole plants. *Plant, Cell & Environment*, 42, 2789-2807.

McGuire, K.L., Zak, D.R., Edwards, I.P., Blackwood, C.B. & Upchurch, R. (2010). Slowed decomposition is biotically mediated in an ectomycorrhizal, tropical rain forest. *Oecologia*, 164, 785-795.

Meng, Y.M., Davison, J., Clarke, J.T., Zobel, M., Gerz, M., Moora, M. *et al.* (2023). Environmental modulation of plant mycorrhizal traits in the global flora. *Ecology Letters*, 26, 1862-1876.

Mueller-Dombois, D. & Ellenberg, H. (1974). *Aims and methods of vegetation ecology*. New York: John Wiley and Sons.

Noblin, X., Mahadevan, L., Coomaraswamy, I.A., Weitz, D.A., Holbrook, N.M. & Zwieniecki, M.A. (2008). Optimal vein density in artificial and real leaves. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 9140-9144.

Peay, K.G., Kennedy, P.G., Davies, S.J., Tan, S. & Bruns, T.D. (2010). Potential link between plant and fungal distributions in a dipterocarp rainforest: community and phylogenetic structure of tropical ectomycorrhizal fungi across a plant and soil ecotone. *New Phytologist*, 185, 529-542.

Phillips, R.P., Brzostek, E. & Midgley, M.G. (2013). The mycorrhizal-associated nutrient economy: a new framework for predicting carbon-nutrient couplings in temperate forests. *New Phytologist*, 199, 41-51.

Poorter, H., Niklas, K.J., Reich, P.B., Oleksyn, J., Poot, P. & Mommer, L. (2012). Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *New Phytologist*, 193, 30-50.

Poorter, H. & Ryser, P. (2015). The limits to leaf and root plasticity: what is so special about specific root length? *New phytologist*, 206, 1188-1190.

Posada, D. (2008). jModelTest: phylogenetic model averaging. *Molecular biology and evolution*, 25, 1253-1256.

Pregitzer, K.S., DeForest, J.L., Burton, A.J., Allen, M.F., Ruess, R.W. & Hendrick, R.L. (2002). Fine root architecture of nine North American trees. *Ecological Monographs*, 72, 293-309.

Raven, J.A. (2002). Selection pressures on stomatal evolution. *New Phytologist*, 153, 371-386.

Reich, P.B. (2014). The world-wide 'fast-slow' plant economics spectrum: a traits

manifesto. *Journal of Ecology*, 102, 275-301.

Reich, P.B. & Oleksyn, J. (2004). Global patterns of plant leaf N and P in relation to
temperature and latitude. *Proceedings of the National Academy of Sciences*, 101,
11001-11006.

Reich, P.B., Tjoelker, M.G., Pregitzer, K.S., Wright, I.J., Oleksyn, J. & Machado, J.L.
(2008). Scaling of respiration to nitrogen in leaves, stems and roots of higher
land plants. *Ecology Letters*, 11, 793-801.

Rieger, M.L., Paula (1999). Root system hydraulic conductivity in species with
contrasting root anatomy. *Journal of Experimental Botany*, 50, 201-209.

Robinson, D., Hodge, A. & Fitter, A. (2003). Constraints on the form and function of
root systems. In: *Root Ecology* (eds. de Kroon, H & Visser, EJW). Springer
Berlin Heidelberg Berlin, Heidelberg, pp. 1-31.

Sack, L. & Holbrook, N.M. (2006). Leaf hydraulics. *Annual Review of Plant Biology*,
57, 361-381.

Sack, L. & Scoffoni, C. (2013). Leaf venation: structure, function, development,
evolution, ecology and applications in the past, present and future. *New
Phytologist*, 198, 983-1000.

Sack, L., Scoffoni, C., McKown, A.D., Frole, K., Rawls, M., Havran, J.C. *et al.* (2012).
Developmentally based scaling of leaf venation architecture explains global
ecological patterns. *Nature Communications*, 3, 837.

Scoffoni, C., Albuquerque, C., Brodersen, C.R., Townes, S.V., John, G.P., Cochard, H.
et al. (2017). Leaf vein xylem conduit diameter influences susceptibility to
embolism and hydraulic decline. *New Phytologist*, 213, 1076-1092.

Sterner, R.W. & Elser, J.J. (2002). *Ecological stoichiometry: the biology of elements
from molecules to the biosphere*. Princeton University Press.

Steudle, E. & Peterson, C.A. (1998). How does water get through roots? *Journal of
experimental Botany*, 49, 775-788.

Tjoelker, M.G., Craine, J.M., Wedin, D., Reich, P.B. & Tilman, D. (2005). Linking leaf
and root trait syndromes among 39 grassland and savannah species. *New
Phytologist*, 167, 493-508.

Tyree, M.T. & Ewers, F.W. (1991). The hydraulic architecture of trees and other woody
plants. *New Phytologist*, 119, 345-360.

Tyree, M.T. & Sperry, J.S. (1989). Vulnerability of xylem to cavitation and embolism.
Annual Review of Plant Biology, 40, 19-36.

Tyree, M.T. & Zimmermann, M.H. (2002). *Xylem structure and the ascent of sap*.
Springer Berlin, Heidelberg.

Valverde-Barrantes, O.J., Freschet, G.T., Roumet, C. & Blackwood, C.B. (2017). A
worldview of root traits: the influence of ancestry, growth form, climate and
mycorrhizal association on the functional trait variation of fine-root tissues in
seed plants. *New Phytologist*, 215, 1562-1573.

Verslues, P.E., Bailey-Serres, J., Brodersen, C.R., Buckley, T.N., Conti, L., Christmann,
A. *et al.* (2022). Burning questions for a warming and changing world: 15
unknowns in plant abiotic stress. *The Plant Cell*, 35, 67-108.

Vleminckx, J., Fortunel, C., Valverde-Barrantes, O., Paine, C.E.T., Engel, J., Petronelli,
P. *et al.* (2021). Resolving whole-plant economics from leaf, stem and root traits
of 1467 Amazonian tree species. *Oikos*, 130, 1193-1208.

Vogelmann, T.C., Nishio, J.N. & Smith, W.K. (1996). Leaves and light capture: light
propagation and gradients of carbon fixation within leaves. *Trends in Plant
Science*, 1, 65-70.

Volder, A., Smart, D.R., Bloom, A.J. & Eissenstat, D.M. (2005). Rapid decline in nitrate

uptake and respiration with age in fine lateral roots of grape: implications for root efficiency and competitive effectiveness. *New Phytologist*, 165, 493-501.

Wang, Z.Q., Gong, H.Y., Sardans, J., Zhou, Q.P., Deng, J.M., Niklas, K.J. *et al.* (2022). Divergent nitrogen and phosphorus allocation strategies in terrestrial plant leaves and fine roots: A global meta-analysis. *Journal of Ecology*, 110, 2745-2758.

Warton, D.I., Wright, I.J., Falster, D.S. & Westoby, M. (2006). Bivariate line-fitting methods for allometry. *Biological Reviews*, 81, 259-291.

Weemstra, M., Kuyper, T.W., Sterck, F.J. & Umaña, M.N. (2023). Incorporating belowground traits: avenues towards a whole-tree perspective on performance. *Oikos*, 2023, e08827.

Weemstra, M., Mommer, L., Visser, E.J., Ruijven, J., Kuyper, T.W., Mohren, G.M. *et al.* (2016). Towards a multidimensional root trait framework: a tree root review. *New Phytologist*, 211, 1159-1169.

Weigelt, A., Mommer, L., Andraczek, K., Iversen, C.M., Bergmann, J., Bruelheide, H. *et al.* (2021). An integrated framework of plant form and function: the belowground perspective. *New Phytologist*, 232, 42-59.

Weigelt, A., Mommer, L., Andraczek, K., Iversen, C.M., Bergmann, J., Bruelheide, H. *et al.* (2023). The importance of trait selection in ecology. *Nature*, 618, E29-E30.

Wen, H.D., Lin, L.X., Yang, J., Hu, Y.H., Cao, M., Liu, Y.H. *et al.* (2018). Species composition and community structure of a 20 hm2 plot of mid-mountain moist evergreen broad-leaved forest on the Mts. Ailaoshan, Yunnan Province, China. *Chinese Journal of Plant Ecology*, 42, 419-429.

Werden, L.K., Averill, C., Crowther, T.W., Calderón-Morales, E., Toro, L., Alvarado, J.P. *et al.* (2023). Below-ground traits mediate tree survival in a tropical dry forest restoration. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 378, 20210067.

Westoby, M., Falster, D.S., Moles, A.T., Vesk, P.A. & Wright, I.J. (2002). Plant ecological strategies: some leading dimensions of variation between species. *Annual Review of Ecology and Systematics*, 33, 125-159.

Westoby, M. & Wright, I.J. (2006). Land-plant ecology on the basis of functional traits. *Trends in Ecology & Evolution*, 21, 261-268.

Whittaker, R.H. (1972). Evolution and measurement of species diversity. *Taxon*, 21, 213-251.

Withington, J.M., Reich, P.B., Oleksyn, J. & Eissenstat, D.M. (2006). Comparisons of structure and life span in roots and leaves aming temperate trees. *Ecological Monographs*, 76, 381-397.

Wolfe, B.T., Detto, M., Zhang, Y.-J., Anderson-Teixeira, K.J., Brodribb, T., Collins, A.D. *et al.* (2023). Leaves as bottlenecks: The contribution of tree leaves to hydraulic resistance within the soil−plant−atmosphere continuum. *Plant, Cell & Environment*, 46, 736-746.

Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F. *et al.* (2004). The worldwide leaf economics spectrum. 428, 821-827.

Xia, M.X., Guo, D.L. & Pregitzer, K.S. (2010). Ephemeral root modules in *Fraxinus mandshurica*. *New Phytologist*, 188, 1065-1074.

Xu, H., Li, Y., Lin, M., Wu, J., Luo, T., Zhou, Z. *et al.* (2015). Community characteristics of a 60 ha dynamics plot in the tropical montane rain forest in Jianfengling, Hainan Island. *Biodiversity Science*, 23, 192-201.

Zeng, Q., Zhou, R., Liu, Y., Wei, Q., Cao, H., Cai, X. *et al.* (1999). The community

1057 characterstics and conservation strategies of endangered species mangelietia
1058 pachyphylla. *Journal of tropical and subtropical botany*, 7, 109-119.
1059 Ziyatdinov, A., Vázquez-Santiago, M., Brunel, H., Martinez-Perez, A., Aschard, H. &
1060 Soria, J.M. (2018). lme4qtl: linear mixed models with flexible covariance
1061 structure for genetic studies of related individuals. *BMC Bioinformatics*, 19, 68.
1062
1063

Consolidated Figures and Figure Legends

Table 1. Summary of the 13 key leaf and root traits and species importance value for woody species in six (sub)-tropical forests.

Trait (abbreviation)	Functions	Units	N	Mean	SE	Max	Min	CV(%)	Blomberg's K	Pagel's λ
Root traits										
Root diameter (RD)	economic & flux-related	μm	101	338	2.0	1010	72.6	58.6	**0.42**	**0.98**
Specific root length (SRL)	economic & flux-related	m g⁻¹	101	57.7	0.4	170	4.4	68.4	**0.18**	**0.87**
Root tissue density (RTD)	economic structural	g cm⁻³	99	0.4	0.0	0.7	0.1	36.1	**0.12**	**0.51**
Root carbon concentration (RootC)	economic composition	mg g⁻¹	96	467	0.5	590	323	9.8	0.05	<0.01
Root nitrogen concentration (RootN)	economic composition	mg g⁻¹	101	18.2	0.1	40.7	5.4	37.7	0.11	**0.77**
Root C:N ratios	economic composition		96	27.2	0.1	77.8	9.8	44.1	**0.10**	**0.71**
Root vessel diameter (VesDiam$_{root}$)	flux-related	μm	101	0.3	0.03	0.5	0.1	22.9	**0.11**	**0.58**
Root vessel density (VesDens$_{root}$)	flux-related	mm⁻²	96	4509	32	18684	1095	67.6	0.04	**0.36**
Root stele:diameter ratio (Stele:Diam)	flux-related		101	8.7	0.0	17.5	2.7	39.7	**0.06**	**0.42**
Leaf traits										
Leaf dry mass per area (LMA)	economic structural	g m⁻²	101	78.2	0.3	159	32.5	33.9	0.05	0.38
Leaf nitrogen concentration (LeafN)	economic composition	mg g⁻¹	101	19.5	0.1	44.6	5.6	32.5	0.04	**0.42**
Leaf vein density (VLA)	flux-related	mm mm⁻²	82	6.1	0.02	13.0	2.3	31.5	0.04	**0.37**
Leaf minor vein diameter (VeinDiam$_{leaf}$)	flux-related	μm	96	9.3	0.03	27.6	5.0	30.3	0.03	0.09
Community structure										
Importance value (IV)		%	95	1.4	0.02	10.8	0.0	135.8		
Relative basal area (RBA)		%	68	1.8	0.06	24.8	0.0	222.5		

N, number of species analyzed. SE, standard error. CV (%), the coefficient of variance. Blomberg's *K* and Pagel's *λ* value: bold values indicate traits have significant phylogenetic signal ($p < 0.05$), i.e., show phylogenetic influence.

1072

1073
Figure 1 | The coordination of root and leaf traits across scales from tissues within organs, to
individual species performance to community assembly. a) At the organ scale, plant leaves are
designed for efficient light capture and gas/water exchange, while plant roots and their associated
mycorrhizal fungi support nutrient and water uptake. Water and nutrients enter through roots and are
transported to plant leaves via the vascular system that coordinates whole-plant function. We
hypothesized that the vascular hydraulic flux-related traits are coordinated between roots and leaves
across species. Within organs, certain traits may show compensation trade-offs, such as vein density
(VLA) and vein diameter (Veindiam$_{leaf}$)(Feild & Brodribb 2013). **b)** Yet, each species will show a
unique trait coordination, which would influence its performance in its community. For example,
species of the family Magnoliaceae generally have thick roots with wide root vessels, and thick leaf
minor veins, relative to other species, corresponding to a low abundance value in their home
community. **c)** We hypothesized that across species, the opitmzied coordination between leaves and
roots of vascular flux-related traits, and of economics-related traits, would be linked with mycorrhizal
association and with their relative dominance within forest communities.
1088
1089

Figure 2 | Root-leaf trait coordination and compensation across 101 subtropical/tropical forest species. a, Principal component analysis highlights independent axes of vascular-related traits (PC1) and economic traits (PC2) across 101 woody species in six (sub)-tropical forests. These two orthogonal axes suggest that tradeoffs operate independently along the hydraulic and economic dimensions. PC1 indicates significant correlations among root diameter (RD), Stele:Diam ratio, root vessel diameter (VesDiam$_{root}$), specific root length (SRL), vein length per unit leaf area (VLA), and minor leaf vein diameter (VeinDiam$_{leaf}$). PC2 consists of the "economics traits", showing strong correlations between root nitrogen (RootN) and leaf nitrogen concentration (LeafN), and leaf dry mass per area (LMA). We acquired importance values (relative coverage, relative abundance, relative dominance) for 95 species in the different native communities. Each species' importance value is mapped onto a size scale; bigger circles associated with higher importance value. The color gradient in **a** indicates regions of highest (green) to lowest (white) occurrence probability of species in the trait space, with contour lines indicating 0.5, 0.95, and 0.99 quantiles. **b**, Stele:Diam ratio was positively related to VLA. **c**, SRL (proxy of root conduits density) of first-order roots is positively related to VLA (proxy for leaf conduits density) at the organ level. Both measures of conduit density, SRL and VLA, are key determinants of transport capacity. As the rules of the branching systems, thicker veins and root vessels come at the cost of their length density (inset panels, compensation in VLA *vs.* VeinDiam$_{leaf}$, SRL *vs.* VesDiam$_{root}$), which is important for determining their transport efficiency. **d,** At the terminal ends of the distribution system, there is a positive coordination between VeinDiam$_{leaf}$ and VesDiam$_{root}$. The density of conduits (**b, c**) more than the size of conduits (**d**) is the basis of differences in transport in root or leaf, despite

1111   the long-standing expectation that the size of conduits is the leading determinant of transport capacity.

1112   Economic composition traits show between organ coordination (RootN *vs.* LeafN; LeafN *vs.* 1/LMA;

1113   RootN *vs.* 1/LMA) (**e**, **f**), but economic structural traits show decoupling between organs (**g**).

1114

Figure 3 | Flux-related root-leaf traits link to the species dominance in the communities.
Different species with multiple water-use and light-capture strategies among different mycorrhizal
types each occupy a unique position within their respective communities, ranging from low to
high dominance indicated by their importance value (IV). **a,** we found a correlation between a
species position within PC1 and their importance value ($R^2 = 0.09$, $P < 0.01$). PC1 is the linear
combination of a range of important hydraulic traits, with low values of PC1 denoting species with
low root vessel diameter, low leaf vein diameter, low root diameter and high SRL (proxy of root
conduits density), high vein leaf per leaf area (proxy of leaf conduits density), high Stele: Diam
(Fig.2a). **b,** We found no correlation between PC2 and species importance value. PC2 is a linear
combination of leaf mass per unit area (LMA), leaf nitrogen concentration (LeafN), and root
nitrogen concentration (RootN). **c, d,** conduit number (i.e., VLA, SRL), indicated that the exchange
surface area, rather than diameter, is the basis of differences in transport in these fine structures,
significantly influencing the performance of species in the communities. **e,** Stele:Diam ratio is
important for determining their transport efficiency, constrained by the phylogenetic history, and
is correlated with the importance value of species in the communities. Yellow and green represent
the mycorrhizal fungi type; the EcM species tend to competitively dominate in seasonal forests.
The x and y-axis were log scaled.

1134
Figure 4 | The influence of key root and leaf traits on the species' importance value (IV). We
performed linear mixed-effects models, with all root-leaf traits as fixed effects and species
phylogeny and sampling sites as random effects. **a**, **b**, Optimal model results for predictors of
species IV using the complete and imputed datasets, respectively. Root diameter (RD), root vessel
diameter (VesDiam$_{root}$), and mycorrhizal type (MYT) were significantly correlated with the IV of
species. We also interpreted the key factors of variation in the species IV in the community by
variance partitioning, based on the complete (**c**) and imputed datasets (**d**). Root traits play a
significant role in community assembly, and the relative importance of VesDiam$_{root}$, a flux-related
trait, is four times greater than Leaf N concentration (LeafN). SRL: specific root length; VLA: leaf
vein density.
1145