TITLE: MULTIPLE, REDUNDANT CARBOXYLIC ACID TRANSPORTERS SUPPORT MITOCHONDRIAL

METABOLISM IN *PLASMODIUM FALCIPARUM*

Authors: Krithika Rajaram[1,2*], Gabriel W. Rangel[3,4*], Justin T. Munro[4,5], Sethu C. Nair[1], Manuel

Llinás[3,4,5#], Sean T. Prigge[1#]

[1] Department of Molecular Microbiology and Immunology, Bloomberg School of Public Health,

Johns Hopkins University, Baltimore, MD 21205

[2] Present address: Department of Microbiology, The Ohio State University, Columbus, OH

43210

[3] Department of Biochemistry and Molecular Biology, Pennsylvania State University, University

Park, PA 16802

[4] Huck Center for Malaria Research, Pennsylvania State University, University Park, PA 16802

[5] Department of Chemistry, Pennsylvania State University, University Park, PA 16802

[*] Equal Contribution

[#] Corresponding Authors

ABSTRACT:

The mitochondrion of the deadliest human malaria parasite, *Plasmodium falciparum,* is

an essential source of cellular acetyl-CoA during the asexual blood-stage of the parasite life cycle.

Blocking mitochondrial acetyl-CoA synthesis leads to a hypoacetylated proteome and parasite

death. We previously determined that mitochondrial acetyl-CoA is primarily synthesized from

glucose-derived pyruvate by α-ketoacid dehydrogenases. Here, we asked if inhibiting the import

of glycolytic pyruvate across the mitochondrial inner membrane would affect acetyl-CoA

production and, thus, could be a potential target for antimalarial drug development. We selected

the two predicted mitochondrial pyruvate carrier proteins (*Pf*MPC1 and *Pf*MPC2) for genetic

knockout and isotopic metabolite tracing via HPLC-MS metabolomic analysis. Surprisingly, we

observed that asexual blood-stage parasites could survive the loss of either or both *Pf*MPCs with

only minor growth defects, despite a substantial reduction in the amount of glucose-derived

isotopic labelling into acetyl-CoA. Furthermore, genetic deletion of two additional mitochondrial

carboxylic acid transporters – DTC (di/tricarboxylic acid carrier) and YHM2 (a putative citrate/α-
ketoglutarate carrier protein) – only mildly affected asexual blood-stage replication, even in the
context of *Pf*MPC deficiency. Although we observed no added impact on the incorporation of
glucose carbon into acetyl-CoA in these quadruple knockout mutants, we noted a large decrease
in glutamine-derived label in tricarboxylic acid cycle metabolites, suggesting that DTC and YHM2
both import glutamine derivatives into the mitochondrion. Altogether, our results expose
redundant routes used to fuel the blood-stage malaria parasite mitochondrion with imported
carbon from two major sources – glucose and glutamine.

SIGNIFICANCE:

The mitochondrion of malaria parasites generates key molecules, such as acetyl-CoA, that
are required for numerous cellular processes. To support mitochondrial biosynthetic pathways,
the parasites must transport carbon sources into this organelle. By studying how the
mitochondrion obtains pyruvate, a molecule derived from glucose, we have uncovered
redundant carbon transport systems that ensure parasite survival in red blood cells. This
metabolic redundancy poses a challenge for drug development, as it enables the parasite to
adapt and survive by relying on alternative pathways when one is disrupted.

INTRODUCTION:

Plasmodium falciparum is a protozoan parasite that causes the deadliest form of malaria
in humans, killing over 600,000 people annually (World Health Organization, 2023). Each malaria
parasite contains a single mitochondrion that is required throughout its life cycle. Despite its
essentiality, only a few critical functions have been attributed to the mitochondrion during the
symptomatic stage of disease when parasites largely replicate asexually within human red blood
cells. The tricarboxylic acid (TCA) cycle is dispensable in the asexual blood stage (ABS) forms (Ke
et al., 2015; Rajaram et al., 2022), and the primary purpose of the mitochondrial electron
transport chain (mETC) is to furnish ubiquinone for dihydroorotate dehydrogenase (DHODH), a
mitochondrial enzyme that participates in *de novo* pyrimidine synthesis (Painter et al., 2007). As
a result, nearly all mitochondrial-specific antimalarials have been developed to target either the

mETC or DHODH. However, in sexual blood-stage parasites, other mitochondrial enzymes are

critical to development including aconitase (Ke et al., 2015) and isocitrate dehydrogenase (Yang

et al., 2022). To expand the druggable mitochondrial proteome, the organelle has been the focus

of intense research to further our understanding of its biology and metabolic output.

In line with these efforts, we recently established that the *P. falciparum* mitochondrion is

responsible for generating cellular acetyl-CoA in ABS parasites (Nair et al., 2023). This production

is largely driven by mitochondrial pyruvate dehydrogenase (mPDH), also known as the branched

chain alpha-keto acid dehydrogenase. The multisubunit mPDH complex uses glucose-derived

carbon, likely in the form of pyruvate, a three-carbon carboxylic acid product of glycolysis, as its

substrate. Although a second PDH complex exists in the apicoplast organelle in *Plasmodium*, it is

dispensable in ABS parasites (Cobbold et al., 2013; Pei et al., 2010; Swift et al., 2020a). When

mPDH is disrupted, parasites become reliant on the TCA cycle enzyme α-ketoglutarate

dehydrogenase (KDH) for survival. KDH is an α-ketoacid dehydrogenase that is closely related to

mPDH and that primarily converts α-ketoglutarate into succinyl-CoA in parasites; however, *in*

vitro, it has also been demonstrated to catalyze the conversion of pyruvate to acetyl-CoA (Chan

et al., 2013). The two enzymes constitute a synthetic lethal pair, and their inactivation results in

the hypoacetylation of histones and other proteins outside the mitochondrion (Nair et al., 2023).

A recent study has shown that 5-15% of the cellular acetate pool in mammalian cells is

derived from ROS-mediated pyruvate decarboxylation (Liu et al., 2018). While non-enzymatic

conversion of pyruvate to acetate may also occur in *Plasmodium* parasites, it is insufficient to

sustain parasites lacking functional mPDH and KDH, which instead require supplementation with

millimolar quantities of acetate for survival (Nair et al., 2023). A nuclear acetyl-CoA synthetase

(ACS) is responsible for scavenging and ligating free acetate to CoA (Nair et al., 2023; Prata et al.,

2021; Summers et al., 2021). However, given the high concentration of acetate required for this

bypass to be effective, ACS is unlikely to be a significant generator of acetyl-CoA under

physiological conditions (Nair et al., 2023). Combined with recent observations that pyruvate and

mitochondrial energy metabolism are linked to artemisinin resistance, the acetyl-CoA

biosynthetic pathway has emerged as a promising target for drug discovery (Chen et al., 2014; de

Vries et al., 2022; Mok et al., 2021; Schalkwijk et al., 2019; Zhang et al., 2022).

Pyruvate is generated from the breakdown of glucose during glycolysis and must be
imported from the cytoplasm into the mitochondrial matrix for use by mPDH/KDH (Nair et al.,
2023). In many eukaryotes, the transport of ions, metabolites and proteins across the outer
mitochondrial membrane is mediated by a highly abundant porin called the voltage-dependent
anion channel (VAC) (Hodge and Colombini, 1997; Homblé et al., 2012; Mallo et al., 2021;
Wideman et al., 2013). *P. falciparum* has a putative VAC that was recently demonstrated to be
essential for parasite survival (Anaguano et al., 2023). The protein localizes primarily to a
membrane at the parasite-host interface but also appears to overlap partially with the
mitochondrion. The inner mitochondrial membrane is far more selective, and entry is typically
mediated by specific solute transporters (Cunningham and Rutter, 2020). *P. falciparum* encodes
two proteins that are homologous to the mitochondrial pyruvate carrier proteins MPC1 and
MPC2 found in yeast, mammals, and the related apicomplexan parasite *Toxoplasma gondii*
(Bricker et al., 2012; Herzig et al., 2012; Lyu et al., 2023; Weiner and Kooij, 2016; Wunderlich,
2022). These are small proteins (12 kDa and 15 kDa, respectively) that form part of a heteromeric
complex of about 150 kDa in other organisms (Bricker et al., 2012; Herzig et al., 2012).

In this study, we probed whether the putative MPC proteins in *P. falciparum* (hereafter
referred to as *Pf*MPC) play a role in acetyl-CoA synthesis by transporting pyruvate into the
mitochondrion. We employed CRISPR-based gene deletion methods to discover that both
proteins are in fact dispensable in ABS parasites. However, there is a marked decrease in the
incorporation of ^{13}C glucose into acetyl-CoA and its derivatives in *Pf*MPC-deficient parasites,
confirming the role of the *Pf*MPCs in pyruvate import. Deletion of two additional carboxylic acid
transporters, a mitochondrial di/tricarboxylate carrier (DTC) and a putative citrate/α-
ketoglutarate transporter (YHM2), in the parasite line lacking the two *Pf*MPCs did not further
impact survival or ^{13}C glucose incorporation into known mitochondrial metabolites but did
impact ^{13}C glutamine incorporation into TCA cycle metabolites. Overall, our data indicate that
malaria parasites employ redundant mitochondrial carboxylate transporters to fuel the TCA cycle
with carbon derived from glucose and glutamine.

RESULTS:

Mitochondrial carrier proteins *Pf*MPC1 and *Pf*MPC2 are dispensable for blood-stage growth of

P. falciparum

Pyruvate is conventionally imported into mitochondria for conversion to acetyl-CoA. This

process is mediated by members of the mitochondrial pyruvate carrier (MPC) family. While three

paralogs of this family have been found in yeast, human and *Plasmodium* genomes only encode

two MPC proteins (Bricker et al., 2012; Herzig et al., 2012; Wunderlich, 2022). *Pf*MPC1 and

*Pf*MPC2 have been assigned strong localization scores by two different computational methods

(MitoProtII and *Plasmodium* Mitocarta) (Claros and Vincens, 1996; Lyu et al., 2023). To confirm

these predictions, we examined the cellular location of the putative *Pf*MPC1 (PF3D7_1340800)

and *Pf*MPC2 (PF3D7_1470400) proteins in *P. falciparum* by generating parasite lines with second

copies of *Pf*MPC1 or *Pf*MPC2 fused to a C-terminal SFG (superfolder GFP) tag (Pédelacq et al.,

2006; Roberts et al., 2019) (**Fig. S1A-D**). Unfortunately, the *Pf*MPC1-SFG signal was too faint to

be measured by live-cell imaging and was undetectable by immunofluorescence microscopy.

However, an MPC1 homolog in the closely related parasite *T. gondii* has been experimentally

established as a mitochondrial protein (Lyu et al., 2023). Live microscopy analysis of the

transgenic parasites showed that *Pf*MPC2-SFG colocalized with the MitoTracker probe,

confirming that *Pf*MPC2 is a mitochondrial protein (**Fig. 1A**).

To determine if *Pf*MPC1 and *Pf*MPC2 have essential roles in blood-stage parasites, we

used CRISPR/Cas9 to knock out the genes individually in an NF54[attB] parasite line (Adjalley et al.,

2010) (**Fig. S2**). Successful gene deletions were confirmed by diagnostic PCR (**Fig. 1B-C**), indicating

that both proteins are dispensable in ABS parasites. In yeast, MPC1 forms a heteromeric complex

with MPC2 or MPC3, and deletion of either participating subunit renders the transporter non-

functional (Bender et al., 2015; Tavoulari et al., 2019). However, it has recently been found that

homotypic complexes of human MPC2 are capable of transporting pyruvate in the absence of

MPC1 (Nagampalli et al., 2018). To determine if one MPC protein can substitute for the other in

P. falciparum, we attempted to create a double-knockout line by targeting *Pf*MPC1 for deletion

in Δ*Pf*MPC2 parasites. To provide a possible metabolic bypass, we divided the transfected

parasite culture into two flasks; one flask contained media that was supplemented with 5 mM

acetate in the event that the deletion of both *Pf*MPC proteins blocked all acetyl-CoA production

in the mitochondrion.

To our surprise, we obtained parasites lacking *Pf*MPC1 and *Pf*MPC2 (hereafter referred to

as Δ2) in media with or without acetate (**Fig. S3A-C**). We compared the growth of Δ2 parasites

against the parental line and observed only a moderate ABS fitness defect (**Fig. 1D**). These results

suggest that *Pf*MPC1 and *Pf*MPC2 either do not play a major role in supporting acetyl-CoA

synthesis in *P. falciparum*, or that a different mechanism is able to partially compensate for their

absence.

Fig. 1. *Pf*MPC proteins are dispensable in ABS parasites. (A) Representative immunofluorescence

microscopy images of *Pf*MPC2-SFG parasites show that *Pf*MPC2-SFG (green) colocalizes with MitoTracker

(red) [Mander's coefficient M1 (green in red) = 0.97, SD = 0.015, n = 19]. Images represent fields that are

10 µm long by 10 µm wide. Gene deletion of (B) *Pf*MPC1 and (C) *Pf*MPC2 in NF54[attB] (parental) parasites

was confirmed by PCR amplification (Δ5' and Δ3' products). Intact gene loci (5' and 3' products) were

detected only in the parental (blue) and not in the mutant lines (red). (D) Growth of Δ2 and parental

parasites was monitored by flow cytometry over an 8-day period. Cultures were seeded at 0.5%

parasitemia on day 0 and diluted 1:8 every other day. Two biological experiments were conducted in

quadruplicate (two-way ANOVA computed using Prism 10 [GraphPad Software, Inc], with Šídák's multiple

comparison test; *, $p \leq 0.05$; **, $p \leq 0.01$). Error bars represent the standard deviations from the mean.

***Pf*MPCs are required for optimal incorporation of glucose carbon into acetyl-CoA**

Once pyruvate enters the mitochondrion, it is acted upon by α-ketoacid dehydrogenases

to yield the two-carbon-donor compound acetyl-CoA. Acetyl-CoA then enters the TCA cycle by

combining with the four-carbon molecule oxaloacetate to form six-carbon citrate. Citrate is

converted in two steps to α-ketoglutarate, which also serves as an entry point for carbon derived

from glutamine into the TCA cycle (**Fig. 2A**). Our previously published results indicate that acetyl-

CoA or an unknown derivative is also exported out of the mitochondrion to acetylate proteins

and metabolites in the cytosol and nucleus (Nair et al., 2023). To determine if the levels of acetyl-

CoA and downstream products are affected in Δ2 compared to parental parasites, we replaced

unlabeled glucose or glutamine in the culture media with isotopically labeled $^{13}C_6$-glucose or $^{13}C_5$-

glutamine, respectively, and traced carbon incorporation into various metabolites using HPLC-

MS metabolomics (**Fig. 2A**).

We found that over 86% of hexose phosphate and over 60% of α-ketoglutarate were

derived from ^{13}C-glucose and ^{13}C-glutamine, respectively, over the course of the 2.5-hour labeling

period in both parasite lines (**Fig. 2B**, **Fig. S4A**). As expected, the fraction of labeling from ^{13}C-

glutamine in TCA cycle intermediates and N-acetyl-glutamate, an acetylated amino acid derived

from acetyl-CoA, did not vary significantly between the two lines, ruling out any intrinsic

variations in these metabolic pathways between the parental and Δ2 parasites (**Fig. 2C-E**, **Fig. S4**).

We next examined the incorporation of ^{13}C label from glucose into acetyl-CoA and N-

acetyl-glutamate (**Fig. 2C-D**). In parental parasites, both metabolites were labeled with two

carbons from $^{13}C_6$-glucose (M+2); by contrast, this labeling was highly reduced, though not

absent, in Δ2 parasites. The integrity of this pathway is also evident from the levels of two-carbon

and five-carbon (M+5; with three oxaloacetate carbons derived from anaplerotic sources) labeled

citrate species, which are only slightly reduced in Δ2 compared to parental parasites (**Fig. 2E**).

Similarly, negligible reductions are observed in the M+2 glucose label in downstream TCA cycle

metabolites (**Fig. S4**). These results suggest that *Pf*MPC1 and *Pf*MPC2 are the primary

mitochondrial pyruvate carriers, and their loss negatively impacts acetyl-CoA synthesis and

parasite growth. Despite this, *Pf*MPC deficiency is not lethal likely because some glucose-derived

carbon is still available in the mitochondrion to produce sufficient levels of acetyl-CoA.

Fig. 2. *Pf*MPC deletion reduces the synthesis of acetyl-CoA from glucose. (A) Schematic depicting the labeling pattern of mitochondrial metabolites in Δ2 parasites after short-term incubation with uniformly labeled ¹³C-glucose (solid red circles) or ¹³C-glutamine (solid blue circles). Black circles represent unlabeled atoms. Circles colored with both red and blue indicate carbons that may be derived from either ¹³C-glucose or ¹³C-glutamine. Arrows represent enzymatic reactions and are colored based on the known flow (dark green), reduced flow (light green) or predicted flow (dotted green) of carbon atoms. Abbreviations: PEP – phosphoenolpyruvate; AAT – aspartate aminotransferase; MDH – malate dehydrogenase; mPDH –

mitochondrial pyruvate dehydrogenase; KDH – α-ketoglutarate dehydrogenase; SDH – succinate

dehydrogenase; FH – fumarate dehydrogenase; MQO – malate-quinone oxidoreductase. (B-E) Fraction of

isotopically labeled metabolites when parental or Δ2 parasites are incubated with labeled glucose (Gluc)

or glutamine (Glut). Color coding denotes the mass (M) of the parent compound plus the mass shift from

the incorporation of heavy carbon isotope (^{13}C). Labeling data are expressed as the fraction of the total

metabolite pool observed in 'n' independent experiments. Error bars represent the standard deviations

from the mean.

DTC and YHM2 are dispensable individually and together with *Pf*MPCs

While *Pf*MPC1 and *Pf*MPC2 appear responsible for the vast majority of pyruvate import

into the *P. falciparum* mitochondrion, a limited amount of glucose carbon still enters the

organelle in their absence (**Fig. 2**, **Fig. S4**). To determine if other mitochondrial carboxylate

transporters can substitute for *Pf*MPCs, we searched the annotated *P. falciparum* genome using

resources on PlasmodDB.org for putative mitochondrial carboxylate carriers and found two

candidates (Alvarez-Jarreta et al., 2024; Aurrecoechea et al., 2009; Harb et al., 2024). The first, a

di/tricarboxylic acid transporter (DTC; PF3D7_0823900), is a known mitochondrial protein that

has been shown to exchange several carboxylates in reconstituted liposomes (Nozawa et al.,

2011; van Dooren et al., 2006); however, its affinity for pyruvate, a monocarboxylate, has not

been evaluated. The second candidate, YHM2 (PF3D7_1223800), a homolog of the yeast

citrate/α-ketoglutarate transporter, has not been characterized in *P. falciparum* (Castegna et al.,

2010). To determine its subcellular location, we generated transgenic parasites expressing a

second copy of SFG-tagged YHM2 (**Fig. S1A-B**, **Fig. S5**) and confirmed the colocalization of YHM2-

SFG with MitoTracker by live microscopy (**Fig. 3A**).

To determine the essentiality of DTC and YHM2 in *P. falciparum*, we attempted to delete

the genes individually in the parental NF54^{attB} line (**Fig. S2A**). Parasites lacking DTC or YHM2 were

confirmed by PCR (**Fig. 3B-C**, **Fig. S6**). Next, we used the same CRISPR plasmids to sequentially

disrupt the coding regions of DTC and YHM2 in Δ2 parasites. As before, we split the transfected

parasite culture into selective media with and without acetate. Drug-resistant parasites were

obtained in both acetate-free and supplemented media. We verified DTC and YHM2 deletions in

clonal lines using diagnostic PCR as well as by whole genome sequencing (**Fig. S7**, **Fig. S8**). Growth

comparisons of the triple (Δ*Pf*MPC1/*Pf*MPC2/DTC or Δ3) and quadruple

(Δ*Pf*MPC1/*Pf*MPC2/DTC/YHM2 or Δ4) mutants with parental parasites revealed only moderate

differences in replication rates (**Fig. 3D-E**). These results indicate that DTC and YHM2 are both

dispensable in ABS parasites and their absence does not confer large fitness costs to parasites.

Fig. 3. Multiple transporters are dispensable in ABS parasites. (A) Representative immunofluorescence

microscopy images of YHM2-SFG parasites show that YHM2-SFG (green) colocalizes with MitoTracker

(red) [Mander's coefficient M1 (green in red) = 0.97, SD = 0.026, n = 16]. Images represent fields that are

10 μm long by 10 μm wide. Gene deletion of (B) DTC and (C) YHM2 in NF54^attB (parental) parasites was

confirmed by PCR amplification (Δ5' and Δ3' products). Intact gene loci (5' and 3' products) were detected

only in the parental (blue) and not in the mutant lines (red). Growth of parental, (D) Δ3, and (E) Δ4

parasites was compared by measuring their initial and final parasitemia over the 48-h developmental cycle

within red blood cells. The average fold change in parasitemia was calculated from the ratio of these

values to provide a measure of fitness. The data are from eight biological experiments with quadruplicate

samples. Error bars represent standard deviations from the mean. Statistics: Unpaired t-test using Prism

10 [GraphPad Software, Inc]; ns – not significant. Note: The growth data for the Δ3 line was collected

alongside Δ2, so the control parental data shown in Fig. 3D is identical to that in Fig. 1D.

DTC and YHM2 transport α-ketoglutarate

Although DTC and YHM2 are dispensable in ABS parasites, their absence could still affect

carbon import into the mitochondrion. To determine if the deletion of these genes in the Δ2

background confers an additive decrease in glucose carbon incorporation into acetyl-CoA, we

incubated Δ3, Δ4 and parental parasites in growth media containing $^{13}C_6$-glucose. As expected,

we observed a significant loss of two-carbon label (M+2) in acetyl-CoA and N-acetyl-glutamate

(**Fig. S9B-C**) in Δ3 and Δ4 mutants when compared with parental parasites. However, this

reduction was similar to what we observed with the Δ2 parasites (**Fig. 2C-D**). This suggests that

DTC and YHM2 do not play a major role in importing glucose carbon into the mitochondrion. As

observed with the double mutant, some two- and five-carbon label was still being incorporated

into citrate in Δ3 and Δ4 parasites (**Fig. S9D**).

The *P. falciparum* DTC protein has been shown to mediate efficient counter-transport of

α-ketoglutarate and other carboxylates *in vitro* (Nozawa et al., 2011). While the substrate

specificity of YHM2 has not been determined in malaria parasites, its homolog in yeast is a

citrate/α-ketoglutarate antiporter (Castegna et al., 2010). To test if *P. falciparum* DTC and YHM2

import glutamine carbon as α-ketoglutarate to feed the TCA cycle, we supplemented the growth

media of Δ3, Δ4, and parental parasites with $^{13}C_5$-glutamine. N-acetyl-glutamate and α-

ketoglutarate were fully labeled with $^{13}C_5$-glutamine carbon to a similar extent in all parasite lines

(**Fig. S9C**, **Fig. 4A-B**). On the other hand, we observed a substantial reduction in $^{13}C_5$-glutamine-

labeled succinate, fumarate, and malate in Δ3 parasites, and this defect was even more

pronounced in Δ4 parasites (**Fig. 4C-E**). Taken together, these results suggest that DTC and YHM2

are both responsible for the transport of α-ketoglutarate from the cytosol into the

mitochondrion.

Interestingly, succinate was found to contain three labeled carbons from glucose (M+3)

in Δ3 and Δ4 parasites but not in the parental or Δ2 lines (**Fig. 4C**, **Fig. S4B**). This M+3 succinate is

likely derived from an anaplerotic pathway in the cytosol that produces M+3 oxaloacetate from

the glycolytic intermediate phosphoenolpyruvate (Cobbold et al., 2013; MacRae et al., 2013;

Storm et al., 2014). Oxaloacetate can be converted into malate or aspartate in the cytoplasm,

with the latter generating fumarate during purine salvage (**Fig. 4A**) (Bulusu et al., 2011; Storm et

al., 2014). Anaplerotic glucose carbon may enter the TCA cycle as malate or fumarate. We

speculate that the reduced level of α-ketoglutarate in the mitochondria of Δ3 and Δ4 parasites

leads to a reversal in the direction of TCA cycle reactions downstream of KDH, with malate

successively converted to fumarate and succinate by fumarate hydratase and succinate

dehydrogenase, respectively. The end result is an accumulation of M+3 succinate in the Δ3 and

Δ4 parasite lines (**Fig. 4A**).

Fig. 4. DTC and YHM2 control glutamine carbon incorporation into TCA cycle metabolites. (A) Schematic

depicting the labeling pattern of mitochondrial metabolites in Δ4 parasites after short-term incubation

with uniformly labeled ^{13}C-glucose (solid red circles) or ^{13}C-glutamine (solid blue circles). Black circles

represent unlabeled atoms. Circles colored with both red and blue denote carbons that may be derived

from either $^{13}C_6$-glucose or $^{13}C_5$-glutamine, respectively. Arrows represent enzymatic reactions and are

colored based on the known flow (dark green), reduced flow (light green), predicted flow (dotted green)

or reverse flow (purple) of carbon atoms. Abbreviations: PEP – phosphoenolpyruvate; AAT – aspartate

aminotransferase; MDH – malate dehydrogenase; mPDH – mitochondrial pyruvate dehydrogenase; KDH

– α-ketoglutarate dehydrogenase; SDH – succinate dehydrogenase; FH – fumarate dehydrogenase; MQO

– malate-quinone oxidoreductase. (B-E) Fraction of isotopically labeled metabolites when parental, Δ3 or

Δ4 parasites are incubated with labeled glucose (Gluc) or glutamine (Glut). Color coding denotes the mass

(M) of the parent compound plus the mass shift from the incorporation of heavy carbon isotope (^{13}C).

Labeling data are expressed as the fraction of the total metabolite pool observed in 'n' independent

experiments. Error bars represent the standard deviations from the mean.

Glucose limitation does not exacerbate growth defect of parasites lacking multiple carboxylate

transporters

Despite the observed perturbations in central carbon metabolism, we did not see a large

fitness loss in the transporter mutants (**Fig. 1D**, **Fig. 3D-E**). We wondered if this was an artifact of

culturing parasites in medium with roughly twice the glucose concentration (11 mM) compared

to that found in human blood (4.0 – 6.5 mM). We therefore compared the growth of Δ4 and

parental parasites in regular culture medium (11 mM glucose) and low glucose (3.67 mM glucose)

medium over an 8-day period, with media replenishment every other day. While glucose levels

did not alter growth kinetics during the first cycle, both parasite lines exhibited similarly reduced

multiplication rates over the next three cycles in low glucose medium (**Fig. 5**). We therefore

conclude that the high levels of glucose in regular culture medium are not masking fitness defects

in Δ4 parasites.

315
316
Fig. 5. Carboxylate transporter-deficient parasites do not exhibit heightened sensitivity to low glucose

conditions. Growth of parental and Δ4 parasites in low (3.67 mM) and high (11 mM) glucose media was

monitored by flow cytometry over an 8-day period. Cultures were seeded at 0.5% parasitemia on day 0

and diluted 1:8 every other day. Two biological experiments were conducted in quadruplicate. Error bars

represent the standard deviations from the mean. Statistics: Two-way ANOVA using Prism 10 [Graphpad

Software, Inc], with Šídák's multiple comparison test; *, $p \leq 0.05$; **, $p \leq 0.01$.

Deletion of mitochondrial transporters does not alter parasite responsiveness to

dihydroartemisinin

The mitochondrion of the malaria parasite is a key target of artemisinins. These drugs

exert their antimalarial effects in part by inducing rapid mitochondrial membrane depolarization

(Peatey et al., 2015; Wang et al., 2010). Arteminisin-resistant parasites have been found to

exhibit altered mitochondrial energy metabolism, with a broad upregulation of genes linked to

mitochondrial transport and pathways related to pyruvate, glutamate, and acetyl-CoA

metabolism (Chen et al., 2014; Mok et al., 2021; Yu et al., 2023). In particular, transcription of

the YHM2 transporter gene is upregulated in several clinically resistant isolates; however,

whether this single-handedly alters responsiveness to artemisinin has not been established

(Bonive-Boscan et al., 2023; Tripathi et al., 2023; Zhang et al., 2022; Zhu et al., 2022).

We utilized the ΔYHM2 and the dual copy parasite lines to determine the impact of

increased gene dosage of YHM2 versus no expression on sensitivity to dihydroartemisinin (DHA).

Employing standard 72-h proliferation assays, we found that the two transgenic strains and the

NF54[attB] parent had comparable IC_{50} values for DHA (Parental – 1.9 nM; YHM2-SFG – 1.8 nM;

ΔYHM2 – 2.5 nM) (**Fig. 6A-B**). Since the Δ4 transporter mutant exhibits altered pyruvate and

glutamate metabolism, we also assessed its sensitivity to DHA. We did not observe a notable shift

in the IC_{50} value (Parental – 1.7 nM; Δ4 – 1.4 nM) (**Fig. 6C**).

Artemisinin resistance is characterized by delayed parasite clearance in patients and

increased survival of early ring stage parasites *in vitro* (Amaratunga et al., 2012; Dondorp et al.,

2009; Witkowski et al., 2010). The latter can be assessed through a ring-stage survival assay (RSA)

in which 0-3 h rings are exposed to 650 nM DHA for 6 h and their recovery is measured after 66

h (Witkowski et al., 2013). We conducted parallel RSAs for the mutant and parent parasites and

found that neither YHM2 overexpression nor deletion of one or multiple mitochondrial

transporters influenced the ring-stage survival percentage (**Fig. 6D**). Therefore, our data suggest

that changes in mitochondrial transporter expression alone do not alter the susceptibility of *P.*

falciparum to DHA.

Fig. 6. Mitochondrial transporter mutants do not show altered susceptibility to dihydroartemisinin (DHA). Parasites that (A) express a second copy of YHM2 under the constitutively active Cam/HOP promoter, (B) contain a YHM2 gene deletion, or (C) contain deletions in PfMPC1, PfMPC2, DTC and YHM2 genes, were exposed to various concentrations of dihydroartemisinin (DHA) for 72 h followed by measurement of parasitemia via flow cytometry. Parental parasites were employed as control. Two independent biological experiments with quadruplicate samples were conducted. Error bars represent the standard deviation from the mean. The parental data are identical in Fig. 6A and 6B since these experiments were conducted concurrently. The calculated IC_{50} values for DHA (with 95% confidence intervals in brackets) are listed below the graphs. C) Survival percentages from a ring-stage assay for parental, YHM2-SFG, ΔYHM2, and Δ4 parasites are depicted. Two independent biological experiments

were conducted with quadruplicate samples and parasitemia was measured by flow cytometry at 66 h.

Error bars represent the standard deviations from the mean. Statistics: IC_{50} values were obtained using a

nonlinear variable slope four-parameter regression curve fitting model in Prism 10 [GraphPad Software

Inc.]. One-way ANOVA with Dunnett's correction in Prism 10 was used to compare differences between

RSA survival percentages; ns – not significant.

DISCUSSION:

370   Acetyl-CoA is a universally required metabolite that orchestrates numerous cellular

processes mainly by donating its two-carbon acetyl group to recipient molecules (Shi and Tu,

2015). Because it is membrane-impermeable, distinct pools of acetyl-CoA are typically

maintained by enzymes in various subcellular compartments. In the mitochondria of most

eukaryotes, acetyl-CoA is derived from glucose, fatty acid, and branched-chain amino acid

catabolism, and it undergoes complete oxidation through the TCA cycle to generate ATP and CO_2.

Cytosolic acetyl-CoA, which is required for lipid and isoprenoid synthesis, can be sourced in at

least two ways: from the export and cleavage of mitochondrial citrate by ATP-citrate lyase, or

from acetate by an acetyl-CoA synthetase (ACS). Acetyl-CoA can enter the nucleus through the

nuclear pore or can be additionally generated by a resident ACS to regulate gene expression via

acetylation of histones and transcription factors. The *P. falciparum* genome appears to encode

some, but not all, of the enzymes responsible for acetyl-CoA production. During asexual blood

stage development, the mitochondrion serves as the principal source of acetyl-CoA for the entire

parasite (Nair et al., 2023). The pathways for BCAA degradation and fatty acid oxidation are either

missing or incomplete, which leads us to surmise that glycolytic pyruvate is the primary substrate

for acetyl-CoA synthesis in the mitochondrion (Cobbold et al., 2013; Gardner et al., 2002; MacRae

et al., 2013; van Dooren et al., 2006). Indeed, inactivation of mPDH and KDH, the enzymes

required for the conversion of pyruvate to acetyl-CoA, is lethal for parasites (Nair et al., 2023).

Given the critical role of pyruvate in the mitochondrion, this study aimed to determine how

pyruvate is mobilized from the cytoplasm into the organelle in malaria parasites.

390   We targeted two putative mitochondrial pyruvate carrier proteins – *Pf*MPC1 and *Pf*MPC2

–for genetic and metabolic studies in *P. falciparum*. While gene deletion of one or both proteins

surprisingly did not strongly affect parasite viability in red blood cells, our isotopic labeling

analyses revealed that the ^{13}C label from glucose was considerably diminished in acetyl-CoA.

However, this reduction was not as severe as seen previously in the mPDH/KDH double-deletion

mutant, which suggested that one or more redundant mechanisms facilitate glucose carbon

entry into the mitochondrion (Nair et al., 2023). To determine if other mitochondrial carriers can

transport pyruvate, we turned our attention to two carboxylate carrier proteins, DTC and YHM2.

Single and combinatorial deletion mutants were successfully obtained with modest fitness costs.

Treatment of Δ3 (Δ*Pf*MPC1/*Pf*MPC2/DTC) and Δ4 (ΔMPC1/*Pf*MPC2/DTC/YHM2) mutants with

^{13}C$_6$-glucose demonstrated that the absence of DTC and YHM2 does not noticeably affect the

levels of pyruvate-derived acetyl-CoA. Taken together, these results establish that *Pf*MPCs are

the primary mitochondrial pyruvate carriers, but in their absence, an alternative path exists for

some cytosolic pyruvate to enter the mitochondrion.

Although additional details of acetyl-CoA synthesis in the *P. falciparum* mitochondrion

remain to be established, we were able to gain new insights into how another major carbon

source, α-ketoglutarate, is imported into the organelle. Our ^{13}C$_5$-glutamine labeling data show

that DTC plays a key role in facilitating glutamine carbon entry into the TCA cycle as α-

ketoglutarate. In *P. falciparum*, glutamine is initially converted by glutamine synthetase to

glutamate, which can then be converted to α-ketoglutarate in the cytosol by aspartate

aminotransferase or by glutamate dehydrogenases (Storm et al., 2011). A putative GABA

transaminase is speculated to convert glutamate to α-ketoglutarate in the mitochondrion, but

our results indicate that the bulk of glutamine carbon reaches the organelle as α-ketoglutarate,

not as glutamate (Cobbold and McConville, 2014). This is evidenced by the observation that DTC

deletion did not impact glutamine incorporation into α-ketoglutarate, but it affected the labeling

of other TCA cycle metabolites. Additional deletion of YHM2 exacerbated this metabolic

phenotype, consistent with both transporters importing α-ketoglutarate and conversion of

glutamine to α-ketoglutarate occurring prior to mitochondrial import.

The anaplerotic influx of glucose carbon into the TCA cycle has been attributed to DTC,

which can transport oxaloacetate, malate and to a lesser extent, fumarate *in vitro* (Ke et al., 2015;

Nozawa et al., 2011). Interestingly, DTC deletion did not block anaplerotic M+3 carbon

assimilation into TCA cycle intermediates. On the contrary, we noted a large increase in M+3

succinate species when compared with parental parasites. We attribute this buildup to a reversal

in the flow of glucose carbon from M+3 malate/fumarate in response to low levels of α-

ketoglutarate in the mitochondrion. So, if DTC is not the relevant transporter, how does

anaplerotic glucose carbon get into the mitochondrion, and in what form? A previous study

indicated that M+3 oxaloacetate does not enter the organelle; the unknown mitochondrial

carrier must therefore import anaplerotic fumarate or malate (Ke et al., 2015). If the carrier

exhibits broad specificity for carboxylates, it might also serve as the redundant pyruvate

transporter.

Although YHM2 is annotated as a citrate/α-ketoglutarate antiporter, it is difficult to

determine whether the deletion of YHM2 impacts citrate levels in the mitochondrion and the

cytoplasm, since our data reflect metabolite levels in the infected red blood cell as a whole. Even

if YHM2 exports citrate from the mitochondrion, the fate of citrate in the cytoplasm is uncertain

since malaria parasites do not appear to contain an ATP-citrate lyase to transform citrate into

acetyl-CoA (Gardner et al., 2002). The yeast YHM2 homolog supports biosynthetic reactions and

antioxidant defense by coupling citrate/α-ketoglutarate transport to NADPH generation

(Castegna et al., 2010). While it is unclear if NADPH can be formed from citrate interconversion

in the cytoplasm of *P. falciparum*, YHM2 has been recently linked to the artemisinin resistance

phenotype, with such parasites demonstrating increased tolerance to drug-induced oxidative

damage (Rocamora et al., 2018). The YHM2 gene is one of several genes associated with

mitochondrial energy metabolism that are upregulated in resistant clinical isolates (Tripathi et

al., 2023; Zhu et al., 2022). We found that deleting or expressing a second copy of YHM2 did not

alter parasite susceptibility to DHA, indicating that YHM2 is not sufficient to significantly alter

artemisinin resistance.

The metabolite data collected from our series of combinatorial deletion mutants allowed

us to propose substrate specificities for several mitochondrial carriers. Our results also

underscore the high degree of functional redundancy and plasticity in *P. falciparum* parasites, as

evidenced by their ability to survive the loss of four carrier proteins. The gene deletion results

presented in this study are not entirely concordant with a forward genetic screen for essentiality

in *P. falciparum*, in which *Pf*MPC1, *Pf*MPC2 and DTC were all assigned low mutability scores

(**Table 1**) (Zhang et al., 2018). However, the relatively small sizes of the coding sequences of these

genes (*Pf*MPC1: 318 bp, *Pf*MPC2: 387 bp, DTC: 954 bp) decrease the predictive capability of the

screen. On the other hand, YHM2 was identified as dispensable in both *P. falciparum* and *P.*

berghei screens (Schwach et al., 2015; Zhang et al., 2018). Therefore, targeted gene deletion

studies remain important to determine the functional role of genes expressed during the asexual

blood stage.

Gene ID (PF3D7_)	Gene name	Essentiality in asexual blood-stage *Plasmodium*				Mitochondrial Localization	
		P. falciparum (a)		*P. berghei* (b)	*P. falciparum* (this study)	MitoProtII (c)**	*Plasmodium* MitoCarta (d)**
		MFS*	MIS*				
1340800	Mitochondrial pyruvate carrier 1 (*Pf*MPC1)	-3.235	0.119	Not studied	Dispensable	0.8878	5.97197
1470400	Mitochondrial pyruvate carrier 2 (*Pf*MPC2)	-2.167	0.331	Not studied	Dispensable	0.6315	6.87333
0823900	Di/tricarboxylic acid carrier (DTC)	-2.799	0.162	Not studied	Dispensable	0.2626	3.54392
1223800	Citrate/α-ketoglutarate transporter (YHM2)	-0.31	1	Dispensable	Dispensable	0.1695	1.31981

Table 1. Essentiality of mitochondrial transporter genes from forward and reverse genetic approaches.

a – Zhang et al., 2018; b – Schwach et al., 2015; c – Claros and Vincens, 1996; d – Esveld et al., 2021.

* The *P. falciparum* genome-wide screen for essential genes (Zhang et al., 2018) utilized MFS (mutagenesis

fitness score) and MIS (mutagenesis index score) ranging from 0 to 1 to predict gene mutability. Low MFS

and MIS scores indicate gene essentiality.

** MitoProtII scores (ranging from 0 to 1) and *Plasmodium* MitoCarta scores (ranging from -17.2747 to

12.486) predict mitochondrial localization of *Plasmodium* proteins. Proteins with low MitoProtII scores

and high *Plasmodium MitoCarta* scores have a higher probability of localizing to the mitochondrion.

In summary, our study demonstrated that pyruvate is imported into the mitochondrion

of *P. falciparum* parasites via *Pf*MPC1 and *Pf*MPC2 and at least one additional path. We also found

that α-ketoglutarate is transported into the mitochondrion by DTC and YHM2 to feed the TCA

cycle. It remains to be determined how acetyl-CoA is exported from the mitochondrion for

utilization in other intracellular compartments. The importance of acetyl-CoA metabolism in

malaria parasites warrants further experimentation to identify and characterize novel

mitochondrial carriers that support the synthesis and utilization of this critical metabolite.

METHODS:

Parasite culture

P. falciparum NF54[attB] parasites were cultured in red blood cells (RBCs) at 2% hematocrit

in complete medium with AlbuMAX II (CMA). CMA is prepared from RPMI-1640 medium with L-

glutamine (USBiological Life Sciences, R8999) and is supplemented with 20 mM HEPES, 0.2%

sodium bicarbonate, 12.5 µg/mL hypoxanthine, 5 g/L AlbuMAX II (ThermoFisher Scientific,

11021037) and 25 µg/mL gentamicin. Parasite cultures were maintained at 37°C in a gas mixture

containing 94% N_2, 3% O_2 and 3% CO_2.

Plasmid construction

Localization constructs were derived from the previously published p15-EcDPCK-mCherry

plasmid (Swift et al., 2021). The mCherry tag was removed by digesting the plasmid with BsiWI

and AflII, and replaced with an SFG sequence that was amplified using primers listed in **Table S1**

and the p15-Mev-aSFG plasmid as template (Swift et al., 2020b). The SFG amplicon was digested

with BsiWI and AflII and ligated with the cut p15-EcDPCK plasmid using T4 DNA Ligase (NEB,

M0202). The EcDPCK gene was replaced with *Pf*MPC1, *Pf*MPC2, YHM2 or DTC coding sequences

by digesting the plasmid with AvrII and BsiWI. The coding sequences were amplified from *P.*

falciparum NF54[attB] genomic DNA (mRNA for *Pf*MPC1 and *Pf*MPC2 to exclude introns) with

primers included in **Table S1**. The amplicons were digested with BsiWI and AflII and ligated into

the digested p15-SFG plasmid.

Gene deletion constructs for *Pf*MPC1, *Pf*MPC2, DTC and YHM2 were generated using

several versions of pRSng repair plasmids with different drug resistance cassettes (Rajaram et al.,

2022; Swift et al., 2020b). Homology arms 1 (HA1) and 2 (HA2) for the genes were amplified using

primers listed in **Table S1**. The HA1 sequence was inserted into the NotI site and HA2 into the

NgoMIV site of the repair plasmid. For individual knockouts of the *Pf*MPC genes, we used the

original pRSng plasmid which contains a hDHFR (human dihydrofolate reductase) resistance

cassette that confers resistance to the drug WR99210 (Swift et al., 2020b). For deletion of

*Pf*MPC1 in the Δ*Pf*MPC2 background, hDHFR was replaced with blasticidin deaminase (BSD) to

generate a *Pf*MPC1-pRSngBD plasmid. Homology arms for DTC were cloned into a pRSng-Neo

plasmid containing a neomycin cassette, providing resistance to G418 (Rajaram et al., 2022).

Finally, homology arms for YHM2 were cloned into a pRSng-Mev plasmid which contains a
mevalonate cassette that provides resistance to fosmidomycin. To create the pRSng-Mev
plasmid, the pRSng and p15-Mev-aSFG plasmids were digested with BamHI and HindIII to replace
the hDHFR cassette in pRSng with the Mev-aSFG cassette from p15-Mev-aSFG using ligation
(Swift et al., 2020b).

To generate plasmids with guide RNA sequences that target PfMPC1, PfMPC2, DTC and
YHM2 for gene deletion, the pCasG-LacZ vector was digested with the type IIS restriction enzyme
BsaI (Rajaram et al., 2020). Guide RNAs were synthesized as oligonucleotides (**Table S1**),
annealed, and inserted into the cut plasmid. Unless specified, plasmid construction relied on
ligation-independent cloning with In-Fusion (Takara, 638944). Restriction enzymes were sourced
from New England Biolabs.

Parasite transfections

To determine the subcellular localization of _P. falciparum_ transporter proteins, 350 µL
volumes of RBCs were electroporated using the GenePulser XCell system (Bio-Rad) with: 50 µg of
the pINT plasmid (Nkrumah et al., 2006) that expresses the Bxb1 integrase, and 65 µg of the p15-
SFG plasmid expressing either PfMPC1, PfMPC2, DTC or YHM2 fused to a C-terminal SFG tag. The
electroporated RBCs were mixed with 1 mL of NF54^{attB} parasite culture at 2-3% parasitemia and
maintained in 10 mL of CMA for 2 days. They were then transferred to selective media containing
2.5 nM WR99210 for 7 days. The cultures were then returned to CMA until blood smears turned
parasite-positive, whereupon WR99210 was added back to the medium.

To generate deletion strains, 350 µL volumes of RBCs were electroporated with 65 µg
each of the repair plasmid and the appropriate pCasG-gRNA plasmid. The electroporated RBCs
were mixed with 1 mL of parasite culture at 2-3% parasitemia and maintained in 10 mL of CMA
for 2 days. They were then transferred to selective media containing 1.5 µM DSM1 and either 2.5
nM WR99210 (for individual deletions of PfMPC1 and PfMPC2), 0.5 mg/mL G418 (for DTC
knockout), or 25 µM fosmidomycin plus 50 µM mevalonolactone (for YHM2 knockout) for 7 days.
Cultures were then switched to CMA until parasites reappeared, after which they were
transferred to CMA containing WR99210 (ΔPfMPC1 and ΔPfMPC2) or G418 (ΔDTC). To delete

*Pf*MPC1 in the Δ*Pf*MPC2 line, RBCs were electroporated with the pRSngBD-*Pf*MPC1 plasmid and
the same pCasG-gRNA plasmid employed earlier. Two days post transfection, the culture was
selected with 2.5 µg/mL blasticidin for 7 days. Culture media then was replaced with CMA, and
maintained as such until parasite reappearance, at which point blasticidin was reintroduced into
the media.

All transgenic parasite lines were cloned by limiting dilution. Drugs were acquired from
the following vendors: DSM1 – Calbiochem (CAS 92872-51-0); WR99210 – Jacobus
Pharmaceutical Company, Inc.; Blasticidin – Corning (30-100-RB); G418 – Corning (61-234-RG),
Fosmidomycin – ThermoFisher Scientific (FR-31564), Mevalonolactone – Sigma-Aldrich (M4667).

Genotyping PCR

Three µL volumes of parasite culture were used in 25 µL PCR reactions with Phusion High-
Fidelity DNA polymerase (NEB, M0530). Primer pairs and anticipated sizes of PCR products are
indicated in **Fig. S1** and **Fig. S2**. Primer sequences can be found in **Table S1**.

Growth assays

The growth rates of gene knockout lines were compared against the parental line to
determine if gene deletions imparted fitness costs. Parasites were used to seed 200 µL culture
volumes/well in a 96-well plate (ThermoFisher Scientific, 267427) at a 0.5% starting parasitemia
and 2% hematocrit. The plates were incubated in chambers gassed with 94% N_2, 3% O_2, 3% CO_2
at 37°C. After 48 h (the duration of an asexual intraerythrocytic developmental cycle), final
parasitemia was determined by flow cytometry. This process was repeated over at least four
growth cycles, with an eight-fold dilution of cultures every 48 h to prevent overgrowth. Samples
were prepared for flow cytometry as described previously (Swift et al., 2020b). Briefly, parasites
were diluted 1:5 in PBS and stored in a 96-well plate at 4 °C until analysis by flow cytometry.
Parasites were stained by transferring 10 µL of the 1:5 dilutions to a 96-well plate containing 100
µL of 1x SYBR Green (Invitrogen, S7563) in PBS, and incubated for 30 minutes in the dark while
shaking. Post-incubation, 150 µL of PBS was added to each well to dilute unbound SYBR Green
dye. Stained cells were analyzed with an Attune NxT Flow Cytometer (ThermoFisher Scientific) at

a running speed of 25 μL/minute to acquire 10,000 events. Experiments were conducted with

quadruplicate samples.

P. falciparum purification, isotopic labeling, and metabolomics analysis

Heparin-synchronized NF54[attB] parental (Adjalley et al., 2011) and associated knockout

strains of *P. falciparum* parasites were cultured in standard conditions. At approximately 24-30

hours post invasion, healthy trophozoite stage parasites were purified via magnet activated cell

sorting using a SuperMACS™ II Separator (Miltenyi Biotec, 130-044-104) and a D column (Miltenyi

Biotec, 130-041-201). The samples of ~95-100% pure trophozoite-infected erythrocytes were

plated at 0.2% hematocrit in standard culture media and incubated for a one-hour recovery

period. After recovery, the parasites were pelleted, the standard culture media supernatant was

discarded, and parasites were resuspended in complete media made with either glucose-dropout

(Gibco, 11879-020) or glutamine-dropout (Gibco, 21870-076) base RPMI-1640 supplemented

with 25mM HEPES pH 7.4 (Millipore Sigma, H4034), 2 g/L sodium bicarbonate (Fisher Scientific,

BP328), 10 mg/L hypoxanthine (Fisher Scientific, AC122010250), 2.5 g/L AlbuMAX II (Thermo

Fisher Scientific, 11021029), 50 mg/L gentamicin sulfate (Bio Basic, GB0217). 11 mM $^{13}C_6$-glucose

(Cambridge Isotope Laboratories, CLM-1396-1) or 2 mM $^{13}C_5$-glutamine (Cambridge Isotope

Laboratories, CLM-1822-H) was added to respective dropout media. Parasites were cultured in

6-well culture plates at $1x10^8$ cells/5 mL media in technical triplicate for a 2.5 hour labeling

period.

After pelleting and washing in ice cold PBS pH 7.4, the infected erythrocytes were then

resuspended in 1 mL of 90% HPLC-grade methanol with 0.5 μM [$^{13}C_4,^{15}N_1$] aspartate (Millipore

Sigma, 607835) as an internal control. Supernatants were dried under nitrogen gas flow and

stored at -70°C before being resuspended to 10^6 former cells/μL in HPLC-grade water with 1μM

chlorpropamide (Alfa Aesar, J64110) as an additional internal control for reverse phase UHPLC-

MS analysis on a Thermo Exactive Plus Orbitrap. For each sample, 10 μL were injected through a

XSelect HSS T3 2.5 μM C18 column (Waters, 186006151) and eluted using a 25 min gradient of

3% aqueous methanol, 15 mM acetic acid (Millipore Sigma, A6283), 10 mM tributylamine

(Millipore Sigma, 90781), 2.5 μM medronic acid ion pairing agent (Millipore Sigma, M9508) (A)

and 100% HPLC-grade methanol (B). Negative-ion mode, using a scan range of 85 to 1,000 m/z

and a resolution of 140,000 at m/z 200 was used for ion detection.

Resulting .raw files were converted to .mzML format and centroided using the MSConvert

software of the ProteoWizard package (Chambers et al., 2012). The .mzML files were then

aligned, peaks picked, and peaks annotated (including isotopic variants) with El-MAVEN (Agrawal

et al., 2019). The chlorpropamide internal standard intensity as well as pooled quality control

(QC) samples were used to evaluate technical reproducibility of ion detection across the run. EL-

MAVEN-called peaks were then manually curated based on the proximity to the standard

retention time in the targeted library, signal/blank ratio, and peak shape. Integrated areas of

acceptable peaks were exported from EL-MAVEN in a.csv file and further processing was

performed using Microsoft Excel.

Chlorpropamide standard abundance was used to correct the peak areas per metabolite

for instrument variation in ion detection during the run. Subsequently, the average blank signals

were subtracted (or substituted to avoid negative abundances), and metabolites were filtered

based on the relative standard deviation (RSD, Standard Deviation/Mean) of QC samples run

regularly throughout data collection and the metabolites were excluded from analysis if RSD >

30% (**Supplemental File 2**). Raw data for all metabolites are available at the NIH workbench under

submission Project ID: PR002196.

Whole genome sequencing of clonal parasite lines

Parasites used for genomic DNA (gDNA) extraction were grown to 8 - 10 % schizonts in a

50-mL culture at 2% hematocrit. Cells were isolated by centrifugation (400 g for 3 minutes) before

aspirating the supernatant and adding 50 mL of 0.1% saponin in PBS. Following 10 minutes of

lysis, parasites were isolated by centrifugation (1,500 g for 10 minutes) and washed twice with

50 mL of PBS. The gDNA was extracted using a Qiagen DNeasy Blood and Tissue kit (69504)

according to the provided protocol. Extracted gDNA was quantified using a Nanodrop

spectrophotometer or a Qubit 2.0 fluorometer. Sequencing libraries for Δ2 and Δ3 lines were

prepared with 1 µg of gDNA per sample using the TruSeq® DNA PCR-Free Low-Throughput Library

Prep Kit (Illumina, 20015962) according to the manufacturer's directions. The Δ2 and Δ3 libraries

were sequenced on a HiSeq2500 Illumina platform. The sequencing library for the Δ4 line was

prepared with 100 ng of gDNA using the Celero PCR Workflow and an Enzymatic Fragmentation

DNA Seq Kit (Tecan, 30238472). This library was sequenced on the iSeq 100 Illumina platform.

Genomic sequence data for the Δ2, Δ3 and Δ4 transgenic lines are available at the NCBI Sequence

Read Archive under submission BioProject PRJNA1181605.

IC_{50} and ring-stage survival assays for DHA

To perform dose-response experiments with DHA, late-stage parasites from the parental

NF54^{attB}, ΔYHM2, YHM2-SFG and Δ4 lines were magnetically sorted from mixed-stage cultures

using MACS LS columns (Miltenyi Biotech, 130-042-401). The purified parasites were diluted to

1% parasitemia in 1.5% hematocrit and incubated with serial dilutions of DHA resuspended in

DMSO to achieve a final DMSO concentration of 0.1%. After 72 h of incubation in standard

culturing conditions, parasitemia was measured by flow cytometry as described above.

For ring-stage survival assays, cultures were synchronized by magnetic column

enrichment of late-stage parasites over two consecutive ABS cycles. After a 3 h period for

reinvasion, parasites were treated with 5% sorbitol before initiation of the assay to remove any

late-stage parasites. The highly synchronized ring-stage parasites from NF54^{attB}, ΔYHM2, YHM2-

SFG and Δ4 lines were maintained for 6 hours in CMA containing 650 nM of DHA in 0.1% DMSO.

Simultaneously, parasites were incubated in CMA with 0.1% DMSO only to serve as controls.

Following the 6 h incubation, the cultures were washed three times in CMA to remove drug or

DMSO. The washed parasites were further cultured for 66 h before measurement of parasitemia

by flow cytometry. Survival rates for each parasite line were determined by calculating the

percentage of surviving DHA-treated parasites to DMSO-treated parasites.

ACKNOWLEDGEMENTS

This work was supported by NIH R01 AI168183 (awarded to S.T.P. and M.L.), the Johns

Hopkins Malaria Research Institute, and the Bloomberg Family Foundation. K.R. was supported

by the Johns Hopkins Discovery Award, G.R. was supported by a Penn State Eberly Research

Fellowship, and J.T.M. was supported by training grant T32 DK120509. M.L. was also supported

by the Eberly College of Science and the Huck Institutes of the Life Sciences at The Pennsylvania

State University. The authors would like to acknowledge the Huck Institutes' Metabolomics Core

Facility (RRID:SCR_023864) for maintenance of the Thermo Exactive Plus.

Adjalley, S.H., Johnston, G.L., Li, T., Eastman, R.T., Ekland, E.H., Eappen, A.G., Richman, A., Sim,
B.K.L., Lee, M.C.S., Hoffman, S.L., Fidock, D.A., 2011. Quantitative assessment of
Plasmodium falciparum sexual development reveals potent transmission-blocking
activity by methylene blue. Proc Natl Acad Sci U S A 108, E1214-1223.
https://doi.org/10.1073/pnas.1112037108
Adjalley, S.H., Lee, M.C.S., Fidock, D.A., 2010. A Method for Rapid Genetic Integration into
Plasmodium falciparum Utilizing Mycobacteriophage Bxb1 Integrase, in: Braman, J.
(Ed.), In Vitro Mutagenesis Protocols, Methods in Molecular Biology. Humana Press,
Totowa, NJ, pp. 87–100. https://doi.org/10.1007/978-1-60761-652-8_6
Agrawal, S., Kumar, S., Sehgal, R., George, S., Gupta, R., Poddar, S., Jha, A., Pathak, S., 2019. El-
MAVEN: A Fast, Robust, and User-Friendly Mass Spectrometry Data Processing Engine
for Metabolomics, in: D'Alessandro, A. (Ed.), High-Throughput Metabolomics, Methods
in Molecular Biology. Springer New York, New York, NY, pp. 301–321.
https://doi.org/10.1007/978-1-4939-9236-2_19
Alvarez-Jarreta, J., Amos, B., Aurrecoechea, C., Bah, S., Barba, M., Barreto, A., Basenko, E.Y.,
Belnap, R., Blevins, A., Böhme, U., Brestelli, J., Brown, S., Callan, D., Campbell, L.I.,
Christophides, G.K., Crouch, K., Davison, H.R., DeBarry, J.D., Demko, R., Doherty, R.,
Duan, Y., Dundore, W., Dyer, S., Falke, D., Fischer, S., Gajria, B., Galdi, D., Giraldo-
Calderón, G.I., Harb, O.S., Harper, E., Helb, D., Howington, C., Hu, S., Humphrey, J.,
Iodice, J., Jones, A., Judkins, J., Kelly, S.A., Kissinger, J.C., Kittur, N., Kwon, D.K.,
Lamoureux, K., Li, W., Lodha, D., MacCallum, R.M., Maslen, G., McDowell, M.A., Myers,
J., Nural, M.V., Roos, D.S., Rund, S.S.C., Shanmugasundram, A., Sitnik, V., Spruill, D.,
Starns, D., Tomko, S.S., Wang, H., Warrenfeltz, S., Wieck, R., Wilkinson, P.A., Zheng, J.,
2024. VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource
center in 2023. Nucleic Acids Res 52, D808–D816.
https://doi.org/10.1093/nar/gkad1003
Amaratunga, C., Sreng, S., Suon, S., Phelps, E.S., Stepniewska, K., Lim, P., Zhou, C., Mao, S.,
Anderson, J.M., Lindegardh, N., Jiang, H., Song, J., Su, X., White, N.J., Dondorp, A.M.,
Anderson, T.J.C., Fay, M.P., Mu, J., Duong, S., Fairhurst, R.M., 2012. Artemisinin-resistant
Plasmodium falciparum in Pursat province, western Cambodia: a parasite clearance rate
study. Lancet Infect Dis 12, 851–858. https://doi.org/10.1016/S1473-3099(12)70181-0
Anaguano, D., Dedkhad, W., Brooks, C.F., Cobb, D.W., Muralidharan, V., 2023. Time-resolved
proximity biotinylation implicates a porin protein in export of transmembrane malaria
parasite effectors. J Cell Sci 136, jcs260506. https://doi.org/10.1242/jcs.260506
Aurrecoechea, C., Brestelli, J., Brunk, B.P., Dommer, J., Fischer, S., Gajria, B., Gao, X., Gingle, A.,
Grant, G., Harb, O.S., Heiges, M., Innamorato, F., Iodice, J., Kissinger, J.C., Kraemer, E., Li,
W., Miller, J.A., Nayak, V., Pennington, C., Pinney, D.F., Roos, D.S., Ross, C., Stoeckert,
C.J., Treatman, C., Wang, H., 2009. PlasmoDB: a functional genomic database for malaria
parasites. Nucleic Acids Res 37, D539-543. https://doi.org/10.1093/nar/gkn814
Bender, T., Pena, G., Martinou, J., 2015. Regulation of mitochondrial pyruvate uptake by
alternative pyruvate carrier complexes. The EMBO Journal 34, 911–924.
https://doi.org/10.15252/embj.201490197

Bonive-Boscan, A.D., Acosta, H., Rojas, A., 2023. Metabolic changes that allow *Plasmodium*
falciparum artemisinin-resistant parasites to tolerate the oxidative stress.
https://doi.org/10.1101/2023.10.21.560494
Bricker, D.K., Taylor, E.B., Schell, J.C., Orsak, T., Boutron, A., Chen, Y.-C., Cox, J.E., Cardon, C.M.,
Van Vranken, J.G., Dephoure, N., Redin, C., Boudina, S., Gygi, S.P., Brivet, M., Thummel,
C.S., Rutter, J., 2012. A mitochondrial pyruvate carrier required for pyruvate uptake in
yeast, Drosophila, and humans. Science 337, 96–100.
https://doi.org/10.1126/science.1218099
Bulusu, V., Jayaraman, V., Balaram, H., 2011. Metabolic fate of fumarate, a side product of the
purine salvage pathway in the intraerythrocytic stages of Plasmodium falciparum. J Biol
Chem 286, 9236–9245. https://doi.org/10.1074/jbc.M110.173328
Castegna, A., Scarcia, P., Agrimi, G., Palmieri, L., Rottensteiner, H., Spera, I., Germinario, L.,
Palmieri, F., 2010. Identification and functional characterization of a novel mitochondrial
carrier for citrate and oxoglutarate in Saccharomyces cerevisiae. J Biol Chem 285,
17359–17370. https://doi.org/10.1074/jbc.M109.097188
Chambers, M.C., Maclean, B., Burke, R., Amodei, D., Ruderman, D.L., Neumann, S., Gatto, L.,
Fischer, B., Pratt, B., Egertson, J., Hoff, K., Kessner, D., Tasman, N., Shulman, N., Frewen,
B., Baker, T.A., Brusniak, M.-Y., Paulse, C., Creasy, D., Flashner, L., Kani, K., Moulding, C.,
Seymour, S.L., Nuwaysir, L.M., Lefebvre, B., Kuhlmann, F., Roark, J., Rainer, P., Detlev, S.,
Hemenway, T., Huhmer, A., Langridge, J., Connolly, B., Chadick, T., Holly, K., Eckels, J.,
Deutsch, E.W., Moritz, R.L., Katz, J.E., Agus, D.B., MacCoss, M., Tabb, D.L., Mallick, P.,
2012. A cross-platform toolkit for mass spectrometry and proteomics. Nat Biotechnol
30, 918–920. https://doi.org/10.1038/nbt.2377
Chan, X.W.A., Wrenger, C., Stahl, K., Bergmann, B., Winterberg, M., Müller, I.B., Saliba, K.J.,
2013. Chemical and genetic validation of thiamine utilization as an antimalarial drug
target. Nat Commun 4, 2060. https://doi.org/10.1038/ncomms3060
Chen, N., LaCrue, A.N., Teuscher, F., Waters, N.C., Gatton, M.L., Kyle, D.E., Cheng, Q., 2014.
Fatty acid synthesis and pyruvate metabolism pathways remain active in
dihydroartemisinin-induced dormant ring stages of Plasmodium falciparum. Antimicrob
Agents Chemother 58, 4773–4781. https://doi.org/10.1128/AAC.02647-14
Claros, M.G., Vincens, P., 1996. Computational method to predict mitochondrially imported
proteins and their targeting sequences. Eur J Biochem 241, 779–786.
https://doi.org/10.1111/j.1432-1033.1996.00779.x
Cobbold, S.A., McConville, M.J., 2014. The Plasmodium Tricarboxylic Acid Cycle and
Mitochondrial Metabolism, in: Hommel, M., Kremsner, P.G. (Eds.), Encyclopedia of
Malaria. Springer New York, New York, NY, pp. 1–18. https://doi.org/10.1007/978-1-
4614-8757-9_13-1
Cobbold, S.A., Vaughan, A.M., Lewis, I.A., Painter, H.J., Camargo, N., Perlman, D.H.,
Fishbaugher, M., Healer, J., Cowman, A.F., Kappe, S.H., Llinas, M., 2013. Kinetic flux
profiling elucidates two independent acetyl-CoA biosynthetic pathways in Plasmodium
falciparum. The Journal of biological chemistry 288, 36338–50.
https://doi.org/10.1074/jbc.M113.503557

Cunningham, C.N., Rutter, J., 2020. 20,000 picometers under the OMM: diving into the vastness
of mitochondrial metabolite transport. EMBO Rep 21, e50071.
https://doi.org/10.15252/embr.202050071

de Vries, L.E., Jansen, P.A.M., Barcelo, C., Munro, J., Verhoef, J.M.J., Pasaje, C.F.A., Rubiano, K.,
Striepen, J., Abla, N., Berning, L., Bolscher, J.M., Demarta-Gatsi, C., Henderson, R.W.M.,
Huijs, T., Koolen, K.M.J., Tumwebaze, P.K., Yeo, T., Aguiar, A.C.C., Angulo-Barturen, I.,
Churchyard, A., Baum, J., Fernández, B.C., Fuchs, A., Gamo, F.-J., Guido, R.V.C., Jiménez-
Diaz, M.B., Pereira, D.B., Rochford, R., Roesch, C., Sanz, L.M., Trevitt, G., Witkowski, B.,
Wittlin, S., Cooper, R.A., Rosenthal, P.J., Sauerwein, R.W., Schalkwijk, J., Hermkens,
P.H.H., Bonnert, R.V., Campo, B., Fidock, D.A., Llinás, M., Niles, J.C., Kooij, T.W.A.,
Dechering, K.J., 2022. Preclinical characterization and target validation of the
antimalarial pantothenamide MMV693183. Nat Commun 13, 2158.
https://doi.org/10.1038/s41467-022-29688-5

Dondorp, A.M., Nosten, F., Yi, P., Das, D., Phyo, A.P., Tarning, J., Lwin, K.M., Ariey, F.,
Hanpithakpong, W., Lee, S.J., Ringwald, P., Silamut, K., Imwong, M., Chotivanich, K., Lim,
P., Herdman, T., An, S.S., Yeung, S., Singhasivanon, P., Day, N.P., Lindegardh, N., Socheat,
D., White, N.J., 2009. Artemisinin resistance in Plasmodium falciparum malaria. The New
England journal of medicine 361, 455–67. https://doi.org/10.1056/NEJMoa0808859

Gardner, M.J., Hall, N., Fung, E., White, O., Berriman, M., Hyman, R.W., Carlton, J.M., Pain, A.,
Nelson, K.E., Bowman, S., Paulsen, I.T., James, K., Eisen, J.A., Rutherford, K., Salzberg,
S.L., Craig, A., Kyes, S., Chan, M.-S., Nene, V., Shallom, S.J., Suh, B., Peterson, J., Angiuoli,
S., Pertea, M., Allen, J., Selengut, J., Haft, D., Mather, M.W., Vaidya, A.B., Martin, D.M.A.,
Fairlamb, A.H., Fraunholz, M.J., Roos, D.S., Ralph, S.A., McFadden, G.I., Cummings, L.M.,
Subramanian, G.M., Mungall, C., Venter, J.C., Carucci, D.J., Hoffman, S.L., Newbold, C.,
Davis, R.W., Fraser, C.M., Barrell, B., 2002. Genome sequence of the human malaria
parasite Plasmodium falciparum. Nature 419, 498–511.
https://doi.org/10.1038/nature01097

Harb, O.S., McDowell, M.A., Roos, D.S., 2024. VEuPathDB Resources: A Platform for Free Online
Data Exploration, Integration, and Analysis. Methods Mol Biol 2802, 573–586.
https://doi.org/10.1007/978-1-0716-3838-5_19

Herzig, S., Raemy, E., Montessuit, S., Veuthey, J.-L., Zamboni, N., Westermann, B., Kunji, E.R.S.,
Martinou, J.-C., 2012. Identification and Functional Expression of the Mitochondrial
Pyruvate Carrier. Science 337, 93–96. https://doi.org/10.1126/science.1218530

Hodge, T., Colombini, M., 1997. Regulation of metabolite flux through voltage-gating of VDAC
channels. J Membr Biol 157, 271–279. https://doi.org/10.1007/s002329900235

Homblé, F., Krammer, E.-M., Prévost, M., 2012. Plant VDAC: facts and speculations. Biochim
Biophys Acta 1818, 1486–1501. https://doi.org/10.1016/j.bbamem.2011.11.028

Ke, H., Lewis, I.A., Morrisey, J.M., McLean, K.J., Ganesan, S.M., Painter, H.J., Mather, M.W.,
Jacobs-Lorena, M., Llinás, M., Vaidya, A.B., 2015. Genetic investigation of tricarboxylic
acid metabolism during the Plasmodium falciparum life cycle. Cell Rep 11, 164–174.
https://doi.org/10.1016/j.celrep.2015.03.011

Liu, X., Cooper, D.E., Cluntun, A.A., Warmoes, M.O., Zhao, S., Reid, M.A., Liu, J., Lund, P.J.,
Lopes, M., Garcia, B.A., Wellen, K.E., Kirsch, D.G., Locasale, J.W., 2018. Acetate

Production from Glucose and Coupling to Mitochondrial Metabolism in Mammals. Cell
175, 502-513.e13. https://doi.org/10.1016/j.cell.2018.08.040

Lyu, C., Chen, Y., Meng, Y., Yang, J., Ye, S., Niu, Z., El-Debs, I., Gupta, N., Shen, B., 2023. The
Mitochondrial Pyruvate Carrier Coupling Glycolysis and the Tricarboxylic Acid Cycle Is
Required for the Asexual Reproduction of Toxoplasma gondii. Microbiol Spectr e05043-
22. https://doi.org/10.1128/spectrum.05043-22

MacRae, J.I., Dixon, M.W., Dearnley, M.K., Chua, H.H., Chambers, J.M., Kenny, S., Bottova, I.,
Tilley, L., McConville, M.J., 2013. Mitochondrial metabolism of sexual and asexual blood
stages of the malaria parasite Plasmodium falciparum. BMC Biol 11, 67.
https://doi.org/10.1186/1741-7007-11-67

Mallo, N., Ovciarikova, J., Martins-Duarte, E.S., Baehr, S.C., Biddau, M., Wilde, M.-L., Uboldi,
A.D., Lemgruber, L., Tonkin, C.J., Wideman, J.G., Harding, C.R., Sheiner, L., 2021.
Depletion of a Toxoplasma porin leads to defects in mitochondrial morphology and
contacts with the endoplasmic reticulum. J Cell Sci 134, jcs255299.
https://doi.org/10.1242/jcs.255299

Mok, S., Stokes, B.H., Gnädig, N.F., Ross, L.S., Yeo, T., Amaratunga, C., Allman, E., Solyakov, L.,
Bottrill, A.R., Tripathi, J., Fairhurst, R.M., Llinás, M., Bozdech, Z., Tobin, A.B., Fidock, D.A.,
2021. Artemisinin-resistant K13 mutations rewire Plasmodium falciparum's intra-
erythrocytic metabolic program to enhance survival. Nat Commun 12, 530.
https://doi.org/10.1038/s41467-020-20805-w

Nagampalli, R.S.K., Quesñay, J.E.N., Adamoski, D., Islam, Z., Birch, J., Sebinelli, H.G., Girard,
R.M.B.M., Ascenção, C.F.R., Fala, A.M., Pauletti, B.A., Consonni, S.R., de Oliveira, J.F.,
Silva, A.C.T., Franchini, K.G., Leme, A.F.P., Silber, A.M., Ciancaglini, P., Moraes, I., Dias,
S.M.G., Ambrosio, A.L.B., 2018. Human mitochondrial pyruvate carrier 2 as an
autonomous membrane transporter. Sci Rep 8, 3510. https://doi.org/10.1038/s41598-
018-21740-z

Nair, S.C., Munro, J.T., Mann, A., Llinás, M., Prigge, S.T., 2023. The mitochondrion of
Plasmodium falciparum is required for cellular acetyl-CoA metabolism and protein
acetylation. Proc Natl Acad Sci U S A 120, e2210929120.
https://doi.org/10.1073/pnas.2210929120

Nkrumah, L.J., Muhle, R.A., Moura, P.A., Ghosh, P., Hatfull, G.F., Jacobs, W.R., Fidock, D.A.,
2006. Efficient site-specific integration in Plasmodium falciparum chromosomes
mediated by mycobacteriophage Bxb1 integrase. Nat Methods 3, 615–621.
https://doi.org/10.1038/nmeth904

Nozawa, A., Fujimoto, R., Matsuoka, H., Tsuboi, T., Tozawa, Y., 2011. Cell-free synthesis,
reconstitution, and characterization of a mitochondrial dicarboxylate-tricarboxylate
carrier of Plasmodium falciparum. Biochem Biophys Res Commun 414, 612–617.
https://doi.org/10.1016/j.bbrc.2011.09.130

Painter, H.J., Morrisey, J.M., Mather, M.W., Vaidya, A.B., 2007. Specific role of mitochondrial
electron transport in blood-stage Plasmodium falciparum. Nature 446, 88–91.
https://doi.org/10.1038/nature05572

Peatey, C.L., Chavchich, M., Chen, N., Gresty, K.J., Gray, K.-A., Gatton, M.L., Waters, N.C., Cheng,
Q., 2015. Mitochondrial Membrane Potential in a Small Subset of Artemisinin-Induced

823   Dormant Plasmodium falciparum Parasites In Vitro. J Infect Dis 212, 426–434.
824   https://doi.org/10.1093/infdis/jiv048
825 Pédelacq, J.-D., Cabantous, S., Tran, T., Terwilliger, T.C., Waldo, G.S., 2006. Engineering and
826   characterization of a superfolder green fluorescent protein. Nat Biotechnol 24, 79–88.
827   https://doi.org/10.1038/nbt1172
828 Pei, Y., Tarun, A.S., Vaughan, A.M., Herman, R.W., Soliman, J.M.B., Erickson-Wayman, A., Kappe,
829   S.H.I., 2010. Plasmodium pyruvate dehydrogenase activity is only essential for the
830   parasite's progression from liver infection to blood infection. Mol Microbiol 75, 957–
831   971. https://doi.org/10.1111/j.1365-2958.2009.07034.x
832 Prata, I.O., Cubillos, E.F.G., Krüger, A., Barbosa, D., Martins, J., Setubal, J.C., Wunderlich, G.,
833   2021. Plasmodium falciparum Acetyl-CoA Synthetase Is Essential for Parasite
834   Intraerythrocytic Development and Chromatin Modification. ACS Infect Dis 7, 3224–
835   3240. https://doi.org/10.1021/acsinfecdis.1c00414
836 Rajaram, K., Liu, H.B., Prigge, S.T., 2020. Redesigned TetR-Aptamer System To Control Gene
837   Expression in Plasmodium falciparum. mSphere 5.
838   https://doi.org/10.1128/mSphere.00457-20
839 Rajaram, K., Tewari, S.G., Wallqvist, A., Prigge, S.T., 2022. Metabolic changes accompanying the
840   loss of fumarate hydratase and malate–quinone oxidoreductase in the asexual blood
841   stage of Plasmodium falciparum. Journal of Biological Chemistry 298, 101897.
842   https://doi.org/10.1016/j.jbc.2022.101897
843 Roberts, A.D., Nair, S.C., Guerra, A.J., Prigge, S.T., 2019. Development of a conditional
844   localization approach to control apicoplast protein trafficking in malaria parasites.
845   Traffic 20, 571–582. https://doi.org/10.1111/tra.12656
846 Rocamora, F., Zhu, L., Liong, K.Y., Dondorp, A., Miotto, O., Mok, S., Bozdech, Z., 2018. Oxidative
847   stress and protein damage responses mediate artemisinin resistance in malaria
848   parasites. PLoS Pathog 14, e1006930. https://doi.org/10.1371/journal.ppat.1006930
849 Schalkwijk, J., Allman, E.L., Jansen, P.A.M., de Vries, L.E., Verhoef, J.M.J., Jackowski, S., Botman,
850   P.N.M., Beuckens-Schortinghuis, C.A., Koolen, K.M.J., Bolscher, J.M., Vos, M.W., Miller,
851   K., Reeves, S.A., Pett, H., Trevitt, G., Wittlin, S., Scheurer, C., Sax, S., Fischli, C., Angulo-
852   Barturen, I., Jiménez-Diaz, M.B., Josling, G., Kooij, T.W.A., Bonnert, R., Campo, B.,
853   Blaauw, R.H., Rutjes, F.P.J.T., Sauerwein, R.W., Llinás, M., Hermkens, P.H.H., Dechering,
854   K.J., 2019. Antimalarial pantothenamide metabolites target acetyl–coenzyme A
855   biosynthesis in *Plasmodium falciparum*. Sci. Transl. Med. 11, eaas9917.
856   https://doi.org/10.1126/scitranslmed.aas9917
857 Schwach, F., Bushell, E., Gomes, A.R., Anar, B., Girling, G., Herd, C., Rayner, J.C., Billker, O.,
858   2015. PlasmoGEM, a database supporting a community resource for large-scale
859   experimental genetics in malaria parasites. Nucleic Acids Res 43, D1176-1182.
860   https://doi.org/10.1093/nar/gku1143
861 Shi, L., Tu, B.P., 2015. Acetyl-CoA and the regulation of metabolism: mechanisms and
862   consequences. Curr Opin Cell Biol 33, 125–131.
863   https://doi.org/10.1016/j.ceb.2015.02.003
864 Storm, J., Perner, J., Aparicio, I., Patzewitz, E.-M., Olszewski, K., Llinas, M., Engel, P.C., Müller, S.,
865   2011. Plasmodium falciparum glutamate dehydrogenase a is dispensable and not a drug

866    target during erythrocytic development. Malar. J. 10, 193.
867    https://doi.org/10.1186/1475-2875-10-193
868 Storm, J., Sethia, S., Blackburn, G.J., Chokkathukalam, A., Watson, D.G., Breitling, R., Coombs,
869    G.H., Müller, S., 2014. Phosphoenolpyruvate carboxylase identified as a key enzyme in
870    erythrocytic Plasmodium falciparum carbon metabolism. PLoS Pathog 10, e1003876.
871    https://doi.org/10.1371/journal.ppat.1003876
872 Summers, R.L., Pasaje, C.F.A., Pisco, J.P., Striepen, J., Luth, M.R., Kumpornsin, K., Carpenter,
873    E.F., Munro, J.T., Lin, D., Plater, A., Punekar, A.S., Shepherd, A.M., Shepherd, S.M.,
874    Vanaerschot, M., Murithi, J.M., Rubiano, K., Akidil, A., Ottilie, S., Mittal, N., Dilmore,
875    A.H., Won, M., Mandt, R.E.K., McGowen, K., Owen, E., Walpole, C., Llinás, M., Lee,
876    M.C.S., Winzeler, E.A., Fidock, D.A., Gilbert, I.H., Wirth, D.F., Niles, J.C., Baragaña, B.,
877    Lukens, A.K., 2021. Chemogenomics identifies acetyl-coenzyme A synthetase as a target
878    for malaria treatment and prevention. Cell Chem Biol S2451-9456(21)00350–0.
879    https://doi.org/10.1016/j.chembiol.2021.07.010
880 Swift, R.P., Rajaram, K., Keutcha, C., Liu, H.B., Kwan, B., Dziedzic, A., Jedlicka, A.E., Prigge, S.T.,
881    2020a. The NTP generating activity of pyruvate kinase II is critical for apicoplast
882    maintenance in Plasmodium falciparum. Elife 9, e50807.
883    https://doi.org/10.7554/eLife.50807
884 Swift, R.P., Rajaram, K., Liu, H.B., Dziedzic, A., Jedlicka, A.E., Roberts, A.D., Matthews, K.A., Jhun,
885    H., Bumpus, N.N., Tewari, S.G., Wallqvist, A., Prigge, S.T., 2020b. A mevalonate bypass
886    system facilitates elucidation of plastid biology in malaria parasites. PLoS Pathog 16,
887    e1008316. https://doi.org/10.1371/journal.ppat.1008316
888 Swift, R.P., Rajaram, K., Liu, H.B., Prigge, S.T., 2021. Dephospho-CoA kinase, a nuclear-encoded
889    apicoplast protein, remains active and essential after Plasmodium falciparum apicoplast
890    disruption. EMBO J 40, e107247. https://doi.org/10.15252/embj.2020107247
891 Tavoulari, S., Thangaratnarajah, C., Mavridou, V., Harbour, M.E., Martinou, J.-C., Kunji, E.R.,
892    2019. The yeast mitochondrial pyruvate carrier is a hetero-dimer in its functional state.
893    EMBO J 38, e100785. https://doi.org/10.15252/embj.2018100785
894 Tripathi, J., Stoklasa, M., Nayak, S., Low, K.E., Lee, E.Q.H., Rénia, L., Malleret, B., Bozdech, Z.,
895    2023. The artemisinin-induced dormant stages of *Plasmodium falciparum* exhibit
896    hallmarks of cellular senescence and drug resilience.
897    https://doi.org/10.1101/2023.01.29.526019
898 van Dooren, G.G., Stimmler, L.M., McFadden, G.I., 2006. Metabolic maps and functions of the
899    Plasmodium mitochondrion. FEMS Microbiol Rev 30, 596–630.
900    https://doi.org/10.1111/j.1574-6976.2006.00027.x
901 Wang, J., Huang, L., Li, J., Fan, Q., Long, Y., Li, Y., Zhou, B., 2010. Artemisinin directly targets
902    malarial mitochondria through its specific mitochondrial activation. PLoS One 5, e9582.
903    https://doi.org/10.1371/journal.pone.0009582
904 Weiner, J., Kooij, T.W.A., 2016. Phylogenetic profiles of all membrane transport proteins of the
905    malaria parasite highlight new drug targets. Microb Cell 3, 511–521.
906    https://doi.org/10.15698/mic2016.10.534
907 Wideman, J.G., Gawryluk, R.M.R., Gray, M.W., Dacks, J.B., 2013. The ancient and widespread
908    nature of the ER-mitochondria encounter structure. Mol Biol Evol 30, 2044–2049.
909    https://doi.org/10.1093/molbev/mst120

Witkowski, B., Amaratunga, C., Khim, N., Sreng, S., Chim, P., Kim, S., Lim, P., Mao, S., Sopha, C.,
Sam, B., Anderson, J.M., Duong, S., Chuor, C.M., Taylor, W.R.J., Suon, S., Mercereau-
Puijalon, O., Fairhurst, R.M., Menard, D., 2013. Novel phenotypic assays for the
detection of artemisinin-resistant Plasmodium falciparum malaria in Cambodia: in-vitro
and ex-vivo drug-response studies. Lancet Infect Dis 13, 1043–1049.
https://doi.org/10.1016/S1473-3099(13)70252-4
Witkowski, B., Lelièvre, J., Barragán, M.J.L., Laurent, V., Su, X., Berry, A., Benoit-Vical, F., 2010.
Increased tolerance to artemisinin in Plasmodium falciparum is mediated by a
quiescence mechanism. Antimicrob. Agents Chemother. 54, 1872–1877.
https://doi.org/10.1128/AAC.01636-09
World Health Organization, 2023. World malaria report 2023. World Health Organization,
Geneva.
Wunderlich, J., 2022. Updated List of Transport Proteins in Plasmodium falciparum. Front Cell
Infect Microbiol 12, 926541. https://doi.org/10.3389/fcimb.2022.926541
Yang, M., Ke, H., Morrisey, J.M., Mather, M.W., Vaidya, A.B., 2022. Genetic disruption of
isocitrate dehydrogenase arrests the full development of sexual stage parasites in
Plasmodium falciparum. https://doi.org/10.1101/2022.08.05.502965
Yu, X., Wang, C., Zhao, Y., Tang, J., Zhu, M., Platon, L., Culleton, R., Zhu, G., Ménard, D., Zhang,
Q., Cao, J., 2023. Ring-stage growth arrest: Metabolic basis of artemisinin tolerance in
Plasmodium falciparum. iScience 26, 105725.
https://doi.org/10.1016/j.isci.2022.105725
Zhang, H., Guo, J., Li, H., Guan, Y., 2022. Machine learning for artemisinin resistance in malaria
treatment across in vivo-in vitro platforms. iScience 25, 103910.
https://doi.org/10.1016/j.isci.2022.103910
Zhang, M., Wang, C., Otto, T.D., Oberstaller, J., Liao, X., Adapa, S.R., Udenze, K., Bronner, I.F.,
Casandra, D., Mayho, M., Brown, J., Li, S., Swanson, J., Rayner, J.C., Jiang, R.H.Y., Adams,
J.H., 2018. Uncovering the essential genes of the human malaria parasite Plasmodium
falciparum by saturation mutagenesis. Science 360, eaap7847.
https://doi.org/10.1126/science.aap7847
Zhu, L., van der Pluijm, R.W., Kucharski, M., Nayak, S., Tripathi, J., White, N.J., Day, N.P.J., Faiz,
A., Phyo, A.P., Amaratunga, C., Lek, D., Ashley, E.A., Nosten, F., Smithuis, F., Ginsburg,
H., von Seidlein, L., Lin, K., Imwong, M., Chotivanich, K., Mayxay, M., Dhorda, M.,
Nguyen, H.C., Nguyen, T.N.T., Miotto, O., Newton, P.N., Jittamala, P., Tripura, R.,
Pukrittayakamee, S., Peto, T.J., Hien, T.T., Dondorp, A.M., Bozdech, Z., 2022. Artemisinin
resistance in the malaria parasite, Plasmodium falciparum, originates from its initial
transcriptional response. Commun Biol 5, 274. https://doi.org/10.1038/s42003-022-
03215-0
947