Cervicovaginal microbial features predicts *Chlamydia trachomatis* spread to the upper genital tract of infected women

Sangmi Jeong[1], Tammy Tollison[1], Hayden Brochu[1], Hsuan Chou[1], Ian Huntress[1], Kacy S. Yount[2], Xiaojing Zheng[2], Toni Darville[2], Catherine M. O'Connell[2], Xinxia Peng[1,3]

[1]Department of Molecular Biomedical Sciences, North Carolina State University, Raleigh, North Carolina, USA; [2]Department of Pediatrics, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA; [3]Bioinformatics Research Center, North Carolina State University, Raleigh, North Carolina, USA

Corresponding author email : Xinxia Peng, PhD, North Carolina State University, Raleigh, NC 27695, phone (919) 515-4481, fax: (919)-513-6464, xpeng5@ncsu.edu

ABSTRACT

Introduction: *Chlamydia trachomatis (CT)* infection can lead to pelvic inflammatory

disease, infertility and other reproductive sequelae when it ascends to the upper genital

tract. Factors including chlamydial burden, co-infection with other sexually-transmitted

bacterial pathogens and oral contraceptive use influence risk for upper genital tract

spread. Cervicovaginal microbiome composition influences CT susceptibility and we

investigated if it contributes to spread by analyzing amplicon sequence variants (ASVs)

derived from the V4 region of 16S rRNA genes in vaginal samples collected from

women at high risk for CT infection and for whom endometrial infection had been

determined.

Results: Participants were classified as CT negative (CT-, n=77), CT positive at the

cervix (Endo-, n=77), or CT positive at both cervix and endometrium (Endo+, n=66).

Although we were unable to identify many significant differences between CT infected

and uninfected women, differences in abundance of ASVs representing *Lactobacillus*

iners and *L. crispatus* subspecies but not dominant lactobacilli were detected. Twelve

informative ASVs predicted endometrial chlamydial infection (AUC=0.74), with CT ASV

abundance emerging as a key predictor. We also observed a positive correlation

between levels of cervically secreted cytokines previously associated with CT ascension

and abundance of the informative ASVs.

Conclusion: Our findings suggest that vaginal microbial community members may

influence chlamydial spread directly by nutrient limitation and/or disrupting endocervical

epithelial integrity and indirectly by modulating pro-inflammatory signaling and/or

homeostasis of adaptive immunity. Further investigation of these predictive microbial

factors may lead to cervicovaginal microbiome biomarkers useful for identifying women

at increased risk for disease.

INTRODUCTION

Chlamydia trachomatis (CT) is the most common, sexually transmitted, bacterial infection globally (1), with young adults at greatest risk for infection. During childbirth, CT can also be transmitted to a newborn through contact with infected cervical tissue and secretions, resulting in infection of mucous membranes of the eye, oropharynx, urogenital tract, and rectum (2). In addition, CT has been extensively studied as a potential co-factor for human papillomavirus (HPV) enhancing cervical cancer risk through cellular transformation, viral load enhancement, and oncogene overexpression (3). When CT ascends from the cervix to the endometrium or uterus, it can lead to upper genital tract infection, increasing risk for pelvic inflammatory disease and reproductive sequelae such as ectopic pregnancy, or infertility (2). Despite the availability of diagnostics and effective treatments, the asymptomatic nature of CT infection often leads to undiagnosed cases, contributing to its importance as a global health burden (4).

The cervicovaginal microbiome (CVM) is frequently classified into community state types (CSTs), based on the dominant bacterial species present. Using 16s rRNA gene sequencing, Ravel et al. defined the most common CSTs (I-V) (5). CSTs I, II, III, and V are dominated by specific *Lactobacillus* species with overall lower diversity of other microbiota and are generally associated with healthy reproductive health outcomes (6, 7). However, CST IV is comprised of heterogeneous, predominantly anaerobic bacterial genera and *Lactobacillus* species are sparse (5). CST IV is associated with adverse health outcomes such as bacterial vaginosis (BV) (8), pregnancy failure (9), and pre-term birth (10). Vaginal dysbiosis, strongly associated

with CST IV, has also been linked to increased STI vulnerability (7, 8, 11-13). In particular, *in vitro* (14, 15) and *in vivo* (16, 17) studies have associated indole-producers within the CVM with CT evasion of IFN-mediated depletion of tryptophan, an essential chlamydial nutrient (18, 19). Anaerobes also produce biogenic amines and proinflammatory short-chain fatty acids, promoting symptoms and disease (20, 21). In contrast, *Lactobacillus crispatus*-dominated CST I is considered protective against CT infection (8, 13, 22). Mechanisms contributing to protection include production of lactic acid that can directly kill CT (23) while D(-) lactic acid produced by some *Lactobacillus sp.* inhibits CT infection by down-regulating host epithelial cell proliferation (6). CVM-derived metabolites may also act on their host to stimulate DC activation and accelerate immunity (24). STI co-pathogens may also act directly, or in conjunction with microflora, to influence chlamydial infection. We previously observed that women with PID caused by *Neisseria gonorrhoeae* (NG) and CT carried systemic blood transcriptional signatures with greatest activation of cell death pathways and suppression of responses essential for adaptive immunity and CT infection clearance (25). Although these studies have established that CVM composition influences risk for chlamydial cervical infection (8, 13, 22, 26), the extent to which it influences ascending infection has yet to be investigated.

In this study, we sought to investigate the contribution of the CVM to ascending chlamydial infection using high-throughput amplicon sequencing based on 16S ribosomal RNA (16S rRNA) genes, by determining if bacterial members, especially those previously associated with susceptibility to cervical infection, can be implicated in chlamydial spread to the upper genital tract. We analyzed the amplicon sequence

variants (ASVs) derived from the V4 region of 16S rRNA genes to investigate the CVM

of study participants enrolled in T cell Response Against Chlamydia (TRAC), a highly

characterized cohort of women at high risk for CT infection and for whom endometrial

infection had been determined. When participants were categorized into CSTs based on

their CVM composition, CST III (dominated by *Lactobacillus iners*; 30.0%) and CST IV

(characterized as diverse; 54.5%) predominated. At the cervical level, ASV analysis

revealed no significant overall differences in CVM composition between the CT-, Endo-,

and Endo+ groups. Nevertheless, our analysis revealed that certain cervicovaginal

microbial features were predictive of ascended chlamydial infection (Endo+). Including

CT ASV abundance in the analysis significantly improved prediction accuracy, indicating

that CT burden at the cervix is a critical factor in the progression of infection.

Furthermore, we detected a positive correlation between microbial features predicting

CT ascension and levels of selected cytokines in cervical secretions, indicating that

these CVM members may influence ascending infection through their interaction with

host immune responses, acting on CT indirectly and directly. This study underscores

the potential of developing mucosal biomarkers to identify women at higher risk of CT

upper genital tract infection, which could be valuable for clinical practice and future

vaccine trials.

RESULTS

Classification of participants based on STI clinical diagnostic tests

Cervicovaginal swab samples and endometrial tissue biopsy specimens were

collected from 246 women aged 18-35 years in the TRAC cohort (27, 28).

Demographics and sexual history of the cohort were previously described (27). Cervical and endometrial infection status of study participants was determined using nucleic acid amplification tests (NAATs) on relevant samples (27). Twenty-six vaginal swab-derived samples were excluded from this study for the following reasons: three had quality concerns identified during DNA extraction, while the infection status of the remaining twenty-three samples was undetermined because they yielded equivocal results regarding chlamydial detection at the cervix or endometrium during clinical diagnostic testing.

The remaining 220 participants were classified into the following three groups: Endo+, Endo-, and CT- (**Supplementary Table S1**). A total of 143 women who tested positively for CT at their cervix (CT+) were subsequently divided into two groups based on confirmed upper genital tract infection. Women testing positively for CT infection at their cervix and endometrium (n=66) were defined as Endo-positive (Endo+) and the remaining CT infected women who tested positive for CT at their cervix but tested negative for endometrial CT infection (n=77) were grouped as Endo-negative (Endo-). The control group (n=77) was comprised of women testing negatively for cervical chlamydial infection (CT-). However, among the 77 uninfected women, seven samples from participants that were initially diagnosed as CT- by NAAT clinical testing but subsequently identified as CT+ by the detection of CT reads in 16S analysis, were later confirmed to be CT+ by separate quantitation of CT burden in cervical samples by qPCR (29). Consequently, these participants were excluded from 16S microbiome analyses comparing CT+ to uninfected, comparing Endo+ to Endo-, and were reserved for a proof-of-concept analysis. Clinical diagnostic testing at enrollment additionally

determined cervical infection with the bacterial STI pathogens *Neisseria gonorrhoeae*

(NG) and/or *Mycoplasma genitalium* (MG) in all participants (**Supplementary Table S1**)

(27).

Characteristics of cervicovaginal microbiota in the TRAC cohort

The CVM of TRAC participants were profiled using 16S rRNA gene V4 region

sequencing and ASV-based analysis. ASVs identifying common STI bacterial

pathogens were detected in cervicovaginal 16S rRNA including CT, NG, and MG

(**Figure 1A**). CT ASV vaginal abundance amongst participants testing positively for CT

infection positively correlated with CT cervical burden (Spearman correlation coefficient

0.575) (29) previously assayed by qPCR (27). CT ASV sequences were detected in 83

of the 143 CT-infected women (58%) and in 7 of 77 women who were diagnosed as CT-

(9.1%) (**Figure 1A, Supplementary Table S1**). Of 20 NG infected women identified by

clinical diagnostic testing, NG ASV sequences were detected in 17 of 20 (85%), while

NG ASV sequences were detected in 5 out of 200 women testing negatively (2.5%).

Assaying via qPCR indicated these discrepancies most likely were a consequence of

low burden and/or differential site sampling (29). In contrast, MG-identifying ASV

sequences were detected in only 6 of the 38 participants clinically diagnosed with MG

(15.8%), an observation explained by the overall low burden of MG detected in infected

TRAC participants (30). MG abundance in cervical specimens, previously determined

by qPCR (30), was significantly associated with detection of MG ASV in sample libraries

(Mann-Whitney, P=0.0206). Overall, these results support the high quality of this 16S

rRNA based sequencing dataset.

Based on ASV relative abundance from 213 women, the most prevalent genera

in these women were *Lactobacillus, Gardnerella, Prevotella*, and *Megasphaera* (**Figure**

1A, Supplementary Figure 1A). The three most abundant ASVs observed overall were

Lactobacillus iners, Lactobacillus crispatus, and *Gardnerella vaginalis* (**Supplementary**

Figure 1B). Within each group (CT-, Endo-, and Endo+), individuals exhibited a

spectrum of *Lactobacillus* abundance, from *Lactobacillus*-predominant compositions to

Lactobacillus-sparse (**Figure 1A, Supplementary figure 1A-B**). Alpha diversity indices

were assessed at the ASV level to investigate whether diversities of cervicovaginal

microbial communities differed among the Endo+, Endo-, and CT- groups. Neither

Shannon nor Inverse Simpson indices were significantly different between among three

groups (**Supplementary Figure 1C;** Wilcoxon rank-sum test p-values > 0.05).

Community State Types (CSTs) represented within TRAC cohort.

The cervicovaginal microbial composition of TRAC participants was further

evaluated by categorizing each sample according to CST, taxonomic profiles that

represent discrete vaginal microbial communities (**Figure 1B**) (5, 31). The TRAC cohort

exhibits a high prevalence of CT infection (65%, 143 of 220 participants) with CST IV

(characterized as diverse) as the predominant CST type (54.5%, 120 of 220, **Figure 1**).

Almost all the remaining participants were assigned as CST I (dominated by *L. crispatus;*

14.1%, 31 of 220) or CST III (dominated by *L. iners*; 30.0%, 66 of 220) (**Figure 1**). Two

participants were assigned to CST II (dominated by *L. gasseri*) and one participant was

assigned to CST V (dominated by *L. jensenii*) (**Figure 1**). The seven participants that

tested false CT negative were assigned to CST I, III, and IV (**Supplementary Table S1**).

To assess a potential relationship between CST and epidemiological information

in the TRAC cohort, we first evaluated the association between CST and cervical CT

infection status for 213 women excluding seven false negatives (**Supplementary Table**

S1, Supplementary Table S2). However, the association between CT infection status

and CST was not significant (**Supplementary Table S2;** Fisher's exact test P-value >

0.05). Second, the association between CST and demographics and physiologic

information was assessed for 220 women (**Supplementary Table S1, Supplementary**

Table S2). The CST type was significantly associated with Nugent score or race

(**Supplementary Table S2;** Fisher's exact test p-value < 0.05). Of the 119 women in

CST IV, 107 (89.9%) were classified as having BV based on the Nugent score,

compared to lower proportions of BV in the other CST types. CST IV was more common

among Black and Multiracial participants, whereas CST I was more common among

Caucasian participants, consistent with previous studies (5, 12, 32-35).

Significant differences in abundance were observed between the CT+ and CT-

groups for *Lactobacillus* and *Prevotella* species.

Since it has been reported that cervicovaginal microbial profiles differ between

CT- and CT+ individuals (36-40), we investigated CVM compositions of CT+ and CT-

TRAC participants. We used sparse partial least squares discriminant analysis (sPLS-

DA) (41) because this method is effective for supervised classification and feature

selection, aiming to identify the best discriminating features between groups while

handling high-dimensional data. The statistical discrimination performance for

classifying CT- and CT+ women using the first two sPLS-DA components was

estimated by a mean area under the curve (AUC) of 0.70 (**Figure 2A**), indicating that

overall CVM profiles in the presence of CT ASV tended to differ between CT+ and CT-

women. However, when CT ASV was excluded from the discrimination analysis,

performance dropped to a mean AUC of 0.51 (**Figure 2B**).

We also performed sPLS-DA analysis within CSTs, to examine if differences in

CVM composition (excluding CT ASV) could be detected between CT+ and CT-

individuals. In CST I (dominated by *L. crispatus*), CT- women appeared to cluster in a

manner visually distinct from CT+ women but discrimination performance was

statistically inconsistent, with a mean AUC of 0.49, likely due to the small number of

samples assigned to CST I (n=30) (**Supplementary Figure 2A**). Within CST III

(dominated by *L. iners*), cervical CT infection status was moderately discriminated (AUC

mean = 0.71, AUC s.d. = 0.04, n = 63) (**Supplementary Figure 2B**) selecting six low-

abundant *L. crispatus* ASVs for the first sPLS-DA component (**Supplementary Table**

S3). Within CST IV-assigned participants, the discrimination between CT+ and CT-

groups was weak (AUC mean = 0.50, AUC s.d. = 0.04, n = 117) (**Supplementary**

Figure 2C).

To identify cervicovaginal microbiome features that are differentially abundant

between CT- and CT+ groups, we used Microbiome Multivariate Association with Linear

Models (MaAsLin2) (42), which produces a list of significant microbial features with false

discovery rate correction. A total of 28 ASVs differed in abundance between CT+ and

CT- groups, including ASVs assigned to *Fannyhessea vaginae*, *Gardnerella vaginalis*

and multiple *Prevotella* spp, microorganisms previously associated with enhanced

sensitivity to STI infection and BV (7, 8, 11-13) (**Supplementary Table S4**, unadjusted

P-value < 0.05). CT (ASV sequence # ZOtu76) and NG (ASV sequence # ZOtu95) were

more abundant in CT+ women, with Benjamini Hochbergh (BH)-adjusted P-values <

0.0001 and unadjusted P-values < 0.05, respectively. Two *Dialister* ASVs (ASV

sequence #'s ZOtu105 and ZOtu220) were (**Supplementary Table S4**, unadjusted P-

value < 0.05). Four of the 28 ASVs, including NG and 3 ASVs assigned to *Dialister*

invisus, a member of the gut microbiome that has been associated with infections at

other mucosal sites*, Prevotella sp.*, and *L. crispatus* (ASV sequence #'s Zotu96,

Zotu220, Zotu304, and Zotu2052), were also identified as informative ASVs

discriminating cervical CT infection status by sPLS-DA analysis (**Supplementary Table**

S3 and S4).

Although no significant differences in abundance of dominant *Lactobacillus spp.*

ASVs (ASV sequence #s ZOtu1, ZOtu2, ZOtu11, ZOtu32, and ZOtu374) were detected

between CT+ and CT- participants in the overall cohort based on their predominant

ASVs, differences were observed in 7 uncommon ASVs assigned to *L. iners*, *L.*

crispatus, *L. jensenii*, *L. gasseri* and *L. colini* (**Supplementary Table S4,**

Supplementary Figure 3). To investigate if these low-abundance ASVs were

sequencing artifact, we conducted multiple sequence alignments to identify the position

of mismatches between these low-abundant *Lactobacillus* ASV sequences and their

corresponding reference sequence, reasoning that errors generated by merging of

paired-end reads would be disproportionately concentrated in the middle of sequences.

The 7 ASV sequences were aligned with the corresponding ASV sequences of their

most abundant *Lactobacillus sp.* ASVs listed in **Supplementary Table S4**, and V4

region sequences of *Lactobacillus sp.* curated in the RDP 16S No18 reference

database (43) (**Supplementary Figure 4**). The most abundant *Lactobacillus* ASVs

(ASV sequence #s ZOtu1, ZOtu2, ZOtu11, ZOtu32, and ZOtu374) exactly matched their

corresponding V4 region reference sequences (**Supplementary Figure 4**). In contrast,

the 7 low-abundant *Lactobacillus* ASV sequences contained mismatches at 20 of 253

nucleotides in the V4 region, except for one ASV (ASV sequence # ZOtu459).

Intriguingly, these nucleotide variants frequently occurred at the same 20 positions

along the *Lactobacillus* V4 region, rather than being concentrated in the middle of ASV

sequences (**Supplementary Figure 4**). These results suggested that the observed

sequence variants in these low-abundant *Lactobacillus* ASVs reflected biological

variation in *Lactobacillus* subspecies or strains that composed the CVM of TRAC

participants.

We investigated whether CVM features could predict CT infection status within

this high STI risk cohort, the majority of whom had BV at enrollment (27)

(**Supplementary Table S1**). We performed a random forest-based analysis using ASV

abundance, while excluding the CT ASV. In brief, we randomly partitioned the samples

from CT+ and CT- groups into a training dataset and a separate test dataset. We built a

predictive model using the training dataset and evaluated its performance by predicting

the CT infection status using the independent test dataset (see Methods for details).

The prediction performance was assessed using the average receiver operating

characteristic (ROC) curves across 100 replicates. The corresponding average AUC

was found to be 0.61 for the prediction using all ASVs excluding the CT ASV

(**Supplementary Figure 5**). Thirteen ASVs contributed to prediction of absent CT

infection (**Supplementary Table S5**). NG ASV (ASV sequence# ZOtu95), more

abundant in CT+ women (**Supplementary Table S4**), was the strongest contributor to

the prediction. NG is not considered a member of the CVM, thus its contribution to this

prediction likely derives from the frequency of co-infection with CT in women reporting

high risk behaviors for STI acquisition. Four *Syntrophococcus sucromutans*-assigned

ASVs and four *Prevotella spp.* ASVs, which showed differential abundance between

CT+ and CT- women (**Supplementary Table S4**), were found to be predictors of absent

CT infection (**Supplementary Table S5**). However, the ASVs of *S. sucromutans* (ASV

sequence #s ZOtu709, ZOtu1481, ZOtu1814, and ZOtu2372) have low confidence

genus-level assignments, indicating that these strains, currently classified as members

of the class *Clostridia*, require further investigation to resolve their taxonomic

classification.

Cervicovaginal microbial features are predictive of susceptibility to ascended

chlamydial infection

Given the severe sequelae that can arise after CT infection ascends to involve

the endometrium and fallopian tubes and the absence of diagnosing this outcome non-

invasively, we investigated if CVM profiles of CT+ women were reflective of CT

endometrial infection. We first examined the overall differences in CVM compositions

between Endo- and Endo+ patients using sPLS-DA analysis, as before (see also

Methods). Endo+ and Endo- women showed distinguishable separations by the first two

sPLS-DA components including CT ASV (**Figure 2C**; AUC mean = 0.76, AUC s.d. =

0.01). The CT ASV (ASV sequence # ZOtu76) and two *Lactobacillus* ASVs (ASV

sequence #s ZOtu915 and ZOtu914) contributed the most to the first component, while

more than 50 ASVs were identified as important features for the second component

discriminating Endo+ and Endo- women (**Supplementary Table S6**). Despite the high

number of important features identified, in the absence of CT ASV, discrimination

between Endo+ and Endo- weakened with an AUC of 0.55 (**Figure 2D**).

We also examined the CVM profiles within CST I, CST III, and CST IV types

separately for CT endometrial infection status using sPLS-DA analysis. Endo- women

were visually separable from Endo+ women within CST I with the first sPLS-DA

component including 8 *Lactobacillus* ASVs (**Supplementary Table S6**), but the

statistical performance was very low, with a mean AUC of 0.47 (**Supplementary Figure**

2D), again likely due to the small number of women within CST I (n=23). The

discrimination of endometrial infection status within CST III was also weak (AUC mean

= 0.58, AUC s.d. = 0.09, n=43) (**Supplementary Figure 2E**). However, endometrial

infection status within CST IV (characterized as diverse) can be discriminated (AUC

mean = 0.71, AUC s.d. = 0.03, n=76) (**Supplementary Figure 2F**) with the CT ASV

solely contributing to the first sPLS-DA component and the interaction of 19 ASVs

including 4 *Prevotella* ASVs and a *Mycoplasma microti*-assigned ASV (# ZOtu45)

contributing the second sPLS-DA component (**Supplementary Table S6**).

To identify CVM features that would be most informative in predicting CT

ascension, we performed random forest-based analysis using ASV abundances as

features to predict individual patients in Endo+ or Endo- groups. Since CT is not only a

key factor of ascended infection but potentially modulates the cervical and endometrial

environment for other microbes, we assessed the predictive performance of the model

in three scenarios: (A) prediction with all ASVs including CT ASV (# ZOtu76) (**Figure**

3A), (B) prediction with all ASVs but excluding CT ASV (**Figure 3B**), and (C) prediction

with only CT ASV (**Figure 3C**). Average AUCs were 0.74 for the model with all ASVs,

and 0.57 for the model that excludes the CT ASV (**Figure 3A-B**), indicating the

importance of CT ASV abundance for prediction. When the model was trained by the

CT ASV alone it resulted in an average AUC of 0.74 (**Figure 3C**).

In the prediction with all ASVs including CT ASV, CT and 11 additional ASVs

were identified as informative features using selection frequency based on mean

decrease in accuracy (**Table 1**; feature selection frequency 80 out of 100 replicates).

These 11 ASVs were also selected as informative features in the prediction using all

ASVs but excluding CT ASV (**Table 1**; feature selection frequency $\geq$ 80 out of 100

replicates). Notably, there was no overlap between these features and those selected

as predictors of absent CT cervical infection (**Supplementary Table S5**). Among the

identified 12 ASVs, four were identified with higher abundance in Endo+ women (ASV

sequence #s ZOtu76, ZOtu233, ZOtu1808, ZOtu567) (**Table 1**). Of the four? highly

abundant ASVs in Endo+ women, three were also identified as important features

discriminating Endo+ and Endo- patients in the above sPLS-DA analysis (ASV

sequence #s ZOtu76, ZOtu233, and ZOtu1808) (**Table 1** and **Supplementary Table**

S6). All three ASVs assigned to *Sutterella stercoricanis* (ASV sequence #'s ZOtu110,

ZOtu184, and ZOtu180) exhibited lower abundance in Endo+ women (**Table 1**).

Actinotignum schaalii ASV (ASV sequence # Zotu281) and *Haemophilus haemolyticus*

ASV (ASV sequence # Zotu103) were also identified as predictors, which were less

abundant in Endo+. *Mycoplasma microti* (ASV sequence # Zotu45), a species

previously isolated from the respiratory tracts of prairie voles (43), seemed an unlikely

component of the CVM. Subsequent examination determined that the assigned ASV

shared 100% identity with uncultured *Mycoplasma sp.* clone Mnola (44), now

Candidatus Malacoplasma girerdii (45), a strict endosymbiont of *Trichomonas vaginalis*

(TV). Within TRAC, we observed that detection of *Ca.* M. girerdii ASV reads coincided

with a positive TV clinical diagnostic finding in 14 of 37 participants (37.8%) but was

only detected in 7 of 183 (3.8%) of participants who tested negatively for TV suggesting

that this was the more correct assignment. Overall, these results suggest that multiple

CVM features together can be predictive of susceptibility to CT ascension but the

amount of CT at the cervix is a key factor.

Cervicovaginal microbial features associated with ascending CT infection are

positively correlated with cytokines in cervical secretions associated with

endometrial CT infection

To better understand possible functional consequences of these 12 CT

ascension-related microbial features on host immune responses, we explored their

associations with levels of 7 cytokines in TRAC cervical secretions that had been

previously associated with CT ascension (44). The following cytokines, CXCL10, TNF- α,

IL17A, CXCL9, CXCL11, CCL4, and CXCL13, were positively associated with

endometrial infection (unadjusted $P<0.05$) in a univariable regression model. Using

canonical correlation analysis (CCA) (45), we investigated whether there was an overall

correlation between these cytokine levels and the predictive microbial features. With 12

ASVs, including CT, we observed a positive correlation between microbial abundances

and cytokine levels (**Figure 4A**; canonical correlation = 0.54). There was also a positive

but weaker association between cytokine levels and CVM-derived ASVs when CT ASV

was excluded (**Figure 4B**; canonical correlation = 0.43). CT ASV abundance also

correlated weakly with the seven cytokines (**Figure 4C**; canonical correlation = 0.49).

Together, these results indicate that host immune responses are not exclusively

modulated by CT abundance. The CCA-based analyses suggest that the predictive

CVM microbial features influence CT ascension directly and indirectly, through

interaction with host immune responses.

A proof-of-concept analysis for predicting ascending CT infection using the

twelve informative ASVs

We identified twelve ASVs that are predictive of susceptibility to ascending CT

infection using random forest-based prediction and ASV abundance of Endo+ and

Endo- women (**Table 1 and Figure 3A**). Four false negative CT+ participants

(**Supplementary Table S1**) had residual endometrium-derived samples for additional

testing and were used for proof-of-concept prediction of Endo+ vs Endo- here. An

optimal threshold of 0.55 was derived from the averaged ROC curve of the model

informed by all ASVs (**Figure 3A**) and was used to determine between Endo+ and

Endo- (**Supplementary Figure 6**). The model correctly predicted CT ascension status

for two out of four participants based on confirmed detection of CT DNA (29). CT ASV

abundance was considered as a key factor for the prediction of CT ascension but was

unable to predict Endo+ vs Endo- status alone (data not shown). This proof-of-concept

analysis of four false negative CT+ participants could not validate the 12 ASVs including

CT ASV due to very small sample size. Validation of the 12 CVM features identified

requires testing in a larger, independent cohort.

DISCUSSION

Recent literature has revealed that cervicovaginal microbiota may modulate

susceptibility to infections such as human papillomavirus (HPV), CT, NG and MG

infections (46). CVM composition has been associated with increased risk of incident

CT infection (16, 22, 46, 47). In this study, we sought to determine whether CVM

composition facilitates CT ascension from the cervix to the upper genital tract by

profiling 16S rRNA genes obtained from vaginal swabs collected at enrollment from the

TRAC cohort. Although we were unable to identify many significant differences between

CT infected and uninfected women, we identified 12 cervicovaginal microbial features

that predicted active upper tract infection. Chlamydial cervical burden was a key driver

of ascent because vaginal CT ASV abundance alone, effectively predicted this outcome

with AUC of 0.74 for sensitivity and specificity. We also observed positive correlation

between levels of 7 cytokines previously associated with CT ascension and abundance

of CVM-derived ASVs, indicating that local host immune responses during infection

were not exclusively modulated by pathogen load.

TRAC is comprised of women at elevated risk for STI infection and a striking

feature of this cohort is the very high rate of CT infection, 67%, amongst enrollees,

reflective of highly targeted recruitment strategies (27). This may have blunted our

ability to detect significant differences in the composition of the CVM, at the level of CST,

for individual genera or species that associated with protection against CT infection.

Unsurprisingly, most participants were assigned to CST III or CST IV, previously

recognized as neutral or favorable for CT infection (7, 8, 11-13). Within CST III, CT-

women were separable from CT+ using sPLS-DA components containing *L. crispatus*-

assigned ASVs. Amongst CST I-assigned participants, the CST most highly associated

with protection from CT infection (8, 13, 22), CVM composition did not differ sufficiently

to cluster CT infected and uninfected participants effectively. TRAC was designed as a

longitudinal study to investigate T cell responses important for protection from incident

chlamydial infection, but the high rate of CT positivity at enrollment combined with a

stringent requirement that participants have completed a minimum of three follow up

assessments, provided insufficient power to confirm previously identified associations of

CST with infection susceptibility. Nevertheless, a random forest-based analysis using all

ASVs (excluding CT) was moderately predictive of CT- women **(Supplementary Table**

S5), while MaAsLin2 analysis identified differentially abundant community members

previously associated with susceptibility **(Supplementary Table S4)**. Conserved

between both analyses were ASVs assigned to *Prevotella sp.* and *S. sucromutans* that

associated with CT infection. Studies, *in vitro* (14, 15) and *in vivo* (16, 17) have

associated indole-producers, e.g. *Prevotella sp.*, with CT evasion of IFNγ-mediated

tryptophan depletion (18, 19). Of the *Prevotella sp.* discriminating CT infection in our

analysis (48), only *P. brunnea* is an indole producer (48, 49), but it is also possible that

they modulate host factors, indirectly influencing chlamydial susceptibility. *S.*

sucromutans, in association with *Prevotella sp.*, serves as a biomarker of gut barrier

dysfunction (leaky gut) (50) where increased luminal abundance of Zonulin-1, an

endogenous tight-junction protein, was associated with decreased abundance of these

species. Directly relevant to our study, recent work by Hinderfield and colleagues (51), demonstrated that CST IV-associated microorganisms promote paracellular permeability of human ectocervical monolayers with coincident down-regulation of Zonulin and Occludin expression, leading them to hypothesize that undermining cervical epithelium integrity enhances susceptibility to TV and other STI pathogens (51). Thus, a CVM that contributes essential substrates and improves pathogen access to the cervical epithelium can increase susceptibility to chlamydial infection.

In contrast, lactic acid produced by *Lactobacillus sp.* directly kills CT (23) while D(-) lactic acid produced by some *Lactobacillus sp.* inhibits CT infection by down-regulating host epithelial cell proliferation (6). In this study, we were unable to detect associations of the most abundant *Lactobacillus* species with CT positivity but identified differences in low abundance strains **(Supplementary Table S4)**. We questioned whether the low-abundant *Lactobacillus* ASVs that were differentially abundant by MaAsLin2 analysis were genuine *Lactobacillus* species or accidental assignments of artificial sequences to *Lactobacillus* species, given the taxonomic complexity among *Lactobacillus* species (52) and the limitations of characterizing these species through 16S rRNA gene sequencing (53). Although ASVs have intrinsic biological meaning independent of reference database or the specific context of a study (54), the taxonomic resolution is much lower compared to the full length of 16S rRNA sequences or metagenome sequences (55). Recent genomic studies identifying genetic elements that control functional differences between and within *Lactobacillus species*, such as the role of bile salt hydrolases that promote colonization of vertebrate-adapted niches (56), and the ability to metabolize extracellular glycogen important for microbial growth in the

female genital tract (57, 58) as well as identification of novel urogenital tract species (59)

suggest that it is premature to conclude that the differences we detected between CT

infected and uninfected women are artifactual, and future investigations should consider

implementing typing approaches with superior resolution that will enable us to

determine if further investigation into the biological roles of individual *Lactobacillus*

subspecies in cervical CT infection is needed.

The TRAC cohort provided a unique opportunity to investigate the CVM

contribution to chlamydial ascension from the cervix to the upper genital tract. Using

sPLS-DA, we were able to observe overall differences in CVM compositions between

Endo- and Endo+ participants, although discriminatory resolution dropped considerably

when the CT ASV was excluded, emphasizing the importance of CT cervical abundance

in driving ascension. At the level of CST, discriminatory performance was highest in

CST IV, but much reduced for CST I and CST III participants, suggesting that CVM-

derived factors that promote susceptibility to CT infection also contribute to its ascent.

However, we did not detect much overlap in the specific CVM features differentially

abundant between CT- and CT+ participants by MaAsLin2 analysis and those selected

by random forest analysis as predictive of spread with the obvious exception of the CT

ASV.

Many of the CVM features selected by our analysis have not previously been

associated with human chlamydial infection but have been investigated in the context of

the respiratory or gut microbiome. Through "nutritional immunity", the host may limit iron

availability as a means of protection against invading pathogens (60). CT requires iron

(61, 62) and may experience additional competition for this essential nutrient from CVM

members. Iron restriction also impacts host cellular metabolism, leading to differential

host gene expression with downstream consequences for availability of other key

substrates required for chlamydial developmental success (63). Antithetically,

transcription of the *trpBA* operon essential for chlamydial scavenging of exogenous

indole to tryptophan is iron-regulated (64) leading to the proposal that a low iron

environment could increase risk for sequelae (64). *H. haemolyticus* is a respiratory tract

commensal that limits infection with the pathogenic *H. influenzae* through competition

(65), because it uses its superior ability to sequester iron to facilitate occupation of their

shared niche (66, 67). Its relatively lower abundance in Endo+ participants may indicate

that chlamydial need for iron predominates during natural infection and that *H.*

haemolyticus abundance modulates ascended infection through its negative influence

on chlamydial growth.

Also less abundant in Endo+ participants, *S. stercoricanis* has not been

previously described as a significant member of CVM but within the human gut

microbiome, its reduced abundance has been identified as a predictor of enhanced

susceptibility to respiratory infection (68). *Sutterella sp.* are epithelium-adherent (69),

but do not contribute to gut dysbiosis or loss of gut epithelial integrity. In the context of

STI infection, their abundance is significantly increased in the rectum of HIV negative,

STI uninfected men, and CD4/CD8 ratio positively correlated with *Sutterella* in HIV-

infected patients, consistent with an association with T cell-mediated adaptive

responses independent of active bacterial infection (70). They are mildly pro-

inflammatory, and it has been proposed that their association with their human host is

mutualistic through their contribution to steady state induction of Th17 cell differentiation

(71, 72). We previously reported that IL-17A transcript was significantly increased in

blood of participants with high cervical *C. trachomatis* burden compared to those with

low cervical burden and uninfected participants (25) and that cervical secretion of IL-

17A was associated with increased odds of endometrial infection (44). Nevertheless, we

observed increased frequencies of CD4 Th17 cells in peripheral blood of CT-infected

TRAC participants compared to uninfected participants, and a further increase in

participants without recurring infection compared to those who were reinfected with CT

within a year after enrollment (73). Together, these results suggest that CT infection

elicits local and systemic Th17 responses, and that the systemic Th17 responses

detected were associated with reduced reinfection. It is possible that occupation of a

niche by benign *S. stercoricanis* within the CVM of Endo- participants protects against

expansion of members that positively contribute to chlamydial burden by provision of

necessary intermediates or by compromise of epithelial integrity. It is also possible that

its reduced abundance in Endo+ women contributed to reduced CD4 T cell activity,

ultimately delaying resolution of chlamydial infection. Future studies that are more

informative of function, e.g. transcriptomics, metabolomics, may identify common

metabolites or pathways that can be associated with CT spread to the upper

reproductive tract, while profiling of adaptive immune cell populations recruited to the

infection site will lend further insight into mechanisms of chlamydial clearance.

We previously observed the importance of CT/NG coinfection on risk for CT

reinfection (27). We also reported altered bloodborne transcriptional signatures pointing

to suppressed host adaptive responses with CT/NG coinfection in women with pelvic

inflammatory disease (25). However, we did not observe TV nor MG co-infection

associating with ascended CT in TRAC participants (27, 30). In this study, an ASV that

we ultimately re-assigned to *Ca*. M. girerdii (74, 75) was identified as a CVM feature

predictive of ascended infection. Members of this large and diverse class of *Mollicutes,*

distinguished by a small genome and lacking a cell wall, such as *M. hominis* and

Ureaplasma sp., have been identified as commensals and/or opportunistic genital tract

pathogens, although meta-analyses have not effectively resolved the extent of their

impact on female infertility or other reproductive outcomes including preterm birth (76,

77). Intriguingly, *M. hominis* and *Ca. M. girerdii*, alone or together, can live intracellularly

or on the surface of TV, accelerating the parasite's growth (78, 79) and modulating its

pro-inflammatory potential. Relevant to CT infection because of the importance of cell-

mediated responses for chlamydial clearance, TV harboring *M. hominis*, but not alone,

synergistically upregulates IL-8, IL-1β, and TNF-α production as well as inducing

production of the Th-17 polarizing cytokine IL-23 by human monocytic THP-1 cells with

coincubation (80). If, as we speculate, recruitment of chlamydia-specific Th1 and Th17

cells to the genital tract is essential for control, TV coinfection, in some but not all

instances, might promote this outcome, explaining why *Ca.* M. girerdii ASV was

associated with Endo- women. Confusingly, ASVs assigned to the other TV

endosymbiont *M. hominis* were also detected in many TRAC participants but were not

selected as predictive features. We were not able to immediately resolve this

contradictory observation because rRNA from TV was not captured in this analysis

since it is a protozoan and the clinical diagnostic was not quantitative, but it highlights

the potential for factors that influence pathogen success to be mediated through

interactions between microbiome members, including those that cannot be detected

using a 16S rRNA profiling approach.

ASVs contributing to prediction of ascended CT infection that were more

abundant in Endo+ women included *Actinobaculum massiliense,* a urinary tract

commensal or opportunistic pathogen (81), and reportedly a rare cause of PID (82). A

recent proteomic analysis identified two highly expressed subtilisin-like proteases which

may cleave cytokines (83), providing potential to modulate the host inflammatory

response to itself and CT. ASVs assigned to *Butyrivibrio hungatei* and the genus

Acetothermia were also more abundant in Endo+ participants. Neither bacterium is well

documented as a member of CVM. In the context of gut microflora, *B. hungatei* is

thought to act positively because reduced abundance of this butyrate producer has

been associated with dysbiosis and disorders of the gut-brain axis such as Alzheimer's

Disease (84). Butyrate is a short chain fatty acid (SCFA) that modulates colon health by

multiple mechanisms. In low amounts it serves as a carbon source for other

commensals, while at higher doses it can be toxic, targeting other members of normal

flora or pathogens such as *Salmonella sp.* (85). Colonic enterocytes transport butyrate

as a preferred substrate (86) promoting epithelial health and integrity. However, in the

female genital tract, SCFAs are associated with the production of pro-inflammatory

cytokines and perturbation of epithelial integrity (87).

With potential translational impact, random-forest analysis revealed the

importance of cervical CT abundance in prediction of concomitant upper genital tract

spread. Using the CT ASV alone is a highly informative and simple feature, capable of

predicting chlamydial ascension with equivalent AUC performance to the prediction

using 12 ASV CVM biomarkers. A major barrier to preventing CT disease is delayed diagnosis, because infection is often asymptomatic (88), but both clinical and subclinical upper-tract inflammation can lead to chronic oviduct damage (89). Diagnostic biomarkers of ascended infection would be particularly useful for evaluation of therapeutics and vaccines (90-92) and improve case management, since women with clinical PID receive a longer antibiotic course compared to asymptomatic women with CT (93). Abundance of the CT ASV appears to be a better predictor of ascended CT infection than blood-borne transcriptional biosignatures (94) or profiling circulating immune cells (73) for this cohort of highly exposed women but was unsuccessful when applied to our small proof-of-concept set. Nevertheless, CT ASV with the CVM-derived selective features provided the best correlation with local cytokines, indicating the potential for further improved prediction with inclusion of host immune features. We previously performed a small pilot study that indicated that whole transcriptome sequencing analysis of ribosomal RNA-depleted total RNA isolated from cervical swab samples contained pathogen-specific sequences from women with confirmed sexually transmitted bacterial pathogens (95). Simultaneously, we identified and quantified their active microbial communities and after integration with their associated host-derived reads, we detected clustered host transcriptional profiles reflecting microbiome differences and STI infection. Together, these findings indicate that cervical sampling can yield sensitive and robust biomarkers for evaluation of candidate chlamydial vaccines, such as those currently in development or advancing towards Phase I (96) and Phase II trials.

High throughput sequencing of 16S rRNA enabled us to efficiently identify CVM

communities in the TRAC cohort, but taxonomic resolution for microbial species or

strains was frequently too low to confidently determine their taxonomy as cervicovaginal

microbiome. Nonetheless, we were able to detect STI pathogenic microbes, as well as

highly variable microbes such as *Lactobacillus sp.* and *Mycoplasma sp.* using ASV

sequences. Copy number of the 16S rRNA gene can vary widely, ranging from 1 to 37

in bacteria and 1 to 5 in archaea (97). This variation in 16S rRNA gene copy number

(GCN) limits the accuracy of quantifying 16S rRNA genes and CT 16S reads, potentially

introducing biases in subsequent analyses (98). Indeed, this effect may be reflected in

our variable success detecting ASVs associated with NG (four copies) versus CT (two

copies) and MG (single copy) (99) in samples from women testing positively with highly

sensitive clinical diagnostic tests. Although novel methods have been developed to

correct for 16S rRNA GCN (100-103), we did not apply 16S rRNA GCN prediction tools

for precise quantification of 16S rRNA genes in this study. In future work, we can adopt

full-length 16S rRNA sequencing as an alternative to short reads of 16S rRNA to

improve taxonomic resolution and quantification accuracy.

This study had several additional limitations. For example, the participants of

TRAC cohort were highly exposed to CT infection, which may have introduced bias in

the representation of the CVM communities in general. Vaginal swab samples for

microbiome sequencing were collected only at enrollment when participants were

actively infected, so there were no baseline samples to investigate temporal changes in

CVM features related to the development of CT ascending infection. Even if we had

chosen to sample participants longitudinally, all received antibiotic treatment at

enrollment, which could have affected CVM composition. Also, 16S rRNA-based

sequencing provided information only on the composition of CVM profiles. This limits

the ability to understand the functional roles that these microbial communities may play

in modulating host immune resistance or susceptibility to CT ascension. We did not

investigate other factors such as host genetic heterogeneity and the effect of innate

immunity on ascended CT infection.

In future studies, the integration of other techniques such as transcriptomics and

metabolomics with 16S rRNA based sequencing, studies that have been initiated with

establishment of an additional cohort of CT-infected women (TRAC2) (104) are

expected to facilitate the development of deeper functional insights and identification of

relevant biological pathways. Additionally, expanding the analysis with other relevant

members of the CVM including parasites such as TV and viruses could also enhance

our understanding of microbial interactions with the host and improve the prediction of

upper genital tract infection spread. These multi-omics analyses may also contribute to

the identification of biomarkers of protection or disease risk that could contribute to

assessment of vaccine efficacy in future clinical trials.

ACKNOWLEDGEMENTS. We thank the women who agreed to participate in this study;

Ingrid Macio, Melinda Petrina, Carol Priest, Abi Jett, and Lorna Rabe under direction of

Harold Wiesenfeld and Sharon Hillier, for their efforts in the clinic and the microbiology

laboratory; and the staff at the Allegheny County Health Department STD Clinic, for

their efforts. We would also like to acknowledge the contribution of Dr. Ian Huntress,

who sadly passed away during the performance of these studies.

Availability of data and material

Raw 16S rRNA gene sequencing data supporting the conclusions of this article is publicly available at NCBI's Short Read Archive (SRA) under BioProject accession number PRJNA1136868 (https://www.ncbi.nlm.nih.gov/bioproject/). **Supplementary Figures S1–S8** and **Supplementary Tables S1–S3, S5**, and **S7** are included in **Additional File 1: Supplementary Materials**. **Supplementary Table S4** is provided in **Additional File 2**, and **Supplementary Table S6** is provided in **Additional File 3.** The legends of **Supplementary Tables S4** and **S6** are provided in **Additional File 1: Supplementary Materials.**

Financial support. This work was supported by the National Institute of Allergy and Infectious Diseases via U19 AI084024 and R01AI170959.

METHODS

T Cell Response Against Chlamydia (TRAC) Cohort

This cohort was composed of mainly young (median age, 21 years; range, 18-35 years), single (89%), African American (66%) women at high risk for the acquisition of CT (**Supplementary Table S1**). Inclusion criteria indicating high-risk status included > 3 sexual partners in the previous 6 months, ≤ 14 years of age at sexual debut, history of pelvic inflammatory disease (PID), or presentation to the recruitment site with any of the following: presence of mucopurulent cervicitis on exam, or sexual contact with a partner known to be infected with CT or NG or non-gonococcal non-chlamydial urethritis.

Women with a current diagnosis of PID according to the Centers for Disease Control

and Prevention guidelines were excluded. Additional exclusion criteria were pregnancy,

uterine procedure or miscarriage in the preceding 60 days, menopause, hysterectomy,

antibiotic therapy in the preceding 14 days, and allergy to study medications. A total of

246 TRAC participants were enrolled into a longitudinal study designed to investigate T

cell responses important for protection from incident chlamydial infection over 12

months of follow-up between February 2011 and August 2014 and were recruited from

the Allegheny County Health Department's Sexually Transmitted Diseases Clinic,

Magee-Womens Hospital (MWH) Ambulatory Care Clinic, and the Reproductive

Infectious Disease Research Unit at MWH in Pittsburgh, PA. This study complied with

the Declaration of Helsinki guidelines and all study participants provided informed

consent prior to initiation of study procedures. The Institutional Review Boards at the

University of Pittsburgh and the University of North Carolina approved the study.

Participants completed questionnaires regarding obstetric/gynecologic history,

behavioral practices, sexual histories, contraceptive methods, and symptoms according

to study protocols. Clinical, histological, and microbiological testing was performed, and

blood and endometrial samples were obtained at enrollment, after which all participants

received single-dose agents to treat *gonorrhoea* (ceftriaxone, 250 mg intramuscularly)

and chlamydia (azithromycin, 1 g orally). Participants in this cohort were assessed for

cervical and endometrial infection using Aptima Combo 2 (AC2) (Hologic, Marlborough,

MA) NAAT with overall infection rates of 67% and 8.5% for CT and NG, respectively.

MG and TV infection was determined using Aptima MG and Aptima TV diagnostics

(Hologic) respectively.

16S library preparation and sequencing

DNA was extracted from cervicovaginal swabs stored at -80 ℃ using the AllPrep DNA/RNA Mini Kit (Qiagen, Valencia, CA). Before extraction, the manufacturer's protocol was optimized with minor modifications to achieve the best performance in terms of DNA/RNA yield and equal recovery of six different species comprising the ATCC Vaginal Microbiome Whole Cell Mix (MSA-2007). Average yields were 52.8 ng/µl yield of DNA in a final volume of 100µl per sample. The V4 regions of 16S rRNA gene were amplified using the Illumina 16S V4 primer set of 515F [GTGYCAGCMGCCGCGGTAA] and 806R [GGACTACNVGGGTWTCTAAT] (105, 106) designed for dual indexing as described by the Earth Microbiome Project protocol (https://earthmicrobiome.org/protocols-and-standards/16s/). Sequencing libraries were prepared according to the Illumina's 16S Metagenomic Sequencing Library Preparation Guide (https://support.illumina.com/downloads/16s_metagenomic_sequencing_library_prepara tion.html; Part # 15044223 Rev. B), and sequenced on an Illumina NextSeq 500 Sequencer (Illumina, San Diego, CA).

16S rRNA-seq data processing

The 16S rRNA data analysis pipeline is outlined in **Supplementary Figure 7.** Raw fastq files containing paired-end read sequences per each sample were demultiplexed and preprocessed into amplicon sequence variants (ASVs) using bioinformatics software, USEARCH v11.0 (107) and PEAR (108). The fastq_eestats2

function in USEARCH was used to determine if trimming was needed to optimize read

merging, based on expected errors for 150bp forward and reverse reads across

different length cutoffs (100, 110, 120, 130, 140, and 150bp), which was 150bp for our

data. Merged reads were expected to be >252bp with short overlap nucleotides (<10bp).

PEAR, known for its robustness with short overlapping regions, was employed for read

merging, utilizing a minimum overlap size of 8 bases (108). Following read merging,

primer sequences were trimmed from both ends, and low-quality reads were filtered at a

maximum expected error rate of 1.0.

Sequence denoising (error-correcting) and the generation of zero-radius OTUs

(zOTUs), referred to as amplicon sequence variants (ASVs), were achieved using

unoise3 function in USEARCH (107). The unoise3 algorithm prioritizes read abundance

over sequence differences by processing sequences based on descending order of

read abundance (109). As a result, a rare sequence can be incorporated into a highly

abundant centroid sequence even if its difference is relatively high and may be

discarded by chance depending on the order of pooled unique sequences (107, 109).

To reproduce the same set of ASV sequences in TRAC cohort despite this limitation of

randomly discarding low-abundant sequences, we added additional steps to the

standard denoising and ASV generation process. We reordered unique sequences

based on read frequency, read length, entropy, and confidence level of taxonomy

assignment, all in decreasing order except for read length. Considering the optimal

length of V4 region of 16S rRNA DNA to be 253bp, we calculated the distances

between 253bp and the length of read, and then arranged the reads from the smallest

distance to the longest distance away from 253bp. Using the rearranged ~995,000

unique read sequences, we consistently reproduced the same total set of 2,777 ASVs

from an average of ~151,000 reads per sample. Of 2,777 ASV sequences, the Burrows-

Wheeler Alignment tool (110) aligned 569 to the human reference genome (GRCh38)

and these were excluded. We also removed ASV sequences shorter than 252bp, with

1,990 ASVs retained for further analysis. The inferred ASVs were assigned taxonomy

using the RDP Classifier 16S trainset No. 18 raw training database

(https://sourceforge.net/projects/rdp-classifier/files/RDP_Classifier_TrainingData/).

For analyses based on ASV abundances of CT+ and CT- women, 783 out of

1,990 ASVs were excluded because they had fewer than 50 read counts across CT+

and CT- groups because these low-abundance ASVs represented experimental artifacts

generated during sequencing. Similarly, of 1,990 ASVs, 1,043 with fewer than 50 reads

across Endo+ and Endo- women were excluded from analyses based on ASV

abundances of Endo+ and Endo- women.

Multiple sequence alignment of 7 low-abundant *Lactobacillus sp.* ASV sequences

(**Supplementary Table S4**) with their corresponding sequences of the most dominant

Lactobacillus sp. ASVs and the V4 region sequences of *Lactobacillus sp.* obtained from

the RDP 16S No18 reference database was performed using MUSCLE 3.8.31v (111).

The alignment was visualized using UGENE (112) to with the goal of identifying

mismatches in the central region of the 252bp ASV sequences (**Figure S4**).

Modification of reference sequences for V4 regions in RDP Classifier

RDP Classifier 16S trainset No18 raw training database provides both bacterial

and archaeal 16S rRNA reference sequences. Among a total of 21,195 reference

sequences, nearly 40% had identical V4 region sequences for multiple species ranging

from 2 to 65 different species in the RDP Classifier database (**Supplementary Table**

S7). In such cases, sintax function for taxonomic assignment in USEARCH randomly

chooses a species among multiple species given by the reference annotation (113). In

addition, since the RDP Classifier database is not limited to human-related microbial

taxonomy annotation, assigned species may include those not typically observed in the

human cervicovaginal microbiota. For example, although *L. crispatus* and *L. jensenii* are

commonly found in the cervicovaginal environment, taxonomic assignment based on

the V4 region of the RDP Classifier database might incorrectly identify *L. acidophilus* as

the predominant *Lactobacillus* species instead (114). To enhance the reliability of

species assignments, we implemented a two-step approach. From the ASV abundance

table, we selected a single species among those with identical V4 reference sequences

based on the highest abundant species driven from the RDP Classifier database.

Selected species are indicated with stars (**) in **Tables 1** and **2**. Remaining species and

their corresponding V4 reference sequences were removed from the reference

database. Using this reduced set of V4 region sequences as our reference database,

we repeated taxonomy assignment for ASVs. Despite random, sintax algorithm-driven

selection of a species from among multiple alternatives, we were able to track the

species chosen from among those with identical V4 sequences.

Statistical analyses

Alpha diversity. To investigate significant differences in alpha diversity among the

three groups (Endo+, Endo-, and CT-), we standardized the total read counts per

sample, which ranged from 34,571 to 417,336, by rarefying the reads across samples to

the minimum read counts (34,571 reads) using rarefy_even_depth function in phyloseq

R package. We assessed Shannon (richness) and Inverse Simpson (evenness) using

phyloseq and microbiome R packages. Phylogenetic diversity was calculated using

picante R package. The phylogenetic tree for phylogenetic diversity was generated from

the 16S rRNA V4 region sequences of swab samples by using MUSCLE (111).

Statistical significance in alpha diversity between each pair of groups was determined

using the Wilcoxon rank-sum test.

The centered log-ratio transformation. Microbiome-derived read count data obtained

through amplicon sequencing is analyzed as compositional data (115) where a

proportional change for any microorganism will affect relevant abundances of others,

rendering ASV read abundances mutually dependent. Centered log-ratio (CLR)

transformation is one approach that generalizes relative abundances with respect to the

geometric mean of all sequences in the sample (116). CLR-transformation of read

counts enables meaningful comparisons of microbial abundances by mitigating

compositional data biases (117). Since CLR-transformation depends on logarithms,

handling zero counts is challenging. We treated absolute zero counts by replacing them

with the smallest CLR-transformed values calculated by from our read counts.

Community state type (CST) classification of TRAC cohort and discrimination

analysis for CT infection status associated with CST. TRAC participants (n=220)

were categorized into CST types according to their vaginal microbial compositions using

VALENCIA (VAginaL community state typE Nearest CentroId clAssifier) (118).

VALENCIA classifies samples for CST types based on their similarity to a set of

reference centroids, and the VALENCIA tool and reference centroids were sourced from

GitHub (https://github.com/ravel-lab/VALENCIA). Where necessary, the taxon names

from RDP Classifier database were manually adapted to correspond with the

VALENCIA taxonomy of the reference centroids.

A sparse partial least squares discriminant analysis (sPLS-DA) (41) was

employed to discriminate between CT infection status across all women, and to

differentiate between CT infection status within CST I, CST III, and CST IV types. The

sPLS-DA was executed using splsda function in mixOmics R package (119). For

training across 213 women, 10-fold cross-validation (CV) with 1,000 iterations was used,

while within assigned CSTs, sPLS-DA utilized 5-fold CV repeated 1,000 times. To

assess the sPLS-DA model, AUC scores were computed from the training CV sets and

averaged over 1,000 iterations, yielding the mean and standard deviation (s.d.). The

ASVs listed in **Supplementary Table S3** and **Supplementary Table S4** were extracted

using the perf function in the mixOmics R package with a threshold of frequencies

greater than 0.80 for each component.

Microbiome Multivariate Association with Linear Models (MaAsLin2) for

differential abundance analysis. To investigate if CVM of TRAC cohort differ in

abundance between CT+ and CT- women, MaAsLin2 (42) was performed using

Maaslin2 R package with the embedded CLR transformation method (normalization =

"CLR", transform = "NONE", analysis_method = "LM", correction = "BH",

min_prevalence = 0).

Random forest classification. Random forest classification was used to determine if

CVM excluding CT is predictive of the lack of CT infection. This method was also

utilized to predict chlamydial ascending infection and to identify microbial biomarkers.

The random forest was performed using randomForest R package (120), and the

processes of parameter selection, and biomarker identification, are outlined in

Supplementary Figure 8. In the parameter selection, we selected the optimal

hyperparameters to improve prediction performance. For model evaluation and

biomarker identification, a random forest classification model was generated and

evaluated with K-fold CV, simultaneously choosing the microbial biomarkers of

chlamydial ascension.

The following parameters were examined: (1) number of folds for CV (K=5 or

K=10), (2) level of phylogeny used to aggregate ASVs into species or genus, (3)

significance level of T-tests and Wilcoxon rank-sum tests used for aggregated microbial

features (0.05 or 0.10), (4) whether to use the default setting of randomForest function

or a hyperparameter grid search. Tuning parameters in the randomForest function

included three options: mtry (number of trees randomly sampled as candidates at each

split), nodesize (minimum size of terminal nodes), and sampsize (sizes of sample to

grow). To accomplish this, we split our dataset into training and test datasets using K-

fold CV, ensuring that each fold included samples from both the Endo+ and Endo-

groups. With training data, we performed two-sample T-tests and Wilcoxon rank-sum

tests on ASVs to aggregate ASVs having p-value < 0.05 and the same direction in

abundances within the same genus or species between Endo+ and Endo- groups.

Specifically, when ASVs have a p-value less < 0.05, ASVs demonstrating the same

directional trend in abundances (i.e., either positive or negative t-statistics) within a

species or genus were aggregated into a single microbiome feature. ASVs with p-values

>0.05 but with consistent directional trends, were also aggregated into a single

microbiome feature. These aggregated microbiome features were tested again using T-

tests and Wilcoxon-tests to choose microbial features having p-values less than the

significance level (0.05 or 0.10). The aggregated microbiome features were

subsequently employed as features of training data and test data for random forest

classification. For every combination of the four parameters, AUC values were

calculated from 20 replicates of random forest, and parameters were selected based on

the highest sum of the mean and median AUC values across the 20 replicates. As a

result, the optimal parameters chosen included a significance level of 0.10, a 10-fold

cross-validation (CV), aggregation of read counts at the genus level for ASVs, and the

default settings of the randomForest R function. These settings were applied twice,

once including CT ASV as a microbiome feature, then excluding CT ASV.

With optimal parameters, the process re-implemented for model evaluation and

biomarker identification with 100 iterations. Prediction accuracy was assessed by

averaging the ROCs and AUCs obtained from the test sets over the 100 iterations.

Optimal thresholds for specificity and sensitivity were obtained by pROC R package

(121). For biomarker identification, ASVs were ranked based on their mean decrease in

accuracy, which indicates the importance of each feature in classifying samples. The

higher the mean decrease in accuracy, the more important the variable is in the model.

Using this information, ASVs were identified as biomarkers predicting chlamydial

ascension.

For validation of the 12 informative ASVs indicating risk for chlamydial ascension,

we performed random forest prediction using the same approach by which the 12

informative ASVs were identified (**Table 1**). A total of 143 samples (both Endo+ and

Endo-) were randomly split into training and test datasets using 10-fold cross-validation

(CV). The test dataset, including the four samples, was predicted using the CLR-

transformed abundance of the 12 ASVs. This entire process was repeated 100 times to

ensure consistency in the predictions across different subsets of the 143 samples used

as the training dataset. We then evaluated the predictions by calculating the mean,

standard deviation, and standard error of Endo+ probabilities for the seven samples, as

well as for the Endo+ and Endo- samples.

Canonical Correlation Analysis. Canonical correlation analysis (CCA) focuses on

finding linear combinations maximizing the correlation between two sets of multiple

variables. To assess the association between cervical cytokine levels and microbial

features, CCA was implemented by CCA R package (45).

Bibliography

1.   Cheong HC, Lee CYQ, Cheok YY, Tan GMY, Looi CY, Wong WF.

Chlamydiaceae: Diseases in Primary Hosts and Zoonosis. Microorganisms. 2019;7(5).

2.   O'Connell CM, Ferone ME. Chlamydia trachomatis Genital Infections. Microb Cell.

2016;3(9):390-403.

3.   Silva J, Cerqueira F, Medeiros R. Chlamydia trachomatis infection: implications

for HPV status and cervical cancer. Arch Gynecol Obstet. 2014;289(4):715-23.

4.   Paavonen J. Chlamydia trachomatis infections of the female genital tract: state of

the art. Ann Med. 2012;44(1):18-28.

5.   Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, et al. Vaginal

microbiome of reproductive-age women. Proc Natl Acad Sci U S A. 2011;108 Suppl

1(Suppl 1):4680-7.

6.   Edwards VL, Smith SB, McComb EJ, Tamarelle J, Ma B, Humphrys MS, et al.

The Cervicovaginal Microbiota-Host Interaction Modulates Chlamydia trachomatis

Infection. mBio. 2019;10(4).

7.   Borgdorff H, Tsivtsivadze E, Verhelst R, Marzorati M, Jurriaans S, Ndayisaba GF,

et al. Lactobacillus-dominated cervicovaginal microbiota associated with reduced

HIV/STI prevalence and genital HIV viral load in African women. ISME J.

2014;8(9):1781-93.

8.   van Houdt R, Ma B, Bruisten SM, Speksnijder A, Ravel J, de Vries HJC.

Lactobacillus iners-dominated vaginal microbiota is associated with increased

susceptibility to Chlamydia trachomatis infection in Dutch women: a case-control study.

Sex Transm Infect. 2018;94(2):117-23.

9.   Tsai HW, Tsui KH, Chiu YC, Wang LC. Adverse effect of lactobacilli-depauperate

cervicovaginal microbiota on pregnancy outcomes in women undergoing frozen-thawed

embryo transfer. Reprod Med Biol. 2023;22(1):e12495.

10.   Tabatabaei N, Eren AM, Barreiro LB, Yotova V, Dumaine A, Allard C, et al.

Vaginal microbiome in early pregnancy and subsequent risk of spontaneous preterm

birth: a case-control study. BJOG. 2019;126(3):349-58.

11.   Gosmann C, Handley SA, Farcasanu M, Abu-Ali G, Bowman BA, Padavattan N,

et al. Lactobacillus-deficient cervicovaginal bacterial communities are associated with

increased HIV acquisition in young South African women. Immunity. 2017;46(1):29-37.

12.   Anahtar MN, Byrne EH, Doherty KE, Bowman BA, Yamamoto HS, Soumillon M,

et al. Cervicovaginal bacteria are a major modulator of host inflammatory responses in

the female genital tract. Immunity. 2015;42(5):965-76.

13.   van der Veer C, Bruisten SM, van der Helm JJ, de Vries HJ, van Houdt R. The

Cervicovaginal Microbiota in Women Notified for Chlamydia trachomatis Infection: A

Case-Control Study at the Sexually Transmitted Infection Outpatient Clinic in

Amsterdam, The Netherlands. Clin Infect Dis. 2017;64(1):24-31.

14.   Ziklo N, Huston WM, Taing K, Katouli M, Timms P. In vitro rescue of genital

strains of Chlamydia trachomatis from interferon-gamma and tryptophan depletion with

indole-positive, but not indole-negative Prevotella spp. BMC Microbiol. 2016;16(1):286.

15. Beatty WL, Belanger TA, Desai AA, Morrison RP, Byrne GI. Tryptophan

depletion as a mechanism of gamma interferon-mediated chlamydial persistence.

Infection and immunity. 1994;62(9):3705-11.

16. Ziklo N, Vidgen ME, Taing K, Huston WM, Timms P. Dysbiosis of the Vaginal

Microbiota and Higher Vaginal Kynurenine/Tryptophan Ratio Reveals an Association

with Chlamydia trachomatis Genital Infections. Front Cell Infect Microbiol. 2018;8:1.

17. Jordan SJ, Olson KM, Barnes S, Wilson LS, Berryhill TF, Bakshi R, et al. Lower

Levels of Cervicovaginal Tryptophan Are Associated With Natural Clearance of

Chlamydia in Women. J Infect Dis. 2017;215(12):1888-92.

18. Ziklo N, Huston WM, Hocking JS, Timms P. Chlamydia trachomatis Genital Tract

Infections: When Host Immune Response and the Microbiome Collide. Trends Microbiol.

2016;24(9):750-65.

19. Aiyar A, Quayle AJ, Buckner LR, Sherchand SP, Chang TL, Zea AH, et al.

Influence of the tryptophan-indole-IFNgamma axis on human genital Chlamydia

trachomatis infection: role of vaginal co-infections. Front Cell Infect Microbiol. 2014;4:72.

20. Yeoman CJ, Thomas SM, Miller ME, Ulanov AV, Torralba M, Lucas S, et al. A

multi-omic systems-based approach reveals metabolic markers of bacterial vaginosis

and insight into the disease. PLoS One. 2013;8(2):e56111.

21. Ness RB, Kip KE, Hillier SL, Soper DE, Stamm CA, Sweet RL, et al. A cluster

analysis of bacterial vaginosis-associated microflora and pelvic inflammatory disease.

Am J Epidemiol. 2005;162(6):585-90.

22. Parolin C, Foschi C, Laghi L, Zhu C, Banzola N, Gaspari V, et al. Insights Into

Vaginal Bacterial Communities and Metabolic Profiles of Chlamydia trachomatis

Infection: Positioning Between Eubiosis and Dysbiosis. Front Microbiol. 2018;9:600.

23. Gong Z, Luna Y, Yu P, Fan H. Lactobacilli inactivate Chlamydia trachomatis

through lactic acid but not H2O2. PLoS One. 2014;9(9):e107758.

24. Pulendran B, Maddur MS. Innate Immune Sensing and Response to Influenza.

Curr Top Microbiol Immunol. 2014.

25. Zheng X, O'Connell CM, Zhong W, Nagarajan UM, Tripathy M, Lee DA, et al.

Discovery of blood transcriptional endotypes in women with pelvic inflammatory disease.

The Journal of Immunology. 2018;200(8):2941-56.

26. Filardo S, Di Pietro M, Porpora MG, Recine N, Farcomeni A, Latino MA, et al.

Diversity of Cervical Microbiota in Asymptomatic Chlamydia trachomatis Genital

Infection: A Pilot Study. Front Cell Infect Microbiol. 2017;7:321.

27. Russell AN, Zheng X, O'Connell CM, Taylor BD, Wiesenfeld HC, Hillier SL, et al.

Analysis of Factors Driving Incident and Ascending Infection and the Role of Serum

Antibody in Chlamydia trachomatis Genital Tract Infection. J Infect Dis.

2016;213(4):523-31.

28. Russell AN, Zheng X, O'Connell CM, Wiesenfeld HC, Hillier SL, Taylor BD, et al.

Identification of Chlamydia trachomatis Antigens Recognized by T Cells From Highly

Exposed Women Who Limit or Resist Genital Tract Infection. J Infect Dis.

2016;214(12):1884-92.

29.   Jeong S, Tollison T, Brochu H, Chou H, Yu T, Baghaie P, et al. Aptima Combo2-

avoiding variants detected in cervical and endometrial specimens from a cohort of

sexually active cisgender women during 16S microbiome profiling. medRxiv. 2024.

30.   Taylor BD, Zheng X, O'Connell CM, Wiesenfeld HC, Hillier SL, Darville T. Risk

factors for Mycoplasma genitalium endometritis and incident infection: a secondary data

analysis of the T cell Response Against Chlamydia (TRAC) Study. Sex Transm Infect.

2018;94(6):411-3.

31.   Zhou X, Brown CJ, Abdo Z, Davis C, Hansmann MA, Joyce P, et al. Disparity in

the vaginal microbial community composition of healthy Caucasian and black woman.

ISME J. 2007;1:121-33.

32.   Fettweis JM, Brooks JP, Serrano MG, Sheth NU, Girerd PH, Edwards DJ, et al.

Differences in vaginal microbiome in African American women versus women of

European ancestry. Microbiology (Reading). 2014;160(Pt 10):2272-82.

33.   Vodstrcil LA, Twin J, Garland SM, Fairley CK, Hocking JS, Law MG, et al. The

influence of sexual activity on the vaginal microbiota and Gardnerella vaginalis clade

diversity in young women. PLoS One. 2017;12(2):e0171856.

34.   Borgdorff H, van der Veer C, van Houdt R, Alberts CJ, de Vries HJ, Bruisten SM,

et al. The association between ethnicity and vaginal microbiota composition in

Amsterdam, the Netherlands. PLoS One. 2017;12(7):e0181135.

35.   Brotman RM, He X, Gajer P, Fadrosh D, Sharma E, Mongodin EF, et al.

Association between cigarette smoking and the vaginal microbiota: a pilot study. BMC

Infectious diseases. 2014;14(471):1-11.

36.   Ceccarani C, Foschi C, Parolin C, D'Antuono A, Gaspari V, Consolandi C, et al.
Diversity of vaginal microbiome and metabolome during genital infections. Sci Rep.
2019;9(1):14095.

37.   Chiu SF, Huang PJ, Cheng WH, Huang CY, Chu LJ, Lee CC, et al. Vaginal
Microbiota of the Sexually Transmitted Infections Caused by Chlamydia trachomatis
and Trichomonas vaginalis in Women with Vaginitis in Taiwan. Microorganisms.
2021;9(9).

38.   Cheong HC, Yap PSX, Chong CW, Cheok YY, Lee CYQ, Tan GMY, et al.
Diversity of endocervical microbiota associated with genital Chlamydia trachomatis
infection and infertility among women visiting obstetrics and gynecology clinics in
Malaysia. PLoS One. 2019;14(11):e0224658.

39.   Di Pietro M, Filardo S, Porpora MG, Recine N, Latino MA, Sessa R.
HPV/Chlamydia trachomatis co-infection: metagenomic analysis of cervical microbiota
in asymptomatic women. New Microbiologica. 2018;41(1):34-41.

40.   Borgogna JC, Shardell MD, Yeoman CJ, Ghanem KG, Kadriu H, Ulanov AV, et al.
The association of Chlamydia trachomatis and Mycoplasma genitalium infection with the
vaginal metabolome. Sci Rep. 2020;10(1):3420.

41.   Cao K-AL, Boitard S, Besse P. Sparse PLS discriminant analysis: biologically
relevant feature selection and graphical displays for multiclass problems. BMC
Bioinformatics. 2011;12:1-17.

42.   Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, et al.
Multivariable association discovery in population-scale meta-omics studies. PLoS
Comput Biol. 2021;17(11):e1009442.

43.   Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. Appl Environ Microbiol.

2007;73(16):5261-7.

44.   Poston TB, Lee DE, Darville T, Zhong W, Dong L, O'Connell CM, et al. Cervical

Cytokines Associated With Chlamydia trachomatis Susceptibility and Protection. J Infect

Dis. 2019;220(2):330-9.

45.   González, Déjean, Martin, Baccini. CCA: An R package to extend canonical

correlation analysis. Journal of Statistical Software. 2008;23(12):1-14.

46.   Tamarelle J, Thiébaut AC, De Barbeyrac B, Bebear C, Ravel J, Delarocque-

Astagneau E. The vaginal microbiota and its association with human papillomavirus,

Chlamydia trachomatis, Neisseria gonorrhoeae and Mycoplasma genitalium infections:

a   systematic   review   and   meta-analysis.   Clinical   Microbiology   and   Infection.

2019;25(1):35-47.

47.   Filardo S, Di Pietro M, Tranquilli G, Latino MA, Recine N, Porpora MG, et al.

Selected Immunological Mediators and Cervical Microbial Signatures in Women with

Chlamydia trachomatis Infection. mSystems. 2019;4(4).

48.   Sasaki-Imamura T, Yoshida Y, Suwabe K, Yoshimura F, Kato H. Molecular basis

of indole production catalyzed by tryptophanase in the genus Prevotella. FEMS

Microbiology Letters. 2011;322(1):51-9.

49.   Buhl M, Dunlap C, Marschal M. Prevotella brunnea sp. nov., isolated from a

wound of a patient. International Journal of Systematic and Evolutionary Microbiology.

2019;69(12):3933-8.

50.   Horvath A, Durdevic M, Leber B, di Vora K, Rainer F, Krones E, et al. Changes in

the Intestinal Microbiome during a Multispecies Probiotic Intervention in Compensated

Cirrhosis. Nutrients. 2020;12(6).

51.   Hinderfeld AS, Phukan N, Bär A-K, Roberton AM, Simoes-Barbosa A.

Cooperative Interactions between Trichomonas vaginalis and Associated Bacteria

Enhance Paracellular Permeability of the Cervicovaginal Epithelium by Dysregulating

Tight Junctions. Infection and immunity. 2019;87(5).

52.   Goldstein EJ, Tyrrell KL, Citron DM. Lactobacillus species: taxonomic complexity

and controversial susceptibilities. Clin Infect Dis. 2015;60 Suppl 2:S98-107.

53.   O'Callaghan JL, Willner D, Buttini M, Huygens F, Pelzer ES. Limitations of 16S

rRNA Gene Sequencing to Characterize Lactobacillus Species in the Upper Genital

Tract. Frontiers in Cell and Developmental Biology. 2021;9:641921.

54.   Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace

operational taxonomic units in marker-gene data analysis. ISME J. 2017;11(12):2639-43.

55.   Johnson JS, Spakowicz DJ, Hong BY, Petersen LM, Demkowicz P, Chen L, et al.

Evaluation of 16S rRNA gene sequencing for species and strain-level microbiome

analysis. Nat Commun. 2019;10(1):5029.

56.   O'Flaherty S, Briner Crawley A, Theriot CM, Barrangou R. The Lactobacillus Bile

Salt Hydrolase Repertoire Reveals Niche-Specific Adaptation. mSphere. 2018;3(3).

57.   Navarro S, Abla H, Delgado B, Colmer-Hamood JA, Ventolini G, Hamood AN.

Glycogen availability and pH variation in a medium simulating vaginal fluid influence the

growth of vaginal Lactobacillus species and Gardnerella vaginalis. BMC Microbiol.

2023;23(1):186.

58.   Hertzberger R, May A, Kramer G, van Vondelen I, Molenaar D, Kort R. Genetic

Elements Orchestrating Lactobacillus crispatus Glycogen Metabolism in the Vagina. Int

J Mol Sci. 2022;23(10).

59.   Putonti C, Shapiro JW, Ene A, Tsibere O, Wolfe AJ. Comparative genomic study

of Lactobacillus jensenii and the newly defined Lactobacillus mulieris species identifies

species-specific functionality. mSphere. 2020;5(4):00560-20.

60.   Hood MI, Skaar EP. Nutritional immunity: transition metals at the pathogen-host

interface. Nat Rev Microbiol. 2012;10(8):525-37.

61.   Raulston JE. Response of Chlamydia trachomatis serovar E to iron restriction in

vitro and evidence for iron-regulated chlamydial proteins. Infection and Immunity.

1997;65(11):4539-47.

62.   Pokorzynski ND, Thompson CC, Carabeo RA. Ironing Out the Unconventional

Mechanisms of Iron Acquisition and Gene Regulation in Chlamydia. Frontiers in Cellular

and Infection Microbiology. 2017;7:394.

63.   Pokorzynski ND, Alla MR, Carabeo RA. Host cell amplification of nutritional

stress contributes to persistence in Chlamydia trachomatis. mBio. 2022;13(6):e02719-

22.

64.   Pokorzynski ND, Brinkworth AJ, Carabeo R. A bipartite iron-dependent

transcriptional regulation of the tryptophan salvage pathway in Chlamydia trachomatis.

Elife. 2019;8.

65.   Pickering JL, Prosser A, Corscadden KJ, De Gier C, Richmond PC, Zhang G, et

al. Haemophilus haemolyticus Interaction with Host Cells Is Different to Nontypeable

Haemophilus influenzae and Prevents NTHi Association with Epithelial Cells. Frontiers in cellular and infection microbiology. 2016;6:50.

66.   Fulte S, Atto B, McCarty A, Horn KJ, Redzic JS, Eisenmesser E, et al. Heme sequestration by hemophilin from Haemophilus haemolyticus reduces respiratory tract colonization and infection with non-typeable Haemophilus influenzae. mSphere. 2024;9(3):e00006-24.

67.   Latham RD, Torrado M, Atto B, Walshe JL, Wilson R, Guss JM, et al. A heme-binding protein produced by Haemophilus haemolyticus inhibits non-typeable Haemophilus influenzae. Mol Microbiol. 2020;113(2):381-98.

68.   Woodall CA, Hammond A, Cleary D, Preston A, Muir P, Pascoe B, et al. Oral and gut microbial biomarkers of susceptibility to respiratory tract infection in adults: A feasibility study. Heliyon. 2023;9(8):e18610.

69.   Hiippala K, Kainulainen V, Kalliomaki M, Arkkila P, Satokari R. Mucosal Prevalence and Interactions with the Epithelium Indicate Commensalism of Sutterella spp. Front Microbiol. 2016;7:1706.

70.   Ceccarani C, Marangoni A, Severgnini M, Camboni T, Laghi L, Gaspari V, et al. Rectal Microbiota Associated With Chlamydia trachomatis and Neisseria gonorrhoeae Infections in Men Having Sex With Other Men. Front Cell Infect Microbiol. 2019;9:358.

71.   Atarashi K, Tanoue T, Ando M, Kamada N, Nagano Y, Narushima S, et al. Th17 Cell Induction by Adhesion of Microbes to Intestinal Epithelial Cells. Cell. 2015;163(2):367-80.

72.   Gaboriau-Routhiau V, Rakotobe S, Lecuyer E, Mulder I, Lan A, Bridonneau C, et

al. The key role of segmented filamentous bacteria in the coordinated maturation of gut

helper T cell responses. Immunity. 2009;31(4):677-89.

73.   Yount KS, Kollipara A, Liu C, Zheng X, O'Connell CM, Bagwell B, et al. Unique T

cell signatures are associated with reduced Chlamydia trachomatis reinfection in a

highly exposed cohort. bioRxiv. 2023.

74.   Fettweis JM, Serrano MG, Huang B, Brooks JP, Glascock AL, Sheth NU, et al.

An emerging mycoplasma associated with trichomoniasis, vaginal infection and disease.

PLoS One. 2014;9(10):e110943.

75.   Ioannidis A, Papaioannou P, Magiorkinis E, Magana M, Ioannidou V, Tzanetou K,

et al. Detecting the Diversity of Mycoplasma and Ureaplasma Endosymbionts Hosted by

Trichomonas vaginalis Isolates. Front Microbiol. 2017;8:1188.

76.   Ma C, Du J, Dou Y, Chen R, Li Y, Zhao L, et al. The Associations of Genital

Mycoplasmas with Female Infertility and Adverse Pregnancy Outcomes: a Systematic

Review and Meta-analysis. Reprod Sci. 2021;28(11):3013-31.

77.   Jonduo ME, Vallely LM, Wand H, Sweeney EL, Egli-Gany D, Kaldor J, et al.

Adverse pregnancy and birth outcomes associated with Mycoplasma hominis,

Ureaplasma urealyticum and Ureaplasma parvum: a systematic review and meta-

analysis. BMJ Open. 2022;12(8):e062990.

78.   Margarita V, Rappelli P, Dessi D, Pintus G, Hirt RP, Fiori PL. Symbiotic

association with Mycoplasma hominis can influence growth rate, ATP production,

cytolysis and inflammatory response of Trichomonas vaginalis. Frontiers in Microbiology.

2016;7:953.

79.   Margarita V, Bailey NP, Rappelli P, Diaz N, Dessì D, Fettweis JM, et al. Two

Different Species of Mycoplasma Endosymbionts Can Influence Trichomonas vaginalis

Pathophysiology. mBio. 2022;13(3):e00918-22.

80.   Fiori PL, Diaz N, Cocco AR, Rappelli P, Dessi D. Association of Trichomonas

vaginalis with its symbiont Mycoplasma hominis synergistically upregulates the in vitro

proinflammatory response of human monocytes. Sex Transm Infect. 2013;89(6):449-54.

81.   Greub G, Raoult D. "Actinobaculum massiliae," a new species causing chronic

urinary tract infection. J Clin Microbiol. 2002;40(11):3938-41.

82.   Carrillo-Avila JA, Bonilla-Garcia L, Navarro-Mari JM, Gutierrez-Fernandez J. The

first reported case of pelvic inflammatory disease caused by Actinobaculum massiliense.

Anaerobe. 2019;55:93-5.

83.   Yu Y, Tsitrin T, Singh H, Doerfert SN, Sizova MV, Epstein SS, et al.

Actinobaculum massiliense Proteome Profiled in Polymicrobial Urethral Catheter

Biofilms. Proteomes. 2018;6(4).

84.   Reynoso-García J, Miranda-Santiago AE, Meléndez-Vázquez NM, Acosta-Pagán

K, Sánchez-Rosado M, Díaz-Rivera J, et al. A complete guide to human microbiomes:

Body niches, transmission, development, dysbiosis, and restoration. Frontiers in

Systems Biology. 2022;2:951403.

85.   Lamas A, Regal P, Vázquez B, Cepeda A, Franco CM. Short chain fatty acids

commonly produced by gut microbiota influence Salmonella enterica motility, biofilm

formation, and gene expression. Antibiotics. 2019;8(4):265.

86. Ozma MA, Abbasi A, Akrami S, Lahouty M, Shahbazi N, Ganbarov K, et al.

Postbiotics as the key mediators of the gut microbiota-host interactions. Infez Med.

2022;30(2):180-93.

87. Mirzaei R, Kavyani B, Nabizadeh E, Kadkhoda H, Asghari Ozma M, Abdi M.

Microbiota metabolites in the female reproductive system: Focused on the short-chain

fatty acids. Heliyon. 2023;9(3):e14562.

88. Wiesenfeld HC, Sweet RL, Ness RB, Krohn MA, Amortegui AJ, Hillier SL.

Comparison of Acute and Subclinical Pelvic Inflammatory Disease. Sexually transmitted

diseases. 2005;32(7):400-5.

89. Wiesenfeld HC, Hillier SL, Meyn LA, Amortegui AJ, Sweet RL. Subclinical pelvic

inflammatory disease and infertility. Obstet Gynecol. 2012;120(1):37-43.

90. Dodet B. Current barriers, challenges and opportunities for the development of

effective STI vaccines: point of view of vaccine producers, biotech companies and

funding agencies. Vaccine. 2014;32(14):1624-9.

91. Broutet N, Fruth U, Deal C, Gottlieb SL, Rees H. Vaccines against sexually

transmitted infections: the way forward. Vaccine. 2014;32(14):1630-7.

92. Poston TB, Gottlieb SL, Darville T. Status of vaccine research and development

of vaccines for Chlamydia trachomatis infection. Vaccine. 2019;37(50):7289-94.

93. Workowski KA, Bolan GA, Prevention CfDCa. Sexually transmitted diseases

treatment guidelines, 2015. MMWR Recomm Rep. 2015;64(3).

94. Zheng X, O'Connell CM, Zhong W, Poston TB, Wiesenfeld HC, Hillier SL, et al.

Gene Expression Signatures Can Aid Diagnosis of Sexually Transmitted Infection-

Induced Endometritis in Women. Front Cell Infect Microbiol. 2018;8:307.

95.   O'Connell CM, Brochu H, Girardi J, Harrell E, Jones A, Darville T, et al.

Simultaneous profiling of sexually transmitted bacterial pathogens, microbiome, and

concordant host response in cervical samples using whole transcriptome sequencing

analysis. Microb Cell. 2019;6(3):177-83.

96.   Abraham S, Juel HB, Bang P, Cheeseman HM, Dohn RB, Cole T, et al. Safety

and immunogenicity of the chlamydia vaccine candidate CTH522 adjuvanted with

CAF01 liposomes or aluminium hydroxide: a first-in-human, randomised, double-blind,

placebo-controlled, phase 1 trial. The Lancet infectious diseases. 2019;19(10):1091-100.

97.   Pan P, Gu Y, Sun D-L, Wu QL, Zhou N-Y. Microbial diversity biased estimation

caused by intragenomic heterogeneity and interspecific conservation of 16s rrna genes.

Applied and Environmental Microbiology. 2023;89(5):15.

98.   Kembel SW, Wu M, Eisen JA, Green JL. Incorporating 16S gene copy number

information improves estimates of microbial diversity and abundance. PLoS Comput

Biol. 2012;8(10):e1002743.

99.   Klappenbach JA, Saxman PR, Cole JR, Schmidt TM. rrndb: the Ribosomal RNA

Operon Copy Number Database. Nucleic Acids Research. 2001;29(1):181–4.

100.   Douglas GM, Maffei VJ, Zaneveld JR, Yurgel SN, Brown JR, Taylor CM, et al.

PICRUSt2 for prediction of metagenome functions. Nat Biotechnol. 2020;38:685–8.

101.   Miao J, Chen T, Misir M, Lin Y. Deep learning for predicting 16S rRNA gene copy

number. Sci Rep. 2024;14(1):14282.

102.   Angly FE, Dennis PG, Skarshewski A, Vanwonterghem I, Hugenholtz P, Tyson

GW. CopyRighter: a rapid tool for improving the accuracy of microbial community

profiles through lineage-specific gene copy number correction. Microbiome. 2014;2(11).

103. Stoddard SF, Smith BJ, Hein R, Roller BR, Schmidt TM. rrnDB: improved tools

for interpreting rRNA gene abundance in bacteria and archaea and a new foundation for

future development. Nucleic Acids Res. 2015;43(Database issue):D593-8.

104. Liu C, Hufnagel K, O'Connell CM, Goonetilleke N, Mokashi N, Waterboer T, et al.

Reduced Endometrial Ascension and Enhanced Reinfection Associated With

Immunoglobulin G Antibodies to Specific Chlamydia trachomatis Proteins in Women at

Risk for Chlamydia. J Infect Dis. 2022;225(5):846-55.

105. Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb

Ecol. 2015;75:129-37.

106. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. Environ Microbiol. 2016;18(5):1403-14.

107. Edgar RC. UNOISE2: improved error-correction for Illumina 16S and ITS

amplicon sequencing. BioRxiv. 2016:081257.

108. Zhang J, Kobert K, Flouri T, Stamatakis A. PEAR: a fast and accurate Illumina

Paired-End reAd mergeR. Bioinformatics. 2014;30(5):614-20.

109. Antich A, Palacin C, Wangensteen OS, Turon X. To denoise or to cluster, that is

not the question: optimizing pipelines for COI metabarcoding and metaphylogeography.

BMC Bioinformatics. 2021;22(1):177.

110. Li H. Aligning sequence reads, clone sequences and assembly contigs with

BWA-MEM. arXiv. 2013.

111.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic acids research. 2004;32(5):1792–7.

112.   Okonechnikov K, Golosova O, Fursov M. Unipro UGENE: a unified bioinformatics

toolkit. Bioinformatics. 2012;28(8):1166-7.

113.   Edgar RC. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS

sequences. biorxiv. 2016.

114.   Antonio MAD, Hawes SE, Hillier SL. The Identification of Vaginal Lactobacillus

Species and the Demographic and Microbiologic Characteristics of Women Colonized

by These Species. Journal of Infectious Diseases. 1999;180(6):The Journal of

Infectious Diseases.

115.   Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome Datasets

Are Compositional: And This Is Not Optional. Front Microbiol. 2017;8:2224.

116.   Aitchison J. The statistical analysis of compositional data. Royal Statistical

Society. 1982;44(2):139-77.

117.   Quinn TP, Erb I, Gloor G, Notredame C, Richardson MF, Crowley TM. A field

guide for the compositional analysis of any-omics data. Gigascience. 2019;8(9).

118.   France MT, Ma B, Gajer P, Brown S, Humphrys MS, Holm JB, et al. VALENCIA:

a nearest centroid classification method for vaginal microbial communities based on

composition. Microbiome. 2020;8(1):166.

119.   Rohart F, Gautier B, Singh A, Le Cao KA. mixOmics: An R package for 'omics

feature   selection   and   multiple   data   integration.   PLoS   Comput   Biol.

2017;13(11):e1005752.

120.   Liaw A, Wiener M. Classification and Regression by randomForest. R News.

2002;2(3):18-22.

121.   Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez J-C, et al. pROC: an

open-source package for R and S+ to analyze and compare ROC curves. BMC

Bioinformatics. 2011;12:77.

1225

1226

TABLES

Table 1. ASVs predictive of ascended CT infection. The ASVs selected for predicting susceptibility to ascending CT infection were determined based on their selection frequency threshold derived from random forest predictions. The selection frequency indicates the number of times an ASV was detected with non-NA values of mean decrease in accuracy, occurring in 9 or 10 out of 10 folds, out of 100 replications. From the RF classification results, 12 ASVs were identified as microbial biomarkers because their selection frequencies exceeded 80. Notably, despite considering two different scenarios (A: prediction with all ASVs including CT ASV and B: prediction with all ASVs but excluding CT ASV), 11 out of 12 ASVs were consistently identified across both scenarios and indicated by "Shared" in Prediction scenario column of **Table 1**. The relative abundance of these ASVs was determined using t statistics, indicating whether an ASV was higher in Endo+ or lower in Endo+. Total read count was calculated for all 220 women, and details of the %ID and Confidence level columns were provided in **Table 1**.

Relative abundan-ce	Pred. Scena-rio	Species	ASV	Selection frequency	Total read count	%ID	Genus	Confidence level	
								Genus	Species
	w/ CT	*Chlamydia_trachomatis*	ZOtu76	100	15,683	100	*Chlamydia*	1	0.96
Higher in Endo+	**Shared**	*Acetothermia_genera_incertae_sedis_clone*	ZOtu233	100	312	79	*Acetothermia_genera_incertae_sedis*	0.0064	0.0005
		Butyrivibrio_hungatei	ZOtu1808	98	325	89.4	*Butyrivibrio*	0.2048	0.0266
		Actinobaculum_massiliense	ZOtu567	100	63	100	*Actinobaculum*	0.89	0.6141
Lower in Endo+	**Shared**	*Actinotignum_schaalii***	ZOtu281	100	200	100	*Actinotignum*	0.96	0.768
		Haemophilus_haemolyticus	ZOtu103	100	6,697	100	*Haemophilus*	0.96	0.7584
		Candidatus Mycoplasma girerdii	ZOtu45▯	100	68,190	100	*Mycoplasma*		
		Porphyromonas_bennonis	ZOtu373	100	132	99.2	*Porphyromonas*	1	1
		Propionibacterium_lymphophilum	ZOtu333	100	142	100	*Propionibacterium*	1	1
		*Sutterella_stercoricanis***	ZOtu110	100	4,343	96.4	*Sutterella*	0.894	0.5453
		*Sutterella_stercoricanis***	ZOtu184	99	690	99.6	*Sutterella*	0.9801	0.9605
		*Sutterella_stercoricanis***	ZOtu180	87	756	97.6	*Sutterella*	0.8748	0.6998

** ASVs were assigned to a specific species from among different species that share

the same sequence in the V4 region of the 16S rRNA.

⍰ ZOtu45 ASV was initially assigned as *Malacoplasma microti* based on RDP Classifier

DB with 89% identity (Confidence level: 0.8209 at the genus level and 0.4351 at the

species level), but it was later identified as *Candidatus Mycoplasma girerdii* with 100%

identity.

Figure 1. Profiles of community state types from TRAC cohort associated with Chlamydial infection. A. The taxonomic composition of 220 women, incorporating CST types and pathogen infection status (CT = *Chlamydia trachomatis*, NG = *Neisseria gonorrhoeae* (NG), and MG = *Mycoplasma genitalium*) were presented on the heat map. The testing results for pathogenic infection status at enrollment (CT.Test, NG.Test, MG.Test) were provided as bar graphs on the top of the heatmap. The reassigned

results of seven false negatives for CT infection are labeled as CT.FN. The term

'Equivocal' indicates that the reassignment of these cases as either Endo+ or Endo-

was unclear, despite them being confirmed as CT+. CT ASV read counts were

converted into logarithmic scale (CT_ASV; log10(read count + 1)). The observations of

three pathogenic ASV sequences were illustrated with 'Existence of ASVs' at the bottom

of the heatmap (CT.ASV, NG.ASV, MG.ASV). **B.** The table shows the definitions of CST

types and the number of women belonging to each CST type.

Figure 2. Discrimination of chlamydial infection status at the cervix or endometrium in TRAC cohort using sPLS-DA. Discrimination of chlamydial infection status at cervix (A and B) and endometrium (C and D) was visualized using the first two components of sPLS-DA. The discriminations were performed using ASVs including CT ASV (A and C) or excluding CT ASV (B and D). Each component's contribution to the discrimination was indicated at their axis. The performance was evaluated using the mean Area Under the Curve (AUC) values. **A.** The first two sPLS-DA components discriminate CT+ from CT- women based on ASVs with CT ASV (AUC mean = 0.70, AUC s.d. = 0.02) **B.** The discrimination of CT+ from CT- women based on without CT ASV (AUC mean = 0.51, AUC s.d. = 0.03) **C.** The first two sPLS-DA components discriminate Endo+ from Endo- women (AUC mean = 0.76, AUC s.d. = 0.01) **D.** The discrimination of Endo+ from Endo- women based without CT ASV (AUC mean = 0.55, AUC s.d. = 0.03).

1277

Figure 3. ROC curves and corresponding AUCs from random forest prediction for susceptibility to ascending chlamydial infection. Prediction accuracies were evaluated by averaging AUCs over 100 replicates for each scenario. **A.** The prediction performance for the susceptibility to CT ascension using all ASVs, including CT ASV resulted in an AUC of 0.74. **B.** The prediction performance for excluding CT ASV resulted in an AUC of 0.57. **C.** The susceptibility of ascending CT infection, predicted using only CT ASV, led to an AUC of 0.74.

Figure 4. The Correlations between canonical variates of microbiome biomarkers and cervical cytokines. The canonical correlations between cervical cytokines related to chlamydial ascension and microbial biomarkers identified for chlamydial ascension risk were performed under two scenarios: (A) correlating the 7 cervical cytokine levels with 12 informative ASVs, including CT ASV, listed in **Table 1**, and (B) with 11 informative ASVs, excluding CT ASV. **A.** The canonical correlation between the seven cervical cytokine levels and the 12 ASVs, including CT ASV, was visualized using two CCA variates, resulting in a correlation coefficient of 0.57. **B.** The canonical correlation between the seven cervical cytokine levels and the 11 ASVs, excluding CT ASV, yielded a correlation coefficient of 0.44. **C.** Only CT ASV abundance was correlated with the seven cytokine levels, resulting in a correlation coefficient of 0.49.