Identification of a Highly Functional Effector CD8[+] T Cell Program after Transplantation in Mice

and Humans

Gregory S. Cohen,[1†] Joel S. Freibaum,[1†] Riley P. Leathem,[1†] Marina W. Shirkey,[2] Robin Welsh,[1]

Sanniddhya B. Bardhan,[1] Ryo Hatano,[3] Chikao Morimoto,[3] Byoung Chol Oh,[4] Daniela

Čiháková,[1,5] Jonathan S. Bromberg,[6] Scott M. Krummey[1]*

[1]Department of Pathology, Johns Hopkins School of Medicine, Baltimore, MD, 21205, USA

[2]University of Maryland School of Medicine, Department of Surgery, Center for Vascular and

Inflammatory Diseases, Baltimore, MD, 21201, USA

[3]Department of Therapy Development and Innovation for Immune Disorders and Cancers,

Graduate School of Medicine, Juntendo University, Tokyo 113-8421, Japan

[4]Department of Plastic and Reconstructive Surgery, Johns Hopkins School of Medicine,

Baltimore, MD, 21205, USA

[5]W. Harry Feinstone Department of Microbiology and Immunology, Bloomberg School of Public

Health, Johns Hopkins University, Baltimore, MD, 21205, USA

[6]University of Maryland School of Medicine, Departments of Surgery and Microbiology and

Immunology, Center for Vascular and Inflammatory Diseases, Baltimore, MD, 21201, USA

†These authors contributed equally to this work.

*Corresponding Author: Scott M. Krummey, Ross Research Building Room 664B, 720 Rutland

Avenue, Baltimore, MD 21205, 410-955-3600, skrumme1@jhmi.edu

Word count: 3,967

Keywords: effector CD8[+] T cells, alloimmunity, costimulation blockade, tolerance

ABSTRACT

T cell mediated allograft rejection leads to early graft loss for kidney transplant patients.

To better understand the mechanism by which T cells mediate rejection, we investigated the fate

and function of graft-specific CD8$^+$ T cells expressing the activated isoform of CD43 in mice and

humans. Agonism of CD43 1B11 in vitro induced CD8$^+$ T cell proliferation in the presence of sub-

threshold antigen stimulation, and CD43 1B11 mAb treatment in vivo overcame costimulation-

blockade induced tolerance to skin grafts. Relative to CD43 1B11$^-$ populations, CD43 1B11$^+$ CD8$^+$

T cells maintained high T-bet expression along with stem-like molecules IL-7Rα and TCF-1 at

both effector and memory timepoints, and were more persistent following adoptive transfer. In

kidney transplant patients, graft-infiltrating CD8$^+$ T cells that expressed CD43 and the

glycosyltransferase GCNT1 had an effector phenotype that includes high expression of *IFNG*,

ICOS, and perforins/granzymes. In healthy human donors and transplant candidates, the CD43

1D4 mAb clone defined antigen-experienced cytokine-producing CD8$^+$ T cells. In sum, these data

support a progressive differentiation model by which highly proliferative effector CD43 1B11$^+$

CD8$^+$ T cells infiltrate allografts also efficiently persist into memory after antigen clearance.

1. INTRODUCTION

T cell mediated allograft rejection is a major contributor to early graft loss.[1–3] Multiple

studies have highlighted that CD8[+] T cells, which have potent cytolytic and cytokine-producing

functions, mediate rejection in pre-clinical models and patients.[4–8] In the widely held early

precursor model of T cell differentiation, CD8[+] T cells become either short-lived effector cell

(SLEC) populations or longer-lived memory precursor populations in the initial cell divisions after

antigen priming. Following viral and bacterial infection, the transcription factor T-bet is essential

for CD8[+] T cell effector function of short-lived effectors, but also sustains memory cell

homeostasis.[10,16,17] In parallel, self-renewing stem-like features of CD8[+] T cell memory are defined

by responsiveness to IL-7 and the transcription factor T cell factor 1 (TCF-1).[18–20] While many

features of effector and memory CD8[+] T cells have been described, the allogeneic priming

environment differs significantly from that of pathogen infection.[11,13,21,22] The overarching model of

CD8[+] T cell differentiation in transplantation is not well understood.

CD43 is a transmembrane sialomucin which is broadly expressed on leukocytes, and has

complex functions in cellular adhesion and trafficking, as well as cellular signaling.[23–26] However,

T cell activation elicits the core-2 O-glycosylation of CD43 by the glycosyltransferase GCNT1, [27,28]

which is denoted by the binding of the CD43 1B11 antibody clone. Our group and others have

shown that effector and memory CD8[+] T cells upregulate expression of the activated CD43

receptor.[23,29,30] We found that after transplantation, CD43 1B11 CD8[+] T cells were potent effectors

that expressed highly levels of Granzyme B, the cytokines IFN-γ/TNF-α, and rapidly infiltrated

allografts.[29] In the present study, we further delineated key aspects of the fate and function of this

CD8[+] T cell population in mice and humans in order to better understand the population dynamics

driving T cell mediated rejection.

2. MATERIALS AND METHODS

Animals. C57BL/6J, C57BL/6J Ly5.2-Cr (CD45.1), C57Bl/6-Tg(TcraTcrb)1100Mjb/J (OT-I), and

C567BL/6-Tg(CAG-OVAL)916Jen/J (Act-mOVA) were obtained from The Jackson Laboratory

(Bar Harbor, Maine) and used at 6-12 weeks of age. Balb/cJ, were obtained from The Jackson

Laboratory and bred at Johns Hopkins University. This study was conducted in accordance with

the recommendations in the Guide for the Care and Use of Laboratory Animals, and was approved

by the Animal Care and Use Committee at Johns Hopkins University. Animals were housed in

specific pathogen-free animal facilities at Johns Hopkins University.

Skin transplantation and in vivo treatments. Full-thickness tail and ear skins were transplanted

onto the dorsal thorax of recipient mice and secured with adhesive bandages as previously

described [31]. In some experiments, mice were treated with 250 μg/dose of CTLA-4 Ig (BioXCell)

and anti-CD40L (MR1 clone, BioXCell) via i.p. injection as indicated. CD43 1B11 mAb (Biolegend)

was provided at 200 μg/dose via i.p. injection.

Heart transplantation and graft-infiltrating T cell isolation. Heterotopic heart allografts were

performed as previously described via anastomosis to the abdominal aorta and pulmonary

artery.[32] To isolate graft-infiltrating cells, heart grafts were perfused with 1x PBS for 5 min, then

minced and digested using Collagenase II (5000 U/mL) and DNase I (500 U/mL) (Worthington

Biochemicals) in HBSS (Corning) for 30 min at 37 C with gentile agitation. Tissue was then

dissociated using a GentleMACS Tissue Dissociator (Miltenyi).

Flow cytometric cell sorting and adoptive transfer. To obtain effector CD8[+] T cells,

splenocytes and graft draining lymph nodes (inguinal, brachial, axial) from CD45.1[+] mice grafted

with Balb/c skin grafts 10-14 days prior were collected and processed to single cell suspension.

For proliferation analysis, 2×10^6 CD8$^+$ T cells were isolated using CD8$^+$ T cell negative selection

kit (Miltenyi), and stained with CellTrace Violet (Invitrogen) prior to adoptive transfer. For fate

experiments, effector CD8$^+$ T cells were flow cytometrically sorted into CD44^{hi}CD62L^{lo}CD8$^+$ T cells

that were CD43 1B11$^+$ and CD43 1B11$^-$ using a MoFlo XDP instrument. $0.9\text{-}1.2\times10^5$ purified cells

were transferred via tail vein into C57BL/6J mice. Mice were grafted the following day with Balb/c

skin and analyzed by flow cytometry on the indicated day post-graft as described below.

In vitro T cell stimulation. 96-well flat-bottom tissue culture plates were coated with CD3ε, CD28,

or CD43 1B11 mAb (Biolegend) at the indicated concentration in 60 μL of PBS/well for 90 min at

37 C. Naïve CD8$^+$ T cells isolated using CD8$^+$ T Cell Isolation Kit with LS columns (Miltenyi) were

stained with CellTrace Violet (Invitrogen). 2×10^5 CD8$^+$ T cells were plated per well and incubated

at 37 C for 4 days in Complete R10 media comprised of RPMI 1640 with L-glutamine (Corning),

10% FBS, 100 mM HEPES, 500 uM β-mercaptoethanol, 100 U/mL penicillin/stremptomycin and

analyzed by flow cytometry as described below.

Immunofluorescence microscopy. Draining lymph nodes (dLN) were flash frozen in OCT

compound (Sakura Finetek, Torrance, CA). Six µm sections were cut in triplicate using a Microm

HM 550 cryostat (Thermo Fisher Scientific), fixed in cold 1:1 acetone:methanol for 5 minutes, and

washed in PBS. Slides were incubated in primary antibodies and secondary antibodies diluted in

PBST (PBS + 0.03% Triton-X-100 + 0.5% BSA) for 1 hour each. Slides were imaged using an

Accu-Scope EXC-500 fluorescent microscope (Nikon, Tokyo, Japan). Images were analyzed with

Volocity software (PerkinElmer, Waltham, MA). Mean of mean fluorescence intensity (MFI) was

calculated within demarcated high endothelial venule (HEV) region. Groups were compared using

quantitation of MFI multiplied by percent area (sum of HEV area with fluorescence over total HEV

area x MFI).

Murine T cell enrichment and flow cytometry. Flow cytometry and magnetic bead enrichment

were performed in 1X PBS (without Ca^{2+}/Mg^{2+}, pH 7.2), 0.25% BSA, 2 mM EDTA, and 0.09%

azide. $CD45.1^+$ or L^d $QL9^+$ $CD8^+$ T cells were enriched using magnetic beads specific for

fluorophores with LS columns (Miltenyi). MHC tetramers with human β2-microglobulin specific for

QL9 (H-2L^d QLSPFPFDL) were obtained from the NIH Tetramer Core facility. Single cell

suspensions were stained with Live/Dead NIR Zombie (Biolegend) and incubated with the

indicated antibodies for 30-60 min at room temperature. Intracellular staining was performed using

the Transcription Factor Staining Kit (eBiosciences). Samples were acquired on a Cytek Aurora

(4L or 5L configuration).

Human peripheral blood cell analysis. Healthy donor peripheral blood was collected in

Vacutainer tubes containing sodium heparin (BD Biosciences) and processed according to

manufacturer's instructions to isolate mononuclear cells. For cytokine production, $1.0x10^6$ PBMC

were plated in 96 well U-bottom plates in complete RPMI with L-glutamine (10% FBS, 10 mM

HEPES, 10 μM β-mercaptoethanol) for 20 h at 37 C. T cells were stimulated with 0.5 μg/mL CD3ϵ

mAb (OKT3; ThermoFisher) for 20 hrs at 37 C in the presence of 10 μg/mL GolgiStop

(ThermoFisher). For flow cytometry, $0.5-1.0x10^6$ PBMC were stained with Live/Dead NIR Zombie

(Biolegend), incubated with supernatant from 1D4 hybridoma for 30 min, followed by anti-mouse

IgG PE, and the remaining surface antibodies as indicated. Intracellular staining was performed

using the Transcription Factor Staining Kit (eBiosciences) with TNF-α and IFN-γ antibodies.

Samples were acquired on a Cytek Aurora (4L and 5L) instrument.

Human scRNAseq kidney transplant biopsy analysis. Biopsies were collected from kidney

transplant patients experiencing acute rejection as described.[33] Gene count matrices for the

samples were generated by CellRanger count, and analyzed in R (4.3.0) using Seurat (4.3.0.1).

Cells with excluded with fewer than 200, more than 2500 gene counts, or greater than 20%

mitochondrial DNA. Analysis was filtered on mature $CD8^+$ T cells defined as cells expressing at

least one chain of CD3, CD8A, and not expressing CD4. The samples were integrated using the

top 2000 genes and 50 dimensions. Dimensionality reductions and clustering analysis were

generated using the first 12 dimensions and a resolution of 0.5. Relevant figures were

generated with ggplot2 (3.4.3) and pheatmap (1.0.12).

Quantification and Statistical Analysis

Statistical Analysis. All data points represent individual mice, and where individual data points

are not depicted the value of n is provided in the corresponding figure legend. For analysis of

absolute numbers and expression levels, paired or unpaired Student's t-tests (two-tailed) were

performed between two groups; one-way or two-way ANOVA with multiple comparison tests were

used to compare multiple groups. Log-rank (Mantel-cox) test was used to evaluate graft survival

between groups. Error bars represent standard error measurements (SEM). Statistics were

performed using GraphPad Prism 9 or 10. Significance was determined as $*p<0.05$, $**p<0.01$,

$***p<0.001$, $****p<0.0001$.

Study Approval. Healthy donors and kidney transplant candidates were isolated following

protocols approved by the Johns Hopkins Institutional Review Board.

3. RESULTS

3.1 CD43 1B11 agonism with mAb provides costimulatory signal to CD8$^+$ T cells

Published studies with multiple experimental systems and reagents have demonstrated that the CD43 receptor can provide either costimulatory or inhibitory signals to T cells.[25,34,35] We stimulated purified CD8$^+$ T cells with a range of plate-bound CD3ε and CD28 mAb in the presence or absence of plate-bound CD43 1B11 mAb **(Figure 1A)**. We found that a concentration of 10 μg/mL CD3ε/CD28 elicited uniform proliferation of CD8$^+$ T cells **(Figure 1A)**, while 0.625 μg/mL of CD3ε/CD28 was below a threshold necessary to induce CD8$^+$ T cell proliferation **(Figure 1B)**. However, the addition of CD43 1B11 mAb agonism elicited robust proliferation of CD8$^+$ T cells **(Figure 1B)**. These data demonstrate that the CD43 1B11 pathway can provide a costimulatory signal to CD8$^+$ T cells when a sub-threshold strength of antigen stimulation is present.

3.2 CD43 mAb treatment overcomes tolerance to induce graft rejection.

We reasoned that because CD43 1B11 can provide activating signals to CD8$^+$ T cells and CD43 1B11$^+$ CD8$^+$ T cells are potent effectors, that signaling through this receptor could elicit graft rejection. We induced tolerance with combined CD28/CD40 costimulation blockade (CTLA-4 Ig and CD40L mAb; **Figure 1C**), and treated mice with CD43 1B11 mAb on day 8-14, which corresponds to the initial expression of CD43 1B11 epitope on CD8$^+$ T cells [29]. We found that while costimulation blockade induced graft survival to 30 days, treatment with CD43 1B11 elicited graft rejection in 63% of mice by day 20 **(Figure 1D)**. While costimulation blockade elicited few activated effector CD8$^+$ T cells, mice treated with CD43 1B11 had a higher frequency of CD62L^{lo}CD27^{lo} T cell population **(Figure 1E)**. This population is highly proliferative, as denoted by expression of Ki67 **(Figure 1F)**. Together, these data demonstrate that activation of

the CD43 1B11 signaling pathway is critical for acute rejection, and can overcome

transplantation tolerance.

3.3 The CD43$^+$ICOS$^+$ phenotype specifically identifies proliferative graft-specific CD8$^+$ T

cells

As transplantation elicits relatively weak expansion of graft-specific CD8$^+$ T cells,[29,36] we

next evaluated whether the CD43 1B11$^+$ICOS$^+$ phenotype can be used to identify relatively rare

proliferative CD8$^+$ T cells after transplantation by evaluating expression of Ki67, which denotes

recently divided or actively proliferating T cells in H-2^b C57Bl/6 mice that received an H-2^d

Balb/c skin graft on day 14 post-grafting **(Figure 2A)**.

Among the canonical activation markers CD44^{hi} or CD62L^{lo}, many CD44^{hi} or CD62L^{lo}

cells were Ki67$^-$, indicating that these markers are highly sensitive, but not specific, for

proliferative graft-specific CD8$^+$ T cells **(Figure 2B)**. As previously published, we found that very

few KLRG-1$^+$ cells were elicited during acute rejection **(Figure 2B)**, precluding further analysis

of KLRG-1$^+$Ki67$^+$ cells. We found that 57.9$\pm$4.7% and 32$\pm$4.3% of CXCR3$^+$ cells were Ki67$^+$ in

the L^d QL9$^+$ and bulk CD8$^+$ T cell compartments, respectively **(Figure 2B)**. However, both CD43

1B11$^+$ and ICOS$^+$ populations were highly enriched for Ki67$^+$ CD8$^+$ T cells **(Figure 2B)**. Among

ICOS$^+$, 81.1$\pm$4.1% of L^d QL9$^+$ and 66.0$\pm$3.8% of bulk CD8$^+$ T cells were Ki67$^+$; among CD43

1B11$^+$ cells, 87.5$\pm$2.9% L^d QL9$^+$ and 64.5$\pm$4.7% bulk CD8$^+$ T cells were Ki67$^+$.

To directly visualize the proliferation of CD8$^+$ T cells during acute rejection, we adoptively

transferred polyclonal CellTrace Violet (CTV) labeled naïve CD45.1 CD8$^+$ T cells naïve C57Bl/6

(CD45.2) hosts, followed by Balb/c skin grafting **(Figure 2C)**. In parallel, we transferred CTV

labeled Thy1.1$^+$ OT-I CD8$^+$ T cells into C57Bl/6 hosts and provided Act-mOVA skin grafts as a

comparison high-affinity antigen-specific response. After 14 days, we identified the graft-specific

L^d QL9$^+$ CD8$^+$ T cells, bulk CD8$^+$ T cells, or OT-I cells using magnetic column enrichment.

Expression of ICOS and CD43 1B11 were largely restricted to divided cells in each population

(Figure 2D-E). Among divided bulk $CD8^+$ T cells, the frequency of $ICOS^+$ cells was 41.6±4.9%,

while 73.6±4.8% were CD43 $1B11^+$. L^d $QL9^+$ $CD8^+$ T cells displayed a similar profile, with

divided cells being 84.2±7.5% $ICOS^+$ and 91.8±4.6% CD43 $1B11^+$. Thus, at this timepoint, while

both ICOS and CD43 1B11 are specific for proliferating $CD8^+$ T cells, CD43 1B11 is a slightly

more sensitive marker for proliferating graft-specific $CD8^+$ T cells.

3.4 CD43 $1B11^+$ CD8+ T cells rapidly infiltrate heart allografts during acute rejection

We evaluated whether the functional potency of CD43 $1B11^+$ $CD8^+$ T cells identified in

the secondary lymphoid organs corresponded to infiltration into fully vascularized heart

allografts during acute rejection. In this fully allogeneic heart graft model, acute rejection occurs

within 1 week post-graft [37,38]. We evaluated the phenotype of heart graft-infiltrating $CD8^+$ T cells

relative to the spleen on day 5 post-graft **(Figure 2G)**, and found that while only 7.4±3.6% of

$CD8^+$ T cells were CD43 $1B11^+$ $ICOS^+$ in the spleen at this early timepoint, 53.4±14.7% of graft

infiltrating $CD8^+$ T cells were CD43 $1B11^+ICOS^+$ **(Figure 2H)**. These data demonstrate that

CD43 $1B11^+$ $ICOS^+$ are a significant portion of the graft-infiltrating $CD8^+$ T cells during acute

rejection.

3.5 CD43 $1B11^+$ $CD8^+$ T cells maintain expression of IL-7Rα and TCF1 during acute

rejection

We next evaluated the expression of key effector and stem-like markers of graft-specific

$CD44^{hi}$ $CD8^+$ T cells at effector and memory timepoints **(Figure 3A)**. CD43 $1B11^+$ $CD8^+$ T cells

were nearly universally positive for Ki67 **(Figure 3B-C).** However, CD43 $1B11^+$ $CD8^+$ T cells also

expressed similarly high levels of IL-7Rα to naïve, $CD62L^{hi}$ and $CD62L^{lo}$ effector populations

(Figure 3B-C). CD43 $1B11^+$ effectors expressed significantly higher levels of T-bet than either

CD43 1B11$^-$ effector population (85±2.9% CD43 1B11$^+$ vs 48±3.1% CD62L$^+$CD43 1B11$^-$ and

52±5.1% CD62L$^-$CD43 1B11$^+$; **Figure 3B-C**). Finally, CD43 1B11$^+$ CD8$^+$ T cells also maintained

relatively high levels of TCF-1 (69.9 ± 4.3%) similar to naïve and CD43 1B11$^-$ effector CD8$^+$ T

cell populations (**Figure 3B-C**). A similar expression profile of Ki67, IL-7Rα, T-bet, and TCF-1

was observed for bulk CD44^{hi} CD8$^+$ T cell populations **(Figure S1A-B)**.

248   At memory timepoint (>35 days post-graft), we found that all three populations of CD8$^+$ T

cells were high for IL-7Rα and TCF-1 **(Figure 3D-E)**. However, CD43 1B11$^+$ CD8$^+$ T cells

maintained higher expression of T-bet (72.6±3.9% CD43 1B11$^+$ vs 48.0±5.0% CD62L$^+$CD43

1B11$^-$ and 47.2±6.2% CD62L$^-$CD43 1B11$^+$; **Figure 3D-E**). Together, these data demonstrate that

CD43 1B11$^+$ CD8$^+$ T cells have an effector and memory program characterized by high T-bet

expression versus CD43 1B11$^-$ populations, but also largely maintain expression of stemness

markers IL-7Rα and TCF-1.

3.6 Effector CD43 1B11 CD8$^+$ T cells persist relative to CD43 1B11$^-$ CD8$^+$ T cells after

transplantation

258   We used an adoptive transfer approach to assess the fate, and potential persistence, of

259   CD43 1B11$^+$ CD8$^+$ T cells relative to CD43 1B11$^-$ CD8$^+$ T cells. We isolated CD62L^{lo} CD43 1B11$^-$

260   and CD43 1B11$^+$ CD8$^+$ T cells from CD45.1 recipients of Balb/c skin grafts using flow cytometric

261   sorting **(Figure 4A)**. We adoptively transferred equal numbers of these cells into CD45.2

262   C57Bl/6 mice, challenged them with a Balb/c skin graft, and evaluated the number and

263   phenotype of transferred populations on day 21 post-graft.

264   We found that relative to CD43 1B11$^-$ T cells, greater numbers of CD43 1B11$^+$ T cells

were recovered from secondary lymphoid tissue **(Figure 4B)**. While CD43 1B11$^+$ T cells

remained CD43 1B11$^+$ after skin grafting, a fraction of CD43 1B11$^-$ T cells converted to CD43

1B11$^+$ T cells following the second skin graft challenge **(Figure 4C)**. Evaluation of the

localization of these transferred populations in the draining lymph node revealed that CD43

1B11+ population localizes to the high endothelial venule relative to the CD43 1B11- CD8+ T

cells **(Figure 4D)**. Together, these data demonstrate that CD43 1B11+ CD8+ T cells appear to

be a terminal phenotype, as they are formed from CD43 1B11- precursors, and are persistent *in*

vivo.

3.7 CD8+ T cells that are *SPN+GCNT1+* are present in the rejecting allografts of kidney

transplant patients experiencing acute rejection

In order to investigate whether a population of CD43+ CD8+ T cells could be identified in

the allografts of kidney transplant patients during acute rejection, we explored the single cell

RNA-sequencing data generated by Shi et al [33] **(Figure 5A).** Among kidney transplant patients

undergoing acute cellular rejection, we identified 8 clusters of CD8+ T cells **(Figure 5B-C)**. All

clusters expressed the CD43 gene *SPN*. The because the glycosyltransferase GCNT1 is

responsible for glycosylating the CD43 receptor [13,27,28], we reasoned that *SPN+GCNT1+* clusters

could be a surrogate for CD43 1B11+ CD8+ T cells observed using flow cytometry. We found that

clusters 1, 2, 4, and 6 were *SPN+GCNT1+* **(Figure 5D)**. The *SPN+GCNT1+* clusters were higher

for Granzyme B (*GZB*) and lower for CD62L (*SELL*), which is consistent with the CD43 1B11+

CD8+ T cell population in mice. The *SPN+GCNT1+* clusters expressed high levels of granzymes

and perforin (*GZMA, GZMH, GZMK, PRF1*). Relative to GCNT1- clusters present in the allograft,

they expressed elevated transcripts for IFN-γ (*IFNG*), the chemokine receptor CXCR3, and the

co-receptors PD-1 and ICOS, which are markers of recent antigen encounter. Overall, these

results reveal that populations of CD8+ T cells that are *SPN+GCNT1+* are present in rejecting

kidney allografts, and these clusters have features of highly potent effectors that have recently

encountered antigen.

293 **3.8 CD43 1D4 mAb identifies a population of effector CD8$^+$ T in healthy human donors**

294 **and kidney transplant candidates**

295 In humans, high and low molecular weight forms of the CD43 receptor have been

296 identified [39]. The CD43 1D4 mAb clone identifies a large core-2 O-glycosylated isoform of CD43

297 on human T cells [39]. In healthy donors, we found that a fraction of CD14$^+$ monocytes and CD8$^+$

298 T cells stained for CD43 1D4 **(Figure 6A)**. We investigated whether the CD43 1D4$^+$ CD8$^+$ T

299 cells corresponded with the canonical naïve (CCR7$^+$CD45RA^{hi}) or antigen-experienced T cell

300 populations: central memory (T$_{CM}$, CCR7$^+$CD45RA^{lo}), effector memory (T$_{EM}$, CCR7$^-$CD45RA^{lo}),

301 or T effector memory RA$^+$ (T$_{EMRA}$, CCR7$^-$CD45RA$^+$) populations. While CD43 ID4$^-$ CD8$^+$ T cells

302 were predominantly found within the naïve compartment, CD43 1D4$^+$ CD8$^+$ T cells were found

303 predominantly within antigen-experience fractions **(Figure 6B).** Further exploration of the

304 phenotype of these cells revealed that CD43 1D4$^+$ CD8$^+$ T cells had lower expression of CD27

305 and higher expression of CXCR3 and ICOS than CD43 1D4$^+$ CD8$^+$ T cells **(Figure 6C)**.

306 We next investigated whether CD43 1D4 was expressed on CD8$^+$ T cells from patients

307 with end-stage renal disease who are waitlisted for kidney transplantation. Similar to healthy

308 donors, a population of CD8$^+$ T cells expressed CD43 1D4 **(Figure 6D)**. We found that relative

309 to CD43 1D4$^-$, CD43 1D4$^+$ CD8$^+$ T cells had variable and statistically similar expression of

310 CD45RA and CCR7, demonstrating that this population did not correspond to a canonical T$_{EM}$,

311 T$_{CM}$, or T$_{EMRA}$ populations **(Figure 6E).** CD43 1D4$^+$ CD8$^+$ T cells from these patients also

312 displayed lower expression of CD27 and higher expression of ICOS and CXCR3 **(Figure 6F).**

313 Overall, these data demonstrate that CD43 1D4 identifies a population of antigen-experienced

314 CD8$^+$ T cells in human healthy donors and renal transplant candidates.

315

316 **3.9 CD43 1D4$^+$ CD8$^+$ T cells are efficient cytokine producers relative to CD43 1D4$^-$ CD8$^+$ T**

317 **cells**

To evaluate the functional capacity of CD43 1D4$^+$ and CD43 1D4$^-$ CD8$^+$ T cells, we

stimulated PBMC from healthy donors with CD3ϵ mAb for 18-20 hours and evaluated CD8$^+$ T

cells based on CD45RA and CD43 1D4 expression **(Figure S2A and Figure 6G).** We found that

CD3ϵ mAb stimulation increased CD43 1D4 expression among both CD45RA^{hi} and CD45RA^{lo}

populations **(Figure S2B)**. This change was very modest among CD45RA^{lo} cells, and was more

pronounced among CD45RA^{hi} cells, presumably due to activation of a subset of naïve CD8$^+$ T

cells. When we compared the cytokine production among CD43 1D4$^+$ and 1D4$^-$ fractions, we

found that both CD45RA^{hi} and CD45RA^{lo} CD43 1D4$^+$ fractions produced elevated levels of TNF-

α and IFN-γ relative to the CD43 1D4$^-$ populations **(Figure 6D).** In total, these data demonstrate

that CD43 1D4 is a marker of antigen-experienced and acute effector CD8$^+$ cells that are

functional as determined by cytokine production.

4. DISCUSSION

In this study, we built upon previous work that showed graft-specific L^d QL9$^+$ CD8$^+$ T cells express the activated glycoform of the siaolomucin CD43, defined by the 1B11 mAb.[29] Specifically, we sought to evaluate whether CD43 1B11 a marker of short-lived, highly proliferative CD8$^+$ T cells, analogous to KLRG-1$^+$ short-lived effector CD8$^+$ T cells.

We found that treatment with CD43 1B11 mAb *in vivo* was able to break costimulation-blockade induced tolerance and elicit skin graft rejection in the majority of mice after Balb/c skin grafting. To our knowledge, prior studies using CD43 mAb have not distinguished between global signaling through the CD43 receptor and the specific function of activated CD43 1B11$^+$ receptor.[25,34,35] Our results demonstrate that CD43 1B11 agonism provides a costimulatory signal to CD8$^+$ T cells that can overcome sub-threshold antigen signaling, and that this pathway is critical for alloimmunity *in vivo*.

We found that CD43 1B11$^+$ status was highly sensitive for proliferating CD8$^+$ T cells during acute rejection, and that CD43 1B11$^+$ CD8$^+$ T cells are a significant fraction of graft-infiltrating CD8$^+$ T cells during acute rejection of heart allografts. Consistent with a highly potent effector phenotype, CD43 1B11$^+$ CD8$^+$ T cells had higher levels of T-bet expression versus CD43 1B11$^-$ CD8$^+$ T cell populations. However, in contrast to terminally differentiated effector CD8$^+$ T cells, CD43 1B11$^+$ CD8$^+$ T cells also maintained expression of memory-precursor and stem-like makers TCF-1 and IL-7Rα, and were more persistent in adoptive transfer experiments. TCF1 is associated with the survival and proliferative capacity of T cells in a variety of settings,[18,19,43,44] and TCF1 expression in CD4$^+$ T cells corresponds with self-renewal capacity of effectors in transplantation.[43] Further work is needed to understand the importance of the heterogeneity of TCF-1 expression within the CD43 1B11$^+$ population, including whether this population is predominantly self-renewing or fed by CD43 1B11$^-$ cells.

In total, these results demonstrate that CD43 1B11$^+$ CD8$^+$ T cells possess a combination

of effector and stem-like features: potent effector capacity (cytokine production, rapid graft

infiltration, localization near the HEV), robust proliferation, and enhanced survival. In contrast to

the early memory-effector precursor model after viral infection, our findings point to a

progressive differentiation model of CD8$^+$ T cell activation in which CD43 1B11$^+$ CD8$^+$ T cells

dominate the proliferative cytotoxic response during the acute phase, but also efficiently persist

into memory after antigen clearance. Additional studies are needed to further dissect the

population dynamics of the CD43 1B11$^+$ population and elucidate the full transcriptional program

driving its function. Future studies are needed to explore the potential approaches to

therapeutically target them in mice and humans.

367
368

ACKNOWLEDGEMENTS/FUNDING

We thank members of the Johns Hopkins Department of Pathology Division of Immunology for helpful discussions. We appreciate Hao Zhang and the Bloomberg School of Public Health Flow Cytometry and Cell Sorting Core for excellent cell sorting, the Johns Hopkins Ross Flow Cytometry Core, and the NIH Tetramer Core Facility for tetramer reagents and technical advice. We thank the Johns Hopkins Collaborative Outcomes Research in Endocrine Surgery (CORES) for human subject sample collection and storage. This work was supported by National Institute of Allergy and Infection Disease (NIAID) R56 AI179856 (S.M.K.), American Society of Transplantation Research Network Career Transition Grant (S.M.K.), KidneyCure Carl W. Gottschalk Research Scholar Grant (S.M.K.), NIAID R01 AI114496 (J.S.B.), National Heart, Lung, and Blood Institute (NLBI) R01HL118183 (D.C.), NHLBI R01HL136586 (D.C.), National Cancer Institute R01CA289729 (D.C.), The American Heart Association 23EIA1040103 (D.C.), Global Autoimmune Institute (D.C.), and the Matthew Poyner MVP Memorial Myocarditis Research Fund (D.C.).

DISCLOSURE

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

Further information and requests for resources and reagents should be directed to the lead

contact, Scott Krummey (skrumme1@jhmi.edu)

AUTHOR CONTRIBUTIONS

Conceptualization, S.M.K., G.S.C., J.S.B., R.P.L., J.S.F., B.O., D.C.; methodology, G.S.C., J.S.F.,

M.W.S., R.H., S.M.K., C.M., J.S.B., R.W.; formal analysis, S.M.K., G.S.C., M.W.S.; investigation,

G.S.C., J.S.F., R.P.L., M.W.S., R.H., R.W.; writing, G.S.C., S.M.K.; visualization, S.M.K., G.S.C.,

J.S.F., R.P.L.; supervision, S.M.K., J.S.B., C.M., D.C.

 FIGURE LEGENDS

Figure 1. Treatment with CD43 1B11 mAb can break costimulation-blockade induced

tolerance. (A) C57Bl/6 mice were provided with Balb/c skin grafts and treated with tolerance-

inducing costimulation-blockade with CTLA-4 Ig and CD154 mAb (250 μg/dose, post-graft day 0,

2, 4, 6). Groups of mice were treated with IgG2a isotype control or CD43 1B11 mAb (200 μg/dose)

on post-graft day 8, 11, and 14. (B) On post-graft day 14, the phenotype of CD8$^+$ T cells was

assessed in the blood of skin grafted mice. (C) The expression of Ki67 was assessed on

populations of CD8$^+$ T cells in the blood from B. (D) Skin graft survival was assessed on mice

treated as described in A. Each data point depicts an individual mouse. Summary data depicts

pooled results from 2 independent experiments (n=6-8 mice/group) and depicts mean ± SEM.

Statistics performed by (B) Student's t-test (2-tailed), (C) (C) 1-way ANOVA or (E) mixed-effects

model (REML) with Tukey's multiple comparisons test or (E) unpaired Student's t-test. Error bars

depict SEM. **p<0.01, ***p<0.001, ****p<0.0001.

Figure 2. Proliferative CD8$^+$ T cells are defined by CD43 1B11$^+$ and ICOS$^+$ phenotype after

transplantation. (A) C57Bl/6 mice were provided with Balb/c skin grafts and CD8$^+$ T cells were

assessed on day post-graft 14 in the spleen and graft draining lymph nodes. (B) Representative

flow and summary data depicting the frequency of Ki67$^+$ cells among CD8$^+$ T cells expressing the

indicated marker. Minimal expression of KLRG-1 precluded analysis. (C) 2x10^6 naïve CD45.1 or

1x10^6 Thy1.1 OT-I CD8$^+$ T cells labeled with CellTrace Violet were adoptively transferred into

CD45.2 C57Bl/6 mice, who were provided with Balb/c or Act-mOVA skin grafts, respectively.

Transferred cells were enriched using paramagnetic beads specific for APC, which stained with

Thy1.1 or L^d QL9 tetramers. (D) Among the adoptively transferred CD8$^+$ T cell populations, gates

were drawn around the undivided (gray) and proliferation (orange) populations. The minimal

number of undivided OT-I cells precluded further analysis. (E) The frequency of ICOS$^+$ cells

among each indicated adoptively transferred CD8$^+$ T cell population. (F) Frequency of CD43

1B11$^+$ cells among each indicated adoptively transferred CD8$^+$ T cell population. (G-H) C57Bl/6

mice were provided with heterotopic Balb/c heart grafts and the phenotype of CD8$^+$ T cells was

assessed in the spleen and heart graft on day 5 post-graft. Each data point depicts an individual

mouse. Summary data depicts pooled results from 2-5 independent experiments (n=5-10

mice/group) and depicts mean $\pm$ SEM. Statistics performed by (B) 1-way ANOVA with Dunnett's

multiple comparison test versus CD44^{hi} and (H) unpaired Student's t-test (2-tailed). Error bars

depict SEM. N.D, no data, due to low cell numbers. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 3. Graft-specific CD43 1B11$^+$ CD8$^+$ T cells have both effector and stem-like

phenotypic features. (A) C57Bl/6 mice were provided with Balb/c skin grafts and CD8$^+$ T cells

were assessed on day 14 post-graft (Acute Effector) or day >35 post-graft (Memory) in pooled

spleen and graft draining lymph nodes. Graft-specific cells were identified after enrichment using

L^d QL9 tetramers and APC microbeads, and gated based on CD62L and CD43 1B11 expression

as depicted. (B) Representative flow plots and (C) summary data depicting expression of Ki67,

IL-7Rα, TCF-1, and T-bet among CD44^{lo} or CD44^{hi} L^d QL9$^+$ CD8$^+$ T cell populations at the Acute

Effector timepoint. See also **Figure S1.** (D) Representative flow plots and (E) summary data

depicting expression of Ki67, IL-7Rα, TCF-1, and T-bet among CD44^{lo} or CD44^{hi} L^d QL9$^+$ CD8$^+$ T

cell populations at the Memory timepoint. Summary data depicts pooled results from 2-3

independent experiments (n=8-14 mice/group) and depicts mean $\pm$ SEM. Statistics performed by

(C, E) 1-way ANOVA with Tukey's multiple comparison test (excluding CD44^{lo} population used as

a staining control). Error bars depict SEM. *p<0.05, **p<0.01, ***p<0.001, ***p<0.0001.

Figure 4. CD43 1B11$^+$ CD8$^+$ T cells are persistent after grafting relative to CD43 1B11$^-$ CD8$^+$ T cells. (A) Effector CD45.1 CD8$^+$ T cells (post-graft day 10-14 after Balb/c skin grafts) were flow cytometrically sorted into CD43 1B11$^+$ and CD43 1B11$^-$ populations. 1x10^5 were adoptively transferred into C57Bl/6 (CD45.2) hosts, who were provided with Balb/c skin grafts the following day. Transferred cells were assessed on post-graft day 14. (B) Recovery of CD43 1B11$^+$ and CD43 1B11$^-$ CD8$^+$ T cells in the secondary lymphoid organs. (C) Frequency of CD43 1B11$^+$ CD8$^+$ T cells among adoptively transferred populations. (D) Immunofluorescence histology of draining lymph nodes depicting adoptively transferred CD45.1 CD8$^+$ T cells (red) and the HEV marker ER-TR7 (green). Arrows depict localization of adoptively transferred CD45.1 CD8$^+$ T cells outside (CD43 1B11$^-$) or within (CD43 1B11$^+$) HEV. Summary data depicts pooled results from 3 independent experiments (n=6-10 mice/group) and depicts mean $\pm$ SEM. Statistics performed by unpaired Student's t-test (2-tailed). Error bars depict SEM. HEV, high endothelial venule. *p<0.05, **p<0.01, ***p<0.001, ***p<0.0001.

Figure 5. A population of effector-like graft-infiltrating CD8$^+$ T cells from kidney transplant patients are *SPN$^+$GCNT1$^+$*. (A) Biopsies from kidney transplant patients with clinical signs of acute rejection were analyzed by single cell RNA-sequencing. (B) Hierarchical clustering and U-MAP analysis of CD8$^+$ T cells from kidney allograft biopsies. (C) Expression of *SPN, GCNT1, GZMB,* and *SEL*L among CD8$^+$ T cell clusters. (D) Heatmap depicting expression genes for T cell of co-receptor and effector proteins.

Figure 6. A population of human CD8$^+$ T cells are CD43 1D4$^+$ and have an antigen-experienced effector phenotype. (A) The frequency of peripheral healthy donor CD14$^+$ monocytes and CD8$^+$ T cells that express the CD43 epitope 1D4. (B-C) Expression of CD45RA and CCR7 among CD43 1D4$^+$ and 1D4$^-$ CD8$^+$ T cells from healthy donors. (C) Representative flow plots and summary data showing expression of CD27, CXCR3, and ICOS among CD43 1D4$^+$

and 1D4$^-$ CD8$^+$ T cell populations. (D) Frequency of CD43 1D4$^+$ CD8$^+$ T cells among kidney

transplant candidates. (E) Expression of CD45RA, CCR7 among CD43 1D4$^+$ and CD43 1D4$^-$

CD8$^+$ T cells from kidney transplant candidates. (F) Expression of CD27, CXCR3, and ICOS,

among CD43 1D4$^+$ and CD43 1D4$^-$ CD8$^+$ T cells from kidney transplant candidates. (G) Healthy

donor PBMC were stimulated for 20-24 h with CD3ε mAb (OKT3) and assessed for TNF-α and

IFN-γ production. See also **Figure S2**. (H) Summary data depicting the proportion of TNF-α (left)

or IFN-γ (right) produced by each CD45RA and CD43 1D4 quadrant population. Each data point

depicts an individual human donor. Summary data depicts pooled results from (A-C) 3

independent experiments (n=12-13 individuals/group), (D-F) 2 independent experiments (n=10

individuals/group), (H) 3 independent experiments (n=11 individuals/group). Summary data and

depicts mean $\pm$ SEM. Statistics performed by (B) 2-way ANOVA with Sidak's multiple comparisons

test and (C, E) paired Student's t-test (2-tailed), (G) 1-way ANOVA with Dunnett's multiple

comparisons test. Error bars depict SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure S1. CD43 1B11$^+$ CD8$^+$ T cells have both effector and stem-like phenotypic features.

Related to **Figure 3.** C57Bl/6 mice were provided with Balb/c skin grafts and CD8$^+$ T cells were

assessed on day 14 post-graft (Acute Effector) in the pooled spleen and graft draining lymph

nodes. (A) Representative flow plots and (B) summary data depicting expression of Ki67, IL-7Rα,

TCF-1, and T-bet among CD44^{lo} or CD44^{hi} L^d QL9$^+$ CD8$^+$ T cell populations at the Acute Effector

timepoint. Summary data depicts pooled results from 2-3 independent experiments (n=8-14

mice/group) and depicts mean $\pm$ SEM. Statistics performed by 1-way ANOVA with Tukey's multiple

comparison test (excluding CD44^{lo} population used as staining control). Error bars depict SEM.

*p<0.05, **p<0.01, ***p<0.001, ***p<0.0001.

Figure S2. Evaluation of cytokine production by human CD43 1D4$^+$ CD8$^+$ T cell populations.

Related to **Figure 6**. (A) Healthy donor PBMC were stimulated for 20-24 h with CD3ϵ mAb (OKT3)

and assessed for TNF-α and IFN-γ production. (B) Representative flow cytometry plots and

summary data showing expression of CD43 1D4 expression among CD45RA^{hi} and CD45RA^{lo}

populations of CD8$^+$ T cells. Summary data depicts pooled results from 3 independent

experiments (n=11 individuals/group), and depict mean $\pm$ SEM. Statistics performed by (B) paired

Student's t-test (2-tailed), **p<0.01, ****p<0.0001

SUPPLEMENTAL INFORMATION

Additional supporting information may be found online in the Supporting Information section.

REFERENCES

1.   Callemeyn J, Lamarthée B, Koenig A, Koshy P, Thaunat O, Naesens M. Allorecognition

and the spectrum of kidney transplant rejection. *Kidney Int.* Published online December

2021. doi:10.1016/j.kint.2021.11.029

2.   Cherukuri A, Salama AD, Mehta R, et al. Transitional B cell cytokines predict renal

allograft outcomes. *Sci Transl Med.* 2021;13(582). doi:10.1126/scitranslmed.abe4929

3.   Cole EH, Johnston O, Rose CL, Gill JS. Impact of Acute Rejection and New-Onset

Diabetes on Long-Term Transplant Graft and Patient Survival. *Clinical Journal of the*

American Society of Nephrology. 2008;3(3). doi:10.2215/CJN.04681107

4.   Zaitsu M, Issa F, Hester J, Vanhove B, Wood KJ. Selective blockade of CD28 on human

T cells facilitates regulation of alloimmune responses. *JCI Insight.* 2017;2(19).

doi:10.1172/jci.insight.89381

5.   Liu D, Badell IR, Ford ML. Selective CD28 blockade attenuates CTLA-4–dependent

CD8+ memory T cell effector function and prolongs graft survival. *JCI Insight.* 2018;3(1).

doi:10.1172/jci.insight.96378

6.   Crepeau RL, Ford ML. Challenges and opportunities in targeting the CD28/CTLA-4

pathway in transplantation and autoimmunity. *Expert Opin Biol Ther.* 2017;17(8):1001-

1012. doi:10.1080/14712598.2017.1333595

7.   Ford ML, Adams AB, Pearson TC. Targeting co-stimulatory pathways: transplantation and

autoimmunity. *Nat Rev Nephrol.* 2014;10(1):14-24. doi:10.1038/nrneph.2013.183

8.   Krummey SM, Ford ML. Heterogeneity within T Cell Memory: Implications for Transplant

Tolerance. *Front Immunol*. 2012;3. doi:10.3389/fimmu.2012.00036

9.   Jameson SC, Masopust D. Understanding Subset Diversity in T Cell Memory. *Immunity*.

2018;48(2):214-226. doi:10.1016/j.immuni.2018.02.010

10.   Chung HK, McDonald B, Kaech SM. The architectural design of CD8+ T cell responses in

acute and chronic infection: Parallel structures with divergent fates. *Journal of*

Experimental Medicine. 2021;218(4). doi:10.1084/jem.20201730

11.   Ford ML. T Cell Cosignaling Molecules in Transplantation. *Immunity*. 2016;44(5).

doi:10.1016/j.immuni.2016.04.012

12.   Zehn D, Thimme R, Lugli E, de Almeida GP, Oxenius A. 'Stem-like' precursors are the

fount to sustain persistent CD8+ T cell responses. *Nat Immunol*. 2022;23(6):836-847.

doi:10.1038/s41590-022-01219-w

13.   Nolz JC. Molecular mechanisms of CD8+ T cell trafficking and localization. *Cellular and*

Molecular Life Sciences. 2015;72(13):2461-2473. doi:10.1007/s00018-015-1835-0

14.   Zehn D, Lee SY, Bevan MJ. Complete but curtailed T-cell response to very low-affinity

antigen. *Nature*. 2009;458(7235):211-214. doi:10.1038/nature07657

15.   King CG, Koehli S, Hausmann B, Schmaler M, Zehn D, Palmer E. T Cell Affinity

Regulates Asymmetric Division, Effector Cell Differentiation, and Tissue Pathology.

Immunity. 2012;37(4):709-720. doi:10.1016/j.immuni.2012.06.021

16.   Joshi NS, Cui W, Chandele A, et al. Inflammation Directs Memory Precursor and Short-

Lived Effector CD8+ T Cell Fates via the Graded Expression of T-bet Transcription

Factor. *Immunity*. 2007;27(2). doi:10.1016/j.immuni.2007.07.010

17. Kaech SM, Cui W. Transcriptional control of effector and memory CD8+ T cell

differentiation. *Nat Rev Immunol*. 2012;12(11):749-761. doi:10.1038/nri3307

18. Broomfield BJ, Groom JR. Defining the niche for stem-like CD8+ T cell formation and

function. *Curr Opin Immunol*. 2024;89:102454. doi:10.1016/j.coi.2024.102454

19. Im SJ, Obeng RC, Nasti TH, et al. Characteristics and anatomic location of PD-1 [+] TCF1 [+]

stem-like CD8 T cells in chronic viral infection and cancer. *Proceedings of the National*

Academy of Sciences. 2023;120(41). doi:10.1073/pnas.2221985120

20. Im SJ, Hashimoto M, Gerner MY, et al. Defining CD8+ T cells that provide the proliferative

burst after PD-1 therapy. *Nature*. 2016;537(7620):417-421. doi:10.1038/nature19330

21. Braza F, Brouard S, Chadban S, Goldstein DR. Role of TLRs and DAMPs in allograft

inflammation and transplant outcomes. *Nat Rev Nephrol*. 2016;12(5):281-290.

doi:10.1038/nrneph.2016.41

22. Colvin MM, Smith CA, Tullius SG, Goldstein DR. Aging and the immune response to

organ transplantation. *Journal of Clinical Investigation*. 2017;127(7):2523-2529.

doi:10.1172/JCI90601

23. Onami TM, Harrington LE, Williams MA, et al. Dynamic Regulation of T Cell Immunity by

CD43. *The Journal of Immunology*. 2002;168(12):6022-6031.

doi:10.4049/jimmunol.168.12.6022

24. Mody PD, Cannon JL, Bandukwala HS, et al. Signaling through CD43 regulates CD4 T-

cell trafficking. *Blood*. 2007;110(8):2974-2982. doi:10.1182/blood-2007-01-065276

25.   Sperling AI, Green JM, Mosley RL, et al. CD43 is a murine T cell costimulatory receptor

581   that functions independently of CD28. *Journal of Experimental Medicine*.

582   1995;182(1):139-146. doi:10.1084/jem.182.1.139

583   26.   Clark MC, Baum LG. T cells modulate glycans on CD43 and CD45 during development

584   and activation, signal regulation, and survival. *Ann N Y Acad Sci*. 2012;1253(1):58-67.

585   doi:10.1111/j.1749-6632.2011.06304.x

586   27.   Perkey E, Maurice De Sousa D, Carrington L, et al. GCNT1-Mediated O-Glycosylation of

587   the Sialomucin CD43 Is a Sensitive Indicator of Notch Signaling in Activated T Cells. *J

588   Immunol*. 2020;204(6):1674-1688. doi:10.4049/jimmunol.1901194

589   28.   Nolz JC, Harty JT. IL-15 regulates memory CD8+ T cell O-glycan synthesis and affects

590   trafficking. *Journal of Clinical Investigation*. 2014;124(3):1013-1026.

591   doi:10.1172/JCI72039

592   29.   Cohen GS, Kallarakal MA, Jayaraman S, et al. Transplantation elicits a clonally diverse

593   CD8+ T cell response that is comprised of potent CD43+ effectors. *Cell Rep*.

594   2023;42(8):112993. doi:10.1016/j.celrep.2023.112993

595   30.   Hikono H, Kohlmeier JE, Takamura S, Wittmer ST, Roberts AD, Woodland DL. Activation

596   phenotype, rather than central– or effector–memory phenotype, predicts the recall

597   efficacy of memory CD8+ T cells. *Journal of Experimental Medicine*. 2007;204(7):1625-

598   1636. doi:10.1084/jem.20070322

599   31.   Trambley J, Bingaman AW, Lin A, et al. Asialo GM1+ CD8+ T cells play a critical role in

600   costimulation blockade–resistant allograft rejection. *Journal of Clinical Investigation*.

601   1999;104(12):1715-1722. doi:10.1172/JCI8082

32.   Liu F, Kang SM. Heterotopic Heart Transplantation in Mice. *Journal of Visualized*

Experiments. 2007;(6). doi:10.3791/238

33.   Shi T, Burg AR, Caldwell JT, et al. Single-cell transcriptomic analysis of renal allograft

rejection reveals insights into intragraft TCR clonality. *J Clin Invest*. 2023;133(14).

doi:10.1172/JCI170191

34.   Walker J, Green JM. Structural requirements for CD43 function. *J Immunol*.

1999;162(7):4109-4114.

35.   Manjunath N, Correa M, Ardman M, Ardman B. Negative regulation of T-cell adhesion

and activation by CD43. *Nature*. 1995;377(6549):535-538. doi:10.1038/377535a0

36.   Bozeman AM, Laurie SJ, Haridas D, Wagener ME, Ford ML. Transplantation

preferentially induces a KLRG-1lo CD127hi differentiation program in antigen-specific

CD8+ T cells. *Transpl Immunol*. 2018;50. doi:10.1016/j.trim.2018.06.003

37.   Young JS, Khiew SHW, Yang J, et al. Successful Treatment of T Cell-Mediated Acute

Rejection with Delayed CTLA4-Ig in Mice. *Front Immunol*. 2017;8.

doi:10.3389/fimmu.2017.01169

38.   Schwarz C, Unger L, Mahr B, et al. The Immunosuppressive Effect of CTLA4

Immunoglobulin Is Dependent on Regulatory T Cells at Low But Not High Doses.

American Journal of Transplantation. 2016;16(12):3404-3415. doi:10.1111/ajt.13872

39.   Mukasa R, Homma T, Ohtsuki T, et al. Core 2-containing O-glycans on CD43 are

preferentially expressed in the memory subset of human CD4 T cells. *Int Immunol*.

1999;11(2):259-268. doi:10.1093/intimm/11.2.259

40.   Ford ML, Onami TM, Sperling AI, Ahmed R, Evavold BD. CD43 Modulates Severity and

624   Onset of Experimental Autoimmune Encephalomyelitis. *The Journal of Immunology*.

625   2003;171(12):6527-6533. doi:10.4049/jimmunol.171.12.6527

626   41.   Fay KT, Chihade DB, Chen CW, et al. Increased mortality in CD43-deficient mice during

627   sepsis. *PLoS One*. 2018;13(9):e0202656. doi:10.1371/journal.pone.0202656

628   42.   Ford ML, Evavold BD. Modulation of MOG 37–50-specific CD8+ T cell activation and

629   expansion by CD43. *Cell Immunol*. 2006;240(1):53-61.

630   doi:10.1016/j.cellimm.2006.06.007

631   43.   Zou D, Yin Z, Yi SG, et al. CD4+ T cell immunity is dependent on an intrinsic stem-like

632   program. *Nat Immunol*. 2024;25(1):66-76. doi:10.1038/s41590-023-01682-z

633   44.   Mo F, Yu Z, Li P, et al. An engineered IL-2 partial agonist promotes CD8+ T cell

634   stemness. *Nature*. 2021;597(7877):544-548. doi:10.1038/s41586-021-03861-0

635   45.   Duckworth BC, Lafouresse F, Wimmer VC, et al. Effector and stem-like memory cell fates

636   are imprinted in distinct lymph node niches directed by CXCR3 ligands. *Nat Immunol*.

637   2021;22(4):434-448. doi:10.1038/s41590-021-00878-5

Figure 1

Figure 2

Figure 3

A

B Acute Effector
L^d QL9^+ CD8^+ T Cells

C

D Memory
L^d QL9^+ CD8^+ T Cells

E

Figure 4

A

B Adoptive Transferred
CD45.1 CD8+ T cells

No. Transferred
CD45.1 Cells

C
Frequency
CD43 1B11+

D Draining Lymph Node
% of CD45.1+
in HEV

Figure 5

A Renal Biopsy → scRNA-Seq CD8+ T cells

Acute Rejection

B Graft-Infiltrating CD8⁺ T Cells

C *SPN* *GCNT1*

GZMB *SELL*

D

Cluster: 5 7 3 8 4 6 1 2

GZMA
GZMH
PRF1
SPN
CXCR3
HAVCR2
IFNG
ICOS
PDCD1
SELL
TNF
TBX21
PRDM1
GCNT1
TOX
GZMK
GZMB
LAG3
TIGIT

Figure 6

Figure S1

Figure S2

A

B

% CD43 1D4+   % CD43 1D4+

CD45RA^hi 52.5%   CD45RA^hi 64.4%

CD45RA^lo 73.6%   CD45RA^lo 76.9%