Title: Divergent Representation and Processing of Task Cues in Sensory and Prefrontal Cortices

of Preterm-Born Mice

Authors: Emily McCoy[1,3,4], Vyshnavi Pendala[4], Mona Fariborzi[1,2,4], Lara Y. Demir[4], Olivia P.

Buell[4], Samuel C. Fedde[4], Jacqueline B. Stinger[4], Luciano Elbaum[4], Troy D. Holsworth[4], Phillip

Amenyo-Awude[4], Xin Tong[4], Adema Ribic[1,2,3,4]*

Affiliations: [1]Program in Fundamental Neuroscience, [2]Interdisciplinary Fellowship in

Quantitative Neurobiology of Behavior, [3]Interdisciplinary Fellowship in Reintegrating the

Phenotype, [4]Department of Psychology, College and Graduate School of Arts and Sciences,

University of Virginia, Charlottesville, VA 22904

*Correspondence: adema.ribic@virginia.edu

Keywords: preterm, prefrontal, visual, task, representation, processing

Abstract: Preterm birth is a leading risk factor for atypicalities in cognitive and sensory

processing, but it is unclear how prematurity impacts circuits that support these functions. To

address this, we trained adult male and female mice born a day early (preterm mice) on a visual

discrimination task and found that they fail to achieve high levels of performance due to

increased responding to the non-rewarded cue (false alarms). While the representation of task

cues measured with *in vivo* electrophysiology is intact in the primary visual cortex (V1) of

trained preterm mice, the representation of the non-rewarded cue is significantly weaker in

regular spiking, putative pyramidal neurons in the prefrontal cortex (PFC), a brain area that

mediates response inhibition. Responses to both task cues are blunted in electrophysiologically

and optogenetically identified fast-spiking Parvalbumin interneurons in preterm mice, indicating

impaired processing of task cues in their PFC. Indeed, single trial neuronal responses evoked by

the non-rewarded cue predict the behavioral outcome more accurately in term than in preterm

mice. Similar cue representation and processing is present in the PFC of adolescent term-born

mice, suggesting that preterm birth impedes prefrontal maturation. Surprisingly, environmental

enrichment, a well-established paradigm that promotes sensory maturation, fails to improve the

performance of preterm mice. Altogether, our study describes the long-term impact of preterm

birth on prefrontal and visual circuits and suggests a limited capacity of early interventions for

reducing the risk of cognitive deficits after preterm birth.

Introduction

Children born prematurely (<37 weeks of gestation) are at 4-5x higher risk for

developing intellectual disability (ID) and attention-deficit/hyperactivity disorder (ADHD) [1–3], at

10x higher risk for developing autism spectrum disorder (ASD) [4,5], and display persistent deficits

in executive function and academic performance [6–8]. As preterm labor occurs in >10% of all

births in US and worldwide, preterm birth is the leading risk factor for cognitive and

neurodevelopmental conditions [6,9–11]. Yet, the neural circuits whose function may be disrupted by

preterm birth are still unknown.

Impairments in the structure and function of the frontal cortex are frequently associated

with executive dysfunction and neurodevelopmental conditions with high incidence in the

preterm population [9,12]. Unsurprisingly, neuroimaging and neurophysiological studies in preterm

infants and children revealed reduced volume of and connectivity between frontal, thalamic and

sensory areas [11,13–16]. On a cellular level, studies of postmortem prefrontal cortex samples

obtained from preterm fetuses demonstrated reduced density of inhibitory interneurons and

altered expression of genes related to inhibitory neurotransmission [17,18]. Some of the changes in

brain structure can be recapitulated in animal models of prematurity, which also display

disrupted layer-specific distribution and proportion of cortical interneurons [18–21], and alterations

in the expression of multiple neurotransmission markers [22,23]. Behaviorally, animal models of

prematurity often display persistent hyperactivity [23–26], and some cognitive deficits [18,27], but the

association between preterm birth-related changes in the frontal cortex and high incidence of

cognitive dysfunction in the preterm population remains unclear.

To address this, we performed a detailed neurophysiological study of prefrontal function

in adult mice born prematurely that were trained on a prefrontal cortex (PFC)-dependent visual

discrimination task. While adult mice born 1 day early do not display any changes in baseline

visual processing and locomotor activity, they show a significantly impaired learning trajectory.

Preterm mice respond more often than term mice to the non-rewarded cue, have weaker

representation of the non-rewarded cue in the PFC, and show disinhibition of neuronal firing

during task performance. Cue representation in regular-spiking (RS) and evoked activity of fast-

spiking (FS) neurons in the adult preterm PFC resemble those of adolescent term born mice,

suggesting that preterm birth impairs prefrontal maturation. Surprisingly, postnatal

environmental enrichment-a classical paradigm for improving the outcomes of impaired

neurodevelopment [21,28–32]-fails to improve the task performance in preterm mice, indicating

limited efficacy of early interventions in improving negative neurodevelopmental outcomes after

preterm birth [33]. Together, our study describes circuit- and cell-type specific impairments in

prefrontal function and top-down sensory processing after preterm birth, paving a path towards

uncovering the circuit mechanisms of cognitive dysfunction in the preterm born population.

Methods

Mice. Mice were maintained on C57BL/6 background (The Jackson Laboratory, Bar Harbor,

ME) on reverse 12:12 light:dark cycle (lights off 11 AM-11 PM), with food and water *ad libitum,*

except during visual discrimination training (see below). Animals of both sexes were used

between postnatal day 21 and 7 months of age. Preterm mice were generated through timed

breedings, where the day after the pairing was considered as gestational day (GD) 0. Once the

pregnancy was confirmed (>1.5g increase in weight at GD 10), pregnant dams were habituated to

handlers by daily handling sessions. Mifepristone (MFP, Milipore Sigma, Burlington, MA or

HelloBio, Bristol, UK) was dissolved in DMSO (Milipore Sigma) and 150 µg was injected

subcutaneously on GD 17. Preterm mice were delivered on GD 18. The cage with preterm mice

was occasionally supplemented with external heat and oxygen to prevent hypothermia and

hypoxia, commonly observed in preterm mice. Control term mice were obtained from timed

pregnant dams injected with DMSO on GD 17, and with MFP on GD 18. All dams received

continuous enrichment in form of plastic igloos and nesting material, as well as sunflower seeds

(Bio-Serv, Flemington, NJ) beginning 3-4 days before parturition. Animals were treated in

accordance with the University of Virginia Institutional Animal Care and Use Committee

guidelines.

Environmental enrichment. Nursing dams with their litters were transferred to a standard rat cage

when the pups reached postnatal day (P) 5. Dams were co-housed with a nursing dam and its

litter that was age matched and on BALB/c strain (purchased from Jackson Laboratory) to

facilitate social enrichment. Cages contained multiple enrichment objects: InnoDome and

InnoWheels from BioServ, gnawing sticks, multiple nesting pads, shredded paper and hay for

building nests, as well as a variety of chew toys (pumice chews, willow twigs, willow balls,

veggie chews, and hay tunnels, hideaways and sticks). Large enrichment objects were rearranged

in the cage every 2 days, while chew toys were replaced with new ones every 2-3 days. Once the

litters reached P25, they were weaned and group-housed in standard cages at 2-3 mice/cage to

allow for placement of an InnoDome and InnoWheel, as well as 2-3 small chew toys. When the

mice reached P60, they were prepared for visual discrimination task as described below.

Viruses. Following viruses were purchased from Addgene: AAV-S5E2-ChR2-mCherry (135634-

AAV1; 2×10^{13} vg/ml), AAV-EF1a-double floxed-hChR2(H134R)-mCherry-WPRE-HGHpA

(20297-AAV9; 3.2×10^{13} vg/ml) and AAV-hSyn-HI-eGFP-Cre-WPRE-SV40 (105540-AAVrg;

1.7×10^{13} vg/ml). All viruses were diluted 1:10 in sterile PBS before the injection.

Surgeries. Animals were anesthetized with isoflurane in oxygen (2-2.5% induction, 1–1.5%

maintenance), warmed with a heating pad at 38°C and given subcutaneous injections of

Buprenorphine SR (1 mg/kg) or Rimadyl (5 mg/kg) and 0.25% Bupivacaine (beneath their

scalp). Eyes were covered with Puralube (Decra, Northwich, UK). Scalp and fascia from Bregma

to behind lambda were removed, and the skull was cleaned, dried and covered with a thin layer

of Scotchbond adhesive (3M, Maplewood, MN). Skin edges were sealed with VetBond (3M).

For virus injections, the head was immobilized in a custom-made stereotactic apparatus and

small craniotomies were made on the left hemisphere with a dental drill at the following

coordinates: primary visual cortex (V1): 0.5 mm anterior to lambda, 2-3 mm lateral to midline;

anterior cingulate cortex (ACC): 2-2.5 mm anterior to bregma, 0.2-0.5 mm lateral to midline.

NanoFil syringe (35 G beveled needle; WPI, Sarasota, FL) was then used to inject the virus at 1

nl/s using a Syringe Pump (KD Scientific, Holliston, MA). Volume of injected virus was 300 nl,

injected in 2 150 nl increments at 2 different depths: 500 and 250 µm beneath the brain surface

for V1 (targeting layers 2/3 and 5), and 700 and 800 µm for the PFC. The syringe was left in

place for 5-10 minutes after the injection to allow the virus solution to diffuse and then very

slowly raised to the surface to prevent the backflow. Muscimol (TMR-X conjugate; HelloBio,

Princeton, NJ) was injected in the same manner, volume and at the same PFC coordinates of

trained term mice at 1.6 mM concentration [34].

Head plates were attached after the Scotchbond adhesive application for animals that received no

virus injection, or immediately after the virus injection for those that did. The head plate

(stainless steel, SendCutSend, Reno, NV) was attached with dental cement (RelyX Ultimate,

3M). After the cement was cured, the well of the head plate was filled with silicone elastomer

(Reynold Advanced Materials, Brighton, MA) to protect the skull.

If the mice received Rimadyl for analgesia, they were given Rimadyl dissolved in hydrogel (1%

food-grade agar in distilled water) during the recovery [35]. Animals were group-housed after the

implantation and monitored daily for signs of shock or infection. On the day of

electrophysiological recording, the animals were anesthetized as above and craniotomies (~0.5

mm in diameter) were made above V1, PFC and cerebellum with 18G needles. If mice were

injected with viruses, the existing craniotomies were reopened. The brain surface was covered in

2-3% low melting point agarose (Promega, Madison, WI) in sterile saline and then capped with

silicone elastomer. Animals were allowed to recover for 1.5-2 h before the recording.

Electrophysiology and optogenetics. For the recording sessions, mice were placed in the head-

plate holder above an aluminum mesh treadmill and allowed to habituate for 5-10 minutes. The

silicone plug was removed, the reference (insulated silver wire electrode; A-M Systems,

Carlsborg, WA) was placed in cerebellum and the well was covered with warm sterile saline. A

multisite electrode spanning all cortical layers (A1x16-5mm-50-177-A16; Neuronexus

Technologies, Ann Arbor, MI) was coated with DiI (Invitrogen) to allow post hoc insertion site

verification and then inserted in the V1 through the craniotomy. For PFC, a 4x4 shank probe

(A4x4-3mm-50-125-177-A16) was used for the recordings, coated with DiI, DiO or DiD based

on the type of fluorophore used in the experiment (mCherry or EGFP; listed in Viruses). The

electrodes were slowly (5-10 µm/s) lowered to the appropriate depth: 800 µm for the PFC and

until the uppermost recording site had entered the brain for the V1 and allowed to settle for 15-30

minutes. For optogenetics, a 473 nm laser diode with 0.4 µm fiber tip (0.22 NA, Doric Lenses,

Quebec, Canada) was lowered to the craniotomy. 10 ms square pulses were delivered at 3-5

mW/mm^2 (200 pulses at 0.5 Hz) using Spike2 (CED, Cambridge, UK) after the last recording

session for optogenetic identification of neuronal subtypes (optotagging). The well was filled

with 3% agarose prior to the recordings to stabilize the electrode and the whole region was kept

moist with surgical gelfoam soaked in sterile saline (Pfizer, MA). The signals from the recording

probes were fed into a 16-channel amplifier (Model 3500; A-M Systems, Sequim, WA) and

amplified 200x, before being sampled at 25 kHz using Spike2 and Power 1401-3 data acquisition

unit (CED). After the recording ended, the electrodes were slowly retracted, and the well was

cleaned and protected with silicone elastomer.

Visual stimuli. For visual stimuli, blank screen was generated with MATLAB (MathWorks,

Natick, MA) using the PsychToolBox extension (Brainard, 1997) and presented on a gamma

corrected 27" LCD. The screen was centered 20-25 cm from the mouse's right eye, covering

~80° of visual space. For calculating baseline orientation selectivity, sinusoidal gratings at 6

orientations 30º apart were presented at 100% contrast at 0.15 cpd, with 1 s long stimuli and 1 s

interstimulus interval (blank screen). For visual discrimination training, visual stimuli (120º and

60º gratings) were presented at 0.15 cpd and 100% contrast in a random sequence for 3 seconds,

followed by a 15 second long interstimulus interval (blank screen). Presentation of 120º triggered

the delivery of 10 µl of water through a syringe pump (New Era Pump Systems, Farmingdale,

NY), while the presentation of 60º had no consequence.

Behavior. 4-7 days after the headpost implantation, mice were gradually water restricted (from 3

ml of water/day to 1 ml of water/day) for 5-7 days. Food was available *ad libitum*. Mice were

weighed daily and their weight was maintained at 85% of initial weight to prevent dehydration.

During water restriction, mice were gradually habituated to handling by the experimenters, the

treadmill, and the water delivery spout during daily habituation sessions [36]. After water

restriction, the visual discrimination training began, where mice were headfixed above the

treadmill and a stainless-steel gavage needle (18G) was positioned near the mouse's mouth for

water delivery. The pump was controlled by the Power1401-3 data acquisition interface (CED)

for lick detection [37]. Mice were trained in 2 daily sessions consisting of 50 120º and 60º

presentations each, for a total of 200 trials per day. Licks of the spout during the 10 s after the

onset of stimulus presentation were quantified according to the signal detection theory as Hits

(the fraction of trials in which the mouse licked to the 120º orientation), Misses (the fraction of

trials in which the mouse failed to lick to the 120º orientation), False Alarms (the fraction of

trials in which the mouse licked to the 60º orientation) and Correct Rejections (the fraction of

trials in which the mouse failed to lick to the 60º orientation), where the performance was

measured as discriminability (d')=z(H) - z(FA). Mice were trained until they achieved a d' of 2

or above for 3 consecutive sessions or up to 4000 trials if they failed to reach the criterion.

For open field, mice were placed in a 24x24 inch plexiglass arena after a brief habituation to the

room and the experimenters. Their activity was recorded for 10 minutes and analyzed using an

open-source toolbox [38].

For measuring locomotor behavior during task performance, a small piezoelectric disc was

placed in contact with the treadmill and the signal was digitized using Power 1401-3 and Spike2.

Waveform averages in response to both cues were constructed using Spike2 and used to compare

the amount of treadmill movement during the training.

Animals were monitored for signs of distress for the duration of the experiment. Animals that

developed corneal damage, cataracts, and symptoms that had the potential to impact their

behavior or physiology (such as immobility and other signs of pain, sudden weight loss or

impeded weight gain) were excluded from the study. All mice except those in the MFP Term and

Environmental Enrichment groups underwent electrophysiology either at the beginning or at the

end of the training, or both. If the animals were recorded at the onset of training, that was

counted as the first training session.

Histology and imaging. After the last training session or electrophysiological recording, mice

were anesthetized with a mixture of ketamine and xylazine and transcranially perfused with

warm 0.1 M phosphate buffer, followed by warm 4% paraformaldehyde (Electron Microscopy

Sciences, Hatfield, PA). Brains were postfixed 1 hr at room temperature, followed by overnight

fixation at 4°C. Brains were sectioned into 40-80 µm sections using a vibratome and stored in 1x

phosphate buffered saline (PBS) and 0.01% sodium azide. For immunohistochemistry, sections

were rinsed in PBS, non-specific binding was blocked with 3% normal horse serum (heat

inactivated, ThermoFisher, Waltham, MA) and 0.3% Triton-X 100 (Sigma-Aldrich) in PBS

(sterile filtered). Antibodies were incubated overnight at 4°C. Parvalbumin (anti-goat) was used

at 1/200 (Swant, Belinzona, Switzerland) and detected with donkey anti-goat Alexa 647

(ThermoFisher). After staining, sections were rinsed in distilled water, mounted on glass slides,

briefly dried, and coverslipped with Aquamount (Polysciences, Warrington, PA). Images were

acquired using Leica Stellaris 8 at 2048x2048 resolution using 10x HC PLAN FLUOTAR air

(NA=0.3 for tiling) or 40x HC PL APO CS2 (NA=1.3) oil immersion objective.

Electrophysiological and behavioral data analysis. Locomotion measured with piezoelectric

sensors was analyzed in Spike2 (CED) using waveform averaging and cues as triggers. Spikes

were isolated from the recordings using template matching in Spike2 (CED). Briefly, the

recordings were band-pass filtered (0.7-7 kHz) and smoothed (1 ms). Threshold for detection

was set at 3 SDs of the mean baseline (blank screen) and a window of 0.9-1 ms. Isolated units

were clustered using waveform properties (amplitude, spike half-width and slope of

repolarization), and the clusters were checked manually for quality in addition to confirming

there were no spikes during the refractory period (2 ms) using interval histograms. Units with

refractory period violations or units with large variations in amplitude (>10%) were excluded

from analysis. For orientation selectivity, custom-written scripts in Spike2 were used to plot the

baseline-subtracted responses of isolated units to the presented orientations (10 ms bin) and

construct a tuning curve, which was then fitted with a Gaussian to determine the preferred (O_{pref})

and orthogonal (O_{orth}) orientations, and calculate the orientation selectivity index (OSI) as (R_{pref}-

R_{orth})/(R_{pref}+R_{orth}), where R is the firing rate at preferred and orthogonal orientations. We

considered the neurons with at least 3x elevated firing rate to the preferred orientation (averaged

over 0.8 s and compared to averaged 0.2 s of baseline) as visually responsive. Tuning width was

calculated as half-width at half-maximum of the O_{pref}.

For optotagging, spikes were isolated from optogenetic recording sessions and PSTHs (0.1-1 ms

bin) were constructed in response to the laser pulses. Units with reliable responses (>50%) of

trials and short latencies (within 10 ms of the laser pulse) were considered optotagged and their

waveforms were used to identify their spiking during the task performance. Unit waveforms

were further compared between optotagging and task recordings for quality assurance. Units

whose waveforms differed by more than 10% in amplitude and width between the two sessions

were discarded from analysis.

Selectivity indices (SIs) were calculated from 10 ms peristimulus time histogram (PSTHs) as

(R_{cue}-$R_{baseline}$)/(R_{cue}+$R_{baseline}$), where R_{cue} is the firing rate during the initial 1-1.5s presentation of

120º and 60º stimuli. Behavioral responses during the task were analyzed using custom scripts in

Spike2 by converting the detected licks into events and constructing PSTHs in response to the

non-rewarded cue during the 2 possible outcomes (Correct Rejection and False Alarm).

Stationary and moving stages were not analyzed separately as animals were typically walking in

short bouts during the recordings. Recordings where large motion artifacts were present when the

animal was simply balancing on the treadmill or grooming were excluded from analysis, along

with recordings from locations outside of the PFC or V1 (as determined through post-hoc

mapping of insertion sites).

Quantification and statistical analysis. All analyses were performed with the researchers blind to

the condition. Statistical analyses were performed in GraphPad Prism 9.0 (GraphPad Inc., La

Jolla, USA), R and JASP, as indicated in text and figure legends. N refers to the number of

animals or single units, as indicated in figure legends or in text. For orientation selectivity

analysis, OSIs, proportions of selective cells (OSI>0.3) and tuning width (OSI>0.3) means

(normal data distribution, Shapiro-Wilk test) or medians (non-normal distribution) were

calculated per mouse before comparison. For comparison of preferred orientations, selective

neurons (OSI>0.3) were only segregated by group (Term/Preterm). For all other analyses, units

significantly modulated by cues were determined through comparison of pre-cue and post-cue

firing using Wilcoxon signed rank test. The proportions of significantly modulated units

(positively and negatively) were determined and reported per animal, and PSTHs and SIs

represent significantly modulated units. PSTHs were compared using permutation test of area

under the curve (AUC) calculated from 0.5 s precue and 1 s (V1) or 3 s (PFC) postcue activity

(10 ms bins). Cue representation was determined as (%R-%NR)/(%R+%NR), where %R and

%NR are the proportion of significantly positively modulated units by the rewarded and non-

rewarded cue, respectively. All unit data was tested for normality using Shapiro-Wilk test and

data violating the assumption of normality was tested using non-parametric tests (Mann-Whitney

t-test, Kolmogorov-Smirnoff for comparison of distributions or non-parametric ANOVA).

Normally distributed data was tested using t-tests and ANOVA with post-hoc tests for comparing

groups as indicated in text and figure legends. Levene's test was used to test homoscedasticity

with appropriate corrections used for heteroscedasticity. For linear mixed model (LMM) in

Figure 2e, an LMM with Animal Group as a fixed effect and Subject and Training Day as

random effects was fitted to the data using the following formula and maximum likelihood

method: d` ~ Group + (1 + Group || Subject) + (1 + Group || Training Day). To test the sex

effects, LMM with Sex and Animal Group as fixed effects and Subject and Training Day as

random effects was fitted using the following formula and maximum likelihood method: d` ~

Sex + Animal Group+ (1 + Sex + Animal Group | Subject) + (1 + Sex + Animal Group | Training

Day). For Figure 2i, lick counts were averaged per 20 trials and an LMM with Animal Group

and Trial Block as fixed effects and Subject as the random effect was fitted to the data using the

following formula and maximum likelihood method: Lick count` ~ Group + `Trial block` + (1 +

Group + `Trial Block` | Animal). For Figure 6b, an LMM with Age as a fixed effect and Subject

and Training Day was fitted to the data using the following formula and maximum likelihood

method: `d`` ~ `Age` + (1 + `Age` | Training Day) + (1 + `Age` | Animal). For Figure 7b, an

LMM with Enrichment and Birth as fixed effects and Subject and Training Day as the random

effects was fitted to the data using the following formula and maximum likelihood method: `d'`

~ Enrichment + Birth + (1 + Enrichment + Birth | Subject) + (1 + Enrichment + Birth | Training

Day). For linear regression, SI values from all neurons per animal were averaged and used as

predictors of d' as described in text and figure legends. For logistic regression, trial by trial spike

time stamps were used to calculate the evoked rate per trial (number of spikes in 3 s window

postcue-number of spikes in 3 s window precue). As behavioral outcomes were biased towards

Correct Rejection, random oversampling was used to balance the datasets. Data are reported as

mean ± SEM (standard error of mean) or median±MAD (median absolute deviation), as

indicated in text or figure legends, where N represents the number of animals or the number of

single units. Target power for all sample sizes was 0.8 and alpha was set to 0.05.

Results

Adult preterm mice commit more errors during visual discrimination

Children born preterm have an increased rate of cognitive deficits and

neurodevelopmental conditions with unclear circuit origins [9]. As we previously found that

preterm birth similarly impacts the resting neural activity in the primary visual cortex (V1) of

juvenile mice and infants [39], we hypothesized that mice born preterm would also display

cognitive deficits. To test this, we chose to train the mice in a visual discrimination task, a

classical learning task that facilitates electrophysiological probing of neural circuits in awake,

behaving mice [40,41]. Preterm mice were generated through subcutaneous injections of

mifepristone (MFP) to timed-pregnant C57BL/6 dams on gestational day (GD) 17 (Figure 1A),

resulting in birth of viable preterm pups on GD 18. Term pups in our colony are born on GD 19

with very little variability, as previously published [42,43]. Before training, adult (2-7 months old)

male and female term and preterm mice were implanted with headposts for head fixation and

their baseline visual function in form of orientation selectivity was tested using *in vivo*

electrophysiology [44]. We collected neuronal responses to sinusoidal oriented gratings from all

cortical layers (6 orientations 30º apart), at 100% contrast and 0.15 cycles per degree spatial

frequency; Figures 1B), and determined the orientation selectivity index (OSI; Figure 1C), the

number of orientation selective neurons (Figure 1D), orientation tuning width (Figure 1E), and

the distribution of preferred frequencies [45]. While neurons in the V1 of naïve preterm mice had

elevated baseline firing rates (during the presentation of the blank screen; median±MAD:

Term=2.41±1.01 spikes/s; Preterm=7.49±3.65 spikes/s, p=0.061, Mann-Whitney t-test U=82,

414 neurons from N=11 term mice and 306 neurons from N=10 preterm mice), we found no

significant differences in any of the orientation selectivity measures between term and preterm

mice (OSI median±MAD: Term=0.78±0.05, Preterm=0.81±0.08, Mann Whitney t-test p=0.86,

U=52; % selective neurons mean±SEM: Term=77.59±3.81, Preterm=72.04±5.22, t-test p=0.4,

t=0.86, df=16.84; tuning width (º) median±MAD: Term=14.02±1.08, Preterm=15.19±1.73, Mann

Whitney t-test p=0.29, U=70.5; distribution of orientation preferences p=0.92, X^2 test). Increase

in neuronal firing at the preferred orientation (calculated as the ratio between firing at the

preferred orientation and baseline firing) was not different between term and preterm mice

(median±MAD Term=6.4±1.75, Preterm=6.59±2.3, Mann-Whitney t-test p=1, U=55). Our

results hence demonstrated that preterm birth in mice does not negatively impact baseline

orientation selectivity and that they can be trained in an orientation discrimination task.

To prepare the mice for task training, adult term and preterm mice were implanted with

headposts, and then gradually water restricted and habituated to the experimental setup and

handlers over 7-10 days [36], after which the training began. To account for the effects of prenatal

mifepristone exposure on behavior, we included an additional group of mice that were born term

to dams injected with mifepristone on GD 18 (MFP Term, Figure 2A). Mice were trained on the

task in 2 daily sessions of 100 trials, where each session consisted of 50 randomized

presentations of 120º and 60º oriented gratings each. 120º was paired with the delivery of 10 µl

of water, while 60º had no consequence (Figure 2B and C). Licks of the spout were recorded and

quantified as Hits (licks to 120º), Misses (no licks to 120º), False Alarms (licks to 60º) and

Correct Rejections (no licks to 60°), which were then used to quantify the discriminability

[d'=z(Hits)-z(False Alarms)]. Mice were considered trained after reaching a d' of 2 or more for 3

sessions in a row, and they were trained for a maximum of 4000 trials (20 days of training) if

they did not cross the threshold within that timeframe. In our task, Hits remained high

throughout the training, while False Alarms gradually decreased (Figure 2D and Supplementary

Figure 1), driving an increase in d'. In term mice from both groups, d' increased in a linear

fashion and most mice crossed the threshold within 2 weeks of training (Figure 2E and 2F).

While preterm mice also displayed an increase in d', the learning trajectory was significantly

different from that of term born mice (LMM, effect of Animal Group p=0.009; Term vs Preterm

p=0.007, MFP Term vs Preterm p=0.02, Term vs MFP Term p=0.78; estimated marginal means

(EMM)±SE Term=2.06±0.16, MFP Term=2.01±0.18, Preterm=1.6±0.13; N=22 Term, 29 Preterm

and 17 MFP Term mice; Supplementary Table 1), and a fraction of them (17.24%) failed to reach

the criterion within the 4000 trial limit (Figure 2E and 2F; Kolmogorov-Smirnov test of d' values

from the last training session Term vs Preterm: D=0.43, p=0.009, Term vs MFP Term: D=0.34,

p=0.14, MFP Term vs Preterm: D=0.2893, p=0.23), confirming that preterm birth negatively

impacts learning. Naïve preterm mice had significantly lower Hit Rates at the onset of training,

which recovered as the training progressed (Figure 2G and Supplementary Figure 1; ANOVA

birth x training p=0.016, $F_{(2, 65)}$=4.37; Bonferroni post-hoc Naïve Term vs. Preterm p=0.005,

Term vs MFP Term p>0.9, MFP Term vs Preterm p=0.0013; all p>0.9 among Trained groups).

However, their False Alarm rates remained high throughout the training (Supplementary Figure

1) and were significantly higher at the end of the training compared to both groups of term mice

(Figure 2H; ANOVA birth x training p<0.0001, $F_{(2, 65)}$=10.66; Bonferroni post-hoc Naïve Term

vs. Preterm p>0.9, Term vs MFP Term p=0.13, MFP Term vs Preterm p=0.04; Trained Term vs

Preterm p=0.036, Term vs MFP Term p=0.53, MFP Term vs Preterm p=0.0007). To test if the

effect on the Hit Rates and False Alarms was driven by preterm mice that failed to reach the

criterion, we repeated the analyses while excluding those animals. We found that the interaction

between birth timing and training remained for both Hit Rates and False Alarms (ANOVA birth x

training HR p=0.04, $F_{(2, 60)}$=3.2; FA p=0.001, $F_{(2, 60)}$=7.7), demonstrating that preterm birth

impacts task performance regardless of whether the animal learned the task.

Our experiments included both male and female mice of sufficient numbers to segregate

the data by sex and test if birth has a sex-specific effect (Supplementary Figure 2). LMM with

Animal Group and Sex as fixed effects was fitted to the learning trajectories, which revealed that

sex has a significant impact on the learning trajectory (p=0.021), with female mice in all groups

reaching lower d' values (Supplementary Table 1). Analyses of Hits and False Alarm rates in

trained mice (Supplementary Figure 2B and C) revealed no effect of sex, while the effect of birth

was still present (ANOVA Hits: birth x sex p=0.48, $F_{(2,62)}$=0.74; sex p=0.42, birth p=0.02; False

Alarms: birth x sex p=0.12, $F_{(2,62)}$=2.2; sex p=0.38, birth p=0.03). Our results hence demonstrate

that preterm birth negatively impacts the learning trajectory of both sexes, as well as that female

mice from all groups reach lower d' values during training.

To test if the poor performance of preterm mice in the task is due to a deficit in

motivation, we compared the lick rates to the rewarded cue (binned in 20 trials) during the last

training session (Figure 2I). Lick frequencies in response to task cues gradually dropped during

the training session, which has been proposed as a measure of motivation [46]. An LMM was fitted

to the data using Animal Group and Trial Block as fixed effects, which confirmed that the

frequency of licks to the rewarded cue is reduced at the end of the session (Trial Block p<0.0001;

Supplementary Tables 2), and that lick frequencies do not differ significantly between animal

groups (p=0.15). Lick frequencies were different in naïve mice, with preterm mice showing less

licks in response to the rewarded cue (Supplementary Tables 3; Animal Group p=0.01).

However, as in trained mice, all groups showed a reduction in lick frequencies towards the end

of the first session (Trial Block p=0.04). Together, these results indicate that poor performance of

preterm mice (at least at the end of the training) is not due to deficits in motivation.

Our task is performed in head-fixed mice (on a treadmill), which may mask locomotion

deficits in preterm mice. To test this, we tested the locomotor activity of term and preterm mice

in an open field using an automated approach [38] (Figure 2J). ANOVA (Welch's) revealed a

significant and small reduction in locomotor activity of MFP term mice when compared to term

mice, while the level of locomotor activity in preterm mice was comparable to term mice

($F_{(2,56.98)}$=3.9, p=0.03; Dunnett's T3 multiple comparisons test Term vs. MFP Term p=0.03, Term

vs. Preterm p=0.83, MFP Term vs. Preterm p=0.41). While ANOVA found a significant

difference between groups of term mice in their preference for moving close to the walls of the

open field arena (Figure 2K, thigmotaxis; $F_{(2,51.62)}$=4.08, p=0.03), post hoc tests revealed only

marginal differences (Dunnett's T3 multiple comparisons test Term vs. MFP Term p=0.07, Term

vs. Preterm p=0.06, MFP Term vs. Preterm p=0.87), indicating largely preserved exploratory
activity in preterm mice.

Altogether, our results demonstrate that preterm mice display impaired learning in a
visual discrimination task, despite visual function, locomotor activity and motivation comparable
to term born mice. As learning impairments in mouse models of neurodevelopmental conditions
are often associated with impaired visual processing during task performance [47,48], we next
characterized the processing of task cues in the V1 of trained term and preterm mice while the
mice were engaged in the task.

Selectivity for task cues is weaker in putative pyramidal neurons in preterm V1

Visual discrimination training increases the representation of and responsiveness to
trained cues in the V1, both thought to support task acquisition [40,41,49–51]. As preterm mice display
impairments in task acquisition (Figure 2), we asked if the processing and representation of task
cues in the V1 of preterm mice are altered compared to term mice. To address this, we used *in*
vivo electrophysiology to isolate neuronal responses to task cues in trained mice while they were
engaged in the task (Figure 3A and B). Firing of neurons in the V1 of trained term and preterm
mice displayed the typical transient (early, 100 ms after the onset of lick) and sustained (late)
components, where the peak of the sustained component typically preceded the onset of licks
(Figure 3C). To test the relationship between V1 spiking and licks, we computed event
correlations in 1 ms time bins for all isolated units while using licks as the reference (Figure 3D).
As trained mice lick to the rewarded cue in bouts (Figure 2D), with 100-200 ms on average
between licks [52], we restricted the correlation window to 200 ms. We found that the bulk of V1
unit firing occurs prior to lick events, with a peak at 55 ms and a significantly higher number of
spiking events preceding the licks (Figure 2D; Wilcoxon matched-pairs signed rank test,
p=0.018, W=-1354).

To compare the representation of task cues between term and preterm V1, we calculated
the percentiles of neurons (per mouse) whose activity was significantly modulated by task cues
(as determined by Wilcoxon signed-rank test of pre- and postcue firing rates). We calculated the
direction of the modulation as the selectivity index (SI): $(R_{cue}-R_{baseline})/(R_{cue}+R_{baseline})$, with
positive values indicating cue-evoked increase in firing and negative values indicating cue-
evoked suppression. We further split the units into broad spiking (putative pyramidal neurons)

and narrow spiking (putative fast-spiking inhibitory interneurons) based on the width of their

action potential waveform (Supplementary Figure 3; 406 neurons from 12 term mice and 253

neurons from 9 preterm mice) [53].

Both regular-spiking (RS) and fast-spiking (FS) neurons in term and preterm mice had

comparable representation of task cues in the V1 with a slight overrepresentation of the rewarded

cue (RM ANOVA with birth and the direction of modulation as factors; Figure 3E: RS Reward:

birth x modulation p=0.93, $F_{(2,38)}$=0.07; RS No Reward: birth x modulation p=0.7, F=0.36;

Figure 3F: 2-tailed t-test, p=0.9, t=0.0046, df=19; Figure 3G: FS Reward: birth x modulation

p=0.95, F=0.09; FS No Reward: birth x modulation p=0.24, F=1.46; Figure 3H: p=0.66, t=0.45).

However, when we compared the peristimulus time histograms (PSTHs) of neurons whose firing

was significantly modulated by task cues (% of all RS neurons: $Term_R$=54.2, $Preterm_R$=59.59,

$Term_{NR}$=42.05, $Preterm_{NR}$=44.89; % of all FS neurons: $Term_R$=50.67, $Preterm_R$=78.14,

$Term_{NR}$=36.7, $Preterm_{NR}$=54.96), we found starkly elevated firing in preterm mice [Figures 3I-L;

R and NR: p=0.000, permutation test of PSTH area under the curve (AUC)].

To test the magnitude and the direction of neuronal responses to task cues, we measured

selectivity index (SI) as $(R_{cue}-R_{baseline})/(R_{cue}+R_{baseline})$, where R_{cue} is the firing rate during cure

presentation. SI ranges from -1 to 1, where the indices of suppressed neurons fall below 0, and

those of excited neurons above it. As the distribution of SI values in a population of neurons can

provide more information about cue evoked activity than population averages, we compared the

cumulative distributions of SI values from neurons significantly modulated by cues in term and

preterm mice. Despite increased firing, RS neurons in preterm V1 had reduced selectivity for

task cues, evident in the distribution of selectivity indices (SI) shifted towards lower values

(Figure 3M; SI median±MAD $Term_R$=0.34±0.12, $Preterm_R$=0.19±0.09, K-S test: D=0.35,

p=0.0007; $Term_{NR}$=0.21±0.1, $Preterm_{NR}$=0.08±0.04, K-S test, D=0.46, p=0.00005). On the other

hand, a large fraction of FS neurons in preterm mice had high SI values, with no changes in the

selectivity for the non-rewarded cue (Figure 3N; $Term_R$=0.36±0.13, $Preterm_R$=0.43±0.23, K-S

test: D=0.29, p=0.000006; $Term_{NR}$=0.17±0.1, $Preterm_{NR}$=0.13±0.09, D=0.11, p=0.36).

As locomotion can increase V1 firing rates [54], we measured the amount of movement on

the treadmill during the task (Supplementary Figure 4A, N=8 term and 5 preterm mice).

Locomotion during the task was very variable in both groups of mice and overall not statistically

different (p=0.37, permutation test of AUC). Elevated firing in the V1 was task-specific, as

PTSHs revealed similar levels of activity in both groups of mice outside of task context (cues

presented in the absence of the reward delivery spout, Supplementary Figure 4B, compare with

Figure 3G). Further, licking activity in preterm mice was slightly elevated (Supplementary

Figure 4B; # of licks per bout $Term_R$=23.65±2.23, $Preterm_R$=28.48±1.49, $Term_{NR}$=8.37±1.28,

$Preterm_{NR}$=13.31±3.056; birth x cue p=0.97, $F_{(1,\ 22)}$=0.00065; birth p=0.024, F=5.85; cue

p<0.0001, F=46.26), but the level of cue-evoked sustained activity in the V1 [mean sustained

response (0.1-1.5 s)/mean transient response (0-0.1s) [55]] was not significantly correlated with

lick frequency in either group of mice (Supplementary Figure 4E; Pearson's correlation, $Term_{RS}$

p=0.97, $Term_{FS}$ p=0.22, $Preterm_{RS}$ p=0.28, $Preterm_{FS}$ p=0.08).

Neurons in the V1 display activity related to behavioral outcomes [56]. While the

behavioral relevance of this activity is unclear, we checked if this activity can be detected in the

V1 of preterm mice. Trained mice typically had a 100% hit rate during the last session

(Supplementary Figure 1), so we compared the responses of RS and FS neurons during the

Correct Rejections (CR) and False Alarms (FA) between term and preterm mice. RS neurons

represented outcomes to a similar degree in term and preterm mice (Supplementary Figure 5A;

ANOVA cue x modulation: Correct Rejection p=0.24, $F_{(2,38)}$=1.47; False Alarms p=0.7, F=0.36),

despite elevated neuronal activity in preterm mice and weakened SIs for non-rewarded cue

during both outcomes (Supplementary Figure 5B and C; p=0.000, permutation test of PSTH

AUC; SI $Term_{CR}$=0.2±0.09, $Preterm_{CR}$=0.09±0.06; K-S test, D=0.48, p=0.0002;

$Term_{FA}$=0.29±0.09, $Preterm_{FA}$=0.09±0.25; D=0.37, p=0.02). While FS neurons of term and

preterm mice represented the behavioral outcomes to a comparable degree (Supplementary

Figure 5D; ANOVA cue x modulation: Correct Rejection p=0.36, F=1.03; False Alarms p=0.75,

F=0.28), neuronal activity was elevated in preterm mice and the cue-evoked peak in activity was

barely detectable during the False Alarms (Supplementary Figures 5E and F; p=0.000,

permutation test of PSTH AUC). Interestingly, selectivity for the non-rewarded cue in FS

neurons displayed a significant shift towards lower values once it was segregated into behavioral

outcomes; SI $Term_{CR}$=0.26±0.12, $Preterm_{CR}$=0.17±0.11; K-S test, D=0.23, p=0.009;

$Term_{FA}$=0.36±0.18, $Preterm_{FA}$=0.23±0.32; D=0.31, p=0.001).

Altogether, our data demonstrates that preterm mice have intact representation of task

cues and behavioral outcomes in the V1, despite significantly elevated neuronal activity and

overall weakened selectivity for the task cues. Increase in neuronal activity in preterm mice

appeared to be specific to task context, suggesting the involvement of top-down modulation.

Therefore, we next examined the neuronal responses to task cues in term and preterm prefrontal

cortex (PFC), a top-down area that sends major glutamatergic monosynaptic input to the V1 and

modulates its activity during discrimination tasks [57,58].

Representation of task cues is altered in the PFC of preterm mice

Within the PFC, anterior cingulate cortex (ACC) sends and receives dense glutamatergic

projections to and from the V1 [57,59,60]. Neurons in the ACC are not responsive to sensory cues in

naïve mice, but develop robust context-specific responses to cues during training in sensory

discrimination tasks [61–63]. Further, optogenetic activation of ACC neurons that project to the V1

(ACC→V1) evokes a robust increase in V1 firing rates and promotes behavioral performance in

visual tasks [57,58], while inactivating ACC impairs visual discrimination (Supplementary Figure 6)

[64], suggesting that ACC exerts top-down modulation of cue processing in the V1 of trained mice.

We therefore hypothesized that elevated activity and altered cue selectivity in preterm V1 during

task performance reflects the activity and selectivity in their PFC. To test this, we recorded the

activity of the PFC while the mice were engaged in the task (Figure 4A and B) and sorted the

isolated neurons by their waveform width as in the V1. While most of the units recorded were

from deep cortical layers (5 and 6), our recordings collected units from layer 2/3 as well (Figure

4B). As expected, putative FS interneurons had higher precue firing rates overall in both groups

of mice (median±MAD FS: Term=8.61±6.97, Preterm=7.57±5.75 spikes/s 143 and 183 units

respectively from 10 and 9 preterm mice; RS: Term=2.7±1.08, Preterm=2.46±1.2 spikes/s, 70

and 170 units respectively), but pre-cue firing rates between term and preterm mice were

comparable (Mann-Whitney t-test, FS: p=0.6, U=43; RS: p=0.96, U=25). Similar to V1, the

firing of PFC neurons preceded the licks, with event correlogram revealing peaks at 14 and 65

ms prior to lick events, as well as a significantly higher number of spiking events prior to lick

events (Figure 4C; Wilcoxon matched-pairs signed rank test, p<0.0001, W=-2770).

We next evaluated cue representation and found that the non-rewarded cue had a

significantly weaker representation in RS neurons of preterm mice, shifting the overall

representation in favor of the rewarded cue (Figure 4D; ANOVA Reward: birth x modulation,

p=0.6, $F_{(2,34)}$=0.52; No Reward: p=0.014, F=4.88; Figure 4E: Mann-Whitney test p=0.045,

U=20.5). Significantly modulated neurons in both groups of mice had an overall higher firing

rate than the entire recorded population (median±MAD; precue firing Term=4.3±2.84,

Preterm=3.27±1.73 spikes/s; % of all neurons $Term_R$=78.7, $Term_{NR}$=52.9, $Preterm_R$=88.1,

$Preterm_{NR}$=23.01), and their PSTHs revealed a significantly increased pre- and postcue firing

rate in preterm mice (Figure 4F and G; p=0.000, permutation test of PSTH AUC). Additionally,

the PSTH to the non-rewarded cue in preterm mice had a very irregular shape with no clear peak,

suggesting weakened selectivity of prefrontal RS neurons for the non-rewarded cue in preterm

mice (Figure 4G). Indeed, while the SI values for the rewarded cue were not different between

term and preterm mice (Figure 4H: median±MAD, $Term_R$=0.72±0.13, $Preterm_R$=0.71±0.1; K-S

test, D=0.13, p=0.36), the SI values for the non-rewarded cue were significantly lower in preterm

mice (Figure 4H: $Term_{NR}$=0.33±0.09, $Preterm_{NR}$=0.16±0.07; K-S test, D=0.4, p=0.000014).

Previous work demonstrated a strong correlation between prefrontal responsivity to task

cues during training and the progressive increase in performance [61]. As the SI reflects the

responsiveness of neurons to task cues, we tested the correlation between the SI values (averaged

per animal) and d' values during the recording session, as well as the strength of the SI values in

predicting behavioral performance using linear regression when a significant correlation was

detected. In term mice, SI values for the rewarded and non-rewarded cues were not significantly

correlated with d' values (Pearson's correlation, Figure 4I). In contrast, SI values for the

rewarded cue were significantly negatively correlated with d' in preterm mice (r=-0.75, p=0.02)

and were a predictor of performance (Figure 4J and Supplementary Tables 4). The correlation

between SI values for the non-rewarded cue and d' in preterm mice was not significant (Figure

4J).

Our bulk recordings did not isolate the activity of PFC neurons that project to the V1,

making it difficult to draw connections between the activity patterns in the V1 and PFC. To do

that, we used an intersectional viral approach to optogenetically label the V1-projecting PFC

neurons, where we injected the retrograde AAV2 variant encoding the Cre-recombinase into the

V1 [65] and DIO-ChR2-mCherry into the PFC (Supplementary Figure 7A and B). We used 10 ms

pulses at 473 nm after the behavioral sessions ended to elicit the ChR2-mediated responses and

isolate the activity of PFC→V1 neurons (short latency responses with >50% reliability,

Supplementary Figure 7C) [66]. Interestingly, we found a significantly increased representation of

the rewarded cue in PFC→V1 neurons of preterm mice (Supplementary Figure 7D; 28 neurons

from N=9 term mice, 35 neurons from N=6 preterm mice; ANOVA; Reward: birth x modulation,

p=0.0017, $F_{(2,26)}$=8.27, No Reward: birth x modulation, p=0.38, F=0.99). PSTHs of neurons

significantly modulated by the cue (% of all neurons: $Term_R$=66.67, $Preterm_R$=98.33,

$Term_{NR}$=55.21, $Preterm_{NR}$=58.45) revealed elevated activity of PFC→V1 neurons in preterm

mice after the presentation of the rewarded cue, especially in the late stages of their response

(Supplementary Figure 7F; p=0.000, permutation test of AUCs). SI values for the rewarded cue

were also substantially higher in preterm than in term mice (median±MAD, $Term_R$=0.31±0.31,

$Preterm_R$=0.57±0.09; K-S test, D=0.79, p=3.15x10^{-6}). PSTHs of PFC→V1 neuron responses to

the non-rewarded cue revealed irregular firing in preterm mice, in line with weakened activity

evoked by the non-rewarded cue in V1 and PFC (Supplementary Figure 7G; p=0.000,

permutation test of AUCs). The SI values for the non-rewarded cue were also elevated in preterm

mice ($Term_{NR}$=0.11±0.19, $Preterm_{NR}$=0.2±0.07), but the differences in their distribution between

term and preterm mice were marginally significant (K-S test, D=0.41, p=0.06), likely due to low

number of significantly modulated neurons (Term=16 and Preterm=23).

The representation of cues in FS neurons was similar to RS neurons, although the

difference in the representation of the non-rewarded cue was not statistically significant (Figure

4K; ANOVA Reward: birth x modulation, p=0.71, $F_{(2,34)}$=0.34; No Reward: p=0.2, F=1.67). As in

RS neurons, the representation of cues was slightly shifted in favor of the rewarded cue (Figure

4L), but it was not statistically different between term and preterm mice (Mann-Whitney test

p=0.17, U=28). However, when we compared the PSTHs of neurons significantly modulated by

task cues between term and preterm mice (% of all neurons $Term_R$=86.42, $Term_{NR}$=56.37,

$Preterm_R$=90.8, $Preterm_{NR}$=37.6), we found that the responses to both cues in preterm mice were

significantly blunted (Figure 4M and N; p=0.000, permutation test of PSTH AUC). Similar to V1

FS neurons, the median SI of prefrontal FS neurons for the rewarded cue in preterm mice was

marginally larger than in term mice, while the SI for the non-rewarded cue was lower (Figure

4O; median±MAD SI: $Term_R$=0.52±0.17, $Preterm_R$=0.58±0.11, K-S test: D=0.19, p=0.02;

$Term_{NR}$=0.21±0.06, $Preterm_{NR}$=0.092±0.07, K-S test, D=0.36, p=0.00097). None of the SI values

in either group of mice correlated with d' (Figures 4P and Q; Pearson's correlation).

FS interneurons can remain undetected during spike sorting of bulk electrophysiological recordings due to their low spontaneous firing rates [67]. To ensure that our spike sorting approach reliably captured the weaker responsivity to task cues in preterm mice, we used a transcriptional enhancer S5E2 [68] to transduce FS interneurons with ChR2 for optogenetic identification ("optotagging", Supplementary Figure 8). S5E2-mCherry only partially colocalized with Parvalbumin (PV) signal in the ACC, which was significantly weaker compared to other cortical areas (Supplementary Figure 8B), as previously reported [69]. As expected, 10 ms pulses of blue light evoked strong rhythmic firing in transduced neurons (Supplementary Figure 8C) [66], with all isolated neurons having a narrow waveform (average spike half-width: $0.33 \pm 3.2 \times 10^{-6}$ ms, N=50 units from 5 term mice and 52 units from 4 preterm mice). Indeed, the pre-cue firing rate of optotagged FS units was significantly lower than that of FS units isolated through spike sorting of bulk electrophysiological data (Figure 4B and 4C; median±MAD, N=units; Term/Preterm narrow-spiking units=5.9±4.79 spikes/s vs Term/Preterm optotagged $S5E2^+$ units=1.45±1.45 spikes/s; Mann-Whitney t-test, p<0.0001, U=11297), with no significant differences between term and preterm optotagged FS units (Term=0.63±0.63 spikes/s, Preterm=1.69±1.52; Mann-Whitney t-test, p=0.18, U=1104). While the fractions of neurons significantly modulated by task cues was overall lower in the optotagged population (Supplementary Figure 8D; compare with 4K), they were not significantly different between term and preterm mice (ANOVA Reward: birth x modulation: p=0.91, $F_{(2\ 14)}$=0.094; No Reward: p=0.99, F=0.014). PSTHs of significantly modulated optotagged neurons (% of all neurons: $Term_R$=61.56, $Preterm_R$=71.18, $Term_{NR}$=52.66, $Preterm_{NR}$=53.26) revealed their weak responsivity to task cues in preterm mice (p=0.000, permutation test of PSTH AUC), similar to what we found in FS neurons isolated through bulk recordings (Supplementary Figure 8E and F; compare with Figure 4M and N). However, the selectivity for the rewarded cue was significantly weakened in preterm mice, while the selectivity for the non-rewarded cue appeared comparable between term and preterm mice (median±MAD SI: $Term_R$=0.78±0.22, $Preterm_R$=0.38±0.15, K-S test: D=0.79, p=0.000003; $Term_{NR}$=0.31±0.06, $Preterm_{NR}$=0.35±0.13, K-S test: D=0.41, p=0.057).

Our results hence demonstrate a significant shift in the representation of task cues towards the rewarded one in putative principal neurons of preterm PFC, along with pronounced disinhibition of their responses to task cues. This shift comes at the expense of the non-rewarded cue, which elicits blunted responses in both putative pyramidal neurons and interneurons.

Surprisingly, the selectivity of the population responses to the rewarded cue was predictive of

performance in preterm, but not in term mice. As the preterm mice that underwent

electrophysiological recordings had comparable d' values to term mice (Figure 4I and J;

mean±SEM Term=2.56±0.16, Preterm=2.83±0.13, N=9 mice/group), our results indicate

divergent processing of task cues in preterm mice and further suggest that preterm mice use

different strategies to encode behavioral outcomes. To test this further, we next compared the

representation of behavioral outcomes to the non-rewarded cue in prefrontal RS and FS neurons

of term and preterm mice and modeled their ability to classify trial-by-trial behavioral outcomes.

Cue-evoked activity of prefrontal neurons is a weaker predictor of behavioral outcomes

in preterm mice

Prefrontal cortex is critical for many forms of goal-directed behavior and its activity is

predictive of behavioral outcomes across different tasks [70–72]. Surprisingly, the population

selectivity for the rewarded cue was predictive of performance in preterm but not in term mice

(Figure 4), suggesting that preterm mice use a strategy different from term mice to encode task-

relevant information. Indeed, both the increase in licks to the rewarded cue and suppression of

licks to the non-rewarded cue drive the increase in performance in preterm mice, while the

suppression of behavioral responses to the non-rewarded cue drives the performance increase in

term mice (Figures 2 and Supplementary Figure 1). We therefore hypothesized that the PFC of

term and preterm mice uses divergent strategies to encode behavioral responses to task cues, and

to the non-rewarded cue in particular as preterm mice show increased False Alarm rates (Figure

2H). As almost 100% of behavioral responses to the rewarded cue in trained mice from both

groups are Hits (Figure 2 and Supplementary Figure 1), we focused our analysis on neuronal

responses to the non-rewarded cue, i.e. Correct Rejections (CRs) and False Alarms (FAs; Figure

2C).

We first tested the representation of behavioral outcomes in prefrontal neurons and

surprisingly found no differences between term and preterm mice in either regular spiking

(Figure 5A; ANOVA: birth x modulation CR: p=0.81, $F_{(2,34)=}0.21$, FA: p=0.6, F=0.51) or fast-

spiking neurons (Figure 5B; CR: p=0.65, F=0.43, FA: p=0.45, F=0.82). PSTHs of significantly

modulated RS neurons (% of all neurons $Term_{CR}$=33.5, $Term_{FA}$=41.42, $Preterm_{CR}$=35,

$Preterm_{FA}$=49.03) revealed elevated firing and irregular cue-evoked activity in preterm mice

during both outcomes (Figure 5C; p=0.000, permutation test of AUCs). While cue-evoked

activity in preterm FS neurons also appeared irregular, analysis revealed that term and preterm

FS responses to the non-rewarded cue are comparable during CR but not during FA trials (Figure

5D, CR p=0.125, FA p=0.022, permutation test of AUCs; % of all neurons $Term_{CR}$=48.53,

$Term_{FA}$=53.34, $Preterm_{CR}$=46.34, $Preterm_{FA}$=56.94). Selectivity indices were shifted towards

lower values in both cell populations of preterm mice during both behavioral outcomes (Figure

5E; median±MAD, K-S test for all; RS CR Term=0.49±0.21, Preterm=0.13±0.09, D = 0.44,

p=0.000042; RS FA Term=0.4±0.16, Preterm=0.29±0.18, D = 0.26, p=0.014; FS CR

Term=0.14±0.07, Preterm=0.09±0.06, D=0.24, p=0.044; FA Term=0.37±0.11,

Preterm=0.08±0.19, D=0.57, p=$1.2x10^{-10}$). Further, FS neurons in term mice showed higher SI

values during FA trials, while the same was true for RS neurons in preterm mice (Mann-Whitney

test, Term FA vs CR: RS p=0.82, U=150, FS p<0.0001, U=526.5; Preterm FA vs CR: RS

p<0.0001, U=1593, FS p=0.7, U=2517).

To test if neuronal responses can predict behavioral outcomes on a single trial basis, we

used logistic regression with CR and FA as binary outcomes and cue-evoked (baseline

subtracted) spike rates (Figure 5F and G) as classifiers. We included all isolated neurons in the

analysis as the number of neurons significantly modulated by the non-rewarded cue was low for

some mice. Cue evoked spike rate distributions of all isolated neurons mirrored the trends seen in

SI values of neurons significantly modulated by task cues, with FS and RS neurons in term and

preterm mice, respectively, preferentially responding to the non-rewarded cue during the FA

trials and weaker cue-evoked firing in preterm FS neurons during both trial outcomes (Figure

5K; median±MAD spikes/s, RS: $Term_{CR}$=1.5±1.43, $Preterm_{CR}$=0.65±1, K-S test p=0.011,

D=0.23; $Term_{FA}$=2±1.15, $Preterm_{FA}$=2.3±3.31, p=0.0008, D=0.28; FS: $Term_{CR}$=1.28±1.27,

$Preterm_{CR}$=0.34±0.89, p=0.0001, D=0.26; $Term_{FA}$=4.41±4.29, $Preterm_{FA}$=1.25±1.86, p=0.93x10^{-}

^{6}, D=0.31). We constructed Receiver Operating Characteristic (ROC) curves for each group of

mice and compared their performance first to shuffled outcomes from each group, and then to

each other (Figure 5H) [73]. We found that neuronal responses to the non-rewarded cue in both

groups of mice classified trial outcomes significantly better than shuffled data (au(ROC)±SE:

Term=0.703±0.0058, 7928 trials; Preterm=0.65±0.005, 11164 trials; Shuffled term=0.5±0.006,

Shuffled preterm=0.5±0.005; Term vs Shuffled term, z=23.26, p<0.0001, Preterm vs shuffled

preterm, z=19.93, p<0.0001). However, firing rates of term mice performed significantly better

as classifiers than those of preterm mice (z=6.79, p<0.0001). ROC values for each mouse did not

correlate with the fraction of significantly modulated neurons nor with their SI values (Pearson's

r=-0.24, p=0.42 and r=0.25, p=0.4, respectively), indicating that including all recorded neurons

in the model did not impair the classification.

Our results hence provide further evidence for weak selectivity of prefrontal neurons for

the non-rewarded cue in preterm mice, indicating divergent processing of task cues and encoding

of behavioral outcomes in preterm PFC. Preterm birth is a significant risk for delays in cognitive

development, especially in the development of response inhibition [74], suggesting that the

impairments in the processing and representation of task cues in preterm PFC reflect a

maturational deficit. As prior research found weak representation of non-rewarded cue in the

brains of adolescent mice trained on an auditory task [75], we next tested if the representation and

processing of cues in our task is similar between adult preterm mice and adolescent term mice.

Similar encoding of behavioral outcomes to the non-rewarded cue in adolescent term and

adult preterm mice

To test for impaired maturation of preterm PFC, we examined the representation and

processing of task cues in adolescent, term born male and female mice and compared it to adult

preterm mice (Supplementary Figure 9). To prepare the adolescent mice for the task, they were

implanted with a headpost between postnatal days 21 and 24 and begun habituation and water

restriction 4 days after the implantation (Figure 6A). The training started between postnatal days

32 and 35 and lasted until mice crossed the criterion (all mice learned the task, N=18).

Adolescent mice performed comparable to term mice (6B, LMM with Age as Fixed Factor and

Animal and Training Day as Random Factors, p=0.42, Supplementary Tables 5), and showed a

significant increase reduction in FA rate between naïve and trained stages (Figure 6C; RM

ANOVA: response x training p<0.0001, $F_{(1,17)}$=40.5; HR p=0.07, FA p<0.0001), indicating that a

decrease in response inhibition drives behavioral performance in our task regardless of age

tested.

We next recorded the activity of prefrontal neurons in adolescent mice during task

performance (N=324 neurons from 9 mice), which we classified into regular and fast-spiking

neurons as in Figures 3 and 4 (Figure 6D and E). As previously reported [75], the representation of

the non-rewarded cue was significantly weaker than that of the rewarded cue (RM ANOVA cue x

modulation Figure 6E: p<0.0001, $F_{(2.2,17.79)}$=37.11; Figure 6F: p<0.0001, $F_{(2.5,20.04)}$=18.37).

Similar to preterm mice, the representation of the non-rewarded cue was significantly weaker

only in RS neurons when compared to adult term mice (Supplementary Figure 9). While the

rewarded cue elicited substantially higher activity in both RS and FS neurons (Figure 6G;

p=0.000, permutation test of AUCs), the PSTHs of neurons whose activity was significantly

modulated by task cues revealed that cue-evoked activity was lower in adolescents than in adult

term mice (% of all neurons: RS R=72.98, NR=29.42, FS R=93.11, NR=35; Supplementary

Figure 9). Preference for the rewarded cue was also evident in SI values, with higher selectivity

for the rewarded cue in both RS and FS of adolescent mice (Figure 6H; median±MAD RS

R=0.4±0.16, NR=0.25±0.16; K-S test, D=0.31, p=0.005; FS R=0.41±0.12, NR=0.26±0.18;

D=0.27, p=0.01).

In terms of behavioral outcomes to the non-rewarded cue, both CR and FA were equally

represented in the adolescent PFC (RM ANOVA outcome x modulation Figure 6E: p=0.17,

$F_{(2.16)}$=1.96; Figure 6F: p=0.95, F=0.04). PSTHs of neurons significantly modulated by the non-

rewarded cue revealed slightly higher neuronal responses during FA trials in both neuronal

populations (Figure 6I; p=0.000, permutation test of AUCs; % of all neurons: RS CR=15.71,

FA=34.03, FS CR=41.73 FA=44.75), but their SI values were comparable (Figure 6J;

median±MAD RS CR=0.28±0.16, FA=0.3±0.22; K-S test, D=0.21, p=0.47; FS CR=0.25±0.16,

FA=0.29±0.21; D=0.12, p=0.89). We next isolated single trial firing for adolescent neurons and

tested if their evoked firing can accurately classify the two behavioral outcomes to the non-

rewarded cue (Figure 6K; term and preterm data from Figure 5M added for comparison). While

the logistic model built on adolescent evoked firing rates performed significantly better than the

shuffled data (p<0.0001, z= 22.92), there was no significant difference between the auROC of

preterm mice and that of adolescent mice (adolescent vs preterm p=0.17, z=1.38; adolescent vs

term adult p<0.0001, z=-5.76).

Life-long environmental enrichment fails to improve the learning trajectory of preterm mice

Adult preterm mice and adolescent term mice display similarly weak representation of

the non-rewarded cue in the PFC (Supplementary Figure 10), as well as similar encoding of its

behavioral outcomes (Figure 6K), suggesting a maturational delay in the PFC of preterm mice.

We therefore tested if environmental enrichment (ENR), a classical paradigm that promotes

sensory maturation and rescues cognitive deficits in animal models of prematurity-related brain

injury, improves learning in preterm mice [21,28,31,76,77]. To do this, we housed timed-pregnant dams

with their term and preterm litters from postnatal day 5 (P5) to weaning (P28) in standard rat

cages supplemented with a running wheel, multiple play and chew objects (such as gnawing

sticks and tunnels), as well as a non-related dam with an age-matched litter for social

enrichment. The object location was shuffled in the cage every 2-3 days. Prior to P5, pups were

housed in a standard cage with enrichment for the dams in the form of igloos, extra nesting

material and sunflower seeds. After weaning, term and preterm mice were housed in groups of 2-

3 in standard cages supplemented with an igloo, small running platform and gnawing sticks.

When mice reached adulthood (>2 months of age), they were prepared for the visual

discrimination task, water-restricted and trained to criterion or to 4000 trials if they did not pass

the criterion (Figure 7A). While term mice had a steep learning curve, the enrichment had a

negative impact of their performance, lowering the overall d' values when compared to term

mice that received no enrichment [LMM with Enrichment (ENR) and Birth as Fixed Effects, and

Animal and Training Day as Random Effects; ENR p<0.0001, Estimated Marginal Means

(EMM) Term ENR=1.6±0.2, Term=2.22±0.19; Supplementary Tables 6]. However, just like term

mice that received no enrichment, all ENR mice learned the task within 15 days (Figure 7B). In

contrast, preterm mice that received ENR had a substantially impaired learning curve (LMM:

Birth p<0.0001) and performed worse than preterm mice that received no enrichment (EMM

Preterm ENR=1.003±0.68, Preterm=1.6±0.14). While a significant fraction of ENR preterm mice

failed to learn the task within 4000 trials (27.27%), the distribution of final d' values was not

significantly different between ENR term and preterm mice (Figure 7C; K-S test, D=0.46,

p=0.16).

When we analyzed Hit Rates and False Alarms in ENR mice and compared them to

standardly reared term and preterm mice (STD; Figure 7D and E, compare with Figure 2G and

H), we found a significant interaction between training, enrichment and birth timing (RM

ANOVA, p=0.002, F=28.49, Supplementary Tables 7). Post-hoc tests revealed no significant

differences between naïve ENR term and preterm mice in Hit Rates and a significant reduction in

trained preterm mice (naïve ENR Term vs Preterm p=0.19, trained p=0.006; Figure 7D). Both

groups of mice had significantly reduced HR compared to STD mice at the onset of training, and

this trend persisted in preterm mice only (Supplementary Tables 7). There were no significant

differences in FA rates between term and preterm mice raised in ENR (Figure 7E; naïve ENR

Term vs Preterm p=0.99, trained p=0.99; Supplementary Tables 7), but trained ENR preterm

mice had comparable FA rates to STD trained term mice (p=0.99).

Discussion

Through extensive electrophysiological characterization of primary visual (V1) and

prefrontal cortices in adult preterm mice trained on a visual associative task, our study identified

impaired representation and processing of sensory cues in the prefrontal cortex (PFC) as a

potential driver of cognitive impairments common in preterm born children and adolescents.

Diminished representation of the non-rewarded cue was present in putative pyramidal neurons of

the preterm PFC, while both putative pyramidal neurons and interneurons displayed reduced

selectivity for the non-rewarded cue. Activity evoked by the non-rewarded cue more accurately

predicted behavioral outcomes in term mice, pointing to their divergent encoding in term and

preterm PFC. Indeed, our study revealed substantial similarities in the representation of the non-

rewarded cue and encoding of its behavioral outcomes between adult preterm mice and term

adolescent mice, indicating that preterm birth disrupts the maturational trajectory of the PFC.

Environmental enrichment, however, did not improve the learning deficit of preterm mice.

Unexpectedly, enrichment impaired learning in both term and preterm mice, highlighting

previous findings on the variability of its outcomes in laboratory animals [78,79].

Our study identified substantial hyperactivity of cue-modulated neurons in the PFC and

V1 of preterm mice. In the PFC, the hyperactivity was specific to regular-spiking (RS), putative

pyramidal neurons, while the response of prefrontal fast-spiking (FS) interneurons to both cues

was blunted in preterm mice. Our findings hence suggest that elevated activity of prefrontal

pyramidal neurons is a consequence of dysfunctional perisomatic inhibition, but do not exclude a

dysfunction of other interneuronal subtypes: task-related neurons in preterm PFC show elevated

firing even before the onset of cue presentation, when FS interneurons do not show reduced

firing rates. Indeed, abnormalities in multiple interneuron subtypes have been reported in

postmortem studies of prematurely born fetuses and in animal models of prematurity [17,19,20,80].

Future electrophysiological studies can now test the function of other interneuronal subtypes in

the preterm PFC.

While cue-evoked activity of FS interneurons in preterm mice may contribute to the

elevated firing of prefrontal RS neurons, it is unclear which inputs fail to sufficiently activate the

FS interneurons upon cue presentation. These inputs are unlikely to come from local pyramidal

neurons-their responses to the rewarded cue in preterm mice are robust. Another major source of

excitation to prefrontal FS interneurons is the mediodorsal thalamus (MDT), whose role in

learning is well established [81–83]. Optogenetic activation of MDT results in a reduction of

putative pyramidal neuron firing rates due to strong feedforward inhibition mediated by MDT

inputs onto FS interneurons [84,85]. Our results point to these inputs being dysfunctional in preterm

mice, resulting in elevated activity of prefrontal pyramidal neurons during cue presentation. This

notion is in line with impairments in thalamic structure observed in preterm infants and

adolescents [13,14] and the hypothesized role of thalamic dysfunction as a driver of impaired

cortical function after preterm birth [86]. In sensory areas of mice, preterm birth accelerates the

refinement of thalamocortical inputs through serotonin signaling [87], but it is unclear if

prematurity has the same effect on higher order thalamic nuclei, such as MDT, whose cortical

target is the PFC. We plan to address this in our future studies.

We previously found that juvenile preterm mice show sparser neuronal firing rates and

increased size of putative inhibitory synapses in the V1 compared to term-born mice [39]. Our

current results indicate that these changes in the activity of V1 in preterm mice are transient,

given the comparable neuronal firing rates of V1 neurons outside of task context in term and

preterm mice. Furthermore, the baseline orientation selectivity is intact in preterm mice and the

firing of V1 neurons is elevated only during task performance, indicating that visual processing

deficits common after preterm birth are due to top-down modulation and not local V1

computations [88,89]. A previous study demonstrated that optogenetic activation of V1-projecting

prefrontal neurons results in robust elevation of V1 firing [58]. Indeed, our results found increased

selectivity for the rewarded cue in V1-projecting PFC neurons, as well as their heightened

activity, supporting that the increase in V1 neuron firing during task performance originates from

top-down projections. Our finding that both RS and FS neurons show similarly elevated task-

related activity further supports the involvement of top-down inputs in altered processing of task

cues in the V1 of preterm mice and warrant further investigation, especially given the previous

reports of top-down sensory processing impairments in preterm infants [90–92]. The firing rates of

V1 neurons in our study are somewhat higher than those previously reported in literature [53] and it

is possible our analysis failed to consistently separate single units from multiunits. On the other

hand, extracellular electrophysiology is biased towards neurons that fire action potentials at

higher frequencies (such as layer 5 pyramidal neurons) as they are easier to isolate from the

background [93,94]. Indeed, this is demonstrated in our study by comparison of optogenetically

identified FS neurons in the PFC and those identified through spike sorting of bulk signals where

we found that optotagged FS neurons show lower median firing rates before cue onset. As our

unit isolation criteria was kept constant for all groups of mice in this study, the presence of

multiunit data is unlikely to bias the results towards elevated firing in preterm mice, especially as

their prefrontal FS neurons display weak cue-evoked firing.

While we have not addressed the precise circuit mechanisms that drive behavioral

impairments and altered processing of task cues in preterm mice in this study, we attempted to

test for the presence of developmental delay in the preterm PFC as they are common in the

preterm population [95,96]. Interestingly, adolescent term-born mice show similarly reduced

representation of the non-rewarded cue in putative prefrontal pyramidal neurons, and their

neurons classify its behavioral outcomes comparable to preterm mice. As the preterm mice used

in our study were adults that ranged in age from 2 to 7 months, our findings indicate that

preterm-birth associated impairments in prefrontal function persist through adulthood, arguing

for a disrupted (instead of arrested) development. This argument is strengthened by our findings

that environmental enrichment, a commonly used developmental intervention that rescues

arrested sensory maturation [77], is ineffective in mitigating the deficits seen in preterm mice. In

fact, enrichment impaired performance in both term and preterm mice, highlighting the

variability in its effects on behavior, particularly in mouse models of neurological conditions

[79,97]. Previous work demonstrated that enrichment negatively impacts motivation in certain tasks

[98]. Indeed, ENR mice in our study had significantly reduced Hit Rates and False Alarms,

suggesting that their motor output was reduced, impairing their performance. Future studies can

test if the type of enrichment used in our study is optimal for promoting the maturation of

prefrontal areas, especially given recent studies that demonstrated the positive effects of tactile

enrichment on sensory information coding in the somatosensory and motor areas of mice [30].

An important finding of our study is that adolescent PFC encodes task-relevant

information differently to adult PFC. Our results confirmed a previous report that the non-

rewarded cue has weak representation in adolescent PFC, tilting the overall cue representation

towards the rewarded cue [75]. Our study, however, identified significantly weaker cue-evoked

activity in adolescent PFC when compared to adults, especially in FS interneurons, which

showed remarkable similarities in firing to adult preterm mice. Our results hence support that

adolescence is a sensitive period for the development of prefrontal function in the context of

goal-directed behaviors [12]. Adolescents in our study ranged 41-48 days of age at the time of the

recording, while adult term mice were 80-90 days old, indicating that task-related activity in the

PFC rapidly and substantially changes between late adolescence and adulthood. Similar to the

previous report [75], we also found that adolescents performed in our task as well as adult mice,

and that they learned to inhibit the licking after the presentation of the non-rewarded cue as

successfully as adult mice. Altogether, these findings imply that behavioral inhibition in the

adolescent brain is either mediated by other brain areas or by the PFC in a manner distinct from

that in adult brain. It will be important to address this in future studies, especially in the context

of prematurity, given the similarities in the encoding of the non-rewarded cue in the PFC of adult

preterm and adolescent term mice.

While our study identified potential circuit loci of impaired neurodevelopmental

outcomes after preterm birth, it did not test if optogenetic or chemogenetic manipulations of

circuit- or cell type-specific activity improved behavioral performance of preterm mice in our

task. Chemogenetic activation of prefrontal FS interneuron promotes behavioral response

inhibition [99], but it is unlikely that chemogenetic or pharmacological approaches would have the

same effect in adult preterm mice given the cue- and outcome-specific deficits in their activity.

Such specificity could be achieved via optogenetic manipulation, but prolonged optogenetic

stimulation (such as during task training) can result in neurotoxicity and tissue damage [100].

Previous research highlighted windows of increased plasticity and vulnerability during prefrontal

development beginning shortly after birth [101]. It is possible that targeted activity and/or circuit

manipulations during these windows could improve neurocognitive outcomes after preterm birth,

but the mechanistic impact of prematurity on the developmental trajectory of the PFC is

unknown. Mice are considered premature at term birth already, and the first 7-10 postnatal days

is typically thought to be the equivalent of third trimester in fetal development [102]. While we

cannot reliably estimate what would be the equivalent of human fetal age of prematurely born
mice, our study nevertheless highlights the long-term impact of advanced birth on prefrontal
function. As most models of prematurity use term born mice subjected to neonate
hypoxia/ischemia [103], future studies need to address the differences and similarities between
different models of prematurity, especially in the context of prefrontal function in prematurely
born humans.

In conclusion, our results identify impairments in the representation and processing of
sensory cues in the visual and prefrontal cortices of preterm mice, and provide further evidence
of interneuron dysfunction in the preterm brain [17,18,103]. Our study highlights the profound and
persistent impact of preterm birth on the function of top-down circuits, further stressing the
vulnerability of the PFC to perinatal and postnatal stressors. [12]

Funding

This study was supported by the Department of Psychology at the University of Virginia, the
National Center For Advancing Translational Sciences of the National Institutes of Health under
Award Numbers KL2TR003016/ULTR003015, the Brain Institute at the University of Virginia,
and R01MH140184 to A.R. L.Y. D. was in part supported by Summer Research Internship
Program (SRIP), O.P.B. by the Harrison Undergraduate Research Award, and S.C.F. by the
College Council Student Research Grant at the University of Virginia.

Acknowledgements

The authors would like to thank the Cang and Liu labs at the Departments of Biology and
Psychology, as well as the Program in Fundamental Neuroscience, for generous access to their
confocal microscopes. The authors would like to acknowledge Dr. Francesca Sciaccotta and Alia
Minaya for their help with data analysis.

Inclusion & ethics

All authors of this study have fulfilled the criteria for authorship required by Nature Portfolio
journals, as their participation was essential for the design and implementation of the study.

Data Availability

All data from this study is available upon reasonable request. Source data is provided with this
paper.

Code Availability

All analysis codes used in this study are available online (CED website).

Author Contributions

E.M. and A.R. performed all electrophysiological experiments and confocal imaging, and E.M.,
V.P. and A.R. performed all behavioral experiments, with assistance from M.F., L.Y.D., O.B.,
S.F., J.S., L.E., T.D.H. and P. A.-A. All authors contributed to the analysis of the behavioral data.
Electrophysiological data was analyzed by X.T. and A.R. A.R. conceived the study and wrote the
manuscript with input from all authors.

Figure Legends

Figure 1. Adult preterm mice have intact orientation selectivity. a) Preterm mice were
generated through subcutaneous injection of mifepristone (MFP) to timed pregnant dams at
gestational day (GD) 17. Preterm mice were delivered at GD18, a day early. Control term mice
were delivered at GD19 by dams injected with vehicle (DMSO) at GD17. **b)** Primary visual
cortex (V1) of mice was recorded while mice were presented with 6 different orientations 30º
apart. Preterm mice had no significant differences in **c)** orientation selectivity index (OSI
median±MAD : Term=0.78±0.05, Preterm=0.81±0.08, Mann Whitney t-test p=0.86, U=52), **d)**
fraction of neurons with OSI<0.3 (mean±SEM: Term=77.59±3.81, Preterm=72.04±5.22, t-test
p=0.4, t=0.86, df=16.84), **e)** tuning width of selective neurons (median±MAD:
Term=14.02±1.08, Preterm=15.19±1.73, Mann Whitney t-test p=0.29, U=70.5), or **f)** the
distribution of preferred frequencies (p=0.92, X^2 test). N=414 neurons from 11 term mice and
306 neurons from 10 preterm mice.

Figure 2. Preterm mice show impaired learning and behavioral inhibition in a visual
discrimination task. a) Schematics of gestation length for experimental groups. Preterm mice
were generated through subcutaneous injection of MFP to timed pregnant dams at GD17, and
control term mice were delivered at GD19 by dams injected with vehicle (DMSO) at GD17 or
with MFP at GD18. **b)** Head-fixed term and preterm mice were trained to discriminate two

orientations while locomoting on a treadmill positioned in front of a screen displaying the task
cues. The reward (water) was delivered through a spout positioned near the mouth. **c)** Task
structure: a single session consisted of 50 randomized, 3 s long presentations of rewarded and
non-rewarded cue each, with an intertrial interval of 15 s. The detection window was 10 s. Four
possible outcomes are Hit, False Alarm (FA), Miss and Correct Rejection (CR), as indicated. **d)**
Representative lick raster of naïve (top) and trained (bottom) mice to the rewarded (left) and
non-rewarded (right) cues, indicated with a blue and grey rectangle, respectively. Note the
reduced reaction times after the presentation of the rewarded cue and withholding of licks during
and after the non-rewarded cue in trained mice. **e)** Behavioral performance was measured as d',
which showed linear improvements in both groups of term mice, but not in preterm mice (LMM,
Term vs Preterm p=0.007, MFP Term vs Preterm p=0.02, Term vs MFP Term p=0.78; N=22
Term, 29 Preterm and 17 MFP Term mice). **f)** A significant fraction of preterm mice (17.24%)
failed to reach the criterion, in contrast to term mice (Kolmogorov-Smirnov test of d' values
from the last training session Term vs Preterm: D=0.43, p=0.009, Term vs MFP Term: D=0.34,
p=0.14, MFP Term vs Preterm: D=0.2893, p=0.23). **g)** Hit rates of naïve preterm mice are
significantly lower at the onset of training (ANOVA birth x training p=0.016, $F_{(2, 65)}$=4.37;
Bonferroni post-hoc Naïve Term vs. Preterm p=0.005, Term vs MFP Term p>0.9, MFP Term vs
Preterm p=0.0013; all p>0.9 among Trained groups). **h)** False Alarm rates were significantly
higher in trained preterm mice, indicating impaired response inhibition (ANOVA birth x training
p<0.0001, $F_{(2, 65)}$=10.66; Bonferroni post-hoc Naïve Term vs. Preterm p>0.9, Term vs MFP Term
p=0.13, MFP Term vs Preterm p=0.04; Trained Term vs Preterm p=0.036, Term vs MFP Term
p=0.53, MFP Term vs Preterm p=0.0007). **i)** Comparison of lick frequencies (binned in 20 trial
blocks) during the last session revealed a significant effect of time in session (LMM; Trial Block
p<0.0001), with lick frequencies dropping towards the end of the session at the same level across
groups (Animal Group p=0.15). **j)** Schematics of the open field test. Mice were released into an
open arena and their locomotor activity was recorded and quantified. **k)** Term and preterm mice
crossed a similar distance in the open field, confirming similar levels of locomotor activity
(Welch's ANOVA ($F_{(2,56.98)}$=3.9, p=0.03; Dunnett's T3 multiple comparisons test Term vs. MFP
Term p=0.03, Term vs. Preterm p=0.83, MFP Term vs. Preterm p=0.41; N=30 Term, 25 MFP
Term and 35 Preterm mice). **l)** Term and preterm mice had comparable preference for the walls
of the open field arena, indicating comparable exploratory drive (Welch's ANOVA $F_{(2,51.62)}$=4.08,

p=0.03; Dunnett's T3 multiple comparisons test Term vs. MFP Term p=0.07, Term vs. Preterm
p=0.06, MFP Term vs. Preterm p=0.87).

Figure 3. Processing of task cues is impaired in the V1 of trained preterm mice. a) V1
activity of trained term and preterm mice was recorded while the mice were engaged in the task.
b) Representative example of averaged field potential response (visually evoked potential/VEP)
to task cues used for confirming the recording location. Scale bars: 100 ms and 50 µV. **c)** Top:
lick raster, and Bottom: peristimulus time histogram (PSTH) of a representative V1 neuron
showing the transient and sustained components of the response during the cue presentation. **d)**
Correlogram of V1 unit activity and licks revealed a peak in firing of V1 neurons prior to lick
onset (Wilcoxon matched-pairs signed rank test, p=0.018, W=-1354). **e-h)** Representation of task
cues is intact in regular-spiking (RS, e and f) and fast-spiking (FS, g and h) neurons in the V1 of
preterm mice (RM ANOVA with birth and the direction of modulation as factors; RS Reward:
birth x modulation p=0.93, $F_{(2,38)}$=0.07; RS No Reward: birth x modulation p=0.7, F=0.36; f: 2-
tailed t-test, p=0.9, t=0.0046, df=19; FS Reward: birth x modulation p=0.95, F=0.09; FS No
Reward: birth x modulation p=0.24, F=1.46; h: p=0.66, t=0.45; bar graphs represent mean±SEM
of % of positively modulated/yellow, negatively modulated/magenta and unmodulated/white
neurons, box plots indicate median and quartiles). **i-l)** PSTHs of V1 neurons significantly
modulated by task cues revealed profoundly elevated firing in both RS (i and j) and FS (k and l)
V1 neurons in preterm mice (permutation tests of area under the PSTH curve, p=0.000 for all).
Data are represented as mean±SEM. **m-n)** Distributions of selectivity indices (SI) of neurons
significantly modulated by task cues revealed a significant shift to lower values for both cues in
RS neurons in preterm mice (m) and an increase in higher values for the rewarded cue in FS
neurons (n). K-S test for all: RS Reward D=0.35, p=0.0007; RS No Reward D=0.46, p=0.00005,
FS Reward D=0.29, p=0.000006; No Reward D=0.11, p=0.36; Ns are indicated in main text).

Figure 4. Representation and processing of task cues is impaired in the PFC of preterm
mice. a) PFC activity of trained term and preterm mice was recorded while the mice were
engaged in the task. **b)** Representative electrode tracks in the coronal section of the PFC. DiI
(green) marks the tracks, PFC areas and layers are outlined (Allen Brain Atlas). Scale bar: 500
µm. **c)** Correlogram of PFC unit activity and licks revealed peaks in firing of PFC neurons prior
to lick onset (Wilcoxon matched-pairs signed rank test, p<0.0001, W=-2770). **d)** Representation

of the non-rewarded cue is significantly weakened in preterm mice (ANOVA Reward: birth x

modulation, p=0.6, $F_{(2,34)}$=0.52; No Reward: p=0.014, F=4.88. Data represents mean±SEM of %

of positively modulated/yellow, negatively modulated/magenta and unmodulated/white neurons

of 9 mice per group. **e)** Overall cue representation is shifted in favor of the rewarded cue in

preterm mice (Mann-Whitney test p=0.045, U=20.5; medians and interquartile range are

indicated in box plots). **f-g)** PSTHs of RS neurons significantly modulated by task cues revealed

significant hyperactivity in preterm PFC. Permutation test of PSTH AUCs, p=0.000, Ns are

indicated in main text. Data represents mean±SEM. **h)** Distributions of SIs indicated that RS

neurons in preterm mice are significantly less responsive to the non-rewarded cue, with several

neurons completely suppressed (values close to -1). Reward: K-S test, D=0.13, p=0.36; No

Reward: D=0.4, p=0.000014; medians are indicated. **i)** Linear regression of SI values for term

and **(j)** preterm mice revealed that the SI for the rewarded cue are correlated with and predictive

of performance (d') in preterm but not term mice (Pearson's correlation and linear regression; r

values, 95% confidence intervals and regression are shown in graph). **k-l)** Representation of cues

is not impaired in FS neurons of preterm PFC (Reward: birth x modulation, p=0.71,

F(2,34)=0.34; No Reward: p=0.2, F=1.67; Mann-Whitney test p=0.17, U=28). m-n) PSTHs of

FS neurons significantly modulated by task cues revealed significantly blunted cue-evoked

activity in preterm PFC. Permutation test of PSTH AUCs, p=0.000, Ns are indicated in main

text. **o)** SI values of preterm mice were shifted towards higher values for the rewarded cue and

towards lower for the non-rewarded cue (K-S test, Reward D=0.19, p=0.02; No Reward D=0.36,

p=0.00097). **p-r)** SI values of FS neurons in either group of mice did not correlate with d'

(Pearson's correlation, r values indicated in graph).

Figure 5. Encoding but not representation of behavioral outcomes to the non-rewarded

cues is altered in the PFC of preterm mice. a-b) Representation of behavioral outcomes to the

non-rewarded cue is not altered in prefrontal RS (a) or FS (b) neurons of preterm mice (ANOVA:

birth x modulation RS CR: p=0.81, $F_{(2,34)}$=0.21, FA: p=0.6, F=0.51; FS CR: p=0.65, F=0.43, FA:

p=0.45, F=0.82). Data represents mean±SEM of % of positively modulated/yellow, negatively

modulated/magenta and unmodulated/white neurons of 9 mice per group. **c)** PSTHs of RS

neurons significantly modulated by the non-rewarded cue during CR (left) and FA (right)

revealed elevated firing of RS neurons during both behavioral outcomes in preterm mice

(permutation test of PSTH AUCs, p=0.000; Ns are indicated in main text). **d)** PSTHs of FS

neurons significantly modulated by the non-rewarded cue during CR (left) and FA (right)

revealed blunted and irregular firing of FS neurons during both behavioral outcomes in preterm

mice that was significantly different to term mice only during FA trials (permutation test of

PSTH AUCs, CR p=0.125, FA p=0.022; Ns are indicated in main text). **e)** SI values for both RS

and FS neurons were shifted towards lower values during both behavioral outcomes (K-S test;

RS CR D = 0.44, p=0.000042; FA D = 0.26, p=0.014; FS CR D=0.24, p=0.044; FA D=0.57,

p=1.2×10^{-10}). **f)** Non-rewarded cue-evoked firing of all neurons isolated from term and preterm

mice during CR and FA revealed overall significant differences in evoked activity in preterm

PFC, with a pronounced reduction of cue-evoked firing of FS neurons (K-S test; RS CR

p=0.011, D=0.23; FA, p=0.0008, D=0.28; FS CR p=0.0001, D=0.26; FA, p=0.93×10^{-6}, D=0.31).

g) Logistic regression model was fit with trial-by-trial firing rates evoked by the non-rewarded

cue as classifiers and CR/FA as outcomes, where N=number of spikes in stimulus/baseline

window. **h)** ROC curves indicated that firing rate data of term mice performs significantly better

in the model than the firing rates of preterm mice (z=6.79, p<0.0001).

Figure 6. Non-rewarded cue has a significantly reduced representation in the PFC of

adolescent term mice. a) Experimental timeline: mice were prepared for training starting at

postnatal day 25, and the training began one week later. **b)** Adolescent term mice learn the task

comparable to adult term mice (LMM, age p=0.42; data represents mean±SEM of N=18 male

and female mice, details indicated in main text). **c)** Increase in the performance of adolescent

mice is due to a reduction in False Alarm rate (RM ANOVA: response x training p<0.0001,

$F_{(1,17)}$=40.5; HR p=0.07, FA p<0.0001). **d)** Schematic of electrophysiological recordings used to

collect the data. **e-f)** Representation of the non-rewarded cue is significantly weaker than that of

the rewarded cue in both RS (e; RM ANOVA cue x modulation p<0.0001, $F_{(2.2,17.79)}$=37.11) and

FS (f; p<0.0001, $F_{(2.5,20.04)}$=18.37) neurons. Data represents mean±SEM of % of positively

modulated/yellow, negatively modulated/magenta and unmodulated/white neurons of 9 mice per

group. **g)** PSTHs of significantly modulated neurons revealed weaker activity evoked by the non-

rewarded cue in both RS (left) and FS neurons (right; p=0.000, permutation test of PSTH AUCs).

Data represents mean±SEM. **h)** SI indices of neurons in (g) confirmed that the selectivity for the

non-rewarded cue is shifted towards lower values in both RS and FS neurons (K-S test RS:

D=0.31, p=0.005; FS: D=0.27, p=0.01). Medians are indicated. **i)** PSTHs of neurons significantly

modulated by the non-rewarded cue showed a significantly higher activity during FA trials in

both RS (left) and FS (right) neurons of adolescent PFC (p=0.000, permutation test of PSTH

AUCs). Data represents mean±SEM. **j)** Distribution of SI values during CR and FA trials was not

significantly different in RS or FS neurons (K-S test; RS: D=0.21, FS: D=0.12, p=0.89). Medians

are indicated. **k)** Logistic regression model (as in Figure 5g-h) of single trial firing rates evoked

by the non-rewarded cue as classifiers of CR/FA revealed that the ROCs of adolescent term and

adolescent preterm mice substantially overlap (adolescent vs preterm p=0.17, z=1.38; adolescent

vs term adult p<0.0001, z=-5.76). All Ns are indicated in the main text.

Figure 7. Lifelong environmental enrichment fails to improve the learning trajectory of

preterm mice. a) Experimental timeline. Term and preterm mice were housed in an enriched

environment from postnatal day 5 (P5) until the end of the training. **b)** The learning trajectory of

term mice reared in an enriched environment was steep, but they performed worse than term

mice from the standard environment. Preterm mice reared in an enriched environment also

performed worse than preterm mice from standard environment [LMM with Enrichment (ENR)

and Birth as Fixed Effects; ENR p<0.0001, Birth p<0.0001) **c)** There are no differences in the

distribution of final d' values between term and preterm mie reared in an enriched environment

(K-S test, D=0.46, p=0.16**). d)** Hit rates during the final training session were significantly

different between term and preterm mice reared in enriched environment (ANOVA training x

birth x enrichment p=0.002, F=28.49; naïve ENR Term vs Preterm p=0.19, trained p=0.006. **e)**

False Alarm rates were not significantly different between term and preterm mice reared in

enriched environment (naïve ENR Term vs Preterm p=0.99, trained p=0.99; N=9 term and 11

preterm mice, statistical details are indicated in Supplementary Materials).

References

1067

1. Heuvelman, H. *et al.* Gestational age at birth and risk of intellectual disability without a

common genetic cause. *Eur. J. Epidemiol.* **33**, 667–678 (2018).

2. Perapoch, J. *et al.* Prematurity and ADHD in Childhood: An Observational Register-

Based Study in Catalonia. *J. Atten. Disord.* **25**, 933–941 (2021).

3.   Crump, C., Sundquist, J. & Sundquist, K. Preterm or early term birth and risk of

attention-deficit/hyperactivity disorder: a national cohort and co-sibling study. *Ann.*

Epidemiol. **86**, 119-125.e4 (2023).

4.   Allen, L., Leon-Attia, O., Shaham, M., Shefer, S. & Gabis, L. V. Autism risk linked to

prematurity is more accentuated in girls. *PLoS ONE* **15**, e0236994 (2020).

5.   Crump, C., Sundquist, J. & Sundquist, K. Preterm or early term birth and risk of autism.

Pediatrics **148**, (2021).

6.   Husby, A., Wohlfahrt, J. & Melbye, M. Gestational age at birth and cognitive outcomes

in adolescence: population based full sibling cohort study. *BMJ* **380**, e072779 (2023).

7.   van Houdt, C. A., Oosterlaan, J., van Wassenaer-Leemhuis, A. G., van Kaam, A. H. &

Aarnoudse-Moens, C. S. H. Executive function deficits in children born preterm or at low

birthweight: a meta-analysis. *Dev. Med. Child Neurol.* **61**, 1015–1024 (2019).

8.   Schnider, B. *et al.* Executive function deficits mediate the association between very

preterm birth and behavioral problems at school-age. *Early Hum. Dev.* **146**, 105076

(2020).

9.   Johnson, S. & Marlow, N. Preterm birth and childhood psychiatric disorders. *Pediatr.*

Res. **69**, 11R–8R (2011).

10.   Johnson, S. & Marlow, N. Early and long-term outcome of infants born extremely

preterm. *Arch. Dis. Child.* **102**, 97–102 (2017).

11.   Tokariev, A. *et al.* Preterm birth changes networks of newborn cortical activity. *Cereb.*

1092   *Cortex* **29**, 814–826 (2019).

1093   12.   Chini, M. & Hanganu-Opatz, I. L. Prefrontal Cortex Development in Health and Disease:

1094   Lessons from Rodents and Humans. *Trends Neurosci.* **44**, 227–240 (2021).

1095   13.   Ball, G. *et al.* The effect of preterm birth on thalamic and cortical development. *Cereb.*

1096   *Cortex* **22**, 1016–1024 (2012).

1097   14.   Nosarti, C. *et al.* Preterm birth and structural brain alterations in early adulthood.

1098   *Neuroimage Clin.* **6**, 180–191 (2014).

1099   15.   Wehrle, F. M. *et al.* Multimodal assessment shows misalignment of structural and

1100   functional thalamocortical connectivity in children and adolescents born very preterm.

1101   *Neuroimage* **215**, 116779 (2020).

1102   16.   Yrjölä, P., Stjerna, S., Palva, J. M., Vanhatalo, S. & Tokariev, A. Phase-Based Cortical

1103   Synchrony Is Affected by Prematurity. *Cereb. Cortex* **32**, 2265–2276 (2022).

1104   17.   Lacaille, H., Vacher, C.-M. & Penn, A. A. Preterm birth alters the maturation of the

1105   gabaergic system in the human prefrontal cortex. *Front. Mol. Neurosci.* **14**, 827370

1106   (2021).

1107   18.   Lacaille, H. *et al.* Impaired interneuron development in a novel model of neonatal brain

1108   injury. *eNeuro* **6**, (2019).

1109   19.   Panda, S. *et al.* Estrogen Treatment Reverses Prematurity-Induced Disruption in Cortical

1110   Interneuron Population. *J. Neurosci.* **38**, 7378–7391 (2018).

1111   20.   Tibrewal, M. *et al.* Disruption of Interneuron Neurogenesis in Premature Newborns and

1112   Reversal with Estrogen Treatment. *J. Neurosci.* **38**, 1100–1113 (2018).

1113   21.   Komitova, M. *et al.* Hypoxia-induced developmental delays of inhibitory interneurons

1114   are reversed by environmental enrichment in the postnatal mouse forebrain. *J. Neurosci.*

1115   **33**, 13375–13387 (2013).

1116   22.   Moloney, R. A. *et al.* Ongoing effects of preterm birth on the dopaminergic and

1117   noradrenergic pathways in the frontal cortex and hippocampus of guinea pigs. *Dev.*

1118   *Neurobiol.* **84**, 93–110 (2024).

1119   23.   Moloney, R. A., Palliser, H. K., Pavy, C. L., Shaw, J. C. & Hirst, J. J. Zuranolone therapy

1120   protects frontal cortex neurodevelopment and improves behavioral outcomes after

1121   preterm birth. *Brain Behav.* **14**, e70009 (2024).

1122   24.   Gramatté, T. & Schmidt, J. The effect of early postnatal hypoxia on the development of

1123   locomotor activity in rats. *Biomed. Biochim. Acta* **45**, 523–529 (1986).

1124   25.   Speiser, Z., Korczyn, A. D., Teplitzky, I. & Gitter, S. Hyperactivity in rats following

1125   postnatal anoxia. *Behav. Brain Res.* **7**, 379–382 (1983).

1126   26.   van der Kooij, M. A. *et al.* Mild neonatal hypoxia-ischemia induces long-term motor- and

1127   cognitive impairments in mice. *Brain Behav. Immun.* **24**, 850–856 (2010).

27.   Domnick, N.-K. *et al.* Neonatal hypoxia-ischemia impairs juvenile recognition memory

by disrupting the maturation of prefrontal-hippocampal networks. *Exp. Neurol.* **273**, 202–

214 (2015).

28.   Forbes, T. A. *et al.* Environmental enrichment ameliorates perinatal brain injury and

promotes functional white matter recovery. *Nat. Commun.* **11**, 964 (2020).

29.   Korkhin, A., Zubedat, S., Aga-Mizrachi, S. & Avital, A. Developmental effects of

environmental enrichment on selective and auditory sustained attention.

Psychoneuroendocrinology **111**, 104479 (2020).

30.   Zheng, H. J. V. *et al.* Environmental enrichment sharpens sensory acuity by enhancing

information coding in barrel cortex and premotor cortex. *eNeuro* **8**, (2021).

31.   Cancedda, L. *et al.* Acceleration of visual system development by environmental

enrichment. *J. Neurosci.* **24**, 4840–4848 (2004).

32.   Schuch, C. P. *et al.* Early environmental enrichment affects neurobehavioral development

and prevents brain damage in rats submitted to neonatal hypoxia-ischemia. *Neurosci.*

Lett. **617**, 101–107 (2016).

33.   Cheong, J. L. Y., Burnett, A. C., Treyvaud, K. & Spittle, A. J. Early environment and

long-term outcomes of preterm infants. *J. Neural Transm.* **127**, 1–8 (2020).

34.   Allen, T. A. *et al.* Imaging the spread of reversible brain inactivations using fluorescent

muscimol. *J. Neurosci. Methods* **171**, 30–38 (2008).

35.   Ingrao, J. C. *et al.* Aqueous stability and oral pharmacokinetics of meloxicam and

carprofen in male C57BL/6 mice. *J. Am. Assoc. Lab. Anim. Sci.* **52**, 553–559 (2013).

36.   Guo, Z. V. *et al.* Procedures for behavioral experiments in head-fixed mice. *PLoS ONE* **9**,

e88678 (2014).

37.   Hayar, A., Bryant, J. L., Boughter, J. D. & Heck, D. H. A low-cost solution to measure

mouse licking in an electrophysiological setup with a standard analog-to-digital

converter. *J. Neurosci. Methods* **153**, 203–207 (2006).

38.   Patel, T. P. *et al.* An open-source toolbox for automated phenotyping of mice in

behavioral tasks. *Front. Behav. Neurosci.* **8**, 349 (2014).

39.   Witteveen, I. F. *et al.* Preterm birth accelerates the maturation of spontaneous and resting

activity in the visual cortex. *Front. Integr. Neurosci.* **17**, 1149159 (2023).

40.   Jurjut, O., Georgieva, P., Busse, L. & Katzner, S. Learning Enhances Sensory Processing

in Mouse V1 before Improving Behavior. *J. Neurosci.* **37**, 6460–6474 (2017).

41.   Liu, D. *et al.* Orbitofrontal control of visual cortex gain promotes visual associative

learning. *Nat. Commun.* **11**, 2784 (2020).

42.   Murray, S. A. *et al.* Mouse gestation length is genetically determined. *PLoS ONE* **5**,

e12418 (2010).

43.   Castillo-Ruiz, A., Hite, T. A., Yakout, D. W., Rosen, T. J. & Forger, N. G. Does birth

trigger cell death in the developing brain? *eNeuro* **7**, (2020).

1166 44. Ribic, A., Crair, M. C. & Biederer, T. Synapse-Selective Control of Cortical Maturation

1167   and Plasticity by Parvalbumin-Autonomous Action of SynCAM 1. *Cell Rep.* **26**, 381-

1168   393.e6 (2019).

1169 45. Niell, C. M. & Stryker, M. P. Highly selective receptive fields in mouse visual cortex. *J.*

1170   *Neurosci.* **28**, 7520–7536 (2008).

1171 46. Berditchevskaia, A., Cazé, R. D. & Schultz, S. R. Performance in a GO/NOGO

1172   perceptual task reflects a balance between impulsive and instrumental components of

1173   behaviour. *Sci. Rep.* **6**, 27389 (2016).

1174 47. Del Rosario, J. *et al.* Diminished cortical excitation and elevated inhibition during

1175   perceptual impairments in a mouse model of autism. *Cereb. Cortex* **31**, 3462–3474

1176   (2021).

1177 48. Goel, A. *et al.* Impaired perceptual learning in a mouse model of Fragile X syndrome is

1178   mediated by parvalbumin neuron dysfunction and is reversible. *Nat. Neurosci.* **21**, 1404–

1179   1411 (2018).

1180 49. Poort, J. *et al.* Learning Enhances Sensory and Multiple Non-sensory Representations in

1181   Primary Visual Cortex. *Neuron* **86**, 1478–1490 (2015).

1182 50. Henschke, J. U. *et al.* Reward Association Enhances Stimulus-Specific Representations

1183   in Primary Visual Cortex. *Curr. Biol.* **30**, 1866-1880.e5 (2020).

1184   51.   Corbo, J., McClure, J. P., Erkat, O. B. & Polack, P.-O. Dynamic Distortion of Orientation

1185   Representation after Learning in the Mouse Primary Visual Cortex. *J. Neurosci.* **42**,

1186   4311–4325 (2022).

1187   52.   Boughter, J. D., St John, S. J., Noel, D. T., Ndubuizu, O. & Smith, D. V. A brief-access

1188   test for bitter taste in mice. *Chem. Senses* **27**, 133–142 (2002).

1189   53.   Durand, S. *et al.* A comparison of visual response properties in the lateral geniculate

1190   nucleus and primary visual cortex of awake and anesthetized mice. *J. Neurosci.* **36**,

1191   12144–12156 (2016).

1192   54.   Niell, C. M. & Stryker, M. P. Modulation of visual responses by behavioral state in

1193   mouse visual cortex. *Neuron* **65**, 472–479 (2010).

1194   55.   Xing, D., Yeh, C.-I., Burns, S. & Shapley, R. M. Laminar analysis of visually evoked

1195   activity in the primary visual cortex. *Proc Natl Acad Sci USA* **109**, 13871–13876 (2012).

1196   56.   Stringer, C., Michaelos, M., Tsyboulski, D., Lindo, S. E. & Pachitariu, M. High-precision

1197   coding in visual cortex. *Cell* **184**, 2767-2778.e15 (2021).

1198   57.   Zhang, S. *et al.* Selective attention. Long-range and local circuits for top-down

1199   modulation of visual cortex processing. *Science* **345**, 660–665 (2014).

1200   58.   Norman, K. J. *et al.* Post-error recruitment of frontal sensory cortical projections

1201   promotes attention in mice. *Neuron* **109**, 1202-1213.e5 (2021).

59. Leinweber, M., Ward, D. R., Sobczak, J. M., Attinger, A. & Keller, G. B. A sensorimotor

circuit in mouse cortex for visual flow predictions. *Neuron* **95**, 1420-1432.e5 (2017).

60. Zhang, S. *et al.* Organization of long-range inputs and outputs of frontal cortex for top-

down control. *Nat. Neurosci.* **19**, 1733–1742 (2016).

61. Le Merre, P. *et al.* Reward-Based Learning Drives Rapid Sensory Signals in Medial

Prefrontal Cortex and Dorsal Hippocampus Necessary for Goal-Directed Behavior.

Neuron **97**, 83-91.e5 (2018).

62. Peters, A. J., Marica, A.-M., Fabre, J. M. J., Harris, K. D. & Carandini, M. Visuomotor

learning promotes visually evoked activity in the medial prefrontal cortex. *Cell Rep.* **41**,

111487 (2022).

63. Wal, A., Klein, F. J., Born, G., Busse, L. & Katzner, S. Evaluating visual cues modulates

their representation in mouse visual and cingulate cortex. *J. Neurosci.* **41**, 3531–3544

(2021).

64. Kim, J.-H., Ma, D.-H., Jung, E., Choi, I. & Lee, S.-H. Gated feedforward inhibition in the

frontal cortex releases goal-directed action. *Nat. Neurosci.* **24**, 1452–1464 (2021).

65. Tervo, D. G. R. *et al.* A designer AAV variant permits efficient retrograde access to

projection neurons. *Neuron* **92**, 372–382 (2016).

66. Lima, S. Q., Hromádka, T., Znamenskiy, P. & Zador, A. M. PINP: a new method of

tagging neuronal populations for identification during in vivo electrophysiological

recording. *PLoS ONE* **4**, e6099 (2009).

67.   Cardin, J. A. Dissecting local circuits in vivo: integrated optogenetic and

1223   electrophysiology approaches for exploring inhibitory regulation of cortical activity. *J.*

1224   *Physiol. Paris* **106**, 104–111 (2012).

1225   68.   Vormstein-Schneider, D. *et al.* Viral manipulation of functionally distinct interneurons in

1226   mice, non-human primates and humans. *Nat. Neurosci.* **23**, 1629–1636 (2020).

1227   69.   Courcelles, E. J. *et al.* Association cortical areas in the mouse contain a large population

1228   of fast-spiking GABAergic neurons that do not express parvalbumin. *Eur. J. Neurosci.*

1229   **59**, 3236–3255 (2024).

1230   70.   Le Merre, P., Ährlund-Richter, S. & Carlén, M. The mouse prefrontal cortex: Unity in

1231   diversity. *Neuron* **109**, 1925–1944 (2021).

1232   71.   Jeong, J. H. & Choi, J.-S. Population analyses reveal heterogenous encoding in the

1233   medial prefrontal cortex during naturalistic foraging. (2025) doi:10.7554/eLife.93994.2.

1234   72.   Lui, J. H. *et al.* Differential encoding in prefrontal cortex projection neuron classes across

1235   cognitive tasks. *Cell* **184**, 489-506.e26 (2021).

1236   73.   Hanley, J. A. & McNeil, B. J. The meaning and use of the area under a receiver operating

1237   characteristic (ROC) curve. *Radiology* **143**, 29–36 (1982).

1238   74.   Réveillon, M., Hüppi, P. S. & Barisnikov, K. Inhibition difficulties in preterm children:

1239   Developmental delay or persistent deficit? *Child Neuropsychol.* **24**, 734–762 (2018).

1240   75.   Nieves, G. M. & Liston, C. Divergent reward cue representations in the prefrontal cortex

1241   drive reward motivation in adolescence and adulthood. *BioRxiv* (2023)

1242   doi:10.1101/2023.11.07.565069.

1243   76.   Orso, R. *et al.* Early environmental enrichment rescues memory impairments provoked

1244   by mild neonatal hypoxia-ischemia in adolescent mice. *Behav. Brain Res.* **407**, 113237

1245   (2021).

1246   77.   Bartoletti, A., Medini, P., Berardi, N. & Maffei, L. Environmental enrichment prevents

1247   effects of dark-rearing in the rat visual cortex. *Nat. Neurosci.* **7**, 215–216 (2004).

1248   78.   Akhund-Zade, J., Ho, S., O'Leary, C. & de Bivort, B. The effect of environmental

1249   enrichment on behavioral variability depends on genotype, behavior, and type of

1250   enrichment. *J. Exp. Biol.* **222**, (2019).

1251   79.   Martínez-Cué, C. *et al.* Differential effects of environmental enrichment on behavior and

1252   learning of male and female Ts65Dn mice, a model for Down syndrome. *Behav. Brain*

1253   *Res.* **134**, 185–200 (2002).

1254   80.   Stolp, H. B. *et al.* Interneuron development is disrupted in preterm brains with diffuse

1255   white matter injury: observations in mouse and human. *Front. Physiol.* **10**, 955 (2019).

1256   81.   Parnaudeau, S. *et al.* Mediodorsal thalamus hypofunction impairs flexible goal-directed

1257   behavior. *Biol. Psychiatry* **77**, 445–453 (2015).

1258   82.   Marton, T. F., Seifikar, H., Luongo, F. J., Lee, A. T. & Sohal, V. S. Roles of prefrontal

1259   cortex and mediodorsal thalamus in task engagement and behavioral flexibility. *J.*

1260   *Neurosci.* **38**, 2569–2578 (2018).

1261   83.   Wimmer, R. D. *et al.* Thalamic control of sensory selection in divided attention. *Nature*

1262   **526**, 705–709 (2015).

1263   84.   Mukherjee, A. *et al.* Variation of connectivity across exemplar sensory and associative

1264   thalamocortical loops in the mouse. *eLife* **9**, (2020).

1265   85.   Delevich, K., Tucciarone, J., Huang, Z. J. & Li, B. The mediodorsal thalamus drives

1266   feedforward inhibition in the anterior cingulate cortex via parvalbumin interneurons. *J.*

1267   *Neurosci.* **35**, 5743–5753 (2015).

1268   86.   Molnár, Z. & Rutherford, M. Brain maturation after preterm birth. *Sci. Transl. Med.* **5**,

1269   168ps2 (2013).

1270   87.   Toda, T. *et al.* Birth regulates the initiation of sensory map formation through serotonin

1271   signaling. *Dev. Cell* **27**, 32–46 (2013).

1272   88.   Leung, M. P., Thompson, B., Black, J., Dai, S. & Alsweiler, J. M. The effects of preterm

1273   birth on visual development. *Clin. Exp. Optom.* **101**, 4–12 (2018).

1274   89.   Hunt, B. A. E. *et al.* Disrupted visual cortex neurophysiology following very preterm

1275   birth. *Biol. Psychiatry Cogn. Neurosci. Neuroimaging* (2019)

1276   doi:10.1016/j.bpsc.2019.08.012.

90.   Emberson, L. L., Boldin, A. M., Riccio, J. E., Guillet, R. & Aslin, R. N. Deficits in Top-

Down Sensory Prediction in Infants At Risk due to Premature Birth. *Curr. Biol.* **27**, 431–

436 (2017).

91.   Boldin, A. M., Geiger, R. & Emberson, L. L. The emergence of top-down, sensory

prediction during learning in infancy: A comparison of full-term and preterm infants.

Dev. Psychobiol. **60**, 544–556 (2018).

92.   Jaffe-Dax, S., Boldin, A. M., Daw, N. D. & Emberson, L. L. A Computational Role for

Top-Down Modulation from Frontal Cortex in Infancy. *J. Cogn. Neurosci.* **32**, 508–514

(2020).

93.   Siegle, J. H. *et al.* Reconciling functional differences in populations of neurons recorded

with two-photon imaging and electrophysiology. *eLife* **10**, (2021).

94.   Wei, Z. *et al.* A comparison of neuronal population dynamics measured with calcium

imaging and electrophysiology. *PLoS Comput. Biol.* **16**, e1008198 (2020).

95.   Kerstjens, J. M., de Winter, A. F., Bocca-Tjeertes, I. F., Bos, A. F. & Reijneveld, S. A.

Risk of developmental delay increases exponentially as gestational age of preterm infants

decreases: a cohort study at age 4 years. *Dev. Med. Child Neurol.* **54**, 1096–1101 (2012).

96.   Pierrat, V. *et al.* Neurodevelopmental outcome at 2 years for preterm children born at 22

to 34 weeks' gestation in France in 2011: EPIPAGE-2 cohort study. *BMJ* **358**, j3448

(2017).

97.   Tang, Y. P., Wang, H., Feng, R., Kyin, M. & Tsien, J. Z. Differential effects of

1297   enrichment on learning and memory function in NR2B transgenic mice.

1298   *Neuropharmacology* **41**, 779–790 (2001).

1299   98.   Imperio, C. G. *et al.* Exposure to environmental enrichment attenuates addiction-like

1300   behavior and alters molecular effects of heroin self-administration in rats.

1301   *Neuropharmacology* **139**, 26–40 (2018).

1302   99.   Jendryka, M. M. *et al.* Control of sustained attention and impulsivity by Gq-protein

1303   signalling in parvalbumin interneurons of the anterior cingulate cortex. *Transl. Psychiatry*

1304   **13**, 243 (2023).

1305   100.   Cardin, J. A. *et al.* Targeted optogenetic stimulation and recording of neurons in vivo

1306   using cell-type-specific expression of Channelrhodopsin-2. *Nat. Protoc.* **5**, 247–254

1307   (2010).

1308   101.   Bitzenhofer, S. H., Pöpplau, J. A., Chini, M., Marquardt, A. & Hanganu-Opatz, I. L. A

1309   transient developmental increase in prefrontal activity alters network maturation and

1310   causes cognitive dysfunction in adult mice. *Neuron* **109**, 1350-1364.e6 (2021).

1311   102.   Colonnese, M. T. *et al.* A conserved switch in sensory processing prepares developing

1312   neocortex for vision. *Neuron* **67**, 480–498 (2010).

1313   103.   Salmaso, N., Jablonska, B., Scafidi, J., Vaccarino, F. M. & Gallo, V. Neurobiology of

1314   premature brain injury. *Nat. Neurosci.* **17**, 341–346 (2014).

a Within task

b

c Licks

Trial #

V1 firing

Spikes/s

Time from cue onset (s)

d

Spike count

**

Time from lick onset (s)

Regular-spiking/putative pyramidal neurons

Fast-spiking/putative inhibitory neurons

e R: 120°   NR: 60°

% of neurons with FR changes to R and NR

ns   ns

T PT   T PT

f

Representation

R

NR

ns

T PT

g R: 120°   NR: 60°

% of neurons with FR changes to R and NR

ns   ns

T PT   T PT

h

Representation

R

NR

ns

T PT

i

Spikes/s

R   Term ***
     Preterm

Time from cue onset (s)

j

NR   Term
     Preterm

Time from cue onset (s)

k

Spikes/s

R   Term ***
     Preterm

Time from cue onset (s)

l

NR   Term ***
     Preterm

Time from cue onset (s)

m

Reward   T]**
          PT

No Reward   T]***
             PT

Selectivity index (SI)

n

Reward   T]***
          PT

No Reward   T]ns
             PT

Selectivity index (SI)

a Within task

b DiI, MO, ACC, L2/3, L5, PLA, L6

c Spike count vs Time from lick onset (s), ***

PFC

Regular-spiking/putative pyramidal neurons

d R: 120° (ns), NR: 60° (*), % of neurons with FR changes to R and NR, T PT

e Representation (R/NR), T PT, *

f R, Term***, Preterm, Spikes/s vs Time from cue onset (s)

g NR, Term***, Preterm, Spikes/s vs Time from cue onset (s)

h Reward (ns, SI), No Reward (***, SI)

i R: r=-0.29, p=0.45; NR: r=0.12, p=0.75; Selectivity index vs Behavioral performance (d')

j R: r=-0.75|R²=0.56, p=0.02; NR: r=-0.02, p=0.975; Selectivity index vs Behavioral performance (d')

Fast-spiking/putative inhibitory neurons

k R: 120° (ns), NR: 60° (ns), % of neurons with FR changes to R and NR, T PT

l Representation (R/NR), T PT, ns

m R, Term***, Preterm, Spikes/s vs Time from cue onset (s)

n NR, Term***, Preterm, Spikes/s vs Time from cue onset (s)

o Reward (*, SI), No Reward (***, SI)

p R: r=-012, p=0.78; NR: r=-0.6, p=0.097; Selectivity index vs Behavioral performance (d')

r R: r=-0.5, p=0.22; NR: r=0.02, p=0.978; Selectivity index vs Behavioral performance (d')