Measuring Microbial Community-Wide Antibiotic Resistance Propagation via Natural

Transformation in the Human Gut Microbiome

Nadratun N. Chowdhury, Samuel P. Forry, Stephanie L. Servetas, Monique E. Hunter, Jennifer

N. Dootz, Joy P. Dunkers, Scott A. Jackson

National Institute of Standards and Technology, Gaithersburg, MD 20878, United States

1) Abstract

*Created with Biorender.com

This work explored the role of natural transformation – a mechanism by which bacteria uptake

and express extracellular genes - in driving antibiotic resistance propagation in the human gut

microbiome. The model extracellular antibiotic resistance gene (eARG) – a plasmid containing a

kanamycin resistance (*kanR*) gene and a green fluorescence protein (GFP) gene – was dosed into

pooled and homogenized human stool and incubated anaerobically. Cellular uptake of the eARG

was assessed via droplet digital PCR, the expression of newly acquired genes was assessed by

culturing on selective media and fluorescent microscopy, newly resistant isolates were identified

by long-read Nanopore sequencing and the impacts on the taxonomy of the gut microbiome was

assessed using shotgun Illumina sequencing. Significant gene uptake of both *kanR* and GFP was

quantified in gut microbes, and extent of gene accumulation correlated with background

kanamycin levels. Gut microbes dosed with background kanamycin expressed kanamycin

resistance acquired by the eARG (as quantified by CFU on kanamycin-containing media). Newly

resistant isolates, identified as *Enterococcus faecium* by long-read sequencing, also expressed

green fluorescence acquired from the eARG. Though compositional changes of the kanamycin-

resistant subpopulation were observed in the gut microbiome in response to eARG and antibiotic

exposure, these changes were not reproducible among replicates and trends in taxonomy due to

transformation could not be identified. This comprehensive analysis therefore establishes the

significant propagation of antibiotic resistance within the human gut microbiome due to eARG

exposure, while evaluating the utility of various measurements in characterizing transformation

in a complex microbial community.

2) Importance

Infections from antibiotic resistant bacteria in the human gut microbiome are a growing public

health concern. Antibiotic resistance may develop in gut microbiota from exposure to

environmentally prevalent extracellular antibiotic resistance genes (eARGs). This work explores

the impact of eARG exposure on a complex human gut microbial community. It quantifies

significant accumulation and expression of eARG-borne genes by endogenous gut

microorganisms, thereby demonstrating that natural transformation may play a role in resistance

propagation in the human gut. It also demonstrates the highly variable changes in gut taxonomy

in response to eARG exposure, implying that eARGs may impact gut composition and therefore

downstream human health effects. These data may be useful in characterizing and mitigating

resistance propagation in the human gut, and in general, the suite of genotypic and phenotypic

measurements used constitute a quantitative framework to characterize the effects of

perturbations on complex microbiomes.

3) Introduction

The rapid evolution of antimicrobial resistance has led to a major global public health crisis.

Infections due to antibiotic-resistant bacteria have contributed to worldwide increases in

mortality - estimated at 35,000 deaths/year in the US, 30,000 deaths per year in the EU and even

greater numbers in low and middle-income countries in the global south. [1-3] Such deadly

infections may occur in the human gut microbiome - the highly complex and diverse community

of bacteria, archaea, viruses, and eukaryotic microbes within the human intestinal tract. The

human gut microbiome is host to various opportunistic pathogens, including *Escherichia coli,*

Klebsiella pneumoniae, Enterococcus faecalis and *Enterococcus faecium.* Infections from

antibiotic-resistant *E. coli*, *E. faecium* and *K. pneumoniae* have been on the rise in last several

years. [4, 5] To improve public health outcomes, there is a growing need to prevent or mitigate the

propagation of antibiotic-resistant bacteria in the human gut microbiome. To do this, the

mechanisms by which resistance spreads in complex microbial communities must be thoroughly

evaluated and understood.

The dissemination of widely used antibiotics (for human health and anthropogenic

purposes) provides the main pressure for bacterial populations to mutate and develop resistance.

[6-8] However, mobile genetic elements coding for antibiotic resistance – such as plasmids,

transposons, integrons, and extracellular chromosomal DNA - may also be agents of spread.

Bacteria may acquire mobile genes coding for antibiotic resistance from other bacteria (of the

same or different species) through horizontal gene transfer (HGT). HGT may occur via various

mechanisms – such as conjugation (in which mobile genetic elements are transferred through a

pilus connecting two bacterial cells) and transduction (in which a bacteriophage transfers DNA

between bacteria). Perhaps the least studied mechanism, however, is natural transformation - in

which bacterial cells uptake extracellular DNA, incorporate this DNA into their genomes and

express traits encoded by the DNA. Transformation is an ecologically significant process that has

been detected on the bench scale and in situ. [9-13] Unlike other forms of gene transfer,

transformation does not require the presence of a living donor cell, a close genetic relationship

between donor and recipient cells, or physical closeness between donor and recipient cells.

Therefore, gene networks in a microbial community may be significantly altered based on

environmental exposures via transformation.

The transformation process requires bacteria to activate proteins that enable the binding

of exogenous DNA to the cell surface, uptake of DNA fragments into the cell, homologous

recombination of DNA fragments to the host chromosome (via a recombinase protein called

recA), and the translation of exogenous genetic material. [14-22] Bacteria with this capability are

considered naturally competent, and over 80 species of bacteria have been determined on the

bench-scale to exhibit natural competence. [23] In most species, the mechanism of competence is

inducible, and bacteria may induce competence to repair damaged DNA, obtain nutrients, or

incorporate genes that improve their fitness. [10, 24-26] Various environmental factors – such as

nutrient limitation or antibiotic concentration – can trigger the activation of competence proteins.

[13, 14, 17] Some naturally competent species – such as *E. coli* –typically inhabit the human gut. [27]

In addition, the human gut may contain extracellular antibiotic resistance genes (eARGs)

released from living and dead resistant bacteria. eARGs are prevalent in the environment and

have been quantified in areas that are potential sources for human exposure– including

freshwater [28-30], sediment [31, 32], animal feed operations [8, 33], drinking water treatment plants [8, 34],

and wastewater treatment plants [35-38]. Though most extracellular DNA quickly degrades when

exposed to ubiquitous DNases, detectable levels of dosed eARGs have been isolated and

quantified within simulated human gut microbiomes in rats. [39] This implies that some eARGs

may persist long enough in the gut to transform endogenous microbes. Transformed microbes

may then disseminate resistance genes to their progeny through vertical gene transfer or to

unrelated bacteria via other forms of HGT. Therefore, natural transformation is potentially a

significant pathway of resistance acquisition for human gut microbes, but its contribution has not

yet been quantified.

Culture-based methods have been previously used to demonstrate that eARGs can

genetically transform and confer resistance within bacterial monocultures. [40-48] Some studies

have quantified transformation in cultured engineered consortia or biofilms of select bacterial

strains. [16, 49, 50] Culture-based methods, while sensitive, are limited in that they are low-

throughput and may only be used to study cultivable strains. In recent years, rapidly developing

high throughput culture-independent methods, such as next generation sequencing, have

provided the opportunity to collect large amounts of information and comprehensively profile

eARGs in environmental microbiomes. These methods are being utilized to study genetic

integration of resistance genes in diverse microbial populations– such as in a recent study that

evaluated the contribution of eARGs to antibiotic resistance propagation in activated sludge. [43] In

the human gut, conjugative transfer of antibiotic resistance genes has been observed in situ using

such culture-independent methods. [51, 52] But there remains a need to make similar community-

wide observations of resistance transfer via natural transformation.

In this work, the contribution of natural transformation to antibiotic resistance

propagation in the human gut microbiome was quantified by dosing a representative eARG into

homogenized human stool. Human stool, meant to represent the contents of the gut microbiome,

was pooled from multiple donors and verified for homogeneity and stability between samples.

The dosed eARG was a synthetic extracellular plasmid deprived of conjugative ability containing

a kanamycin resistance gene (*kanR*) and a gene encoding green fluorescence protein (GFP). The

microbial community was simultaneously exposed to kanamycin to determine impacts of

antibiotic exposure on transformation efficiencies. Genotypic and phenotypic acquisition of

kanamycin resistance in gut microbes was assessed using a combination of culture-based and

molecular methods. Taxonomic shifts due to antibiotic and eARG exposure was evaluated using

next generation sequencing methods, and emergent resistant species were characterized via

microscopy and sequencing. This comprehensive analysis informs the current understanding of

antibiotic resistance development in the human gut while evaluating the utility of various

measurements in characterizing transformation in a complex microbial community.

4) Methods

*Created with Biorender.com

Figure 1. Summary of methods to transform the fecal microbiome with pBAV1K-T5-GFP under

kanamycin exposure, and characterize phenotypic and genotypic effects across three culture

steps.

1A) In the first culture step, homogenized human stool is aliquoted into BHI media with and

without pBAV1K-T5-GFP (eARG) and kanamycin dosing. Liquid culture is grown

anaerobically at 37 °C for 24 hours. 1B) DNA is extracted from stool samples grown in liquid

culture. 1C) DNA is sequenced via whole genome sequencing for taxonomic profiling of the

treated gut microbiome and analyzed via droplet digital PCR to quantify the dosed eARG.

2A) In the second culture step, liquid cultures grown in step 1 are plated on BHI agar plates, one

set on selective plates (with kanamycin) and one set on non-selective plates. Plates are incubated

anaerobically at 37 °C for 72 hours. 2B) Agar plates are prepared for molecular analysis by

swabbing whole plates into a liquid culture and extracting DNA from this culture, 2C) Samples

from the second culture step are analyzed directly from agar plates by counting CFU to

determine how treatment with eARG and kanamycin impacted the quantity of resistant cells.

Extracted DNA from swabbed whole plates are analyzed via whole genome sequencing to

determine the taxonomic profile of the resistant gut microbial population.

3A) Colonies of distinct morphologies are isolated from agar plates containing kanamycin-

treated samples from culture step 2, 3B) DNA from isolates are extracted to prepare for

molecular analysis. 3C) Isolates, prior to DNA extraction, are imaged via fluorescent microscopy

to examine how treatment with eARG impacted expression of green fluorescence. Extracted

DNA from isolates are sequenced via long-read sequencing for molecular identification of

isolates that emerge after treatment with eARG and kanamycin.

4.1) Preparation of materials

4.1.1) Homogenized stool

Human whole stool was obtained from multiple volunteer donors by BioIVT (https://bioivt.com/,

Hicksville, NY, USA). All whole stool samples were collected after informed consent under

approved IRB protocols at BioIVT. Prior to donation, donors were screened for HIV, Hepatitis B

and C, and Syphilis, West Nile virus and *Trypasnosoma cruzi*. Stool samples were collected

from multiple volunteer donors on site and immediately stored at -80˚ C until processing.

Upon receipt of all donations, samples were combined with molecular grade water to a final

concentration of 100 mg/mL wet weight and homogenized. Samples were then aliquoted in 1mL

volumes, shipped to NIST on dry ice, and then stored at -80˚ C until use.

4.1.2) eARG

The rolling circle plasmid pBAV1K-T5-GFP was used as the model eARG in this study. This

plasmid contains genes coding resistance to kanamycin (referred to as *kanR* genes) and a green

fluorescent protein (GFP). The *kanR* gene is an aminoglycoside 3'-phosphotransferase (*aph(3')-*

IIIa), which catalyzes the addition of phosphate from ATP to the 3'-hydroxyl group of

kanamycin, thereby inactivating antibiotic activity. [53] The GFP gene codes for the emission of

green fluorescence. The *kanR* gene is used to select for kanamycin resistance and GFP gene is

used as an additional marker for phenotypic expression of the plasmid. This plasmid backbone

has been previously verified to participate in natural transformation in inter-species studies and

shown the ability to positively replicate in a variety of species – such as *Streptococcus pyogens*

and *Acinetobacter baumanii*. It also is deprived of its ability to conjugate once inside of bacterial

cells. [50] Therefore, it was considered an ideal model eARG that may quantifiably transform

competent gut microorganisms without confounding effects from conjugation. The plasmid was

prepared as described in Text S1 in the Supplementary Information (SI).

4.1.3) Verification of transformability & extractability

E. coli was used as a preliminary control to verify the eARG's transformability. Chemically

competent *E. coli* was transformed with pBAV1K-T5-GFP and verified to have the kanamycin

resistant phenotype and express green fluorescence, as described in Text S2 and shown in Figure

S1 in the SI. The transformed *E. coli* and purified plasmid were then systematically dosed into

the stool samples, extracted, and quantified (as described in Text S2 in SI) to verify the ability to

adequately extract and measure intracellular plasmid in subsequent transformation experiments.

pBAV1K-T5-GFP DNA was successfully recovered from stool samples dosed with transformed

E. coli. (Figure S2 in SI)

4.2) Natural transformation assay

Homogenized stool samples were thawed and 250 µL aliquots were prepared for each sample.

Kanamycin sulfate was dissolved into water to make a stock concentration of 5000 μg/mL

(stored at 4 °C, covered from light). Kanamycin solution was added to stool aliquots, with two

aliquots at each of the final kanamycin concentrations – 0, 0.5, 5 and 50 µg/mL. 10 µL of

prepped pBAV1K-T5-GFP was added to half of the stool aliquots, such that for each of the four

kanamycin concentrations, there was one eARG-free and one eARG-containing sample. This

was done to determine the extent to which background antibiotic concentration drove uptake and

expression of the eARG. The eARG-free samples with kanamycin were prepared to determine

the extent to which the presence of antibiotic caused gut microorganisms to evolve without

horizontal gene transfer. The samples for each condition were prepared in triplicate.

Once prepared, the samples were well mixed via vortex. They were then incubated at 37 °C

for 24 hours. This incubation period was chosen to be similar to a realistic human gut transit

time, which may range between 10 and 73 hours. [54] Samples were incubated in an anaerobic

chamber, which mimics the gut environment. After 24 hours, 1 µL of DNase I was added to each

sample to quench transformation, and the samples were incubated another 15 minutes. Samples

were then pelleted via centrifugation and resuspended in 100 µL PBS. Intracellular DNA was

extracted using the Quick-DNA Fecal/Soil Microbe Miniprep Kit by Zymo Research (Irvine,

CA) according to the manufacturer's instructions, with bead beating on a vortex adaptor at full

speed for 40 minutes.

4.3) Analysis of gut microbiome upon eARG exposure

4.3.1) Quantification of eARG uptake via droplet digital PCR

Droplet digital PCR (ddPCR) was performed to quantify the GFP, *kanR* and 16S genes in

transformed gut microorganisms. The GFP gene was quantified to assess the extent of plasmid

presence as an indication of cellular eARG uptake. The *kanR* gene was also quantified to assess

eARG uptake but is additionally used to determine the quantity of *kanR* genes present that were

not related to eARG exposure. Gene copy numbers of GFP and *kanR* were normalized to copy

numbers of 16S, which is a proxy for the abundance of cells present. Because antibiotic

treatment likely affects the abundance of cells over time, it was vital to normalize gene copy

numbers to 16S gene copies to assess impacts of the eARG or antibiotic. DNA extracted from

transformed stool samples was used as template DNA. All experiments were performed on the

QX200 Droplet Digital PCR System by Bio-Rad (Hercules, CA). All ddPCR reactions for

quantifying GFP and *kanR* genes were performed in 22.5 µL, including the ddPCR Supermix for

Probes (No dUTP) by Bio-Rad. The thermal cycling protocol used for these two genes was: 95

°C for 10 min followed by 40 cycles at 94 °C for 30 s and 59 °C for 60 s, followed by 98 °C for

10 min. All ddPCR reactions for quantifying 16S genes were performed in 25 µL, including the

QX200™ ddPCR™ EvaGreen Supermix by Bio-Rad. The thermal cycling protocol used for this

gene was: 95 °C for 10 min followed by 59 cycles at 94 °C for 30 s and 60 °C for 60 s, followed

by 98 °C for 10 min. The sets of forward and reverse primers used are given in the SI (Table S1).

4.3.2) Microbiome profiling by metagenomics (WGS)

Changes in microbial composition in response to eARG and antibiotic exposure was assessed by

performing whole genome (shotgun) sequencing. Samples were prepared for sequencing using

the NEB Ultra II DNA Library Prep kit for Illumina library preparation (New England Biolabs,

Ipswich, Massachusetts). Samples were combined by mass and prepared for sequencing

following the MiSeq System Denature and Dilute Libraries Guide (Document # 15039740 v10,

Protocol A). Denatured libraries were diluted to a final concentration of 12 pM and combined

with 5 % PhiX control (V3 cat# 15017666 from Illumina). Paired-end sequencing was performed

on an Illumina MiSeq with 2×150 bp reads (MiSeq Reagent Kit v2 600-cycle, cat #: MS-102-

3003). Adapter trimming was done as part of the Illumina MiSeq Generate FASTQ workflow.

Fastq files were assembled and taxonomy was classified using Centrifuge WoL: Reference

Phylogeny for Microbes Release 1 (available on Github; https://biocore.github.io/wol/) built

using 10,575 genomes x 381 genes. [55]

4.4) Post-transformation outgrowths

Transformation assays were also followed up by outgrowths of transformed samples on agar

plates containing kanamycin. This was done to determine the quantity of gut microbes that

displayed phenotypic expression of kanamycin resistance, and to identify specific strains that

may have been transformed. First, homogenized stool samples were prepared with pBAV1K-T5-

GFP and kanamycin as described in Section 4.2. For this experiment, samples were prepared

only at 0 µg/mL and 50 µg/mL kanamycin, each with and without eARG. Kanamycin

concentrations of 0.5 µg/mL and 5 µg/mL were not tested here because 50 µg/mL kanamycin

appeared to drive transformation most effectively. Each sample type was prepared in triplicate.

All incubations were performed in an anaerobic chamber. After 24 hours, the samples were

removed from the incubator and serially diluted. Aliquots of 100 μL of samples from each

condition were each spread plated in triplicate on both selective agar plates (Brain Heart Infusion

(BHI) agar plates with 2.5% sheep's blood and 50 μg/mL kanamycin) and non-selective agar

plates (BHI-blood plates without kanamycin). Counts on non-selective agar plates provided the

baseline count of stool organisms, and counts on selective agar plates provided the count of

kanamycin-resistant organisms. Agar plates were incubated in the anaerobic chamber at 37 °C

for 72 hours. Ample incubation time was given to allow slow-growing microbes to form visible

colonies on the plate.

4.4) Analysis of cultured gut microbiome after eARG exposure

4.4.1) CFU count

After 72 hours, plates were removed from the incubator and CFU (colony-forming unit) counts

were obtained on both selective and non-selective plates. The fraction of organisms that were

kanamycin resistant was determined by taking the ratio of CFU counts on selective agar plates to

CFU counts on non-selective plates. Agar plates were visualized on the Scan 4000 by

Interscience (Saint-Nom-la-Bretèche, France) and images are presented in SI (Figure S3).

4.4.2) Identifying transformed organisms by metagenomics (WGS)

To determine the taxonomic composition of kanamycin-resistant gut microorganisms, and

potentially gain insight into the identities of gut microorganisms subject to natural

transformation, colonies from the outgrowths were analyzed by shotgun sequencing. Selective

agar plates were swabbed after CFU counts, and cells were inoculated in 500 μL PBS. DNA was

extracted from cells using the Quick-DNA Fecal/Soil Microbe Miniprep Kit from Zymo

Research (Irvine, CA) and sequenced according to the method described in Section 4.3.2.

4.4.3) Visualization & isolation of colonies

Colonies on selective agar plates were also isolated to identify specific taxa in stool that may

have participated in transformation. The outgrowths described in Section 4.4 were performed

again, but only on kanamycin-containing selective agar plates. After growth, selective agar plates

were inspected for colonies with varying morphologies. Distinct colonies were selected for each

sample type, and streak-plated on Blood-BHI agar plates with kanamycin. Streak plates were

swabbed into a 50% glycerol solution and stored at -80 °C for further analysis.

4.4.4) Fluorescence microscopy

Isolates from the frozen stock (prepared as described in Section 4.4.3) were inoculated in BHI

liquid culture with 50 μg/mL kanamycin and grown overnight. Cultures for each isolate were

examined via fluorescence microscopy to determine the morphology of the cells, and whether

they expressed green fluorescence as another phenotypic indicator of transformation.

For microscopic imaging, 3 μl sample volumes for each sample were pipetted onto a

microscope slide cleaned with KOH and treated with poly-L-lysine. A coverslip was placed on

top, tamped down lightly and sealed. The images were collected with a Nikon Eclipse Ti2-E

Inverted Microscope equipped with a Hamamatsu (Shizuoka, Japan) ORCA-Flash 4.0 LT+

CMOS camera and a SOLA SE U-nIR solid state light source (Lumecor, Beaverton, OR).

Imaging was performed using a 60X, 1.2 NA plan apochromat water immersion objective with

Immersol W (Zeiss, Oberkochen, Germany) as the index matching fluid. Two channel images

were acquired using manual focus for each image with these settings: GFP channel (ORCA

detector: 2 s integration time, SOLA source: 40%), Differential Inferential Contrast (DIC)

channel (ORCA detector: 90 ms integration time, LED source: 10%). The Nikon format files

were processed in ImageJ.

4.4.5) Identification of stool isolates

To determine the identity of resistant stool microbes isolated from the transformed community,

isolates from the frozen stock (prepared as described in Section 4.4.3) were inoculated in BHI

liquid culture with 50 μg/ml kanamycin and grown overnight at 37 ℃. DNA was extracted from

each culture using the Quick-DNA Fecal/Soil Microbe Miniprep Kit from Zymo Research

(Irvine, CA) per manufacturers protocol, and genomic DNA was prepared for nanopore

sequencing using the Native Barcoding Expansion 1-12 kit (Oxford Nanopore Technologies,

EXP-NBD104). Libraries were sequenced using the Ligation Sequencing kit on the MinION

Mk1C (Flow Cell R9.4.1) with the option 'basecalling' enabled, with a 48 hour run limit, 1.5

hour pore scan frequency, and 200 bp minimum read length. Sequences were demultiplexed

during the run. Barcoded reads were loaded into the EPI2ME Desktop Agent and analyzed using

FASTQ WIMP which utilizes Centrifuge version 1.0.3-Beta and a RefSeq database (built

06/08/2021) that includes 56,044 sequences that represent bacteria, archaea, viral, and fungi.

FASTQ WIMP was run with the default settings.

4.5) Data analysis & visualization

All data points were collected in triplicate. Statistical analysis and data visualization was

performed in RStudio. Significant differences between gene copy numbers via ddPCR and CFU

counts were determined using R functions for the Kruskal-Walis test (among antibiotic and

plasmid treatments) due to the lack of normal distribution. Non-metric multidimensional scaling

R functions (Bray-Curtis) were used to compare metagenomic composition (among antibiotic

and eARG treatments).

5) Results

5.1) Accumulation of eARG genes in gut microbial community measured by ddPCR

Figure 2. Gene copies of target genes (GFP and *kanR)* from pBAV1K-T5-GFP quantified via

droplet digital PCR in DNA extracted from gut microorganisms dosed with pBAV1K-T5-GFP

(plasmid) (x = no plasmid dose, + = plasmid dose) and kanamycin, grown anaerobically.

The abundance of GFP gene copies extracted from transformed cells, normalized to 16S gene

copies, significantly increased with plasmid addition in all conditions. A significant increase in

GFP abundance in eARG-dosed samples compared to eARG-free samples was observed when

no kanamycin was added in the background ($p< 0.01$*). Increasing levels of background

kanamycin, however, impacted the magnitude of GFP gene copy number increase between

eARG-dosed and eARG-free samples. The impact of eARG addition on normalized GFP gene

copy number was significant at all kanamycin concentrations ($p< 0.025$* at 0.5 ng/µl Kan;

$p<0.028$* at 5 ng/µl Kan; $p<0.006$** at 50 ng/µl Kan), and the magnitude of this effect

increased with increasing concentrations of background kanamycin. The largest increase was

observed when 50 ng/µL of kanamycin was present. Of note is that the population of the gut

microbial community, for which 16S gene copies were measured as a proxy, decreased with

increasing concentrations of background kanamycin. Therefore, even when accounting for the

decrease in the total microbial population by normalizing to 16S gene copy number, the

magnitude of GFP gene accumulation correlated with eARG-dosing increased with higher

antibiotic doses.

The *kanR* gene was measured at higher levels than GFP in all samples, even those that

were not dosed with the eARG. The addition of the eARG led to similar magnitudes of increase

in *kanR* gene copies as GFP gene copies, which is to be expected because a copy of each gene

is present on the dosed plasmid. Unlike GFP, however, the quantity of *kanR* genes in eARG-

free samples increased with increasing concentrations of background antibiotic ($p<0.07$).

5.2) Increase in kanamycin resistant organisms due to eARG exposure

Figure 3. Ratio of CFU on selective kanamycin-containing agar plates to CFU on non-selective

agar plates (Ratio Resistant CFU) grown for 72 hours after incubation in liquid culture with

pBAV1K-T5-GFP (x = no plasmid dose, + = plasmid dose) and kanamycin.

All stool samples transformed in the presence of 0 and 50 ng/µL of kanamycin showed

significant growth on agar plates, and the Ratio Resistant CFU (CFU on selective kanamycin-

containing agar plates/CFU on non-selective agar plates) is reported for each sample. The Ratio

Resistant CFU is a measurement that describes the number of kanamycin-resistant culturable

microorganisms as a fraction of the total culturable microorganisms. There was no significant

effect of the eARG dosing on the Ratio Resistant CFU when transformation was performed

without background kanamycin (p<0.512). Therefore, eARG-dosing did not lead to significant

differences in the population of culturable kanamycin-resistant microbes in the absence of the

antibiotic.

The overall Ratio Resistant CFU decreased with 50 ng/µL kanamycin treatment as

compared to samples with no kanamycin treatment. However, upon treatment with kanamycin,

significant differences in the Ratio Resistant CFU between eARG-dosed and eARG-free samples

were observed ($p < 0.049*$). The fraction of cells that expressed kanamycin resistance increased

nearly 100-fold upon eARG dosing. Therefore, the amount of culturable gut microbes expressing

kanamycin resistance, relative to the total gut microbial community, increased upon exposure to

the eARG in the presence of background antibiotic.

5.3) Identification of strains that may undergo natural transformation in gut microbiome

Table 1. Visualization of bacterial isolates of three distinct morphology types grown on

kanamycin-containing agar plates after treatment with kanamycin, with and without pBAV1K-

T5-GFP Isolates are visualized on a Differential Inferential Contrast (DIC) channel and a

fluorescent channel. Isolates are indicated as "Not visible" when they were present on agar plates

but not visible under the microscope, and "Not present" when they were not present on agar

plates.

Morp-	DIC		Fluorescent	
hology	**Plasmid**	**No Plasmid**	**Plasmid**	**No Plasmid**
Type				

Table 2. Identification of bacterial isolates grown on kanamycin-containing agar plates after

treatment with kanamycin, with and withoutpBAV1K-T5-GFP, by Nanopore sequencing.

Morphology Type	eARG	Top Species Classification	Total Reads	Reads Matched	Percent Reads Matched
1	Yes	*Saccharomyces cerevisiae*	456,632	431,802	94.6%
1	No	*Saccharomyces cerevisiae*	553,713	529,229	95.6%
2	Yes	*Enterococcus faecium*	349,470	256,858	73.5%
2	No	*Enterococcus faecium*	300,884	217,620	72.3%
3	Yes	*Enterococcus faecium*	413,122	294,711	71.3%

Outgrowths of transformed stool microorganisms on kanamycin-containing agar plates

led to colonies with various appearances (Figure S3). Three visually distinct types of colonies

were discernable. Each of the isolates were imaged, and as their colonies appeared distinct, their

morphologies did as well (Table 1). In total, three distinct colony morphologies were identified

from the outgrowths. Two of these morphologies appeared to be present in both eARG-dosed

and eARG-free samples, and one distinct morphology appeared only in eARG-dosed samples.

Colonies of similar morphologies between eARG-free and eARG-dosed samples were placed in

the same morphology type category.

Isolates from colonies of the first morphology type were imaged, and appeared to consist

of cells that were circular and large (with a diameter 5-7 μm).These cells most closely identified

with a yeast, *Saccharomyces cerevisiae,* with 94.6% and 95.6% of Nanopore sequencing reads

matching for the eARG and eARG-free samples, respectively (Table 2). Green fluorescence was

not detected in either case, indicating that while this organism expressed resistance to

kanamycin, it did not express green fluorescence.

Isolates from the second colony morphology type visualized under the microscope

appeared to consist of smaller, circular microorganisms with diameters of about 1-2 μm.

Morphology type 2 was also found in both eARG-dosed and eARG-free samples. Isolates of this

morphology type most closely matched the genus group *Enterococcus/Lactococcus,* with 81.4%

and 80.3% of Nanopore sequencing reads matching for eARG-dosed and eARG-free samples,

respectively. The closest species match was *Enterococcus faecium,* which had a 73.5% and

72.3% read match for eARG-dosed and eARG-free samples. Green fluorescence was detected

for isolates of this strain. However, the isolate from the eARG-dosed sample expressed

significantly brighter green fluorescence than the isolate from the eARG-free sample. Though

these strains express both kanamycin resistance (as indicated by growth on the selective agar

plate) and green fluorescence (as indicated by fluorescence microscopy), the levels of green

fluorescence expression appear to increase upon eARG exposure.

The third morphology type was found only on agar plates containing eARG-dosed

samples. This isolate was imaged and found to be a rod-like microorganism about 5-10 µm in

length. This isolate also most closely matched the genus group *Enterococcus/Lactococcus*, with

79.1% of reads matching, and the species *Enterococcus faecium*, with 71.3% of reads matching.

Isolates of morphology type 2 and 3 are possibly different strains of the same genus, which may

explain their varied morphology but close genetic match. The expression of bright green

fluorescence for the isolate of morphology type 3 was detectable, but by a minority of the

microorganisms visualized. Bacteria of this strain, which appear only when treated with the

eARG, express resistance to kanamycin, and to a lesser extent, green fluorescence.

(A)

(B)

Figure 4. Metagenomic composition changes at a species level in stool population upon 24-hour anaerobic incubation with pBAV1K-T5-GFP (x = no plasmid dose, + = plasmid dose) and kanamycin (Kan) (concentration ng/uL), as shown by A) relative abundance and B) Bray-Curtis principal coordinate analysis (for three replicates).

When observing the total gut microbial community exposed to kanamycin and the eARG,

some reproducible effects on composition were observed with increasing kanamycin addition –

such as an increased relative abundances of *Ruminococcus spp, Blautia* spp., and *Eubacterium*

rectale, and a decreased relative abundance of *Ligilactobacillus animalis*, *Shigella* spp. and

Enteroccocus faecium (Figure 4A). Effects of the eARG on the overall gut microbial community

composition post-transformation were not discernible via metagenomics. Principal coordinate

analysis (PCoA) demonstrated distinct clustering based on kanamycin concentration (Figure 4B).

This is to be expected, as antibiotic treatment typically alters the composition of the gut

microbiome. Some weak clustering may have occurred based on plasmid treatment, but this

effect was marginal. Though changes in gene target accumulation due to plasmid addition were

detectable directly after transformation via ddPCR, such changes did not lead to distinct effects

on the microbial composition of the stool.

(A)

(B)

Figure 5. Metagenomic composition changes, at a species level of kanamycin-resistant

subpopulation of stool microorganisms (grown anaerobically for 72 hours on kanamycin-

containing agar plates) after treatment with pBAV1K-T5-GFP (x = no plasmid dose, + = plasmid

dose) and kanamycin (Kan) (concentration ng/uL) (incubated 24 hours anaerobically), as shown

by A) relative abundance and B) Bray-Curtis principal coordinate analysis (for three replicates).

When observing only the kanamycin-resistant gut microorganisms (swabbed off

kanamycin-containing agar plates containing transformed cells), the microbiome did not appear

to shift significantly due to eARG addition in samples that were transformed without background

kanamycin. (Figure 5A) All samples that were transformed without background kanamycin had

similar compositions and clustered together in principle coordinate analysis. (Figure 5B)

The addition of kanamycin at 50 ng/µL led to discernable changes in the taxonomic

profile of kanamycin-resistant colonies. As compared to gut microbiome samples not exposed to

antibiotic, new taxa were observed in kanamycin treated samples, whether they were dosed with

the eARG or not. Such strains may include species like *Lactobacillus gasseiri, Lactobacillus*

animalis, Paeniclostridium sordellii, Turicibacter sanguinis, and *Bifidobacterium animalis*,

where each showed increases in relative abundance in some kanamycin-treated samples.

Specifically, *Ligilactobacillus animalis* appears in the case of the eARG-treated samples, while

Lactobacillus gasseiri appears in the case of the eARG-free samples. Regardless, kanamycin-

resistant strains do not appear to develop in a reproducible manner, as each replicate

demonstrated different compositional makeups (e.g., increases in *Campylobacter* and

Bavilorococcus). Kanamycin-treated samples did not appear to cluster together in principle

coordinate analysis, by eARG-treatment, or overall.

6) Discussion

ddPCR results showed significant increases in normalized abundance of GFP and *kanR* gene copies extracted from eARG-dosed transformed cells. Measurable quantities of eARG in the DNA of the gut microbial community imply that gut microorganisms were taking up and accumulating the eARG. The initial uptake of the dosed plasmid likely occurred in cells that were induced into a state of natural competence. This may have occurred in response to environmental factors, such as the limitation of nutrients during the 24-hour incubation period without addition of media.[24, 56] Once taken up, plasmid-borne genes may have further propagated through the microbial community through other forms of horizontal gene transfer or passed from parent to progeny via vertical gene transfer. GFP and *kanR* accumulation is observed even in samples without background antibiotic, implying that cellular uptake of the extracellular DNA may occur even when the exogenous genes don't provide a selective advantage.

The extent of GFP and *kanR* accumulation increased with increasing levels of background kanamycin. It is possible that the presence of antibiotic stimulates competence mechanisms in gut microorganisms. Antibiotic exposure has been shown to induce competence in a variety of bacterial species [57-61], including *S. pneumoniae* exposed to kanamycin and streptomycin.[62] An increase in competence amongst gut microbes may have led to an increase of eARG uptake, and the resistance trait may have allowed transformed bacteria to have a competitive advantage in propagation. Further, antibiotics may mediate bacterial SOS responses and increase membrane permeability, which could potentially facilitate extracellular gene uptake.[63-66]

Unlike GFP, the *kanR* gene was measured in significant quantities in samples with no eARG addition, possibly because kanamycin resistance genes are present naturally in gut

microbiota, as has been demonstrated in previous studies.[67-70] Of note is that the *kanR* gene

quantity in eARG-free samples increased with increasing concentrations of background

antibiotic. This implies that antibiotic exposure drives the cellular acquisition of the *kanR* genes

through methods other than uptake of the dosed plasmid. Gut microbes may obtain resistance

traits via random mutation or conjugative transfer of existing *kanR* genes, and such microbes

would have selective advantage. In the presence of antibiotic, it is possible that a complex

microbial community will activate various mechanisms to develop resistance.

While ddPCR results measured the likely cellular uptake of the dosed eARG, the

translation and expression of newly acquired genes was confirmed by CFU counts of the

kanamycin-resistant portion of the population. Without background antibiotic, there was no

significant effect of the eARG dose on the CFU counts of resistant organisms, which implies that

cells that took up the eARG did not activate mechanisms to express the acquired genes without

selective pressure. However, when background kanamycin was present (at 50 ng/µL), eARG-

dosed samples contained significantly higher numbers of resistant organisms compared to

eARG-free samples. This implies that in the presence of antibiotic, natural transformation may

occur in the human gut microbiome and reproducibly lead to higher amounts of antibiotic-

resistant bacteria.

Though the dosed eARG was expressed by certain gut microbes, observing isolates of

kanamycin-resistant strains post-transformation revealed that different strains may react to the

dosed eARG in distinct ways. First, there is a subset of inherently kanamycin-resistant organisms

in the gut microbial community that do not express genes on the dosed eARG. *Saccharomyces*

cerevisiae (Table 2) is an example, as this strain grew on kanamycin-containing media in all

conditions and did not express green fluorescence when dosed with the eARG. It is likely that

this strain was not driven to a state of competence, as *S. cerevisiae* typically requires artificial

methods, such as electroporation or addition of a chemical such as lithium or polyethylene

glycol, to become competent.[71] Also, because this strain had intrinsic kanamycin resistance, it

may not have expended the metabolic energy needed to retain and express plasmid-borne genes

that would not provide any additional selective advantage.

Not all strains with intrinsic kanamycin-resistance seem to behave this way, however.

The *Enterococcus/Lactococcus* strain of morphology type 2 (Table 2) included kanamycin-

resistant organisms in both eARG-treated and eARG-free samples, but still showed an increased

(brighter) expression of green fluorescence in eARG-treated samples. This implies that whilst

this strain had inherently kanamycin-resistant members, it still preferred to retain and express the

dosed eARG. It is possible that this strain was more easily activated into a state of natural

competence that *S. cerevisiae - Lactococcus* and *Enterococcus* are the genera with the highest

observed numbers of predicted competence genes among lactic acid bacteria [72 73, 74] Further, the

increased expression of GFP in this isolate also implies that some transformed bacteria may

phenotypically express traits obtained from exogenous DNA, even when such traits don't

provide a selective advantage. When the eARG is taken up into the cell, recombined, and

translated – it is possible that the GFP gene is translated within the same segment as the *kanR*

gene. This may explain the co-occurrence of these traits. This is realistic, as exogenous DNA

segments integrated into host chromosomes by recA are on average about 10,000 bp long[75] and

the dosed plasmid is 3,953 bp long.

Observations of the *Enterococcus/Lactococcus* strain of morphology type 3, which grew

on kanamycin-containing plates only if treated with eARG, imply that previously non-resistant

strains may possibly turn kanamycin-resistant via transformation by the dosed eARG. Only a

small fraction of this strain's kanamycin-resistant population appears to express green

fluorescence, which suggests that transformed cells may preferentially express the *kanR* gene on

the plasmid as compared to the GFP gene. This strain is an example of a newly resistant strain

that emerges in the human gut microbiome in response to eARG exposure. Whilst enterococci

have shown the ability to effectively transfer foreign genes via conjugation[74], it is unknown

whether this strain is particularly likely to be transformed compared to other gut microbes.

Overall, according to metagenomic analysis, the addition of the eARG did not lead to

significant changes in the composition of the gut microbial community post-transformation

(Figure 4). It is possible that newly transformed cells exist in small abundances and are not

detectable amongst the millions of members in the complex microbial community. eARG-

induced taxonomic shifts may potentially be more prominent after the acquired gene is able to

propagate through the community across time and multiple generations, and effects are therefore

not detectable after only 24 hours of incubation. Unlike more sensitive methods such as ddPCR

or CFU counts, which may theoretically detect as little as 1 gene copy or 1 cell, metagenomics

has less utility in detecting changes in a small subset of the microbial community.

Subsequent culturing of the transformed population, on kanamycin-containing agar for 72

hours, led to compositional shifts in the gut microbial community (Figure 5), and particularly led

to a significant reduction in microbial diversity. This reduction may be partly due to the fact that

many gut taxa are not culturable on blood-BHI agar. However, this is also likely reflective of the

fact that many gut taxa, while tolerant to kanamycin exposure in liquid culture throughout the

transformation assay (as shown in Figure 4), do not develop phenotypic kanamycin resistance,

and therefore do not grow on kanamycin-containing agar. *E. coli,* for example, did not appear to

demonstrate phenotypic resistance to kanamycin as it went from being a dominant taxon in the

post-transformation liquid culture community (Figure 4) to not being detected in the agar-grown

kanamycin resistant community (Figure 5). In contrast, *E. faecium* becomes particularly

dominant in the agar-grown kanamycin resistant population, when it was not present in high

relative abundances in the post-transformation liquid culture. *E. faecium* is known to have

intrinsically high levels of kanamycin-resistance, but some *E. faecium* strains may have obtained

the resistant phenotype from the dosed eARG – as is likely the case with isolates of morphology

type 3.

Some new taxa in the agar-grown kanamycin-resistant subpopulation appear to have

resulted from eARG exposure (i.e., *L. animalis)*, and some from kanamycin exposure (i.e., *L.*

gasseiri) (Figure 5). However, the appearance of these taxa was not reproducible amongst the

three replicates of the same treatment (with and without eARG dosing) (Figure 5). Therefore,

whether these new taxa are the result of eARG exposure or of antibiotic selective pressure is not

discernible. It is possible that taxonomic changes in a microbial community exposed to eARG

and antibiotic occur stochastically, and the variability in potential outcomes is too high to capture

trends within three experimental replicates. This is reasonable, as antibiotic resistance

development in response to selection pressure is subject to chance effects of mutation and

genetic drift,[76, 77] and natural transformation relies on inherently stochastic molecular and cellular

events – such as contact between the eARG and the cell surface.[78]Additionally, it is possible that

longer-term compositional effects of eARG exposure on gut microbial composition are more

reproducible, but not found here because the community was evaluated after only 24 hours of

incubation. Therefore, though this study identifies various gut taxa (such as *L. animalis* or *E.*

faecium) that express phenotypic kanamycin resistance after eARG exposure, reproducible

trends in the evolution of gut taxa in response to kanamycin exposure could not be identified

within the limited replicates and incubation time allowed.

Regardless of replicate and time limitations, this work develops a methodological

framework for comprehensively assessing the effects of perturbations in a complex microbiome

using a suite of accessible microbial measurements. A combination of molecular and culture-

based methods – including ddPCR, CFU count, microscopy, short-read and long-read

metagenomic sequencing - was required to reveal the complex dynamics of the gut microbiome

exposed to extracellular elements. Each method provided insights that overcame the limitations

of other methods. For example, while some non-cultivable strains were revealed as potential

transformants via short-read sequencing (*L. animalis* or *L. gasseiri*), some resistant strains did

not appear at high enough relative abundances to be identified by metagenomics and instead

were identified as cultured isolates via microscopy and long-read sequencing (*S. cerevisiae*). And

in samples where transformation did not lead to community-wide compositional changes as

detected via metagenomics, ddPCR was able to identify trends of significant gene uptake.

Therefore, the combination of techniques used in this study results in a comprehensive profile of

natural transformation in the human gut microbiome, as has not been previously recorded.

One limitation of the method suite used, however, is that the DNA extraction and

metagenomic sequencing platforms used resulted in highly fragmented DNA, which did not

allow linkage of mobile genetic elements to their host cells. This is why, in this study, identity of

transformed gut taxa could only be confirmed in cultured isolates, and only two transformed

strains were identified (Morphology Type 2 and 3 in Table 2). However, it is possible that there

are several more transformed organisms that are not culturable. Future studies on HGT dynamics

within complex microbiomes could benefit from recently developed platforms (such as Hi-C

sequencing – a method in which DNA is cross-linked prior to lysis and extraction[79]) that allow

matching of host cells to their mobile ARGs. This would further characterize the extent and

direction of resistance evolution in complex microbiomes and expand a comprehensive analysis

of antibiotic resistance propagation.

7) Conclusions

Currently, understanding of complex HGT dynamics in the gut microbiome is limited and only

beginning to develop. Within this landscape, this work provides critical measurements of the

frequency of resistance acquisition in response to eARG exposure. Accounting for other

potential biological factors by using a homogenous and stable material in a highly controlled

experimental design - exposure to an eARG, in tandem with antibiotics, is shown to increase the

number of organisms that possess both genotypic and phenotypic resistance traits. Quantifying

HGT due to eARG exposure is an essential piece of understanding the multi-faceted evolutionary

behavior of the human gut microbiome, particularly regarding antibiotic resistance evolution.

Whilst the experimental design included large doses of both eARGs and antibiotics into the gut

microbiome to effectively characterize bacterial responses, it is reasonable to assume that similar

processes could occur in humans, as both eARGs and antibiotics are environmentally prevalent,

and antibiotics are administered regularly to patients. In addition, because ARGs encoded on

mobile genetic elements may persist in the microbial community for up to several months of

time post-antibiotic regimen[80, 81], environmental eARG exposure to the gut microbiome could

potentially cause downstream effects, such as facilitating the occurrence of gastrointestinal

diseases or other health effects that warrant further study.

Because this study demonstrates that transformation can impact traits expressed within a complex microbiome like the human gut, it is also reasonable to predict that resistance may propagate in similar ways in other relevant microbiomes. The impact of eARGs on evolving resistant populations in other clinical (i.e. skin, vaginal) or environmental (i.e. wastewater, indoor spaces in hospitals) microbiomes is also paramount to mitigating the rise of antibiotic resistance. This work provides a framework of quantitative genotypic and phenotypic measurements with a model eARG, that can practically be used to characterize resistance propagation and could be applied to future microbiome transformation studies.

NIST Disclaimer

Certain equipment, instruments, software, or materials are identified in this paper in order to specify the experimental procedure adequately. Such identification is not intended to imply recommendation or endorsement of any product or service by NIST, nor is it intended to imply that the materials or equipment identified are necessarily the best available for the purpose.

Ethics approval and Consent to Participate

All work was reviewed and approved by the U. S. National Institute of Standards and Technology (NIST) Research Protections Office. This study (protocol #: MML-2019-0135) was determined to be "not human subjects research" as defined in the Common Rule (45 CFR 46, Subpart A).

Data availability

Metagenome sequencing, ddPCR, culture, and imaging results and all code used for analysis are available online (https://data.nist.gov/od/id/mds2-3579).

Declaration of Competing Interest

The authors declare no conflict of interest.

Acknowledgements

We would like to acknowledge NIST's microbiology team, mainly Lisa Stabryla, Ishi Keenum, Jason Kralj, Kirsten Parratt, and Sheng-Lin Gibson for consulting, reviewing, and providing technical expertise for this project. This material is based upon work supported by the National Research Council's Research Associateship Program (Grant 1644868).

Authors Contributions

N.N.C, S.P.F., and S.L.S. conceptualized the project and designed experiments. N.N.C, M.E.H, J.N.D. and J.P.D. analyzed samples. N.N.C, S.P.F., and S.L.S. analyzed data. N.N.C wrote the manuscript and prepared figures. All authors reviewed and approved the final manuscript.

References

1.   CDC, *Antibiotic resistance threats in the United States, 2019*. US Department of Health and Human Services, Centres for Disease Control and …: 2019.

2.   Cassini, A.; Högberg, L. D.; Plachouras, D.; Quattrocchi, A.; Hoxha, A.; Simonsen, G. S.; Colomb-Cotinat, M.; Kretzschmar, M. E.; Devleesschauwer, B.; Cecchini, M.; Ouakrim, D. A.; Oliveira, T. C.; Struelens, M. J.; Suetens, C.; Monnet, D. L.; Strauss, R.; Mertens, K.; Struyf, T.; Catry, B.; Latour, K.; Ivanov, I. N.; Dobreva, E. G.; Tambic Andraševic, A.; Soprek, S.; Budimir, A.; Paphitou, N.; Žemlicková, H.; Schytte Olsen, S.; Wolff Sönksen, U.; Märtin, P.; Ivanova, M.; Lyytikäinen, O.; Jalava, J.; Coignard, B.; Eckmanns, T.; Abu Sin, M.; Haller, S.; Daikos, G. L.; Gikas, A.; Tsiodras, S.; Kontopidou, F.; Tóth, Á.; Hajdu, Á.; Guólaugsson, Ó.; Kristinsson, K. G.; Murchan, S.; Burns, K.; Pezzotti, P.; Gagliotti, C.; Dumpis, U.; Liuimiene, A.; Perrin, M.; Borg, M. A.; de Greeff, S. C.; Monen, J. C. M.; Koek, M. B. G.; Elstrøm, P.; Zabicka, D.; Deptula, A.; Hryniewicz, W.; Caniça, M.; Nogueira, P. J.; Fernandes, P. A.; Manageiro, V.; Popescu, G. A.; Serban, R. I.; Schréterová, E.; Litvová, S.; Štefkovicová, M.; Kolman, J.; Klavs, I.; Korošec, A.; Aracil, B.; Asensio, A.; Pérez-Vázquez, M.; Billström, H.; Larsson, S.; Reilly, J. S.; Johnson, A.; Hopkins, S., Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-

resistant bacteria in the EU and the European Economic Area in 2015: a population-level modelling analysis. *The Lancet Infectious Diseases* **2019,** *19* (1), 56-66.

3.   Stewardson, A. J.; Marimuthu, K.; Sengupta, S.; Allignol, A.; El-Bouseary, M.; Carvalho, M. J.; Hassan, B.; Delgado-Ramirez, M. A.; Arora, A.; Bagga, R.; Owusu-Ofori, A. K.; Ovosi, J. O.; Aliyu, S.; Saad, H.; Kanj, S. S.; Khanal, B.; Bhattarai, B.; Saha, S. K.; Uddin, J.; Barman, P.; Sharma, L.; El-Banna, T.; Zahra, R.; Saleemi, M. A.; Kaur, A.; Iregbu, K.; Uwaezuoke, N. S. C.; Abi Hanna, P.; Feghali, R.; Correa, A. L.; Munera, M. I.; Le, T. A. T.; Tran, T. T. N.; Phukan, C.; Phukan, C.; Valderrama-Beltrán, S. L.; Alvarez-Moreno, C.; Walsh, T. R.; Harbarth, S., Effect of carbapenem resistance on outcomes of bloodstream infection caused by Enterobacteriaceae in low-income and middle-income countries (PANORAMA): a multinational prospective cohort study. *The Lancet Infectious Diseases* **2019,** *19* (6), 601-610.

4.   Dunn, S. J.; Connor, C.; McNally, A., The evolution and transmission of multi-drug resistant Escherichia coli and Klebsiella pneumoniae: the complexity of clones and plasmids. *Current Opinion in Microbiology* **2019,** *51*, 51-56.

5.   Guzman Prieto, A. M.; van Schaik, W.; Rogers, M. R. C.; Coque, T. M.; Baquero, F.; Corander, J.; Willems, R. J. L., Global Emergence and Dissemination of Enterococci as Nosocomial Pathogens: Attack of the Clones? *Frontiers in Microbiology* **2016,** *7*.

6.   Goossens, H.; Ferech, M.; Stitchele, R. V.; Elseviers, M., Outpatient antibiotic use in Europe and association with resistance: a cross national database study. *The Lancet* **2005,** *365* (9459), 579-587.

7.   Khachatourians, G. G., Agricultural use of antibiotics and the evolution and transfer of antibiotic-resistant bacteria *Canadian Medical Association Journal* **1998,** *159*, 1129-1136.

8.   Pruden, A.; Pei, R.; Storteboom, H.; Carlson, K. H., Antibiotic resistance genes as emerging contaminants: studies in northern Colorado. *Environmental Science and Technology* **2006,** *40*, 7445-7450.

9.   Stewart, G. J.; Sinigalliano, C. D., Detection of horizontal gene transfer by natural transformation in native and introduced species of bacteria in marine and synthetic sediments. *Applied Environmental Microbiology* **1990,** *56* (6), 1818-1824.

10.   Lorenz, M. G.; Wackernagel, W., Bacterial Gene Transfer by Natural Genetic Information in the Environment. *Microbiological Reviews* **1994,** *58* (3), 563-602.

11.   Lorenz, M. G.; Wackernagel, W., High Frequency of Natural Genetic Transformation of Pseudomonas stutzeri in Soil Extract Supplemented with a Carbon/Energy and Phosphorus Source. *Applied and environmental microbiology* **1991,** *57* (4), 1246-1251.

12.   Nagler, M.; Insam, H.; Pietramellara, G.; Ascher-Jenull, J., Extracellular DNA in natural environments: features, relevance and applications. *Applied Microbiology and Biotechnology* **2018,** *102*, 6343+.

13.   Pietramellara, G.; Ascher, J.; Bogogni, F.; Ceccherini, M.; Guerri, G.; Nanipieri, P.,
Extracellular DNA In Soil And Sediment: Fate And Ecological Relevance. *Biology and Fertility*
of Soils **2008,** *45*, 219-235.

14.   Dubnau, D., Genetic competence in Bacillus subtilis. *Microbiological Reviews* **1991,** *55*
(3), 395-424.

15.   Blokesch, M., Natural competence for transformation. *Current Biology* **2016,** *26* (21),
R1126-R1130.

16.   Domenech, A.; Brochado, A. R.; Sender, V.; Hentrich, K.; Henriques-Normark, B.;
Typas, A.; Veening, J.-W., Proton Motive Force Disruptors Block Bacterial Competence and
Horizontal Gene Transfer. *Cell Host & Microbe* **2020,** *27* (4), 544-555.e3.

17.   Solomon, J. M.; Grossman, A. D., Who's competent and when: regulation of natural
genetic competence in bacteria. *Trends in Genetics* **1996,** *12* (4), 150-155.

18.   Smith, H. O.; Danner, D. B.; Deich, R. A., Genetic Transformation. *Annual Review of*
Biochemistry **1981,** *50* (1), 41-68.

19.   Morrison, D. A.; Guild, W. R., Structure of deoxyribonucleic acid on the cell surface
during uptake by pneumococcus. *Journal of bacteriology* **1973,** *115* (3), 1055-1062.

20.   Mortier-Barrière, I.; Velten, M.; Dupaigne, P.; Mirouze, N.; Piétrement, O.;
McGovern, S.; Fichant, G.; Martin, B.; Noirot, P.; Le Cam, E.; Polard, P.; Claverys, J.-P., A
Key Presynaptic Role in Transformation for a Widespread Bacterial Protein: DprA Conveys
Incoming ssDNA to RecA. *Cell* **2007,** *130* (5), 824-836.

21.   Chen, Z.; Yang, H.; Pavletich, N. P., Mechanism of homologous recombination from the
RecA–ssDNA/dsDNA structures. *Nature* **2008,** *453* (7194), 489-494.

22.   Thomas, C. M.; Nielsen, K. M., Mechanisms and barriers to horizontal gene transfer
between bacteria. *Nature Reviews Microbiology* **2005,** *3*, 711-721.

23.   Moralez, J.; Szenkiel, K.; Hamilton, K.; Pruden, A.; Lopatkin, A. J., Quantitative
analysis of horizontal gene transfer in complex systems. *Current Opinion in Microbiology* **2021,**
62, 103-109.

24.   Lorenz, M.; Wackernagel, W., High Frequency Of Natural Genetic Transformation Of
Pseudomonas Stutzeri In Soil Extract Supplemented With A Carbon/Energy And Phosphorus
Source. *Applied and Environmental Microbiology* **1991,** *57*, 1246-1251.

25.   Page, W. J.; von Tigerstrom, M., Optimal conditions for transformation of Azotobacter
vinelandii. *Journal of bacteriology* **1979,** *139* (3), 1058-1061.

26.   Dubnau, D., DNA uptake in bacteria. *Annual Reviews Microbiology* **1999,** *53*, 217-244.

27.   McInnes, R. S.; McCallum, G. E.; Lamberte, L. E.; van Schaik, W., Horizontal transfer
of antibiotic resistance genes in the human gut microbiome. *Current Opinion in Microbiology*
2020, *53*, 35-43.

28.   Guo, X. P.; Yang, Y.; Lu, D. P.; Niu, Z. S.; Feng, J. N.; Chen, Y. R.; Tou, F. Y.;
Garner, J.; Xu, J.; Liu, M.; Hochella, M. F. J., Biofilms as a sink for antibiotic resistance genes
(ARGs) in the Yangtze Estuary. *Water Research* **2017,** *129*, 277-286.

29.   Wang, D. N.; Liu, L.; Qiu, Z. G.; Shen, Z. Q.; Guo, X.; Yang, D.; Li, J.; Liu, W. I.;
Jin, M.; Li, J. W., A new adsorption-elution technique for the concentration of aquatic
extracellular antibiotic resistance genes from large volumes of water. *Water Research* **2016,** *92*,
188-198.

30.   Bien, T. L. T.; Thao, N. V.; Kitamura, S. I.; Obayashi, Y.; Suzuki, S., Release and
constancy of an antibiotic resistance gene in seawater under grazing stress by ciliates and
heterotrophic nanoflagellates. *Environmental Microbiology* **2017,** *32*, 174-179.

31.   Mao, D.; Luo, Y.; Mathieu, J.; Wang, Q.; Feng, L.; Mu, Q.; Feng, C.; Alvarez, P. J.
J., Persistence of Extracellular DNA in River Sediment Facilitates Antibiotic Resistance Gene
Propagation *Environmental Science and Technology* **2014,** *48*, 71-78.

32.   Yuan, K.; Wang, X.; Chen, X.; Zhao, Z.; Fang, L.; Chen, B.; Jiang, J.; Luan, T.;
Chen, B., Occurrence of antibiotic resistance genes in extracellular and intracellular DNA from
sediments collected from two types of aquaculture farms. *Chemosphere* **2019,** *234*.

33.   McKinney, C. W.; Loftin, K. A.; Meyer, M. T.; Davis, J. G.; Pruden, A., tet and sul
Antibiotic Resistance Genes in Livestock Lagoons of Various Operation Type, Configuration,
and Antibiotic Occurrence. *Environmental Science & Technology* **2010,** *44* (16), 6102-6109.

34.   Xi, C.; Zhang, Y.; Marrs, C. F.; Ye, W.; Simon, C.; Foxman, B.; Nriagu, J.,
Prevalence of antibiotic resistance in drinking water treatment and distribution systems. *Applied*
and environmental microbiology **2009,** *75* (17), 5714-5718.

35.   Gardner, C.; Volkoff, S.; Gunsch, C., Examining the behavior of crop-derived antibiotic
resistance genes in anaerobic sludge batch reactors under thermophilic conditions. *Biotechnology*
and Bioengineering **2019,** *116* (11), 3063-3071.

36.   Lu, Y.; Xiao, Y.; Zheng, G.; Lu, J.; Zhou, L., Conditioning with zero-valent iron or
Fe2+ activated peroxydisulfate at an acidic initial sludge pH removed intracellular antibiotic
resistance genes but increased extracellular antibiotic resistance genes in sewage sludge. *Journal*
of Hazardous Materials **2020,** *386*.

37.   Zhang, Y.; Li, A.; Dai, T.; Li, F.; Xie, H.; Chen, L.; Wen, D., Cell-free DNA: a
neglected source for antibiotic resistance genes spreading from WWTPs. *Environmental Science*
and Technology **2018,** *52*, 248-257.

38.   Rizzo, L.; Manaia, C.; Merlin, C.; Schwartz, T.; Dagot, C.; Ploy, M. C.; Michael, I.;
Fatta-Kassinos, D., Urban wastewater treatment plants as hotspots for antibiotic resistant bacteria

and genes spread into the environment: a review. *Science of the Total Environment* **2013,** *447*,
345-360.

39. Wilcks, A.; van Hoek, A. H. A. M.; Joosten, R. G.; Jacobsen, B. B. L.; Aarts, H. J. M.,
Persistence of DNA studied in different ex vivo and in vivo rat models simulating the human gut
situation. *Food and Chemical Toxicology* **2004,** *42* (3), 493-502.

40. Chowdhury, N. N.; Hicks, E.; Wiesner, M. R., Investigating and Modeling the
Regulation of Extracellular Antibiotic Resistance Gene Bioavailability by Naturally Occurring
Nanoparticles. *Environmental Science & Technology* **2022,** *56* (21), 15044-15053.

41. Khanna, M.; Stotzky, G., Transformation of Bacillus Subtilis by DNA Bound on
Montmorillonite and Effect of Dnase on the Transforming Ability of Bound DNA. *Applied and*
Environmental Microbiology **1992,** *58*, 1930-1939.

42. Cai, P.; Huang, Q.; Chen, W.; Zhang, D.; Wang, K.; Jiang, D.; Liang, W., Soil
colloids-bound plasmid DNA: Effect on transformation of E. coli and resistance to DNase I
degradation. *Soil Biology and Biochemistry* **2007,** *39* (5), 1007-1013.

43. Calderón-Franco, D.; van Loosdrecht, M. C. M.; Abeel, T.; Weissbrodt, D. G., Catch
me if you can: capturing microbial community transformation by extracellular DNA using Hi-C
sequencing. *Antonie van Leeuwenhoek* **2023,** *116* (7), 667-685.

44. Chamier, B.; Lorenz, M. G.; Wackernagel, W., Natural Transformation of Acinetobacter
calcoaceticus by Plasmid DNA Adsorbed on Sand and Groundwater Aquifer Material. *Applied*
and environmental microbiology **1993,** *59* (5), 1662-1667.

45. Contente, S.; Dubnau, D., Characterization of plasmid transformation in Bacillus subtilis:
kinetic properties and the effect of DNA conformation. *Molecular & General Genetics* **1979,**
167 (3), 251.

46. de Vos, W. M.; Venema, G.; Canosi, U.; Trautner, T. A., Plasmid transformation in
Bacillus subtilis. *Molecular & General Genetics* **1981,** *181* (4), 424 - 433.

47. Overballe-Petersen, S.; Harms, K.; Orlando, L. A. A.; Mayar, J. V. M.; Rasmussen, S.;
Dahl, T. W.; Rosing, M. T.; Poole, A. M.; Sicheritz-Ponten, T.; Brunak, S.; Inselmann, S.; de
Vries, J.; Wackernagel, W.; Pybus, O. G.; Nielsen, R.; Johnsen, P. J.; Nielsen, K. M.;
Willerslev, E., Bacterial natural transformation by highly fragmented and damaged DNA.
Proceedings of the National Academy of Sciences **2013,** *110* (49), 19860-19865.

48. Romanowski, G.; Lorenz, M. G.; Wackernagel, W., Plasmid DNA in a groundwater
aquifer microcosm -adsorption, DNAase resistance and natural genetic transformation of
Bacillus subtilis. *Molecular Ecology* **1993,** *2* (3), 171-181.

49. Nordgård, L.; Nguyen, T.; Midtvedt, T.; Benno, Y.; Traavik, T.; Nielsen, K. M., Lack
of detectable DNA uptake by bacterial gut isolates grown in vitro and by Acinetobacter baylyi
colonizing rodents in vivo. *Environ. Biosafety Res.* **2007,** *6* (1-2), 149-160.

50.   Cooper, R. M.; Tsimring, L.; Hasty, J., Inter-species population dynamics enhance
microbial horizontal gene transfer and spread of antibiotic resistance. *eLife* **2017,** *6*, e25950.

51.   Porse, A.; Schou, T. S.; Munck, C.; Ellabaan, M. M. H.; Sommer, M. O. A.,
Biochemical mechanisms determine the functional compatibility of heterologous genes. *Nature*
Communications **2018,** *9* (1), 522.

52.   Lester Camilla, H.; Frimodt-Møller, N.; Sørensen Thomas, L.; Monnet Dominique, L.;
Hammerum Anette, M., In Vivo Transfer of the vanA Resistance Gene from an Enterococcus
faecium Isolate of Animal Origin to an E. faecium Isolate of Human Origin in the Intestines of
Human Volunteers. *Antimicrobial Agents and Chemotherapy* **2006,** *50* (2), 596-599.

53.   Fong Desiree, H.; Berghuis Albert, M., Structural Basis of APH(3′)-IIIa-Mediated
Resistance to N1-Substituted Aminoglycoside Antibiotics. *Antimicrobial Agents and*
Chemotherapy **2009,** *53* (7), 3049-3055.

54.   Lee, Y. Y.; Erdogan, A.; Rao, S. S., How to assess regional and whole gut transit time
with wireless motility capsule. *J Neurogastroenterol Motil* **2014,** *20* (2), 265-70.

55.   Zhu, Q.; Mai, U.; Pfeiffer, W.; Janssen, S.; Asnicar, F.; Sanders, J. G.; Belda-Ferre,
P.; Al-Ghalith, G. A.; Kopylova, E.; McDonald, D.; Kosciolek, T.; Yin, J. B.; Huang, S.;
Salam, N.; Jiao, J.-Y.; Wu, Z.; Xu, Z. Z.; Cantrell, K.; Yang, Y.; Sayyari, E.; Rabiee, M.;
Morton, J. T.; Podell, S.; Knights, D.; Li, W.-J.; Huttenhower, C.; Segata, N.; Smarr, L.;
Mirarab, S.; Knight, R., Phylogenomics of 10,575 genomes reveals evolutionary proximity
between domains Bacteria and Archaea. *Nature Communications* **2019,** *10* (1), 5477.

56.   Finkel, S.; Kolter, R., DNA As A Nutrient: Novel Role For Bacterial Competence Gene
Homologs. *Journal of Bacteriology* **2001,** *183*, 6288-6293.

57.   Charpentier, X.; Polard, P.; Claverys, J.-P., Induction of competence for genetic
transformation by antibiotics: convergent evolution of stress responses in distant bacterial
species lacking SOS? *Current Opinion in Microbiology* **2012,** *15* (5), 570-576.

58.   Charpentier, X.; Kay, E.; Schneider, D.; Shuman Howard, A., Antibiotics and UV
Radiation Induce Competence for Natural Transformation in Legionella pneumophila. *Journal of*
Bacteriology **2011,** *193* (5), 1114-1121.

59.   Dorer, M. S.; Fero, J.; Salama, N. R., DNA Damage Triggers Genetic Exchange in
Helicobacter pylori. *PLOS Pathogens* **2010,** *6* (7), e1001026.

60.   Sexton, J. A.; Vogel, J. P., Regulation of Hypercompetence in <i>Legionella
pneumophila</i>. *Journal of Bacteriology* **2004,** *186* (12), 3814-3825.

61.   Prudhomme, M.; Attaiech, L.; Sanchez, G.; Martin, B.; Claverys, J.-P., Antibiotic
Stress Induces Genetic Transformability in the Human Pathogen <i>Streptococcus
pneumoniae</i>. *Science* **2006,** *313* (5783), 89-92.

62.   Stevens, K. E.; Chang, D.; Zwack, E. E.; Sebert, M. E., Competence in Streptococcus
pneumoniae Is Regulated by the Rate of Ribosomal Decoding Errors. *mBio* **2011,** *2* (5),
10.1128/mbio.00071-11.

63.   Beaber, J. W.; Hochhut, B.; Waldor, M. K., SOS response promotes horizontal
dissemination of antibiotic resistance genes. *Nature* **2004,** *427* (6969), 72-74.

64.   Fishbein, S. R. S.; Mahmud, B.; Dantas, G., Antibiotic perturbations to the gut
microbiome. *Nature Reviews Microbiology* **2023,** *21* (12), 772-788.

65.   Lu, J.; Wang, Y.; Li, J.; Mao, L.; Nguyen, S. H.; Duarte, T.; Coin, L.; Bond, P.;
Yuan, Z.; Guo, J., Triclosan at environmentally relevant concentrations promotes horizontal
transfer of multidrug resistance genes within and across bacterial genera. *Environment*
International **2018,** *121*, 1217-1226.

66.   Zhang, P.-Y.; Xu, P.-P.; Xia, Z.-J.; Wang, J.; Xiong, J.; Li, Y.-Z., Combined treatment
with the antibiotics kanamycin and streptomycin promotes the conjugation of Escherichia coli.
FEMS Microbiology Letters **2013,** *348* (2), 149-156.

67.   Wang, W.-L.; Xu, S.-Y.; Ren, Z.-G.; Tao, L.; Jiang, J.-W.; Zheng, S.-S., Application
of metagenomics in the human gut microbiome. *World journal of gastroenterology* **2015,** *21* (3),
803-814.

68.   Bag, S.; Ghosh, T. S.; Banerjee, S.; Mehta, O.; Verma, J.; Dayal, M.; Desigamani, A.;
Kumar, P.; Saha, B.; Kedia, S.; Ahuja, V.; Ramamurthy, T.; Das, B., Molecular Insights into
Antimicrobial Resistance Traits of Commensal Human Gut Microbiota. *Microbial Ecology* **2019,**
77 (2), 546-557.

69.   Broaders, E.; Gahan, C. G. M.; Marchesi, J. R., Mobile genetic elements of the human
gastrointestinal tract. *Gut Microbes* **2013,** *4* (4), 271-280.

70.   Levy, S. B.; Marshall, B.; Schluederberg, S.; Rowse, D.; Davis, J., High frequency of
antimicrobial resistance in human fecal flora. *Antimicrobial Agents and Chemotherapy* **1988,** *32*
(12), 1801-1806.

71.   Kawai, S.; Pham, T. A.; Nguyen, H. T.; Nankai, H.; Utsumi, T.; Fukuda, Y.; Murata,
K., Molecular insights on DNA delivery into Saccharomyces cerevisiae. *Biochemical and*
Biophysical Research Communications **2004,** *317* (1), 100-107.

72.   Connell, L. M.; Kelleher, P.; van Rijswijck, I. M. H.; de Waal, P.; van Peij, N. N. M.
E.; Mahony, J.; van Sinderen, D., Natural Transformation in Gram-Positive Bacteria and Its
Biotechnological Relevance to Lactic Acid Bacteria. *Annual Review of Food Science and*
Technology **2022,** *13* (Volume 13, 2022), 409-431.

73.   Clewell, D. B.; Weaver, K. E.; Dunny, G. M.; Coque, T. M.; Francia, M. V.; Hayes, F.,
Extrachromosomal and Mobile Elements in Enterococci: Transmission, Maintenance, and
Epidemiology. In *Enterococci: From Commensals to Leading Causes of Drug Resistant*

Infection, Gilmore, M. S.; Clewell, D. B.; Ike, Y.; Shankar, N., Eds. Massachusetts Eye and Ear
Infirmary: Boston, 2014.

74.   García-Solache, M.; Rice, L. B., The Enterococcus: a Model of Adaptability to Its
Environment. *Clin Microbiol Rev* **2019,** *32* (2).

75.   Bodmer, W. F., Integration of deoxyribonuclease-treated DNA in bacillus subtilis
transformation. *The Journal of general physiology* **1966,** *49* (6), 233-258.

76.   Santos-Lopez, A.; Marshall, C. W.; Haas, A. L.; Turner, C.; Rasero, J.; Cooper, V. S.,
The roles of history, chance, and natural selection in the evolution of antibiotic resistance. *Elife*
2021, *10*, e70676.

77.   Rebolleda-Gómez, M.; Travisano, M., Adaptation, chance, and history in experimental
evolution reversals to unicellularity. *Evolution* **2019,** *73* (1), 73-83.

78.   Mao, J.; Lu, T., Population-Dynamic Modeling of Bacterial Horizontal Gene Transfer by
Natural Transformation. *Biophysical Journal* **2016,** *110* (1), 258-268.

79.   Kent, A. G.; Vill, A. C.; Shi, Q.; Satlin, M. J.; Brito, I. L., Widespread transfer of
mobile antibiotic resistance genes within individual gut microbiomes revealed through bacterial
Hi-C. *Nature Communications* **2020,** *11* (1), 4379.

80.   Palleja, A.; Mikkelsen, K. H.; Forslund, S. K.; Kashani, A.; Allin, K. H.; Nielsen, T.;
Hansen, T. H.; Liang, S.; Feng, Q.; Zhang, C.; Pyl, P. T.; Coelho, L. P.; Yang, H.; Wang, J.;
Typas, A.; Nielsen, M. F.; Nielsen, H. B.; Bork, P.; Wang, J.; Vilsbøll, T.; Hansen, T.;
Knop, F. K.; Arumugam, M.; Pedersen, O., Recovery of gut microbiota of healthy adults
following antibiotic exposure. *Nature Microbiology* **2018,** *3* (11), 1255-1265.

81.   Yassour, M.; Jason, E.; Hogstrom, L. J.; Arthur, T. D.; Tripathi, S.; Siljander, H.;
Selvenius, J.; Oikarinen, S.; Hyöty, H.; Virtanen, S. M., Strain-level analysis of mother-to-child
bacterial transmission during the first few months of life. *Cell host & microbe* **2018,** *24* (1), 146-
154. e4.

903