Uncovering the electrical synapse proteome in retinal neurons via *in vivo* proximity labeling

Stephan Tetenborg[1,8], Eyad Shihabeddin[1,9], Elizebeth Olive Akansha Manoj Kumar[1,9], Crystal L. Sigulinsky[2,9], Karin Dedek[3,4], Ya-Ping Lin[1], Fabio A. Echeverry[5], Hannah Hoff[5], Alberto E. Pereda[5], Bryan W. Jones[2], Christophe P. Ribelayga[1], Klaus Ebnet[6], Ken Matsuura[7] & John O'Brien[1,8]

[1]College of Optometry, University of Houston, Houston, TX, United states.
[2]Moran Eye Center/Ophthalmology, University of Utah, Salt Lake City, Utah, United states.
[3]Animal Navigation/ Neurosensorics, Institute for Biology and Environmental Sciences, University of Oldenburg, Oldenburg, Germany.
[4]Research Center Neurosensory Science, University of Oldenburg, Oldenburg, Germany.
[5]Dominick P. Purpura Department of Neuroscience, Albert Einstein College of Medicine, Bronx, New York, United States.
[6]Institute-Associated Research Group: Cell Adhesion and Cell Polarity, Institute of Medical Biochemistry, ZMBE, University of Münster, Münster, Germany.
[7]Cell Signal Unit, Okinawa Institute of Science and Technology, Onna-son 904-0495, Japan.
[8]To whom correspondence should be addressed
[9]Contributed equally

Correspondence: stetenbo@Central.UH.EDU (S.T.) and jobrien3@Central.UH.EDU (J.OB.)

Research article, neuroscience

Keywords: TurboID, AII amacrine cell, Electrical Synapse, Connexin 36

The authors declare no competing interests.

Abstract

Electrical synapses containing Connexin 36 (Cx36) represent the main means for direct electrical communication among neurons in the mammalian nervous system. However, little is known about the protein complexes that constitute these synapses. In the present study, we applied different BioID strategies to screen the interactomes of Connexin 36 and its zebrafish orthologue Cx35b in retinal neurons. For *in vivo* proximity labeling in mice, we took advantage of the Cx36-EGFP strain and expressed a GFP-nanobody-TurboID fusion construct selectively in AII amacrine cells. For *in vivo* BioID in zebrafish, we generated a transgenic line expressing a Cx35b-TurboID fusion under control of the Cx35b promoter. Both strategies allowed us to capture a plethora of molecules that were associated with electrical synapses and showed a high degree of evolutionary conservation in the proteomes of both species. Besides known interactors of Cx36 such as ZO-1 and ZO-2 we have identified more than 50 new proteins, such as scaffold proteins, adhesion molecules and regulators of the cytoskeleton. Moreover, we determined the subcellular localization of these proteins in mouse retina and tested potential binding interactions with Cx36. Amongst these new interactors, we identified signal induced proliferation associated 1 like 3 (Sipa1l3), a protein that has been implicated in cell junction formation and cell polarity, as a new scaffold of electrical

42 synapses. Interestingly, Sipa1l3 was able to interact with ZO-1, ZO-2 and Cx36, suggesting a
43 pivotal role in electrical synapse function. In summary, our study provides the first detailed view
44 of the electrical synapse proteome in retinal neurons, which is likely to apply to electrical synapses
45 elsewhere.
46

Introduction

With the discovery of acetylcholine as the first neurotransmitter at the beginning of the 20th

century, it was widely accepted that chemical synaptic transmission is the only means of

communication for neurons. Decades later, several authors independently demonstrated

evidence for the existence of electrical synapses in invertebrates as well as in vertebrates, and

subsequent studies identified gap junctions, which are composed of clusters of intercellular

channels, as the morphological substrates of these synapses (1, 2). Today it is considered

common knowledge among neuroscientists that both types of synapses coexist in the nervous

system and that each of them provides unique features essential for accurate signal processing.

Yet, compared to the efforts that have been made to study protein complexes underlying chemical

synapse function and regulation, we know fairly little about the proteome that constitutes an

electrical synapse (3).

In contrast to chemical synapses, electrical synapses don't rely on neurotransmitters and specific

receptor molecules but instead make use of intercellular channels to convey neural signals

bidirectionally in the form of ionic currents (4). This seemingly simple mode of communication

allows for an instantaneous signal transmission that synchronizes the activities of electrically

coupled neurons giving rise to complex brain functions such as network oscillation, hormone

secretion or coordination of motor functions (5, 6).

In the mammalian nervous system, electrical synapses are mainly, but not exclusively formed by

Connexin 36 (Cx36), a connexin isoform that is highly specific for neurons (7-9) and often referred

to as the "major neuronal connexin". One exception to the rule are the insulin producing beta cells

in the pancreas (10). In general, Cx36 gap junction channels are known to exhibit a very low

single channel conductance (11) but a remarkable degree of plasticity that allows neurons to

regulate the number of open channels and hence the strength of synaptic transmission in order

to adapt to environmental stimuli (12). A prime example of a neuron that is heavily coupled via

Cx36 and in which the signal transduction pathways that regulate electrical coupling have been

well described is the AII amacrine cell (13-15). The AII cell is a glycinergic interneuron in the

mammalian retina that is electrically coupled to neighboring AII amacrine cells and to ON Cone

bipolar cells (16). The main task of this cell is to collect neural signals that originate in rods and

relay them into the cone pathway. This circuit is referred to as the 'primary rod pathway' and it

forms the neurophysiological basis for scotopic (dim light) vision. Within the primary rod pathway,

electrical coupling of AII amacrine cells is regulated in an activity dependent manner (13). The

opening of Cx36 channels in AII amacrine cells requires the activation of calcium calmodulin

dependent kinase II (CaMKII), which is triggered by the influx of calcium through non-synaptic

NMDA receptors. Once CaMKII is activated, it binds to Cx36 and phosphorylates the intracellular

domains causing the channel to open (13, 17, 18). These activity driven changes clearly resemble

long-term potentiation as it has been described for chemical synapses in hippocampal neurons

(19). Another, almost obvious, but not less important, parallel between electrical and chemical

synapses, is the their basic "architecture": 1) Glutamate receptors, for instance, interact with

PSD95 (20), a membrane associated guanylate kinase (MAGUK) that is similar in function and

domain structure to ZO-1, another MAGUK and well-known interactor of Cx36. 2) Electrical

synapses are surrounded by adhesions (21). 3) As described for glutamatergic synapses,

electrical synapses also harbor electron dense structures that appear to consist of a dense

assemblage of synaptic proteins (22, 23). However, the composition of this density or the type of

adhesion molecule that is necessary to specify where an electrical synapse is formed still remain

a mystery and require sophisticated proteomics to solve.

The task of identifying compartment specific proteomes or interactomes for a given protein of

interest (POI) with standard techniques has always been challenging due to the abundance of

non-specific background and the requirement to preserve intact protein complexes. To overcome

these limitations, several laboratories have developed novel proximity techniques such as

TurboID (24, 25), which utilize promiscuous biotin ligases to biotinylate neighboring proteins within

a radius of 5 nm of the POI. Like any other tag, TurboID can be fused to the POI and is then

expressed in cells, where it is allowed to biotinylate proximal proteins that are subsequently

isolated with streptavidin beads under stringent buffer conditions. The ability to directly label

"associated" proteins with an affinity tag combined with the fast enzyme kinetics have made

TurboID a valuable tool for modern *in vivo* proteomics. Previously, BioID and TurboID have been

used to study the proteomes of chemical synapses and even contact sites of astrocytes and

neurons in nematodes and mice (26-28).

In the present study, we have applied two different TurboID approaches to uncover the

electrical synapse proteome in the mouse and the zebrafish retina. Contrasting these two

model organisms allowed us to expose essential components of the molecular architecture of
vertebrate electrical synapses. Beyond known components of electrical synapses, such as ZO-1
and ZO-2, our screen identified a plethora of new molecules that were not previously shown to
associate with Cx36/Cx35b. These include components of the endocytic machinery, adhesion
molecules, regulators of the cytoskeleton, membrane trafficking, as well chemical synapse
proteins. Overall, our screen showed that many components of electrical synapses in zebrafish
and mouse retinas were highly conserved, in particular the ZO proteins and the endocytosis
components, indicating a high degree of evolutionary conservation that suggests a critical
functional role for these proteins.

Results

Two TurboID approaches allow a glimpse at the electrical synapse proteome in zebrafish
photoreceptors and AII amacrine cells of the mouse retina.

In this study, we have adopted two *in vivo* BioID approaches to gain a more comprehensive
understanding of protein complexes that are associated with the neuronal connexin Cx36 and its
zebrafish orthologue Cx35b in retinal neurons. To implement BioID in zebrafish, we generated a
transgenic line expressing a Cx35b-V5-TurboID fusion protein (we used an internal insertion site
as previously described (29)) under the control of Cx35b promoter (Figure 1 A). As shown in
control experiments, the expression of this construct recapitulated the expression pattern of
wildtype Cx35b in photoreceptors (30) which were labelled with ZPR1, indicating that Cx35b-V5-
TurboID is correctly targeted to synapses (Figure 1B). The inner retina, however, displayed much
weaker expression of the fusion construct, which is why we anticipated that most of the candidates
we detected with this strategy are proteins that are associated with Cx35b in photoreceptors. To
induce efficient biotinylation in zebrafish we injected 10mM biotin solution intraperitoneally on
three consecutive days and isolated biotinylated proteins for mass spectrometry (Figure 1B and
C). For the analysis of mass spec data sets we compared the abundance of candidate proteins
from the Cx35b-V5-TurboID strain to wildtype samples that did not express TurboID. In Cx35b-
V5-TurboID samples, but not in wild type retinas, we found several known interactors of the Gjd2
family such as scaffold proteins including Tjp1a, Tjp1b, Tjp2a, Tjp2b, Mupp1, Psd95 and Af6
(Figure 1D) (31-34). In addition to these candidates, we identified proteins that are involved in
membrane trafficking including Syt4, Sec22B, components of the endosomal trafficking
machinery Snap91 and Eps15I1a, and an additional Cx36 orthologue termed Cx34.7, which has

previously been localized to photoreceptors in bass retina (35). The detection of this Gjd1 isoform

via BioID was somewhat surprising, as Cx34.7 was shown to form gap junctions that are

separated from Cx35b containing clusters in bass photoreceptors. To determine the subcellular

localization of Cx34.7 and its connection to Cx35b in zebrafish photoreceptors, we generated a

transgenic Cx34.7-GFP strain that expressed an internally GFP-tagged fusion protein under the

control of the Cx34.7 promoter. In this strain, we found that Cx34.7 was only expressed in the

outer plexiform layer (OPL) where it was associated with Cx35b at some gap junctions (Figure

1E). Thus, besides known interactors of the Gjd family, our *in vivo* BioID approach identified an

additional connexin isoform that is connected to Cx35b in photoreceptors.

To unravel the Cx36 interactome in the mouse retina we initially delivered a Cx36-V5-TurboID

fusion via intravitreal AAV injections into retinal neurons. This construct was similarly designed to

the Cx35b clone but regulated by a short human synapsin promoter to drive expression in the

neural retina (Figure 2A). Unfortunately, this strategy produced massive overexpression artefacts

and resulted in an accumulation of Cx36-V5-TurboID in ganglion cell somas, which would have

made an interactome analysis of a synaptic protein impossible. To solve this issue and to ensure

that TurboID is targeting electrical synapses, we took advantage of the GFP-directed TurboID

approach (36) and the Cx36-EGFP transgenic mouse strain (37-39). The GFP-directed TurboID

strategy was initially developed as a modular system for zebrafish by Xiong et al., and utilizes and

a destabilized GFP nanobody that is fused to TurboID (36) to direct the biotin ligase towards a

given protein of interest carrying a GFP-Tag. To further increase the specificity of our approach

and identify the Cx36 interactome within a single type of inhibitory interneuron, the AII amacrine

cell, we expressed V5-TurboID-dGBP under control of the HKamac promoter (40). As shown

previously, in retinas that were infected with AAVs carrying an HKamac GFP vector, we observed

GFP expression mainly in AII amacrine cells but not in bipolar cells (Figure 2B, lack of GFP

expression in SCGN labelled neurons). In our hands, we also observed GFP labeling in horizontal

cells, which was neglectable for our paradigm, since these neurons do not express Cx36 (41) and

because dGBP is degraded unless it is bound to GFP (42). In an initial experiment we injected

AAVs carrying HKamac_V5-TurboID-dGBP intravitreally into Cx36-EGFP mice and tracked the

localization of the construct two weeks post injection. As expected, all V5 labeled puncta

colocalized with Cx36-EGFP, indicating that V5-TurboID-dGBP reached electrical synapses in AII

amacrine cells. In the next experiment, we injected a 5 mM biotin solution subcutaneously on 4

consecutive days, 3 1/2 weeks post infection. This treatment was sufficient to induce efficient

biotinylation (Figure 1B, lower panel and C), and allowed us to capture a plethora of molecules

that were associated with Cx36. To distinguish these proteins from background, we included wild

type mice injected with AAV HKamac_V5-TurboID-dGBP throughout the entire experiment and

compared the abundance of all proteins we detected. A mass spec hit that was three times or

even more abundant in the Cx36-EGFP condition was considered a candidate that is likely to be

associated with Cx36. We illustrated the relationship of all proteins that fell into this category in a

string diagram (Figure 2D). Among the most abundant proteins in our screen, we identified the

ZO proteins ZO-1 and ZO-2 and signal induced proliferation associated 1 like 3 (Sipa1l3), a PDZ

domain containing protein that has been implicated in the regulation of cell adhesion and cell

polarity. In addition to these candidates, we also identified proteins that are involved in

endocytosis, membrane trafficking, regulation of the cytoskeleton, cell adhesion and chemical

synapses. All of these hits covered different functional categories of proteins that have been

related to connexins (43-48) and thus appear to represent an authentic interactome for a neuronal

gap junction protein.

Localization of BioID hits at Cx36 containing gap junctions in the AII amacrine cell

In the next set of experiments, we validated the most abundant candidates from our list in GFP

expressing AII amacrine cells. We categorized the proteins based on the cell biological function

they serve. In line with the mass spec data, we observed the most frequent colocalization for

Cx36 with scaffold proteins including ZO-1, ZO-2, Cingulin and Sipa1l3 (Figure 3A, arrows).

Although ZO-1, ZO-2 and Cingulin are known components of electrical synapses (34, 44), Sipa1l3

has not been shown before to associate with Cx36, yet it colocalized with the connexin to a similar

extent as the ZO proteins, suggesting that it could be essential for synapse formation or

stabilization. In addition to scaffolding proteins, we confirmed an association with several adapter

proteins of the endocytic machinery, including Eps15l1, Snap91, Hip1r and Itsn1 (Figure 3B,

arrows). In comparison to the scaffold proteins, however, the colocalization of Cx36 with each of

these endocytic components was clearly less frequent and more heterogenous, which appears to

reflect different stages in the life cycle of Cx36. As most proteins we identified in this initial screen

broadly labeled the entire inner plexiform layer (IPL – the inner synaptic layer of the retina), we

performed a colocalization analysis to exclude a random overlap with Cx36. We quantified the

number of Cx36 plaques that contained Cgn, Sipa1l3, Eps15l1 and Hip1r (Figure 3C & D) and

compared the extent of colocalization to a control in which we flipped the Cx36 channel

horizontally. For each candidate, we found a significant reduction in colocalization when the

original images were compared to the flipped control, indicating that the proteins we have

identified are actual components of electrical synapses.

Besides synaptic scaffolds and endocytic adapter proteins we tested a variety of additional hits,

which include synaptojanin 2 binding protein (Sj2bp), synaptotagmin 4 (categorized as

"trafficking") and several proteins that regulate actin dynamics such as G-protein regulated

inducer of neurite outgrowth 1 (Gprin1), dedicator of cytokinesis 7 (Dock7) and microtubule

associated protein 6 (Map6). All of these proteins showed considerable colocalization with Cx36

in AII amacrine cell dendrites (Figure 4A, B). We also tested components of chemical synapses

including GluR2-3, the scaffold Shank2 (Figure 4D) and Bai1 (Figure 4C) a synaptic adhesion

molecule implicated in excitatory synapse formation (49). We found that these proteins showed a

partial overlap with Cx36 at the periphery of each junction.

Identification of novel Cx36 binding partners

Thus far, we have gained significant insight into the Cx36 interactome and identified several new

molecules as components of electrical synapses in AII amacrine cells. To understand how these

molecules are organized at the synapse we tested protein-protein interactions in co-transfected

HEK293T cells. Interestingly, two of the novel proteins we detected in AII cells contain PDZ

domains and thus may directly bind to Cx36 like ZO-1 and ZO-2 (Figure 5B, GFP-Trap IP with

both ZO proteins served as a positive control); these proteins are Sipa1l3 and Sj2bp. We found

that Myc-Sipa1l3 interacted with all major components of electrical synapses: ZO-1, ZO-2 and

Cx36 in co-transfected HEK293T cells. The overlap of Sipa1l3 and the ZO proteins was confined

to cell cortices and to actin like fibers in the cytoplasm (Figure 5A). Additionally, we performed IP

experiments and demonstrated that Cx36 binds to Sipa1l3 in a PDZ dependent manner. A

truncated version of Cx36 lacking the PDZ binding motif (Cx36/S318Ter) failed to interact with

Sipa1l3 in IP experiments. In co-transfected cells we observed that Sj2bp, which contains a single

PDZ domain and a transmembrane domain (50), colocalized with Cx36 but not with the

Cx36/S318Ter, indicating that the interaction of these proteins requires the PDZ domain. In IP

experiments using FLAG tagged Sj2bp as a bait we were unable to coprecipitate Cx36, which

was rather surprising as both proteins clearly colocalized in co-transfected HEK293T cells. One

possible explanation for the lack of a detectable interaction is that Sj2bp displays a low affinity for

the PDZ binding motif (PBM) of Cx36, as it is the case for ZO-1 (45). We have previously shown

that binding of Cx36 to ZO-1 can be artificially enhanced by a simple deletion of amino acids 313-

319 within the C-terminus, creating a new PBM with the following sequence: RTYV (51). We

repeated the IP with this mutant and observed substantial binding to FLAG-Sj2bp (Figure 5B),

suggesting that Cx36 and Sj2bp are compatible interactors. In addition to the candidates we

overexpressed to verify direct binding interactions, we found that Cx36 colocalized with several

endogenous HEK293T proteins that also occurred in our AII cell specific data set, including

Eps15l1, Gprin1 and Sec22b (Figure 5C).

Molecular architecture of the AII amacrine cell/On cone bipolar cell gap junction.

AII amacrine cells form two different sets of gap junctions, with neighboring AII amacrine cells and

with ON cone bipolar cells to convey neural signals that originate in rod photoceptors into the

cone pathway (Illustrated in figure 6A) (9). Ultrastructural studies have revealed an asymmetry in

the cytoplasm of AII/ON cone bipolar cell junctions (AII/ONCBC), which exhibits an electron dense

structure (termed fluffy material) in the AII amacrine cell but not in the cone bipolar cell, hinting at

differences in the molecular composition of both sites of the synapse (23, 52). Moreover, apart

from accessory proteins that bind to the channel, AII/ONCBC gap junctions are likely to be distinct

from AII/AII gap junctions since they recognize ON cone bipolar cells as the correct synaptic

partner. These two characteristics of the AII/ONCBC gap junction can only be explained by a

unique composition of the synapse proteome. We reasoned that the electron density in the AII

cell could harbor several proteins we identified in this study. To test this hypothesis, we triple

labelled AAV/HKamac_GFP infected retinas with antibodies to Scgn, Cx36 and the main

candidates from our screen. With this strategy it was possible for us to visualize AII/ONCBC

contacts (Figure 6 D, arrows) and the precise localization of each candidate. To illustrate the

localization of AII/ONCBC gap junctions in a comprehensive way, we reconstructed the

morphology of AII amacrine cells using serial section transmission EM imaging. As seen in figure

6B, most AII/ONCBC gap junctions (shown in yellow) were located above homologous AII/AII

contacts (Cyan) and contained electron densities on both sides (Figure 6C).

The scaffold proteins ZO-1, ZO-2, Cgn and Sipa1l3 localized at AII/ONCBC contacts suggesting

that they are a part of the electron density in AII cells (Figure 6D, arrows). In addition to these

proteins, we also confirmed that AII/Cone bipolar cell gap junctions contained Syt4, Eps15l1,

Sj2bp and Gprin1. Finally, we addressed the exact localization of the G protein coupled adhesion

receptor Bai1, a synaptic adhesion molecule that has been implicated in excitatory synapse

formation (49). Interestingly, we observed Bai1 at AII/ONCBC contacts, in the vicinity of Cx36,

which is consistent with the notion of synaptic adhesion molecules as an integral part of electrical
synapses (21, 53). This finding further implies that Bai1 might play a role in the formation of
AII/ONCBC contacts. However, further knock-out studies will be necessary to address the
concrete function of this adhesion molecule. Finally, we addressed the subcellular distribution of
Sipa1l3 at AII/ONBC gap junctions. We noticed that these junctions often displayed a ring like
shape, that contained Sipa1l3 in the center of each ring (Figure 6D). One possible interpretation
of this distinct sub-synaptic distribution is that Sipa1l3 is interacting with a type of adhesion
molecule that is directly adjacent to the gap junction.

Electrical synapse scaffolds are targeted to AII amacrine cell/ON cone bipolar cell contacts
in the absence of Cx36.

Previous studies have shown that ZO-1 localizes to neuronal contact sites even in the absence
of connexins (33). This finding is consistent with a study by Meyer et al., (2014), who reported
that Cx36-EGFP containing gap junctions colocalize with ZO-1 at AII/ONCBC contacts despite a
C-terminal tag in the fusion protein that is known to interfere with PDZ domain mediated
interactions (38, 51). One possible interpretation of these observations is that ZO-1, a protein with
multiple protein-protein interaction domains, is connected to additional components of the gap
junction such as adhesion molecules, the actin skeleton or other MAGUKs that retain the scaffold
at the synapse. We wondered if a similar principle applies to the scaffold proteins that were
identified in this study and tested the localization of ZO-2 and Sipa1l3 in Cx36 KO retinas that
were infected with AAV/HKamac_GFP to visualize AII/ONCBC contacts. Here, we observed that
ZO-2 and Sipa1l3 showed a similar effect as ZO-1 and still localized to AII/ONCBC contacts in
Cx36 KO retinas (Figure 7A-C). Additionally, we quantified the density and size of immunoreactive
Sipa1l3 and ZO-1 in the inner plexiform layer of WT and Cx36 KO mouse retina. While the size
of ZO-2 and Sipa1l3 puncta were unchanged in Cx36 KO mice, the density of both ZO-2 and
Sipa1l3 puncta were reduced and the colocalization of ZO-2 and Sipa1l3 was reduced in Cx36KO
retina. Thus, loss of Cx36 resulted in a quantitative defect in formation of electrical synapse
density complexes.

Discussion

In the present study, we have used TurboID to uncover the electrical synapse proteome in retinal neurons. Our screen identified a plethora of molecules that were associated with the neuronal connexins Cx36/Cx35b in zebrafish and mice such as adhesion molecules, scaffold proteins, chemical synapse proteins, components of the endocytic machinery and cytoskeleton associated proteins (Illustrated in Figure 7 E-F). The presence of ZO proteins and endocytosis components in the Cx36 and the Cx35b interactome data suggests a certain degree of conservation in the proteome of electrical synapses. An apparent difference between species, however, is the existence of additional ZO variants and Cx36 homologues in the zebrafish genome (33), which are also represented in our Cx35b interactome. In contrast to the bass retina, where it was shown that Cx35 and Cx34.7 form two distinct circuits (35), we found that zebrafish photoreceptors formed gap junctions that contained both connexins. Cell culture experiments have confirmed that Cx35 and Cx34.7 are able to form heterotypic channels (54) that exhibit slightly altered voltage sensitivities compared to the homotypic channels. How exactly these gap junction channels in zebrafish photoreceptors are configured and which exact cell types they connect remains to be determined.

Among other proteins that were detected in this study, we identified Sipa1l3 as a novel scaffold protein for electrical synapses. Interestingly, Sipa1l3 was also described as a component of post synaptic densities of glutamatergic synapses in hippocampal neurons, and is known interact with Fezzins, which occur in a complex with Shank3 (55). Additionally, we found that Sipa1l3 was able to interact with Cx36 and the ZO proteins. This raises the question of what potential function Sipa1l3 might serve given the abundance of colocalization with Cx36 in AII amacrine cells? Genetic screens have shown that mutations in the Sipa1l3 gene result in abnormal eye and lens development and a decrease in the formation of cell adhesions. Like other members of the Sipa1l family, Sipa1l3 also contains a RAP GTPase activating protein (GAP) domain that is known to regulate the activity of RAP proteins, which in the GTP-bound form impacts processes such as adhesion and actin dynamics (56). One way in which Sipa1l3 might influence the formation of electrical synapses might be as a regulator of Rap1 and its effector proteins. To determine the exact function of Sipa1l3 further studies on KO mice will be required.

One striking aspect of the Cx36 interactome in AII amacrine cells is the abundance of the endocytosis machinery. Considering the short half life time of Cx36 of 3.1 h (29) and the fact that

347 the turnover of electrical synapse proteins is a steady-state process (43), it makes sense that the
348 endocytic machinery is that well represented in our interactome data. Thus, besides
349 phosphorylation, it seems possible that the cellular regulation of turnover mechanisms could
350 function as an additional means to adjust the strength of electrical coupling between AII amacrine
351 cells.

353 Is Bai1 necessary to form electrical synapses in the primary rod pathway?

How and when it is determined where electrical synapses are formed is currently unknown. It has
been suggested that neuronal adhesion molecules serve as cues that wire the correct neurons
together (3). This task also has to be accomplished in the primary rod pathway, where AII
amacrine cells form synapses among each other and with ON Cone bipolar cells. Interestingly,
we identified the brain specific angiogenesis inhibitor 1 (Bai1) between AII amacrine and ON cone
bipolar cell gap junctions, which could suggest that Bai1 is necessary to connect these cells.
Recent reports have described Bai1 as a synaptogenic enzyme capable of regulating intracellular
signaling pathways that determine synapse formation. (49, 57). Thus, besides merely functioning
as a cell adhesion molecule, Bai1 might actively support synaptogenesis in AII amacrine cells at
a level that is hierarchical to Cx36 and its associated scaffolds, which would explain why these
cells form contacts with ON Cone bipolar cells even in the absence of Cx36.

366 Cell type-specific differences in the Cx36 interactome?

Interestingly, our interactome in AII amacrine cells showed no signs of the scaffolds Mupp1 and
Af6, although these candidates were shown to colocalize and interact with Cx36 in mammalian
retinas and several other brain regions (31). Moreover, both proteins also showed up in our Cx35b
interactome in zebrafish retina. One feasible explanation is that AII cell gap junctions simply lack
these scaffolds and that they may instead be used for electrical synapses of other neurons like
bipolar cells, which our approach did not consider. In general, it will be interesting to understand
how well certain components of the electrical synapse proteome are preserved between cell
types. Endocytosis and ZO proteins, for instance, are unlikely to be cell-type specific interactors
of gap junction channels and are more likely to be ubiquitous constituents of electrical synapses,
given their critical function in the constant turnover of Cx36. This is also reflected in the
evolutionary conservation of these components in the proteomes from zebrafish and mice
(supplementary figure 3). Bai1, on the other hand, a protein that is needed to establish specific
synaptic contacts, may only occur in certain electrical synapses.

Material and Methods

Animal husbandry of zebrafish

Maintenance and breeding of zebrafish was conducted under standard conditions (58). Wild type (AB) zebrafish were purchased from the Zebrafish International Resource Center (ZIRC, Eugene, OR, USA). Fish were maintained on a 14 h light/10 h dark cycle. All procedures were performed in accordance with the ARVO statement on the use of animals in ophthalmic and vision research and US Public Health Service guidelines and have been reviewed and approved by the Institutional Animal Care and Use Committees at the University of Texas Health Science Center at Houston and the University of Houston.

Tol2 mediated transgenesis

Transgenic Cx35b-V5-TurboID fish were generated using the Tol2 system and a donor vector containing 4.5 kb of the Cx35b promoter, exon1 of the Cx35b gene and the open reading frame of Cx35-V5-TurboID. The coding sequence of V5-TurboID (25) was cloned into a region of the Cx35b gene encoding the C-terminal tail. V5-TurboID was inserted between leucine 285 and proline 286 to expose the 19 C-terminal amino acids of Cx35b. Additionally, the donor vector contained a Myl7 promoter driving mCherry expression in the heart to select larvae in which the construct was integrated into genome. Transgenic Cx34.7-oxGFP fish were generated in the same way using a donor vector containing 2.8 kb of the Cx34.7 promoter and the coding sequence of Cx34.7 with oxGFP inserted between aspartate 283 and methionine 284 in the C-terminus. To generate transgenic fish, the donor vector was co-injected with Tol2 transposase mRNA into embryos in the 1 cell stage as described previously (59). Transgenic individuals were raised to adulthood and mated.

Animal Husbandry of mice

The experiments in this study were conducted with wildtype mice (C57BL/6J), the Cx36-EGFP strain (37-39) (Kindly provided by Dr. Hannah Monyer) and Cx36 Knock out mice (60) (Kindly provided by Dr. David Paul). Animals of ages 2-12 months were used for experiments. All procedures were performed in accordance with the ARVO statement on the use of animals in ophthalmic and vision research and US Public Health Service guidelines and were approved by the Institutional Animal Care and Use Committee at the University of Houston.

AAV constructs and intravitreal injections

All AAV constructs used in this study were generated with the pAAV-9-(5)-hSYN-CAMKII-GFP vector kindly provided by Dr. Neal Waxham. The coding sequence of the AII amacrine cell specific AII promoter Hkamac was previously described by Khabou et al., (2023), and synthesized as a 654 base pair gBlocks® fragment by IDT genomics. The hSYN promoter and the coding sequence of CaMKII in the pAAV-9-(5)-hSYN-CAMKII-GFP vector were excised via XbaI and EcoRV and the coding sequence of V5-TurboID-dGBP (36, 51, 61) and GFP were fused to the AII promoter and integrated into the vector via Gibson assembly. Titers of AAV particles for different AAV constructs varied ranging from 8×10^{12} to 1×10^{13} gc/ml. For each experimental condition similar amounts of virus particles were injected. All virus particles were generated at the Baylor College of Medicine gene vector core facility. All AAV vectors that were used in this study will be made available via Addgene. Prior to each virus injection, mice were anesthetized with 3.5-3.9% isoflurane using the SomnoSuite/Mouse STAT Pulse Oximeter (SS-01-03, Kent Scientific Corporation). Once the animal was anesthetized, a small incision next to the pupil was made using a 30 G needle. Afterwards, 1.5 µl of the virus solution was carefully injected intravitreally. Eyes were collected 2-4 weeks after each experiment.

DNA constructs

FLAG-SJ2BP and Myc-SIPA1L3 constructs were used in previous studies (50, 62). Venus-ZO-1 (#56394) and GFP-ZO-2 (#27422) were purchased from Addgene. The FLAG-Gprin1 (Clone: OMu09339D) and FLAG-EPS15L1 (Clone: OMu22488D) constructs were purchased from Genscript. All Cx36 pcDNA3.1 expression vectors were generated in previous studies (51, 61, 63).

***In vivo* Biotinylation**

In vivo biotinylation in zebrafish. Prior to biotin injections, adult zebrafish were anesthetized with 0.02% Tricaine and immobilized in a small wax mold. A small incision was made into the abdomen and 30 µl of a 5mM biotin (solved in PBS) solution was injected intraperitoneally using a Hamilton syringe. This procedure was repeated for 2 consecutive days. Afterwards the fish were sacrificed by a cold shock and the retinas were extracted. The streptavidin pull down was carried out as previously described (51, 61). 80 retinas for each condition (Wild type vs. Cx35b-V5-TurboID) were used.

447 *In vivo biotinylation in mouse.* 3 ½ weeks after intravitreal virus injections mice were anesthetized
448 with isoflurane and 1ml of a 5mM biotin was injected subcutaneously for 4 consecutive days. 17
449 retinas of each condition (Cx36-EGFP vs. wild type) were used for the streptavidin pull down.
450

LC/MS/MS analysis
For mass spectrometry analysis of streptavidin pull downs, samples were concentrated on a
acrylamide/bis-acrylamide stacking gel and processed via In-gel digestion. An aliquot of the tryptic
digest (in 2% acetonitrile/0.1% formic acid in water) was analyzed by LC/MS/MS on an Orbitrap
Fusion™ Tribrid™ mass spectrometer (Thermo Scientific™) interfaced with a Dionex UltiMate
3000 Binary RSLCnano System. Peptides were separated onto an analytical C18 column (100µm
ID x 25 cm, 5 µm, 18Å Reprosil-Pur C18-AQ beads from Dr Maisch, Ammerbuch-Entringen,
Germany) at flow rate of 350 nl/min. Gradient conditions were: 3%-22% B for 90 min; 22%-35%
B for 10min; 35%-90% B for 10 min; 90% B held for 10 min, (solvent A, 0.1 % formic acid in water;
solvent B, 0.1% formic acid in acetonitrile). The peptides were analyzed using data-dependent
acquisition method, Orbitrap Fusion was operated with measurement of FTMS1 at resolutions
120,000 FWHM, scan range 350-1500 m/z, AGC target 2E5, and maximum injection time of 50
ms; During a maximum 3 second cycle time, the ITMS2 spectra were collected at rapid scan rate
mode, with HCD NCE 34%, 1.6 m/z isolation window, AGC target 1E4, maximum injection time
of 35 ms. Dynamic exclusion was employed for 20 seconds.
466

Data processing and analysis
The raw data files were processed using Thermo Scientific™ Proteome Discoverer™ software
version 1.4. Spectra were searched against the Uniprot-Mouse or Uniprot-Zebrafish database
using Sequest. Search results were filtered to 1% FDR for strict conditions and 5% for relaxed
conditions using Percolator. Trypsin was set as the enzyme with maximum missed cleavages set
to 2. For peptide mapping, MS tolerance was set to 10 ppm, and MS/MS tolerance to 0.6 Da.
Carbamidomethylation of cysteine residues was used as a fixed modification, while oxidation of
methionine, N-terminal acetylation, and phosphorylation of serine and threonine were set as
variable modifications.
476

Bioinformatics
A ratio of Cx36 conditions to the control was used to identify any proteins that were enriched
greater than 3-fold in mouse retina and 2-fold in zebrafish retina. Proteins above this threshold
were run through the Cytoscape (V3.10.2) plugin ClueGo (V2.5.10) to identify the GO Biological

Processes that were upregulated in each Cx36 condition across both zebrafish and mouse
species (64, 65). Pathways that show a p-value of less than 0.05 were used for analysis. To
determine protein-protein interaction networks, proteins above threshold were also run through
the standard STRING (V12.0) pipeline (66).

Cell culture

Human embryonic kidney 293 T cells (HEK293T/17; catalog #CRL-11268; ATCC, Manassas, VA,
USA) were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal
bovine serum (FBS), 1% penicillin and streptomycin, and 1% non-essential amino acids (all
Thermo Fisher Scientific, Rockford, IL, USA) at 37 °C in a humidified atmosphere with 5% CO_2.
For pull-down experiments 1 million cells were plated in 60mm dishes. Transfections were carried
out with Geneporter 2 as previously described (51).

Immunocytochemistry

Immunolabeling of transfected HEK293T cells was carried out as previously described (61, 63).
Transfected HEK293T expressing the proteins of interest were briefly rinsed in PBS and fixed in
2% paraformaldehyde (PFA) solved in PBS for 15 min at RT. After the fixation step, the coverslips
were washed three times in PBS for 10 min and incubated with the primary antibody solution
containing 10% normal donkey serum in 0.5% Triton X-100 in PBS overnight at 4 ºC .The following
antibodies were used: mouse anti Cx36, 1:500 (MAB3045, Millipore), rabbit anti SIPA1l3, 1:200
(30544-1-AP, Proteintech), anti GFP, 1:250 (A10262, Thermofisher), rabbit anti EPS15L1, 1:250
(PA5-65940, Thermofisher) and mouse anti V5, 1:500 (R960-2, Thermofisher), rabbit anti myc
(16286-1-AP, Proteintech). At the next day, the coverslips were washed 3x10 min in PBS and
incubated with the secondary antibodies diluted in 10% normal donkey serum in 0.5% Triton X-
100 in PBS at RT under light protected conditions. The following secondary antibodies were used:
from donkey, 1:500, conjugated with Cy3, Alexa488, Alexa568, or Alexa647 (Jackson
Immunoresearch, West Grove, PA). Afterwards coverslips were washed 3x10 min in PBS and
mounted with Vectashield containing DAPI (H-2000, Vector Laboratories Inc.)

Immunohistochemistry

Immunolabeling of vertical sections was carried out as previously described (67, 68). Mice were
anesthetized with isoflurane and sacrificed via cervical dislocation. The eyes were removed from
the animal and opened with a circular cut around the *ora serrata*. The lens was removed, eyes
were fixed with 2% PFA solved in PBS for 20 min at room temperature and washed 3x10 min in

PBS. Afterwards eyecups were stored in 30% sucrose at 4ºC overnight. On the next day eyecup preparations were embedded in Tissue-Tek O.C.T. cryomatrix (Sakura Finetek, Torrance, CA, USA) and stored at -20ºC. Eyecups were cut into 20 µm thin section and incubated at 37ºC for 30 min. Afterwards the slides were washed with PBS (2x5 min) and incubated in the primary antibody solution containing 10% normal donkey serum in 0.5% Triton X-100 in PBS overnight. The following primary antibodies were used: Cx36, 1:250 (Clone: 8F6, MAB3045, Millipore); Cx36, 1:250 (Clone: 1E5H5, 37-4600, Thermofisher); GFP, 1:250 (A10262, Thermofisher); ZO-1, 1:250 (Clone: ZO1-1A12, 33-9100, Thermofisher); ZO-2, 1:250 (71-1400, Thermofisher); CGN, 1:250 (PA5-5561, Thermofisher); SIPA1L3, 1:100 (30544-1-AP, Proteintech); EPS15L1, 1:250 (PA5-65940, Thermofisher); HIP1R, 1:250 (AB9882,Sigma); GluR2-3, 1:200 (07-598, Sigma); Gprin1, 1:100 (13771-1-AP, Proteintech); DOCK7, 1:100 (13000-1-AP, Proteintech); MAP6, 1:100 (NBP2-14220, Novus Biologicals); Synaptotagmin4, 1:100 (105 143, Synaptic Systems); SJ2BP, 1:100 (15666-1-AP, Proteintech); BAI1, 1:100 (NB110-81586, Novus Biologicals); SHANK2, 1:500 (Synaptic Systems, 162204); SCGN, 1:250 (CSB-PA020821LA01HU, ARP); SCGN, 1:500 (RD184120100, Biovendor); Vglut1, 1:500 (AB5905, Sigma); ITNS1, 1:100-250 (PA5-115432, Thermofisher). The next day, the slides were washed 3x10 min and incubated with secondary antibodies diluted in 10% normal donkey serum in 0.5% Triton X-100 in PBS for 1h at RT under light protected conditions. The following secondary antibodies were used: from donkey, 1:250, conjugated with Cy3, Alexa488, Alexa568, or Alexa647 or Dylight 405 (Jackson Immunoresearch, West Grove, PA). Afterwards the slides were washed 3x10 min and mounted with Vectashield (H-1900, Vector Laboratories, Inc.).

Immunoprecipitation

Immunoprecipitation experiments were carried out as previously described (46). 1.5 $\times 10^6$ HEK293T cells were plated on 60mm dishes and co-transfected at the next day with 2 µg of each vector using Geneporter2 (Amsbio) transfection reagent. 24h after transfection, cells were rinsed in phosphate buffered saline (pH 7.4) and transferred to a reaction tube. The cells were centrifuged at 5000g for 5 min and afterwards homogenized in immunoprecipitation buffer (IP buffer) containing 200mM NaCl, 2m EDTA, 1% Tx-100, 50mM Tris HCL (pH 7.5) supplemented with protease inhibitors (Roche) and 1mM DTT. The lysate was sonicated 3x for 30 seconds and incubated on ice for 30 min. Afterwards, the lysate was centrifuged at 10,000g for 10 min. 20 µl of ChromoTek GFP-Trap® agarose beads (gta, Proteintech) were applied to the supernatant and incubated overnight at 4ºC on a rotating platform. The next day the sample was centrifuged at 2500g for 5 min and washed three times in IP buffer. Bound proteins were eluted with 60 µl of 1x

Laemmli buffer (BioRad, 1610747) for 5 min at 95ºC. IP experiments for Myc-Sipa1l3 and FLAG-
Sj2bp were conducted under a slightly modified protocol. For IP experiments with Myc- Sipa1l3
0.5 µg of Sipa1l3 antibody (30544-1-AP, Proteintech) was applied to the cell lysate and incubated
overnight on a rotating platform. The next day, 40 µl of prewashed protein A/G beads (Pierce)
were applied to the cell lysate and incubated for 1h at 4 ºC. Afterwards the beads were washed 3
times and bound proteins were eluted with 1x Laemmli buffer. IP experiments with FLAG-Sj2bp
were performed with 20 µl ChromoTek DYDDDDK Fab-Trap® (ffa, Proteintech). Prior to the IP
cell lysates were centrifuged at 20,000g for 30 min at 4ºC. Afterwards, cell lysates were incubated
with agarose overnight and washed 3 times at 4ºC. Bound proteins were eluted with 60 µl of 1x
Laemmli buffer.
559
Western blot
Protein samples were separated via SDS-PAGE containing 7.5% (for Sipa1l3 blots) or 10%
polyacrylamide. Proteins were transferred to nitrocellulose membranes using the transblot turbo
system as previously described (51). Blots were blocked in 5% dried milk powder in TBST (20 mM
Tris pH 7.5, 150 mM NaCl, 0.2% Tween 20) for 30 min at RT and incubated overnight in the
primary antibodies diluted in blocking solution at 4ºC on a rotating platform. The following primary
antibodies were used: Cx36, 1:500 (Clone: 8F6, MAB3045, Millipore); Cx36, 1:500 (Clone:
1E5H5, 37-4600, Thermofisher), SIPA1L3, 1:4000 (30544-1-AP, Proteintech), GFP, 1:1000 (2956,
Cell Signaling), V5 1:1000 (R960-2, Thermofisher). Biotinylated proteins detected with high
sensitivity streptavidin-HRP, 1:1000 (Abcam). At the next day, blots were washed 3x10 min in
TBST and incubated with the secondary antibodies, diluted in the blocking solution as described
previously (51). The following secondary antibodies were used: goat anti mouse HRP, 1:1000
(34130) and goat anti rabbit, 1:1000 (34160, Thermofisher). Afterwards, the blots were washed
3x10 min in TBST and incubated with the ECL solution (32106, Thermofisher) for detection. ECL
was detected with the Biorad ChemicDoc™ MP Imaging System.
575
Confocal microscopy and image analysis
Fluorescence images were acquired with a confocal laser scanning microscope (Zeiss LSM 800)
using a 60x oil objective and the Airy scan function (Pixel size: 50 nm). Confocal scans were
processed using Fiji (ImageJ2 version 2.14.0/1.54f) software (69). The threshold was adjusted
using the *Triangle* function for the entire z-stack (7 slices; 1.08 µm). Two regions of interest (ROI)
with the size of 20.07 x 32.91 μm^2 were positioned to cover the ON and OFF IPL layers. Frequency
and size of the immunoreactive puncta were measured for each channel using the "analyze

particles" function. Colocalization of ZO-1 and Sipa1l3 was quantified using the *Colocalization*

plugin in Fiji. The number of colocalized puncta and the colocalization was measured using the

Analyze particle function with a size an exclusion criterion of >100^2 pixels. Following the

quantification, the size, density and percentage of the colocalized puncta were averaged for each

protein.

Volume construction

Retinal Connectome 2 (RC2) is an ultrastructural dataset acquired from the retina of a light-

adapted 5-month-old female C57BL/6J mouse (70). Methods concerning tissue acquisition and

processing, volume assembly, visualization, and annotation following that extensively detailed for

RC1 (71, 72). In short, the RC2 dataset spans the ganglion cell layer through outer nuclear layer

of a 0.25 mm-diameter field of retina. RC2 was constructed from 1335 serial TEM sections (70

nm thick), captured at 2.18 nm/pixel via Automated Transmission Electron Microscopy at 5000X

and combined with 31 optical sections intercalated throughout the volume. These optical sections

were probed for small-molecule signals for computational molecular phenotyping (73). Sections

were aligned into a single volume using the NCR ToolSet, which has since been replaced by

Nornir (RRID:SCR_016458). All protocols were in accord with Institutional Animal Care and Use

protocols of the University of Utah, the ARVO Statement for the Use of Animals in Ophthalmic

and Visual Research, and the Policies on the Use of Animals and Humans in Neuroscience

Research of the Society for Neuroscience.

Statistical analysis

Data sets acquired in this study were analyzed with GraphPad Prism 8. Data are shown as

mean ± SEM. Normality was tested using the Anderson–Darling and the D'Agostino-Pearson test.

Significance was tested using the two-tailed Mann–Whitney U test. For multiple comparisons a

one-way ANOVA was performed.

Acknowledgements

This project was supported by NIH grants R01EY012857 (J.O.), RF1MH120016 (A.E.P., J.O.),
and R21NS085772 (A.E.P., J.O.) and core grant P30EY007551. S.T. was funded by the *Deutsche*
Forschungsgemeinschaft (DFG) (TE 1459/1-1, Walter Benjamin stipend). E.S. was supported by
NIH training grant TL1TR003169 and individual grant F31EY034793. B.W.J. and C.L.S. are
supported by NIH grants R01EY028927 (to B.W.J.) and P30EY014800 (to the Moran Eye Center
Core); NSF grant 2014862; an Unrestricted Research Grant from Research to Prevent Blindness,
New York, NY to the Department of Ophthalmology & Visual Sciences, University of Utah; and an
Unrestricted Research Grant from Gabe Newell (to B.W.J.). This work is supported in part by the

Clinical and Translational Proteomics Service Center at the University of Texas Health Science
Center We would like to thank Li Li from the UT health mass spec. facility We would like to thank
Dr. Adam C. Miller and Dr. Jen Michel for the helpful discussions. We appreciate Nikki Brantley's
support in the cell culture lab.
623

Author contributions

625   S.T and J.O designed the research. S.T, E.S, E.O.A.M.K, H.H, F.A.E, C.L.S, Y.P.L and K.D
626   performed experiments. K.E and C.R contributed reagents and analytic tools. S.T wrote the paper.
627   All authors edited the paper.
628

Figure legends

Figure 1: Generation of a Cx35b-V5-TurboID zebrafish line. (A) Cartoon illustrating the generation of Cx35-V5-TurboID fish via Tol2-mediated transgenesis and outline of *in vivo* biotinylation experiments. To induce efficient proximity labeling, zebrafish were intraperitoneally injected with 30µl of 5mM biotin (PBS) for three consecutive days. Afterwards the animals were sacrificed, and the retinas were isolated for streptavidin pull-downs. **(B)** Confocal scans of entire zebrafish retinas, the outer retina including ZPR1 labeled cone photoreceptors to confirm successful targeting of Cx35b-V5-TurboID to photoreceptor gap junctions. Neutravidin Oregon Green labeling was used to validate efficient proximity biotinylation of Cx35b and surrounding molecules in biotin injected Cx35b-V5-TurboID fish. Reagent or antibody used for labeling is shown in the gray box above each image. Scale 10µm. **(C)** Western blot of streptavidin pull-downs probed with V5 antibodies and streptavidin-HRP. The arrows indicate the position of the Cx35-V5-TurboID construct, which is detected with streptavidin-HRP and a V5 antibody. **(D)** String diagram illustrating the protein-protein network surrounding Cx35b. Proteins that were two times or more abundant in comparison to the control condition were included in the string diagram. **(E)** Colocalization of Cx35b and Cx34.7-GFP in the outer plexiform layer of the zebrafish retina suggests that a subset of photoreceptor gap junctions contain both connexins. Scale: 5 µm; inset scale: 0.5 µm.

Figure 2: A GFP-directed TurboID approach uncovers the Cx36 interactome in AII amacrine cells. (A) Cartoon illustrating different *in vivo* BioID approaches used to target Cx36. In strategy 1 we expressed an AAV Cx36-V5-TurboID construct with an internal insertion site under the control of the human synapsin promoter. This strategy greatly resulted in overexpression artefacts causing aberrant protein localization. In strategy 2 we adapted the GFP-directed TurboID strategy

developed by Xiong et al., (36) to shuttle a V5-dGBP-TurboID construct to Cx36-EGFP containing

gap junctions in retinas of the Cx36-EGFP transgenic strain. **(B)** Vertical sections of AAV infected

retinas confirming cell type-specificity of our AAV vectors. The HKamac promoter developed by

Khabou et al. (40) is mainly active in AII amacrine cells and does not show any expression in

bipolar cells (SCGN, magenta). V5-dGBP-TurboID colocalizes with Cx36-EGFP in AAV

transduced retinas, confirming the successful delivery of TurboID to the gap junction. Scale: 5µm.

(C) Western blot confirming successful biotinylation and capture of Cx36-EGFP, biotinylated

proteins and V5-TurboID-dGBP.   **(D)** String diagram illustrating the protein-protein interaction

network associated with Cx36 in AII amacrine cells. Proteins that were three times or more

abundant compared to the control condition were included in the string diagram.

Figure 3: Localization of scaffold and endocytosis proteins identified by BioID at Cx36 gap

junctions in AAV transduced AII amacrine cells expressing GFP. (A) Colocalization of Cx36

and scaffold proteins including known interactors such as ZO-1, ZO-2 and cingulin. Besides these

known interactors, we identified Sipa1l3, a PDZ domain containing protein implicated in cell

adhesion and cytoskeletal organization. Sipa1l3 shows abundant colocalization with Cx36. **(B)**

Colocalization of Cx36 and components of the endocytosis machinery. Among all proteins we

have tested, we found frequent colocalization for EpsS15l1, an endocytic adapter protein, and

Cx36. Scale: 10 µm. Magnified inset: 1 µm. **(C)** Bar graphs showing the degree of colocalization

as the percentage of Cx36 puncta that overlap with Cgn and Sipa1l3. Colocalization was

quantified for the original image and a horizontally flipped image to exclude random overlap. Bar

graphs with SEM, p <0.05 *** for Cx36 and Cgn and p <0.0001 **** for Cx36 and Sipa1l3.

Significance for Cx36 and Cgn was determined using a Wilcoxon matched-pairs signed rank test.

Significance for Cx36 and Sipa1l3 was determined using a two-tailed paired t-test. **(D)** Bar graphs

showing the degree of colocalization as the percentage of Cx36 puncta that overlap with

endocytosis proteins the Ep15l1 and Hip1r. Bar graphs with SEM, p <0.0001 **** for Cx36 and

Eps15l1 and p <0.05 *** for Cx36 and Hip1r. Significance was determined using a two-tailed

paired t-test.

Figure 4: Localization of trafficking, cytoskeleton-associated, adhesion and synaptic

proteins identified by BioID at Cx36 gap junctions in AAV transduced AII amacrine cells

expressing GFP. (A) Proteins implicated in membrane trafficking Sj2bp and Syt4 colocalize with

Cx36 in AII amacrine cells. **(B)** Several cytoskeleton associated proteins and regulators such as

Map6, Dock7 or Gprin1 colocalize with Cx36 in AII cell dendrites. **(C)** The adhesion molecule Bai1

was often colocalized with Cx36 in AII cell dendrites. **(D)** Often components of chemical synapses

such as Shank2 and Glur2-3 were found in the periphery of gap junction plaques in AII amacrine

cells. Scale: 10 µm. Magnified inset: 1 µm.

Figure 5: Sipa1l3, Sj2bp and Eps15l1 interact with Cx36. (A) Coexpression of Sipa1l3 with

ZO-1, ZO-2 and Cx36. In transfected HEK293T cells Sipa1l3 colocalizes with all three proteins.

Colocalization of Sj2bp with Cx36 but not with a Cx36 mutant that lacks the PDZ binding motif.

(B) Domain organization of constructs used for IP experiments. Co-precipitation of Cx36 and

Sjbp2 and Sipa1l3 from lysates of co-transfected HEK293T cells. The PDZ binding deficient

Cx36/S318Ter fails to bind Sjbp2 and Sipa1l3. IP experiments with ZO-1 and ZO-2 serve as a

positive control. The proteins that were captured in each IP experiment are indicated with arrows

in magenta. Cx36 dimers in IP samples are indicated with red arrows and monomers are indicated

with blue arrows. **(C)** In transfected HEK293T cells, Cx36 is associated with endogenous proteins

that were also detected in AII amacrine cells. Line scans next to each panel illustrate the close

association of these proteins (56) with Cx36 (magenta). These proteins were also captured via

GFP directed proximity biotinylation. Scale: 10 µm. Magnified inset: 1µm.

Figure 6: Localization of BioID "hits" at AII amacrine cell/ON cone bipolar cell junctions.

(A) Cartoon illustrating the neurons involved in the primary rod pathway and the subcellular

localization of AII/Cone bipolar cell junctions (dashed rectangle). **(B)** 3D reconstruction of AII

amacrine cells from the RC2 connectomics dataset illustrating the localization of AII-AII gap

junctions (cyan plaques in second image) and AII/ON Cone bipolar cell gap junctions (yellow

plaques in second image). **(C)** Transmission electron micrographs of AII/ON Cone bipolar cell gap

junctions. Pseudo colored: Gap junction (Cyan), AII cell (Magenta) and ON cone bipolar cell

(Yellow). White box indicates area shown to right at 40,000X magnification (0.27 nm/px resolution)

with 20 degree tilt. The density profile for the ROI indicated by the yellow box confirms the

pentalaminar (dark-light-dark-light-dark) structure of a gap junction. Also note the asymmetric

cytoplasmic densities on the AII versus ON Cone bipolar cell sides. Scale: 0.5 µm. **(D)** Several of

the proteins we identified in AII amacrine cells colocalize with Cx36 at AII/cone bipolar cell

junctions. The right plot for each panel depicts an intensity scan of a horizontal region of interest

in the middle of each gap junction (arrow). Scale: 10 µm. Magnified inset: 1 µm. **(E)** High resolution

scan highlighting the subsynaptic distribution of SIPA1L3. Scale: 5 µm. Magnified inset: 1 µm.

Figure 7: AII amacrine cell/ON cone bipolar cell contacts remain in Cx36 KO retinas and contain electrical synapse scaffolds. AII amacrine cell/ON cone bipolar cell contacts were visualized with GFP and Scgn. In Cx36 KO retinas, ZO-1, ZO-2 and Sipa1l3 still localize to AII/CBC contacts. **(D)** Both Sipa1l3 and ZO-1 puncta density in the inner plexiform layer were reduced in Cx36KO, but puncta size was unchanged. Colocalization of Sipa1l3 and ZO-1 was reduced in Cx36KO. **(E-F)** Cartoon illustrating electrical synapse configurations of AII amacrine cells and identified components of the electrical synapse proteome of the AII amacrine cell. Scale: 10 μm. Magnified inset: 1μm.

Figure S1: Localization of additional proteins captured with BioID. (A) Ajm1 labels the base of AII cell somas but does not directly colocalize with Cx36. **(B)** Cap1 is associated with Cx36 in AII cell dendrites. **(C)** Nbea does not directly associate with Cx36 but is often found close Cx36 clusters. Scale: 10 μm.

Figure S2: Localization of Ncam1 and Rtn4 clusters in proximity to Cx36 in AII amacrine cells. (A-B) Ncam1 and Rrn4 labeling does overlap with Cx36 but is found adjacent to gap junctions. Scale**:** 10 μm.

Figure S3: Summary table showing a comparison of the most abundant proteins in the electrical synapse proteomes of mice and zebrafish retinas. ZO proteins and endocytosis proteins occur in the electrical synapse proteomes of both species.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary figure 1

965   **Supplementary figure 2**

Supplementary figure 3

Protein	Danio rerio	Mus Musculus
Tjp1a	✓	✓
Tjp1b	✓	✓
Tjp2a	✓	✓
Sec22b	✓	✓
Eps15l1	✓	✓
Snap91	✓	✓
Synergin	✓	✓
Picalmb	✓	☐
Hip1r	☐	✓
MUPP1	✓	☐
Afadin	✓	☐
Cingulin	☐	✓
Sipa1l3	☐	✓
Gprin1	☐	✓
Myo6	✓	✓
PSD95	✓	✓
Sj2bp	☐	✓
Usp53b	✓	☐

References

1. M. V. Bennett, Electrical synapses, a personal perspective (or history). *Brain research reviews* **32**, 16-28 (2000).
2. C. Sotelo, The history of the synapse. *The Anatomical Record* **303**, 1252-1279 (2020).
3. E. A. Martin, A. M. Lasseigne, A. C. Miller, Understanding the molecular and cell biological mechanisms of electrical synapse formation. *Frontiers in neuroanatomy* **14**, 12 (2020).
4. T. C. Süudhof, Neurotransmitter release. *Pharmacology of Neurotransmitter Release*, 1-21 (2008).
5. M. J. Vaughn, J. S. Haas, On the diverse functions of electrical synapses. *Frontiers in Cellular Neuroscience* **16**, 910015 (2022).
6. S. G. Hormuzdi, M. A. Filippov, G. Mitropoulou, H. Monyer, R. Bruzzone, Electrical synapses: a dynamic signaling system that shapes the activity of neuronal networks. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **1662**, 113-137 (2004).
7. G. Söhl, K. Willecke, Gap junctions and the connexin protein family. *Cardiovascular research* **62**, 228-232 (2004).
8. D. F. Condorelli, N. Belluardo, A. Trovato-Salinaro, G. Mudò, Expression of Cx36 in mammalian neurons. *Brain Research Reviews* **32**, 72-85 (2000).
9. S. A. Bloomfield, B. Völgyi, The diverse functional roles and regulation of neuronal gap junctions in the retina. *Nature Reviews Neuroscience* **10**, 495-506 (2009).
10. V. Serre-Beinier *et al.*, Cx36 makes channels coupling human pancreatic β-cells, and correlates with insulin expression. *Human molecular genetics* **18**, 428-439 (2009).
11. M. Srinivas *et al.*, Functional properties of channels formed by the neuronal gap junction protein connexin36. *Journal of Neuroscience* **19**, 9848-9855 (1999).
12. J. O'brien, The ever-changing electrical synapse. *Current opinion in neurobiology* **29**, 64-72 (2014).
13. W. W. Kothmann *et al.*, Nonsynaptic NMDA receptors mediate activity-dependent plasticity of gap junctional coupling in the AII amacrine cell network. *Journal of Neuroscience* **32**, 6747-6759 (2012).
14. W. W. Kothmann, S. C. Massey, J. O'Brien, Dopamine-stimulated dephosphorylation of connexin 36 mediates AII amacrine cell uncoupling. *Journal of Neuroscience* **29**, 14903-14911 (2009).
15. A. Feigenspan, B. Teubner, K. Willecke, R. Weiler, Expression of neuronal connexin36 in AII amacrine cells of the mammalian retina. *Journal of Neuroscience* **21**, 230-239 (2001).
16. R. E. Marc, J. R. Anderson, B. W. Jones, C. L. Sigulinsky, J. S. Lauritzen, The AII amacrine cell connectome: a dense network hub. *Frontiers in neural circuits* **8**, 104 (2014).
17. C. Del Corsso, R. Iglesias, G. Zoidl, R. Dermietzel, D. C. Spray, Calmodulin dependent protein kinase increases conductance at gap junctions formed by the neuronal gap junction protein connexin36. *Brain research* **1487**, 69-77 (2012).
18. C. Alev *et al.*, The neuronal connexin36 interacts with and is phosphorylated by CaMKII in a way similar to CaMKII interaction with glutamate receptors. *Proceedings of the National Academy of Sciences* **105**, 20964-20969 (2008).
19. J. Lisman, H. Schulman, H. Cline, The molecular basis of CaMKII function in synaptic and behavioural memory. *Nature Reviews Neuroscience* **3**, 175-190 (2002).

20.   A. A. Coley, W.-J. Gao, PSD95: A synaptic protein implicated in schizophrenia or autism?
Progress in Neuro-Psychopharmacology and Biological Psychiatry **82**, 187-194 (2018).

21.   S. P. Cárdenas-García, S. Ijaz, A. E. Pereda, The components of an electrical synapse as
revealed by expansion microscopy of a single synaptic contact. *Elife* **13**, e91931 (2024).

22.   A. C. Miller, A. E. Pereda, The electrical synapse: molecular complexities at the gap and
beyond. *Developmental neurobiology* **77**, 562-574 (2017).

23.   E. Strettoi, E. Raviola, R. F. Dacheux, Synaptic connections of the narrow-field, bistratified
rod amacrine cell (AII) in the rabbit retina. *Journal of Comparative Neurology* **325**, 152-168
(1992).

24.   K. J. Roux, D. I. Kim, M. Raida, B. Burke, A promiscuous biotin ligase fusion protein identifies
proximal and interacting proteins in mammalian cells. *Journal of cell biology* **196**, 801-810
(2012).

25.   T. C. Branon *et al.*, Efficient proximity labeling in living cells and organisms with TurboID.
Nature biotechnology **36**, 880-887 (2018).

26.   T. Takano *et al.*, Chemico-genetic discovery of astrocytic control of inhibition in vivo. *Nature*
588, 296-302 (2020).

27.   M. Artan *et al.*, Interactome analysis of Caenorhabditis elegans synapses by TurboID-based
proximity labeling. *Journal of Biological Chemistry* **297** (2021).

28.   A. Uezu *et al.*, Identification of an elaborate complex mediating postsynaptic inhibition.
Science **353**, 1123-1129 (2016).

29.   H. Y. Wang, Y.-P. Lin, C. K. Mitchell, S. Ram, J. O'Brien, Two-color fluorescent analysis of
connexin 36 turnover: relationship to functional plasticity. *Journal of cell science* **128**, 3888-
3897 (2015).

30.   H. Li, A. Z. Chuang, J. O'Brien, Photoreceptor coupling is controlled by connexin 35
phosphorylation in zebrafish retina. *Journal of Neuroscience* **29**, 15178-15186 (2009).

31.   X. Li, B. Lynn, J. Nagy, The effector and scaffolding proteins AF6 and MUPP1 interact with
connexin36 and localize at gap junctions that form electrical synapses in rodent brain.
European Journal of Neuroscience **35**, 166-181 (2012).

32.   S. Tetenborg *et al.*, Phosphorylation of Connexin36 near the C-terminus switches binding
affinities for PDZ-domain and 14–3–3 proteins in vitro. *Scientific reports* **10**, 18378 (2020).

33.   A. M. Lasseigne *et al.*, Electrical synaptic transmission requires a postsynaptic scaffolding
protein. *Elife* **10**, e66898 (2021).

34.   C. Ciolofan *et al.*, Association of connexin36 and zonula occludens-1 with zonula
occludens-2 and the transcription factor zonula occludens-1-associated nucleic acid-
binding protein at neuronal gap junctions in rodent retina. *Neuroscience* **140**, 433-451
(2006).

35.   J. O'Brien, H. B. Nguyen, S. L. Mills, Cone photoreceptors in bass retina use two connexins
to mediate electrical coupling. *Journal of Neuroscience* **24**, 5632-5642 (2004).

36.   Z. Xiong *et al.*, In vivo proteomic mapping through GFP-directed proximity-dependent biotin
labelling in zebrafish. *Elife* **10**, e64631 (2021).

37.   I. Helbig *et al.*, In vivo evidence for the involvement of the carboxy terminal domain in
assembling connexin 36 at the electrical synapse. *Molecular and Cellular Neuroscience* **45**,
47-58 (2010).

38.   A. Meyer *et al.*, AII amacrine cells discriminate between heterocellular and homocellular
locations when assembling connexin36-containing gap junctions. *Journal of cell science*
127, 1190-1202 (2014).

39.   J. M. Christie *et al.*, Connexin36 mediates spike synchrony in olfactory bulb glomeruli.
Neuron **46**, 761-772 (2005).

40.   H. Khabou *et al.*, Optogenetic targeting of AII amacrine cells restores retinal computations
performed by the inner retina. *Molecular Therapy-Methods & Clinical Development* **31**
(2023).
41.   A. Feigenspan *et al.*, Expression of connexin36 in cone pedicles and OFF-cone bipolar cells
of the mouse retina. *Journal of Neuroscience* **24**, 3325-3334 (2004).
42.   J. C. Tang *et al.*, Detection and manipulation of live antigen-expressing cells using
conditionally stable nanobodies. *Elife* **5**, e15312 (2016).
43.   C. E. Flores *et al.*, Trafficking of gap junction channels at a vertebrate electrical synapse in
vivo. *Proceedings of the National Academy of Sciences* **109**, E573-E582 (2012).
44.   B. Lynn, X. Li, J. Nagy, Under construction: building the macromolecular superstructure and
signaling components of an electrical synapse. *The Journal of membrane biology* **245**, 303-
317 (2012).
45.   C. E. Flores, X. Li, M. V. Bennett, J. I. Nagy, A. E. Pereda, Interaction between connexin35 and
zonula occludens-1 and its potential role in the regulation of electrical synapses.
Proceedings of the National Academy of Sciences **105**, 12545-12550 (2008).
46.   S. Tetenborg *et al.*, Trafficking of Connexin36 (Cx36) in the early secretory pathway. *bioRxiv*
(2024).
47.   E. Leithe, S. Sirnes, T. Fykerud, A. Kjenseth, E. Rivedal, Endocytosis and post-endocytic
sorting of connexins. *Biochimica et Biophysica Acta (BBA)-Biomembranes* **1818**, 1870-1879
(2012).
48.   E. A. Martin *et al.*, Neurobeachin controls the asymmetric subcellular distribution of
electrical synapse proteins. *Current Biology* **33**, 2063-2074. e2064 (2023).
49.   Y.-K. Tu, J. G. Duman, K. F. Tolias, The adhesion-GPCR BAI1 promotes excitatory
synaptogenesis by coordinating bidirectional trans-synaptic signaling. *Journal of*
Neuroscience **38**, 8388-8406 (2018).
50.   C. Hartmann *et al.*, The mitochondrial outer membrane protein SYNJ2BP interacts with the
cell adhesion molecule TMIGD1 and can recruit it to mitochondria. *BMC Molecular and Cell*
Biology **21**, 1-17 (2020).
51.   S. Tetenborg *et al.*, Regulation of Cx36 trafficking through the early secretory pathway by
COPII cargo receptors and Grasp55. *Cellular and Molecular Life Sciences* **81**, 1-17 (2024).
52.   Y. Tsukamoto, N. Omi, Classification of mouse retinal bipolar cells: type-specific
connectivity with special reference to rod-driven AII amacrine pathways. *Frontiers in*
neuroanatomy **11**, 92 (2017).
53.   J. Nagy, B. Lynn, Structural and intermolecular associations between connexin36 and
protein components of the adherens junction–neuronal gap junction complex.
Neuroscience **384**, 241-261 (2018).
54.   J. O'Brien, R. Bruzzone, T. W. White, M. R. Al-Ubaidi, H. Ripps, Cloning and expression of two
related connexins from the perch retina define a distinct subgroup of the connexin family.
Journal of Neuroscience **18**, 7625-7637 (1998).
55.   A. Dolnik *et al.*, Sipa1l3/SPAR3 is targeted to postsynaptic specializations and interacts with
the Fezzin ProSAPiP1/Lzts3. *Journal of Neurochemistry* **136**, 28-35 (2016).
56.   R. Greenlees *et al.*, Mutations in SIPA1L3 cause eye defects through disruption of cell
polarity and cytoskeleton organization. *Human molecular genetics* **24**, 5789-5804 (2015).
57.   J. G. Duman *et al.*, The adhesion-GPCR BAI1 regulates synaptogenesis by controlling the
recruitment of the Par3/Tiam1 polarity complex to synaptic sites. *Journal of Neuroscience*
33, 6964-6978 (2013).
58.   M. Westerfield, *The zebrafish book: A guide for the laboratory use of zebrafish Danio ("*
Brachydanio Rerio") (University of Oregon, 2007).

59.   A. Santhanam *et al.*, A zebrafish model of retinitis pigmentosa shows continuous
1163   degeneration and regeneration of rod photoreceptors. *Cells* **9**, 2242 (2020).
1164   60.   N. Jin *et al.*, Molecular and functional architecture of the mouse photoreceptor network.
1165   *Science advances* **6**, eaba7232 (2020).
1166   61.   S. Tetenborg *et al.*, Intralumenal docking of connexin 36 channels in the ER isolates
1167   mistrafficked protein. *Journal of Biological Chemistry* **299** (2023).
1168   62.   K. Matsuura *et al.*, SIPA1L1/SPAR1 interacts with the neurabin family of proteins and is
1169   involved in GPCR signaling. *Journal of Neuroscience* **42**, 2448-2473 (2022).
1170   63.   S. Tetenborg, E. Martinez-Soler, O. John, Characterizing ER Retention Defects of PDZ Binding
1171   Deficient Cx36 Mutants Using Confocal Microscopy. *Bio-protocol* **14**, e5034 (2024).
1172   64.   P. Shannon *et al.*, Cytoscape: a software environment for integrated models of biomolecular
1173   interaction networks. *Genome research* **13**, 2498-2504 (2003).
1174   65.   G. Bindea *et al.*, ClueGO: a Cytoscape plug-in to decipher functionally grouped gene
1175   ontology and pathway annotation networks. *Bioinformatics* **25**, 1091-1093 (2009).
1176   66.   D. Szklarczyk *et al.*, The STRING database in 2023: protein–protein association networks
1177   and functional enrichment analyses for any sequenced genome of interest. *Nucleic acids*
1178   *research* **51**, D638-D646 (2023).
1179   67.   S. Tetenborg *et al.*, Differential distribution of retinal Ca2+/calmodulin-dependent kinase II
1180   (CaMKII) isoforms indicates CaMKII-β and-δ as specific elements of electrical synapses
1181   made of connexin36 (Cx36). *Frontiers in molecular neuroscience* **10**, 425 (2017).
1182   68.   S. Tetenborg *et al.*, Localization of retinal Ca2+/calmodulin-dependent kinase II-β (CaMKII-
1183   β) at bipolar cell gap junctions and cross-reactivity of a monoclonal anti-CaMKII-β antibody
1184   with Connexin36. *Frontiers in Molecular Neuroscience* **12**, 206 (2019).
1185   69.   J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nature*
1186   *methods* **9**, 676-682 (2012).
1187   70.   C. L. Sigulinsky, R. L. Pfeiffer, B. W. Jones, Retinal Connectomics: A Review. *Annual Review*
1188   *of Vision Science* **10**, 263-291 (2024).
1189   71.   J. R. Anderson *et al.*, A computational framework for ultrastructural mapping of neural
1190   circuitry. *PLoS biology* **7**, e1000074 (2009).
1191   72.   J. R. Anderson *et al.*, Exploring the retinal connectome. *Molecular vision* **17**, 355 (2011).
1192   73.   R. Marc, R. Murry, S. Basinger, Pattern recognition of amino acid signatures in retinal
1193   neurons. *Journal of Neuroscience* **15**, 5106-5129 (1995).

1194