A contextual fear conditioning paradigm in head-fixed mice exploring virtual reality.

Seetha Krishnan[1*], Can Dong[1,2*], Heather Ratigan[1], Denisse Morales-Rodriguez[1,3], Chery Cherian[1], Mark Sheffield[1]

[1] Department of Neurobiology and Institute for Neuroscience, University of Chicago, Chicago, IL 60637, USA.

[2] Present address: Department of Neurobiology, Stanford University

[3] Present address: University of California, San Francisco

* equal contribution

Correspondence to: seethakrishnan@uchicago.edu, sheffield@uchicago.edu

ABSTRACT

Contextual fear conditioning is a classical laboratory task that tests associative memory formation and recall. Techniques such as multi-photon microscopy and holographic stimulation offer tremendous opportunities to understand the neural underpinnings of these memories. However, these techniques generally require animals to be head-fixed. There are few paradigms that test contextual fear conditioning in head-fixed mice, and none where the behavioral outcome following fear conditioning is freezing, the most common measure of fear in freely moving animals. To address this gap, we developed a contextual fear conditioning paradigm in head-fixed mice using virtual reality (VR) environments. We designed an apparatus to deliver tail shocks (unconditioned stimulus, US) while mice navigated a VR environment (conditioned stimulus, CS). The acquisition of contextual fear was tested when the mice were reintroduced to the shock-paired VR environment the following day. We tested three different versions of this paradigm and, in all of them, observed an increased conditioned fear response characterized by increased freezing behavior. This was especially prominent during the first trial in the shock-paired

VR environment, compared to a neutral environment where the mice received no shocks. Our results demonstrate that head-fixed mice can be fear conditioned in VR, discriminate between a feared and neutral VR context, and display freezing as a conditioned response, similar to freely behaving animals. Furthermore, using a two-photon microscope, we imaged from large populations of hippocampal CA1 neurons before, during, and following contextual fear conditioning. Our findings reconfirmed those from the literature on freely moving animals, showing that CA1 place cells undergo remapping and show narrower place fields following fear conditioning. Our approach offers new opportunities to study the neural mechanisms underlying the formation, recall, and extinction of contextual fear memories. As the head-fixed preparation is compatible with multi-photon microscopy and holographic stimulation, it enables long-term tracking and manipulation of cells throughout distinct memory stages and provides subcellular resolution for investigating axonal, dendritic, and synaptic dynamics in real-time.

INTRODUCTION

Large-scale multi-photon imaging and holographic stimulation offers the ability to accurately track and record from the same neuronal populations over extended periods at high spatiotemporal resolution, record from axons, dendrites, and dendritic spines, and stimulate neurons with single-cell precision (Bando et al., 2021; Chaigneau et al., 2016; Dombeck et al., 2010; Dong et al., 2021; Emiliani et al., 2015; Fan et al., 2023; Hattori & Komiyama, 2022; Heer & Sheffield, 2024; Hernandez et al., 2016; Krishnan et al., 2022; Russell et al., 2022; Sheffield & Dombeck, 2015). These techniques facilitate the study of neural circuit dynamics, synaptic and plasticity mechanisms and enable the real-time manipulation of neural activity, all of which are crucial for understanding the mechanisms underlying memory formation and recall in the brain. However, head-fixation of animals, which is generally required to utilize these techniques, imposes constraints on the range of testable behaviors compared to freely moving conditions (Guo et al., 2014; Juczewski et al., 2020). This limitation is particularly evident in the Pavlovian contextual fear conditioning (CFC) task, a laboratory method for testing associative learning, fear memory formation, and recall (M. E. Bouton, 1993; Michael S. Fanselow, 1980, 1990; J. J. Kim & Fanselow, 1992; LeDoux, 2000; Maren, 2001; Maren et al., 2013).

In the traditional CFC paradigm for freely moving animals, an animal is placed in an environment (conditioned stimulus, CS) that is paired with an aversive stimulus, such as a mild foot shock (unconditioned stimulus, US). Animals are then removed from the environment. When reintroduced to the environment, animals exhibit a species-specific conditioned response, such as fearful freezing behavior in rodents, if they successfully associate the shock-paired environment with the aversive stimulus. The CFC paradigm

is one of the most basic conditioning procedures for freely moving animals— it's simple

and robust (Michael S. Fanselow, 1980, 1990). Despite its longstanding use in studies

involving freely behaving animals to explore questions on learning and memory

(Anagnostaras et al., 2001; M. E. Bouton & King, 1983; Mark E. Bouton & Bolles, 1980;

Debiec et al., 2002; M. S. Fanselow & Bolles, 1979; Fendt & Fanselow, 1999; Gewirtz et

al., 2000; Holland & Bouton, 1999; Jeansok J. Kim & Jung, 2006; Maren et al., 2013;

Stephen Maren & Holt, 2000; Pezze & Feldon, 2004), it has been noticeably absent in

studies involving head-fixed animals, with some exceptions.

Two prior studies (Lovett-Barron et al., 2014; Rajasethupathy et al., 2015) reported a

version of CFC in head-fixed mice with aversive air puffs as the US. These studies

reported the conditioned response animals displayed as lick suppression, not freezing.

Although lick suppression after fear conditioning has also been observed in freely moving

animals (Leaf & Leaf, 1966; Leaf & Muller, 1965), it has been shown to result from

increased freezing (Mark E. Bouton & Bolles, 1980). Therefore, relying solely on lick

suppression in the head-fixed preparation makes it more challenging to compare with the

extensive previous work in freely moving animals, where freezing is measured as the

predominant conditioned fear response. Furthermore, the aforementioned head-fixed

CFC studies also had the limitation of simultaneously having both a water reward and an

aversive stimulus within the context, potentially confounding the results and complicating

the interpretation. Hence, we aimed to develop a paradigm where freezing is the

conditioned response and without the presence of a reward within the conditioned

context, making it directly comparable to freely moving animals.

Here, we describe three different versions of a contextual fear-conditioning paradigm for

head-fixed mice that resulted in freezing as the conditioned response. The context or CS

was a virtual reality (VR)-based environment, which head-fixed mice explored by running

on a cylindrical treadmill. In a previous study (Ratigan et al., 2023), we demonstrated the

feasibility of this approach by combining it with imaging of thalamic axons innervating the

hippocampus using a two-photon microscope. The present manuscript provides a

comprehensive description and fundamentals of the behavioral framework, to enable the

research community to reproduce and adapt our behavior paradigm. Our main aim is to

show that VR-based CFC can elicit freezing responses similar to those seen in freely

moving animals, while also examining how key parameters affect this response. Though

we explored some important variables, this paper does not attempt to comprehensively

investigate all parameters necessary for successful VR-based CFC. Additionally, we

performed two-photon imaging of a large population of hippocampal CA1 cells during fear

conditioning and tracked the same cells during memory recall across days. We analyzed

the impact of our paradigm on place cells in the hippocampus and compared these

findings to results from CFC in freely moving animals.

RESULTS

A paradigm for contextual fear conditioning in head-fixed mice exploring virtual

reality environments

To establish the contextual fear conditioning protocol (**Figure 1A-C**), head-fixed mice

were water-restricted and trained to run on a treadmill and navigate VR environments for

water rewards, as we have demonstrated previously (Dong et al., 2021; Krishnan et al.,

2022; Krishnan & Sheffield, 2023; Ratigan et al., 2023). Forward movement on the

treadmill translates to forward movement in VR, allowing mice to navigate the VR

environment. The environments used were 2m-long linear tracks rich in visual cues

(**Figure 1**). Reaching the track's end triggered a water reward and defined a completed

trial/lap. Mice were then virtually teleported back to the track's start for the next lap. Unlike

freely moving mice that tend to explore new environments, head-fixed mice placed on a

treadmill do not automatically start exploring VR environments. However, they can be

trained through reinforcement learning using water restriction and water rewards

(Dombeck et al., 2010; Dong et al., 2021; Krishnan et al., 2022; Ratigan et al., 2023;

Sheffield & Dombeck, 2015).

Training to navigate VR environments on a treadmill (Stage 1), using water rewards, was

achieved in approximately 10-14 days, after which well-trained mice could run several

laps per minute. We set a criterion of >3 laps per minute (i.e., a lap velocity of 0.1 - 0.2

m/s) for mice to move on to the next stage (Stage 2). Once the mice reached this stage

(Stage 2), the following day, we added a coat to their tails (**Figure 1B**). We developed a

lightweight, wearable conductive apparatus for the mouse's tail to ensure minimal

discomfort for the mouse. We named this apparatus the "tail-coat" (**Figure 1D**). This tail-

coat enabled us to administer mild electric shocks to the mouse's tail (see Methods).

When the tail-coat was added, some mice reduced their running speed or refused to

move. Some recovered the next day, but if their running speed did not reach at least three

laps per minute (~20% of mice failed to meet this criterion), they were not advanced to

the next stage. Mice that maintained this running speed advanced to Stage 3. Data from

mice that advanced with minimal changes to their lap running behavior after adding the

tail-coat are shown in **Supplementary Figure 1A**.

The following day (Stage 3), mice were placed in the same VR environment with the tail-

coat, but the water reward was removed (**Figure 1B**). Mice that continued to run at least

three laps per minute with the tail-coat and without water reward advanced to the

experiment stage (Stage 4). Around 40% of the mice dropped out at this stage. High levels

of lap running during training were crucial to ensure mice were undeterred by both the

tail-coat and the reward removal (Krishnan et al., 2022; Krishnan & Sheffield, 2023). Our

criteria for consistent running behavior with minimal pauses throughout these three

stages of training ensured that any baseline freezing behavior was minimal, similar to

freely moving animals, and allowed us to attribute freezing after CFC to fear conditioning.

In summary, our criteria for selecting mice for the experimental stage of contextual fear

conditioning required them to run consistently (>3 laps per minute) with 1) rewards, then

2) with a tail-coat, and then 3) without rewards plus the tail-coat. This threshold resulted

in about 40% of the mice proceeding to the experimental stage. Once mice qualified for

the experimental stage, they were no longer excluded from analysis due to their behavior.

We tested three versions of the head-fixed CFC paradigm. In Paradigm 1, the neutral

environment was the training VR environment, and a new VR environment was paired

with shocks. In Paradigms 2 and 3, both the neutral and shock-paired environments were

new and distinct from the training environment to better control for environmental

familiarity. In Paradigms 1 and 2, the tail-coat was removed on the days following CFC,

whereas in Paradigm 3, it remained on the mouse on all post-CFC days. With these

paradigms, we varied some parameters that may influence fear memory recall, contextual

discrimination, fear extinction, and their associated behaviors in head-fixed VR-based

CFC paradigms. We detail these variations in the following sections, starting with

Paradigm 1.

Figure 1. A contextual fear conditioning paradigm for head-fixed mice navigating virtual reality
environments. (A) Experimental setup created with BioRender.com. Mice were head-restrained with their
feet resting on a cylindrical treadmill. Five large monitors surrounded the mice that displayed virtual reality
(VR) environments. Movement on the treadmill advanced the VR display, allowing for context exploration.
(B) Mice were water-restricted and trained to run laps in the VR for water rewards. VR environments were
2m-long linear tracks. Mice were trained to achieve >3 laps per minute, which took ~10 to 14 days (Stage
1). Once well-trained, a "tail-coat" was added to their tails (Stage 2), followed by the removal of the water
reward the next day (Stage 3). **(C)** For Paradigm 1, mice underwent the fear conditioning protocol the
following day after the water reward was removed. On the first experiment day (Day 0), mice spent 10
minutes in the training VR (Familiar VR) and then another 10 minutes in a new VR (CFC VR). After the
initial exploration, mice received mild electric shocks on the tail (4-12 shocks, 0.5-1.2mA in amplitude, 1s

long). The next day (Day 1), mice were tested for memory recall by placing them in the Familiar VR and the CFC VR for 5 minutes each in a counterbalanced manner. **(D)** Schematic of the tail-coat used for delivering mild electric shocks to the mouse's tail. (Top) View from the top (Bottom) side view. Dimensions are provided for a typical 12-week-old male mouse weighing ~30g before water restriction (see Methods for more details).

When tested for contextual fear memory recall, mice displayed increased freezing behavior in the shocked-paired VR environment.

On Day 0 of the experimental stage in Paradigm 1 (CFC day), mice (n = 27, 25 males and 2 females) were exposed to the same VR environment we used for training, referred to as the Familiar VR from here onwards (**Figure 1C**), for 10 minutes. We then switched them to a new VR environment (CFC VR). Following baseline exploration of the CFC VR for 10 minutes, we repeatedly applied mild-electric shocks (4-12 shocks, 0.5-1mA in amplitude, 1s long) to the mouse's tail. We call the period when shocks were delivered "during CFC". Shocks were pseudo-randomly administered with respect to the animal's position in the CFC VR, with an inter-shock interval (ISI) of 1 minute. To ensure that the shocks were associated with the context and not a specific position in the environment, no visual cues were explicitly paired with the shock. We quantified the animal's response to the shocks and observed an immediate increase in running speed following each shock (**Supplementary Figure 1B**). This was consistent across all mice and shocks and was a reliable indicator that the apparatus was working correctly. Sixty seconds after the last shock was administered, the VR screens were switched off, the tail-coat was removed, and mice were returned to their home cages. The next day (Fear memory recall, Day 1), we tested for recall of the learned association between the VR environment and the fearful stimulus. Mice were exposed to both VRs (familiar VR and CFC VR) for five minutes each in a counterbalanced manner. Water reward remained absent on recall days; however, in this paradigm, the tail-coat was no longer present.

Before fear conditioning, mice exhibited similar behavior in both the Familiar and CFC

VRs. We evaluated the time taken to complete a lap and the freezing behavior (defined

as time points when the animal's instantaneous velocity dropped was 0 cm/s). Neither

metric revealed a significant difference between the two VRs before shock administration

(**Figure 2A**, **Supplementary Figure 1D-E**). However, after CFC, when the animals were

reintroduced to the VRs on recall Day 1, we observed a significant increase in freezing

and time taken to complete a lap in the CFC VR compared to the Familiar VR (**Figure**

2A-D). As soon as mice transitioned to the CFC VR (either from dark or after the Familiar

VR) on the recall day, we observed behaviors such as slowing down, freezing, moving

backward, and hesitation to move forward (**Supplementary Videos 1-3**). This response

was markedly different from their behavior in the Familiar VR. This difference in behavior

can be observed clearly in most mice on the very first lap (**Figure 2A-B**). Since we

removed the reward, we didn't find licking behavior to be indicative of recall behavior

(**Supplementary Figure 1C**). However, some mice displayed licking behavior in the

Familiar VR on Day 0 before fear conditioning, which was absent on Day 1. This is likely

because mice had previously received a water reward in the Familiar VR. We will address

this caveat in Paradigms 2 and 3.

To quantify the distinct conditioned responses observed (**Figure 2A-B, Supplementary**

Videos 1-3), we used two metrics: freezing behavior, defined as periods when the

animal's instantaneous velocity was 0 cm/s, and time taken to complete a lap, which

captured the avoidance-like behaviors (such as slowing down, moving backward, and

hesitation to move forward) as well as freezing. When we quantified the freezing behavior

in mice, we found it to be, on average, high throughout the five minutes that mice spent

in the CFC VR on recall Day 1 (**Figure 2C-D,** average freezing (%), mean $\pm$ 95%

Confidence Intervals (CI), Familiar VR: 17.52 $\pm$ 3.25, CFC VR: 23.54 $\pm$ 3.26, p=0.001,

Paired *t*-test, n = 27). We also observed a significant increase in freezing during the first

lap in the CFC VR compared to the familiar VR (**Figure 2B-C,** average first lap freezing

(%), Familiar VR: 15.94 $\pm$ 4.03, CFC VR: 25.11 $\pm$ 3.41, p<0.001, Paired *t*-test). Mice also

took longer to complete a lap in the CFC VR compared to the Familiar VR, both in the

first lap (average time taken to complete first lap (s), Familiar VR: 14.81 $\pm$ 3.74, CFC VR:

41.16 $\pm$ 9.33, p<0.001, Paired *t*-test) and on average (**Figure 2E-F, the** average time

taken to complete all laps (s), Familiar VR: 16.17 $\pm$ 3.05, CFC VR: 38.82 $\pm$ 11.06, p<0.001,

Paired *t*-test). The increase in freezing and time taken to complete a lap in the CFC VR

compared to the Familiar VR was significantly higher on Day 1 than Day 0, indicating that

it results from fear conditioning (**Figure 2C-F**, p<0.05, Paired *t*-test).

Based on studies in freely moving animals (Michael S. Fanselow, 1980, 1990;

Ramanathan et al., 2018; Russo & Parsons, 2021; Trott et al., 2022), we settled on an

intensity of 1 mA and six shocks as a reasonable protocol, which we tested on a sizeable

cohort of mice (n = 12). We found that freezing at this intensity and shock number was

higher on average in the CFC VR; however, mice showed variability in their responses

(**Supplementary Figure 2A-B**). The majority of mice (7/12) displayed appropriate

memory recall (**Supplementary Figure 2A-B)**, indicated by more freezing in the CFC VR

compared to the Familiar VR (delta > 0), while 3/12 mice froze more in the Familiar VR

(delta < 0), and 2/12 showed equal freezing levels in both VRs (delta $\approx$ 0). Such variability

has been regularly documented in freely moving animals and may not be unique to the

VR setup used here (Chu et al., 2024; Dos Santos Corrêa et al., 2019; Navarro-Sánchez

et al., 2024; Poulos et al., 2016; Russo & Parsons, 2021; Totty et al., 2021).

Next, in a small group of mice (n = 15), we investigated whether varying the number of

shocks or the shock amplitude would affect the conditioned fear response

(**Supplementary Figure 2A-B**). Though our parameter exploration was not exhaustive,

we assessed shock amplitude more extensively than the number of shocks.

(**Supplementary Figure 2C-E**). Overall, most shock amplitudes and numbers of shocks

increased freezing behavior in the CFC VR on recall day (**Supplementary Figure 2A-B**).

The observed variability was within the same range as that of the larger cohort of mice

we tested using the 1 mA, 6-shock protocol (**Supplementary Figure 2A-B**). Notably,

increasing the shock amplitude from 0.6 mA to 1 mA, while keeping the number of shocks

consistent at 6 shocks, significantly enhanced both fear discrimination and freezing

behavior (**Supplementary Figure 2C-E**). A more detailed analysis with a larger sample

size may be necessary to understand further how variations in shock parameters affect

the conditioned response, as has been done in studies with freely moving animals (Baldi

et al., 2004; Cordero et al., 1998; Daviu et al., 2024; Poulos et al., 2016; Quinn et al.,

2008).

Finally, we examined the evolution of this conditioned response across days

(**Supplementary Figure 2F-I**). We placed the animals in both the Familiar and the CFC

VR for five minutes over four days (Recall Days 1-4) in a counterbalanced manner and

tested their recall behavior. We quantified both the degree of fear discrimination across

days (**Supplementary Figure 2 F-G**) and the extinction of the fear response within the

CFC VR compared to baseline (**Supplementary Figure 2 H-I**).

Fear discrimination was most prominent on Recall Day 1. Mice displayed the most

substantial increase in freezing in the CFC VR compared to the Familiar VR on Recall

Day 1 (**Supplementary Figure 2F-G**). Freezing on the first lap in the CFC VR was also

most pronounced on Recall Day 1. Both average and first lap freezing in the CFC VR

decreased on Recall Day 2 compared to Recall Day 1, becoming similar to the Familiar

VR. While first lap freezing remained similar between the Familiar and the CFC VR on

subsequent Recall Days 3 and 4, we observed some variability in the average freezing

behavior. Average freezing increased significantly again on Recall Day 3 in the CFC VR

but was similar in both VRs on Recall Day 4.

To determine the extinction of the fear response, we compared freezing within each VR

to baseline freezing levels on Day 0 (**Supplementary Figure 2H-I**). We quantified this

using a Linear Mixed Effects (LME) Model, treating recall days as fixed effects and mouse

as a random effect, with baseline freezing on Day 0 as the reference (**Tables 1-2**). In the

CFC VR, both first-lap and average freezing were most prominent on Recall Day 1 and

returned to baseline levels by Recall Day 2. While we observed a small increase in

average freezing on Recall Day 3, it remained lower than Recall Day 1 levels and returned

to baseline by Recall Day 4. This pattern suggests that the pronounced first-lap freezing

observed on Recall Day 1 extinguishes quickly, while average freezing, though more

variable, also extinguishes by Recall Day 4.

In the Familiar VR, freezing behavior—both first lap and average—did not differ

significantly from baseline on most days. However, we observed an unexpected increase

in average freezing compared to baseline on Recall Day 4. This increase could reflect

either fear generalization or decrease in motivation - a response to the prolonged absence

of reward in a previously rewarded environment. To distinguish between these

possibilities, we examined freezing behavior in control mice that underwent the same

paradigm without shocks (no-shock controls, n = 7, black lines in **Supplementary Figure**

2H-I). Average freezing in the Familiar VR exceeded that of no-shock controls on both

Recall Days 1 and 4. Since no-shock controls explored the environments under no-reward

conditions, this suggests that the increase in freezing in the Familiar VR stems from fear

conditioning and may indicate generalization rather than motivational changes. This

aligns with findings from freely moving animals, where specific fear memories have been

shown to generalize over time with repeated exposures to both neutral and conditioned

contexts (Biedenkapp & Rudy, 2007; Wiltgen & Silva, 2007).

In summary, these results demonstrate that head-fixed mice can acquire fear associations

in VR environments. Most mice displayed freezing as a conditioned response specifically

to the shock-paired VR and showed robust fear discrimination—behavior that mirrors that

of freely moving animals after fear conditioning. By the fourth recall day, freezing had

largely extinguished. Notably, beyond freezing, other avoidance-like defensive behaviors

were prominent in the shock-paired VR, demonstrating the range of behavioral responses

our paradigm can capture.

Paradigm 1

Figure 2. Paradigm 1: Head-fixed mice show increased freezing behavior following contextual fear conditioning in the CFC VR. (A) A single example mouse's lap running behavior on experiment days, Day 0 and Day 1. Behavior is shown for approximately 3 minutes in all sessions except during CFC, which is shown for the 6 minutes that the session lasted. Frames where freezing was detected (instantaneous velocity 0cm/s) are marked with black dots. A zoomed-in portion on the right in CFC VR highlights these freezing epochs as periods of minimal movement on the treadmill (i.e. instantaneous velocity 0cm/s), which are less visible in the full-scale view. The traces on the right show that this mouse increased freezing, decreased velocity, and moved backward (shown in gray) in the CFC VR (red traces) but not in the Familiar VR (blue traces) on Recall Day. This mouse received six shocks at 1mA intensity at 60 s inter-stimulus interval (ISI). **(B)** First two minutes of recall behavior in more mice (n = 5) in Familiar VR versus CFC VR. **(C-D)** (Left Average freezing percentage on recall day in the very first lap **(C)** and all laps **(D)** during the 5 minutes that mice explored the Familiar (blue) and CFC (red) VR. Freezing (%) was calculated as the number of frames where freezing was detected in a lap divided by the total number of frames in each lap. (Right) Delta calculated as the difference in the amount of freezing in the CFC VR compared to the Familiar

VR before CFC (Day 0) and after CFC (Day 1). The dashed line represents 0. **(E-F)** Same as C-D but for the time taken to complete the first lap **(E)** and all laps **(F).** Mice displayed an increase in freezing and in time taken to complete a lap in the first lap and, on average, in the CFC VR. In C-F, circles and pluses represent individual mice (n = 27, 25 male and two female mice). In C-F, data was pooled from mice receiving different numbers of shocks (4, 6, 12) at varying intensities (0.5 mA, 0.6mA, 0.8 mA, 1 mA, and 1.2 mA), which is separately displayed in **Supplementary Figure 2**. Lines join data from the same mouse. P-values were calculated using a *paired t-test.*

Day 0: Contextual Fear Conditioning

Supplementary Figure 1. Running and licking behaviors in Paradigm 1. (A) Box plot of average time taken to complete a lap before and after the addition of a tail-coat recorded in a randomly chosen subset of mice (circles, n=8). **(B)** Instantaneous velocity increased immediately after shock onset. This increased running speed also served as a reliable measure to confirm that mice received the tail shock. The grey line indicates the average velocity calculated by shuffling the instantaneous velocity 100 times. **(C)** Licking behavior doesn't stop immediately in the familiar VR without water, as mice previously received water rewards in the familiar VR. However, it was low in the other sessions and across environments. There was no significant difference between the Familiar and CFC VR on Day 1. Licking behavior was collected only in a subset of mice (circle, n=5). **(D-E)** The amount of freezing **(D)** and time taken to complete a lap **(E)** were not significantly different between the Familiar and CFC VR before fear conditioning (n=25). While not significantly different, there was a trend towards longer time taken in the first lap of the CFC VR **(E)**, which we have observed before when mice enter novel environments from familiar ones (Dong et al., 2021). The significantly lesser freezing during this first lap **(D)** suggests that animals were exploring slowly rather than remaining immobile. The boxplots (in A, C-E) range from the first quartile (25th percentile) to the third quartile (75th percentile), and the box shows the interquartile range (IQR). The line across the box represents the median (50th percentile). The whiskers extend to 1.5*IQR on either side of the box, and anything above this range is defined as an outlier. P-values were calculated using a paired *t-test.*

Supplementary Figure 2. Change in freezing with shock amplitude, number of shocks, and across recall days in Paradigm 1. (A-B) The left panel shows freezing in the first lap **(A)** and across all laps **(B).** The right panel illustrates delta freezing, calculated as the difference in freezing between the CFC VR and the Familiar VR. The colors represent the shock parameters used. Number of animals used is indicated in the figure legend. The dashed line represents the average freezing percentage in a control paradigm, where mice experienced the same conditions without shocks (n = 7). In most cases, we observed an increase in freezing in the CFC VR compared to the Familiar VR (delta>0). **(C-D)** Freezing in the first lap **(C)** and across

all laps **(D)** by shock amplitude. In all cases, mice received six shocks that were 1 minute apart. Greater fear discrimination could be seen at higher shock amplitudes. **(E)** Median freezing in the CFC VR by shock amplitude for first lab freezing (black) and average freezing across laps (gray). **(F-G)** A subset of animals underwent multiple days of recall tests (n = 25 Day 0, n = 27 Recall Day 1, n = 13 Recall Day 2, n = 8 Recall Day 3, n = 7 Recall Day 4). VR environments were presented in a counterbalanced manner for 5 minutes each day. The increase in freezing in the CFC VR was highest on the first day, both in the first lap **(F)** and across all laps **(G)**. P-values were calculated using a paired *t-test.* **(H-I)** Within-VR comparison of the same dataset. First lap freezing **(H)** and average freezing across all laps **(I)**, separated by Familiar VR (left panels) and CFC VR (right panels), demonstrate fear extinction across days. The black line represents average freezing in control mice that received no shocks. P-values indicate significant differences between experimental and no-shock control groups (t-test). Fear extinction is quantified by the LME model in Tables 1 and 2, which shows the statistical comparison of freezing in recall days versus Day 0 (baseline) within each VR.

Table 1: LME summary (estimate ± standard error, p-value) for first lap freezing in Paradigm 1 with fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2	Recall Day 3	Recall Day 4
Familiar VR	-0.97 ± 1.95, p=0.62	-5.75 ± 2.51, p=0.02	-4.13 ± 3.03, p=0.18	1.08 ± 3.18, p=0.73
CFC VR	12.96 ± 2.02, p<0.001	3.42 ± 2.55, p=0.18	1.73 ± 3.07, p=0.57	1.26 ± 3.23, p=0.70

Table 2: LME summary (estimate ± standard error, p-value) for average freezing in Paradigm 1 with fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2	Recall Day 3	Recall Day 4
Familiar VR	1.00 ± 1.13, p=0.37	-0.43 ± 1.48, p=0.77	0.76 ± 1.80, p=0.67	4.87 ± 1.88, p=0.013
CFC VR	7.55 ± 1.58, p<0.001	2.89 ± 2.04, p=0.16	6.03 ± 2.47, p=0.018	3.78 ± 2.58, p=0.15

A second paradigm where head-fixed mice discriminate between two novel VRs.

As previously mentioned, the incomplete extinction of licking behavior in the familiar VR before fear conditioning or the gradual increases in freezing behavior in the familiar VR during recall tests may indicate that mice still associate the VR with water rewards, which could be a caveat. Furthermore, in traditional CFC paradigms with freely moving mice, fear behavior has been assessed mainly by comparing two novel contexts, one associated with shocks and the other acting as a control. To address this, we tested another paradigm (Paradigm 2), using a novel VR instead of the familiar VR to act as the control (**Figure 3A**). One novel VR was assigned as the Control VR, and a second novel

VR was assigned as the CFC VR, which would later become paired with the shocks. In

this paradigm, we also added an extra habituation day to increase the animal's pre-

exposure time to the novel VRs prior to conditioning (Day -1). Pre-exposure to the context

before fear conditioning can enhance contextual fear, as seen in freely moving animals

(Biedenkapp & Rudy, 2007; Michael S. Fanselow, 1980, 1990; Maren et al., 2013; Rudy

et al., 2002; Wiltgen & Silva, 2007).

The training was done similarly to the previous paradigm. After mice showed consistent

running behavior in the Familiar VR with the tail-coat and without reward, they were

advanced to the experimental stage. On Day -1, mice were exposed to the two novel VRs

for ten minutes each. On Day 0, they were re-exposed to the same two VRs for ten

minutes before receiving six one-second-long mild electric shocks (at 1 mA and 1 minute

ISI) in one of them - the CFC VR. On Day 1, mice were exposed to the two VRs for five

minutes to evaluate memory recall (**Figure 3A-B**).

Compared to Paradigm 1, in the novel VRs, mice showed no differences in licking

behavior across all experimental sessions (**Supplementary Figure 3A**). There was also

no significant difference in freezing behavior and time taken to complete a lap between

the two VRs on Day -1 and Day 0 before fear conditioning (**Supplementary Figure 3B-**

C).

Head-fixed mice show heightened fear discrimination during the first lap when

discriminating between two novel VRs.

We compared the freezing behavior between Paradigm 1 and Paradigm 2 on Recall Day

1 to the same intensity and number of shocks (1 mA, 1 s long, 60 s ISI, six shocks, n=12

mice each paradigm). While first-lap freezing in the CFC VR was markedly higher in

Paradigm 2, significant differences between the Control VR and CFC VR were observed only for first-lap freezing, not for average freezing or lap completion time across all laps (**Figure 3B-D, 4A-E**). The effect sizes for the difference in first-lap freezing between the two VRs were nearly identical between the paradigms (Cohen's d: First-lap freezing: Paradigm 1: 1.038, Paradigm 2: 1.040), but the effect size for average freezing was notably smaller in Paradigm 2 compared to Paradigm 1 (Cohen's d: Average Freezing: Paradigm 1: 0.797, Paradigm 2: 0.093).

When we quantified freezing by lap on Recall Day 1 in Paradigm 2, we found that after the first lap, mice showed similar freezing in both the Control VR and the CFC VR across all subsequent laps (**Supplementary Figure 4**). In contrast, in Paradigm 1, mice consistently froze more in the CFC VR compared to the Familiar VR throughout Recall Day 1 (**Supplementary Figure 4**).

Indeed, when we quantified freezing behavior in the Control VR and CFC VR in Paradigm 2, we found that the most significant increase in freezing and other avoidance-like defensive behaviors in the CFC VR was observed in the first lap (freezing in first lap %: Control VR: 11.04 ± 1.8 CFC VR: 16.05 ± 3.94, p = 0.016, time taken to complete first lap (s): Control VR: 14.27 ± 2.85, CFC VR: 25.68 ± 5.55, p = 0.001, Paired *t*-test, **Figure 3B-E, Figure 4A-E**). Neither average freezing across all laps (**Figure 4C,** average freezing in all laps %: Control VR: 20.54 ± 4.13, CFC VR: 21.15 ± 4.21, p=0.738, Paired *t*-test) nor time taken to complete all laps (**Figure 4E,** the average time taken to complete all laps: Control VR: 21.73 ± 8.11, CFC VR: 24.27 ± 6.36, p=0.558, Paired *t*-test) showed significant differences between the two VRs.

When mice were returned to the two VRs the following day for a second day of recall

(Recall Day 2), they continued to exhibit modest levels of fear discrimination, as indicated

by a small but statistically significant difference in freezing in the first lap and on average

between the CFC VR and Control VR (**Supplementary Figure 5 A-B**). However, when

assessing fear extinction by comparing freezing in the CFC VR across recall days to

baseline levels (**Supplementary Figure 5 C-D, Tables 3-4**), we found no significant

differences beyond the very first lap on Recall Day 1. This suggests that while mice

maintained their ability to discriminate between environments by freezing more in the CFC

VR, their overall fear response had extinguished. Thus, in Paradigm 2, fear extinction

occurs rapidly within the first recall session after the initial lap.

Figure 3. Using a novel VR as the neutral environment instead of a familiar VR in a second paradigm results in modest increases in freezing. (A) The training paradigm for Paradigm 2 is similar to Paradigm 1. On experiment day, mice were introduced to two novel VRs, one of which would be associated with the shock (CFC VR) and the other wouldn't (Control VR). In this paradigm, there was an added habituation day, where mice were exposed to the two VRs for ten minutes. The next day, mice ran in the two VRs again before receiving mild electric shocks in the CFC VR. The recall test occurred the next day. In this cohort, all mice received six shocks at 1 mA intensity. **(B-D)** These panels show the lap running behavior of a single mouse on experiment days -1, 0, and 1. Behavior is displayed for about 3 minutes in all sessions except

**Figure 4. In Paradigm 2, the most significant freezing and reduced speed occurred in the first lap of
the CFC VR compared to the Control VR. (A)** First two minutes of recall behavior in five mice in Control
VR (green) versus CFC VR (red). **(B-C)** The left panels show the amount of freezing on the recall day
in (B) the very first lap and (C) all laps during the 5 minutes when the mice explored the Control (green)
and CFC (red) VR. Freezing (%) is calculated as the number of frames where freezing was detected in a
lap divided by the total number of frames in each lap. On the right, the delta is calculated as the difference
in the amount of freezing in the CFC VR compared to the Control VR before (Day 0) and after CFC (Day
1). The dashed line represents 0. **(D-E)** Same as B-C but for the time taken to complete the first lap **(D)** and
all laps **(E).** In B-E, circles and pluses represent individual mice (n = 12). Lines join data from the same
mouse. P-values were calculated using a *paired t-test.*

Supplementary Figure 3. Licking and running behavior before fear conditioning, in Paradigm 2. (A) Licking behavior remains low in all sessions, unlike Paradigm 1. It does not significantly differ between the Control vs CFC VR before and after CFC. **(B-C)** The amount of freezing and average running speed was not significantly different between the Control and CFC VR before fear conditioning, both on Day -1 **(B)** and Day 0 **(C)**. P-values were calculated using a *paired t-test.*

427
Supplementary Figure 4. Lap-wise freezing behavior in the three paradigms. (A-F) Average freezing
across mice **(A-C)** and the number of freezing epochs (**D-F,** see Methods) in each lap across the two VRs
on the first day of recall in **(A, D)** Paradigm 1, **(B, E)** Paradigm 2 and **(C, F)** Paradigm 3. Lines indicate
mean, and shading indicates 95% confidence intervals. P-values between **(A, D)** Familiar VR and CFC
VR and **(B-C, E-F)** Control VR and CFC VR were calculated using a *paired t-test.* * indicates p-
values<0.01.

434
Supplementary Figure 5. Mice largely extinguished their fear within the first day of recall in
Paradigm 2. (A-B) First lap freezing **(A)** and average freezing across all laps **(B)** across two days of
recall in Paradigm 2. P-values were calculated using a paired t-test. Circles indicate data from individual
mice (n = 12 for Recall Day 1 and n = 8 for Recall Day 2). P-values were calculated using a paired *t-test.*
(C-D) Within-VR comparison of the same dataset. First lap freezing **(C)** and average freezing across all
laps **(D),** separated by Familiar VR (left panels) and CFC VR (right panels), demonstrate fear extinction
across days. Fear extinction is quantified by the LME model in Tables 3 and 4, which shows the statistical
comparison of freezing in recall days versus Day 0 (baseline) within each VR.

Table 3: LME summary (estimate ± standard error, p-value) for first lap freezing in Paradigm 2
with fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2
Control VR	-5.22 ± 2.88, p=0.08	-6.99 ± 3.22, p=0.04
CFC VR	5.02 ± 1.98, p=0.02	4.01 ± 2.24, p=0.09

446 **Table 4: LME summary (estimate ± standard error, p-value) for average freezing in Paradigm 2**
447 **with fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).**

VR	Recall Day 1	Recall Day 2
Control VR	-2.71 ± 2.00, p=0.19	-1.15 ± 2.27, p=0.62
CFC VR	-0.23 ± 2.06, p=0.91	2.76 ± 2.32, p=0.25

<u>A shorter ISI and leaving the tail-coat on during fear memory recall improved fear</u>

<u>discrimination when using two novel VRs.</u>

Freezing behavior in the CFC VR in Paradigm 2 was weaker than in Paradigm 1, with

rapid fear extinction after the first recall lap. In Paradigm 3, two parameters were changed

relative to Paradigm 2 to test whether freezing behavior and fear discrimination could be

enhanced. First, the interval between consecutive shocks was reduced to see if a shorter

ISI could produce a more robust fear response (McNally & Westbrook, 2006; Williams,

1994). Second, the tail-coat was kept on during memory recall, based on evidence that

removing contextual cues can reduce conditioned freezing (González et al., 2003). We

hypothesized that the tail-coat might act as a contextual cue and that its presence during

recall might enhance freezing behavior. To further test these effects, we used a lower

shock intensity of 0.6 mA—which we previously showed as producing only modest fear

responses in Paradigm 1 (**Supplementary Figure 2 C-E**)—to determine whether a

shorter ISI and the retention of the tail-coat during recall could enhance conditioned fear

responses produced by the lower shock intensity.

In Paradigm 3, mice followed the same experimental protocol as Paradigm 2, with two

novel VRs: one assigned as the Control VR and the other as the CFC VR. A habituation

day in the two new VRs (Day -1) was followed by the CFC day (Day 0), where mice

received six shocks (0.6 mA, 20-26 s ISI) in the CFC VR. This was followed by recall days

(Days 1-3), where mice explored the two VRs presented in a counter-balanced manner.

Unlike the other two paradigms, in Paradigm 3, the tail-coat remained on during the recall

days (**Figure 5A**).

We found that mice (n = 20 male mice) showed better fear discrimination between the

Control and CFC VRs in Paradigm 3 compared to Paradigm 2 (**Figure 5B-D**). Mice not

only showed significant freezing in the first lap (freezing in first lap %: Control VR: 13.61

$\pm$ 2.24 CFC VR: 19.73 $\pm$ 5.17, p = 0.021, Paired *t*-test) but average freezing across all

laps was also higher in the CFC VR compared to the Control VR (**Figure 5C-D, Figure**

6A-D, average freezing in all laps %: Control VR: 15.28 $\pm$ 2.00, CFC VR: 21.92 $\pm$ 2.95, p

< 0.001, Paired *t*-test).

While first-lap freezing showed a weaker effect size in Paradigm 3 compared to the other

paradigms, there was a strong effect of average freezing across the session (Cohen's d:

First lap freezing: Paradigm 1: 1.038, Paradigm 2: 1.040, Paradigm 3: 0.719. Average

Freezing: Paradigm 1: 0.797, Paradigm 2: 0.093, Paradigm 3: 1.232). On average, we

found that Paradigm 1 still produced the highest absolute freezing behavior in the CFC

VR (average freezing in all laps %: CFC VR: Paradigm 1: 23.54 $\pm$ 3.26, Paradigm 2: 21.51

$\pm$ 4.21, Paradigm 3: 21.92 $\pm$ 2.95). The difference in freezing behavior between the CFC

VR and the neutral VR was comparable between Paradigms 1 and 3 (Δ average freezing

% (CFC VR – Familiar/Control VR): Paradigm 1: 6.74, Paradigm 2: 0.61, Paradigm 3:

6.63). Thus, unlike Paradigm 2, where the largest differences in freezing between the

CFC VR and the Control VR were restricted to the first lap, the shorter ISIs and the

addition of the tail-coat during the recall day in Paradigm 3 improved contextual fear

discrimination between the two novel VRs. When tested for memory recall in Paradigm

3, mice froze more in the CFC VR than in the Control VR, and this effect persisted beyond

the first lap (**Figure 5C-D, Supplementary Figure 4**). Mice also maintained their ability

to discriminate between the two VRs for the first two recall days, freezing significantly

more in the CFC VR than in the Control VR on both Recall Days 1 and 2 (**Supplementary**

Figure 6A-B). However, by Recall Day 3, this discrimination diminished, as freezing

levels no longer differed significantly between the two VRs.

Unlike the rapid fear extinction observed in Paradigm 2, mice in Paradigm 3 showed

delayed extinction (**Table 5-6**, **Supplementary Figure 6 C-D**). Freezing levels in the CFC

VR remained elevated above baseline during the first lap and across all laps on Recall

Days 1 and 2, with average freezing still elevated above baseline on Recall Day 3. While

freezing levels showed a consistent downward trend across days, these results suggest

that fear had not completely extinguished by Recall Day 3. As in Paradigm 1, the most

pronounced increases in first lap freezing in the CFC VR occurred in early recall days,

while elevated average freezing persisted longer. These findings suggest that the shorter

ISI and presence of the tail-coat during memory recall may enhance conditioned fear and

delay fear extinction in head-fixed mice discriminating between two novel VRs, though

the individual effects of these parameters require further investigation.

The tail-coat's presence during recall days likely contributed to some transfer of fear to

the Control VR, evidenced by elevated freezing in the Control VR compared to baseline.

Nevertheless, freezing levels in the Control VR remained consistently lower than in the

CFC VR across all days (**Supplementary Figure 6, Tables 5-6**).

In summary, our results show that mice exhibited greater average freezing behavior in

the CFC VR in Paradigms 1 and 3 compared to Paradigm 2. In Paradigm 2, the most

noticeable increase in freezing occurred during the very first lap in the CFC VR compared

to the Control VR. Fear extinction was rapid in Paradigm 2 and was back to baseline

levels within the first day of recall. We found that fear discrimination could be improved,

and fear extinction could be delayed with a shorter ISI when presenting tail shocks and

when the tail-coat was present during memory recall, as shown in Paradigm 3. Thus, we

provide three different variations for performing CFC in head-fixed mice that can be used

to understand the neural underpinnings of fear memory.

Figure 5. A third paradigm that uses a novel VR as the neutral environment but keeps the tail-coat on during memory recall led to increased freezing in the CFC Environment. (A) The training and experiment paradigm was similar to Paradigm 2 except that the shocks were administered closer together (20-25 s ISI), and the tail-coat was kept on during recall days. **(B)** First two minutes of recall behavior in Paradigm 3 in five mice in Control VR (green) versus CFC VR (red). **(C-D)** Comparison between freezing in the first lap **(C)** and all laps **(D)** in Paradigm 1 (blue and red) versus Paradigms 2 and 3 (green and red). n = 12 mice were used in Paradigms 1 and 2 (10 male and two female mice), and n = 20 male mice in Paradigm 3. P-values were calculated using a *paired t-test*. In Paradigm 3, by keeping the tail-coat on during the recall days, we observed an increase in freezing in the CFC VR in the first lap and across all laps compared to Paradigm 2.

Paradigm 3
Day 1: Fear Memory Recall

Figure 6. Mice exhibited increased freezing and overall reduced speed in the CFC VR in Paradigm 3 (A-B) The right panels show the amount of freezing on the recall day in **(A)** the very first lap and **(B)** all laps during the 5 minutes when the mice explored the Control (green) and CFC (red) VR. Freezing (%) is calculated as the number of frames where freezing was detected in a lap divided by the total number of frames in each lap. On the left, the delta is calculated as the difference in the amount of freezing in the CFC VR compared to the Control VR before (Day 0) and after CFC (Day 1). The dashed line represents 0. **(C-D)** Same as A-B but for time taken to complete the first lap **(C)** and all laps **(D).** Circles and pluses represent individual mice (n = 20). Lines join data from the same mouse. P-values were calculated using a *paired t-test.*

542
Supplementary Figure 6. Mice displayed better fear discrimination and a delay in extinction in
Paradigm 3 compared to Paradigms 1 and 2 (A-B) Freezing in the first lap **(A)** and average freezing
across all laps **(B)** in the Control VR vs CFC VR on three consecutive recall days in Paradigm 3 (n = 20).
The tail-coat was kept on all recall days. P-values were calculated using a paired *t-test*. **(C-D)** Within-VR
comparison of the same dataset. First lap freezing **(C)** and average freezing across all laps **(D),** separated
by Familiar VR (left panels) and CFC VR (right panels), demonstrate fear extinction across days. Fear
extinction is quantified by the LME model in Tables 5 and 6, which shows the statistical comparison of
freezing in recall days versus Day 0 (baseline) within each VR.

Table 5: LME summary (estimate ± standard error, p-value) for first lap freezing in Paradigm 3 with
fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2	Recall Day 3
Control VR	1.84± 1.92, p=0.34	3.01 ± 1.92, p=0.12	-2.09 ± 1.92, p=0.28
CFC VR	9.67 ± 2.61, p=0.0005	6.84 ± 2.61, p=0.01	3.33 ± 2.61, p=0.21

553
554 **Table 6: LME summary (estimate ± standard error, p-value) for average freezing in Paradigm 3 with**
555 **fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).**

VR	Recall Day 1	Recall Day 2	Recall Day 3
Control VR	2.65 ± 0.97, p=0.008	3.99 ± 0.97, p=0.0001	2.28 ± 0.97, p=0.02
CFC VR	10.30 ± 1.41, p<0.001	8.81 ± 1.41, p<0.001	4.28 ± 1.41, p=0.004

556
Place cells in the CA1 subregion of the hippocampus displayed remapping and

decreased field widths following contextual fear conditioning.

The hippocampus plays a critical role in the encoding, consolidation, and recall of

memories and is necessary for the expression of contextual fear memory (Anagnostaras

et al., 2001; Ji & Maren, 2007; Liu et al., 2012; Maren et al., 2013; Stephen Maren & Holt,

2000; Phillips & LeDoux, 1992; Ramirez et al., 2013; Wiltgen & Silva, 2007). Context is

thought to be represented in the hippocampus by place cells that fire at specific spatial

locations (place fields) in an environment (O'Keefe & Dostrovsky, 1971). These place

cells have been thought to help discriminate between feared and neutral contexts.

Previous studies on freely moving animals have shown that fear conditioning can cause

place cells to remap (Blair et al., 2023; E. J. Kim et al., 2015; Kinsky et al., 2023; Mamad

et al., 2019; Moita et al., 2004; Ormond et al., 2023; Schuette et al., 2020; Wang et al.,

2012, 2015; Wu et al., 2017). Remapping is defined by place cells shifting their firing

fields, indicative of place cells adapting to incorporate new information into memory

(Colgin et al., 2008; Leutgeb et al., 2005; Muller & Kubie, 1987). Additionally, a recent

study (Schuette et al., 2020) that performed calcium imaging of hippocampal cells in freely

moving mice found that fear conditioning narrows the width of place fields in the feared

environment. This suggests that place cells encode the same environment on a finer scale

following fear conditioning, potentially enhancing context discrimination and threat

avoidance (Schuette et al., 2020).

We replicated these findings in our head-fixed version of CFC in VR. To investigate the

effect of fear conditioning on place cell activity in our task, we expressed a calcium

indicator, GCaMP6f (Chen et al., 2013), in hippocampal pyramidal cells and, using two-

photon microscopy, imaged from the same hippocampal neurons across days during fear

conditioning and fear memory recall in Paradigm 1 (**Figure 7A-B,** n=8 mice). We found

that a subset of place cells formed before CFC remained stable, while others remapped

or changed their preferred firing locations after CFC (see examples in **Figure 7C, 7D-G**).

This is similar to previously reported observations (Blair et al., 2023; Moita et al., 2004;

Wang et al., 2012, 2015).

We found remapping to occur in both the Familiar and the CFC VR to similar extents, as

indicated by the lower correlation across sessions/days compared to the within-session

correlation (**Figure 7F-G**). This contradicts previous findings by Moita et al. (2004), who

reported more stability in the control environment compared to the CFC environment after

fear conditioning. This discrepancy may be explained by the different reward protocols

used. Moita et al.'s rats foraged continuously for food pellets during the experimental

sessions, while our mice received no rewards. We removed water rewards from the

previously rewarded Familiar VR, and as we have shown in earlier work, removing

rewards from a previously rewarded environment causes the place map to become more

unstable and likely to remap across days (Krishnan et al., 2022; Krishnan & Sheffield,

2023). Additionally, Moita et al. measured place cell stability within a single day, with just

one hour between conditioning and retrieval sessions, while our recordings were

separated by a day, including intervening sleep periods. This longer interval may have

contributed to more pronounced remapping compared to the shorter interval used by

Moita et al (Cai et al., 2009; Maboudi et al., 2018).

Next, we looked at the place cells in each session. Interestingly, we found an increase in

the number of place cells in the CFC VR during memory recall, a finding not previously

reported (**Figure 8A-B**). We also found that place cells had narrower fields in the CFC

VR during memory recall (**Figure 8C**), as previously shown (Schuette et al., 2020). We

did not find a significant difference in other place cell parameters we examined, such as

lap-by-lap reliability (**Figure 8D**) and out-of-field firing (**Figure 8E**). Thus, our observations

of remapping and narrower place field widths replicate findings from freely moving

animals, validating our paradigm's use to explore further questions using the head-fixed

preparation to study the neural underpinnings of contextual fear memories. We add to

this field by showing an increase in the proportion of place cells following CFC, a possible

neural substrate of fear memory recall.

Figure 7. Fear conditioning results in place cell remapping in both Familiar and CFC VR. (A) A schematic of the procedure for two-photon imaging of large populations of the same CA1 neurons over multiple days, created with BioRender.com **(B)** An example field of view across the two experiment days. The same field of view was aligned and imaged across days. **(C)** Examples of a few place cells in the CFC VR across sessions on Day 0 (before CFC, during CFC) and Day 1 (Memory Recall). White lines separate laps in each session. Some place cells maintain stable fields across days, while others remap. **(D)** Place

fields defined on Day 0 in Familiar VR before CFC are plotted across Day 1 during fear memory recall (n =
8 mice). **(E)** Place fields defined on Day 0 in CFC VR before CFC are plotted across the fear conditioning
session and on Day 1 during fear memory recall. **(F)** On the left is a scatter plot of the center of mass of
place fields defined in Familiar VR before CFC on Day 0 (x-axis) compared to their center of mass on Day
1 during fear memory recall (y-axis). On the right is a boxplot of correlation coefficients between mean place
fields defined in Familiar VR before CFC on Day 0 and memory recall on Day 1 (Day 0: Day 1). The within-
session correlation coefficients serve as control and were calculated between mean place fields in the first
half and second half of the Familiar VR before the CFC session. **(G)** A scatter plot of the center of mass of
place fields defined in CFC VR before CFC on Day 0 (x-axis) compared to their center of mass on (right)
Day 0 during CFC and (middle) Day 1 during memory recall. On the left is a distribution of correlation
coefficients between mean place fields defined in Familiar VR before CFC on Day 0 and during CFC (Day
0: Day 0) and memory recall on Day 1 (Day 0: Day 1). The within-session correlation was calculated
between mean place fields in the first half and second half of the CFC VR before the fear conditioning
session. Asterisk (*) indicates significant *P* values (KS test, *P* < 0.01). Our findings show that place fields
present in the before CFC sessions in both Familiar and CFC VR showed significant remapping following
fear conditioning.

635
Figure 8. The widths of place fields in CA1 narrow during memory recall. (A) Place fields were defined
and sorted by track length in each session, pooled from all mice (*n* = 8 mice). Each cell's activity was
normalized to its peak, and cells were sorted by their center of mass along the track. **(B)** The percentage
of place cells is calculated as the number of place cells divided by the total number of recorded cells. More
place cells were identified by our algorithm in the CFC VR on Day 1. **(C-E)** Parameters of place fields: **(C)**
Width, **(D)** Place Field (PF) Reliability, and **(E)** out/in field firing ratio in both VRs on Day 0 and Day 1. The
width of the place fields in the CFC VR significantly decreased on Day 1. No other parameters significantly
varied across days. P-values were calculated using a *paired t-test*.

DISCUSSION
We present three paradigms for performing CFC in head-fixed mice, using a conductive

"tail-coat" to deliver aversive shocks as mice navigate VR environments on a treadmill.

Our key findings are: 1) Head-fixed mice exhibited freezing behavior in response to an

aversive stimulus in VR, consistent with conditioned responses seen in freely moving

animals. Freezing was especially pronounced during the first lap in the fear-conditioned

environment compared to the neutral environment across all paradigms tested. 2) With

mice on a cylindrical treadmill, additional conditioned responses such as hesitation and

backward movement were detected. 3) The choice of neutral environment influenced fear

discrimination: mice froze more in the fear-conditioned VR when the neutral environment

was familiar rather than novel. Freezing in the fear-conditioned VR could be enhanced by

shorter ISIs and the presence of the tail-coat during recall when the neutral environment

was novel. 4) Following fear conditioning, place cells in the hippocampal CA1 region

showed remapping, increased numbers, and narrower fields, paralleling findings in freely

moving animals and offering new insights.

Here, we experimented with three paradigms for fear conditioning in mice, with the main

difference being the neutral environment where the mice didn't receive any shock. In

Paradigm 1, the neutral VR was familiar, where the mice had previously received rewards,

and the CFC VR was novel. In Paradigms 2 and 3, both the neutral and CFC VRs were

similarly novel. In Paradigm 3, we decreased the ISI and kept the tail-coat on the animal

when testing for memory recall. Thus, we have tested three different versions for

performing CFC in head-fixed mice. On average, mice showed more freezing behavior in

the CFC VR compared to the neutral VR. A common feature across all paradigms was

pronounced freezing during the very first lap in the CFC VR. While mice in Paradigm 2

extinguished fear rapidly after the first lap, freezing behavior persisted up to 3-4 recall

days in Paradigms 1 and 3, to varying degrees. We also observed variability in

conditioned responses across mice. Some displayed an opposite pattern, freezing more

in the neutral VR or equally in both VRs. This variability in behavior, similar to that

observed in freely moving animals (Chu et al., 2024; Dos Santos Corrêa et al., 2019;

Navarro-Sánchez et al., 2024; Poulos et al., 2016; Russo & Parsons, 2021; Totty et al.,

2021), can be insightful when combined with investigations of neural activity, helping to

understand the neural dynamics that contribute to an animal's ability to learn the

association between a context and fearful stimuli. This can reveal the mechanisms by

which such memories may generalize or become maladaptive, as seen in post-traumatic

stress disorders.

We have shown here an example of using Paradigm 1 to study place cells in the

hippocampus and their role in fear memory formation and recall. We recorded from a

large group of the same hippocampal CA1 cells across days and demonstrated that place

cells remap and narrow their place field widths with fear conditioning. This occurs in head-

fixed mice navigating VR environments, similar to freely moving animals. Narrow place

fields may suggest sharper spatial tuning to incorporate a salient emotional memory and

may aid in context discrimination (Schuette et al., 2020). However, how place cells narrow

their widths in response to fear conditioning remains an open question. One possible way

to answer this is by imaging CA1 place cell dendrites to understand how they integrate

inputs before and after fear conditioning. We can explore what dendritic firing and

plasticity mechanisms lead to narrow place field widths. We can then perturb these firing

patterns to attribute causality to the narrowing of place field widths in fear discrimination.

Our head-fixed paradigm enables the recording of dendrites and the study of plasticity

mechanisms (Sheffield et al., 2017; Sheffield & Dombeck, 2015), thus facilitating these

investigations.

We recently used Paradigm 3 to understand the role of the thalamic nucleus reuniens-

hippocampal pathway (Ratigan et al., 2023). Using a two-photon microscope, we

identified neural activity in this pathway by imaging axons from the nucleus reuniens that

project to the CA1 with single-axon resolution. We found this pathway necessary for

suppressing fear memory retrieval, context discrimination, and fear extinction. The robust

freezing response in Paradigm 3 in the fear-conditioned VR, compared to a neutral novel

VR, allowed us to align our findings with the existing literature on the role of the nucleus

reuniens-hippocampal circuit in fear memory in freely moving animals (Ramanathan &

Maren, 2019; Ramanathan et al., 2018; Totty et al., 2023; Troyner et al., 2018; Vetere et

al., 2017; Xu & Südhof, 2013). Furthermore, the subcellular resolution achieved through

two-photon calcium imaging using the head-fixed preparation allowed us to longitudinally

observe and understand the inputs that axons from the reuniens send to the hippocampus

during fear memory formation and retrieval, a level of detail that is difficult to achieve with

current techniques in freely moving preparations.

In another unpublished study, we used Paradigm 2 and identified co-active neurons

between CA1 and CA3 subregions of the hippocampus during fear memory recall. The

difference between the first and subsequent laps has helped us understand the

progression of neural activity in the hippocampus as mice go from retrieving the fear

memory to extinction within one session. Moreover, the variability in response across

mice has allowed us to regress the neural activity to the behavior, identifying neural

differences between those mice that accurately recall the fear memory and those that do

not. These studies showcase the broad applicability of our different CFC paradigms.

In the laboratory, CFC provokes freezing behavior in freely moving rodents, as

demonstrated by several studies that span decades (Michael S. Fanselow, 1980, 1990;

Fendt & Fanselow, 1999; Maren, 2001; Maren et al., 2013; Phillips & LeDoux, 1992;

Tovote et al., 2015). In our study, we found that a mild electric shock applied to the tail

could also trigger freezing behavior as the conditioned response in head-fixed mice,

making our paradigm more comparable to the freely moving paradigms than previously

attempted methods using air-puff stimulation (Lovett-Barron et al., 2014; Rajasethupathy

et al., 2015). There are numerous opportunities for future work to systematically test how

different parameters—such as shock amplitude, number and spacing of shocks, duration

of fear conditioning, context pre-exposure time, the configuration of Control versus CFC

VR environments, and the inclusion of additional sensory cues—may influence both

behavioral outcomes during recall and extinction, as well as the underlying neural

representations (Michael S. Fanselow, 1990; Huckleberry et al., 2016; Ji & Maren, 2007;

Maren & Holmes, 2016; Poulos et al., 2016). Our current study provides only an initial

exploration of this parameter space.

In this manuscript, we focused on freezing as the primary conditioned response to align

with studies of freely moving animals. However, we also see other defensive responses

unique to our treadmill-based task. as well as fear-related hesitation and backward

movements during memory recall—defensive behaviors that have also been observed in

freely moving animals (Chu et al., 2024; Furuyama et al., 2023; Totty et al., 2021; Trott et

al., 2022). Our technique enables us to conduct a more detailed analysis of the animal's

behavior. The head-fixed setup allows us to easily measure physiological responses

including pupil diameter, respiration, vocalization, heart rate, and body temperature.

These parameters are known to change with fear conditioning (Dupin et al., 2020; Godsil

et al., 2000; Korn et al., 2017; Shionoya et al., 2013; Stiedl & Spiess, 1997). Such

physiological measurements can also help differentiate between the various conditioned

responses we observe. There are also opportunities to use machine-learning-empowered

algorithms to parse out new behavior markers for contextual fear memory, such as

gestures or facial expressions (Datta et al., 2019; Mathis & Mathis, 2020; Syeda et al.,

2024; Wiltschko et al., 2020). Thus, we can understand the animal's behavior in a well-

controlled environment, which when combined with neural activity recording, can lead to

new insights into the neural circuits mediating different defensive responses and

emotional states.

Significant insights into memory mechanisms through the CFC task have also come from

recent discoveries of memory engrams in the brain (Josselyn et al., 2015; Josselyn &

Tonegawa, 2020; Liu et al., 2012; Tonegawa et al., 2018). Engram cells, identified in

various brain regions with the help of genetic tagging and optogenetics, play a crucial role

in encoding, consolidating, and retrieving specific memories. Typically, these cells are

identified based on the expression of the immediate early gene *Fos* during fear

conditioning. Their artificial activation with optogenetics can trigger memory recall,

evidenced by increased freezing behavior. Understanding which neurons form these

engram cells, how *Fos* expression is driven during memory formation, how memory

information is stored in these cells, how they evolve with learning, and how their activation

results in memory recall is an active area of research (Frankland et al., 2024; Mocle et

al., 2024; Monasterio et al., 2024; Pettit et al., 2022; Suthard et al., 2024; Uytiepo et al.,

2024). Incorporating our paradigm with simultaneous measurements of cellular and

subcellular activity can help answer some of these questions. For instance, a recent study

(Monasterio et al., 2024) tagged engram cells during fear conditioning while animals were

freely moving and then imaged the spontaneous dynamics of these cells using two-

photon imaging before and after fear conditioning. The study revealed that cells forming

engrams were inherently more active. A limitation of this study was the separate contexts

for imaging versus tagging, which can be addressed with our paradigm. Our CFC

paradigm allows for direct neural recording of these cells before, during, and after tagging

in the same context to examine how activity during memory formation relates to engram

cell emergence, population dynamics, and memory recall. Furthermore, holographic

stimulation techniques can reactivate specific cell ensembles of interest to help

understand the causal relationships between engram cells and memory. Moreover, the

hippocampus contains both place cells and engram cells, both of which appear to be

involved in contextual memory. However, their relationship is not yet fully understood. Our

CFC paradigm offers an opportunity to study both place cells, as we have shown here,

and engram cells simultaneously. This can be achieved by incorporating techniques that

allow for *in vivo* labeling of *Fos* (Pettit et al., 2022). Thus, our head-fixed version of the

CFC paradigm opens new avenues for research into the neural underpinnings of fear

learning and memory.

Another advantage of our method is that it allows for studying changes in neuronal

dynamics in response to an aversive stimulus, such as a shock. Our fields of view during

the shock and subsequent flight-like response periods remained stable, and we could

correct image movements during these periods with Suite2p (Pachitariu et al., 2017). This

is a significant improvement over one-photon and electrophysiological techniques, where

abrupt movements such as those induced by an electric shock can change fields of view

and reduce the likelihood of recording from the same groups of cells, thereby hindering

longitudinal assessments of neural dynamics. In Paradigm 1, where we imaged

hippocampal cells during shock, we did not find cells responding to the shock itself, as

others have reported (Barth et al., 2023; Mocle et al., 2024; Suthard et al., 2024). We

suspect this could be due to the number or duration of shocks administered and our

shocks not being associated with a specific cue or location. In future studies, we plan to

optimize these parameters to understand better how neural dynamics change in response

to an aversive cue.

Finally, we recognize that our head-fixed CFC paradigm has some limitations. Compared

to the freely moving CFC, our setup is less naturalistic. We defined different contexts as

two visually distinct VR environments while odor, sound, and tactile cues remained

constant. This could have made it harder for mice to differentiate between the two

contexts, particularly in Paradigm 2. Unlike freely moving animals in CFC studies, our

mice were water-restricted throughout the experimental process. Because the animals

need to get used to wearing a tailcoat and running in the VRs without a reward, we tend

to 'over-train' the mice. This criterion excludes mice that are less adept at running, limiting

the number that go through the process and requiring larger cohorts of mice. This also

presents scalability challenges compared to freely moving paradigms, as successful

implementation of the head-fixed CFC approach requires weeks of reward-based training,

larger sample sizes, and a technically demanding VR setup that integrates multiple

hardware and software components to precisely monitor behavior. Additionally, while

rewards were removed during experimental sessions, water rewards were still used to

motivate running during training. This reward-associated training could subtly influence

later behavior. For instance, the increased freezing in the Familiar VR on later recall days

and remapping of place cells in that context could reflect the lingering effects of reward-

associated training. While Paradigms 2 and 3 address many of these concerns, our

approach still differs from freely moving paradigms due to this initial reward-based

training. The impact of using rewards during training on recall behavior needs further

investigation. Furthermore, despite our best efforts to include equal numbers of male and

female mice, only some female mice reached the experimental stage. This could be due

to the tail-coat weight and the reduced weight of females after water restriction compared

to males. Female mice of the same age (10-12 weeks) weighed significantly less than

males (20-23 g vs 25-30 g) and lost further weight following water restriction. Although

we observed no behavioral differences between the four tested females and the male

mice, we cannot guarantee similar results in other female animals without further testing.

Studies indicate that different sexes may display varying conditioned responses, as seen

in rats (Gruene et al., 2015; S. Maren et al., 1994). In the future, we aim to address these

limitations.

In conclusion, we have developed a head-fixed version of the classic CFC paradigm. Our

findings demonstrate that mice exhibit freezing behavior when fear-conditioned in a VR

environment. Moreover, we observed that hippocampal place cells remap and narrow

their fields in response to fear conditioning, mirroring observations in freely moving

animals. This innovative paradigm bridges the behavioral task gap for techniques

requiring head-fixation, paving the way for new investigations into neural circuits and

deepening our understanding of the neural foundations of emotional memory.

Acknowledgments: We thank the members of the Sheffield Lab for comments on the manuscript, helpful discussions, and feedback throughout the process.

Funding: This work was supported by The Whitehall Foundation, The Searle Scholars Program, The Sloan Foundation, The University of Chicago Institute for Neuroscience start-up funds and the National Institute for Health (1DP2NS111657-01, 1RF1NS127123-01) awarded to M.S. and a T32 training grant (T32DA043469) and K01 (K01DA060994-01) from National Institute on Drug Abuse awarded to S.K.

Author contributions: S.K. and M.S. conceived the idea. S.K. and M.S. supervised the project. S.K. designed the experiments. S.K., C.D., H.R., D.M., and C.C. collected the data. S.K. and C.D. analyzed the data. S.K., C.D., H.R., and M.S. interpreted the data. S.K. prepared the manuscript draft. All authors reviewed the results, provided feedback, and approved the final version of the manuscript.

Declaration of interests: The authors declare no competing interests.

Data and code availability: Behavior data, imaging data, and scripts used for data analysis are available on GitHub at https://github.com/seethakris/CFCMethodspaper.

Supplementary Video Legend

Supplementary Videos 1, 2 and 3: Example video of a head-fixed mouse navigating on a treadmill in VR. The examples show mouse behavior on Recall Day 1 in Paradigm 1 in: **1)** the Familiar VR and **2 and 3)** the CFC VR. Video 2 shows freezing behavior in CFC VR and Video 3 shows another mouse displaying more avoidance-like behaviors like slowing down and backward movements. Videos were recorded for a few seconds immediately after the mouse was transitioned to each VR environment. Videos were collected at a sampling rate of 30 frames/sec.

Methods

Subjects

All experimental and surgical procedures adhered to the University of Chicago Animal Care and Use Committee guidelines. This study utilized 10-12 week-old C57BL/6J, Thy1GCaMP6s and Tg(Grik4-cre)G32-4Stl/J mice. Following the start of water restriction, the mice were individually housed in a reverse 12-hour light/dark cycle, and behavioral experiments were conducted during the animals' dark cycle.

Head-fixation surgery

Mice were anesthetized with 1-2% isoflurane and given an intraperitoneal injection of 0.5 mL of saline and a subcutaneous injection of 0.45 mL of Meloxicam (1-2 mg/kg). A small incision was made on the skull to attach a metal head-plate (9.1 mm x 31.7 mm, Atlas Tool and Die Works) to head-restrain the mice. Some mice also had an imaging window implanted above the hippocampus, as previously described (Dong et al., 2021; Krishnan et al., 2022; Sheffield & Dombeck, 2015). Post-surgery mice were housed individually. After a recovery day, they were water-restricted to 0.8-1.0 ml/day.

Behavior training

Head-restrained mice were trained to run on a cylindrical treadmill to navigate virtual reality (VR) environments similar to the setups previously described (Dong et al., 2021; Krishnan et al., 2022; Sheffield & Dombeck, 2015). Behavioral training began seven days following the water restriction. Briefly, VR environments were created using the Virtual Reality MATLAB Engine, or VIRMEN (Aronov & Tank, 2014), and projected to five

screens covering the mouse's field of view. Mice were head restrained with their limbs

resting on a freely rotating Styrofoam wheel, which acted as a treadmill. VR environments

were 2 m linear tracks. Mice received a water reward (4 µl) through a waterspout upon

completing each traversal of the linear track, considered a lap. This reward was followed

by a 1.5-second pause in the VR to allow the mouse to consume the water. The mice

were then virtually teleported to the start of the track to begin a new lap. Training sessions

lasted 30 minutes per day. Mice were considered well-trained when they could run at >3

laps per minute. We found that this level of training was necessary to ensure that mice

would continue to run even without rewards (Krishnan et al., 2022; Krishnan & Sheffield,

2023). Generally, about 60% of mice met our criteria, which typically took around 10-14

days to achieve. Behavior data was collected using a PicoScope Oscilloscope

(PICO4824, Pico Technology, v6.13.2).

Tail-coat

A custom-designed "tail-coat" delivered mild electric shocks to a head-fixed mouse's tail

during virtual reality (VR) navigation. A pictorial representation of the tail-coat can be

found in **Figure 1D**. The tail-coat, handmade from wearable conductive fabrics, featured

a lightweight design to ensure the mouse could continue running during baseline VR

exploration (**Table 7**). First, the mouse's tail was measured while anesthetized for head-

plate surgeries. A small denim patch cloth (Michael's) was cut into a rectangle to fit snugly

on the mouse's tail. Marks were made where snap buttons (Michael's) would be sewn,

ensuring they clipped just above the tail. Given that the mouse's tail narrows away from

the body, the buttons closer to the body were placed slightly further apart than those

closer to the tip. Typically, the buttons were set 2 cm apart on the narrow end and 2.5 cm

apart on the broader end. Care was taken to adjust the placement of the buttons so they

weren't too tight or loose for the mouse. We found that these measurements were

generally similar across sexes and for the age group that we tested. After measurements

were taken and the cloth cut, small strips of 8mm wide conductive fabric tape (Adafruit)

were attached to either side of the cloth, with some tape hanging off the edge. A metal

strip (McMaster Carr) of the same width was affixed to this portion of the tape to provide

a secure grip for alligator clips used in fear conditioning. Finally, snap buttons were sewn

with conductive thread at the marked spots. Clear nail polish was applied to secure the

threads in place. The tail-coat creation process should take less than half an hour.

Dimensions of the cloth for a typical male mouse, 12 weeks of age and weighing ~30 g,

are shown in **Figure 1D**. The denim cloth was 4 cm in width and 3.2 cm in length; the

conductive tape was 1.2 cm in width and 5.2 cm in length, with 1.5 cm hanging out at

each side.

Table 7: Items required to for head-fixed contextual fear discrimination		
Item	Manufacturer	Link
Denim patch cloth	Any fabric or hobby shop	https://www.michaels.com/product/loops-threads-denim-patches-assorted-10113795
Snap Buttons	Any fabric or hobby shop	https://www.michaels.com/product/loops-threads-sew-on-snaps-10354228?michaelsStore=1191&inv=14
Conductive nylon fabric tape	Adafruit	https://www.adafruit.com/product/3960
Stainless conductive thread – 3 Ply	Adafruit	https://www.adafruit.com/product/641
Multipurpose 304 Stainless steel sheet	McMaster-Carr	https://www.mcmaster.com/products/304-stainless-steel/multipurpose-304-stainless-steel-6/?s=304-stainless-steel

Shock box	Colbourn precision animal shocker Alternate: Lafayette Instruments Scrambled Grid Current Generator	https://lafayetteinstrument.com

Contextual fear conditioning protocol in head-fixed mice

Once the mice were well-trained and met our criteria of running >3 laps per minute, they

were ready for the next stage. The following day, a tail-coat was attached to the mouse's

tail. Mice that maintained consistent running in the VR, with minimal pauses, while

wearing the tail-coat advanced to the next stage. For mice startled by the tail-coat, we

typically attempted another day with the tail-coat, rewarding them and observing if their

running behavior improved to running at least three laps per minute. If not, these mice

were not advanced to the next stage. In mice that do advance, on the subsequent day,

the tail-coat was kept on, and the water reward was removed. All else remained the same,

including the 1.5 s pause at the end, to allow animals to distinguish the start and end of

a lap. If the mice continued to run as before, they moved on to the experiment stage.

From that day forward, the mice no longer received water reinforcement when running in

the VR environments. Typically, about 40% of mice made it to the experiment stage. Tail-

coats were only added during VR exploration and removed before returning the animals

to their home cage. Once the mice advanced to the experiment stage, they were taken

through the entire process and not excluded from analysis, even if they showed signs of

inconsistent running behavior, which happened with some mice (for instance, see

Supplementary Figure 3B-C).

We implemented three distinct paradigms to assess contextual fear discrimination. In the

first paradigm, we conducted experiments over two days (Paradigm 1). On Day 0, mice

engaged with the same VR in which they were trained, now referred to as the Familiar

VR. After 10 minutes, we transitioned them to a new VR for another 10 minutes; this new

VR would later be associated with shocks (CFC VR). During this phase, the mice explored

both VRs without any rewards while wearing the tail-coat. After the initial exploration in

the CFC VR, they received six mild electric shocks, each lasting 1 second with 1-minute

intervals in between. The shocks were administered using a shock generator (Colbourn

Precision animal shocker, Harvard Bioscience). A TTL pulse was generated using

custom-written codes in MATLAB to turn on the shock generator via an Arduino. The

shock times were recorded by the PicoScope Oscilloscope (PICO4824, Pico Technology,

v6.13.2). We tested different shock intensities ranging from 0.5 to 1.2 mA. After the last

shock, mice were removed from the VR and returned to their home cage. The following

day (Day 1), we placed the mice in the Familiar VR or CFC VR for ten minutes to test

their memory recall. No rewards were delivered during this phase; however, the tail-coat

was not added during memory recall. Some mice underwent additional recall tests over

subsequent days to assess the time required for fear memory extinction. As a control,

some mice underwent the same paradigm but without any shocks.

In the second paradigm (Paradigm 2), the experiment spanned over three days. On Day

- 1, mice explored two novel VRs for ten minutes each, establishing a baseline for context

exploration. One of these VRs served as a control, while the other, the CFC VR, was

where the mice would later receive electric shocks. On Day 0, the mice again spent ten

minutes each in the Control and CFC VRs. After this, they received six mild electric

shocks, each lasting one second, with a strength of 1 mA and an inter-trial interval of one

minute. The mice wore the tail-coat on both Days -1 and 0. On Day 1, the mice were

placed in either the Control VR or CFC VR for ten minutes each, in a counterbalanced

manner, to test for memory recall. No rewards were given on all the experimental days;

however, the tail-coat was not added to the tails on the recall day (Day 1). The third

paradigm (Paradigm 3) was similar to Paradigm 2 and used two novel VRs, except that

a tail-coat was also added to the mice's tails during recall tests.

Additionally, a subset of mice in Paradigms 1 and 2 and all mice in Paradigm 3 went

through multiple days of recall tests. The subset of mice was randomly chosen, not based

on their recall behavior from the previous day. Mice were presented with either the CFC

VR or the Control VR in a counterbalanced manner.

All VRs were randomly assigned as either Control or CFC VR to prevent biasing any

inherent preferences in mice. However, the Familiar VR was always kept the same. We

also used both sexes in our experiments. However, we found female mice less receptive

to the "tail-coat" than their male counterparts. We believe this is due to the low average

weight of female mice (~16 g) following water restriction. Female mice of the same age

(10-12 weeks) weighed significantly less than males (20-23 g vs 25-30 g) and lost further

weight following water restriction. As a result, only four female mice could proceed

through and complete all the behavioral tests. Our ultimate goal is to compare behavior

and neural activity across sexes, considering sex as a biological variable. To achieve this,

we plan to adjust the age and use female mice of appropriate weight to ensure they can

run on the treadmill with the tailcoat in the future.

Behavior measurement

Freezing periods were periods during which the animal's instantaneous velocity was 0 cm/s. Freezing epochs were defined as continuous, uninterrupted freezing periods longer than 1 s. The time taken to complete a lap was calculated as the total time (in seconds) the animal took to run from 0 to 200 cm.

Surgery and two-photon imaging

Stereotaxic surgeries and cannula window implantation above the CA1 subregion of the hippocampus were performed in the same way as previously reported (Krishnan et al., 2022). Briefly, a genetically-encoded calcium indicator, AAV1-CamKII-GCaMP6f (pENN.AAV.CamKII.GCaMP6f.WPRE.SV40 was a gift from James M. Wilson – Addgene viral prep #100834-AAV1; https://www.addgene.org/100834/; RRID: Addgene_100834) was injected (~50 nL at a depth of 1.25 mm below the surface of the dura) using a beveled glass micropipette leading to GCaMP6f expression in a large population of CA1 pyramidal cells. Mice underwent water restriction (1.0 ml/day) following viral injection and, seven days later, were implanted with a hippocampal window and head plate (Dombeck et al., 2010). Mice were allowed to rest from the surgeries for 3-4 days, after which behavioral training began. Imaging was done using a laser scanning two-photon microscope (Neurolabware). The setup is the same as the one previously described (Krishnan et al., 2022). Time-series images were collected through Scanbox (v4.1, Neurolabware), and the PicoScope Oscilloscope (PICO4824, Pico Technology, v6.13.2) was used to synchronize frame acquisition timing with behavior. At the end of the imaging session on Day -1 or Day 0, a 1-minute time-series movie was collected at a higher magnification

and then averaged to aid as a reference frame in finding the same imaging plane on

subsequent days. Across-day images collected were motion-corrected using Suite2p

(Pachitariu et al., 2017) and aligned using ImageJ (v1.53, NIH). Region of interests were

extracted using Suite2p.

Place cell extraction and place field parameters

Place fields were identified as previously described (Dong et al., 2021; Krishnan et al.,

2022). Briefly, the 2 m track was divided into 40 position bins (each 5 cm wide). The

running behavior of the animal was filtered to exclude periods where the animal was

immobile (speed <1 cm/s). Filtering ensured that place cells were defined only during

active exploration and not during freezing bouts. We have previously found that filtering

versus not filtering yields similar results (Krishnan et al., 2022). Extracted place fields

satisfied the following criteria, which were used for all conditions and all mice: 1. The

average $\Delta F/F$ was greater than 10% above the baseline. 2. The cell displayed significant

calcium transients on >30% of laps in the field. 3. The rising phase of the mean transient

was located on the track. 4. Their p-value from bootstrapping was <0.05. Multiple place

fields within the same cell were treated independently. Parameters of place fields were

calculated as described before (Krishnan et al., 2022); this includes the center of mass

(COM), place field reliability, out/in-field firing ratio, and place field widths.

Statistics

For related samples, we performed a paired t-test. Multiple comparisons of related

samples were corrected with Bonferroni post-hoc. To quantify freezing across days, we

employed a linear mixed-effects model with days of recall as the fixed effect and behavior

prior to fear conditioning on Day 0 as baseline. The model also included a random effect

for mice to account for the nesting of observations within individual subjects. Separate

models were used for each VR. Boxplots are plotted to display the entire distribution of

the data. The box in the boxplot ranges from the first quartile (25th percentile) to the third

quartile (75th percentile), and the box shows the interquartile range (IQR). The line across

the box represents the median (50th percentile). The whiskers extend to 1.5*IQR on either

side of the box, and anything above this range is defined as an outlier. A two-tailed

Kolmogorov-Smirnov (KS) Test was used to compare distributions. When not using

boxplots, mean and confidence intervals are displayed.

References

Anagnostaras, S. G., Gale, G. D., & Fanselow, M. S. (2001). Hippocampus and contextual fear conditioning: Recent controversies and advances. *Hippocampus*, *11*(1), 8 17.

Aronov, D., & Tank, D. W. (2014). Engagement of Neural Circuits Underlying 2D Spatial Navigation in a Rodent Virtual Reality System. *Neuron*, *84*(2), 442 456.

Baldi, E., Lorenzini, C. A., & Bucherelli, C. (2004). Footshock intensity and generalization in contextual and auditory-cued fear conditioning in the rat. *Neurobiology of Learning and Memory*, *81*(3), 162–166.

Bando, Y., Wenzel, M., & Yuste, R. (2021). Simultaneous two-photon imaging of action potentials and subthreshold inputs in vivo. *Nature Communications*, *12*(1), 7229.

Barth, A. M., Jelitai, M., Vasarhelyi-Nagy, M. F., & Varga, V. (2023). Aversive stimulus-tuned responses in the CA1 of the dorsal hippocampus. *Nature Communications*, *14*(1), 1–17.

Biedenkapp, J. C., & Rudy, J. W. (2007). Context preexposure prevents forgetting of a contextual fear memory: implication for regional changes in brain activation patterns associated with recent and remote memory tests. *Learning & Memory (Cold Spring Harbor, N.Y.)*, *14*(3), 200–203.

Blair, G. J., Guo, C., Wang, S., Fanselow, M. S., Golshani, P., Aharoni, D., & Blair, H. T. (2023). Hippocampal place cell remapping occurs with memory storage of aversive experiences. *ELife*, *12*. https://doi.org/10.7554/eLife.80661

Bouton, M. E. (1993). Context, time, and memory retrieval in the interference paradigms of Pavlovian learning. *Psychological Bulletin*, *114*(1), 80–99.

Bouton, M. E., & King, D. A. (1983). Contextual control of the extinction of conditioned fear: tests for the associative value of the context. *Journal of Experimental Psychology. Animal Behavior Processes*, *9*(3), 248–265.

Bouton, Mark E., & Bolles, R. C. (1980). Conditioned fear assessed by freezing and by the suppression of three different baselines. *Animal Learning & Behavior*, *8*(3), 429–434.

Cai, D. J., Shuman, T., Harrison, E. M., Sage, J. R., & Anagnostaras, S. G. (2009). Sleep deprivation and Pavlovian fear conditioning. *Learning & Memory* , *16*(10), 595–599.

Chaigneau, E., Ronzitti, E., Gajowa, M. A., Soler-Llavina, G. J., Tanese, D., Brureau, A. Y. B., Papagiakoumou, E., Zeng, H., & Emiliani, V. (2016). Two-Photon Holographic Stimulation of ReaChR. *Frontiers in Cellular Neuroscience*, *10*, 234.

Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., Schreiter, E. R., Kerr, R. A., Orger, M. B., Jayaraman, V., Looger, L. L., Svoboda, K., & Kim, D. S. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, *499*(7458), 295 300.

Chu, A., Gordon, N. T., DuBois, A. M., Michel, C. B., Hanrahan, K. E., Williams, D. C., Anzellotti, S., & McDannald, M. A. (2024). A fear conditioned cue orchestrates a suite of behaviors in rats. *ELife*, *13*. https://doi.org/10.7554/eLife.82497

Colgin, L. L., Moser, E. I., & Moser, M.-B. (2008). Understanding memory through hippocampal remapping. *Trends in Neurosciences*, *31*(9), 469 477.

Cordero, M. I., Merino, J. J., & Sandi, C. (1998). Correlational relationship between shock intensity and corticosterone secretion on the establishment and subsequent expression of contextual fear conditioning. *Behavioral Neuroscience*, *112*(4), 885–891.

Datta, S. R., Anderson, D. J., Branson, K., Perona, P., & Leifer, A. (2019). Computational Neuroethology: A Call to Action. *Neuron*, *104*(1), 11–24.

Daviu, N., Molina, P., Nadal, R., Belda, X., Serrano, S., & Armario, A. (2024). Influence of footshock number and intensity on the behavioral and endocrine response to fear conditioning and cognitive fear generalization in male rats. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, *135*(111112), 111112.

Debiec, J., LeDoux, J. E., & Nader, K. (2002). Cellular and systems reconsolidation in the hippocampus. *Neuron*, *36*(3), 527 538.

Dombeck, D. A., Harvey, C. D., Tian, L., Looger, L. L., & Tank, D. W. (2010). Functional imaging of hippocampal place cells at cellular resolution during virtual navigation. *Nature Neuroscience*, *13*(11), 1433–1440.

Dong, C., Madar, A. D., & Sheffield, M. E. J. (2021). Distinct place cell dynamics in CA1 and CA3 encode experience in new environments. *Nature Communications*, *12*(1), 2977.

Dos Santos Corrêa, M., Vaz, B. D. S., Grisanti, G. D. V., de Paiva, J. P. Q., Tiba, P. A., & Fornari, R. V. (2019). Relationship between footshock intensity, post-training corticosterone release and contextual fear memory specificity over time. *Psychoneuroendocrinology*, *110*(104447), 104447.

Dupin, M., Garcia, S., Messaoudi, B., Doyère, V., & Mouly, A.-M. (2020). Respiration and brain neural dynamics associated with interval timing during odor fear learning in rats. *Scientific Reports*, *10*(1), 17643.

Emiliani, V., Cohen, A. E., Deisseroth, K., & Hausser, M. (2015). All-Optical Interrogation of Neural Circuits. *Journal of Neuroscience*, *35*(41), 13917 13926.

Fan, L. Z., Kim, D. K., Jennings, J. H., Tian, H., Wang, P. Y., Ramakrishnan, C., Randles, S., Sun, Y., Thadhani, E., Kim, Y. S., Quirin, S., Giocomo, L., Cohen, A. E., & Deisseroth, K. (2023). All-optical physiology resolves a synaptic basis for behavioral timescale plasticity. *Cell*, *186*(3), 543-559.e19.

Fanselow, M. S., & Bolles, R. C. (1979). Naloxone and shock-elicited freezing in the rat. *Journal of Comparative and Physiological Psychology*, *93*(4), 736–744.

Fanselow, Michael S. (1980). Conditional and unconditional components of post-shock freezing. *The Pavlovian Journal of Biological Science : Official Journal of the Pavlovian*, *15*(4), 177–182.

Fanselow, Michael S. (1990). Factors governing one-trial contextual conditioning. *Animal Learning & Behavior*, *18*(3), 264–270.

Fendt, M., & Fanselow, M. S. (1999). The neuroanatomical and neurochemical basis of conditioned fear. *Neuroscience and Biobehavioral Reviews*, *23*(5), 743–760.

Frankland, P. W., Josselyn, S. A., & Köhler, S. (2024). Engrams. *Current Biology: CB*, *34*(12), R559–R561.

Furuyama, T., Imayoshi, A., Iyobe, T., Ono, M., Ishikawa, T., Ozaki, N., Kato, N., & Yamamoto, R. (2023). Multiple factors contribute to flight behaviors during fear conditioning. *Scientific Reports*, *13*(1), 10402.

Gewirtz, J. C., McNish, K. A., & Davis, M. (2000). Is the hippocampus necessary for contextual fear conditioning? *Behavioural Brain Research*, *110*(1–2), 83–95.

Godsil, B. P., Quinn, J. J., & Fanselow, M. S. (2000). Body temperature as a conditional response measure for pavlovian fear conditioning. *Learning & Memory* , *7*(5), 353–356.

Gruene, T. M., Flick, K., Stefano, A., Shea, S. D., & Shansky, R. M. (2015). Sexually divergent expression of active and passive conditioned fear responses in rats. *ELife*, *4*. https://doi.org/10.7554/eLife.11352

Guo, Z. V., Hires, S. A., Li, N., O'Connor, D. H., Komiyama, T., Ophir, E., Huber, D., Bonardi, C., Morandell, K., Gutnisky, D., Peron, S., Xu, N.-L., Cox, J., & Svoboda,

K. (2014). Procedures for behavioral experiments in head-fixed mice. *PloS One*, *9*(2), e88678.

Hattori, R., & Komiyama, T. (2022). Longitudinal two-photon calcium imaging with ultra-large cranial window for head-fixed mice. *STAR Protocols*, *3*(2), 101343.

Heer, C. M., & Sheffield, M. E. J. (2024). *Distinct catecholaminergic pathways projecting to hippocampal CA1 transmit contrasting signals during navigation in familiar and novel environments*. https://doi.org/10.7554/elife.95213

Hernandez, O., Papagiakoumou, E., Tanese, D., Fidelin, K., Wyart, C., & Emiliani, V. (2016). Three-dimensional spatiotemporal focusing of holographic patterns. *Nature Communications*, *7*, 11928.

Holland, P. C., & Bouton, M. E. (1999). Hippocampus and context in classical conditioning. *Current Opinion in Neurobiology*, *9*(2), 195–202.

Huckleberry, K. A., Ferguson, L. B., & Drew, M. R. (2016). Behavioral mechanisms of context fear generalization in mice. *Learning & Memory*, *23*(12), 703–709.

Ji, J., & Maren, S. (2007). Hippocampal involvement in contextual modulation of fear extinction. *Hippocampus*, *17*(9), 749–758.

Josselyn, S. A., Köhler, S., & Frankland, P. W. (2015). Finding the engram. *Nature Reviews. Neuroscience*, *16*(9), 521 534.

Josselyn, S. A., & Tonegawa, S. (2020). Memory engrams: Recalling the past and imagining the future. *Science*, *367*(6473). https://doi.org/10.1126/science.aaw4325

Juczewski, K., Koussa, J. A., Kesner, A. J., Lee, J. O., & Lovinger, D. M. (2020). Stress and behavioral correlates in the head-fixed method: stress measurements, habituation dynamics, locomotion, and motor-skill learning in mice. *Scientific Reports*, *10*(1), 12245.

Kim, E. J., Park, M., Kong, M.-S., Park, S. G., Cho, J., & Kim, J. J. (2015). Alterations of hippocampal place cells in foraging rats facing a "predatory" threat. *Current Biology: CB*, *25*(10), 1362–1367.

Kim, J. J., & Fanselow, M. S. (1992). Modality-specific retrograde amnesia of fear. *Science*, *256*(5057), 675 677.

Kim, Jeansok J., & Jung, M. W. (2006). Neural circuits and mechanisms involved in Pavlovian fear conditioning: a critical review. *Neuroscience and Biobehavioral Reviews*, *30*(2), 188–202.

Kinsky, N. R., Orlin, D. O., Ruesch, E. A., Diba, K., & Ramirez, S. (2023). Erasable hippocampal neural signatures predict memory discrimination. *BioRxiv.Org: The Preprint Server for Biology*, 2023.02.02.526824.

Korn, C. W., Staib, M., Tzovara, A., Castegnetti, G., & Bach, D. R. (2017). A pupil size response model to assess fear learning. *Psychophysiology*, *54*(3), 330–343.

Krishnan, S., Heer, C., Cherian, C., & Sheffield, M. E. J. (2022). Reward expectation extinction restructures and degrades CA1 spatial maps through loss of a dopaminergic reward proximity signal. *Nature Communications*, *13*(1), 6662.

Krishnan, S., & Sheffield, M. E. J. (2023). Reward Expectation Reduces Representational Drift in the Hippocampus. In *bioRxiv* (p. 2023.12.21.572809). https://doi.org/10.1101/2023.12.21.572809

Leaf, R. C., & Leaf, S. R. (1966). Recovery time as a measure of degree of conditioned suppression. *Psychological Reports*, *18*(1), 265–266.

Leaf, R. C., & Muller, S. A. (1965). Simple method for CER conditioning and measurement. *Psychological Reports*, *17*(1), 211–215.

LeDoux, J. E. (2000). Emotion circuits in the brain. *Annual Review of Neuroscience*, *23*, 155–184.

Leutgeb, S., Leutgeb, J. K., Barnes, C. A., Moser, E. I., McNaughton, B. L., & Moser, M.-B. (2005). Independent codes for spatial and episodic memory in hippocampal neuronal ensembles. *Science (New York, N.Y.)*, *309*(5734), 619–623.

Liu, X., Ramirez, S., Pang, P. T., Puryear, C. B., Govindarajan, A., Deisseroth, K., & Tonegawa, S. (2012). Optogenetic stimulation of a hippocampal engram activates fear memory recall. *Nature*, *484*(7394), 381.

Lovett-Barron, M., Kaifosh, P., Kheirbek, M. A., Danielson, N., Zaremba, J. D., Reardon, T. R., Turi, G. F., Hen, R., Zemelman, B. V., & Losonczy, A. (2014). Dendritic inhibition in the hippocampus supports fear learning. *Science*, *343*(6173), 857 863.

Maboudi, K., Ackermann, E., de Jong, L. W., Pfeiffer, B. E., Foster, D., Diba, K., & Kemere, C. (2018). Uncovering temporal structure in hippocampal output patterns. *ELife*, *7*. https://doi.org/10.7554/eLife.34467

Mamad, O., Agayby, B., Stumpp, L., Reilly, R. B., & Tsanov, M. (2019). Extrafield activity shifts the place field center of mass to encode aversive experience. *ENeuro*, *6*(2), ENEURO.0423-17.2019.

Maren. (2001). Neurobiology of Pavlovian fear conditioning. *Annual Review of Neuroscience*, *24*(1), 897 931.

Maren, & Holmes, A. (2016). Stress and Fear Extinction. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *41*(1), 58–79.

Maren, Phan, K. L., & Liberzon, I. (2013). The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nature Reviews. Neuroscience*, *14*(6), 417–428.

Maren, S., De Oca, B., & Fanselow, M. S. (1994). Sex differences in hippocampal long-term potentiation (LTP) and Pavlovian fear conditioning in rats: positive correlation between LTP and contextual learning. *Brain Research*, *661*(1–2), 25–34.

Maren, Stephen, & Holt, W. (2000). The hippocampus and contextual memory retrieval in Pavlovian conditioning. *Behavioural Brain Research*, *110*(1–2), 97 108.

Mathis, M. W., & Mathis, A. (2020). Deep learning tools for the measurement of animal behavior in neuroscience. *Current Opinion in Neurobiology*, *60*, 1–11.

McNally, G. P., & Westbrook, R. F. (2006). A short intertrial interval facilitates acquisition of context-conditioned fear and a short retention interval facilitates its expression. *Journal of Experimental Psychology. Animal Behavior Processes*, *32*(2), 164–172.

Mocle, A. J., Ramsaran, A. I., Jacob, A. D., Rashid, A. J., Luchetti, A., Tran, L. M., Richards, B. A., Frankland, P. W., & Josselyn, S. A. (2024). Excitability mediates allocation of pre-configured ensembles to a hippocampal engram supporting contextual conditioned threat in mice. *Neuron*, *112*(9), 1487-1497.e6.

Moita, M. A. P., Rosis, S., Zhou, Y., LeDoux, J. E., & Blair, H. T. (2004). Putting fear in its place: remapping of hippocampal place cells during fear conditioning. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *24*(31), 7015–7023.

Monasterio, A., Lienkaemper, C., Coello, S., Ocker, G. K., Ramirez, S., & Scott, B. B. (2024). CA1 Engram Cell Dynamics Before and After Learning. In *bioRxiv* (p. 2024.04.16.589790). https://doi.org/10.1101/2024.04.16.589790

Muller, R. U., & Kubie, J. L. (1987). The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *Journal of Neuroscience*, *7*(7), 1951 1968.

Navarro-Sánchez, M., Gil-Miravet, I., Montero-Caballero, D., Castillo-Gómez, E., Gundlach, A. L., & Olucha-Bordonau, F. E. (2024). Some key parameters in contextual fear conditioning and extinction in adult rats. *Behavioural Brain Research*, *462*, 114874.

O'Keefe, J., & Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Research*, *34*(1), 171 175.

Ormond, J., Serka, S. A., & Johansen, J. P. (2023). Enhanced Reactivation of Remapping Place Cells during Aversive Learning. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *43*(12), 2153–2167.

Pachitariu, M., Stringer, C., Dipoppa, M., Schröder, S., Rossi, L. F., Dalgleish, H., Carandini, M., & Harris, K. D. (2017). Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *BioRxiv*, 061507.

Pettit, N. L., Yap, E.-L., Greenberg, M. E., & Harvey, C. D. (2022). Fos ensembles encode and shape stable spatial maps in the hippocampus. *Nature*. https://doi.org/10.1038/s41586-022-05113-1

Pezze, M. A., & Feldon, J. (2004). Mesolimbic dopaminergic pathways in fear conditioning. *Progress in Neurobiology*, *74*(5), 301–320.

Phillips, R. G., & LeDoux, J. E. (1992). Differential contribution of amygdala and hippocampus to cued and contextual fear conditioning. *Behavioral Neuroscience*, *106*(2), 274–285.

Poulos, A. M., Mehta, N., Lu, B., Amir, D., Livingston, B., Santarelli, A., Zhuravka, I., & Fanselow, M. S. (2016). Conditioning- and time-dependent increases in context fear and generalization. *Learning & Memory* , *23*(7), 379 385.

Quinn, J. J., Wied, H. M., Ma, Q. D., Tinsley, M. R., & Fanselow, M. S. (2008). Dorsal hippocampus involvement in delay fear conditioning depends upon the strength of the tone-footshock association. *Hippocampus*, *18*(7), 640–654.

Rajasethupathy, P., Sankaran, S., Marshel, J. H., Kim, C. K., Ferenczi, E., Lee, S. Y., Berndt, A., Ramakrishnan, C., Jaffe, A., Lo, M., Liston, C., & Deisseroth, K. (2015). Projections from neocortex mediate top-down control of memory retrieval. *Nature*, *526*(7575), 653 659.

Ramanathan, K. R., & Maren, S. (2019). Nucleus reuniens mediates the extinction of contextual fear conditioning. *Behavioural Brain Research*, *374*(112114), 112114.

Ramanathan, K. R., Ressler, R. L., Jin, J., & Maren, S. (2018). Nucleus Reuniens Is Required for Encoding and Retrieving Precise, Hippocampal-Dependent Contextual Fear Memories in Rats. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *38*(46), 9925–9933.

Ramirez, S., Liu, X., Lin, P. A., Suh, J., Pignatelli, M., Redondo, R. L., Ryan, T. J., & Tonegawa, S. (2013). Creating a False Memory in the Hippocampus. *Science*, *341*(6144), 387 391.

Ratigan, H. C., Krishnan, S., Smith, S., & Sheffield, M. E. J. (2023). A thalamic-hippocampal CA1 signal for contextual fear memory suppression, extinction, and discrimination. *Nature Communications*, *14*(1), 6758.

Rudy, J. W., Barrientos, R. M., & O'Reilly, R. C. (2002). Hippocampal formation supports conditioning to memory of a context. *Behavioral Neuroscience*, *116*(4), 530–538.

Russell, L. E., Dalgleish, H. W. P., Nutbrown, R., Gauld, O. M., Herrmann, D., Fişek, M., Packer, A. M., & Häusser, M. (2022). All-optical interrogation of neural circuits in behaving mice. *Nature Protocols*, *17*(7), 1579–1620.

Russo, A. S., & Parsons, R. G. (2021). Behavioral expression of contextual fear in male and female rats. *Frontiers in Behavioral Neuroscience*, *15*, 671017.

Schuette, P. J., Reis, F. M. C. V., Maesta-Pereira, S., Chakerian, M., Torossian, A., Blair, G. J., Wang, W., Blair, H. T., Fanselow, M. S., Kao, J. C., & Adhikari, A. (2020). Long-Term Characterization of Hippocampal Remapping during Contextual Fear Acquisition and Extinction. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *40*(43), 8329–8342.

Sheffield, M. E. J., Adoff, M. D., & Dombeck, D. A. (2017). Increased Prevalence of Calcium Transients across the Dendritic Arbor during Place Field Formation. *Neuron*, *96*(2), 490 504.e5.

Sheffield, M. E. J., & Dombeck, D. A. (2015). Calcium transient prevalence across the dendritic arbour predicts place field properties. *Nature*, *517*(7533), 200 204.

Shionoya, K., Hegoburu, C., Brown, B. L., Sullivan, R. M., Doyère, V., & Mouly, A.-M. (2013). It's time to fear! Interval timing in odor fear conditioning in rats. *Frontiers in Behavioral Neuroscience*, *7*, 128.

Stiedl, O., & Spiess, J. (1997). Effect of tone-dependent fear conditioning on heart rate and behavior of C57BL/6N mice. *Behavioral Neuroscience*, *111*(4), 703–711.

Suthard, R. L., Senne, R. A., Buzharsky, M. D., Diep, A. H., Pyo, A. Y., & Ramirez, S. (2024). Engram reactivation mimics cellular signatures of fear. *Cell Reports*, *43*(3), 113850.

Syeda, A., Zhong, L., Tung, R., Long, W., Pachitariu, M., & Stringer, C. (2024). Facemap: a framework for modeling neural activity based on orofacial tracking. *Nature Neuroscience*, *27*(1), 187–195.

Tonegawa, S., Morrissey, M. D., & Kitamura, T. (2018). The role of engram cells in the systems consolidation of memory. *Nature Reviews. Neuroscience*, *19*(8), 485 498.

Totty, M. S., Tuna, T., Ramanathan, K. R., Jin, J., Peters, S. E., & Maren, S. (2023). Thalamic nucleus reuniens coordinates prefrontal-hippocampal synchrony to suppress extinguished fear. *Nature Communications*, *14*(1), 6565.

Totty, M. S., Warren, N., Huddleston, I., Ramanathan, K. R., Ressler, R. L., Oleksiak, C. R., & Maren, S. (2021). Behavioral and brain mechanisms mediating conditioned flight behavior in rats. *Scientific Reports*, *11*(1), 8215.

Tovote, P., Fadok, J. P., & Lüthi, A. (2015). Neuronal circuits for fear and anxiety. *Nature Reviews. Neuroscience*, *16*(6), 317–331.

Trott, J. M., Hoffman, A. N., Zhuravka, I., & Fanselow, M. S. (2022). Conditional and unconditional components of aversively motivated freezing, flight and darting in mice. *ELife*, *11*. https://doi.org/10.7554/eLife.75663

Troyner, F., Bicca, M. A., & Bertoglio, L. J. (2018). Nucleus reuniens of the thalamus controls fear memory intensity, specificity and long-term maintenance during consolidation. *Hippocampus*, *28*(8), 602–616.

Uytiepo, M., Zhu, Y., Bushong, E., Polli, F., Chou, K., Zhao, E., Kim, C., Luu, D., Chang, L., Quach, T., Haberl, M., Patapoutian, L., Beutter, E., Zhang, W., Dong, B., McCue, E., Ellisman, M., & Maximov, A. (2024). Synaptic architecture of a memory engram in the mouse hippocampus. *BioRxiv : The Preprint Server for Biology*. https://doi.org/10.1101/2024.04.23.590812

Vetere, G., Kenney, J. W., Tran, L. M., Xia, F., Steadman, P. E., Parkinson, J., Josselyn, S. A., & Frankland, P. W. (2017). Chemogenetic Interrogation of a Brain-wide Fear Memory Network in Mice. *Neuron*, *94*(2), 363-374.e4.

Wang, M. E., Wann, E. G., Yuan, R. K., Álvarez, M. M. R., Stead, S. M., & Muzzio, I. A. (2012). Long-term stabilization of place cell remapping produced by a fearful experience. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *32*(45), 15802 15814.

Wang, M. E., Yuan, R. K., Keinath, A. T., Ramos Álvarez, M. M., & Muzzio, I. A. (2015). Extinction of Learned Fear Induces Hippocampal Place Cell Remapping. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *35*(24), 9122–9136.

Williams, D. A. (1994). Slow extinction of conditioned responding following exposure to two bouts of massed shock. *Animal Learning & Behavior*, *22*(2), 203–213.

Wiltgen, B. J., & Silva, A. J. (2007). Memory for context becomes less specific with time. *Learning & Memory (Cold Spring Harbor, N.Y.)*, *14*(4), 313–317.

Wiltschko, A. B., Tsukahara, T., Zeine, A., Anyoha, R., Gillis, W. F., Markowitz, J. E., Peterson, R. E., Katon, J., Johnson, M. J., & Datta, S. R. (2020). Revealing the structure of pharmacobehavioral space through motion sequencing. *Nature Neuroscience*, *23*(11), 1433–1443.

Wu, C.-T., Haggerty, D., Kemere, C., & Ji, D. (2017). Hippocampal awake replay in fear memory retrieval. *Nature Publishing Group*, *20*(4), 571 580.

Xu, W., & Südhof, T. C. (2013). A neural circuit for memory specificity and generalization. *Science*. https://science.sciencemag.org/content/339/6125/1290.short?casa_token=QtgiZ VpO0vgAAAAA:1evZSg4yR3MKpHHsrcyiJs135dikEsxBXNlOZAipkPQiXbkaIkho kkaVVO3qDwmtWTIkwSzbdcZtbbE