A contextual fear conditioning paradigm in head-fixed mice exploring virtual reality.

Seetha Krishnan[1*], Can Dong[1,2*], Heather Ratigan[1], Denisse Morales-Rodriguez[1,3], Chery Cherian[1], Mark Sheffield[1]

[1] Department of Neurobiology and Institute for Neuroscience, University of Chicago, Chicago, IL 60637, USA.

[2] Present address: Department of Neurobiology, Stanford University

[3] Present address: University of California, San Francisco

* equal contribution

Correspondence to: seethakrishnan@uchicago.edu, sheffield@uchicago.edu

ABSTRACT

Contextual fear conditioning is a classical laboratory task that tests associative memory formation and recall. Techniques such as multi-photon microscopy and holographic stimulation offer tremendous opportunities to understand the neural underpinnings of these memories. However, these techniques generally require animals to be head-fixed. There are few paradigms that test contextual fear conditioning in head-fixed mice, and none where the behavioral outcome following fear conditioning is freezing, the most common measure of fear in freely moving animals. To address this gap, we developed a contextual fear conditioning paradigm in head-fixed mice using virtual reality (VR) environments. We designed an apparatus to deliver tail shocks (unconditioned stimulus, US) while mice navigated a VR environment (conditioned stimulus, CS). The acquisition of contextual fear was tested when the mice were reintroduced to the shock-paired VR environment the following day. We tested three different variations of this paradigm and, in all of them, observed an increased conditioned fear response characterized by increased freezing behavior. This was especially prominent during the first trial in the

shock-paired VR environment, compared to a neutral environment where the mice received no shocks. Our results demonstrate that head-fixed mice can be fear conditioned in VR, discriminate between a feared and neutral VR context, and display freezing as a conditioned response, similar to freely behaving animals. Furthermore, using a two-photon microscope, we imaged from large populations of hippocampal CA1 neurons before, during, and following contextual fear conditioning. Our findings reconfirmed those from the literature on freely moving animals, showing that CA1 place cells undergo remapping and show narrower place fields following fear conditioning. Our approach offers new opportunities to study the neural mechanisms underlying the formation, recall, and extinction of contextual fear memories. As the head-fixed preparation is compatible with multi-photon microscopy and holographic stimulation, it enables long-term tracking and manipulation of cells throughout distinct memory stages and provides subcellular resolution for investigating axonal, dendritic, and synaptic dynamics in real-time.

INTRODUCTION

Large-scale multi-photon imaging and holographic stimulation offers the ability to accurately track and record from the same neuronal populations over extended periods at high spatiotemporal resolution, record from axons, dendrites, and dendritic spines, and stimulate neurons with single-cell precision (Bando et al., 2021; Chaigneau et al., 2016; Dombeck et al., 2010; Dong et al., 2021; Emiliani et al., 2015; Fan et al., 2023; Hattori & Komiyama, 2022; Heer & Sheffield, 2024; Hernandez et al., 2016; Krishnan et al., 2022; Russell et al., 2022; Sheffield & Dombeck, 2015). These techniques facilitate the study of neural circuit dynamics, synaptic and plasticity mechanisms and enable the real-time manipulation of neural activity, all of which are crucial for understanding the mechanisms underlying memory formation and recall in the brain. However, head-fixation of animals, which is generally required to utilize these techniques, imposes constraints on the range of testable behaviors compared to freely moving conditions (Guo et al., 2014; Juczewski et al., 2020). This limitation is particularly evident in the Pavlovian contextual fear conditioning (CFC) task, a laboratory method for testing associative learning, fear memory formation, and recall (M. E. Bouton, 1993; Michael S. Fanselow, 1980, 1990; J. J. Kim & Fanselow, 1992; LeDoux, 2000; Maren, 2001; Maren et al., 2013).

In the traditional CFC paradigm for freely moving animals, an animal is placed in an environment (conditioned stimulus, CS) that is paired with an aversive stimulus, such as a mild foot shock (unconditioned stimulus, US). Animals are then removed from the environment. When reintroduced to the environment, animals exhibit a species-specific conditioned response, such as fearful freezing behavior in rodents, if they successfully associate the shock-paired environment with the aversive stimulus. The CFC paradigm

is one of the most basic conditioning procedures for freely moving animals— it's simple

and robust (Michael S. Fanselow, 1980, 1990). Despite its longstanding use in studies

involving freely behaving animals to explore questions on learning and memory

(Anagnostaras et al., 2001; M. E. Bouton & King, 1983; Mark E. Bouton & Bolles, 1980;

Debiec et al., 2002; M. S. Fanselow & Bolles, 1979; Fendt & Fanselow, 1999; Gewirtz et

al., 2000; Holland & Bouton, 1999; Jeansok J. Kim & Jung, 2006; Maren et al., 2013;

Stephen Maren & Holt, 2000; Pezze & Feldon, 2004), it has been noticeably absent in

studies involving head-fixed animals, with some exceptions.

Two prior studies (Lovett-Barron et al., 2014; Rajasethupathy et al., 2015) reported a

version of CFC in head-fixed mice with aversive air puffs as the US. These studies

reported the conditioned response animals displayed as lick suppression, not freezing.

Although lick suppression after fear conditioning has also been observed in freely moving

animals (Leaf & Leaf, 1966; Leaf & Muller, 1965), it has been shown to result from

increased freezing (Mark E. Bouton & Bolles, 1980). Therefore, relying solely on lick

suppression in the head-fixed preparation makes it more challenging to compare with the

extensive previous work in freely moving animals, where freezing is measured as the

predominant conditioned fear response. Furthermore, the aforementioned head-fixed

CFC studies also had the limitation of simultaneously having both a water reward and an

aversive stimulus within the context, potentially confounding the results and complicating

the interpretation. Hence, we aimed to develop a paradigm where freezing is the

conditioned response and without the presence of a reward within the conditioned

context, making it directly comparable to freely moving animals.

Here, we describe a contextual fear-conditioning paradigm for head-fixed mice that

resulted in freezing as the conditioned response. The context or CS was a virtual reality

(VR)-based environment, which head-fixed mice explored by running on a spherical

treadmill. We previously demonstrated the feasibility of this approach (Ratigan et al.,

2023) and now provide a detailed overview and in-depth analysis of the behavior. Our

main aim is to show that VR-based CFC can elicit freezing responses similar to those

seen in freely moving animals, while also examining how key parameters affect this

response. Though we explored some important variables, this paper does not attempt to

comprehensively investigate all parameters necessary for successful VR-based CFC.

Additionally, we performed two-photon imaging of a large population of hippocampal CA1

cells during fear conditioning and tracked the same cells during memory recall across

days. We analyzed the impact of our paradigm on place cells in the hippocampus and

compared these findings to results from CFC in freely moving animals.

RESULTS

A paradigm for contextual fear conditioning in head-fixed mice exploring virtual

reality environments

To establish the contextual fear conditioning protocol (**Figure 1A-C**), head-fixed mice

were water-restricted and trained to run on a treadmill and navigate VR environments for

water rewards, as we have demonstrated previously (Dong et al., 2021; Krishnan et al.,

2022; Krishnan & Sheffield, 2023; Ratigan et al., 2023). Forward movement on the

treadmill translates to forward movement in VR, allowing mice to navigate the VR

environment. The environments used were 2m-long linear tracks rich in visual cues

(**Figure 1**). Reaching the track's end triggered a water reward and defined a completed

trial/lap. Mice were then virtually teleported back to the track's start for the next lap. Unlike

freely moving mice that tend to explore new environments, head-fixed mice placed on a

treadmill do not automatically start exploring VR environments. However, they can be

trained through reinforcement learning using water restriction and water rewards

(Dombeck et al., 2010; Dong et al., 2021; Krishnan et al., 2022; Ratigan et al., 2023;

Sheffield & Dombeck, 2015).

Training to navigate VR environments on a treadmill (Stage 1), using water rewards, was

achieved in approximately 10-14 days, after which well-trained mice could run several

laps per minute. We set a criterion of >3 laps per minute (i.e., a lap velocity of 0.1 - 0.2

m/s) for mice to move on to the next stage (Stage 2). Once the mice reached this stage

(Stage 2), the following day, we added a coat to their tails (**Figure 1B**). We developed a

lightweight, wearable conductive apparatus for the mouse's tail to ensure minimal

discomfort for the mouse. We named this apparatus the "tail-coat" (**Figure 1D**). This tail-

coat enabled us to administer mild electric shocks to the mouse's tail (see Methods).

When the tail-coat was added, some mice reduced their running speed or refused to

move. Some recovered the next day, but if their running speed did not reach at least three

laps per minute (~20% of mice failed to meet this criterion), they were not advanced to

the next stage. Mice that maintained this running speed advanced to Stage 3. Data from

mice that advanced with minimal changes to their lap running behavior after adding the

tail-coat are shown in **Supplementary Figure 1A**.

The following day (Stage 3), mice were placed in the same VR environment with the tail-

coat, but the water reward was removed (**Figure 1B**). Mice that continued to run at least

three laps per minute with the tail-coat and without water reward advanced to the

experiment stage (Stage 4). Around 40% of the mice dropped out at this stage. High levels

of lap running during training were crucial to ensure mice were undeterred by both the

tail-coat and the reward removal (Krishnan et al., 2022; Krishnan & Sheffield, 2023). Our

criteria for consistent running behavior with minimal pauses throughout these three

stages of training ensured that any baseline freezing behavior was minimal, similar to

freely moving animals, and allowed us to attribute freezing after CFC to fear conditioning.

In summary, our criteria for selecting mice for the experimental stage of contextual fear

conditioning required them to run consistently (>3 laps per minute) with 1) rewards, then

2) with a tail-coat, and then 3) without rewards plus the tail-coat. This threshold resulted

in about 40% of the mice proceeding to the experimental stage. Once mice qualified for

the experimental stage, they were no longer excluded from analysis due to their behavior.

We tested three variations of the head-fixed CFC paradigm. In Paradigm 1, the neutral

environment was the training VR environment, and a new VR environment was paired

with shocks. In Paradigms 2 and 3, both the neutral and shock-paired environments were

new and distinct from the training environment to better control for environmental

familiarity. In Paradigms 1 and 2, the tail-coat was removed on the days following CFC,

whereas in Paradigm 3, it remained on the mouse on all post-CFC days. With these

paradigms, we varied some parameters that may influence fear memory recall, contextual

discrimination, fear extinction, and their associated behaviors in head-fixed VR-based

CFC paradigms. We detail these variations in the following sections, starting with

Paradigm 1.

Figure 1. A contextual fear conditioning paradigm for head-fixed mice navigating virtual reality environments. (A) Experimental setup created with BioRender.com. Mice were head-restrained with their feet resting on a spherical treadmill. Five large monitors surrounded the mice that displayed virtual reality (VR) environments. Movement on the treadmill advanced the VR display, allowing for context exploration. **(B)** Mice were water-restricted and trained to run laps in the VR for water rewards. VR environments were 2m-long linear tracks. Mice were trained to achieve >3 laps per minute, which took ~10 to 14 days (Stage 1). Once well-trained, a "tail-coat" was added to their tails (Stage 2), followed by the removal of the water reward the next day (Stage 3). **(C)** Once the water reward was removed, mice underwent the fear conditioning protocol the following day. On the first experiment day (Day 0), mice spent 10 minutes in the training VR (Familiar VR) and then another 10 minutes in a new VR (CFC VR). After the initial exploration, mice received mild electric shocks on the tail (4-12 shocks, 0.5-1.2mA in amplitude, 1s long). The next day (Day 1), mice were tested for memory recall by placing them in the Familiar VR and the CFC VR for 5 minutes each in a counterbalanced manner. **(D)** Schematic of the tail-coat used for delivering mild electric shocks to the mouse's tail. (Top) View from the top (Bottom) side view. Dimensions are provided for a typical 12-week-old male mouse weighing ~30g before water restriction (see Methods for more details).

<u>When tested for contextual fear memory recall, mice displayed increased freezing</u>

<u>behavior in the shocked-paired VR environment.</u>

On Day 0 of the experimental stage in Paradigm 1 (CFC day), mice (n = 18) were exposed

to the same VR environment we used for training, referred to as the Familiar VR from

here onwards (**Figure 1C**), for 10 minutes. We then switched them to a new VR

environment (CFC VR). Following baseline exploration of the CFC VR for 10 minutes, we

repeatedly applied mild-electric shocks (4-12 shocks, 0.5-1mA in amplitude, 1s long) to

the mouse's tail. We call the period when shocks were delivered "during CFC". Shocks

were pseudo-randomly administered with respect to the animal's position in the CFC VR,

with an inter-shock interval (ISI) of 1 minute. To ensure that the shocks were associated

with the context and not a specific position in the environment, no visual cues were

explicitly paired with the shock. We quantified the animal's response to the shocks and

observed an immediate increase in running speed following each shock (**Supplementary**

Figure 1B). This was consistent across all mice and shocks and was a reliable indicator

that the apparatus was working correctly. Sixty seconds after the last shock was

administered, the VR screens were switched off, the tail-coat was removed, and mice

were returned to their home cages. The next day (Fear memory recall, Day 1), we tested

for recall of the learned association between the VR environment and the fearful stimulus.

Mice were exposed to both VRs (familiar VR and CFC VR) for five minutes each in a

counterbalanced manner. Water reward remained absent on recall days; however, in this

paradigm, the tail-coat was no longer present.

Before fear conditioning, mice exhibited similar behavior in both the Familiar and CFC

VRs. We evaluated the time taken to complete a lap and the freezing behavior (defined

as time points when the animal's instantaneous velocity dropped below 1cm/s). Neither

metric revealed a significant difference between the two VRs before shock administration

(**Figure 2A**, **Supplementary Figure 1D-E**). However, after CFC, when the animals were

reintroduced to the VRs on recall Day 1, we observed a significant increase in freezing

and time taken to complete a lap in the CFC VR compared to the Familiar VR (**Figure**

2A-D). As soon as mice transitioned to the CFC VR (either from dark or after the Familiar

VR) on the recall day, we observed behaviors such as slowing down, freezing, moving

backward, and hesitation to move forward (**Supplementary Videos 1-2**). This response

was markedly different from their behavior in the Familiar VR. This difference in behavior

can be observed clearly in most mice on the very first lap (**Figure 2A-B**). Since we

removed the reward, we didn't find licking behavior to be indicative of recall behavior

(**Supplementary Figure 1C**). However, some mice displayed licking behavior in the

Familiar VR on Day 0 before fear conditioning, which was absent on Day 1. This is likely

because mice had previously received a water reward in the Familiar VR. We will address

this caveat in Paradigms 2 and 3.

When we quantified the freezing behavior in mice, we found it to be, on average, high

throughout the five minutes that mice spent in the CFC VR on recall Day 1 (**Figure 2C-**

D, average freezing (%), mean $\pm$ 95% Confidence Intervals (CI), Familiar VR: 28.95 $\pm$

5.47, CFC VR: 44.13 $\pm$ 7.75, p=0.001, Paired *t*-test). We also observed a significant

increase in freezing during the first lap in the CFC VR compared to the familiar VR (**Figure**

2B-C, average first lap freezing (%), Familiar VR: 24.13 $\pm$ 5.59, CFC VR: 43.71 $\pm$ 6.49,

p<0.001, Paired *t*-test). This freezing behavior also translated to mice taking longer to

complete a lap in the CFC VR compared to the Familiar VR, both in the first lap (average

time taken to complete first lap (s), Familiar VR: 14.90 ± 5.27, CFC VR: 42.78 ± 12.17,

p<0.001, Paired *t*-test) and on average (**Figure 2E-F, the** average time taken to complete

all laps (s), Familiar VR: 17.58 ± 4.18, CFC VR: 42.92 ± 13.96, p=0.001, Paired *t*-test).

The increase in freezing and time taken to complete a lap in the CFC VR compared to

the Familiar VR was significantly higher on Day 1 than Day 0, indicating that it results

from fear conditioning (**Figure 2C-F**, p<0.05, Paired *t*-test).

Based on studies in freely moving animals (Michael S. Fanselow, 1980, 1990;

Ramanathan et al., 2018; Russo & Parsons, 2021; Trott et al., 2022), we settled on an

intensity of 1 mA and six shocks as a reasonable protocol, which we tested on a sizeable

cohort of mice (n = 12). We found that freezing at this intensity and shock number was

higher on average in the CFC VR; however, mice showed variability in their responses

(**Supplementary Figure 2A-B**). The majority of mice (7/12) displayed appropriate

memory recall (**Supplementary Figure 2A-B)**, indicated by more freezing in the CFC VR

compared to the Familiar VR (delta > 0), while 3/12 mice froze more in the Familiar VR

(delta < 0), and 2/12 showed equal freezing levels in both VRs (delta ≈ 0). Such variability

has been regularly documented in freely moving animals and may not be unique to the

VR setup used here (Chu et al., 2024; Dos Santos Corrêa et al., 2019; Navarro-Sánchez

et al., 2024; Poulos et al., 2016; Russo & Parsons, 2021; Totty et al., 2021). Next, in a

small group of mice (n = 6), we investigated whether changing the number of shocks or

the shock amplitude would affect the conditioned fear response (**Supplementary Figure**

2A-B). We found that most shock amplitudes and numbers of shocks increased freezing

behavior on average in the CFC VR on recall day (**Supplementary Figure 2A-B**). The

observed variability was within the same range as that of the larger cohort of mice we

tested using the 1 mA, 6-shock protocol (**Supplementary Figure 2A-B**). A more detailed

analysis with a larger sample size may be necessary to understand further how variations

in shock parameters affect the conditioned response, as has been done in studies with

freely moving animals (Baldi et al., 2004; Cordero et al., 1998; Daviu et al., 2024; Poulos

et al., 2016; Quinn et al., 2008).

Finally, we examined the extinction of this conditioned response (**Supplementary Figure**

2C-D). We placed the animals in both the Familiar and the CFC VR for five minutes over

four days (Recall Days 1-4) in a counterbalanced manner and tested their recall behavior.

Mice displayed the most substantial increase in freezing in the CFC VR on the first day

of recall (**Supplementary Figure 2C-D**, Average freezing (%): Recall Day 1: Familiar VR:

28.95 ± 5.47, CFC VR: 44.13 ± 7.75). Freezing on the first lap in the CFC VR was also

most pronounced on Recall Day 1 (First lap freezing (%): Recall Day 1: Familiar VR: 24.13

± 5.59, CFC VR: 43.71 ± 6.49). Both average (Familiar VR: 27.78 ± 7.86, CFC VR: 33.19

± 10.22) and first lap freezing (Familiar VR: 20.30 ± 6.24, CFC VR: 25.56 ± 8.03) in the

CFC VR decreased on the second day of recall compared to the first recall day, becoming

similar to the Familiar VR. While first lap freezing remained similar between the Familiar

and the CFC VR on subsequent Recall Days 3 and 4, we observed variability in the

average freezing behavior. Average freezing increased significantly again on Recall Day

3 in the CFC VR (Average freezing (%): Recall Day 3: Familiar VR: 29.05 ± 13.32, CFC

VR: 43.89 ± 13.10) and was high in both VRs on Recall Day 4 (Recall Day 4: Familiar

VR: 37.66 ± 15.77, CFC VR: 38.04 ± 13.42).

To quantify freezing levels more precisely, baseline freezing on Day 0 was compared to

recall-day freezing using a Linear Mixed Effects (LME) Model, with recall days as fixed

effects and mouse as a random effect (**Tables 1-2**). In the CFC VR, mice froze

significantly more on Recall Days 1, 3, and 4 compared to Day 0. By Recall Day 4,

freezing levels were elevated in both the Familiar and CFC VRs relative to baseline (**Table**

2). On the other hand, first-lap freezing showed a significant increase only on Recall Day

1 in the CFC VR and remained similar to Day 0 on Recall Days 2-4 in both VRs (**Table**

1). This pattern suggests that freezing in the CFC VR peaks on the first recall day, with a

strong conditioned fear response to the shock-paired VR observed on Recall Day 1. The

marked difference in freezing behavior between the CFC VR and Familiar VR during the

first lap diminishes after the first recall day. On the other hand, average freezing behavior

becomes more variable in subsequent recall days, suggesting that complete fear

extinction in Paradigm 1 may take longer than four days. The increase in freezing on

Recall Day 4 in the Familiar VR may reflect either fear generalization or a response to the

extended lack of reward in a previously rewarded environment. Further research is

needed to determine the exact conditions required for complete fear extinction in this

paradigm.

In summary, these results demonstrate that head-fixed mice can acquire fear associations

in VR environments. Most mice displayed freezing as a conditioned response specifically

to the shock-paired VR—behavior that mirrors that of freely moving animals after fear

conditioning. However, complete fear extinction in this paradigm may require more than

four days of testing.

Figure 2. Head-fixed mice show increased freezing behavior following contextual fear conditioning in the CFC VR. (A) A single example mouse's lap running behavior on experiment days, Day 0 and Day 1. Behavior is shown for approximately 3 minutes in all sessions except during CFC, which is shown for the 6 minutes that the session lasted. Frames where freezing was detected (instantaneous velocity <1cm/s) are marked with black dots. The traces on the right show that this mouse increased freezing, decreased velocity, and moved backward in the CFC VR (red traces) but not in the Familiar VR (blue traces) on Recall Day. This mouse received six shocks at 1mA intensity at 60 s inter-stimulus interval (ISI). **(B)** First two minutes of recall behavior in more mice (n = 5) in Familiar VR versus CFC VR. **(C-D)** (Left Average freezing percentage on recall day in the very first lap **(C)** and all laps **(D)** during the 5 minutes that mice explored the Familiar (blue) and CFC (red) VR. Freezing (%) was calculated as the number of frames where freezing was detected in a lap divided by the total number of frames in each lap. (Right) Delta calculated as the difference in the amount of freezing in the CFC VR compared to the Familiar VR before CFC (Day 0) and after CFC (Day 1). The dashed line represents 0. **(E-F)** Same as C-D but for the time taken to complete the first lap **(E)** and all laps **(F).** Mice displayed an increase in freezing and in time taken to complete a lap in the first lap and, on average, in the CFC VR. In C-F, circles and pluses represent individual mice (n = 18, 16 male and two female mice). In C-F, data was pooled from mice receiving different numbers of shocks (4, 6, 12) at varying intensities (0.5 mA, 0.8 mA, 1 mA, and 1.2 mA), which is separately displayed in

261 **Supplementary Figure 2**. Lines join data from the same mouse. P-values were calculated using a *paired*
262 *t-test.*

Day 0: Contextual Fear Conditioning

263
Supplementary Figure 1. Running and licking behaviors in Paradigm 1. (A) Box plot of average time
taken to complete a lap before and after the addition of a tail-coat recorded in a randomly chosen subset
of mice (circles, n=8). **(B)** Instantaneous velocity increased immediately after shock onset. This increased
running speed also served as a reliable measure to confirm that mice received the tail shock. The grey line
indicates the average velocity calculated by shuffling the instantaneous velocity 100 times. **(C)** Licking
behavior doesn't stop immediately in the familiar VR without water, as mice previously received water
rewards in the familiar VR. However, it was low in the other sessions and across environments. There was
no significant difference between the Familiar and CFC VR on Day 1. Licking behavior was collected only
in a subset of mice (circle, n=5). **(D-E)** The amount of freezing **(D)** and time taken to complete a lap **(E)**
were not significantly different between the Familiar and CFC VR before fear conditioning (n=18). While not
significantly different, there was a trend towards longer time taken in the first lap of the CFC VR, which we
have observed before when mice enter novel environments from familiar ones (Dong et al., 2021). The
boxplots (in A, C-E) range from the first quartile (25th percentile) to the third quartile (75th percentile), and
the box shows the interquartile range (IQR). The line across the box represents the median (50th
percentile). The whiskers extend to 1.5*IQR on either side of the box, and anything above this range is
defined as an outlier. P-values were calculated using a paired *t-test.*

Supplementary Figure 2. Change in freezing with shock amplitude, number of shocks, and across recall days. (A-B) The left panel shows the average freezing in the first lap **(A)** and all laps **(B).** The right panel illustrates delta freezing, calculated as the difference in freezing between the CFC VR and the Familiar VR. The x-axis on the right panel is meaningless. A random jitter parameter separates the data points in the x-axis to prevent them from overlapping. The dashed line represents the average freezing percentage in a control paradigm, where mice experienced the same conditions without shocks. The largest effect sizes were observed in the first lap when the mouse received 0.5 mA, 12 shocks (lime green), or 0.8 mA/1 mA, 6 shocks (orange, purple). Four shocks produced a small effect. Increasing the number of shocks (to 6 or 12) enhanced freezing in the CFC VR. However, increasing shock intensity only led to modest increases in average freezing, as seen with 1.2mA, six shocks (pink) compared to 1mA, six shocks (purple). Number of animals used is indicated in the figure legend. **(C-D)** A subset of animals underwent multiple days of recall tests (n = 18 Day 0, n = 18 Recall Day 1, n = 11 Recall Day 2, n = 8, Recall Day 3, n = 7 Recall Day 4). VR environments were presented in a counterbalanced manner for 5 minutes each day. The increase in freezing in the CFC VR was highest on the first day, both in the first lap **(C)** and across all laps **(D).** P-values were calculated using a paired *t-test.*

Table 1: LME summary (estimate ± standard error, p-value) for first lap freezing in Paradigm 1 with fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2	Recall Day 3	Recall Day 4
Familiar VR	-2.28 ± 4.69, p=0.62	-6.36 ± 5.50, p=0.25	-2.92 ± 6.13, p=0.63	7.76 ± 6.41, p=0.23
CFC VR	24.34 ± 3.94, p<0.001	5.88 ± 4.62, p=0.21	5.50 ± 5.16, p=0.29	5.39 ± 5.40, p=0.32

VR	Recall Day 1	Recall Day 2	Recall Day 3	Recall Day 4
Familiar VR	2.35 ± 2.82, p=0.41	1.14 ± 3.40, p=0.74	3.05 ± 3.83, p=0.43	13.29 ± 3.98, p=0.0019
CFC VR	19.5 ± 3.63, p<0.001	8.44 ± 4.33, p=0.06	18.29 ± 4.86, p=0.0006	12.85 ± 5.07, p=0.016

A second paradigm where head-fixed mice discriminate between two novel VRs.

As previously mentioned, the incomplete extinction of licking behavior in the familiar VR before fear conditioning or the gradual increases in freezing behavior in the familiar VR during recall tests may indicate that mice still associate the VR with water rewards, which could be a caveat. Furthermore, in traditional CFC paradigms with freely moving mice, fear behavior has been assessed mainly by comparing two novel contexts, one associated with shocks and the other acting as a control. To address this, we tested another paradigm (Paradigm 2), using a novel VR instead of the familiar VR to act as the control (**Figure 3A**). One novel VR was assigned as the Control VR, and a second novel VR was assigned as the CFC VR, which would later become paired with the shocks. In this paradigm, we also added an extra habituation day to increase the animal's pre-exposure time to the novel VRs prior to conditioning (Day -1). Pre-exposure to the context before fear conditioning can enhance contextual fear, as seen in freely moving animals (Biedenkapp & Rudy, 2007; Michael S. Fanselow, 1980, 1990; Maren et al., 2013; Rudy et al., 2002; Wiltgen & Silva, 2007).

The training was done similarly to the previous paradigm. After mice showed consistent running behavior in the Familiar VR with the tail-coat and without reward, they were advanced to the experimental stage. On Day -1, mice were exposed to the two novel VRs for ten minutes each. On Day 0, they were re-exposed to the same two VRs for ten

minutes before receiving six one-second-long mild electric shocks (at 1 mA and 1 minute

ISI) in one of them - the CFC VR. On Day 1, mice were exposed to the two VRs for five

minutes to evaluate memory recall (**Figure 3A-B**).

Compared to Paradigm 1, in the novel VRs, mice showed no differences in licking

behavior across all experimental sessions (**Supplementary Figure 3A**). There was also

no significant difference in freezing behavior and time taken to complete a lap between

the two VRs on Day -1 and Day 0 before fear conditioning (**Supplementary Figure 3B-**

C).

Head-fixed mice show heightened fear discrimination during the first lap when

discriminating between two novel VRs.

We compared the freezing behavior between Paradigm 1 and Paradigm 2 on Recall Day

1 to the same intensity and number of shocks (1 mA, 1 s long, 60 s ISI, six shocks, n=12

mice each paradigm). We observed only modest increases in average freezing behavior

and time taken to complete a lap in the CFC VR in Paradigm 2, compared to Paradigm 1

(**Figure 3C-D, 4A-E**). The effect sizes for the difference in freezing behavior between

CFC VR and Control VR in Paradigm 2 were smaller than those between Familiar VR

and CFC VR in Paradigm 1 (Cohen's d: First lap freezing: Paradigm 1: 1.377, Paradigm

2: 0.917. Average Freezing: Paradigm 1: 0.893, Paradigm 2: 0.076). When we quantified

freezing by lap on Recall Day 1 in Paradigm 2, we found that after the first lap, mice

showed similar freezing in both the Control VR and the CFC VR across all subsequent

laps (**Supplementary Figure 4**). In contrast, in Paradigm 1, mice consistently froze more

in the CFC VR compared to the Familiar VR throughout Recall Day 1 (**Supplementary**

Figure 4).

Indeed, when we quantified freezing behavior in the Control VR and CFC VR in Paradigm

2, we found that the most significant increase in freezing in the CFC VR was observed in

the first lap (freezing in first lap %: Control VR: 16.50 ± 3.15 CFC VR: 23.19 ± 5.75, p =

0.026, time taken to complete first lap (s): Control VR: 14.27 ± 2.85, CFC VR: 25.68 ±

5.55, p = 0.001, Paired *t*-test, **Figure 3, Figure 4A-E**). Neither average freezing (**Figure**

4C, average freezing in all laps %: Control VR: 30.32 ± 6.25, CFC VR: 31.06 ± 6.07,

p=0.792, Paired *t*-test) nor time taken to complete all laps (**Figure 4E,** the average time

taken to complete all laps: Control VR: 21.73 ± 8.11, CFC VR: 24.27 ± 6.36, p=0.558,

Paired *t*-test) showed significant differences between the two VRs.

When mice were returned to the two VRs the following day for a second day of recall

(Recall Day 2), we found no significant difference in first lap freezing between the Control

and CFC VR (**Supplementary Figure 5,** First lap freezing (%) Recall Day 2: Control VR:

13.96 ± 4.24, CFC VR: 21.34 ± 6.79, p = 0.083, Paired *t*-test). Average freezing in both

CFC VR and Control VR returned to baseline levels on Recall Day 1 and remained so on

Recall Day 2 (**Supplementary Figure 5, Tables 3-4,** Average freezing (%) Recall Day 1:

Control VR: 30.32 ± 6.25, CFC VR: 31.06 ± 6.07, p = 0.792, Recall Day 2: Control VR:

33.35 ± 6.11, CFC VR: 35.88 ± 5.89, p = 0.045, Paired *t*-test). This indicates that fear

extinction in Paradigm 2 happens quickly, within the first recall session after the first lap.

Figure 3. Using a novel VR as the neutral environment instead of a familiar VR in a second paradigm results in modest increases in freezing. (A) The training paradigm is similar to paradigm 1. On experiment day, mice were introduced to two novel VRs, one of which would be associated with the shock (CFC VR) and the other wouldn't (Control VR). In this paradigm, there was an added habituation day, where mice were exposed to the two VRs for ten minutes. The next day, mice ran in the two VRs again before receiving mild electric shocks in the CFC VR. The recall test occurred the next day. In this cohort, all mice received six shocks at 1 mA intensity. **(B-D)** These panels show the lap running behavior of a single mouse on experiment days 1, 2, and 3. Behavior is displayed for about 3 minutes in all sessions except during

CFC. Frames with freezing detected by a threshold (instantaneous velocity <1cm/s) are marked by black dots. In **(D),** this mouse shows an increase in freezing, a decrease in velocity, and some backward movement when in the CFC VR (red traces). **(E)** Comparison between freezing in the first lap (left) and all laps (right) in paradigm 1 (blue and red) versus paradigm 2 (green and red). In paradigm 1, only animals that received six shocks at 1 mA intensity are included. n = 12 mice were used in both paradigms (10 male and two female mice). The scale bar for panels **B-D** is indicated in **B**. P-values were calculated using a paired *t-test.*

Figure 4. In Paradigm 2, the most significant freezing and reduced speed occurred in the first lap of the CFC VR compared to the Control VR. (A) First two minutes of recall behavior in five mice in Control VR (green) versus CFC VR (red). **(B-C)** The left panels show the amount of freezing on the recall day in (B) the very first lap and (C) all laps during the 5 minutes when the mice explored the Control (green) and CFC (red) VR. Freezing (%) is calculated as the number of frames where freezing was detected in a lap divided by the total number of frames in each lap. On the right, the delta is calculated as the difference in the amount of freezing in the CFC VR compared to the Control VR before (Day 2) and after CFC (Day 3). The dashed line represents 0. **(D-E)** Same as B-C but for the time taken to complete the first lap **(D)** and all laps **(E).** In B-E, circles and pluses represent individual mice (n = 12). Lines join data from the same mouse. P-values were calculated using a *paired t-test.*

Supplementary Figure 3. Licking and running behavior before fear conditioning, in Paradigm 2. (A)
Licking behavior remains low in all sessions, unlike Paradigm 1. It does not significantly differ between the
Control vs CFC VR before and after CFC. **(B-C)** The amount of freezing and average running speed was
not significantly different between the Control and CFC VR before fear conditioning, both on Day 1 **(B)** and
Day 2 **(C)**. However, in this paradigm, some mice paused more on both days than in Paradigm 1. P-values
were calculated using a *paired t-test.*

396
Supplementary Figure 4. Lap-wise freezing behavior in the three paradigms. (A-F) Average freezing
across mice **(A-C)** and the number of freezing epochs **(D-F,** see Methods) in each lap across the two VRs
on the first day of recall in **(A, D)** Paradigm 1, **(B, E)** Paradigm 2 and **(C, F)** Paradigm 3. Lines indicate
mean, and shading indicates 95% confidence intervals. P-values between **(A, D)** Familiar VR and CFC
VR and **(B-C, E-F)** Control VR and CFC VR were calculated using a *paired t-test.* * indicates p-
values<0.01.

403
404 **Supplementary Figure 5. Mice extinguished their fear on the second recall day in Paradigm 2.** (A-
405 B) First lap freezing (A) and average freezing across all laps (B) across two days of recall in Paradigm 2.
406 P-values were calculated using a paired t-test. Circles indicate data from individual mice (n = 12 for Recall
407 Day 1 and n = 8 for Recall Day 2). Mice showed no significant difference in freezing behavior between the
408 two environments on Recall Day 2.

Table 3: LME summary (estimate ± standard error, p-value) for first lap freezing in Paradigm 2 with fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2
Control VR	-8.22 ± 4.48, p=0.08	-10.76 ± 5.01, p=0.05
CFC VR	6.82 ± 2.85, p=0.03	4.91 ± 3.24, p=0.15

Table 4: LME summary (estimate ± standard error, p-value) for average freezing in Paradigm 2 with fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2
Control VR	-3.99 ± 2.97, p=0.20	-1.47 ± 3.37, p=0.67
CFC VR	-0.02 ± 2.8, p=0.99	4.31 ± 3.17, p=0.19

A shorter ISI and leaving the tail-coat on during fear memory recall improved fear discrimination when using two novel VRs.

Freezing behavior in the CFC VR in Paradigm 2 was weaker than in Paradigm 1, with rapid fear extinction after the first recall lap. In Paradigm 3, two parameters were changed relative to Paradigm 2 to test whether freezing behavior and fear discrimination could be enhanced. First, the interval between consecutive shocks was reduced to see if a shorter ISI could produce a more robust fear response (McNally & Westbrook, 2006; Williams, 1994). Second, the tail-coat was kept on during memory recall, based on evidence that removing contextual cues can reduce conditioned freezing (González et al., 2003). We hypothesized that the tail-coat might act as a contextual cue and that its presence during recall might enhance freezing behavior.

In Paradigm 3, mice followed the same experimental protocol as Paradigm 2, with two novel VRs: one assigned as the Control VR and the other as the CFC VR. A habituation day in the two new VRs (Day -1) was followed by the CFC day (Day 0), where mice received six shocks (0.6 mA, 20-26 s ISI) in the CFC VR. This was followed by recall days

(Days 1-3), where mice explored the two VRs presented in a counter-balanced manner.

Unlike the other two paradigms, in Paradigm 3, the tail-coat remained on during the recall

days (**Figure 5A**).

We found that mice (n = 20) showed better fear discrimination between the Control and

CFC VRs in Paradigm 3 compared to Paradigm 2 (**Figure 5B-D**). Mice not only showed

more significant freezing in the first lap (freezing in first lap %: Control VR: 17.83 $\pm$ 2.95

CFC VR: 25.86 $\pm$ 6.51, p = 0.016, Paired *t*-test) but average freezing across all laps was

also higher in the CFC VR compared to the Control VR (**Figure 5C-D, Figure 6A-D,**

average freezing in all laps %: Control VR: 19.86 $\pm$ 2.87, CFC VR: 28.19 $\pm$ 3.66, p < 0.001,

Paired *t*-test).

While first-lap freezing showed a weaker effect size in Paradigm 3 compared to the other

paradigms, there was a strong effect of average freezing across the session (Cohen's d:

First lap freezing: Paradigm 1: 1.377, Paradigm 2: 0.917, Paradigm 3: 0.744. Average

Freezing: Paradigm 1: 0.893, Paradigm 2: 0.076, Paradigm 3: 1.185). On average, we

found that Paradigm 1 still produced the highest absolute freezing behavior in the CFC

VR (average freezing in all laps %: CFC VR: Paradigm 1: 44.13 $\pm$ 7.75, Paradigm 2: 31.06

$\pm$ 6.07, Paradigm 3: 28.19 $\pm$ 3.66). The difference in freezing behavior between the CFC

VR and the neutral VR was greatest in Paradigm 1 (Δ average freezing (CFC VR –

Familiar/Control VR): Paradigm 1: 13.14, Paradigm 2: 0.74, Paradigm 3: 8.33). However,

unlike Paradigm 2, where the largest differences in freezing between the CFC VR and

the Control VR were restricted to the first lap, the shorter ISIs and the addition of the tail-

coat during the recall day in Paradigm 3 improved contextual fear discrimination between

the two novel VRs. When tested for memory recall in Paradigm 3, mice froze more in the

CFC VR than in the Control VR, and this effect persisted beyond the first lap (**Figure 5C-**

D, Supplementary Figure 4).

Additionally, unlike the rapid fear extinction seen in Paradigm 2, mice showed a delay in

extinction in Paradigm 3 (**Table 5-6**, **Supplementary Figure 6**). Average freezing

continued to be higher in the CFC VR than in the Control VR on Recall Day 2 (Average

freezing (%) Recall Day 1: Control VR: 19.86 ± 2.87, CFC VR: 28.19 ± 3.66, $p < 0.001$,

Recall Day 2: Control VR: 21.79 ± 3.70, CFC VR: 26.46 ± 3.34, $p < 0.001$, Paired *t*-test).

By Recall Day 3, there was no significant difference in freezing between the Control VR

and the CFC VR (First lap freezing (%): Recall Day 3: Control VR: 13.10 ± 3.37, CFC VR:

17.37 ± 5.15, $p = 0.151$, Average freezing (%): Recall Day 3: Control VR: 19.43 ± 3.18,

CFC VR: 20.73 ± 3.61, $p = 0.240$, Paired *t*-test).

However, when comparing average freezing to baseline levels on Day 0 using an LME

model, a significant increase in average freezing was observed in both the Control VR

and CFC VR groups. This increase persisted across all three recall days (**Tables 5-6**).

Unlike Paradigm 1, in Paradigm 3, freezing levels showed a downward trend over time.

However, like Paradigm 1, they had not returned to baseline by the third recall day,

suggesting that fear was not entirely extinguished. The presence of the tail-coat during

recall days likely contributed to some degree of fear generalization, as mice showed

elevated freezing relative to baseline even in the Control VR. However, freezing levels in

the Control VR remained much lower than in the CFC VR on all days, indicating that mice

continued to discriminate between the VR environments (**Figure 6, Supplementary**

Figure 6, Tables 5-6).

There was no difference in first-lap freezing in the Control VR compared to baseline on

any of the recall days. First-lap freezing in the CFC VR was still higher than baseline on

Recall Day 2; however, it was back to baseline levels by Recall Day 3 (**Table 5-6**). As we

noticed in Paradigm 1, this indicates that pronounced increases in freezing in the CFC

VR in the very first lap are most prominent in the earlier days of recall. On the other hand,

the increases in average freezing continue to persist and may take longer to extinguish.

Thus, a shorter ISI and the presence of the tail-coat during memory recall may enhance

conditioned fear in head-fixed mice when discriminating between two novel VRs. The

influence of each parameter on its own will need further investigation.

In summary, our results show that mice exhibited greater average freezing behavior in

the CFC VR in Paradigms 1 and 3 compared to Paradigm 2. In Paradigm 2, the most

noticeable increase in freezing occurred during the very first lap in the CFC VR compared

to the Control VR. We found that fear discrimination could be improved with a shorter ISI

when presenting tail shocks and when the tail-coat was present during memory recall, as

shown in Paradigm 3. Furthermore, in Paradigms 1 and 3, the fear was not wholly

extinguished over the days we tested for recall and may take longer. However, fear

extinction was rapid in Paradigm 2 and was back to baseline levels within the first day of

recall. Thus, we provide three different variations for performing CFC in head-fixed mice

that can be used to understand the neural underpinnings of fear memory.

Figure 5. A third paradigm that uses a novel VR as the neutral environment but keeps the tail-coat on during memory recall led to increased freezing in the CFC Environment. (A) The training and experiment paradigm was similar to Paradigm 2 except that the shocks were administered closer together (20-25 s ISI), and the tail-coat was kept on during recall days. **(B)** First two minutes of recall behavior in Paradigm 3 in five mice in Control VR (green) versus CFC VR (red). **(C-D)** Comparison between freezing in the first lap **(C)** and all laps **(D)** in paradigm 1 (blue and red) versus paradigms 2 and 3 (green and red).

501 n = 12 mice were used in paradigms 1 and 2 (10 male and two female mice), and n = 20 mice in
502 paradigm 3. P-values were calculated using a *paired t-test*. In Paradigm 3, by keeping the tail-coat on
503 during the recall days, we observed an increase in freezing in the CFC VR in the first lap and across all
504 laps compared to Paradigm 2.

505
Figure 6. Mice froze more and reduced speed in Paradigm 3 across all laps on Recall Day 1; this
freezing behavior continued into the second day of recall in Paradigm 3. (A-B) The right panels show
the amount of freezing on the recall day in **(A)** the very first lap and **(B)** all laps during the 5 minutes when
the mice explored the Control (green) and CFC (red) VR. Freezing (%) is calculated as the number of
frames where freezing was detected in a lap divided by the total number of frames in each lap. On the left,
the delta is calculated as the difference in the amount of freezing in the CFC VR compared to the Control
VR before (Day 2) and after CFC (Day 3). The dashed line represents 0. **(C-D)** Same as A-B but for time
taken to complete the first lap **(C)** and all laps **(D).** Circles and pluses represent individual mice (n = 20).
Lines join data from the same mouse. P-values were calculated using a *paired t-test.*

515
Supplementary Figure 6. Mice show delay in extinction in Paradigm 3 (A-B) Freezing in the first lap
(A), and average freezing across all laps **(B)** in the Control VR vs CFC VR on three consecutive recall
days in Paradigm 3. The tail-coat was kept on all recall days. P-values were calculated using a *paired t-*
test.

Table 5: LME summary (estimate ± standard error, p-value) for first lap freezing in Paradigm 3 with
fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).

VR	Recall Day 1	Recall Day 2	Recall Day 3
Control VR	2.29 ± 2.51, p=0.37	3.96 ± 2.51, p=0.12	-2.43 ± 2.51, p=0.34
CFC VR	13.21 ± 3.33, p<0.001	9.79 ± 3.34, p=0.005	4.73 ± 3.34, p=0.16

522
523 **Table 6: LME summary (estimate ± standard error, p-value) for average freezing in Paradigm 3 with**
524 **fixed effects (recall days, baseline: before CFC, Day 0) and random effect (mouse).**

VR	Recall Day 1	Recall Day 2	Recall Day 3
Control VR	3.43 ± 1.28, p=0.01	5.35 ± 1.28, p=0.0001	2.99 ± 1.28, p=0.02
CFC VR	13.15 ± 1.78, p<0.001	11.42 ± 1.78, p<0.001	5.69 ± 1.78, p=0.0023

525
Place cells in the CA1 subregion of the hippocampus displayed remapping and

decreased field widths following contextual fear conditioning.

The hippocampus plays a critical role in the encoding, consolidation, and recall of

memories and is necessary for the expression of contextual fear memory (Anagnostaras

et al., 2001; Ji & Maren, 2007; Liu et al., 2012; Maren et al., 2013; Stephen Maren & Holt,

2000; Phillips & LeDoux, 1992; Ramirez et al., 2013; Wiltgen & Silva, 2007). Context is

thought to be represented in the hippocampus by place cells that fire at specific spatial

locations (place fields) in an environment (O'Keefe & Dostrovsky, 1971). These place

cells have been thought to help discriminate between feared and neutral contexts.

Previous studies on freely moving animals have shown that fear conditioning can cause

place cells to remap (Blair et al., 2023; E. J. Kim et al., 2015; Kinsky et al., 2023; Mamad

et al., 2019; Moita et al., 2004; Ormond et al., 2023; Schuette et al., 2020; Wang et al.,

2012, 2015; Wu et al., 2017). Remapping is defined by place cells shifting their firing

fields, indicative of place cells adapting to incorporate new information into memory

(Colgin et al., 2008; Leutgeb et al., 2005; Muller & Kubie, 1987). Additionally, a recent

study (Schuette et al., 2020) that performed calcium imaging of hippocampal cells in freely

moving mice found that fear conditioning narrows the width of place fields in the feared

environment. This suggests that place cells encode the same environment on a finer scale

following fear conditioning, potentially enhancing context discrimination and threat

avoidance (Schuette et al., 2020).

We replicated these findings in our head-fixed version of CFC in VR. To investigate the

effect of fear conditioning on place cell activity in our task, we expressed a calcium

indicator, GCaMP6f (Chen et al., 2013), in hippocampal pyramidal cells and, using two-

photon microscopy, imaged from the same hippocampal neurons across days during fear

conditioning and fear memory recall in Paradigm 1 (**Figure 7A-B**). We found that a subset

of place cells formed before CFC remained stable, while others remapped or changed

their preferred firing locations after CFC (see examples in **Figure 7C, 7D-G**). This is

similar to previously reported observations (Blair et al., 2023; Moita et al., 2004; Wang et

al., 2012, 2015).

We found remapping to occur in both the Familiar and the CFC VR to similar extents, as

indicated by the lower correlation across sessions/days compared to the within-session

correlation (**Figure 7F-G**). This contradicts previous findings by Moita et al. (2004), who

reported more stability in the control environment compared to the CFC environment after

fear conditioning. This remapping in both the Familiar and the CFC VR could be due to

differences in the CFC paradigm used in that study versus ours and will need further

investigation. It could also result from taking the water reward away from a previously

rewarded VR, as is the case in Paradigm 1. We have shown before that this makes the

place map more likely to drift across days (Krishnan et al., 2022; Krishnan & Sheffield,

2023). Whether these results hold the same for the Control VR in Paradigms 2 and 3 will

need further investigation.

Next, we looked at the place cells in each session. Interestingly, we found an increase in

the number of place cells in the CFC VR during memory recall, a finding not previously

reported (**Figure 8A-B**). We also found that place cells had narrower fields in the CFC

VR during memory recall (**Figure 8C**), as previously shown (Schuette et al., 2020). We

did not find a significant difference in other place cell parameters we examined, such as

lap-by-lap reliability (**Figure 8D**) and out-of-field firing (**Figure 8E**). Thus, our observations

of remapping and narrower place field widths replicate findings from freely moving

animals, validating our paradigm's use to explore further questions using the head-fixed

preparation to study the neural underpinnings of contextual fear memories. We add to

this field by showing an increase in the proportion of place cells following CFC, a possible

neural substrate of fear memory recall.

Figure 7. Fear conditioning results in place cell remapping in both Familiar and CFC VR. (A) A
schematic of the procedure for two-photon imaging of large populations of the same CA1 neurons over
multiple days, created with BioRender.com **(B)** An example field of view across the two experiment days.
The same field of view was aligned and imaged across days. **(C)** Examples of a few place cells in the CFC
VR across sessions on Day 0 (before CFC, during CFC) and Day 1 (Memory Recall). White lines separate
laps in each session. Some place cells maintain stable fields across days, while others remap. **(D)** Place

fields defined on Day 0 in Familiar VR before CFC are plotted across Day 1 during fear memory recall (n = 4 mice). **(E)** Place fields defined on Day 0 in CFC VR before CFC are plotted across the fear conditioning session and on Day 1 during fear memory recall. **(F)** On the left is a scatter plot of the center of mass of place fields defined in Familiar VR before CFC on Day 0 (x-axis) compared to their center of mass on Day 1 during fear memory recall (y-axis). On the right is a boxplot of correlation coefficients between mean place fields defined in Familiar VR before CFC on Day 0 and memory recall on Day 1 (Day 0: Day 1). The within-session correlation coefficients serve as control and were calculated between mean place fields in the first half and second half of the Familiar VR before the CFC session. **(G)** A scatter plot of the center of mass of place fields defined in CFC VR before CFC on Day 0 (x-axis) compared to their center of mass on (right) Day 0 during CFC and (middle) Day 1 during memory recall. On the left is a distribution of correlation coefficients between mean place fields defined in Familiar VR before CFC on Day 0 and during CFC (Day 0: Day 0) and memory recall on Day 1 (Day 0: Day 1). The within-session correlation was calculated between mean place fields in the first half and second half of the CFC VR before the fear conditioning session. Asterisk (*) indicates significant *P* values (KS test, *P* < 0.01). Our findings show that place fields present in the before CFC sessions in both Familiar and CFC VR showed significant remapping following fear conditioning.

Figure 8. The widths of place fields in CA1 narrow during memory recall. (A) Place fields were defined and sorted by track length in each session, pooled from all mice (*n* = 4 mice). Each cell's activity was normalized to its peak, and cells were sorted by their center of mass along the track. **(B)** The percentage of place cells is calculated as the number of place cells divided by the total number of recorded cells. More place cells were identified by our algorithm in the CFC VR on Day 1. **(C-E)** Parameters of place fields: **(C)** Width, **(D)** Place Field (PF) Reliability, and **(E)** out/in field firing ratio in both VRs on Day 0 and Day 1. The width of the place fields in the CFC VR significantly decreased on Day 1. No other parameters significantly varied across days. P-values were calculated using a *paired t-test*.

DISCUSSION

We present three paradigms for performing CFC in head-fixed mice, using a conductive

"tail-coat" to deliver aversive shocks as mice navigate VR environments on a treadmill.

Our key findings are: 1) Head-fixed mice exhibited freezing behavior in response to an

aversive stimulus in VR, consistent with conditioned responses seen in freely moving

animals. Freezing was especially pronounced during the first lap in the fear-conditioned

environment compared to the neutral environment across all paradigms tested. 2) With

mice on a spherical treadmill, additional conditioned responses such as hesitation and

backward movement were detected. 3) The choice of neutral environment influenced fear

discrimination: mice froze more in the fear-conditioned VR when the neutral environment

was familiar rather than novel. Freezing in the fear-conditioned VR could be enhanced by

shorter ISIs and the presence of the tail-coat during recall when the neutral environment

was novel. 4) Following fear conditioning, place cells in the hippocampal CA1 region

showed remapping, increased numbers, and narrower fields, paralleling findings in freely

moving animals and offering new insights.

Here, we experimented with three paradigms for fear conditioning in mice, with the main

difference being the neutral environment where the mice didn't receive any shock. In

Paradigm 1, the neutral VR was familiar, where the mice had previously received rewards,

and the CFC VR was novel. In Paradigms 2 and 3, both the neutral and CFC VRs were

similarly novel. In Paradigm 3, we decreased the ISI and kept the tail-coat on the animal

when testing for memory recall. Thus, we have tested three different variations for

performing CFC in head-fixed mice. On average, mice showed more freezing behavior in

the CFC VR compared to the neutral VR. A common feature across all paradigms was

pronounced freezing during the very first lap in the CFC VR. While mice in Paradigm 2

extinguished fear rapidly after the first lap, freezing behavior persisted up to 3-4 recall

days in Paradigms 1 and 3, to varying degrees. We also observed variability in

conditioned responses across mice. Some displayed an opposite pattern, freezing more

in the neutral VR or equally in both VRs. This variability in behavior, similar to that

observed in freely moving animals (Chu et al., 2024; Dos Santos Corrêa et al., 2019;

Navarro-Sánchez et al., 2024; Poulos et al., 2016; Russo & Parsons, 2021; Totty et al.,

2021), can be insightful when combined with investigations of neural activity, helping to

understand the neural dynamics that contribute to an animal's ability to learn the

association between a context and fearful stimuli. This can reveal the mechanisms by

which such memories may generalize or become maladaptive, as seen in post-traumatic

stress disorders.

We have shown here an example of using Paradigm 1 to study place cells in the

hippocampus and their role in fear memory formation and recall. We recorded from a

large group of the same hippocampal CA1 cells across days and demonstrated that place

cells remap and narrow their place field widths with fear conditioning. This occurs in head-

fixed mice navigating VR environments, similar to freely moving animals. Narrow place

fields may suggest sharper spatial tuning to incorporate a salient emotional memory and

may aid in context discrimination (Schuette et al., 2020). However, how place cells narrow

their widths in response to fear conditioning remains an open question. One possible way

to answer this is by imaging CA1 place cell dendrites to understand how they integrate

inputs before and after fear conditioning. We can explore what dendritic firing and

plasticity mechanisms lead to narrow place field widths. We can then perturb these firing

patterns to attribute causality to the narrowing of place field widths in fear discrimination.

Our head-fixed paradigm enables the recording of dendrites and the study of plasticity

mechanisms (Sheffield et al., 2017; Sheffield & Dombeck, 2015), thus facilitating these

investigations.

We recently used Paradigm 3 to understand the role of the thalamic nucleus reuniens-

hippocampal pathway (Ratigan et al., 2023). Using a two-photon microscope, we

identified neural activity in this pathway by imaging axons from the nucleus reuniens that

project to the CA1 with single-axon resolution. We found this pathway necessary for

suppressing fear memory retrieval, context discrimination, and fear extinction. The robust

freezing response in Paradigm 3 in the fear-conditioned VR, compared to a neutral novel

VR, allowed us to align our findings with the existing literature on the role of the nucleus

reuniens-hippocampal circuit in fear memory in freely moving animals (Ramanathan &

Maren, 2019; Ramanathan et al., 2018; Totty et al., 2023; Troyner et al., 2018; Vetere et

al., 2017; Xu & Südhof, 2013). Furthermore, the subcellular resolution achieved through

two-photon calcium imaging using the head-fixed preparation allowed us to longitudinally

observe and understand the inputs that axons from the reuniens send to the hippocampus

during fear memory formation and retrieval, a level of detail that is difficult to achieve with

current techniques in freely moving preparations.

In another unpublished study, we used Paradigm 2 and identified co-active neurons

between CA1 and CA3 subregions of the hippocampus during fear memory recall. The

difference between the first and subsequent laps has helped us understand the

progression of neural activity in the hippocampus as mice go from retrieving the fear

memory to extinction within one session. Moreover, the variability in response across

mice has allowed us to regress the neural activity to the behavior, identifying neural

differences between those mice that accurately recall the fear memory and those that do

not. These studies showcase the broad applicability of our different CFC paradigms.

In the laboratory, CFC provokes freezing behavior in freely moving rodents, as

demonstrated by several studies that span decades (Michael S. Fanselow, 1980, 1990;

Fendt & Fanselow, 1999; Maren, 2001; Maren et al., 2013; Phillips & LeDoux, 1992;

Tovote et al., 2015). In our study, we found that a mild electric shock applied to the tail

could also trigger freezing behavior as the conditioned response in head-fixed mice,

making our paradigm more comparable to the freely moving paradigms than previously

attempted methods using air-puff stimulation (Lovett-Barron et al., 2014; Rajasethupathy

et al., 2015). There are numerous future opportunities to test how various parameters,

such as shock amplitude, number of shocks, duration of fear conditioning, the

configuration of the Control versus CFC VR, or adding more sensory cues, may influence

the animal's recall and extinction behavior and how it is encoded in the brain (Michael S.

Fanselow, 1990; Huckleberry et al., 2016; Ji & Maren, 2007; Maren & Holmes, 2016;

Poulos et al., 2016). Moreover, our technique enables us to conduct a more detailed

analysis of the animal's behavior. For example, besides freezing behavior, we have

observed a sprinting behavior or a 'flight response' in response to each shock delivery

and instances of fear-related backward movements during memory recall; defensive

behaviors that have also been observed in freely moving animals (Chu et al., 2024;

Furuyama et al., 2023; Totty et al., 2021; Trott et al., 2022). As our mice are head-fixed,

our setup allows easy measure of pupil diameter, respiration, vocalization, heart rate, and

body temperature. These have been shown to change with fear conditioning (Dupin et al.,

2020; Godsil et al., 2000; Korn et al., 2017; Shionoya et al., 2013; Stiedl & Spiess, 1997).

There are also opportunities to use machine-learning-empowered algorithms to parse out

new behavior markers for contextual fear memory, such as gestures or facial expressions

(Datta et al., 2019; Mathis & Mathis, 2020; Syeda et al., 2024; Wiltschko et al., 2020).

Thus, we can understand the animal's behavior in a well-controlled environment, which

when combined with neural activity recording, can lead to new insights into the neural

circuits mediating different defensive responses and emotional states.

Significant insights into memory mechanisms through the CFC task have also come from

recent discoveries of memory engrams in the brain (Josselyn et al., 2015; Josselyn &

Tonegawa, 2020; Liu et al., 2012; Tonegawa et al., 2018). Engram cells, identified in

various brain regions with the help of genetic tagging and optogenetics, play a crucial role

in encoding, consolidating, and retrieving specific memories. Typically, these cells are

identified based on the expression of the immediate early gene *Fos* during fear

conditioning. Their artificial activation with optogenetics can trigger memory recall,

evidenced by increased freezing behavior. Understanding which neurons form these

engram cells, how *Fos* expression is driven during memory formation, how memory

information is stored in these cells, how they evolve with learning, and how their activation

results in memory recall is an active area of research (Frankland et al., 2024; Mocle et

al., 2024; Monasterio et al., 2024; Pettit et al., 2022; Suthard et al., 2024; Uytiepo et al.,

2024). Incorporating our paradigm with simultaneous measurements of cellular and

subcellular activity can help answer some of these questions. For instance, a recent study

(Monasterio et al., 2024) tagged engram cells during fear conditioning while animals were

freely moving and then imaged the spontaneous dynamics of these cells using two-

photon imaging before and after fear conditioning. The study revealed that cells forming

engrams were inherently more active. A limitation of this study was the separate contexts

for imaging versus tagging, which can be addressed with our paradigm. Our CFC

paradigm allows for direct neural recording of these cells before, during, and after tagging

in the same context to examine how activity during memory formation relates to engram

cell emergence, population dynamics, and memory recall. Furthermore, holographic

stimulation techniques can reactivate specific cell ensembles of interest to help

understand the causal relationships between engram cells and memory. Moreover, the

hippocampus contains both place cells and engram cells, both of which appear to be

involved in contextual memory. However, their relationship is not yet fully understood. Our

CFC paradigm offers an opportunity to study both place cells, as we have shown here,

and engram cells simultaneously. This can be achieved by incorporating techniques that

allow for *in vivo* labeling of *Fos* (Pettit et al., 2022)*.* Thus, our head-fixed version of the

CFC paradigm opens new avenues for research into the neural underpinnings of fear

learning and memory.

Another advantage of our method is that it allows for studying changes in neuronal

dynamics in response to an aversive stimulus, such as a shock. Our fields of view during

the shock and subsequent flight-like response periods remained stable, and we could

correct image movements during these periods with Suite2p (Pachitariu et al., 2017). This

is a significant improvement over one-photon and electrophysiological techniques, where

abrupt movements such as those induced by an electric shock can change fields of view

and reduce the likelihood of recording from the same groups of cells, thereby hindering

longitudinal assessments of neural dynamics. In Paradigm 1, where we imaged

hippocampal cells during shock, we did not find cells responding to the shock itself, as

others have reported (Barth et al., 2023; Mocle et al., 2024; Suthard et al., 2024). We

suspect this could be due to the number or duration of shocks administered and our

shocks not being associated with a specific cue or location. In future studies, we plan to

optimize these parameters to understand better how neural dynamics change in response

to an aversive cue.

Finally, we recognize that our head-fixed CFC paradigm has some limitations. Compared

to the freely moving CFC, our setup is less naturalistic. We defined different contexts as

two visually distinct VR environments while odor, sound, and tactile cues remained

constant. This could have made it harder for mice to differentiate between the two

contexts, particularly in Paradigm 2. Unlike freely moving animals in CFC studies, our

mice were water-restricted throughout the experimental process. Because the animals

need to get used to wearing a tailcoat and running in the VRs without a reward, we tend

to 'over-train' the mice. This criterion excludes mice that are less adept at running, limiting

the number that go through the process and requiring larger cohorts of mice. Furthermore,

despite our best efforts to include equal numbers of male and female mice, only some

female mice reached the experimental stage. We believe this is due to the tail-coat weight

and the reduced weight of females after water restriction compared to males (see

Methods). Although we observed no behavioral differences between the four tested

females and the male mice, we cannot guarantee similar results in other female animals

without further testing. Studies indicate that different sexes may display varying

conditioned responses, as seen in rats (Gruene et al., 2015; S. Maren et al., 1994). In the

future, we aim to address these limitations.

In conclusion, we have developed a head-fixed version of the classic CFC paradigm. Our

findings demonstrate that mice exhibit freezing behavior when fear-conditioned in a VR

environment. Moreover, we observed that hippocampal place cells remap and narrow

their fields in response to fear conditioning, mirroring observations in freely moving

animals. This innovative paradigm bridges the behavioral task gap for techniques

requiring head-fixation, paving the way for new investigations into neural circuits and

deepening our understanding of the neural foundations of emotional memory.

Acknowledgments: We thank the members of the Sheffield Lab for comments on the manuscript, helpful discussions, and feedback throughout the process.

Funding: This work was supported by The Whitehall Foundation, The Searle Scholars Program, The Sloan Foundation, The University of Chicago Institute for Neuroscience start-up funds and the National Institute for Health (1DP2NS111657-01, 1RF1NS127123-01) awarded to M.S. and a T32 training grant (T32DA043469) and K01 (K01DA060994-01) from National Institute on Drug Abuse awarded to S.K.

Author contributions: S.K. and M.S. conceived the idea. S.K. and M.S. supervised the project. S.K. designed the experiments. S.K., C.D., H.R., D.M., and C.C. collected the data. S.K. and C.D. analyzed the data. S.K., C.D., H.R., and M.S. interpreted the data. S.K. prepared the manuscript draft. All authors reviewed the results, provided feedback, and approved the final version of the manuscript.

Declaration of interests: The authors declare no competing interests.

Data and code availability: Behavior data, imaging data, and scripts used for data analysis are available on GitHub at https://github.com/seethakris/CFCMethodspaper.

Methods

Subjects

All experimental and surgical procedures adhered to the University of Chicago Animal Care and Use Committee guidelines. This study utilized 10-12 week-old C57BL/6J and Tg(Grik4-cre)G32-4Stl/J mice. Following the start of water restriction, the mice were individually housed in a reverse 12-hour light/dark cycle, and behavioral experiments were conducted during the animals' dark cycle.

Head-fixation surgery

Mice were anesthetized with 1-2% isoflurane and given an intraperitoneal injection of 0.5 mL of saline and a subcutaneous injection of 0.45 mL of Meloxicam (1-2 mg/kg). A small incision was made on the skull to attach a metal head-plate (9.1 mm x 31.7 mm, Atlas Tool and Die Works) to head-restrain the mice. Some mice also had an imaging window implanted above the hippocampus, as previously described (Dong et al., 2021; Krishnan et al., 2022; Sheffield & Dombeck, 2015). Post-surgery mice were housed individually. After a recovery day, they were water-restricted to 0.8-1.0 ml/day.

Behavior training

Head-restrained mice were trained to run on a spherical treadmill to navigate virtual reality (VR) environments similar to the setups previously described (Dong et al., 2021; Krishnan et al., 2022; Sheffield & Dombeck, 2015). Behavioral training began seven days following the water restriction. Briefly, VR environments were created using the Virtual Reality MATLAB Engine, or VIRMEN (Aronov & Tank, 2014), and projected to five screens

covering the mouse's field of view. Mice were head restrained with their limbs resting on

a freely rotating Styrofoam wheel, which acted as a treadmill. VR environments were 2 m

linear tracks. Mice received a water reward (4 µl) through a waterspout upon completing

each traversal of the linear track, considered a lap. This reward was followed by a 1.5-

second pause in the VR to allow the mouse to consume the water. The mice were then

virtually teleported to the start of the track to begin a new lap. Training sessions lasted 30

minutes per day. Mice were considered well-trained when they could run at >3 laps per

minute. We found that this level of training was necessary to ensure that mice would

continue to run even without rewards (Krishnan et al., 2022; Krishnan & Sheffield, 2023).

Generally, about 60% of mice met our criteria, which typically took around 10-14 days to

achieve. Behavior data was collected using a PicoScope Oscilloscope (PICO4824, Pico

Technology, v6.13.2).

Tail-coat

A custom-designed "tail-coat" delivered mild electric shocks to a head-fixed mouse's tail

during virtual reality (VR) navigation. A pictorial representation of the tail-coat can be

found in **Figure 1D**. The tail-coat, handmade from wearable conductive fabrics, featured

a lightweight design to ensure the mouse could continue running during baseline VR

exploration (**Table 7**). First, the mouse's tail was measured while anesthetized for head-

plate surgeries. A small denim patch cloth (Michael's) was cut into a rectangle to fit snugly

on the mouse's tail. Marks were made where snap buttons (Michael's) would be sewn,

ensuring they clipped just above the tail. Given that the mouse's tail narrows away from

the body, the buttons closer to the body were placed slightly further apart than those

closer to the tip. Typically, the buttons were set 2 cm apart on the narrow end and 2.5 cm

apart on the broader end. Care was taken to adjust the placement of the buttons so they

weren't too tight or loose for the mouse. We found that these measurements were

generally similar across sexes and for the age group that we tested. After measurements

were taken and the cloth cut, small strips of 8mm wide conductive fabric tape (Adafruit)

were attached to either side of the cloth, with some tape hanging off the edge. A metal

strip (McMaster Carr) of the same width was affixed to this portion of the tape to provide

a secure grip for alligator clips used in fear conditioning. Finally, snap buttons were sewn

with conductive thread at the marked spots. Clear nail polish was applied to secure the

threads in place. The tail-coat creation process should take less than half an hour.

Dimensions of the cloth for a typical male mouse, 12 weeks of age and weighing ~30 g,

are shown in **Figure 1D**. The denim cloth was 4 cm in width and 3.2 cm in length; the

conductive tape was 1.2 cm in width and 5.2 cm in length, with 1.5 cm hanging out at

each side.

Table 7: Items required to for head-fixed contextual fear discrimination		
Item	Manufacturer	Link
Denim patch cloth	Any fabric or hobby shop	https://www.michaels.com/product/loops-threads-denim-patches-assorted-10113795
Snap Buttons	Any fabric or hobby shop	https://www.michaels.com/product/loops-threads-sew-on-snaps-10354228?michaelsStore=1191&inv=14
Conductive nylon fabric tape	Adafruit	https://www.adafruit.com/product/3960
Stainless conductive thread – 3 Ply	Adafruit	https://www.adafruit.com/product/641
Multipurpose 304 Stainless steel sheet	McMaster-Carr	https://www.mcmaster.com/products/304-stainless-steel/multipurpose-304-stainless-steel-6/?s=304-stainless-steel

| Shock box | Colbourn precision animal shocker | https://lafayetteinstrument.com |
| | Alternate: Lafayette Instruments Scrambled Grid Current Generator | |

Contextual fear conditioning protocol in head-fixed mice

Once the mice were well-trained and met our criteria of running >3 laps per minute, they

Once the mice were well-trained and met our criteria of running >3 laps per minute, they

were ready for the next stage. The following day, a tail-coat was attached to the mouse's

tail. Mice that maintained consistent running in the VR, with minimal pauses, while

wearing the tail-coat advanced to the next stage. For mice startled by the tail-coat, we

typically attempted another day with the tail-coat, rewarding them and observing if their

running behavior improved to running at least three laps per minute. If not, these mice

were not advanced to the next stage. In mice that do advance, on the subsequent day,

the tail-coat was kept on, and the water reward was removed. All else remained the same,

including the 1.5 s pause at the end, since we wanted animals to be able to distinguish

the start and end of a lap. If the mice continued to run as before, they moved on to the

experiment stage. From that day forward, the mice no longer received water

reinforcement when running in the VR environments. Typically, about 40% of mice made

it to the experiment stage. Tail-coats were only added during VR exploration and removed

before returning the animals to their home cage. Once the mice advanced to the

experiment stage, they were taken through the entire process and not excluded from

analysis, even if they showed signs of inconsistent running behavior, which happened

with some mice (for instance, see **Supplementary Figure 3B-C**).

We implemented three distinct paradigms to assess contextual fear discrimination. In the

first paradigm, we conducted experiments over two days (Paradigm 1). On Day 0, mice

engaged with the same VR in which they were trained, now referred to as the Familiar

VR. After 10 minutes, we transitioned them to a new VR for another 10 minutes; this new

VR would later be associated with shocks (CFC VR). During this phase, the mice explored

both VRs without any rewards while wearing the tail-coat. After the initial exploration in

the CFC VR, they received six mild electric shocks, each lasting 1 second with 1-minute

intervals in between. The shocks were administered using a shock generator (Colbourn

Precision animal shocker, Harvard Bioscience). A TTL pulse was generated using

custom-written codes in MATLAB to turn on the shock generator via an Arduino. The

shock times were recorded by the PicoScope Oscilloscope (PICO4824, Pico Technology,

v6.13.2). We tested different shock intensities ranging from 0.5 to 1.2 mA. After the last

shock, mice were removed from the VR and returned to their home cage. The following

day (Day 1), we placed the mice in the Familiar VR or CFC VR for ten minutes to test

their memory recall. No rewards were delivered during this phase; however, the tail-coat

was not added during memory recall. Some mice underwent additional recall tests over

subsequent days to assess the time required for fear memory extinction. As a control,

some mice underwent the same paradigm but without any shocks.

In the second paradigm (Paradigm 2), the experiment spanned over three days. On Day

- 1, mice explored two novel VRs for ten minutes each, establishing a baseline for context

exploration. One of these VRs served as a control, while the other, the CFC VR, was

where the mice would later receive electric shocks. On Day 0, the mice again spent ten

minutes each in the Control and CFC VRs. After this, they received six mild electric

shocks, each lasting one second, with a strength of 1 mA and an inter-trial interval of one

minute. The mice wore the tail-coat on both Days -1 and 0. On Day 1, the mice were

placed in either the Control VR or CFC VR for ten minutes each, in a counterbalanced

manner, to test for memory recall. No rewards were given on all the experimental days;

however, the tail-coat was not added to the tails on the recall day (Day 1). The third

paradigm (Paradigm 3) was similar to Paradigm 2 and used two novel VRs, except that

a tail-coat was also added to the mice's tails during recall tests.

Additionally, a subset of mice in Paradigms 1 and 2 and all mice in Paradigm 3 went

through multiple days of recall tests. The subset of mice was randomly chosen, not based

on their recall behavior from the previous day. Mice were presented with either the CFC

VR or the Control VR in a counterbalanced manner.

All VRs were randomly assigned as either Control or CFC VR to prevent biasing any

inherent preferences in mice. However, the Familiar VR was always kept the same. We

also used both sexes in our experiments. However, we found female mice less receptive

to the "tail-coat" than their male counterparts. We believe this is due to the low average

weight of female mice (~16 g) following water restriction. Female mice of the same age

(10-12 weeks) weighed significantly less than males (20-23 g vs 25-30 g) and lost further

weight following water restriction. As a result, only four female mice could proceed

through and complete all the behavioral tests. Our ultimate goal is to compare behavior

and neural activity across sexes, considering sex as a biological variable. To achieve this,

we plan to adjust the age and use female mice of appropriate weight to ensure they can

run on the treadmill with the tailcoat in the future.

Behavior measurement

899   Freezing periods were periods during which the animal's instantaneous velocity was

900   lower than 1 cm/s. Freezing epochs were defined as continuous, uninterrupted freezing

901   periods longer than 1 s. The time taken to complete a lap was calculated as the total time

902   (in seconds) the animal took to run from 0 to 200 cm.

Surgery and two-photon imaging

Stereotaxic surgeries and cannula window implantation above the CA1 subregion of the

hippocampus were performed in the same way as previously reported (Krishnan et al.,

2022). Briefly, a genetically-encoded calcium indicator, AAV1-CamKII-GCaMP6f

(pENN.AAV.CamKII.GCaMP6f.WPRE.SV40 was a gift from James M. Wilson – Addgene

viral prep #100834-AAV1; https://www.addgene.org/100834/; RRID: Addgene_100834)

was injected (~50 nL at a depth of 1.25 mm below the surface of the dura) using a beveled

glass micropipette leading to GCaMP6f expression in a large population of CA1 pyramidal

cells. Mice underwent water restriction (1.0 ml/day) following viral injection and, seven

days later, were implanted with a hippocampal window and head plate (Dombeck et al.,

2010). Mice were allowed to rest from the surgeries for 3-4 days, after which behavioral

training began. Imaging was done using a laser scanning two-photon microscope

(Neurolabware). The setup is the same as the one previously described (Krishnan et al.,

2022). Time-series images were collected through Scanbox (v4.1, Neurolabware), and

the PicoScope Oscilloscope (PICO4824, Pico Technology, v6.13.2) was used to

synchronize frame acquisition timing with behavior. At the end of the imaging session on

Day -1 or Day 0, a 1-minute time-series movie was collected at a higher magnification

and then averaged to aid as a reference frame in finding the same imaging plane on

subsequent days. Across-day images collected were motion-corrected using Suite2p

(Pachitariu et al., 2017) and aligned using ImageJ (v1.53, NIH). Region of interests were

extracted using Suite2p.

Place cell extraction and place field parameters

Place fields were identified as previously described (Dong et al., 2021; Krishnan et al.,

2022). Briefly, the 2 m track was divided into 40 position bins (each 5 cm wide). The

running behavior of the animal was filtered to exclude periods where the animal was

immobile (speed <1 cm/s). Filtering ensured that place cells were defined only during

active exploration and not during freezing bouts. We have previously found that filtering

versus not filtering yields similar results (Krishnan et al., 2022). Extracted place fields

satisfied the following criteria, which were used for all conditions and all mice: 1. The

average $\Delta F/F$ was greater than 10% above the baseline. 2. The cell displayed significant

calcium transients on >30% of laps in the field. 3. The rising phase of the mean transient

was located on the track. 4. Their p-value from bootstrapping was <0.05. Multiple place

fields within the same cell were treated independently. Parameters of place fields were

calculated as described before (Krishnan et al., 2022); this includes the center of mass

(COM), place field reliability, out/in-field firing ratio, and place field widths.

Statistics

For related samples, we performed a paired t-test. Multiple comparisons of related

samples were corrected with Bonferroni post-hoc. To quantify freezing across days, we

employed a linear mixed-effects model with days of recall as the fixed effect and behavior

prior to fear conditioning on Day 0 as baseline. The model also included a random effect

for mice to account for the nesting of observations within individual subjects. Separate

models were used for each VR. Boxplots are plotted to display the entire distribution of

the data. The box in the boxplot ranges from the first quartile (25th percentile) to the third

quartile (75th percentile), and the box shows the interquartile range (IQR). The line across

the box represents the median (50th percentile). The whiskers extend to 1.5*IQR on either

side of the box, and anything above this range is defined as an outlier. A two-tailed

Kolmogorov-Smirnov (KS) Test was used to compare distributions. When not using

boxplots, mean and confidence intervals are displayed.

References

Anagnostaras, S. G., Gale, G. D., & Fanselow, M. S. (2001). Hippocampus and contextual fear conditioning: Recent controversies and advances. *Hippocampus*, *11*(1), 8 17.

Aronov, D., & Tank, D. W. (2014). Engagement of Neural Circuits Underlying 2D Spatial Navigation in a Rodent Virtual Reality System. *Neuron*, *84*(2), 442 456.

Baldi, E., Lorenzini, C. A., & Bucherelli, C. (2004). Footshock intensity and generalization in contextual and auditory-cued fear conditioning in the rat. *Neurobiology of Learning and Memory*, *81*(3), 162–166.

Bando, Y., Wenzel, M., & Yuste, R. (2021). Simultaneous two-photon imaging of action potentials and subthreshold inputs in vivo. *Nature Communications*, *12*(1), 7229.

Barth, A. M., Jelitai, M., Vasarhelyi-Nagy, M. F., & Varga, V. (2023). Aversive stimulus-tuned responses in the CA1 of the dorsal hippocampus. *Nature Communications*, *14*(1), 1–17.

Biedenkapp, J. C., & Rudy, J. W. (2007). Context preexposure prevents forgetting of a contextual fear memory: implication for regional changes in brain activation patterns associated with recent and remote memory tests. *Learning & Memory (Cold Spring Harbor, N.Y.)*, *14*(3), 200–203.

Blair, G. J., Guo, C., Wang, S., Fanselow, M. S., Golshani, P., Aharoni, D., & Blair, H. T. (2023). Hippocampal place cell remapping occurs with memory storage of aversive experiences. *ELife*, *12*. https://doi.org/10.7554/eLife.80661

Bouton, M. E. (1993). Context, time, and memory retrieval in the interference paradigms of Pavlovian learning. *Psychological Bulletin*, *114*(1), 80–99.

Bouton, M. E., & King, D. A. (1983). Contextual control of the extinction of conditioned fear: tests for the associative value of the context. *Journal of Experimental Psychology. Animal Behavior Processes*, *9*(3), 248–265.

Bouton, Mark E., & Bolles, R. C. (1980). Conditioned fear assessed by freezing and by the suppression of three different baselines. *Animal Learning & Behavior*, *8*(3), 429–434.

Chaigneau, E., Ronzitti, E., Gajowa, M. A., Soler-Llavina, G. J., Tanese, D., Brureau, A. Y. B., Papagiakoumou, E., Zeng, H., & Emiliani, V. (2016). Two-Photon Holographic Stimulation of ReaChR. *Frontiers in Cellular Neuroscience*, *10*, 234.

Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., Schreiter, E. R., Kerr, R. A., Orger, M. B., Jayaraman, V., Looger, L. L., Svoboda, K., & Kim,

D. S. (2013). Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, *499*(7458), 295 300.

Chu, A., Gordon, N. T., DuBois, A. M., Michel, C. B., Hanrahan, K. E., Williams, D. C., Anzellotti, S., & McDannald, M. A. (2024). A fear conditioned cue orchestrates a suite of behaviors in rats. *ELife*, *13*. https://doi.org/10.7554/eLife.82497

Colgin, L. L., Moser, E. I., & Moser, M.-B. (2008). Understanding memory through hippocampal remapping. *Trends in Neurosciences*, *31*(9), 469 477.

Cordero, M. I., Merino, J. J., & Sandi, C. (1998). Correlational relationship between shock intensity and corticosterone secretion on the establishment and subsequent expression of contextual fear conditioning. *Behavioral Neuroscience*, *112*(4), 885–891.

Datta, S. R., Anderson, D. J., Branson, K., Perona, P., & Leifer, A. (2019). Computational Neuroethology: A Call to Action. *Neuron*, *104*(1), 11–24.

Daviu, N., Molina, P., Nadal, R., Belda, X., Serrano, S., & Armario, A. (2024). Influence of footshock number and intensity on the behavioral and endocrine response to fear conditioning and cognitive fear generalization in male rats. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, *135*(111112), 111112.

Debiec, J., LeDoux, J. E., & Nader, K. (2002). Cellular and systems reconsolidation in the hippocampus. *Neuron*, *36*(3), 527 538.

Dombeck, D. A., Harvey, C. D., Tian, L., Looger, L. L., & Tank, D. W. (2010). Functional imaging of hippocampal place cells at cellular resolution during virtual navigation. *Nature Neuroscience*, *13*(11), 1433–1440.

Dong, C., Madar, A. D., & Sheffield, M. E. J. (2021). Distinct place cell dynamics in CA1 and CA3 encode experience in new environments. *Nature Communications*, *12*(1), 2977.

Dos Santos Corrêa, M., Vaz, B. D. S., Grisanti, G. D. V., de Paiva, J. P. Q., Tiba, P. A., & Fornari, R. V. (2019). Relationship between footshock intensity, post-training corticosterone release and contextual fear memory specificity over time. *Psychoneuroendocrinology*, *110*(104447), 104447.

Dupin, M., Garcia, S., Messaoudi, B., Doyère, V., & Mouly, A.-M. (2020). Respiration and brain neural dynamics associated with interval timing during odor fear learning in rats. *Scientific Reports*, *10*(1), 17643.

Emiliani, V., Cohen, A. E., Deisseroth, K., & Hausser, M. (2015). All-Optical Interrogation of Neural Circuits. *Journal of Neuroscience*, *35*(41), 13917 13926.

Fan, L. Z., Kim, D. K., Jennings, J. H., Tian, H., Wang, P. Y., Ramakrishnan, C., Randles, S., Sun, Y., Thadhani, E., Kim, Y. S., Quirin, S., Giocomo, L., Cohen, A. E., & Deisseroth, K. (2023). All-optical physiology resolves a synaptic basis for behavioral timescale plasticity. *Cell*, *186*(3), 543-559.e19.

Fanselow, M. S., & Bolles, R. C. (1979). Naloxone and shock-elicited freezing in the rat. *Journal of Comparative and Physiological Psychology*, *93*(4), 736–744.

Fanselow, Michael S. (1980). Conditional and unconditional components of post-shock freezing. *The Pavlovian Journal of Biological Science : Official Journal of the Pavlovian*, *15*(4), 177–182.

Fanselow, Michael S. (1990). Factors governing one-trial contextual conditioning. *Animal Learning & Behavior*, *18*(3), 264–270.

Fendt, M., & Fanselow, M. S. (1999). The neuroanatomical and neurochemical basis of conditioned fear. *Neuroscience and Biobehavioral Reviews*, *23*(5), 743–760.

Frankland, P. W., Josselyn, S. A., & Köhler, S. (2024). Engrams. *Current Biology: CB*, *34*(12), R559–R561.

Furuyama, T., Imayoshi, A., Iyobe, T., Ono, M., Ishikawa, T., Ozaki, N., Kato, N., & Yamamoto, R. (2023). Multiple factors contribute to flight behaviors during fear conditioning. *Scientific Reports*, *13*(1), 10402.

Gewirtz, J. C., McNish, K. A., & Davis, M. (2000). Is the hippocampus necessary for contextual fear conditioning? *Behavioural Brain Research*, *110*(1–2), 83–95.

Godsil, B. P., Quinn, J. J., & Fanselow, M. S. (2000). Body temperature as a conditional response measure for pavlovian fear conditioning. *Learning & Memory* , *7*(5), 353–356.

Gruene, T. M., Flick, K., Stefano, A., Shea, S. D., & Shansky, R. M. (2015). Sexually divergent expression of active and passive conditioned fear responses in rats. *ELife*, *4*. https://doi.org/10.7554/eLife.11352

Guo, Z. V., Hires, S. A., Li, N., O'Connor, D. H., Komiyama, T., Ophir, E., Huber, D., Bonardi, C., Morandell, K., Gutnisky, D., Peron, S., Xu, N.-L., Cox, J., & Svoboda, K. (2014). Procedures for behavioral experiments in head-fixed mice. *PloS One*, *9*(2), e88678.

Hattori, R., & Komiyama, T. (2022). Longitudinal two-photon calcium imaging with ultra-large cranial window for head-fixed mice. *STAR Protocols*, *3*(2), 101343.

Heer, C. M., & Sheffield, M. E. J. (2024). *Distinct catecholaminergic pathways projecting to hippocampal CA1 transmit contrasting signals during navigation in familiar and novel environments.* https://doi.org/10.7554/elife.95213

Hernandez, O., Papagiakoumou, E., Tanese, D., Fidelin, K., Wyart, C., & Emiliani, V. (2016). Three-dimensional spatiotemporal focusing of holographic patterns. *Nature Communications*, *7*, 11928.

Holland, P. C., & Bouton, M. E. (1999). Hippocampus and context in classical conditioning. *Current Opinion in Neurobiology*, *9*(2), 195–202.

Huckleberry, K. A., Ferguson, L. B., & Drew, M. R. (2016). Behavioral mechanisms of context fear generalization in mice. *Learning & Memory* , *23*(12), 703–709.

Ji, J., & Maren, S. (2007). Hippocampal involvement in contextual modulation of fear extinction. *Hippocampus*, *17*(9), 749–758.

Josselyn, S. A., Köhler, S., & Frankland, P. W. (2015). Finding the engram. *Nature Reviews. Neuroscience*, *16*(9), 521 534.

Josselyn, S. A., & Tonegawa, S. (2020). Memory engrams: Recalling the past and imagining the future. *Science*, *367*(6473). https://doi.org/10.1126/science.aaw4325

Juczewski, K., Koussa, J. A., Kesner, A. J., Lee, J. O., & Lovinger, D. M. (2020). Stress and behavioral correlates in the head-fixed method: stress measurements, habituation dynamics, locomotion, and motor-skill learning in mice. *Scientific Reports*, *10*(1), 12245.

Kim, E. J., Park, M., Kong, M.-S., Park, S. G., Cho, J., & Kim, J. J. (2015). Alterations of hippocampal place cells in foraging rats facing a "predatory" threat. *Current Biology: CB*, *25*(10), 1362–1367.

Kim, J. J., & Fanselow, M. S. (1992). Modality-specific retrograde amnesia of fear. *Science*, *256*(5057), 675 677.

Kim, Jeansok J., & Jung, M. W. (2006). Neural circuits and mechanisms involved in Pavlovian fear conditioning: a critical review. *Neuroscience and Biobehavioral Reviews*, *30*(2), 188–202.

Kinsky, N. R., Orlin, D. O., Ruesch, E. A., Diba, K., & Ramirez, S. (2023). Erasable hippocampal neural signatures predict memory discrimination. *BioRxiv.Org: The Preprint Server for Biology*, 2023.02.02.526824.

Korn, C. W., Staib, M., Tzovara, A., Castegnetti, G., & Bach, D. R. (2017). A pupil size response model to assess fear learning. *Psychophysiology*, *54*(3), 330–343.

Krishnan, S., Heer, C., Cherian, C., & Sheffield, M. E. J. (2022). Reward expectation extinction restructures and degrades CA1 spatial maps through loss of a dopaminergic reward proximity signal. *Nature Communications*, *13*(1), 6662.

Krishnan, S., & Sheffield, M. E. J. (2023). Reward Expectation Reduces Representational Drift in the Hippocampus. In *bioRxiv* (p. 2023.12.21.572809). https://doi.org/10.1101/2023.12.21.572809

Leaf, R. C., & Leaf, S. R. (1966). Recovery time as a measure of degree of conditioned suppression. *Psychological Reports*, *18*(1), 265–266.

Leaf, R. C., & Muller, S. A. (1965). Simple method for CER conditioning and measurement. *Psychological Reports*, *17*(1), 211–215.

LeDoux, J. E. (2000). Emotion circuits in the brain. *Annual Review of Neuroscience*, *23*, 155–184.

Leutgeb, S., Leutgeb, J. K., Barnes, C. A., Moser, E. I., McNaughton, B. L., & Moser, M.-B. (2005). Independent codes for spatial and episodic memory in hippocampal neuronal ensembles. *Science (New York, N.Y.)*, *309*(5734), 619–623.

Liu, X., Ramirez, S., Pang, P. T., Puryear, C. B., Govindarajan, A., Deisseroth, K., & Tonegawa, S. (2012). Optogenetic stimulation of a hippocampal engram activates fear memory recall. *Nature*, *484*(7394), 381.

Lovett-Barron, M., Kaifosh, P., Kheirbek, M. A., Danielson, N., Zaremba, J. D., Reardon, T. R., Turi, G. F., Hen, R., Zemelman, B. V., & Losonczy, A. (2014). Dendritic inhibition in the hippocampus supports fear learning. *Science*, *343*(6173), 857 863.

Mamad, O., Agayby, B., Stumpp, L., Reilly, R. B., & Tsanov, M. (2019). Extrafield activity shifts the place field center of mass to encode aversive experience. *ENeuro*, *6*(2), ENEURO.0423-17.2019.

Maren. (2001). Neurobiology of Pavlovian fear conditioning. *Annual Review of Neuroscience*, *24*(1), 897 931.

Maren, & Holmes, A. (2016). Stress and Fear Extinction. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, *41*(1), 58–79.

Maren, Phan, K. L., & Liberzon, I. (2013). The contextual brain: implications for fear conditioning, extinction and psychopathology. *Nature Reviews. Neuroscience*, *14*(6), 417–428.

Maren, S., De Oca, B., & Fanselow, M. S. (1994). Sex differences in hippocampal long-term potentiation (LTP) and Pavlovian fear conditioning in rats: positive correlation between LTP and contextual learning. *Brain Research*, *661*(1–2), 25–34.

Maren, Stephen, & Holt, W. (2000). The hippocampus and contextual memory retrieval in Pavlovian conditioning. *Behavioural Brain Research*, *110*(1–2), 97 108.

Mathis, M. W., & Mathis, A. (2020). Deep learning tools for the measurement of animal behavior in neuroscience. *Current Opinion in Neurobiology*, *60*, 1–11.

McNally, G. P., & Westbrook, R. F. (2006). A short intertrial interval facilitates acquisition of context-conditioned fear and a short retention interval facilitates its expression. *Journal of Experimental Psychology. Animal Behavior Processes*, *32*(2), 164–172.

Mocle, A. J., Ramsaran, A. I., Jacob, A. D., Rashid, A. J., Luchetti, A., Tran, L. M., Richards, B. A., Frankland, P. W., & Josselyn, S. A. (2024). Excitability mediates allocation of pre-configured ensembles to a hippocampal engram supporting contextual conditioned threat in mice. *Neuron*, *112*(9), 1487-1497.e6.

Moita, M. A. P., Rosis, S., Zhou, Y., LeDoux, J. E., & Blair, H. T. (2004). Putting fear in its place: remapping of hippocampal place cells during fear conditioning. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *24*(31), 7015–7023.

Monasterio, A., Lienkaemper, C., Coello, S., Ocker, G. K., Ramirez, S., & Scott, B. B. (2024). CA1 Engram Cell Dynamics Before and After Learning. In *bioRxiv* (p. 2024.04.16.589790). https://doi.org/10.1101/2024.04.16.589790

Muller, R. U., & Kubie, J. L. (1987). The effects of changes in the environment on the spatial firing of hippocampal complex-spike cells. *Journal of Neuroscience*, *7*(7), 1951 1968.

Navarro-Sánchez, M., Gil-Miravet, I., Montero-Caballero, D., Castillo-Gómez, E., Gundlach, A. L., & Olucha-Bordonau, F. E. (2024). Some key parameters in contextual fear conditioning and extinction in adult rats. *Behavioural Brain Research*, *462*, 114874.

O'Keefe, J., & Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Research*, *34*(1), 171 175.

Ormond, J., Serka, S. A., & Johansen, J. P. (2023). Enhanced Reactivation of Remapping Place Cells during Aversive Learning. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *43*(12), 2153–2167.

Pachitariu, M., Stringer, C., Dipoppa, M., Schröder, S., Rossi, L. F., Dalgleish, H., Carandini, M., & Harris, K. D. (2017). Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *BioRxiv*, 061507.

Pettit, N. L., Yap, E.-L., Greenberg, M. E., & Harvey, C. D. (2022). Fos ensembles encode and shape stable spatial maps in the hippocampus. *Nature*. https://doi.org/10.1038/s41586-022-05113-1

Pezze, M. A., & Feldon, J. (2004). Mesolimbic dopaminergic pathways in fear conditioning. *Progress in Neurobiology*, *74*(5), 301–320.

Phillips, R. G., & LeDoux, J. E. (1992). Differential contribution of amygdala and hippocampus to cued and contextual fear conditioning. *Behavioral Neuroscience*, *106*(2), 274–285.

Poulos, A. M., Mehta, N., Lu, B., Amir, D., Livingston, B., Santarelli, A., Zhuravka, I., & Fanselow, M. S. (2016). Conditioning- and time-dependent increases in context fear and generalization. *Learning & Memory*, *23*(7), 379 385.

Quinn, J. J., Wied, H. M., Ma, Q. D., Tinsley, M. R., & Fanselow, M. S. (2008). Dorsal hippocampus involvement in delay fear conditioning depends upon the strength of the tone-footshock association. *Hippocampus*, *18*(7), 640–654.

Rajasethupathy, P., Sankaran, S., Marshel, J. H., Kim, C. K., Ferenczi, E., Lee, S. Y., Berndt, A., Ramakrishnan, C., Jaffe, A., Lo, M., Liston, C., & Deisseroth, K. (2015). Projections from neocortex mediate top-down control of memory retrieval. *Nature*, *526*(7575), 653 659.

Ramanathan, K. R., & Maren, S. (2019). Nucleus reuniens mediates the extinction of contextual fear conditioning. *Behavioural Brain Research*, *374*(112114), 112114.

Ramanathan, K. R., Ressler, R. L., Jin, J., & Maren, S. (2018). Nucleus Reuniens Is Required for Encoding and Retrieving Precise, Hippocampal-Dependent Contextual Fear Memories in Rats. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *38*(46), 9925–9933.

Ramirez, S., Liu, X., Lin, P. A., Suh, J., Pignatelli, M., Redondo, R. L., Ryan, T. J., & Tonegawa, S. (2013). Creating a False Memory in the Hippocampus. *Science*, *341*(6144), 387 391.

Ratigan, H. C., Krishnan, S., Smith, S., & Sheffield, M. E. J. (2023). A thalamic-hippocampal CA1 signal for contextual fear memory suppression, extinction, and discrimination. *Nature Communications*, *14*(1), 6758.

Rudy, J. W., Barrientos, R. M., & O'Reilly, R. C. (2002). Hippocampal formation supports conditioning to memory of a context. *Behavioral Neuroscience*, *116*(4), 530–538.

Russell, L. E., Dalgleish, H. W. P., Nutbrown, R., Gauld, O. M., Herrmann, D., Fişek, M., Packer, A. M., & Häusser, M. (2022). All-optical interrogation of neural circuits in behaving mice. *Nature Protocols*, *17*(7), 1579–1620.

Russo, A. S., & Parsons, R. G. (2021). Behavioral expression of contextual fear in male and female rats. *Frontiers in Behavioral Neuroscience*, *15*, 671017.

Schuette, P. J., Reis, F. M. C. V., Maesta-Pereira, S., Chakerian, M., Torossian, A., Blair, G. J., Wang, W., Blair, H. T., Fanselow, M. S., Kao, J. C., & Adhikari, A. (2020). Long-Term Characterization of Hippocampal Remapping during Contextual Fear Acquisition and Extinction. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *40*(43), 8329–8342.

Sheffield, M. E. J., Adoff, M. D., & Dombeck, D. A. (2017). Increased Prevalence of Calcium Transients across the Dendritic Arbor during Place Field Formation. *Neuron*, *96*(2), 490 504.e5.

Sheffield, M. E. J., & Dombeck, D. A. (2015). Calcium transient prevalence across the dendritic arbour predicts place field properties. *Nature*, *517*(7533), 200 204.

Shionoya, K., Hegoburu, C., Brown, B. L., Sullivan, R. M., Doyère, V., & Mouly, A.-M. (2013). It's time to fear! Interval timing in odor fear conditioning in rats. *Frontiers in Behavioral Neuroscience*, *7*, 128.

Stiedl, O., & Spiess, J. (1997). Effect of tone-dependent fear conditioning on heart rate and behavior of C57BL/6N mice. *Behavioral Neuroscience*, *111*(4), 703–711.

Suthard, R. L., Senne, R. A., Buzharsky, M. D., Diep, A. H., Pyo, A. Y., & Ramirez, S. (2024). Engram reactivation mimics cellular signatures of fear. *Cell Reports*, *43*(3), 113850.

Syeda, A., Zhong, L., Tung, R., Long, W., Pachitariu, M., & Stringer, C. (2024). Facemap: a framework for modeling neural activity based on orofacial tracking. *Nature Neuroscience*, *27*(1), 187–195.

Tonegawa, S., Morrissey, M. D., & Kitamura, T. (2018). The role of engram cells in the systems consolidation of memory. *Nature Reviews. Neuroscience*, *19*(8), 485 498.

Totty, M. S., Tuna, T., Ramanathan, K. R., Jin, J., Peters, S. E., & Maren, S. (2023). Thalamic nucleus reuniens coordinates prefrontal-hippocampal synchrony to suppress extinguished fear. *Nature Communications*, *14*(1), 6565.

Totty, M. S., Warren, N., Huddleston, I., Ramanathan, K. R., Ressler, R. L., Oleksiak, C. R., & Maren, S. (2021). Behavioral and brain mechanisms mediating conditioned flight behavior in rats. *Scientific Reports*, *11*(1), 8215.

Tovote, P., Fadok, J. P., & Lüthi, A. (2015). Neuronal circuits for fear and anxiety. *Nature Reviews. Neuroscience*, *16*(6), 317–331.

Trott, J. M., Hoffman, A. N., Zhuravka, I., & Fanselow, M. S. (2022). Conditional and unconditional components of aversively motivated freezing, flight and darting in mice. *ELife*, *11*. https://doi.org/10.7554/eLife.75663

Troyner, F., Bicca, M. A., & Bertoglio, L. J. (2018). Nucleus reuniens of the thalamus controls fear memory intensity, specificity and long-term maintenance during consolidation. *Hippocampus*, *28*(8), 602–616.

Uytiepo, M., Zhu, Y., Bushong, E., Polli, F., Chou, K., Zhao, E., Kim, C., Luu, D., Chang, L., Quach, T., Haberl, M., Patapoutian, L., Beutter, E., Zhang, W., Dong, B., McCue, E., Ellisman, M., & Maximov, A. (2024). Synaptic architecture of a memory engram in the mouse hippocampus. *BioRxiv : The Preprint Server for Biology*. https://doi.org/10.1101/2024.04.23.590812

Vetere, G., Kenney, J. W., Tran, L. M., Xia, F., Steadman, P. E., Parkinson, J., Josselyn, S. A., & Frankland, P. W. (2017). Chemogenetic Interrogation of a Brain-wide Fear Memory Network in Mice. *Neuron*, *94*(2), 363-374.e4.

Wang, M. E., Wann, E. G., Yuan, R. K., Álvarez, M. M. R., Stead, S. M., & Muzzio, I. A. (2012). Long-term stabilization of place cell remapping produced by a fearful experience. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *32*(45), 15802 15814.

Wang, M. E., Yuan, R. K., Keinath, A. T., Ramos Álvarez, M. M., & Muzzio, I. A. (2015). Extinction of Learned Fear Induces Hippocampal Place Cell Remapping. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *35*(24), 9122–9136.

Williams, D. A. (1994). Slow extinction of conditioned responding following exposure to two bouts of massed shock. *Animal Learning & Behavior*, *22*(2), 203–213.

Wiltgen, B. J., & Silva, A. J. (2007). Memory for context becomes less specific with time. *Learning & Memory (Cold Spring Harbor, N.Y.)*, *14*(4), 313–317.

Wiltschko, A. B., Tsukahara, T., Zeine, A., Anyoha, R., Gillis, W. F., Markowitz, J. E., Peterson, R. E., Katon, J., Johnson, M. J., & Datta, S. R. (2020). Revealing the structure of pharmacobehavioral space through motion sequencing. *Nature Neuroscience*, *23*(11), 1433–1443.

Wu, C.-T., Haggerty, D., Kemere, C., & Ji, D. (2017). Hippocampal awake replay in fear memory retrieval. *Nature Publishing Group*, *20*(4), 571 580.

Xu, W., & Südhof, T. C. (2013). A neural circuit for memory specificity and generalization. *Science*. https://science.sciencemag.org/content/339/6125/1290.short?casa_token=QtgiZVpO0vgAAAAA:1evZSg4yR3MKpHHsrcyiJs135dikEsxBXNlOZAipkPQiXbkaIkhokkaVVO3qDwmtWTIkwSzbdcZtbbE