Reference genome choice compromises population genetic analyses

Maria Akopyan[1,2], Matthew Genchev[1], Ellie E. Armstrong[2], and Jazlyn A. Mooney[1]

1. Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, CA, USA
2. Department of Evolution, Ecology and Organismal Biology, University of California, Riverside, Riverside, CA, USA

Correspondence: J.A.M., jazlynmo@usc.edu

Highlights

- A species-specific reference genome improves read mapping and variant detection
- Reference bias underestimates genetic diversity and differentiation
- Divergent reference genomes distort demographic histories and recombination landscapes
- Unique F_{ST} outliers are detected across references, affecting functional interpretations

Keywords

reference bias; comparative genomics; population genetics; demographic inference; selection scan; conservation; recombination

Summary

Characterizing genetic variation in natural populations is vital to evolutionary biology, however many non-model species lack genomic resources. Here, we demonstrate that reference bias significantly affects population genomic analyses by mapping whole genome sequence data from gray foxes (*Urocyon cinereoargenteus*) to a conspecific reference and two heterospecific canid genomes (dog and Arctic fox). Mapping to the conspecific genome improved read pairing by ~5%, detected 26–32% more SNPs, and 33–35% more singletons. Nucleotide diversity estimates increased over 30%, F_{ST} increased from 0.189 to 0.197, and effective population size estimates were 30-60% higher with the conspecific reference. Recombination rates varied by up to 3-fold at chromosome ends with heterospecific references. Importantly, F_{ST} outlier detection differed markedly, with heterospecific genomes identifying twice as many unique outlier windows. These findings highlight the impact of reference genome choice and the importance of conspecific genomic resources for accurate evolutionary inference.

Graphical Abstract

Introduction

Accurately characterizing genetic variation is essential for understanding the evolutionary
dynamics of populations and informing management strategies for species of conservation
concern. The increasing accessibility of whole genome sequencing (WGS) data for non-model
organisms has led to a proliferation of studies employing WGS data that estimate genetic
diversity, assess population structure and connectivity, and identify adaptive genetic variation.
Recent advancements in analytical techniques for WGS data have enabled sophisticated
analyses that extend beyond classic population genetic diversity metrics to characterize
structural variation[1], infer demographic histories[2], and estimate population recombination rates[3].
These developments provide deeper insights into the historical and contemporary processes
shaping genetic variation in natural populations. However, the reliability of these inferences and
cross-population comparisons depends on the reference genome used. Reference genomes
are essential for comparing individuals and populations, but they do not capture the full
spectrum of genetic diversity. Population-level diversity is missed because a haploid reference
represents variation from only a single individual at each site in the genome.
Reference bias arises because a single reference genome—usually a haploid sequence
from one individual—is used as the coordinate system for mapping. This reference cannot fully
represent the genetic diversity of an entire species. As a result, sequencing reads that closely
match the reference tend to map with higher quality and are retained, while dissimilar reads with

lower quality scores often map poorly and are discarded. This can lead to mapping errors or
missed variant calls, especially for true population-specific genetic variation. Thus, reference
bias will result in missed or mis-called genotypes that will impact estimates of genetic diversity
and downstream analyses with the potential to distort conclusions about the population's
evolutionary history. Reference bias, along with sequencing depth, has been shown to influence
downstream genetic analyses. For example, in simulated low-coverage (2-4x) genomes,
introducing a 2% reference divergence can cause estimates of θ to increase by 0.14- to 0.18-
fold[4]. Similar effects were noted on commonly used neutrality statistics: a small downward bias
in Tajima's D, an upward bias in Fu and Li's D, and a strong downward bias in Fay and Wu's H[4].
Importantly, these shifts were still observed even at moderate coverage (8x) and in empirical
data[4].
It is often the case that researchers will map to a reference genome from related species
when a conspecific reference genome is unavailable. For example, in modern dogs and wolves,
using a heterospecific (wolf) reference instead of a conspecific (dog) reference led to a 10%
difference in heterozygosity estimates between individual dogs[5]. In non-model species like big
cats, using a conspecific reference can alter both heterozygosity (ranging from a 0.25-fold
decrease to a 3.21-fold increase) and demographic history inference[6]. In fishes, reference bias
has been associated with overestimated heterozygosity, with local reference genomes reducing
reference/non-reference heterozygous calls by 0.13-0.31% and non-reference/non-reference
calls by 0.05-0.10%[7]. Studies have also shown that reference bias leads to inaccurate detection
of structural variation[8] and distorted phylogenies[9], potentially skewing conclusions about
population history and genetic diversity in fish species. Lastly, recent work on the impact of
reference bias in two species of conservation concern, the New Zealand kiwi and the beluga
whale, revealed alarming mismatches in demography and inhibited the detection of long runs of
homozygosity[10].
Despite the impact of reference bias on population genetic inference, few studies have
examined how it may affect the characterization of demographic history and population
recombination rates, which are increasingly popular targets of estimation. The gray fox (*Urocyon*
cinereoargenteus) provides a compelling case study in this research. Gray foxes are widely
distributed across North America, and despite their phenotypic similarity, genetic evidence
suggests deep divergence between eastern and western lineages[11]. Previous gray fox genetic
studies have relied on a domestic dog (*Canis lupus familiaris*) reference genome[12], which poses
challenges due to significant karyotypic differences between the two species[13,14]. In fact, despite
most canid study's reliance on the domestic dog reference genome, the clade contains a
number of large karyotypic rearrangements, for example, dogs have 38 pairs of autosomal
chromosomes, gray foxes have 32, and the Arctic fox (*Vulpes lagopus*) has 23-24.
The domestic dog, gray fox, and Arctic fox represent distinct evolutionary lineages within
Canidae[15]. The gray fox represents the most basal lineage within the family, having diverged
approximately 10 million years ago from all other living canids[15]. The remaining canids are
comprised of three major clades: the red fox clade, the South American clade, and the wolf-like
clade. The Arctic fox belongs to the red fox clade, which diverged from the wolf-like clade
approximately 7-8 million years ago. The domestic dog is part of the wolf-like clade and
diverged most recently, having split from other wolf-like canids about 3-4 million years ago.
There are also differences in the genome assembly sizes: the gray fox possesses the largest

genome (2.66 Gb) compared to dog (2.48 Gb) and Arctic fox (2.35 Gb), some of which may be
due to the sex of the individual selected for assembly. While GC content is relatively similar
across all three species (~42%), repeat content is variable, with the gray fox exhibiting the
highest percentage (38.23%), followed by dog (33.67%) and Arctic fox (31.29%). These
genomic differences are likely to bias downstream haplotype-based analyses that rely on
synteny and linkage disequilibrium (e.g., demographic inference and recombination mapping).
Despite this, most previous genomic research within *Canidae*—including studies on common
(gray fox), near-threatened (Channel Island fox)[16,17], and endangered (Ethiopian wolf)[18]
species—has been conducted by mapping to the domestic dog genome, primarily because it
was the only canid reference genome available with annotations and had both high continuity
and contiguity[15].
The recently published gray fox reference genome[14] provides us with a high-quality
conspecific reference for alignment, variant calling, and ultimately for quantifying reference bias.
Here, we re-analyze previously published WGS data from two populations of North American
gray foxes[12]. We examine how inference of demographic histories and recombination
landscapes differ when using the heterospecific domestic dog and Arctic fox genomes in
contrast to the conspecific gray fox genome. To investigate the effects of reference bias on our
analysis, we compare the underlying genetic variation, and the site frequency spectrum (SFS)
estimated for each population using different reference genomes, as well as estimates of
genetic diversity and differentiation. Finally, we examine the effect of reference bias on both
coalescent and SFS based demographic inference, F_{ST} outlier scans for selection, as well as
recombination rate inference.

Results

We analyzed whole-genome resequencing data from 12 gray foxes, including six from the
eastern U.S. and six from the western U.S. populations (Figure S1 and 1A). We examined
reference bias by mapping these data to three reference genomes: the domestic dog genome
CanFam4[19], the Arctic fox genome[20], and the conspecific gray fox genome[14] (Figure 1B). Gray
fox reads showed significantly higher mapping success to the conspecific reference genome
compared to heterospecific genomes (99.7% vs. ~99%; χ^2_{KW}=33.19, P<0.001, Table 1), with
genes in unmapped regions enriched for sensory perception and immunity functions (details in
supplementary text). Additionally, proper read pairing was higher with the conspecific reference
(94.7%) vs. heterospecific references (89.4-90.3%), representing nearly 5% more properly
paired reads (χ^2_{KW}=40.14, P<0.001).
126
127   *A species-matched reference genome yields more SNPs and rare variants*
128
We examined the impact of reference genome choice on the detection and characterization of
genetic variation and allele frequency distributions in gray fox populations. Across all reference
genomes, we detected more SNPs in the western population of gray foxes compared to the
eastern population, with nearly double the number of SNPs identified in the west (Figure 1;
Table 1). Notably, the gray fox reference genome consistently yielded more SNPs compared to

the heterospecific references, with 26-32% more variants identified in each population (Table 1,
Figure 1).

We identified more singletons (i.e., rare variants where an allele is only found once in the
population sample) in the east and west using the gray fox reference (Figure 1,2). Specifically,
the conspecific reference detected 33% more singletons (2.4M vs. 1.8M) in the eastern
population and 35% more singletons (5.4M vs. 4M) in the western population compared to the
heterospecific references. In addition, mean SNP depth varied slightly across genomes, with
gray fox showing higher values (mean: east; 8.5, west; 8.7) compared to Arctic fox and dog
(both east: 8.3, west: 8.6). Missing data on a per-site basis was lowest with the gray fox genome
(east and west: 0.02), while the Arctic fox and dog genomes had higher missingness (east:
0.05, west: 0.03).

After lifting over SNPs in the heterospecific genomes to their corresponding positions in
the gray fox genome, we found that approximately 80% of the variants identified using
heterospecific reference genomes were also identified as variants when using the gray fox
genome (Figure 1). Among these matching variants, non-singleton SNPs made up about 50-
53%, while the remainder were singletons. The west population had a higher proportion of
matching singletons compared to the east (~32% vs. ~24%), whereas the east population had a
slightly higher proportion of matching non-singleton SNPs (~53% vs. ~50%). For the variants
that did not lift over as SNPs, the majority (14-18%) were not identified as any type of variant in
the gray fox genome. Of these, about half mapped to invariant sites, while the other half did not
map at all (Table S1).

A species-matched reference genome results in higher effective population size estimates

We inferred demographic histories using smc++[21] and observed distinct population size
trajectories when using different reference genomes for both the eastern and western
populations. Across all genomes, the western population consistently showed higher effective
population size (N_e) estimates than the eastern population (Figure 2). The gray fox genome
produced trajectories with smaller fluctuations in N_e over time and generally higher N_e
estimates, particularly for the western population. In contrast, the Arctic fox and dog genomes
revealed more variability, especially in the west, where the discrepancies between reference
genomes were more pronounced. For instance, in the most recent time period, approximately
5,000-7,000 years ago, the gray fox genome indicated population growth in the west, although it
should be noted that smc++ typically has higher uncertainty for estimates in the recent past,
<6,000 years ago[21]. Meanwhile, the heterospecific genomes suggested a population decline,
with N_e estimates dropping below 75,000.

To evaluate whether removing variants that failed to map between species improves
demographic inference accuracy, we compared smc++ trajectories before and after masking
non-lifted-over SNPs in both heterospecific genomes and reciprocally in the gray fox genome.
This approach made the demographic trajectories more similar across reference genomes
(Figure S2). We also suspected that the total number of SNPs might affect demographic
inference, so we down-sampled the gray fox SNPs to random subsets matching the number of
variants available when using lifted-over variants from other genomes. These down-sampled
trajectories were also similar to the lifted-over results, indicating that observed differences in

demographic trajectories were driven not only by variant identity differences but also by the total
number of informative variants available for analysis.
To evaluate how reference bias impacts demographic inference methods, we
supplemented smc++ analysis (combining SFS with linkage disequilibrium in coalescent HMMs)
with MSMC2 (coalescent HMM only) and Stairway Plot2 (SFS only) (Figure S3). Both MSMC2
and Stairway Plot 2 exhibited greater inconsistencies across reference genomes for the eastern
population compared to the western population. MSMC2 inferred recent growth for the eastern
population across all references, but heterospecific references exaggerated this growth, yielding
eastern population sizes surpassing western ones—an outcome not observed with other
methods and inconsistent with higher western diversity estimates. In contrast, Stairway Plot 2
revealed larger discrepancies in the eastern population during ancient times. For the western
population, MSMC2 inferred recent growth only when using heterospecific references, whereas
Stairway Plot 2 indicated stability across references, highlighting the varying sensitivities of
demographic inference methods to reference bias. Despite variation across reference genomes,
the methodological approach had a stronger influence on demographic inference outcomes.
The disparities between methods highlight the critical importance of selecting appropriate
demographic reconstruction methods, as method-specific biases can overshadow reference
genome variation and underscores the need for careful method selection
196
197 *Recombination rates vary across reference genomes and populations*
198
We detected significant differences in the distributions of recombination rates inferred using the
different reference genomes in both populations (Anderson-Darling k-sample test, $p<0.001$;
Figure 3). In the eastern population, recombination rates were lower when inferred using the
Arctic fox genome compared to the gray fox genome. The average recombination rate was
0.448 cM/Mb with the Arctic fox genome, which is approximately 31% lower than the 0.650
cM/Mb observed with the gray fox genome. Conversely, when using the CanFam4 genome, the
average increased to 0.872 cM/Mb, about 34% higher than the rate inferred with the gray fox
genome. In the western population, recombination rates were higher when inferred using both
heterospecific genomes compared to the conspecific gray fox genome. Specifically, the average
recombination rate was 1.02 cM/Mb with the Arctic fox genome, approximately 13% higher than
the 0.903 cM/Mb observed with the gray fox genome, and 0.989 cM/Mb with the CanFam4
genome, about 9.5% higher than the rate inferred with the gray fox genome. Overall, these
differences were more pronounced in the east, where the heterospecific genomes led to both
underestimation (Arctic fox) and overestimation (CanFam4) of recombination rates relative to
the gray fox genome. In contrast, in the west, both heterospecific genomes consistently
overestimated recombination rates compared to the gray fox genome, and the differences were
less pronounced.
Additionally, recombination landscapes based on each genome revealed substantial
variation across 50 kb windows, with heterospecific reference genomes resulting in increased
variability, particularly toward chromosome ends. In the eastern population, using the
conspecific gray fox genome, recombination rates ranged from 0.000343 to 7.34 cM/Mb.
However, when using the heterospecific Arctic fox genome, the maximum recombination rate
doubled to 14.9 cM/Mb, and with the CanFam4 genome, it more than tripled to 23.4 cM/Mb.

Similarly, in the western population, recombination rates estimated with the gray fox genome
varied from 0.000598 to 11.6 cM/Mb. With the Arctic fox genome, the maximum recombination
rate more than doubled to 24.9 cM/Mb, and with the CanFam4 genome, it increased to 14.5
cM/Mb. Compared to the gray fox genome, the heterospecific genomes resulted in higher
maximum recombination rates, with a consistent trend of overestimation toward the ends of
chromosomes, suggesting that heterospecific reference genomes can lead to an overestimation
of recombination rates, especially at the higher end of the spectrum.
229
230 *Heterospecific genomes underestimate diversity and differentiation*
231
We detected significant differences in estimates of average nucleotide diversity (π) in 50 kb
windows in both gray fox populations based on the three reference genomes (Figure 4). In the
eastern population, nucleotide diversity was lowest with the CanFam4 genome (π=0.000618),
followed closely by Arctic fox (π=0.000625), while the gray fox genome yielded a higher
estimate of π=0.000812 (χ^2_{KW} = 6663.3, P<0.001). Similarly, in the western population, diversity
was lowest using the heterospecific references (Arctic fox: π=0.00122, CanFam4: π=0.00121),
while the gray fox genome produced a higher estimate of π=0.00164 (χ^2_{KW}=12409, P<0.001).
The variation in estimates of nucleotide diversity was slightly more evident in the west
population, where π was approximately 1.34 times higher using the gray fox genome compared
to the heterospecific references, while in the east, π was 1.31 times higher. While estimates of
π were higher in the west than in the east across all reference genomes, the difference between
populations was greater when using the gray fox genome, where diversity in the west was about
2.02 times higher than in the east. By contrast, diversity based on the Arctic fox genome
showed a 1.95-fold difference between the west and east, and CanFam4 showed a similar 1.96-
fold difference.
Estimates of genetic differentiation (F_{ST}) between the eastern and western populations
also varied depending on the reference genome. The heterospecific references produced
identical F_{ST} values (mean F_{ST}=0.189), while the gray fox genome resulted in a significantly
higher mean F_{ST} of 0.197 (χ^2_{KW}=558.27, P<0.001). Additionally, the correlation between
nucleotide diversity and F_{ST} was more strongly negative in the eastern population (τ=-0.173 to -
0.196) compared to the western population (τ=-0.081 to -0.118), with the strongest correlations
observed using the gray fox reference in both populations.
In both populations, mean Tajima's D was lowest with the gray fox genome (-0.108 east,
-0.356 west), compared to the Arctic fox (-0.056 east, -0.329 west) and CanFam4 (-0.063 east, -
0.328 west) references (χ^2_{KW} = 195.3 east, 144.4 west; P<0.001). Pairwise comparisons
confirmed significantly lower Tajima's D estimates with the gray fox genome (P<0.001), while
estimates did not differ between the two heterospecific references (P=0.17 east, P=1.00 west).
This indicates an excess of low-frequency polymorphisms when mapping to the conspecific
genome, consistent with the higher number of singletons detected in the SFS based on the gray
fox reference.
262
263 *Reference genome choice affects F_{ST} outlier detection*
264

We defined outliers as windows with F_{ST} values exceeding three standard deviations above the
mean for each reference genome, then matched these windows across reference genomes to
identify shared and unique outlier regions. Across comparisons, we observed that heterospecific
references consistently identified more than twice the number of unique outlier windows
compared to the conspecific reference (Figure 5). Notably, the number of windows unique to
each heterospecific reference was similar to the number of shared outliers between the
heterospecific and conspecific references. In the Arctic fox and gray fox comparison, 148
shared outlier windows were identified, with 137 unique to the Arctic fox reference and 63
unique to the gray fox reference (Figure 5A). Between the Canfam4 and gray fox reference
genomes, 165 outlier windows were shared, with 141 unique to Canfam4 and 61 unique to gray
fox (Figure 5B). The majority of windows in both comparisons did not show elevated F_{ST} values
in either reference (33,350 and 31,649 in the Canfam4 and the Arctic fox comparisons,
respectively), reflecting our conservative method of outlier detection.
Between the Arctic fox and Canfam4, we identified 191 shared outlier windows, 158
windows unique to the Arctic fox reference, and 42 windows unique to the Canfam4 reference,
with 35,133 windows showing no elevated F_{ST} values in either reference (Figure S5). Thus, the
Arctic fox and Canfam4 references shared slightly more outliers than either shared with gray fox
(148-165), while the number of unique outliers for each heterospecific reference remained
comparable to their gray fox comparisons.
284
285 *Functional enrichment patterns reveal reference-specific biological processes in F_{ST} outliers*
286
The functional enrichment analysis of F_{ST} outliers revealed both distinct and overlapping gene
ontology (GO) biological process terms across reference genomes (Figure 4C). Gray fox had
the highest number of unique terms with 74, followed by Canfam4 with 71, and Arctic fox with
65, indicating that each reference captured distinct biological processes. Importantly, there were
more unique terms than shared terms across references. Among shared terms, 55 were
common to all three references. Notably, the overlap between gray fox and Canfam4 (48 shared
terms) was larger than the overlap between Arctic fox and either gray fox (25 terms) or Canfam4
(26 terms), suggesting a closer alignment in identified biological processes between the gray fox
and Canfam4 references.
Further, the unique terms identified by each reference genome suggested distinct
thematic focuses (Figure S6). The Arctic fox reference highlighted processes related to cellular
transport, secretion, and signaling, with terms like "positive regulation of catecholamine
secretion" and "golgi to endosome transport," indicating an emphasis on cellular signaling and
developmental regulation. The Canfam4 reference captured growth, differentiation, and
endocrine response, with terms such as "positive regulation of muscle cell differentiation" and
"response to thyroid hormone," pointing to processes involved in physiological development. In
contrast, the gray fox reference emphasized nervous system function, immune modulation, and
cell proliferation, reflected by terms like "positive regulation of nervous system process" and
"calcineurin-mediated signaling."
Gene Ontology (GO) enrichment analysis of genes located exclusively in Arctic fox
reference-specific outlier windows revealed significant enrichment for terms related to muscle
development and metabolism. The top biological process terms included "Muscle Organ

Development" (GO:0007517), "Positive Regulation of Muscle Hypertrophy" (GO:0014742), and
"Positive Regulation of Glycolytic Process" (GO:0045821). In contrast, genes in Canfam4
reference-specific outlier windows showed markedly different functional enrichment patterns,
predominantly associated with chromosomal organization and cellular differentiation. The most
significant terms included "Positive Regulation of Chromosome Organization" (GO:2001252),
"Cell Differentiation in Spinal Cord" (GO:0021515), and "Regulation of Chromosome
Separation" (GO:1905818). These distinct functional enrichment patterns demonstrate how
reference genome choice can lead to substantially different biological interpretations. The Arctic
fox reference-specific outliers suggest selection on genes involved in muscle development and
energy metabolism, which could be interpreted as adaptations related to locomotion, hunting
behavior, or thermal regulation. Conversely, the Canfam4 reference-specific outliers point
toward selection on chromosomal organization and cellular differentiation pathways, potentially
indicating differences in developmental processes or cell cycle regulation.

Discussion

Reference genomes are essential tools in genetic studies, serving as coordinate systems for
annotating sequence features and comparing individuals and populations. However, many non-
model organisms lack annotated chromosome-level assemblies[22], forcing researchers to use
genomes from closely related species[23,24], which can compound reference bias. Reference bias
has been previously studied across the tree of life[5,6,10,25–28] both within species[26,27] and between
species[6]. In this study, we demonstrate that mapping to a heterospecific reference genome
significantly impacts population genomic inferences. This leads to discrepancies in effective
population size estimates, recombination rates, measures of genetic diversity, differentiation,
and selection signals. Our findings highlight the importance of using conspecific genomes for
accurate evolutionary inferences.

Although the average sequencing depth showed only minor discrepancies using a discordant
reference, we observed significant differences in both the percentage of reads mapped and the
proportion of properly paired reads. Reads are not properly paired when the read orientation is
incorrect, one read does not map at all, and/or the gap between the reads are of an unexpected
size. Although thresholds for gap size and mismatch rates can be changed during read
alignment, most studies use inconsistent filters at this stage of the pipeline, including retaining
only reads with a certain MAPQ score, retaining proper pairs, or only retaining the primary
alignment[29]. Likely, the number of properly paired reads decreases when the gray fox is
mapped to the heterospecific genomes because of structural variation between the genomes,
some of which may be smaller indels, and some of which may reflect the large chromosomal
rearrangements between the species. Following common practice, we retained improperly
paired reads, which may explain downstream variant detection effects, though selective filtering
could potentially mitigate this.

To exemplify the inconsistency of pipelines and filters applied, we screened five papers which
used the domestic dog as a reference genome for analyses of a heterospecific canid taxa and
found that four papers did not remove improperly paired reads before variant calling[12,16,18,30],

while one paper did remove improperly paired reads[17], and their remaining mapping and filtering
choices had almost no overlap. Despite many of these studies filtering on MAPQ, mapping
quality is assigned for an individual read by BWA[31], and not as a pair (i.e., the MAPQ is
ambiguous to the read being properly paired). GATK automatically filters reads with a MAPQ of
10 or less during variant calling and other filters such as genotype quality (GQ; accuracy of the
particular variant call at that site) and QUAL (how confident the variant caller is that there is
variation at that site) are also likely to affect the resulting variant calls. Though we do not
explore these as part of this work, a systematic investigation of these filters and their
downstream impacts on mapping and variant calling, especially in the case of divergent
references, is highly needed.
361
We found that the conspecific gray fox genome provided a more accurate representation of
genetic variation, revealing more SNPs and a more reliable SFS compared to the heterospecific
genomes. Heterospecific references underestimated the presence of low-frequency alleles,
which can have profound implications for downstream analyses. For example, underestimating
rare variants can lead to failure to detect recent population expansion in demographic inference,
as the characteristic excess of rare alleles that signals expansion would be artificially reduced.
Additionally, an underestimation of genetic load can occur when assessing deleterious
mutations, since many harmful variants tend to be maintained at low frequencies due to
purifying selection. Furthermore, recombination rate estimates in population-scale maps can
become distorted due to missing variants, as these estimates rely on accurate measurement of
linkage disequilibrium patterns between neighboring loci.
373
The higher mean SNP depth and lower missing data rates with the gray fox reference further
support its superior performance in variant detection, indicating a more accurate and
comprehensive alignment that reduces the likelihood of false negatives. Approximately 20% of
the variants identified with heterospecific references were incorrectly identified when lifted over
to the gray fox genome (Figure 1). Of these, ~5% were misclassified (singleton vs. non-
singleton), and ~15% were unrecognized as variants, with half mapping to monomorphic sites
and half failing to map (Table S1). The 20% discrepancy likely results from sequence
divergence between species, leading to alignment errors or misinterpretation of homologous
regions. The higher proportion of matching non-singleton SNPs suggests that common variants
are more consistently detected across references, while rare variants are more likely to be
missed when using heterospecific genomes. When estimating nucleotide diversity (π), the gray
fox reference produced higher values in both populations, indicating that heterospecific
genomes can underestimate genetic diversity within populations. The greater difference in π
between the western and eastern populations observed with the gray fox genome also suggests
that the conspecific reference provides a more sensitive measure of genetic diversity.
389
Having performed the first demographic inference for the gray fox with a conspecific reference
genome, we sought to compare the eastern and western gray fox populations to gain deeper
insights into their population histories. These two populations had quite distinct demographic
trajectories, with the western population showing overall larger and stable population size with
no recent growth. In contrast, the eastern population showed a stable population with some

recent growth. Genetic diversity was significantly higher in the western gray fox population
compared to the eastern population, suggesting differing evolutionary pressures and histories in
these lineages. Our results differ from previous studies[12] of these populations, which mapped to
CanFam3.1 and used PSMC on a single individual to infer the ancient demographic histories of
the populations. Specifically, we infer smaller and more stable populations throughout time, with
less of a disparity between east and west population sizes. Both our study and the previous
study identified the same pattern of the western population being more diverse than the eastern
population for most timepoints, though the magnitude of our diversity estimates is much lower
than previous estimates[12]. We also did not detect a recent decline in the west, which was found
only when mapped to heterospecific references in our analysis as well as in previous work[12].
405
We used multiple methods for demographic inference to determine if coalescent or SFS-based
approaches were more robust to reference bias. All methods showed that heterospecific
genomes consistently underestimated effective population sizes compared to the conspecific
reference. The conspecific reference produced estimates that align more closely with
expectations based on species biology and similar studies in other mammals[32]. Importantly,
heterospecific genomes produced demographic trajectories inconsistent with the conspecific
reference throughout most of the inferred history, regardless of population or inference method
(Figure S3). This inconsistency is particularly worrying in the case of species threatened with
extinction, because approximately 99% of these species do not have a reference genome[33] and
researchers will ultimately be forced to map to a heterospecific reference. When researchers
must use heterospecific references, several strategies can mitigate reference bias. For instance,
using end-to-end alignment methods that require the entire read to align reduces bias at indels,
compared to local aligners that allow parts of reads that do not align to be ignored (i.e., soft
clipped)[34]. Using local realignment methods that ensure consistent gap placements across all
reads covering the same region, and comparing results across different filtering thresholds can
also help reduce bias[34]. Additionally, constructing consensus references or using multiple
population reference genomes in a "reference flow" approach effectively reduces bias while
requiring fewer computational resources than graph-based methods[35].
424
Since pedigree-based recombination estimates are difficult to obtain and have low resolution,
researchers typically use population-based methods like pyrho[3] to infer recombination maps that
account for demographic histories. Focusing on the recombination rate estimates inferred with
the species-matched reference, we found that landscapes of recombination differed notably
between the eastern and western populations, with the west consistently exhibiting higher
recombination rates, particularly at chromosome ends. A recombination landscape that is fairly
stable with peaks towards the end of the chromosome has been previously observed in dogs[36]
and is particularly interesting given that the family *Canidae* has a psuedogenized copy of
positive-regulatory domain zinc finger protein 9 (PRDM9)[14,37]. PRDM9 is known to initiate
meiotic recombination by specifying the locations of double-strand breaks and creating
recombination hotspots. Previous work highlighted how its pseudogenization in *Canidae*
contributes to observed recombination patterns in dogs, which can be used to identify the
directionality of chromosomes. Chromosome directionality has yet to be established for the new

reference genome, so future work could take advantage of the result to identify directionality of
chromosomes in the gray fox.
440
Conversely in the case of heterospecific reference genomes, where demographic trajectories
were quite discordant from the conspecific reference, the inferred recombination maps and rates
differed significantly in both populations. In the eastern population, recombination rates were
underestimated with the Arctic fox genome and overestimated with the dog genome compared
to the gray fox genome, while in the west, both heterospecific genomes overestimated
recombination rates, with less pronounced differences. The inconsistent direction of bias—
where the Arctic fox genome underestimates recombination rates in the eastern population but
overestimates them in the western population compared to the gray fox genome—suggests that
population-specific patterns may influence how reference bias manifests. This variability can
complicate mitigation efforts, as the potential sources of bias differ between populations, making
uniform corrections ineffective and highlighting the complexities in addressing reference bias
when it varies between populations. In addition, recombination landscapes based on
heterospecific genomes showed greater variability, particularly toward chromosome ends, with
higher maximum rates than those inferred using the gray fox genome. These patterns highlight
how reference genome choice can introduce substantial bias, affecting both the magnitude and
distribution of inferred recombination rates.
457
Potential sources of bias for the inferred recombination rate include chromosomal synteny and
spurious singletons. In the context of synteny, previous work has shown that there have been
large scale chromosomal rearrangements within the evolutionary history of these species[14].
These karyotype differences have resulted in less than half of the dog and Arctic fox
chromosomes being syntenic with gray fox chromosomes (Figure S8; Table S2). Thus, when a
recombination map is estimated using a heterospecific genome with karyotypic differences as a
reference, we are ultimately disrupting linkage patterns and inferring false recombination events
across multiple chromosomes. To investigate this, we examined whether differences in
recombination rates between each heterospecific reference and the conspecific reference
varied between syntenic and non-syntenic chromosomes but found no difference in the mean
recombination rates in either population or genome comparison (Figure S9). The observed
inflation in recombination rates could result from disrupted linkage patterns caused by lack of
synteny, increased sequence divergence leading to spurious singletons, or an interaction of
these factors. Disentangling these contributions is challenging because they are inherently
interconnected: synteny disruptions and sequence divergence both affect how variants are
identified and mapped, while spurious singletons may arise as an artifact of these processes,
obscuring their individual effects. In syntenic regions, pyrho[3] would infer inflated recombination
rates when there are spurious singletons and other incorrectly mapped variants being
interpreted as evidence of recombination events. The misclassification and misidentification of
~20% of the variants as a result of reference bias (Figure 1) is likely what led to skewed
estimates of recombination rates, especially in regions of the genome where alignment is poor.
The observation of non-syntenic regions and lower reference genome quality introducing bias
was also captured in recent work examining an improved olive baboon reference genome[38].
481

Finally, we examined estimates of F_{ST} and used the genome-wide distribution of F_{ST} to perform
an outlier-scan to detect genes that are potentially under selection. We found that estimates of
F_{ST} were higher when using the gray fox genome compared to the values obtained from
heterospecific references. We also observed a stronger negative correlation between π and F_{ST}
when using the gray fox reference, suggesting that regions of low diversity correspond to areas
of high differentiation. The discrepancy among overall π and F_{ST} values across reference
genomes is not concerning in terms of magnitude, suggesting that genome-wide averages are
likely sufficient for broadly characterizing diversity within, and differentiation between,
populations. However, studies typically go beyond genome-wide averages, examining variation
in F_{ST} across the genome to identify regions of high differentiation, which may indicate the
presence of genes under selection or regions contributing to reproductive isolation.
493
Importantly, we found that the choice of reference genome substantially affected F_{ST} outlier
detection. Heterospecific references identified more than twice the number of unique outlier
windows compared to the conspecific reference. This discrepancy could be due to alignment
issues, where regions not well conserved between species may erroneously appear as highly
differentiated regions. The limited overlap of outlier windows between references raises
concerns about the reliability of using heterospecific genomes for detecting genomic regions
under selection. The distinct GO terms associated with each reference further illustrate how
reference genome choice shapes the inferred biological processes underlying F_{ST} outliers. For
instance, unique GO terms in the Arctic fox reference emphasized cellular signaling and
developmental regulation, while Canfam4 highlighted growth and endocrine response, and the
gray fox reference captured nervous system function and immune modulation. These
differences underscore the biological biases introduced by reference genome choice. In sum,
using a heterospecific reference genome would inflate false positives for both the F_{ST} outliers,
genes, and biological pathways that may be under selection. Thus, using a conspecific
reference would certainly provide more accurate insights into the biological processes
influencing genetic differentiation within and between species.
510
This study highlights the critical impact of reference genome choice on population genomic
inferences and emphasizes the value of conspecific genomes in uncovering accurate
evolutionary histories. Quantifying reference bias using *Canidae* has implications for some of
the world's most endangered species, such as the Ethiopian wolf (*Canis simensis*) and the
African wild dog (*Lycaon pictus*), and more common species, like the gray fox and coyote
(*Canis latrans*). Reference bias may also affect our understanding of historic hybridization and
introgression, a phenomenon common in the canid clade[30,39]. Furthermore, genomic data are
increasingly being used to inform conservation management plans. Measures such as adaptive
capacity[40] and differential adaptations between populations[41] are being used to make
recommendations regarding translocations and rewilding. Our selection scans suggest that
these inferences are skewed by reference bias when divergent reference genomes are used.
Our results underscore the necessity of using conspecific reference genomes in conservation
genetics and evolutionary studies, particularly for accurately understanding divergence and
diversity in non-model organisms. However, we acknowledge that developing high-quality
reference genomes for all species is not always feasible due to resource constraints. In such

cases, careful consideration should be given to the phylogenetic proximity and degree of
synteny of the available reference genomes. Our work demonstrates that even closely related
heterospecific references may not adequately capture the genetic landscape of the target
species. Future studies should focus on improving reference genome assemblies for non-model
organisms, leveraging advancements in sequencing technologies and assembly algorithms.
Additionally, methodologies that are less reference-dependent, such as reference-free variant
calling or pangenomic approaches, may help mitigate some of the biases introduced by
heterospecific references.
534
535 *Limitations of the Study*
536
Our work provides important insights into reference bias but has limitations. First, the gray fox
reference genome comes from an east coast individual, potentially introducing bias that future
work could mitigate computationally or by generating a west coast reference. Second, we
quantified reference bias in *Canidae*, a mammalian clade with notable karyotype shifts. Despite
these shifts, *Canidae* species are relatively young, likely moderating reference bias compared to
more divergent taxa. More divergent non-mammalian taxa are likely to experience stronger
reference bias. Since non-mammalian species comprise most of the 99% of species without
reference genomes, this highlights the broader importance of addressing reference bias.
However, karyotype differences occur even between closely related species, making *Canidae*
useful for studying the interplay between karyotype shifts and reference bias. Additionally, our
mapping and filtering parameters, while representing common practices, may influence the
magnitude of reference bias observed. Further analysis is needed to examine how these
choices interact with reference bias and impact other genomic analyses, including inference of
introgression patterns and linkage disequilibrium, which were beyond the scope of this study.
This study does not address reference bias in ancient DNA. Previous research has shown that,
in addition to DNA quality, reference bias is influenced by factors such as sequence length,
genetic divergence[43,44], the alignment tool used[43], and map quality filtering[43,45]. Genetic
summary statistics require careful interpretation in ancient DNA studies, where both human and
non-human extinct lineages will be mapped to divergent interspecific references. The additional
temporal component of ancient DNA will likely amplify the biases observed here.
557
558 *Conclusion*
559
We demonstrate that using a conspecific reference genome enhances the inference of
population size histories and recombination rates, improves the detection of genetic variation,
provides more accurate estimates of nucleotide diversity and genetic differentiation, and
influences the biological interpretation of genomic data. These findings highlight the broader
implications for genomic research in other non-model organisms and stress the need for
continued efforts in generating high-quality, species-specific genomic resources. Considering
these biases, our work demonstrates that results generated from mapping to a highly divergent
reference genome should be interpreted with caution.
568

Acknowledgements

We want to thank Doc Edge, Joshua Schraiber, and Jeffrey Spence for insightful and helpful conversations during this work. J.A.M was supported by the startup funds from Dornsife College of Letters, Arts and Sciences through the Department of Quantitative and Computational Biology and the USC WiSE Gabilan Assistant Professorship. M.G. was supported by USC Provost's Fellowship.

Data accessibility and benefit-sharing

GitHub code for filtering can be found here https://github.com/ellieearmstrong/Gray_Fox_2023/tree/main/filtering and analyses can be found at https://github.com/makopyan/fox/.

Author contributions

M.A. and J.A.M. conceived of the project. J.A.M., M.A., and M.G. performed data processing and analyses. E.A. conducted mapping and generated variant call files used for analyses. J.A.M., M.A., E.A., and M.G. wrote the manuscript. All authors approved of the final manuscript.

STAR☐Methods

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms		
NVIDIA Parabricks v4.1.1.144	O'Connell et al.[46]	https://www.nvidia.com/en-us/clara/genomics/
Mosdepth v0.3.3	Pedersen and Quinlan[47]	https://github.com/brentp/mosdepth
SAMtools v1.16.1	Li and Durbin[31]	http://www.htslib.org/
R v4.2.1/v4.3.2	R Core Team[48]	https://www.r-project.org/
tidyverse	Wickham et al.[49]	https://www.tidyverse.org/
GATK v4.1.4.1	McKenna et al.[50]	https://gatk.broadinstitute.org/
pysam v0.22.1	Li and Durbin[31]	https://github.com/pysam-developers/pysam
bedtools v2.31.1	Quinlan et al.[51]	http://bedtools.readthedocs.io/
BCFtools v1.16	Danecek et al.[52]	http://samtools.github.io/bcftools/
GenMap v1.3.0	Pockrandt et al.[53]	https://github.com/cpockrandt/genmap
TETools v1.7	Lerat et al.[54]	https://github.com/l-modolo/TEtools
GALBA v1.0.9	Brůna et al.[55]	https://github.com/Gaius-Augustus/GALBA
AUGUSTUS v3.3.2	Stanke et al.[56]	https://github.com/Gaius-Augustus/Augustus
SNPRelate v1.36.1	Zheng et al.[57]	https://github.com/zhengxwen/SNPRelate

VCFtools v1.14	Danecek et al.[58]	https://vcftools.github.io/
smc++ v1.15.4	Terhorst et al.[21]	https://github.com/PopSim-LA/smcpp
MSMC2 v2.1.4	Stephan and Ke[59]	https://github.com/stschiff/msmc2
MSMC-tools	Schiffels et al.	https://github.com/stschiff/msmc-tools
Stairway Plot 2 v2.1.2	Liu and Fu[60]	https://github.com/xiaoming-liu/stairway-plot-v2
pyrho v0.1.0	Spence and Song[3]	https://github.com/popgenmethods/pyrho
minimap2 v2.14	Li[61]	https://github.com/lh3/minimap2
Crossmap v0.7.0	Zhao et al.[62]	https://crossmap.sourceforge.net/
GenomicRanges v1.54.1	Lawrence et al.[63]	https://bioconductor.org/packages/GenomicRanges
ComplexUpset v1.3.3	Krassowski et al.[64]	https://github.com/krassowski/complex-upset
Enrichr	Xie et al.[65]	https://maayanlab.cloud/Enrichr/
paf2chain	Guarracino[66]	https://github.com/AndreaGuarracino/paf2chain
SnpEff	Cingolani et al.[67]	http://pcingola.github.io/SnpEff/
bigWigAverageOverBed	Kent et al.[68]	https://genome.ucsc.edu/goldenPath/help/bigWig.html
filterGM.rb	Armstrong and Campana[69]	https://github.com/ellieearmstrong/Gray_Fox_2023
RM2bed.rb	Armstrong and Campana[69]	https://github.com/ellieearmstrong/Gray_Fox_2023
simplify_bed.rb	Armstrong and Campana[69]	https://github.com/ellieearmstrong/Gray_Fox_2023
Deposited Data		
Gray fox whole-genome data	Preckler-Quisquater et al.[12]	PRJNA966176
Gray fox reference genome	Armstrong et al.[14]	GCA_032313775.1
Arctic fox reference genome	Peng et al.[20]	GCF_018345385.1
Domestic dog reference genome (CanFam4)	Wang et al.[19]	GCF_011100685.1
Brown bear genome	Armstrong et al.[70]	GCF_023065955.2
Human genome assembly	Nurk et al.[71]	GCF_009914755.1
PhyloP scores for CanFam4	Meadows et al.[72]	https://doi.org/10.5281/zenodo.8084059

Figure Legends

Figure 1. Reference bias influences variant detection. A) *Sampling localities of gray fox genomes, with a star indicating the origin of the reference genome[14] and circles representing WGS data[12].* ***B)*** *Phylogenetic relationships of three canid reference genomes: Arctic fox (Vulpes lagopus, blue), domestic dog (Canis lupus familiaris, red), and gray fox (Urocyon cinereoargenteus, gold, most basal). Diploid chromosome numbers (2n) shown for each species.* ***C-F)*** *SNP counts and liftover classifications for eastern (C, D) and western (E, F) populations. Bar charts show total SNP counts (millions, above bars) divided into singleton (light) and non-singleton (dark) variants. Alluvial diagrams track variant reclassification when lifting over from heterospecific to gray fox reference genomes, showing transitions between singleton, SNP, invariant ("not snp"), and unmapped ("not map") categories. Percentages show variant proportions and transition flows. Species-matched reference genomes yield higher effective population size estimates.*

Figure 2. Reference bias influences the SFS and demographic trajectories. A) *SFS for eastern and western populations show more singletons using the conspecific gray fox genome.* ***B)*** *Effective population sizes (y-axis) and years from present (x-axis) inferred with smc++ reveal discordant demographic histories of eastern (light) and western (dark) foxes resolved using the species-matched (gold) and heterospecific genomes (red and blue).*

Figure 3. Recombination rate comparisons between reference genomes for gray fox populations. *Histograms of the square root of recombination rates (cM/Mb) across 50 kb windows in (A) east and (B) west populations. Colors represent the three reference genomes used for recombination rate estimates, with vertical lines representing mean values. Loess-smoothed (span=0.1) recombination rates (cm/Mb) per chromosome computed over 50 kb windows in Eastern (C) and Western (D) gray foxes based on the three reference genomes.*

Figure 4. Estimates of diversity and differentiation are higher with gray fox reference in eastern and western gray fox populations. *Nucleotide diversity (π) for eastern (A) and western (B) gray fox populations, and genetic differentiation (F_{ST}) between the populations (C), in 50-kb windows based on three reference genomes. Violin plots with mean estimates and standard deviation error bars are shown. The correlation between diversity and differentiation across genomic windows for the east (D) and west (E) populations is illustrated. Kendall's tau (τ) correlation coefficients are reported, showing significant negative correlations between π and F_{ST} in both populations across all reference genomes.*

Figure 5. Reference-specific patterns revealed by F_{ST} outlier and functional enrichment analyses. *(A) Scatter plot comparing F_{ST} values between gray fox and arctic fox references, with outlier windows unique to the arctic fox (blue) and gray fox (gold) highlighted. (B) Scatter plot comparing F_{ST} values between gray fox and Canfam4 references, highlighting outliers unique to Canfam4 (red) and gray fox (gold). For both (A) and (B), outliers shared between the two references are shown in black, while non-outlier windows are in gray. (C) UpSet plot of significant (p<0.05) GO biological process terms identified as enriched for each reference genome. Bars represent the intersection size (y-axis) of GO terms shared across reference genomes (x-axis).*

Tables

Table 1. *The total number of successfully mapped reads, number of SNPs, and heterozygosity estimates (mean ± standard deviation) for each reference genome and population. The total number of reads (both mapped and unmapped) was 3,886,389,246 for the East and 1,043,628,176 for the West population.*

Genome	Population	Reads Mapped	Total SNPs	Heterozygosity (mean ± sd)
Arctic fox	East	3,836,962,124	5,407,943	0.181 ± 0.057
CanFam4	East	3,848,140,971	5,392,973	0.181 ± 0.058
Gray fox	East	3,865,958,291	6,828,177	0.200 ± 0.051
Arctic fox	West	1032573693	10,012,122	0.163 ± 0.049
CanFam4	West	1035433543	9,969,686	0.163 ± 0.049
Gray fox	West	1040317760	13,199,882	0.174 ± 0.047

Table 2. *Pyrho hyperparameter settings for each reference genome and both populations.*

	Block Penalty		Window Size (bp)	
	East	West	East	West
Gray fox	25	25	50	25
Canfam4	25	50	50	50
Arctic fox	50	25	150	50

References

1. Kosugi, S., Momozawa, Y., Liu, X., Terao, C., Kubo, M., and Kamatani, Y. (2019). Comprehensive evaluation of structural variation detection algorithms for whole genome sequencing. Genome Biol. 20, 117. https://doi.org/10.1186/s13059-019-1720-5.

2. Salmona, J., Heller, R., Lascoux, M., and Shafer, A. (2017). Population genomics, concepts, approaches and applications. Popul. Genom., 511–537. https://doi.org/10.1007/13836_2017_1.

3. Spence, J.P., and Song, Y.S. (2019). Inference and analysis of population-specific fine-scale recombination maps across 26 diverse human populations. Sci. Adv. 5, eaaw9206. https://doi.org/10.1126/sciadv.aaw9206.

4. Nevado, B., Ramos☐Onsins, S.E., and Perez☐Enciso, M. (2014). Resequencing studies of nonmodel organisms using closely related reference genomes: optimal experimental designs and bioinformatics approaches for population genomics. Mol Ecol 23, 1764–1779. https://doi.org/10.1111/mec.12693.

5. Gopalakrishnan, S., Castruita, J.A.S., Sinding, M.-H.S., Kuderna, L.F.K., Räikkönen, J., Petersen, B., Sicheritz-Ponten, T., Larson, G., Orlando, L., Marques-Bonet, T., et al. (2017). The wolf reference genome sequence (Canis lupus lupus) and its implications for Canis spp. population genomics. BMC Genom. 18, 495. https://doi.org/10.1186/s12864-017-3883-3.

6. Armstrong, E.E., Taylor, R.W., Miller, D.E., Kaelin, C.B., Barsh, G.S., Hadly, E.A., and Petrov, D. (2020). Long live the king: chromosome-level assembly of the lion (Panthera leo) using linked-read, Hi-C, and long-read data. BMC Biol. 18, 3. https://doi.org/10.1186/s12915-019-0734-5.

7. Thorburn, D.J., Sagonas, K., Binzer☐Panchal, M., Chain, F.J.J., Feulner, P.G.D., Bornberg☐Bauer, E., Reusch, T.B.H., Samonte☐Padilla, I.E., Milinski, M., Lenz, T.L., et al. (2023). Origin matters: Using a local reference genome improves measures in population genomics. Mol. Ecol. Resour. 23, 1706–1723. https://doi.org/10.1111/1755-0998.13838.

8. Maurstad, M.F., Hoff, S.N.K., Cerca, J., Ravinet, M., Bradbury, I., Jakobsen, K.S., Præbel, K., and Jentoft, S. (2024). Reference genome bias in light of species-specific chromosomal reorganization and translocations. bioRxiv, 2024.06.28.599671. https://doi.org/10.1101/2024.06.28.599671.

9. Rick, J.A., Brock, C.D., Lewanski, A.L., Golcher-Benavides, J., and Wagner, C.E. (2023). Reference genome choice and filtering thresholds jointly influence phylogenomic analyses. Syst. Biol. 73, 76–101. https://doi.org/10.1093/sysbio/syad065.

10. Prasad, A., Lorenzen, E.D., and Westbury, M.V. (2022). Evaluating the role of reference☐genome phylogenetic distance on evolutionary inference. Mol. Ecol. Resour. 22, 45–55. https://doi.org/10.1111/1755-0998.13457.

11. Goddard, N.S., Statham, M.J., and Sacks, B.N. (2015). Mitochondrial analysis of the most basal canid reveals deep divergence between eastern and western North American gray foxes (Urocyon spp.) and ancient roots in Pleistocene California. PLoS ONE 10, e0136329. https://doi.org/10.1371/journal.pone.0136329.

12. Preckler☐Quisquater, S., Kierepka, E.M., Reding, D.M., Piaggio, A.J., and Sacks, B.N. (2023). Can demographic histories explain long☐term isolation and recent pulses of asymmetric gene flow between highly divergent grey fox lineages? Mol. Ecol. 32, 5323–5337. https://doi.org/10.1111/mec.17105.

13. Wayne, R.K., Nash, W.G., and O'Brien, S.J. (1987). Chromosomal evolution of the Canidae.
Cytogenet. Genome Res. *44*, 123–133. https://doi.org/10.1159/000132356.

14. Armstrong, E.E., Bissell, K.L., Fatima, H.S., Heikkinen, M.A., Jessup, A., Junaid, M.O., Lee,
D.H., Lieb, E.C., Liem, J.T., Martin, E.M., et al. (2024). Chromosome-level assembly of the gray
fox (Urocyon cinereoargenteus) confirms the basal loss of PRDM9 in Canidae. G3: Genes,
Genomes, Genet. 14, jkae034. https://doi.org/10.1093/g3journal/jkae034.

15. Lindblad-Toh, K., Wade, C.M., Mikkelsen, T.S., Karlsson, E.K., Jaffe, D.B., Kamal, M.,
Clamp, M., Chang, J.L., Kulbokas, E.J., Zody, M.C., et al. (2005). Genome sequence,
comparative analysis and haplotype structure of the domestic dog. Nature *438*, 803–819.
https://doi.org/10.1038/nature04338.

16. Robinson, J.A., Vecchyo, D.O.-D., Fan, Z., Kim, B.Y., vonHoldt, B.M., Marsden, C.D.,
Lohmueller, K.E., and Wayne, R.K. (2016). Genomic flatlining in the endangered island fox. Curr
Biol 26, 1183–1189. https://doi.org/10.1016/j.cub.2016.02.062.

17. Robinson, J.A., Brown, C., Kim, B.Y., Lohmueller, K.E., and Wayne, R.K. (2018). Purging of
strongly deleterious mutations explains long-term persistence and absence of inbreeding
depression in island foxes. Curr. Biol. 28, 3487-3494.e4.
https://doi.org/10.1016/j.cub.2018.08.066.

18. Mooney, J.A., Marsden, C.D., Yohannes, A., Wayne, R.K., and Lohmueller, K.E. (2022).
Long-term small population size, deleterious variation, and altitude adaptation in the Ethiopian
wolf, a severely endangered canid. Mol. Biol. Evol. *40*, msac*277*.
https://doi.org/10.1093/molbev/msac277.

19. Wang, C., Wallerman, O., Arendt, M.-L., Sundström, E., Karlsson, Å., Nordin, J.,
Mäkeläinen, S., Pielberg, G.R., Hanson, J., Ohlsson, Å., et al. (202*1*). A novel canine reference
genome resolves genomic architecture and uncovers transcript complexity. Commun. Biol. 4,
185. https://doi.org/10.1038/s42003-021-01698-x.

20. Peng, Y., Li, H., Liu, Z., Zhang, C., Li, K., Gong, Y., Geng, L., Su, J., Guan, X., Liu, L., et al.
(2021). Chromosome level genome assembly of the Arctic fox (Vulpes lagopus) using PacBio
sequencing and Hi C technology. Mol. Ecol. Resour. 21, 2093–2108.
https://doi.org/10.1111/1755-0998.13*397*.

21. Terhorst, J., Kamm, J.A., and Song, Y.S. (2017). Robust and scalable inference of
population history from hundreds of unphased whole genomes. Nat. Genet. 49, 303–309.
https://doi.org/10.1038/ng.3748.

22. Lewin, H.A., Richards, S., Aiden, E.L., Allende, M.L., Archibald, J.M., Bálint, M., Barker,
K.B., Baumgartner, B., Belov, K., Bertorelle, G., et al. (2022). The Earth BioGenome Project
2020: Starting the clock. Proc. Natl. Acad. Sci. *119*, e2115635118.
https://doi.org/10.1073/pnas.2115635118.

23. Galla, S.J., Forsdick, N.J., Brown, L., Hoeppner, M., Knapp, M., Maloney, R.F., Moraga, R.,
Santure, A.W., and Steeves, T.E. (2018). Reference genomes from distantly related species
can be used for discovery of single nucleotide polymorphisms to inform conservation
management. Genes 10, 9. https://doi.org/10.3390/genes10010009.

24. Minias, P., Dunn, P.O., Whittingham, L.A., Johnson, J.A., and Oyler-McCance, S.J. (2019).
Evaluation of a chicken 600K SNP genotyping array in non-model species of grouse. Sci. Rep.
9, 6407. https://doi.org/10.1038/s41598-019-42885-5.

25. Valiente-Mullor, C., Beamud, B., Ansari, I., Francés-Cuesta, C., García-González, N., Mejía,

L., Ruiz-Hueso, P., and González-Candelas, F. (2021). One is not enough: On the effects of
reference genome for the mapping and subsequent analyses of short-reads. PLoS Comput.
Biol. 17, e1008678. https://doi.org/10.1371/journal.pcbi.1008678.
26. Yang, X., Lee, W.-P., Ye, K., and Lee, C. (2019). One reference genome is not enough.
Genome Biol. 20, 104. https://doi.org/10.1186/s13059-019-1717-0.
27. Miga, K.H., and Wang, T. (2021). The Need for a Human Pangenome Reference Sequence.
Annu. Rev. Genom. Hum. Genet. 22, 1–22. https://doi.org/10.1146/annurev-genom-120120-
081921.
28. Wu, X., Heffelfinger, C., Zhao, H., and Dellaporta, S.L. (2019). Benchmarking variant
identification tools for plant diversity discovery. BMC Genom. 20, 701.
https://doi.org/10.1186/s12864-019-6057-7.
29. Hemstrom, W., Grummer, J.A., Luikart, G., and Christie, M.R. (2024). Next-generation data
filtering in the genomics era. Nat. Rev. Genet. 25, 750–767. https://doi.org/10.1038/s41576-024-
00738-6.
30. Chavez, D.E., Gronau, I., Hains, T., Dikow, R.B., Frandsen, P.B., Figueiró, H.V., Garcez,
F.S., Tchaicka, L., Paula, R.C. de, Rodrigues, F.H.G., et al. (2022). Comparative genomics
uncovers the evolutionary history, demography, and molecular adaptations of South American
canids. Proc. Natl. Acad. Sci. 119, e2205986119. https://doi.org/10.1073/pnas.2205986119.
31. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler
transform. Bioinformatics 25, 1754–1760. https://doi.org/10.1093/bioinformatics/btp324.
32. Wilder, A.P., Supple, M.A., Subramanian, A., Mudide, A., Swofford, R., Serres-Armero, A.,
Steiner, C., Koepfli, K.-P., Genereux, D.P., Karlsson, E.K., et al. (2023). The contribution of
historical processes to contemporary extinction risk in placental mammals. Science *380*,
eabn5856–eabn5856. https://doi.org/10.1126/science.abn5856.
33. Paez, S., Kraus, R.H.S., Shapiro, B., Gilbert, M.T.P., Jarvis, E.D., Group, V.G.P.C., Al-Ajli,
F.O., Ceballos, G., Crawford, A.J., Fedrigo, O., et al. (2022). Reference genomes for
conservation. Science *377*, 364–366. https://doi.org/10.1126/science.abm8127.
34. Lin, M.-J., Iyer, S., Chen, N.-C., and Langmead, B. (2024). Measuring, visualizing, and
diagnosing reference bias with biastools. Genome Biol. 25, 101. https://doi.org/10.1186/s13059-
024-03240-8.
35. Chen, N.-C., Solomon, B., Mun, T., Iyer, S., and Langmead, B. (2021). Reference flow:
reducing reference bias using multiple population genomes. Genome Biol. 22, 8.
https://doi.org/10.1186/s13059-020-02229-3.
36. Campbell, C.L., Bhérer, C., Morrow, B.E., Boyko, A.R., and Auton, A. (2016). A pedigree-
based map of recombination in the domestic dog genome. G3: Genes, Genomes, Genet. 6,
3517–3524. https://doi.org/10.1534/g3.116.034678.
37. Axelsson, E., Webster, M.T., Ratnakumar, A., Consortium, T.L., Ponting, C.P., and
Lindblad-Toh, K. (2012). Death of PRDM9 coincides with stabilization of the recombination
landscape in the dog genome. Genome Res. 22, 51–63. https://doi.org/10.1101/gr.124123.111.
38. Batra, S.S., Levy-Sakin, M., Robinson, J., Guillory, J., Durinck, S., Vilgalys, T.P., Kwok, P.-
Y., Cox, L.A., Seshagiri, S., Song, Y.S., et al. (2020). Accurate assembly of the olive baboon
(Papio anubis) genome using long-read and Hi-C data. GigaScience 9, giaa134.
https://doi.org/10.1093/gigascience/giaa134.
39. Gopalakrishnan, S., Sinding, M.-H.S., Ramos-Madrigal, J., Niemann, J., Castruita, J.A.S.,

Vieira, F.G., Carøe, C., Montero, M. de M., Kuderna, L., Serres, A., et al. (2018). Interspecific
Gene Flow Shaped the Evolution of the Genus Canis. Curr. Biol. 28, 3441-3449.e5.
https://doi.org/10.1016/j.cub.2018.08.041.
40. Funk, W.C., Forester, B.R., Converse, S.J., Darst, C., and Morey, S. (2019). Improving
conservation policy with genomics: a guide to integrating adaptive potential into U.S.
Endangered Species Act decisions for conservation practitioners and geneticists. Conserv.
Genet. 20, 115–134. https://doi.org/10.1007/s10592-018-1096-1.
41. Hoffmann, A., Griffin, P., Dillon, S., Catullo, R., Rane, R., Byrne, M., Jordan, R., Oakeshott,
J., Weeks, A., Joseph, L., et al. (2015). A framework for incorporating evolutionary genomics
into biodiversity conservation and management. Clim. Chang. Responses 2, 1.
https://doi.org/10.1186/s40665-014-0009-x.
42. Brandt, D.Y.C., Aguiar, V.R.C., Bitarello, B.D., Nunes, K., Goudet, J., and Meyer, D. (2015).
Mapping bias overestimates reference allele frequencies at the HLA genes in the 1000
Genomes Project phase I data. G3: Genes, Genomes, Genet. 5, 931–941.
https://doi.org/10.1534/g3.114.015784.
43. Dolenz, S., Valk, T. van der, Jin, C., Oppenheimer, J., Sharif, M.B., Orlando, L., Shapiro, B.,
Dalén, L., and Heintzman, P.D. (2024). Unravelling reference bias in ancient DNA datasets.
Bioinformatics 40, btae436. https://doi.org/10.1093/bioinformatics/btae436.
44. Swiel, Y., Kelso, J., and Peyrégne, S. (2024). Resolving the source of branch length
variation in the Y chromosome phylogeny. bioRxiv, 2024.07.05.602100.
https://doi.org/10.1101/2024.07.05.602100.
45. Günther, T., and Nettelblad, C. (2019). The presence and impact of reference bias on
population genomic studies of prehistoric human populations. PLoS Genet. 15, e1008302.
https://doi.org/10.1371/journal.pgen.1008302.
46. O'Connell, K.A., Yosufzai, Z.B., Campbell, R.A., Lobb, C.J., Engelken, H.T., Gorrell, L.M.,
Carlson, T.B., Catana, J.J., Mikdadi, D., Bonazzi, V.R., et al. (2023). Accelerating genomic
workflows using NVIDIA Parabricks. BMC Bioinform. 24, 221. https://doi.org/10.1186/s12859-
023-05292-2.
47. Pedersen, B.S., and Quinlan, A.R. (2017). Mosdepth: quick coverage calculation for
genomes and exomes. Bioinformatics 34, 867–868.
https://doi.org/10.1093/bioinformatics/btx699.
48. Team, R.C. (2020). R: 2019. A Language and Environment for Statistical Computing version
3.
49. Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the Tidyverse. J. Open Source Softw.
4, 1686. https://doi.org/10.21105/joss.01686.
50. McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella,
K., Altshuler, D., Gabriel, S., Daly, M., et al. (2010). The genome analysis toolkit: A MapReduce
framework for analyzing next-generation DNA sequencing data. Genome Res 20, 1297–1303.
https://doi.org/10.1101/gr.107524.110.
51. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics 26, 841–842. https://doi.org/10.1093/bioinformatics/btq033.
52. Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of SAMtools and BCFtools.

GigaScience 10, giab008. https://doi.org/10.1093/gigascience/giab008.

53. Pockrandt, C., Alzamel, M., Iliopoulos, C.S., and Reinert, K. (2020). GenMap: ultra-fast

computation of genome mappability. Bioinformatics 36, 3687–3692.

https://doi.org/10.1093/bioinformatics/btaa222.

54. Lerat, E., Fablet, M., Modolo, L., Lopez-Maestre, H., and Vieira, C. (2017). TEtools

facilitates big data expression analysis of transposable elements and reveals an antagonism

between their activity and that of piRNA genes. Nucleic Acids Res. 45, e17–e17.

https://doi.org/10.1093/nar/gkw953.

55. Brůna, T., Li, H., Guhlin, J., Honsel, D., Herbold, S., Stanke, M., Nenasheva, N., Ebel, M.,

Gabriel, L., and Hoff, K.J. (2023). Galba: genome annotation with miniprot and AUGUSTUS.

BMC Bioinform. 24, 327. https://doi.org/10.1186/s12859-023-05449-z.

56. Stanke, M., Diekhans, M., Baertsch, R., and Haussler, D. (2008). Using native and

syntenically mapped cDNA alignments to improve de novo gene finding. Bioinformatics 24,

637–644. https://doi.org/10.1093/bioinformatics/btn013.

57. Zheng, X., Levine, D., Shen, J., Gogarten, S.M., Laurie, C., and Weir, B.S. (2012). A high-

performance computing toolset for relatedness and principal component analysis of SNP data.

Bioinformatics 28, 3326–3328. https://doi.org/10.1093/bioinformatics/bts606.

58. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker,

R.E., Lunter, G., Marth, G.T., Sherry, S.T., et al. (2011). The variant call format and VCFtools.

Bioinformatics 27, 2156–2158. https://doi.org/10.1093/bioinformatics/btr330.

59. Stephan, S., and Ke, W. (2020). MSMC and MSMC2: The multiple sequentially Markovian

coalescent. In Methods in Molecular Biology. (Springer US), pp. 147–166.

https://doi.org/10.1007/978-1-0716-0199-0_7.

60. Liu, X., and Fu, Y.-X. (2020). Stairway Plot 2: demographic history inference with folded

SNP frequency spectra. Genome Biol. 21, 280. https://doi.org/10.1186/s13059-020-02196-9.

61. Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34,

3094–3100. https://doi.org/10.1093/bioinformatics/bty191.

62. Zhao, H., Sun, Z., Wang, J., Huang, H., Kocher, J.-P., and Wang, L. (2014). CrossMap: a

versatile tool for coordinate conversion between genome assemblies. Bioinformatics 30, 1006–

1007. https://doi.org/10.1093/bioinformatics/btt730.

63. Lawrence, M., Huber, W., Pagès, H., Aboyoun, P., Carlson, M., Gentleman, R., Morgan,

M.T., and Carey, V.J. (2013). Software for computing and annotating genomic ranges. PLoS

Comput. Biol. 9, e1003118. https://doi.org/10.1371/journal.pcbi.1003118.

64. Krassowski, M., Arts, M., Heeg, M., and Lagger, C. (2022). ComplexUpset.

65. Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L.,

Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene set

knowledge discovery with Enrichr. Curr. Protoc. 1, e90. https://doi.org/10.1002/cpz1.90.

66. Guarracino, A. (2024). paf2chain.

67. Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X., and

Ruden, D.M. (2012). A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff. Fly 6, 80–92. https://doi.org/10.4161/fly.19695.

68. Kent, W.J., Zweig, A.S., Barber, G., Hinrichs, A.S., and Karolchik, D. (2010). BigWig and

BigBed: enabling browsing of large distributed datasets. Bioinformatics 26, 2204–2207.

https://doi.org/10.1093/bioinformatics/btq351.

69. Armstrong, E.E., and Campana, M.G. (2022). RatesTools: a Nextflow pipeline for detecting
de novo germline mutations in pedigree sequence data. Bioinformatics 39, btac784.
https://doi.org/10.1093/bioinformatics/btac784.
70. Armstrong, E.E., Perry, B.W., Huang, Y., Garimella, K.V., Jansen, H.T., Robbins, C.T.,
Tucker, N.R., and Kelley, J.L. (2022). A beary good genome: Haplotype-resolved, chromosome-
level assembly of the brown bear (Ursus arctos). Genome Biol. Evol. 14, evac125.
https://doi.org/10.1093/gbe/evac125.
71. Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A.V., Mikheenko, A., Vollger, M.R.,
Altemose, N., Uralsky, L., Gershman, A., et al. (2022). The complete sequence of a human
genome. Science *376*, 44–53. https://doi.org/10.1126/science.abj6987.
72. Meadows, J.R.S., Kidd, J.M., Wang, G.-D., Parker, H.G., Schall, P.Z., Bianchi, M.,
Christmas, M.J., Bougiouri, K., Buckley, R.M., Hitte, C., et al. (2023). Genome sequencing of
2000 canids by the Dog10K consortium advances the understanding of demography, genome
function and architecture. Genome Biol. 24, 187. https://doi.org/10.1186/s13059-023-03023-7.

A) Diversity(π) in East B) Diversity(π) in West C) F_{ST} East v West

Legend: east, west, arcticfox, canfam4, grayfox

D)
East — $\tau=-0.173$, $p<0.0001$
East — $\tau=-0.160$, $p<0.0001$
East — $\tau=-0.196$, $p<0.0001$

E)
West — $\tau=-0.095$, $p<0.0001$
West — $\tau=-0.081$, $p<0.0001$
West — $\tau=-0.118$, $p<0.0001$

F_{ST} East v West

Nucleotide diversity (π)