Decontaminating genomic data for accurate species delineation and hybrid detection in the *Lasius* ant genus

Kristine Jecha[1]*, Guillaume Lavanchy[1,2]*, Tanja Schwander[1]

[1]Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland.

[2]Present address: Biodiversity and Evolution Unit, Department of Biology, Lund University. Ekologihuset, Sölvegatan 37, 223 62 Lund, Sweden

*equal contribution

corresponding authors: kristine.jecha@unil.ch; guillaume.lavanchy@gmail.com

Abstract

Species identification and delineation by molecular methods has become a widely used technique and has revealed hybrids between species previously believed to be completely reproductively isolated. However, the application of molecular methods is associated with a risk of DNA contamination, which can result in the identification of false hybrids and generate inaccurate conclusions about the species' identities and characteristics. Here, we generate and analyze a dataset of nuclear SNP data and mitochondrial DNA sequences from over 1,000 *Lasius* ants to accurately investigate the species delineation and hybridization proclivity within

the genus. We describe an approach, based on a combination of competitive mapping and

allelic depth ratio analysis, that allows us to identify DNA contaminations and filter them from

large-scale datasets. By applying this approach to the *Lasius* ants, we are able to remove

interspecific contamination, and to clearly delineate each species genetically as well as identify

a hybrid individual between *L. emarginatus* and *L. platythorax*.

Keywords: competitive mapping, DNA barcoding, hybridization, introgression, RAD

sequencing, Formicidae

Significance

Next-generation sequencing enables large-scale studies of population structure, species

delineation, and introgression for a wide range of taxa. However, the potential for cross-

contamination during sample preparation is rarely considered, despite potential major impacts

on the conclusions. Here, we develop a new decontamination pipeline and apply it to a large-

scale genotyping dataset to study species delineation and hybridization in the ant genus *Lasius*.

We find only one hybrid in a thousand individuals, a very different picture from the contaminated

dataset. Testing for and removing contamination should be part of every population genomics

study.

Introduction

The most widely accepted definition of a species refers to groups of populations that are

reproductively isolated from others (Mayr, 1942). However, in practical applications, species

distinctions rely on morphological criteria without any reproductive behavior information. Yet,

even high-quality morphological identification brings its own practical challenges by being time-

consuming and often requiring experienced taxonomists. These issues can be particularly

apparent in sampling studies in which the identification of many difficult-to-distinguish species is

required for robust analysis. The disconnect between the concept and practical implementation

of the definition of a species underscores the need for a critical examination of widely used

methodologies and increased implementation of more precise ones when possible.

Molecular methods of identification, often referred to as "DNA barcoding", are used to

identify distinct species based on genetic divergence at specific genetic regions (Hebert et al,

2003). The mitochondrial gene cytochrome c oxidase I (COI) has become the standard for

species identification via DNA barcoding as it evolves rapidly enough to clearly distinguish

between species and even diverged populations of the same species, but not too fast as to still

remain reliable in grouping conspecific individuals (Cox & Hebert, 2001). However, relying solely

on a mitochondrial gene can introduce limitations as it is maternally transmitted and therefore

will not reveal any history of hybridization between species. Hybridization has recently become

a rich area of research as it was believed to be a very rare occurrence but is, in fact, common

and widespread across many taxa (Edelman & Mallet, 2021). Analysis of nuclear DNA can not

only identify hybrids but also assess the extent of introgression between the parental species

within each population, providing insight into the hybridization patterns of the species pair. Yet,

the use of these more advanced techniques can introduce a new problem: the risk of sequence

data contamination. This can further obfuscate species delineations and lead to

misinterpretations of hybrid status.

Because of the intricacies associated with methods of species identification, there is a

need for novel techniques to decontaminate complex data and benchmark tests to ensure that

they are precise and accurate. Investigating the ant genus *Lasius,* comprising 86 different

species within the Holarctic, provides an ideal case study (Wilson, 1955; Janda et al., 2004).

These species have been notoriously difficult to distinguish with many cases of cryptic species

that are morphologically similar and cases of misclassifications later leading to synonymizing

(Agosti & Bolton, 1990; Seifert, 1991; Seifert 1992; Wilson, 1955). Additionally, while there have

been several instances of hybridization between *Lasius* species, a complete investigation into

the patterns of hybridization and introgression between the various species within the genus has

never been performed. A clear molecular identification technique would allow these species to

be quickly distinguished and eliminate the risk of mistakenly analyzing the incorrect species.

With a dataset of over 1,000 *Lasius* ants from the Swiss canton of Vaud, we were able to

examine the accuracy of a combination of morphological as well as nuclear and mitochondrial

molecular identification techniques: restriction site-associated DNA sequencing (RADseq) for

genotyping and COI gene analysis. In addition, we were able to use the data to further analyze

these species for cryptic species and hybrid individuals. However, apparent contamination of

the sequenced nuclear DNA required us to develop pipelines to decontaminate our data to

varying degrees in order to ensure accurate conclusions on the evolutionary dynamics of this

genus. This illustrates the necessity for rigorous quality control and decontamination to be

performed when using these types of molecular techniques.

Methods

Study area and sampling

We collected samples of *Lasius* as part of the *Opération Fourmis* citizen sampling

campaign (Freitag et al., 2020; Avril et al., 2019) as well as through a structured sampling

process throughout the Vaud canton of Switzerland in 2019 (Szewczyk et al., 2024). About 10

workers from each colony were collected and stored in 70% EtOH until DNA extraction. In total

3,173 *Lasius* worker ants were collected, 434 from the structured sampling, and 2,739 from

citizen sampling (Figure 1). These samples were identified morphologically by expert

taxonomists to the species level when possible following Seifert (2018).

<u>Figure 1.</u> Map of *Lasius* colony sample locations within the canton of Vaud in

Switzerland (n =3,138). Black points represent ants collected by the citizen science

Opération Fourmis project, red points represent ants collected via structured scientific

sampling. 35 samples missing geographical data and not included on the map.

DNA extraction and sequencing

For each colony, we selected one worker ant at random to undergo genetic sequencing.

For each worker, three legs were used for DNA extraction. We genotyped up to 200 individuals

for each (morphologically identified) species. For species that had more than 200 colonies

collected in the field, we downsampled those from the citizen science campaign that were near

each other in order to ensure geographically widespread sampling, resulting in a total of 1,171

genotyped individuals. The DNA extraction and COI barcoding procedures were outsourced to

the Canadian Center for DNA Barcoding (CCDB) (Guelph, Canada) for 790 samples and to

AllGenetics (A Coruña, Spain) for the remaining 381. A fragment of the COI gene was

successfully amplified using primers LepF1 and LepR1 for 1,111 samples (Hebert et al., 2004).

The specific protocols used for extraction, PCR amplification, and sequencing are available at

the CCDB's website https://ccdb.ca/resources (last checked Nov 17, 2023). We generated

nuclear DNA sequences using RAD sequencing protocols described in Brelsford et al. (2016).

We used restriction enzymes EcoRI and MseI to digest the genomic DNA followed by ligating

custom barcode adaptors to the DNA fragments, which were then amplified in 16 PCR cycles.

Fragments with a size of 280 to 430 bp were selected using 2% agarose cassettes on a

BluePippin (Sage Science). We sequenced these libraries on eight Illumina HiSeq 2500 lanes

(single end, 150 bases) to generate the final read data.

SNP calling

We then demultiplexed the sequenced reads using stacks v2.53 process_radtags

command with the options -c -q -r -t 143 (Catchen et al., 2013) and mapped the reads against a

previously generated reference genome of *Lasius niger* (BioProject: PRJNA1159026) with the

mem algorithm of BWA v0.7.17 (Li & Durbin, 2010). Alignment file formatting and summaries

were created using samtools v1.15.1 (Danecek et al., 2021). We built loci using gstacks with

options --phasing-dont-prune-hets and --ignore-pe-reads and ran populations with the option --

write-single-snp to retain only one SNP per RAD locus. We were then able to filter the data with

vcftools 0.1.14 (Danecek et al., 2011) keeping only genotype calls with a minimum depth of 8

and loci with a minor allele count of two which were present in at least 75% of individuals. Any

samples missing more than 50% of the genotyped loci were excluded. As a quality control, we

looked at the relationship between the proportion of missing genotypes and "relative

heterozygosity" (heterozygosity based on polymorphic loci) per individual. We measured relative

heterozygosity as the proportion of heterozygous alleles to total genotyped SNPs.

Competitive mapping

In order to decontaminate the RADseq reads, we used a competitive mapping technique

based on Feuerborn et al.'s (2020) study in which they used competitive mapping to remove

modern human contaminant DNA from several ancient faunal genomic datasets. This consisted

of concatenating the reference genomes of species representing potential contaminants to the

target reference genome, mapping the RADseq reads to the concatenated genomes, and

removing reads mapping to the non-target genomes. In our case, we combined the *Lasius niger*

genome with ant genomes from seven genera to represent other ant genera also found in Vaud,

Switzerland as part of the *Opération fourmis* sampling (Szewczyk et al., 2024). These included

Camponotus, *Formica*, *Leptothorax*, *Myrmica*, *Tapinoma*, *Temnothorax*, and *Tetramorium*

(Romiguier et al., 2022 and BioProjects: PRJNA445978, PRJNA819254, and PRJNA533534).

We mapped the reads from each sequenced *Lasius* individual to this multi-genus concatenated

genome with the mem algorithm of BWA v0.7.17 (Li & Durbin, 2010). We could then separate

the reads into those preferentially mapping to the target *Lasius* genome and those that were

mapping to a possible contaminant genome using R v4.2.1 (R Core Team, 2021). To filter out

possible contaminants, we kept only reads that mapped either primarily or secondarily to the

Lasius genome using the "filterbyname.sh" function of BBMap v38.63 (Bushnell, 2014). We then

remapped these selected reads to the *Lasius niger* reference genome and performed SNP

calling as previously stated. Any samples missing more than 50% of genotypes were excluded.

154   Allelic depth ratio filtering

Putative contaminations between samples from the same genus cannot be removed by

our competitive mapping approach. To overcome that, we used an additional, complementary

approach based on allelic depth ratio. In this paper, we define allelic depth ratio (ADR) at a

heterozygous locus as the ratio of reads of the allele with most reads to the sum of all reads at

the locus. It can thus range from 0.5 to <1. For a real heterozygote in a diploid individual, it is

expected to be around 0.5 in all single-copy loci (Figure 2). However, if a heterozygous

genotype is caused by contamination, and the amount of contaminant DNA is expected to be

lower than the original sample's DNA, and ADR will be higher than 0.5. We used ADR-based

filtering in two steps. First, we discarded individuals with a highly skewed ADR distribution, as

these most likely represent highly contaminated samples. We estimated the expected

distribution of ADR by sampling from a binomial distribution with locus depth as the number of

observations for all heterozygous sites in all individuals. We then considered that individuals

whose observed median ADR (across all their heterozygous loci) was beyond the 5% percentile

of this expected ADR distribution were highly contaminated and were discarded (Figure 2G).

Second, in retained individuals, we corrected the individual genotypes with ADR beyond

the 25% percentile of the expected ADR distribution. We assumed that the allele with the fewest

reads was derived from a contaminant and considered the individual homozygous for the other

allele (Figure 2H). This filtering was performed using a custom script in R v4.2.1 (R Core Team,

2021), using the package vcfR v1.14.0 (Knaus & Grünwald, 2017).

Figure 2: Overview of the decontamination pipeline. **A** Competitive Mapping. Reads are mapped simultaneously to the focal genome assembly and assemblies of putative contaminant taxa. **B** Typical dataset after SNP calling. Blue and yellow are two alleles, grey is missing data. **B-G** Allelic Depth Ratio filtering approach. **C** Allelic Depth Ratio (ADR) is expected to be around 0.5 for a real heterozygous. **D** Contaminations are expected to lead to ADR higher than 0.5. **E**

180   We first estimate ADR distribution across all heterozygous loci for each individual

181   independently. **F** We then estimate the expected ADR distribution across all heterozygous loci

182   given observed depths. **G** We discard individuals whose median ADR is beyond the top 5%

183   percentile of the expected ADR distribution. **H** We correct genotypes with ADR beyond the top

184   25% percentile of the expected ADR distribution by removing the allele with the fewest reads.

Haploid allele filtering

As an alternative way to remove contaminants and assess the overall robustness of

our findings, we artificially "haploidized" our competitively mapped dataset. At all heterozygous

sites, we artificially removed the allele with the fewest reads. When there was the same number

of reads at both alleles, one was drawn at random. We also considered homozygous sites

haploid. This filtering was performed using a custom script in R v4.2.1 (R Core Team, 2021),

using the package vcfR v1.14.0 (Knaus & Grünwald, 2017).

Molecular-based species identification

To investigate the congruence between the morphological identification and SNP-based

identification, we used the SNP calls to create a MultiDimensional Scaling (MDS) plot with the R

package MASS (Venables & Ripley, 2002) with R v4.2.1 (R Core Team, 2021). In conjunction

with the MDS clustering, we ran ADMIXTURE (Alexander et al., 2009) in 10 replicates for K

values ranging from 1 to 15. We then selected the best value of K and the best replicate based

on Cross-Validation error. We conducted the same analysis in haploid mode with the option --

haploid="*" for the "haploidized" dataset. We then visualized the results with pong (Behr et al.,

2016) and chose the most representative run. In cases where the morphological identification

was ambiguous or different from the genetic identification, the individual was reassigned to its

genetic species identification. We were able to directly identify hybrid individuals using the

ADMIXTURE data results. Any individual that contained 98.4375% or less of their main species

ancestry (expected ancestry proportion after 5 generations of backcrossing) was identified as

being of hybrid ancestry.

With the COI gene sequences, we created a phylogeny of the *Lasius* individuals. We

first aligned the sequences using the L-INS-i algorithm of mafft v7.481 (Katoh et al., 2005). We

then built a Maximum Likelihood tree using IQ-TREE v2.2.0.5 (Minh et al., 2020) using

ModelFinder Plus (Kalyaanamoorthy et al., 2017) to select the best substitution model, and

assessed it with 1000 UltraFast Bootstraps. Species delineation from the COI gene was

compared to combined RAD-seq and morphological delineations to further confirm species

identity for each individual.

Results

Decontamination of RADseq Data

RADseq genotyping was performed on 1,171 *Lasius* individuals. After excluding

individuals missing at least 50% of genotypes, 981 individuals were left. The allelic depth ratio

showed signs of a large amount of nuclear DNA contamination as many individuals exhibited an

ADR peak around 0.9 (Figure 3A). The effect of contamination can also be seen by the

dramatically increased proportion of heterozygous loci, especially in individuals that are missing

a higher proportion of genotypes (Figure 3B). In order to obtain biologically meaningful results

based on the SNP data created from these individuals, it was necessary to decontaminate the

data.

Through competitive mapping, it appeared that the majority of the read contamination

originated from ants from the *Formica* genus as the majority of non-*Lasius* reads mapped to the

Formica genome (supplemental Figure S1). After filtering out the reads that did not preferentially

map to the *Lasius* genome, an average of 64.3% (median 72.6%, standard deviation 18.1%) of

reads remained. After SNP calling and discarding low-data individuals, we retained 974

samples.

The efficiency of competitive mapping as a decontamination technique can be seen

clearly when comparing the decontaminated allelic depth ratio (Figure 3C) and the

decontaminated proportion of heterozygous loci per individual (Figure 3D) to the contaminated

data. The majority of minor allele frequencies are around 0.5, as expected.

However, competitive mapping is only able to remove contaminants outside of the

Lasius genus, intrageneric contamination would remain. To eliminate this contamination, we

compared a technique of artificial "haploidization" of the samples and an allelic depth ratio

(ADR) filtering. For the latter, we discarded individuals considerably outside the expected ADR

and corrected individual genotypes that were slightly beyond the expected distribution (Figure

3E and 3F). After SNP calling and removing low-data individuals, the ADR filtering retained 902

individuals.

Figure 3. A) Allelic depth ratio distribution of contaminated RADseq data. B) Proportion

of heterozygous loci per individual in the contaminated RADseq data. C) Allelic depth

ratio distribution of competitively mapped RADseq data. D) Proportion of heterozygous

loci per individual in the competitively mapped RADseq data. E) Allelic depth ratio

distribution in the filtered RADseq data (filtered based on the allelic depth ratio, ADR).

F) Proportion of heterozygous alleles per individual in the ADR- filtered RADseq data.

In order to identify which filtering technique was successful at eliminating the most

contamination while still being useful for molecular analysis, we compared allelic depth ratio

plots of each filtering technique (Figure 3). Competitive mapping alone was not sufficient to filter

all contaminants, since some individuals still showed signatures of contamination (ADR peaks

>> 0.5). The competitive mapping in conjunction with either ADR filtering or "haploidization"

were the two best options for removing contaminants based on allelic depth frequency and MDS

clustering (Figure 4 B-E). However, the "haploidization" resulted in more loss of information and

would preclude additional analyses such as estimations of heterozygosity. For this reason, we

selected the competitive mapping + ADR filtering method for downstream analyses.

Figure 4. Taxonomic inference based on the different versions of the dataset. **A** Species color code. **B** - **E** MDS plots of original data and each decontamination process. **F** Phylogeny of the COI mitochondrial gene. **G** Final genetic-based species identification. **H** - **K** ADMIXTURE proportions of original data and each decontamination process, corresponding to the adjacent MDS plots. Stars denote individuals inferred to be introgressed. White lines correspond to individuals that were filtered out due to high levels of suspected contamination or low remaining SNP data. **B & H**: Original

267   (contaminated) data. **C & I**: Competitive mapping only. **D & J:** Competitive mapping +

268   haploidization. **E & K**: Competitive mapping + Allelic Depth Ratio filtering.

Species Delineation

Out of the 902 genotyped individuals that had sufficient data after competitive mapping

and ADR filtering, 22 could not be confidently identified to the species level during initial

morphological identification. The availability of nuclear genotype information in conjunction with

COI sequence information allowed us to unambiguously assign these samples to a species.

Additionally, 18 morphological misidentifications could be corrected based on the molecular

analyses, see Supplementary Table 1 for misidentification reassignments.

Three individuals, two morphologically identified as *L. alienus* and one *L. psammophilus*,

could not confidently be assigned to a genetic group in the ADMIXTURE analysis (Figure 5E)

due to the low number of conspecifics. For the two *L. alienus* the ancestry proportions are

unique enough that it is unlikely that they belong to any of the larger species' groups. For the *L.*

psammophilus individual, in the absence of other samples of this species, it remains unclear if

this is truly a *L. psammophilus* or another *Lasius* species not represented in the remaining

dataset.

The combination of the MDS, COI, and ADMIXTURE analyses also allowed us to

investigate the presence of cryptic species. MDS plots based on SNP distances were created

for each species with at least 10 individuals to improve the resolution for detecting cryptic

species (Supplementary Figure 2). However, there was no evidence of genetic subdivision

within any of the species in the MDS plot, COI phylogeny, nor when the K was increased in the

ADMIXTURE analysis, indicating that there are no cryptic species within the sampled *Lasius*

species.

MDS, ADMIXTURE, and COI species delineations were always consistent except in one

case where an individual morphologically identified as *L. platythorax* clustered with other *L.*

platythorax in the MDS and had a majority of *L. platythorax* SNPs according to the ADMIXTURE

analysis; however, this individual was on the *L. emarginatus* branch of the COI phylogeny

(Figure 5A). The ancestry proportion of this individual is 95.06% from *L. platythorax* and 4.94%

from *L. emarginatus* based on ADMIXTURE. This individual appears to have retained the

mitochondria of a female *L. emarginatus* during a hybridization event in its ancestry, followed by

generations of backcrossing into the *L. platythorax* population, resulting in a predominantly *L.*

platythorax nuclear DNA.

Hybrid identification

All individuals with admixture percentages of at least 1.56% (the expected proportion for

a hybridization event followed by five generations of backcrossing) were classified as "admixed"

(Figure 5C-E). Before filtering, 256 individuals were identified as hybrids using this cut-off.

However, read filtering by competitive mapping reduced the number of hybrids to 56 individuals,

artificial haploidization of the data in conjunction with competitive mapping reduced this to 9

individuals, and allelic depth filtering with competitive mapping resulted in only 1 individual

classified as admixed. Indications that this single individual is a hybrid are consistent with the

comparison of nuclear and mitochondrial DNA, in which the majority of this individual's nuclear

DNA is consistent with other *L. platythorax* samples; however, its COI gene appears consistent

with *L. emarginatus* samples. This difference between the number of admixed individuals after

the haploidization method and ADR method of filtering is because false hybrids identified in the

haploidized dataset were removed during the ADR filtering due to falling above the 5%

percentile of expected ADR distribution. The admixed identification of these individuals in the

haploidized dataset indicates that at certain loci, more than half of the reads at that site

contained an allele originating from contamination. However, as the overall allelic depth ratio

frequency of these individuals was high enough for them to be excluded during the ADR

filtering, this is likely a result of contamination at sites with low coverage.

Discussion

While sequencing technologies continue to advance and make genetic information more

accurate and accessible, the risk of contamination in the data is rarely considered. Lack of

proper quality control can easily result in drastic misinterpretations of any results. Here we show

the substantial differences in biological interpretations between contaminated and

decontaminated data.

We developed a method that allowed us to successfully eliminate the contamination

from RADseq data of *Lasius* ants, allowing us to investigate over 900 individuals' genetic

identities. Using molecular analyses, we were able to confidently assign all individuals to a

species when multiple conspecifics were present among the genetically analyzed samples. We

corrected several morphological misidentifications among difficult to distinguish species (n = 18)

and we were able to assign morphologically ambiguous individuals to a clear species (n = 22).

This illustrates the benefits of complementing morphological identification with molecular data to

distinguish similar-looking species in an integrative way (Dayrat, 2005). In extension to this, we

analyzed the known species groups to determine if there were any cryptic species. Based on

our analysis of nuclear genotypes and COI divergence, we did not find evidence of cryptic

species or lineages within any of the *Lasius* species in our sampling that had more than 10

individuals (*L. brunneus, L. emarginatus, L. flavus, L. fuliginosus, L. niger, L. paralienus, L.*

platythorax). Those with less than 10 individuals (*L.alienus, L.mixtus, L.myops,*

L.psammophilus) could not be confidently assessed due to the low sample size. Despite this,

the position of the 2 *L. alienus* individuals on the COI tree assignment as a sister group to *L.*

paralienus is consistent with the fact that *L. alienus* was previously distinguished as a complex

of several cryptic species, two of which being *L. paralienus* and *L. psammophilus*, therefore

making it likely they are all genetically close (Seifert, 1992). However, the sole *L. psammophilus*

is assigned as a sister group to L. brunneus whereas previous studies have determined it to be

a sister group to *L. alienus* (Janda, Folková, & Zrzavý, 2004; Steiner et al., 2004). As this was

the only sample from this species, we cannot be sure if this is truly a *L. psammophilus* or is a

member of a different species.

After thorough decontamination, we were able to identify only one hybrid. Interestingly,

the ancestry proportion of this individual is 95.06% from *L. platythorax* and 4.94% from *L.*

emarginatus; however, the COI phylogeny positions it with *L. emarginatus* indicating that this

individual retains the mitochondria from *L. emarginatus*. As mitochondria can only be passed

from mother to offspring, it appears that the hybridization event between these two species

occurred 4 to 5 generations ago between a female *L. emarginatus* and a male *L. platythorax*

and each generation produced a daughter that continued to mate with male *L. platythorax*. This

is clear evidence of the hybridization between the two species and the ability to produce fertile,

hybrid offspring that are capable of introgression back into their parental species' population.

Previous studies have also found evidence of hybridization within the *Lasius* genus, identifying

several phenotypic intermediates between *L. emarginatus* and *L. platythorax* that were

proposed to be F1 hybrids between the two species (Seifert, 1992; Seifert, 2019). Hybrids and

evidence of introgression between *L. niger* and *L. psammophilus* as well as between *L. niger*

and *L. alienus* have also been previously recorded (Pearson, 1983; Van der Have et al., 2011).

We did not encounter any cases of hybrids from these species, likely as a result of the small

sample sizes of *L. psammophilus* and *L. alienus* in this survey. As our sampling did not

specifically target hybrid individuals, we are able to study an unbiased sampling and observe

the frequency of hybridization between the *Lasius* species in this survey. It appears that

hybridization is relatively rare among *Lasius* species. Surprisingly, we did not find any F_1

colonies, despite previous indications that they might be frequent in this genus (Feldhaar et al.,

2008).

The ecological behaviors of different species can affect the frequency of hybridization

between them. Several species of *Lasius* ants exhibit social parasitism which generally entails a

parasitic queen invading the colony of a host species and manipulating the workers to care for

her and her own brood of eggs (Janda et al., 2004). Among our sampled species, *L. mixtus* and

L. fuliginosus exhibit temporary social parasitism in which the invading queen kills the host

queen, and the host workers are temporarily mixed with the parasite's workers until the hosts

eventually die off and the colony is only made of the parasitic species (Janda et al., 2004). *L.*

mixtus has been observed parasitizing several species including *L. niger, L. platythorax, L.*

alienus and *L. flavus* (Donisthorpe ,1922; Wilson, 1955; Collingwood, 1979; Dekoninck et al.,

2004; Seifert 2018; Seifert, 1988). *L. fuliginosus* has been found to parasitize *L. niger, L.*

alienus, L. brunneus, among others, and is a hyperparasite (a parasite of a parasite) of *L.*

mixtus (Janda et al., 2004; Stärcke, 1944; Wilson,1955; Yamauchi & Hayashida 1968; Seifert,

1996). One would assume that these species that are often in close contact would be more

likely to hybridize. However, we found no hybrids between parasite and host species suggesting

the presence of barriers contributing to reproductive isolation.

The combined use of nuclear and mitochondrial DNA gives us insight into cases that

would have resulted in erroneous species identification if the mitochondrial COI gene had been

used alone. The *L. emarginatus* and *L. platythorax* hybrid individual would have been assigned

to *L. emarginatus* as a misidentification due to its mismatched mitochondria, even though it is

mostly of *L. platythorax* ancestry. This could have resulted in many misinterpretations of

morphology and behavior if a mostly *L. platythorax* individual was studied believing it to be a *L.*

emarginatus individual. To ensure the greatest amount of information can be applied it is

necessary to adequately remove any contaminants from the data to ensure accurate analyses

of both nuclear and mitochondrial sequences. As we have shown, contamination from various

sources can result in the presence of false introgression in the dataset. In this case, it would

have indicated widespread hybridization between various *Lasius* species that has never been

recorded, as well as indicated individuals with ancestry from three species. Such "triple hybrids"

have been reported in a similar dataset (Zbinden et al., 2023). Estimating the potential

contribution of contaminations to this signal would be interesting.

The use of competitive mapping enabled us to exclude reads that most likely belong to

other ant genera. We found that a large amount of the contamination appeared to be a result of

Formica material being introduced into our *Lasius* samples. More precise filtering based on

allelic depth frequency allowed us to further eliminate interspecific contamination within the

Lasius genus. While this resulted in some loss of low-quality samples, the remaining samples

provide a more accurate representation of the genetic makeup of *Lasius* individuals throughout

Vaud, Switzerland. While we cannot be sure how this contamination was introduced into our

samples, it appeared that most of the contaminated samples were those with DNA extraction

performed by the CCDB (see Supplementary Table 2), highlighting the fact that even data

based on wet lab work conducted by professional sequencing facilities should be quality

controlled.

The ADR method we presented in this paper enabled us to drastically reduce the rate of

false positive hybridization signals. It may however be slightly too conservative in the specific

case where two hybridizing species differ in the amount of paralogs in their genomes. Paralogs

are expected to yield signals similar to a contamination. For example, an originally

monomorphic genomic region undergoing duplication, followed by divergence, could result in an

AAAB signal. This would result in an ADR of 0.75, which would likely be filtered out.

RADseq allows for genotyping of non-model organisms to be done quickly and in large

enough quantities for evolutionary research at the population level (Razkin et al., 2016;

Rancihac et al., 2019; Herrera & Shank, 2016; Wagner et al., 2020). Alone, but especially in

conjunction with COI barcoding, one can create a relatively clear picture of how populations or

species intermingle with each other. Yet the presence of contamination can easily lead to

incorrect conclusions. Not only can it create false-positive hybrid signals, but it may also result

in other biases, such as artificially inflating heterozygosity estimates or underestimating

population differentiation. Several studies have generated inconclusive or even incorrect results

as a result of DNA contamination that was not fully examined before analysis (Merchant et al.,

2014; Boothby et al., 2015; Koutsovoulos et al., 2016). While some of these kinds of mistakes

are quickly caught, many others can easily slip through the cracks and go on to inform other

studies with incorrect information. Correctly analyzing and interpreting data relies heavily on the

quality of the data in the first place. Human and technical errors are unavoidable, but

understanding methods in which to minimize these errors can allow for accurate conclusions to

be drawn. The method we developed here can be applied to any type of genotyping and

resequencing data, and we suggest that systematically checking for the presence of

contamination should be part of the default quality-check pipeline of all genomic datasets.

Acknowledgments

We thank Marc Bastardot, Marjorie Labédan, Christine La Mendola, Heather Ryder and Simon Vogel for their help processing samples and Anne Freitag for morphological species identification of samples. This study was supported by funding from a bequest of M. Rullens to the University of Lausanne, the fondation Herbette, the Société Vaudoise des Sciences Naturelles, and the Retraites Populaires.

Data Availability

The code used and scripts created for the data analysis can be found on GitHub: https://github.com/glavanc1/Lasius_species_ID. Demultiplexed RADseq reads can be found on SRA (Bioproject PRJNA1176511). COI sequences can be found on GenBank (accession PQ506036-PQ507160). See Supplementary Table 3 for SRA and GenBank accession numbers for each *Lasius* individual and outcome during the contamination filtering process.

References

Agosti, D., & Bolton, B. (1990). New characters to differentiate the ant genera Lasius F. and Formica L. (Hymenoptera: Formicidae). *Entomologist's Gazette*, *41*(3), 149-156.

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome research*, *19*(9), 1655-1664.

Avril, A., Dépraz, A., Schwander, T., & Freitag, A. (2019). Opération Fourmis, le premier

recensement participatif des fourmis vaudoises-Contexte, méthodologie et bilan

préliminaire. *Bulletin de la Société vaudoise des Sciences naturelles*, *98*, 109-120.

Behr, A. A., Liu, K. Z., Liu-Fang, G., Nakka, P., & Ramachandran, S. (2016). Pong: fast analysis

and visualization of latent clusters in population genetic data. *Bioinformatics*, *32*(18),

2817-2823.

Boothby, T. C., Tenlen, J. R., Smith, F. W., Wang, J. R., Patanella, K. A., Osborne Nishimura,

E., ... & Goldstein, B. (2015). Evidence for extensive horizontal gene transfer from the

draft genome of a tardigrade. Proceedings of the National Academy of Sciences,

112(52), 15976-15981.

Brelsford, A., Dufresnes, C., & Perrin, N. (2016). High-density sex-specific linkage maps of a

European tree frog (Hyla arborea) identify the sex chromosome without information on

offspring sex. *Heredity*, *116*(2), 177-181.

Bushnell, B. (2014). *BBMap: a fast, accurate, splice-aware aligner* (No. LBNL-7065E).

Lawrence Berkeley National Lab.(LBNL), Berkeley, CA (United States).

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an

analysis tool set for population genomics. *Molecular ecology*, *22*(11), 3124-3140.

Collingwood, C. A. (1979). The Formicidae of Fennoscandia and Denmark. Fauna

Entomologica candinavica, 8.

Cox, A. J., & Hebert, P. D. (2001). Colonization, extinction, and phylogeographic patterning in a

freshwater crustacean. *Molecular ecology*, *10*(2), 371-386.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., ... & 1000

Genomes Project Analysis Group. (2011). The variant call format and VCFtools.

Bioinformatics, *27*(15), 2156-2158.

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., ... & Li, H. (2021).

Twelve years of SAMtools and BCFtools. *Gigascience*, *10*(2), giab008.

Dayrat, B. (2005). Towards integrative taxonomy. *Biological journal of the Linnean society*,

85(3), 407-417.

Dekoninck, W., Boer, P., & Maelfait, J. P. (2004). Lasius platythorax Seifert, 1991 as a host of

several Chthonolasius species, with remarks on the colony foundation of the parasites

(Hymenoptera: Formicidae). *Myrmecologische Nachrichten*, *6*, 5-8.

Donisthorpe, H. (1922). The colony founding of Acanthomyops (Dendrolasius) fuliginosus Latr.

The Biological Bulletin, *42*(4), 173-184.

Edelman, N. B., & Mallet, J. (2021). Prevalence and adaptive impact of introgression.

Annual Review of Genetics, 55(1), 265-283.

Feldhaar, H., Foitzik, S., & Heinze, J. (2008). Lifelong commitment to the wrong partner:

hybridization in ants. *Philosophical Transactions of the Royal Society B: Biological

Sciences*, *363*(1505), 2891-2899.

Feuerborn, T. R., Palkopoulou, E., van der Valk, T., von Seth, J., Munters, A. R., Pečnerová, P.,

... & Díez-del-Molino, D. (2020). Competitive mapping allows for the identification and

exclusion of human DNA contamination in ancient faunal genomic datasets. *BMC

genomics*, *21*(1), 1-10.

Freitag A., Schwander T., Broennimann O. & Dépraz A. (2020). Opération Fourmis, les

résultats du premier recensement participatif des espèces de fourmis vaudoises. Bulletin

de la Société Vaudoise des Sciences Naturelles 99: 13-27. https://doi.org/10.5169/seals-

917230

Hebert, P. D., Cywinska, A., Ball, S. L., & DeWaard, J. R. (2003). Biological identifications

through DNA barcodes. *Proceedings of the Royal Society of London. Series B:

Biological Sciences*, *270*(1512), 313-321.

Hebert, P. D., Penton, E. H., Burns, J. M., Janzen, D. H., & Hallwachs, W. (2004). Ten species

in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly

Astraptes fulgerator. *Proceedings of the National Academy of Sciences*, *101*(41), 14812-

14817.

Herrera, S., & Shank, T. M. (2016). RAD sequencing enables unprecedented phylogenetic

resolution and objective species delimitation in recalcitrant divergent taxa. *Molecular*

phylogenetics and evolution, *100*, 70-79.

Janda, M., Folková, D., & Zrzavý, J. (2004). Phylogeny of Lasius ants based on mitochondrial

DNA and morphology, and the evolution of social parasitism in the Lasiini (Hymenoptera:

Formicidae). *Molecular phylogenetics and evolution*, *33*(3), 595-614.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermiin, L. S. (2017).

ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature methods*,

14(6), 587-589.

Katoh, K., Kuma, K. I., Toh, H., & Miyata, T. (2005). MAFFT version 5: improvement in accuracy

of multiple sequence alignment. *Nucleic acids research*, *33*(2), 511-518.

Knaus, B. J., & Grünwald, N. J. (2017). vcfr: a package to manipulate and visualize variant call

format data in R. *Molecular ecology resources*, *17*(1), 44-53.

Koutsovoulos, G., Kumar, S., Laetsch, D. R., Stevens, L., Daub, J., Conlon, C., ... & Blaxter, M.

(2016). No evidence for extensive horizontal gene transfer in the genome of the

tardigrade Hypsibius dujardini. Proceedings of the National Academy of Sciences,

113(18), 5053-5058.

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows–Wheeler

transform. *Bioinformatics*, *26*(5), 589-595.

Mayr, E. (1942). Systematics and the origin of species from the viewpoint of a zoologist.

Systematics and the origin of species from the viewpoint of a zoologist.

Merchant, S., Wood, D. E., & Salzberg, S. L. (2014). Unexpected cross-species contamination

in genome sequencing projects. *PeerJ*, *2*, e675.

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A.,

& Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic

inference in the genomic era. *Molecular biology and evolution*, *37*(5), 1530-1534.

Pearson, B. (1983). Hybridisation between the ant species Lasius niger and Lasius alienus: the

genetic evidence. *Insectes sociaux*, *30*(4), 402-411.

R Core Team. A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, 2021. *Google Scholar There is no corresponding record*

for this reference.

Rancilhac, L., Goudarzi, F., Gehara, M., Hemami, M. R., Elmer, K. R., Vences, M., & Steinfarz,

S. (2019). Phylogeny and species delimitation of near Eastern Neurergus newts

(Salamandridae) based on genome-wide RADseq data analysis. *Molecular*

Phylogenetics and Evolution, *133*, 189-197.

Razkin, O., Sonet, G., Breugelmans, K., Madeira, M. J., Gómez-Moliner, B. J., & Backeljau, T.

(2016). Species limits, interspecific hybridization and phylogeny in the cryptic land snail

complex Pyramidula: the power of RADseq data. *Molecular Phylogenetics and*

Evolution, *101*, 267-278.

Romiguier, J., Borowiec, M. L., Weyna, A., Helleu, Q., Loire, E., La Mendola, C., ... & Keller, L.

(2022). Ant phylogenomics reveals a natural selection hotspot preceding the origin of

complex eusociality. *Current Biology*, *32*(13), 2942-2947.

Seifert, B. (1988). A revision of the European species of the ant subgenus Chthonolasius

(Insecta, Hymenoptera, Formicidae). *Entomologische Abhandlungen*, *51*(2), 143-180.

Seifert, B. (1991). Lasius platythorax n. sp., a widespread sibling species of Lasius niger

(Hymenoptera: Formicidae). *Entomologia Generalis*, *16*(1), 69-81.

Seifert, B. (1992) A taxonomic revision of the Palaearctic members of the ant subgenus Lasius

s.str. (Hymenoptera: Formicidae). Abhandlungen und Berichte des Naturkundemuseums

Görlitz, 66 (5), 1–67.

Seifert, B. (1996). *Ameisen: beobachten, bestimmen*. Augsburg: Naturbuch Verlag.

Seifert, B. (2018). The Ants of Central and North Europe. Lutra Verlags- und

Verttriebsgesellschaft.

Seifert, B. (2019). Lasius nigroemarginatus Forel, 1874 is a F1 Hybrid between L. emarginatus

(Olivier, 1792) and L. platythorax Seifert, 1991 (Hymenoptera, Formicidae). *Beiträge zur*

Entomologie= Contributions to Entomology, *69*(2), 291-300.

Stärcke, A. (1944). Retouches sur quelques Fourmis d'Europe: III. Autres Lasius.

Entomologische Berichten, *11*(255), 153-158.

Szewczyk, T. M., Lavanchy, G., Freitag, A., Depraz, A., Avril, A., Broennimann, O., ... &

Schwander, T. (2024). LEVERAGING CITIZEN SCIENCE TO ASSESS RICHNESS,

DIVERSITY, AND ABUNDANCE IN ANT COMMUNITIES. *bioRxiv*, 2024-06.

Van der Have, T. M., Pedersen, J. S., & Boomsma, J. J. (2011). Mating, hybridisation and

introgression in Lasius ants (Hymenoptera: Formicidae). *Myrmecological News*, *15*, 109-

115.

Venables WN, Ripley BD (2002). *Modern Applied Statistics with S*, Fourth edition. Springer,

New York. ISBN 0-387-95457-0, https://www.stats.ox.ac.uk/pub/MASS4/.

Wagner, F., Ott, T., Schall, M., Lautenschlager, U., Vogt, R., & Oberprieler, C. (2020). Taming

the Red Bastards: Hybridisation and species delimitation in the Rhodanthemum

arundanum-group (Compositae, Anthemideae). *Molecular phylogenetics and evolution*,

144, 106702.

Wilson, E. O. (1955). A monographic revision of the ant genus Lasius. *Bulletin of the museum of*

Comparative Zoology, Harvard, *113*, 2-pls.

Yamauchi, K., & Hayashida, K. (1968). Taxonomic Studies on the Genus Lasius in Hokkaido,

with Ethological and Ecological Notes (Formicidae, Hymenoptera).: I. The Subgenus

Dendrolasius or Jet Black Ants (With 8 Text-figures and 2 Tables). *北海道大學理學部紀*

要, *16*(3), 396-412.

Zbinden, Z. D., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2023). A community genomics

approach to natural hybridization. *Proceedings of the Royal Society B*, *290*(1999),

20230768.

589