Long-term high temperatures affect seed maturation and seed coat integrity in *Brassica napus*

Unnikannan Prabhullachandran[1, 2], Ivana Urbánková[3], Alejandro Medaglia-Mata[4,5], Audrey Creff[6], Aline Voxeur[7], Ivan Petřík[8], Aleš Pěnčík[8], Ondřej Novák[8], Benoit Landrein[6], Jan Hejátko[3], Hélène S. Robert[1]

1. Hormonal Crosstalk in Plant Development, Mendel Center for Plant Genomics and Proteomics, Central European Institute of Technology (CEITEC), Masaryk University, Brno, Czech Republic

2. Laboratory of Functional Genomics and Proteomics, National Centre for Biomolecular Research, Faculty of Science, Masaryk University, Brno, Czech Republic

3. Functional Genomics and Proteomics of Plants, Mendel Center for Plant Genomics and Proteomics, Central European Institute of Technology (CEITEC), Masaryk University, Brno, Czech Republic

4. Bioinformatics Core Facility, Central European Institute of Technology (CEITEC), Masaryk University, Brno, Czech Republic

5. Department of Biology, Faculty of Medicine, Masaryk University, Brno, Czech Republic

6. Laboratoire Reproduction et Développement des Plantes, Université de Lyon, ENS de Lyon, UCB Lyon, CNRS, INRAE, INRIA, Lyon, France

7. Université Paris-Saclay, INRAE, AgroParisTech, Institute Jean-Pierre Bourgin for Plant Science (IJPB), Versailles, France

8. Laboratory of Growth Regulators, Faculty of Science, Palacký University and Institute of Experimental Botany, The Czech Academy of Sciences, Olomouc, Czech Republic

Hélène S. Robert (corresponding author helene.robert.boisivon@ceitec.muni.cz)

ORCID

UK 0000-0001-6430-575X

AMM 0000-0003-3496-0571

AC 0000-0001-7736-5882

AV 0000-0001-9452-6756

IP 0000-0001-5320-7599

AP 0000-0002-1314-2249

33   ON 0000-0003-3452-0154

34   BL 0000-0002-2371-9996

35   JH 0000-0002-2622-6046

36   HSR 0000-0002-6610-836X

SUMMARY

• Temperatures above the optimum growth temperature affect seed development,

producing seeds with ruptured seed coats. This phenotype is associated with accelerated

embryo development. However, the molecular mechanism underlying this effect remains

unclear.

• To investigate the occurrence of temperature-induced seed coat rupture, we combined

detailed phenotyping approaches of oilseed rape seeds with transcriptomics, histology,

immunolabelling, hormone and cell wall profiling.

• Our data suggest that high temperatures accelerate embryo growth, resulting in the

formation of larger embryos but not larger seeds. The formation of large embryos

increased the mechanical pressure exerted by the embryo on the seed coat cells, reducing

their thickness. The seed coat began to mature prematurely with the accumulation of

demethylesterified pectin, possibly making the cell wall stiffer. Application of abscisic acid,

a hormone involved in seed dormancy, did not rescue the seed coat rupture phenotype.

Hormonal and transcriptional profiling indicated that the embryo did not enter dormancy.

• Prolonged high temperatures during seed development accelerated embryo growth by

stimulating cell division, while the seed coat, which depends on cell elongation, could not

withstand the tension exerted by the embryo, started seed maturation and eventually

ruptured.

KEYWORDS

Brassica napus, cell wall, long-term high temperature, pectin, seed coat rupture, seed
development

INTRODUCTION

The adverse impacts of global warming on agricultural production pose a major challenge to
global food security in the 21[st] century. Extreme weather events, such as heat stress, drought
and floods, have been observed to reduce crop yields and shorten growing seasons,
particularly in warm tropical regions. This has the potential to affect the production of staple
crops, such as rice, wheat, maize, and soybean (Zhao *et al.*, 2017; Agnolucci & De Lipsis, 2020;
Jägermeyr *et al.*, 2021; Heino *et al.*, 2023). The mean global temperature increase in 2023 was
1.36°C above preindustrial levels (National Aeronautics and Space Administration, 2024),
highlighting the need to adapt agricultural practices to higher temperatures.

Brassica napus (canola or oilseed rape), an allotetraploid species in the Brassicaceae family,
is the second most widely cultivated oilseed crop, used for the production of vegetable oil,
biofuel, and protein in animal feed. Cultivation of oilseed rape at temperatures above the
optimum growth temperature of 21°C for prolonged periods, especially during the flowering
stage, resulted in several adverse effects, including rapid vegetative growth, reduced viability
of female gametophytes, increased seed abortion rate, accelerated embryo development,
and a reduction in seed oil composition (Young *et al.*, 2004; Mácová *et al.*, 2022; Secchi *et al.*,
2023). One of the distinctive phenotypes observed during seed development of certain canola
cultivars subjected to prolonged heat stress that affects seed yield is the occurrence of the
preharvest sprouting phenotype (PHS) (Mácová *et al.*, 2022). Misregulation of seed dormancy
by abscisic acid and dormancy-related genes is considered to be the main cause of PHS in
many cereal crops (Benech-Arnold & Rodríguez, 2018; Tai *et al.*, 2021). This phenotype is
associated with seed coat rupture (SCR), which is observed in seeds during the early stages of
maturation. The exact molecular mechanism by which this phenotype is produced at elevated
temperatures remains unclear.

In Brassicaceae, seed size is regulated by the interaction between maternal and zygotic signals
that coordinate the proper development of the seed coat, endosperm, and embryo (Li *et al.*,
2019). During the maturation of *Arabidopsis* seeds, the pressure generated by the interaction
between the developing embryo and the endosperm within the seed, driven by embryo
growth, exerts tension on the developing integuments (Creff *et al.*, 2015, 2023). The tension

generated by zygotic tissues is sensed by a specific layer of the seed coat's outer integument, the adaxial epidermis (oi1). This mechanical stimulus triggers a transcriptional response marked by the activation of the mechanosensitive gene *EUI-like P450A1* (*ELA1*), which is believed to promote cell wall stiffening through demethylesterified pectin accumulation regulating seed size (Creff *et al.*, 2023). The dry mass of the primary plant cell wall is composed of cellulose, hemicellulose, and pectin (Cosgrove, 2024). Pectin forms a hydrated gel-like matrix in which cellulose and hemicellulose cross-link to form the cell wall. It is abundant in the middle lamella, an adhesive layer between adjacent cells. The most abundant pectin, homogalacturonan (HG), is produced in a methylesterified state, which can be demethylesterified by the activity of pectin methylesterase (PME) (Turbant *et al.*, 2016; Wu *et al.*, 2018). Blockwise removal of methyl groups can lead to the formation of Ca^{2+} cross-links, which increase the wall stiffness and inhibit growth. Alternatively, random removal can lead to pectin degradation, affecting wall rheology (Wu *et al.*, 2018).

Seed size correlates with endosperm proliferation and development. The two processes are linked by a brassinosteroid-induced cell expansion and increased levels of methylesterified pectin in the seed coat, which helps to regulate seed size (Lima *et al.*, 2024). Mechanical constriction of seed size, which restricts embryo growth, triggers the maturation programme by increasing lipid and protein storage and inhibiting embryonic cell divisions (Rolletschek *et al.*, 2021, 2024). In this study, we used a multi-method approach to investigate the occurrence of the temperature-induced SCR phenotype. The results show that SCR occurs in seeds around 20 days after pollination (20 DAP) when plants are cultivated at elevated temperatures for a prolonged period. The pressure induced by unrestricted embryo growth led to increased tension in the seed coat, characterised by reduced thickness of the seed coat layers, and resulted in early changes in cell wall composition. Accumulation of demethylesterified pectin may stiffen the cell wall of the seed coat, which may break despite being stiffened or may promote pectin degradation making the seed coat more breakable. The precise mechanism by which accelerated embryo development influences heat stress-mediated seed development in oilseed rape plants remains to be elucidated. A deeper understanding of the effects of prolonged elevated temperatures on seed development could prove invaluable in the generation of thermotolerant varieties of canola varieties, which would help to improve crop yields in the context of global warming in the 21st century.

MATERIALS AND METHODS

Plant materials and growth conditions

Two *Brassica napus* spring cultivars, Topas DH4079 and Westar, were used for this study. Seeds were sterilised, germinated, and grown initially in the phytotron (21°C, 16/8 h long day period, 150 µmol/m^2/s) until the emergence of the first flowering buds, as previously mentioned (Mácová *et al.*, 2022). Plants are then transferred to greenhouse chambers with control temperature treatment (CT) and heat treatment (HT) conditions maintained under a long-day regime (16 h light/8 h dark), LED lights with a light intensity of 150 µmol/m^2/s, 35%–45% humidity, 18°C during the night and 21°C (CT) and 32°C (HT) during the day. Seeds used for the experiments are obtained from hand-pollinated flowers. The age of the seeds is counted from the day after pollination (DAP).

Seed phenotyping

Ten hand-pollinated and dried siliques from eight plants in three biological replicates from CT and HT conditions were screened for the total, viable, dead seeds and seeds undergoing SCR (seed coat rupture with radicle or cotyledon protrusion). Fresh seeds from 14 DAP to 20 DAP were collected from Topas and Westar growing at CT and HT for embryo staging. The whole seed and its isolated embryo were imaged using a Levenhuk M500 base digital camera attached to a Zeiss stemi 305 stereomicroscope. Images were used to measure the seed and embryo areas using ImageJ from 14, 18, and 20 DAP from both conditions. The ratio of the embryo-to-seed area was calculated to get information on the space occupied by the embryo within the seed at each stage of development. Statistical significance was accounted for by a two-way ANOVA test with Sidak's multiple comparisons (* $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$, and **** $p < 0.00005$).

Hormonal measurements

Two experiments were performed. First, the hormone profiling was done on 14, 20, 26, and 36 DAP whole seeds collected from CT or HT in five replicates, each pooled from three plants. The 20, 26, and 36 DAP seeds collected from HT showing an SCR phenotype were separately collected and analysed. A second experiment was carried out on isolated embryos and seed coats of 14 and 18 DAP seeds collected from CT or HT in four replicates, each pooled from three plants. Abscisates, auxins (except indole-3-pyruvic acid; IPyA), and gibberellins were

determined using ultra-high-performance liquid chromatograph I-Class (Waters, Milford, CT,
USA) coupled with tandem mass spectrometer Xevo TQ-XS (Waters, Manchester, UK)
according to the "hormonomics" methodology (Šimura *et al.*, 2018). Approximately 10 mg of
fresh weight of each sample was aliquoted into microtubes. One ml of 60% aqueous
acetonitrile (v/v), a mixture of stabile isotopic labelled internal standards, and four ceramic
extraction beads were added into the tube. The samples were shaken in a Retsch MM400
oscillation mill (Haan, Germany) at 27 Hz for 5 min, sonicated for 3 min, and incubated for 30
min in a refrigerator. Allegra 64R benchtop centrifuge (Beckman Coulter, USA) at 20,000 rpm
for 15 minutes was used to separate the pellet from the supernatant. The supernatant was
loaded onto Oasis® HLB 30 mg/1 cc extraction cartridge (Waters, Milford, CT, USA) previously
activated with 1 ml methanol, water, and 60% aqueous acetonitrile (v/v). The cartridge was
finally washed with 60% aqueous acetonitrile (v/v). The flow-through and wash fractions were
collected into borosilicate glass tubes (Thermo Fisher Scientific, USA). The samples were
evaporated in a SpeedVac concentrator (RC1010 Centrivap Jouan, ThermoFisher, USA) and
reconstructed in 40 µl of 25% aqueous acetonitrile (v/v). Five µl were injected onto Acquity
UPLC® CSHTM 1.7 µm, 2.1x150 mm chromatographic column and eluted using a gradient
elution of acidified water and acetonitrile. Data were acquired in a multiple reaction
monitoring mode. The detailed chromatographic and mass spectrometry conditions are
provided by Šimura et al. (2018). Data were processed with MassLynx V4.2 software (Waters,
Manchester, UK), and the concentrations of determined plant hormones were calculated
using the isotope dilution method. Endogenous IPyA was determined following the protocol
described by Pěnčík et al. (2018). Approximately 10 mg (fresh weight) samples were extracted
in 1 ml of cold 50 mmol/l phosphate buffer (pH 7.0) containing 0.1% sodium
diethyldithiocarbamate and isotope-labelled internal standard of IPyA. A 200 µl aliquot was
derivatised with cysteamine, acidified to pH 2.7 with HCl, and purified using in-tip µSPE. Eluted
samples were evaporated under reduced pressure, reconstituted in 10% aqueous methanol,
and analysed using an HPLC system 1260 Infinity II (Agilent Technologies, USA) equipped with
a Kinetex C18 column (50 mmx2.1 mm, 1.7 µm; Phenomenex) and coupled to 6495 Triple
Quad detector (Agilent Technologies, USA).

ABA treatment

ABA treatment was performed by culturing 15 siliques of 14 DAP and 18 DAP from HT and CT

conditions, respectively, collected from three plants in three biological replicates. The siliques

were cut at the pedicel region, collected, surface-sterilised with 70% ethanol, air dried, and

cultured in 50 ml tubes with 20 ml of ½ MS with 0.8% plant agar, 0.05% plant preservative for

tissue culture media (Plant Tissue Culture Technologies) and supplemented with 10 µM ABA

(Duchefa Biochemie) or the same volume of DMSO as negative control. A representative

picture of the cultured siliques is shown in Fig. S1. The silique culture was incubated in the

greenhouse with the other plants to simulate the same temperature conditions. The total,

viable, dead and SCR seed numbers were manually estimated on dried siliques under a Zeiss

stemi 305 stereomicroscope.

RNA extraction and sequencing

RNA was extracted from 50 mg of ground tissue of 14 DAP, 18 DAP, 20 DAP normal, and 20

DAP SCR seeds from CT and HT, collected from three plants in four biological replicates using

a NucleoSpin RNA Plant and Fungi kit (Macherey-Nagel) following the manufacturer's

protocol. The samples were treated with RNase-free rDNase (Macherey-Nagel) and repurified

using the RNeasy MinElute Cleanup Kit (Qiagen). The samples were sequenced by Novogene

Genomic Sequencing Labs. After sample quality control and library preparation, paired-end

short-read RNA sequencing was performed on poly-A RNA using an Illumina Novaseq

Sequencing platform. Quality check of raw paired-end fastq reads was carried out by FastQC

(Andrews, 2010), and their origin was categorised using BioBloomTools v2.3.4 (Chu *et al.*,

2014). The Illumina adapters and quality trimming of raw fastq reads were performed using

Trimmomatic v0.39 (Bolger *et al.*, 2014) with settings CROP:250 LEADING:3 TRAILING:3

SLIDINGWINDOW:4:5 MINLEN:35. Trimmed RNA-Seq reads were mapped to the *Brassica*

napus genome (genome build: Bra_napus_v2.0) with gene annotation (NCBI *Brassica napus*

Annotation Release 101) using STAR v2.7.3a (Dobin *et al.*, 2013) as splice-aware short read

aligner and default parameters except outFilterMismatchNoverLmax 0.1 and twopassMode

Basic. Quality control after alignment concerning the number and percentage of uniquely-

and multi-mapped reads, rRNA contamination, mapped regions, read coverage distribution,

strand specificity, gene biotypes, and PCR duplication was performed using RSeQC v4.0.0

(Wang *et al.*, 2012), Picard toolkit v2.25.6 (Picard toolkit, 2018), Qualimap v.2.2.2

(Okonechnikov *et al.*, 2016). All compatible results and statistics were processed by MultiQC

v1.10.1 (Ewels *et al.*, 2016). The sequence data have been deposited in the NCBI under the

Bio-Project PRJNA1188476 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1188476).

The differential gene expression analysis was calculated based on the gene counts produced

using featureCounts from Subread package v2.0 (Liao *et al.*, 2014) and further analysed by

Bioconductor package DESeq2 v1.34.0 (Love *et al.*, 2014). Because of its conservative features

and to prevent false positive results, data produced by DESeq2 with independent filtering

were chosen for the differential gene expression analysis. Genes were considered to be

differentially expressed based on a cut-off for adjusted p-value ≤ 0.05 and log2FC ≥1 or ≤-1.

Clustered heatmaps were generated from selected top differentially regulated genes using R

package pheatmap v1.0.12 (Kolde, 2015), volcano plots were produced using ggplot2 v3.3.5

package (Wickham, 2011), and MA plots were generated using ggpubr v0.4.0 package

(Kassambara, 2018). GO enrichment is done taking all the Entrez gene IDs and cross-mapping

them against UniProt IDs, then taking all the transcript sequences and mapping them (using

BLAST) against the NCBI Nucleotide database subset for all species under taxons 3701

(A*rabidopsis* genus) and 981071 (*Brassicaceae* tribe) obtaining RefSeq gene IDs which were

cross-mapped against UniProt IDs and then used in GO enrichment. The co-expression cluster

analysis is described in Method S1.

Seed embedding and sectioning

Seeds were fixed in FAA (45% ethanol, 5% acetic acid and 1.85% formaldehyde) for 24 h under

vacuum, followed by another incubation at 4°C for 24 h in fresh FAA. The seeds were washed

once with water. Tissue softening was done with Franklin Maceration solution (1:1 v/v of 30%

H_2O_2 and glacial acetic acid) for 4 h at 60°C. Dehydration was done using 10, 20, 30, 40, 50,

60% ethanol series in 10% glycerine for 1 h at 30°C, followed by 4 days in 70% ethanol at 4°C

and dehydration with 80, 90 and 95% ethanol for 1 h at room temperature. Infiltration and

embedding with Technovit 7100 (Kulzer Technik Gmbh) were performed according to the

manufacturer's protocol with the following modifications. The initial infiltration was

performed in a 50% infiltration solution in ethanol overnight at room temperature, followed

by incubation for one week at 4°C in 100% infiltration solution with 1% w/v Technovit 7100

Hardener I. Embedding was performed with 6.6% w/v Technovit 7100 Hardener II in

infiltration solution and incubated for 8 days at -18°C to delay initial polymerisation. The

samples were then incubated at room temperature to complete the polymerisation process.

The 3-µm sections were prepared with a Leica RM2235 microtome using an attached stainless-steel knife.

Toluidine blue staining

The sections were stained with 0.05% Toluidine blue (Acros Organics) for 1 minute, followed by four washes of 3 minutes each. Mounting was done on dried slides with Entellan (Sigma-Aldrich). Imaging of the sections was performed using a Zeiss AxioImager microscope. The thickness of the seed coat layers was measured in twenty regions per section, for three sections per seed and ten seeds per sample, using Image J. Graphs were plotted with significance calculated using a two-way ANOVA test with Tukey multiple comparisons.

Staining of lignin and pectin

Seven seeds at 14 and 18 DAP from CT and HT and 22 DAP from CT were used. Pectin was detected using a 0.05% w/v Ruthenium Red stain (Sigma-Aldrich) in an aqueous solution. Sections were rehydrated before staining for 7 minutes, followed by two wash steps with water, and mounted in Entellan (Sigma-Aldrich). Lignin was detected with 0.5% Safranin-O (Acros organics) in an aqueous solution followed by three washes in water. Slides were mounted in Entellan (Sigma-Aldrich). Imaging was done using a Zeiss AxioImager microscope.

Immunolabelling

Seeds at 10, 14, and 18 DAP from CT and HT, in two biological replicates, were fixed in 4% w/v paraformaldehyde in ice-cold PEM buffer (50 mM PIPES, 5 mM EGTA, and 5 mM $MgSO_4$, pH 6.9). Samples were incubated on ice under vacuum for twice 1 h with fresh fixative solution, followed by overnight incubation at 4°C. The samples were washed twice with ice-cold PEM buffer, dehydrated through an ethanol series (50%, 70%, 80%, 90% and 100% three times) under vacuum and infiltered with increasing concentrations (30%, 50% and 100%) of LR white resin (Agar Scientific) in absolute ethanol over 8 days before being polymerised at 60°C for 24 h. Six seeds by timepoint per growth condition per biological replicate were sectioned (1 µm thickness) using a glass knife mounted on a Leica RM2265 or EM UC7 microtome. Immunolocalizations were performed with JIM5, LM19, LM20, and LM25 antibodies (Agrisera). For immunolocalisation with LM25, an incubation with 0.1% pectolyase in 1X PBS was done for 25 min at room temperature and washed once in 1X PBS before the blocking

step. For immunolocalisation with JIM5, LM19, and LM25, the sections were blocked with 3% w/v BSA in 1X PBS and incubated for 1 h at room temperature. For the immunolocalisation with LM20, an incubation in CAPS buffer (50 mM CAPS and 2 mM $CaCl_2$) for 1 h followed by three washes with 1x PBS was performed before blocking with 3% skimmed milk in 1X PBS. Sections were incubated with 1:10 v/v dilution of antibody prepared with 1% BSA in 1X PBS for JIM5, LM19 and LM25 and 1% skimmed milk in 1X PBS for LM20 overnight at 4°C in a humid chamber. The sections were brought to room temperature for 1 h, washed three times with the above-mentioned buffers and incubated for 1 h in a humid chamber in the dark with the secondary antibodies. A dilution of 1:100 v/v of secondary antibodies [anti-rat IgG Alexa 488 (Thermo Fisher) or anti-rat IgG Dylight 488 (Agrisera) for JIM5 and anti-rat IgM Dylight 488 (Thermo Fisher) for LM19, LM20 and LM25] in 1% BSA or 1% skimmed milk in 1X PBS was used. Sections were washed in the respective buffers and 1X PBS, counterstained with 0.025 mg/ml Calcofluor white (Sigma-Aldrich), and mounted with VECTASHIELD antifade mounting medium (Eurobio). The images were obtained with the ZEISS Axioimager 2 with a 40X dry objective.

The intensity of the immunofluorescence in the cell walls of the epidermal and palisade layers of the seed coat was quantified using ImageJ software. Images were adjusted using min/max settings for optimal representation of fluorescent signals. For the three epidermal layers, a segmented line with a width of 25 points was used to manually mark the cell walls of adjacent cells of the same layer. The sum of fluorescence intensity in the three epidermal layers was represented as one measurement. A region of interest was marked using the polygon selection tool for the palisade layer. The mean fluorescence value obtained over the marked regions for the different layers was subtracted from the background signal over the same area to obtain the final fluorescence intensity.

Cell wall profiling from the seed coat

The enzymatic fingerprinting of the cell wall of seed coat from 10, 14 and 18 DAP seeds grown under CT and HT conditions was performed in three biological replicates. The seed coats were dissected from the seeds and were submerged in 96% (v/v) ethanol. The ethanol fraction was discarded, and the seed coats were dried in a speed vacuum concentrator at 30°C overnight. Dried seed coats were digested with 5 U/mg DW sample of *Pectobacterium carotovorum* endo-polygalacturonase or *Paenibacillus* sp. xyloglucanase (Megazyme, Bray, Ireland) in 50

mM ammonium acetate buffer (pH 5) at 37°C for 18 h. Samples were centrifuged at 13,000

rpm for 10 min, and 100 µl of the supernatants were transferred into vials. For MS analysis,

10 µl of each fraction was injected. The oligosaccharides released from digestion were

separated according to Voxeur et al. (2019). Xyloglucan and pectin fragments were analysed

using an HPLC system (UltiMate 3000 RS HPLC system, Thermo Scientific, Waltham, MA, USA)

coupled to an Impact II Ultra-High Resolution Qq-Time-Of-Flight (UHR-QqTOF) spectrometer

(Bruker Daltonics, Bremen, Germany) equipped with an electrospray ionisation (ESI) source

in negative mode with the end plate offset set voltage to 500 V, capillary voltage to 4000 V,

nebuliser to 40 psi, dry gas to 8 l/min and dry temperature of 180°C. The Compass 1.8

software (Bruker Daltonics) was used to acquire the data. Chromatographic separation was

performed on an ACQUITY UPLC Protein BEH SEC Column (125A, 1.7 µm, 4.6 mm X 300 mm,

Waters Corporation, Milford, MA, USA) coupled with guard Column BEH SEC Column (125A,

1.7 µm, 4.6 mm X 30 mm). Elution was performed in 50 mM ammonium formate and 0.1%

formic acid at a flow rate of 0.4 ml/min and with a column oven temperature of 40°C.

Space-restricted growth of siliques

A 3-mm cross-section silicone tube was inserted over 10 DAP and 12 DAP siliques growing on

plants under HT and CT conditions, respectively, in three biological replicates with 20 siliques

per replicate. A 10 DAP silique at HT has a similar silique thickness to a 12 DAP silique at CT.

Siliques from the age and growth conditions without a silicone tube were used as a negative

control. The number of total, viable, dead and SCR seeds per dried silique were counted.

Statistical analysis

GraphPad Prism v.8.4.3 was used to plot the graphs and calculate the statistical significance

using either ANOVA or multiple comparison tests. R programming was used to analyse the

data and obtain the statistical significance for the RNA sequencing experiment.

RESULTS

***Brassica napus* seed coat integrity is affected by long-term heat stress**

We have previously reported that in *Brassica napus* cv. Topas the early seed development is

affected under long-term heat treatment (HT) (Mácová *et al.*, 2022). This study focuses on

the seed coat rupture (SCR) observed in seeds of Topas developed under HT, starting at 20

DAP, while this phenomenon has never been observed in seeds grown under control

treatment (CT) (Fig. 1a-e). The SCR phenotype occurred with variable outcomes, from a minor

seed coat break with the embryo remaining intact within the seed to a complete testa rupture

resulting in the embryo root or shoot protruding through the break site. Such embryos often

had distinguishable phenotypes (Fig. 1c,d). The quantification of the phenotypes on dry seeds

revealed a significantly reduced number of total viable seeds produced per silique,

accompanied by a significant increase in the proportion of dead seeds and seeds with an SCR

phenotype per silique in HT-grown siliques. At HT, approximately 50% of the seeds in a silique

underwent SCR (Fig. 1e). Embryo staging of the seeds from 14 to 20 DAP indicated that

embryo development is accelerated in seeds developed under HT compared to seeds grown

under CT (Fig. 1f), which is consistent with our previous observations in seeds at early

developmental stages (Mácová *et al.*, 2022). We estimated that the embryos in the seeds at

HT were four days ahead of the embryos in seeds at CT in their developmental stage, e.g., a

14 DAP embryo at HT would be developmentally similar to an 18 DAP embryo at CT (Fig. 1f).

Based on these observations, we designated 14 and 18 DAP HT seeds as early maturing seeds

and HT seeds after 20 DAP as late maturing seeds.

Fig. 1. SCR phenotype observed in Topas seeds developed under HT. (a, b) 26 DAP seeds (a)

and their isolated embryos (b) grown under CT. (c, d) 26 DAP seeds (c) and their isolated

embryos (d) grown under HT. SCR was observed (*, Δ, +, ●). * Indicates seeds with testa

rupture with intact embryo, Δ indicates testa rupture with radicle protrusion, + indicates testa

rupture with cotyledon protrusion, and ● indicates embryos with abnormal development. (e) Quantification of the number of dry seeds per silique from CT (blue) and HT (red) showing the number of total, viable, SCR and dead seeds. Asterisks indicate statistically significant differences in HT in a multiple t-test using the Holm-Sidak method (**** corresponds to *p*-values of $p < 0.0001$. (f) Embryo stage of seeds developed on CT (top) and HT (bottom) from 14 to 20 DAP. The isolated embryo shown on the right is from the seed on the left. Scale bars: 2 mm (a-d) and 3mm (f).

Unbalanced hormonal profiles of the seeds at HT may affect seed maturation

The observed SCR phenotype led us to hypothesise that HT may deregulate seed dormancy and maturation programs by affecting the homeostasis of the plant hormones abscisic acid, gibberellins, and auxin. GA and its metabolites were below the detection limit, preventing any conclusions about the influence of HT on their levels. In the seeds at 20, 26, and 36 DAP developed under HT, ABA and IAA levels are significantly and age-dependently reduced (Fig. 2a; Fig. S2). The levels of the auxin precursor anthranilate (ANT) were significantly lower in 14 DAP HT embryos. In contrast, the levels of tryptophan (TRP) and indole-pyruvic acid (IPyA) in 14 DAP HT embryos and indole-3-acetonitrile (IAN) in 18 DAP HT embryos were significantly higher compared to CT (Fig. S2). However, the levels of IAA in the CT and HT embryos at 14 or 18 DAP were not significantly different because the degradation products such as 2-oxoindole-3-acetic acid (oxIAA) and IAA-aspartate (IAA-Asp) were significantly higher in the embryos developed under HT. The levels of auxin metabolites in the seed coat were similar to those observed in the embryos, except for a significant increase in IAA levels at HT. The IAA levels were significantly reduced in mature seeds (20, 26 and 36 DAP). However, IAA levels in intact and SCR seeds were not different (Fig. S2).

A significant increase in the levels of active ABA and its catabolic products (7-hydroxy-ABA, phaseic acid (PA) and dihydrophaseic acid (DPA)) was observed in the embryos of 14 and 18 DAP seeds, whereas no significant changes were identified in the isolated seed coat (Fig. 2b). When embryo stage was taken into account and ABA levels in 14 DAP embryos from HT were compared with those in 18 DAP embryos from CT, no significant differences were found. ABA levels were significantly reduced in 26 and 36 DAP mature seeds developed under HT, with no significant difference between intact and SCR seeds (Fig. 2a). These reduced ABA and IAA levels in mature seeds led to the hypothesis of reduced seed dormancy in HT.

404.

Fig. 2. ABA levels and expression of genes involved in ABA metabolism in *Brassica napus* Topas seeds. (a) Seeds were from 14 DAP to 36 DAP of CT (blue) and HT (red, dark red). From 20 DAP, HT seeds were separated into intact seeds (red) and SCR seeds (dark red). Quantification is presented as pmol per g fresh weight. (b) Measurements were performed for ABA, phaseic acid (PA), and dihydrophaseic acid (DPA) from embryos and seed coats from 14 DAP and 18 DAP seeds grown in CT (blue) and HT (red). Asterisks indicate statistically significant differences in HT and between 14 DAP HT and 18 DAP CT in a multiple t-test using the Holm-Sidak method (*, **, *** and **** correspond to *p*-values of 0.05 > *p* > 0.01, 0.01 > *p* > 0.001, 0.001 > *p* > 0.0001 and *p* < 0.0001, respectively); n.s., not significant. (c) 15 siliques of 14 DAP and 18 DAP from HT and CT, respectively, were cultured in the presence or absence of 10 μM ABA (Fig. S1). Dry seeds were quantified for the number of total viable, SCR and dead seeds per silique. Samples with the same letter were not statistically significantly different. (d) Relative expression of selected genes representing DORMANCY ASSOCIATED gene 1 (DRM1) and FRUCTOSE 1,6-BIPHOSPHATE ALDOLASE 6 (FBA6) in 14 DAP (blue), 20 DAP intact (yellow) and 20 DAP SCR (grey) seeds. Data were extracted from the transcriptome dataset (Tables S1 and S2). (e) Heatmap showing the differential expression of selected genes involved in ABA metabolism in 14 DAP, 20 DAP intact, and 20 DAP SCR seeds. The full dataset is provided in Tables S1 and S2.

ABA treatment on the siliques did not restore the SCR phenotype

To test whether the reduced ABA in the seeds was one of the causes of the SCR phenotype

and reduced seed dormancy at HT, we cultured the Topas siliques in the presence of 10 μM

ABA in an *in vitro* silique culture experiment. The number of seeds undergoing SCR at HT in

the siliques cultured with ABA was not significantly different from that in untreated siliques

(Fig. 2c). Moreover, *in vitro* silique culture appeared to partially rescue SCR at HT compared

to seeds developed in siliques attached to the plants, most likely due to the increased

humidity of the culture setup (Fig. S1). Thus, the cultured siliques produced more viable seeds.

Comparable results were observed with different modes of exogenous application of ABA to

individual siliques and in an *in vitro* seed culture (data not shown).

The embryo-to-seed size ratio is increased at HT

We observed that seed development in Topas and Westar was accelerated under long-term

high-temperature growth, with embryos exhibiting a four-day advance in maturity under HT

compared to CT (Fig. 1, Fig. S3, Mácová et al., 2022). The SCR rate for Topas seeds was higher

than that for Westar seeds (Mácová *et al.*, 2022). We hypothesised that this accelerated

embryo growth may interfere with seed development, thereby inducing the SCR phenotype.

We compared seed area and embryo area in seeds from 14 to 20 DAP at CT and HT in both

cultivars. The seed area increased from 14 to 20 DAP, and this increase was reduced from 20

DAP as the seeds entered the maturation phase (Fig. 3a, c). In Topas, the seed size at 18 DAP

was significantly larger under HT than under CT (Fig. 3a). In Westar, seeds were slightly larger

under HT at 14 DAP and smaller at 20 DAP (Fig. 3c). It should be noted that the seeds of the

Westar cultivar were larger on average than those of the Topas cultivar. Embryos at HT were

more than twice the size of those at the same DAP at CT for each DAP in both cultivars (Fig.

3b, d). These observations suggest that the seeds did not undergo a compensatory increase

in size to accommodate the HT-induced enlargement of the embryos. The embryo-to-seed

area ratio provides valuable insight into the space occupied by the embryo within the seed at

a given DAP under the two growth conditions (Fig. 3e). We observed that the embryos

developed under HT filled significantly more space in the seed than those grown under CT.

Furthermore, the embryo-to-seed area ratio in Topas was significantly higher than that in

Westar at 20 DAP. These observations suggest that the accelerated embryo growth in HT

resulted in the formation of larger embryos but not larger seeds.

Fig 3. Embryo and seed size measurements of *Brassica napus* Westar and Topas seeds

developed at CT and HT. (a, b) Seed (a) and embryo (b) area measurements in mm^2 of 14, 18

and 20 DAP Topas seeds developed at CT (blue) and HT (red). (c, d) Seed (c) and embryo (d)

area measurements in mm^2 of 14, 18 and 20 DAP Westar seeds developed at CT (blue) and

HT (red). Asterisks indicate statistically significant differences in HT and between 14 DAP HT

and 18 DAP CT in a two-way ANOVA test with Sidak's multiple comparisons (*, **, *** and

**** correspond to *p*-values of $0.05 > p > 0.005$, $0.005 > p > 0.0005$, $0.0005 > p > 0.0005$ and

$p < 0.00005$, respectively); n.s., not significant. (e) Embryo-to-seed size ratio in 14, 18 and 20

DAP Topas and Westar seeds developed at CT and HT. Samples with the same letter were not

statistically significantly different. Representative pictures of the seeds and embryos are

shown in Fig. S3.

The formation of larger embryos at HT impacts the thickness of the seed coat cells

We wondered whether the presence of larger embryos in these seeds might affect the seed

coat cells, leading to SCR. The seed coat of *B. napus* is composed of three layers of cells (Fig.

4a): an outer epidermal layer of three cell files, a middle palisade layer made of a single layer of cells with substantial secondary cell walls, and an inner endothelial layer surrounding the endosperm (Jiang & Deyholos, 2010). The thickness of these three layers of the seed coat was measured in seeds of 10, 14, 18, 20, and 22 DAP developed under CT and HT (Fig. 4f-h). Seed coat thickness was not affected by HT in 10 DAP seeds (Fig. 4f-h). However, all three seed coat layers were significantly thinner in older seeds developed under HT (Fig. 4b-h). The palisade and the first endothelial layers were the most affected by this reduction, with a 46% reduction in endothelial cell layer thickness induced by HT as the seed aged from 14 DAP to 20 DAP, whereas the thickness reduction was only 12% at CT (Fig. 4i). These results indicate that the HT-induced embryo growth from 14 DAP increased tension on the cells of the developing seed coat, reducing its thickness due to cell stretching.

Fig. 4 The thickness of the seed coat cells is reduced in HT. (a-e) Histological sections of Topas seeds focused on the seed coat, stained with toluidine blue. (a) The seed coat is composed of an epidermal layer, a palisade layer and an endodermal layer. (b-e) Seed coat of 14 DAP seeds

from CT (b) and HT (d) and 18 DAP seeds from CT (c) and HT (e). Scale bars represent 200 µm.

(f-h) Box-whisker plots show the size mean, median, outliers, and SD in µm of the thickness

of the epidermal layer (f), palisade layer (g) and first endothelial cell layer (h) in 10, 14, 18,

and 20 DAP seeds from CT (blue) and HT (red). Samples with the same letter were not

statistically significantly different. (i) Percentage of reduction in thickness as the seed aged

from 14 to 20 DAP.

Transcriptional regulation of the dormancy regulators provides a conflicting picture of the

HT effects on seed dormancy

To gain an insight into the molecular mechanisms of SCR at HT in Topas, we performed

transcriptional profiling of seeds developed under HT and CT conditions at 14, 18 and 20 DAP,

including seeds without (20NSHT samples) and seeds with (20SHT samples) SCR phenotypes

at 20 DAP. Principal component analysis (PCA) using gene expression data from all the

samples was used to test for differences between samples and biological replicates. The PCA

revealed that the 14 DAP HT samples clustered with the 18 and 20 DAP CT samples,

correlating with the advanced embryo stage of the 14 DAP HT seeds (Fig. 5a). High-quality

clean reads were used for the differential expression gene (DEG) and gene ontology (GO)

enrichment analysis. Many misregulated (up- and down-regulated) DEGs, shown as Venn

diagrams (Fig. S4), were identified in our comparisons of samples from the same DAP: 14HT

vs. 14CT, 18HT vs. 18CT, and 20NSHT vs. 20SHT and 20CT, and of samples from the same

embryo stage (14HT vs. 18CT). The significant GO terms associated with the transcriptional

response of the seeds to HT conditions are related to cell cycle, development, hormonal

regulation, heat stress and changes in metabolic pathways (Fig. S5, Table S3).

To better understand the relationship between the HT-induced SCR phenotype, hormonal

deregulation, and seed dormancy, we first focused the analysis on the differential expression

of the dormancy gene markers DORMANCY ASSOCIATED gene 1 (DRM1) (Rae *et al.*, 2013) and

FRUCTOSE 1,6-BIPHOSPHATE ALDOLASE 6 (FBA6) (Lu *et al.*, 2012), genes involved in ABA

metabolism and signalling, and previously identified transcription factors involved in

dormancy regulation. We did not observe any significant difference between the 20SHT and

20NSHT (Fig. 2d,e; Fig. 5b; Table S2), possibly because the molecular information leading to

the SCR might already be encoded in younger seeds. The BnaDRM1 LOC106451187 and

LOC106437808 genes are highly expressed in dormant tissues (Rae *et al.*, 2013) and were

downregulated in 14HT and 20HT seeds (Fig. 2d). The BnaFBA6 LOC106391482 gene, which is involved in the sugar metabolism pathway (Lu *et al.*, 2012), was upregulated in 14HT and 20HT seeds. The misregulation of these two dormancy markers indicates that seeds under HT may have a reduced dormancy state. Expression of the following enzymes involved in ABA biosynthesis (Wu *et al.*, 2023), zeaxanthin epoxidase BnaABA1 (LOC106390346), 9-cis-epoxycarotenoid dioxygenase BnaNCED9 (LOC106354700), dehydrogenase reductase BnaABA2 (LOC106390565), and abscisic aldehyde oxidase BnaAAO3 (LOC106446766), was significantly lower at 14HT but no significant differences were observed as seeds matured from 14 to 20 DAP (Fig. 2e). The expression of BnaCYP707A1 (LOC111208232) and BnaCYP707A4 (LOC106416608), which are involved in the ABA degradation pathway to form PA and DPA, was significantly increased at 20SHT (Fig. 2e), which correlated with the reduced ABA concentration in 20 DAP seeds from HT (Fig. 2a). Although the expression profiles of genes involved in ABA metabolism support the reduced ABA levels in the seeds under HT, the expression of ABA signalling genes BnaABI3 (LOC106361788 and LOC1064188223) and BnaABI4 (LOC106443577), which promote seed dormancy, was increased in 14HT seeds, while BnaABI5 (LOC106451127) was decreased (Fig. 2e). In older seeds, their expression was not affected by HT. We further examined the expression pattern of some transcription factors reported to regulate seed dormancy and seed germination. The expression of the AUXIN RESPONSE FACTORs ARF10 and ARF16, which directly promote the expression of ABI3 (Liu *et al.*, 2013) was not significantly changed in any of the comparisons (Fig. 5b). The AP2 transcription factor CHOTTO 1 (CHO1)/PLETHORA 5 (PLT5)/AINTEGUMENTA-like 5 (AIL5) acts downstream of ABI4 during seed germination (Yamagishi *et al.*, 2009; Yano *et al.*, 2009). Similar to ABI4, its expression was upregulated at 14HT and insignificant at 20 DAP (Fig. 5b). DELAY OF GERMINATION 1 (DOG1) (Carrillo-Barral *et al.*, 2020), REDUCED DORMANCY 5 (RDO5)/DOG18 (Xiang *et al.*, 2014), RGA-like 2 (RGL2), and PHYTOCHROME INTERACTING FACTOR 3-like 5 (PIL5) (Oh *et al.*, 2007), which are reported to promote dormancy, and SPATULA (SPT) (Vaistij *et al.*, 2013), which plays an antagonistic role in dormancy regulation, were up-regulated in 14HT, 18HT and 20HT seeds (Fig. 5b; Fig. S6; Table S4). HONSU, a seed-specific protein phosphatase 2C that negatively regulates seed dormancy (Kim *et al.*, 2013), was up-regulated in 14HT seeds, but down-regulated in later stages (Fig. 5b). The regulators and markers of seed germination were up-regulated in HT seeds, supported by the overall reduction in ABA levels in the whole seeds entering the maturation stage. However, the

regulators of seed dormancy were also up-regulated at all stages. These data suggest that the transcriptional behaviour of seed dormancy and germination regulators in HT-developing seeds is contradictory, making it difficult to draw conclusions.

Fig. 5 Transcriptional changes in Topas seeds associated with growth at HT indicate changes in cell wall composition. (a) Principal component analysis using gene expression data from the samples tested for differences between samples and biological replicates. (b) Heatmap showing differential transcription of selected genes involved in seed dormancy in 14 DAP, 18 DAP, 20 DAP intact and 20 DAP SCR seeds. Expression in 14 DAP HT seeds is compared with expression in 18 DAP CT seeds. (c) GO terms related to cell wall associated with the transcriptional response of Topas seeds to HT are shown. (d) Heatmap showing the differential transcription of selected genes involved in pectin metabolism in 14 and 18 DAP seeds grown at CT and HT. The expression in 14 DAP HT seeds is compared with the expression in 18 DAP CT seeds. The full dataset is presented in Tables S1, S2 and S3.

The composition and properties of the seed coat cell wall are altered in HT-grown seeds

Gene ontology enrichment analysis revealed significant misregulation in many pathways related to plant cell walls (Fig. 5c). Interestingly, these included secondary cell wall biogenesis, cell wall thickening and modification, cellulose, pectin, and xyloglucan metabolism and modification pathways. Looking at the DEGs associated with pectin metabolic pathways, we

identified in 14HT and 18HT seeds the misregulation of enzymes such as galacturonosyltransferases (GAUT) involved in pectin biosynthesis (Mohnen, 2008), pectinmethylesterases and pectinmethylesterase inhibitors, involved in the chemical modifications of pectin (Wormit & Usadel, 2018; Wu *et al.*, 2018; Xu *et al.*, 2022), and polygalacturonases and pectate lyases, involved in pectin degradation (Marin-Rodriguez, 2002; Park *et al.*, 2010; Safran *et al.*, 2023) (Fig. 5d). This suggests that heat stress may regulate the chemical modification of pectin, a major constituent of the cell wall.

To confirm whether the transcriptional changes in the cell wall-modifying enzymes affect the pectin and lignin levels in the seed coat of HT-grown seeds, we used Ruthenium Red, which stains the polyanionic molecules in the deesterified monomers of pectin in red (Nagayama *et al.*, 2019), and Safranin-O, which has a high affinity for the phenolic compounds in lignin and stains the walls bright red (Bond *et al.*, 2008) (Fig. 6a,b). The cells of the palisade layer in the seed coat of 14 and 18 DAP Topas seeds grown under HT were strongly stained red with Ruthenium Red and Safranin-O compared to the 14, 18 and 22 DAP seeds grown under CT. The staining intensity increased with seed age. These observations suggest that seeds entering the maturation phase accumulate more secondary cell wall elements in the seed coat in HT than in CT.

Seed coat cell wall composition determines its mechanical strength (Mendu *et al.*, 2011; Sechet *et al.*, 2018). The presence of pectin and hemicellulose in the seed coat layers was investigated by immunostaining the seed coat with selective antibodies (Fig. 6d-g). For pectin, which is mainly made of homogalacturonan (HG), we used the antibody LM19 that binds preferentially to demethylesterified HG, the antibody LM20 that binds preferentially to methylesterified HG, and the antibody JIM5, which has a high affinity for partially demethylesterified HG (Fig. 6d-f). For hemicellulose, composed mainly of xyloglucan, the LM25 antibody was used and combined with a pectolyase treatment to digest the pectin that masks the hemicellulose (Fig. 6g). Fluorescence was quantified in the epidermal and palisade layers of the seed coat. The seed coats of HT-developed 14 and 18 DAP seeds had significantly higher levels of fluorescence of the LM19 and JIM5 antibody signals in the palisade layer, indicating an increase in demethylesterified HG. In contrast, no significant differences in the fluorescence intensity of the LM20 antibody signal were found in the cells of the palisade layer in the testa of CT and HT seeds. In the epidermal layer of seeds developed under HT, the signal for the JIM5 and LM19 antibodies was significantly higher in 14 DAP seeds, whereas the

signal for the LM20 antibody was significantly reduced at 18 DAP, indicating a reduction in
the amount of methylesterified HG. The results indicate an early enrichment of pectin and an
early shift towards demethylesterified pectin in the cell walls of the palisade and epidermal
cells in 14 DAP seeds grown under HT, with a significant reduction in the amount of
methylesterified HG molecules. LM25 fluorescence was significantly higher only in the
palisade cells of 18 DAP HT-grown seeds compared to CT seeds, suggesting a slight effect of
temperature on the amount of xyloglucan in the seed coat of these seeds (Fig. 6g).

To confirm our microscopic observations, we performed enzymatic fingerprinting of the cell
wall of the isolated seed coats from 10, 14 and 18 DAP seeds of Topas grown at CT and HT.
Digestion of HG by an endo-polygalacturonase between unmethylesterified ester stretches of
the galacturonic acid (GalA) releases several oligogalacturonans (OGs). OGs have been termed
GalAxMeyAcz based on the degree of polymerisation of GalA and the number of methyl and
acetyl groups. In our analysis of the Topas seed coat, we identified GalA3, GalA3ox, GalA3Xyl,
GalA3Me, GalA3MeAc, GalA4Ac, GalA4Me, GalA4Meox, GalA4Me-H_2O, and DPx>4Mey
(methylesterified OGs with degree of polymerisation greater than 4). The total peak area
obtained from the MS analysis was plotted by grouping OGs with and without methyl groups
(Fig. 6h). A significant increase of unmethylesterified OGs was released from the cell walls of
14 DAP seed coat at HT compared to CT, which correlates with the signal intensity of the LM19
antibody binding to the unmethylesterified HGs in the cell walls in the seed coat layers (Fig.
6i,j). Similarly, a significant increase in the methylesterified OGs in the seed coats of 14 DAP
with varying numbers of methyl groups correlates with the signal intensity of JIM5 antibody
binding to partially methylesterified HGs in the cell walls of the epidermal and palisade layers
(Fig. 6i,j). It is worth noting the release of oxidized OGs, which might reflect the presence of
long oxidized OGs trapped with the pectin and not solubilized through alcohol extraction.
Regarding xyloglucan, LXXGAc levels did not change significantly and correlated with LM25
signal in the seed coat (Fig. 6k,l). However, a significant increase in the level of fucosylated
subunits is observed in the 14 DAP seed coats of HT compared to CT (Fig. 6k).

Using a variety of approaches, we found that growth on HT modified the cell wall properties
in the seed coat of seeds from 14 DAP onwards. The cell walls of the epidermal and palisade
cells accumulate more pectin and shift prematurely towards the production of
demethylesterified pectin, possibly making the cell wall stiffer.

Fig. 6 The composition and properties of the seed coat cell wall are altered in HT-grown seeds. (a, b) Histological sections of Topas seeds focused on the seed coat, stained with Ruthenium Red for pectin (a) and Safranin-O for lignin (b). Sections of 14, 18, and 22 DAP CT seeds and 14 and 18 DAP HT seeds are shown. Scale bars represent 200 μm. (d-g) Immunolocalization of pectin and xyloglucans using LM19, JIM5, LM20 and LM25 antibodies on the testa of 14 and 18 DAP HT and CT seeds. The signal is visible in green, the cell structure is marked in magenta with calcofluor white. Scale bars represent 100 μm. Fluorescence intensity was measured in epidermal and palisade cells in 10, 14 and 18 DAP HT and CT seeds. Asterisks indicate statistically significant differences in HT and between seed ages in a one-way ANOVA test with Tukey multiple comparisons (*, **, *** and **** correspond to p-values of $0.05 > p > 0.01$, $0.01 > p > 0.001$, $0.001 > p > 0.0001$ and $p < 0.0001$, respectively); n.s., not significant. (h) Pectin composition of the seed coat cell wall of 10, 14, and 18 DAP seeds from HT and CT. The graph shows the measurements expressed as peak area from the MS analysis for GalA3, GalA3ox, GalA3Xyl, GalA3Me, GalA3MeAc, GalA4Ac, GalA4Me, GalA4Meox, GalA4Me-H_2O and DPx>4Mey (methylesterified OGs with degree of polymerisation greater than 4). (i) Graph showing the total intensity signal for pectin from (d-f). (j) Graph showing the amount grouped as methylesterified versus non-methylesterified OGs from (h). (k) Graph showing the measurements presented as peak area from the MS analysis for xyloglucans. (l) Graph showing the total intensity signal for xyloglucan from (g). Samples with the same letter were not statistically significantly different. (m) A 3-mm silicon tube was applied to siliques from 10 DAP CT and 12 DAP HT. Dry seeds were quantified for the number of total viable, SCR and dead seeds per silique. Asterisks indicate statistically significant differences in HT and between seed ages in a two-way ANOVA test with Tukey multiple comparisons (*, **, *** and **** correspond to p-values of $0.05 > p > 0.01$, $0.01 > p > 0.001$, and $p < 0.001$, respectively); n.s., not significant.

Application of mechanical constraints during silique development partially rescues SCR phenotype

Since the seed coat thickness of seeds developed under HT was significantly reduced, we suspected that such a seed coat would lack the strength to accommodate the developing embryo. The continuous embryo growth would exert additional mechanical pressure on the seed coat, resulting in SCR. To verify this, we applied a silicone tube to developing siliques to

provide an external mechanical constraint on the seeds in an attempt to compensate for the

early embryonic pressure on the seed coat (Fig. 6m). Such an experiment was previously

performed to show that an applied mechanical force on the siliques triggers seed maturation

by arresting embryonic growth (Rolletschek *et al.*, 2024). The presence of the silicone tube on

the siliques of CT plants did not affect the seed viability or promote SCR. In HT, the external

force exerted by the silicone tube on the silique (and thus on the seeds inside) resulted in a

significant decrease in the number of seeds undergoing SCR and an increase in the number of

viable seeds (Fig. 6m). The application of an external mechanical support reinforced the

compromised mechanical strength of the seed coat caused by the faster embryo

development.

DISCUSSION

Rising global temperatures adversely affect the production of major staple crops (Agnolucci

et al., 2020; Jägermeyr *et al.*, 2021), by hindering proper seed development, as in *Brassica*

napus (Mácová *et al.*, 2022). The occurrence of temperature-induced preharvest seed

sprouting was reported in our previous study in three spring cultivars of *B. napus* (Mácová *et*

al., 2022). In particular, prolonged cultivation of Topas at temperatures above the suboptimal

temperature resulted in half of the seeds per silique undergoing SCR. In this study, we have

elucidated the molecular mechanism behind the regulation of the SCR phenotype that occurs

in Topas during seed maturation.

Genetic screens in *Arabidopsis* have identified the molecular components involved in seed

maturation and dormancy (Finkelstein *et al.*, 2002; Holdsworth *et al.*, 2008). Reduced ABA

levels or increased levels of active GAs result in reduced seed dormancy (Benech-Arnold &

Rodríguez, 2018; Tai *et al.*, 2021). Although HT conditions during seed maturation (after 20

DAP) in Topas resulted in reduced ABA levels, suggesting reduced seed dormancy, there were

no differences between SCR and non-SCR seeds. Furthermore, the failure of *in vitro* ABA

treatment to rescue SCR led to the hypothesis of a minimal role for ABA in SCR. No direct

conclusions could be drawn linking the state of dormancy to the SCR phenotype, as even the

genes reported to positively regulate seed dormancy, such as *ABI3, ABI4, DOG1, RDO5, CHO1*,

and *SPT*, have increased expression levels in seeds developed on HT, but still lead to SCR.

Growth at elevated temperatures accelerated embryo development in *B. napus* from early

developmental stages (Mácová *et al.*, 2022). Indeed, most of the embryos from 6 DAP seeds

produced at optimal growth temperature were at the 8-cell stage, whereas at elevated temperatures, they were at the mid to late globular stage. A similar difference was observed in the maturation phase, with a difference of 4 days for equivalent developmental stages. As a result, embryos were larger in seeds that did not enlarge, possibly exerting mechanical pressure on the seed coat. Mechanical constraints applied to developing seeds of *B. napus* cv. Reston promoted seed maturation by activating the storage metabolism and the expression of genes involved in the embryo maturation process (Rolletschek *et al.*, 2024). The higher levels of ABA in 14 and 18 DAP embryos developed under HT and the increased expression pattern of *ABI3* and *DOG1* in HT seeds during early maturation suggest that seed development under elevated growth temperature could promote an early maturation process (Baud *et al.*, 2016; Leprince *et al.*, 2016; Rolletschek *et al.*, 2024).

The proper exchange of signals between the seed coat, endosperm, and embryo interacts to define the final size, shape, and dormancy state of the seed at the end of the maturation phase (Li *et al.*, 2019). The *Arabidopsis iku2-2 zou-4* double mutant has reduced seed viability due to developing seeds undergoing testa rupture (Creff *et al.*, 2015), similar to the SCR phenotype observed in Topas. The mutation in *HAIKU2 (IKU2)* results in increased endosperm pressure, favouring mechanosensitive precocious testa stiffening, possibly dependent on demethylesterified pectin accumulation. This results in small *iku2* seeds (Creff *et al.*, 2023). Seeds of *zhoupi-4 (zou-4)* have a persistent endosperm at the torpedo stage, even as the embryo expands in size, producing large seeds during development (Creff *et al.*, 2015). Testa rupture in *iku-2 zou-4* seeds would be a consequence of early testa stiffening together with increased pressure from the developing embryo and a persistent endosperm. Both *IKU2* and *ZOU* expression were down-regulated in Topas seeds developed at high temperatures (Fig. S6 and Table S2), suggesting that a similar process may occur in *B. napus*.

GO enrichment analysis of the seed transcriptome revealed a temperature-dependent misregulation of cell wall-related pathways in seeds. Pathways related to cellulose, pectin and xyloglucan metabolism, cell wall thickening, and pectin modification were overrepresented. The primary cell wall is composed of cellulose, hemicellulose, and pectin (Cosgrove, 2024), while the secondary cell wall can be stiffened with lignin. Cellulose and hemicellulose are embedded in a gel matrix formed by pectin and influence the extensibility of the cell wall, an essential feature for cell growth and morphogenesis (Peaucelle *et al.*, 2011). Many genes involved in pectin biosynthesis, modification, and degradation were misregulated, which led

us to investigate the methylesterification pattern of pectin, which could influence wall

rheology. Ruthenium Red staining for pectin and Safranin-O staining for lignin indicated that

both pectin and lignin levels were increased in the palisade layer of HT seed coats.

Immunostaining with pectin-specific antibodies and cell wall profiling of the seed coat of early

maturing seeds indicated that the overall pectin composition was altered, particularly in the

palisade layer. The significant enrichment of demethylesterified pectin in the palisade layer

of early maturing seeds developed under HT suggests that the seed coat cell walls may be

stiffer when grown at elevated temperatures.

The embryo stage and size difference observed in HT seeds may exert increased pressure on

the walls of the developing seed coat for the same overall seed size, resulting in increased

tension between adjacent cells within an integument layer. The reduction in the thickness of

the epidermal, palisade and endothelial cell layers of the seed coat observed in HT seeds

suggests that increased tension was exerted between adjacent cells and that the seed coat

cells resisted the tension by cell stretching rather than cell division. Such stress-dependent

stiffening of seed coat cell walls has been shown in *iku2* mutants (Creff *et al.*, 2023). Thus,

increased stiffness of the seed coat cell walls would resist strain between seed coat cells to

compensate for the pressure exerted by the developing embryo. Tight regulation of these

opposing forces governs the wall mechanics required for seed growth. Mechanical support of

Topas seeds developed under HT by applying a silicone tube partially rescued SCR, strongly

suggesting that the seed coat layers lack the strength to withstand the tension from the

pressure exerted by the developing embryo. The exact process by which increased pectin

demethylesterification plays a role in the wall mechanics of *Brassica napus* seeds is unclear.

Our data suggest two possible mechanisms underlying the heat-induced SCR phenotype. First,

exposure of seeds to elevated temperatures leads to changes in the seed coat cell wall,

making it more susceptible to rupture under the pressure exerted by the expanding embryo.

Second, HT-dependent embryo expansion increases the tension on the seed coat and may

promote the observed changes in cell wall composition, similar to what happens in

Arabidopsis. However, these changes may not be sufficient to prevent SCR due to the intense

pressure exerted by the growing embryo.

ACKNOWLEDGEMENTS

This work was financially supported by the project TowArds Next GENeration Crops, reg. no.
CZ.02.01.01/00/22_008/0004581 of the ERDF Programme Johannes Amos Comenius. We
acknowledge the Bioinformatics (supported by the NCLG research infrastructure LM2023067
funded by MEYS CR), Cell Imaging (supported by MEYS CR LM2023050 Czech-BioImaging), and
Plant Sciences Core Facilities of CEITEC Masaryk University for their support in obtaining the
scientific data presented in this paper. This work has benefited from the support of IJPB's
Plant Observatory platforms Po-Chem. The IJPB benefits from the support of Saclay Plant
Sciences SPS (ANR-17-EUR-0007). This work was also supported by a grant from the French
ANR (MECHASEED, ANR-23-CE13-0009) to BL.

COMPETING INTERESTS

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

HSR secured the funding. UK and HSR conceived the project, designed the experiments, and
interpreted the data. UK performed the seed phenotyping, histology, ABA treatments,
transcriptomics, and immunolabelling of the seed coat. IU and JH optimised the protocol for
Technovit embedding of oilseed rape seed and prepared the slides for histology. AMM
analysed the transcriptomic data. AC and BL performed the immunolabelling of the seed coat.
AV performed the cell wall profiling. IP, AP and ON performed the hormonal profiling. UK and
HSR drafted the manuscript. All authors have proofed and approved the manuscript.

DATA AVAILABILITY

The data generated during the current study are included in this published article (and its
Supporting   Information   file)   and   are   available   in   the   Zenodo   repository
[https://zenodo.org/doi/10.5281/zenodo.14224074]. The dataset supporting the conclusions
of this article is deposited to the NCBI repository (BioProject accession number
PRJNA1188476, https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1188476).

REFERENCES

Abu-Jamous B, Kelly S. **2018**. Clust: automatic extraction of optimal co-expressed gene clusters from
gene expression data. *Genome Biology* **19**: 172.

Agnolucci P, De Lipsis V. **2020**. Long-run trend in agricultural yield and climatic factors in Europe.
Climatic Change **159**: 385–405.

Agnolucci P, Rapti C, Alexander P, De Lipsis V, Holland RA, Eigenbrod F, Ekins P. **2020**. Impacts of
rising temperatures and farm management practices on global yields of 18 crops. *Nature Food* **1**:
562–571.

Andrews S. **2010**. FastQC: A Quality Control Tool for High Throughput Sequence Data.

Baud S, Kelemen Z, Thévenin J, Boulard C, Blanchet S, To A, Payre M, Berger N, Effroy-Cuzzi D,
Franco-Zorrilla JM, *et al.* 2016. Deciphering the molecular mechanisms underpinning the
transcriptional control of gene expression by L-AFL proteins in Arabidopsis seed. *Plant Physiology*:
pp.00034.2016.

Benech-Arnold RL, Rodríguez M V. 2018. Pre-harvest Sprouting and Grain Dormancy in Sorghum
bicolor: What Have We Learned? *Frontiers in Plant Science* **9**.

Bolger AM, Lohse M, Usadel B. **2014**. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics **30**: 2114–2120.

Bond J, Donaldson L, Hill S, Hitchcock K. **2008**. Safranine fluorescent staining of wood cell walls.
Biotechnic & Histochemistry **83**: 161–171.

Broad Institute, GitHub Repository. **2018**. Picard Toolkit.

Carrillo-Barral N, Rodríguez-Gacio M del C, Matilla AJ. **2020**. Delay of Germination-1 (DOG1): A Key
to Understanding Seed Dormancy. *Plants* **9**: 480.

Chu J, Sadeghi S, Raymond A, Jackman SD, Nip KM, Mar R, Mohamadi H, Butterfield YS, Robertson
AG, Birol I. **2014**. BioBloom tools: fast, accurate and memory-efficient host species sequence
screening using bloom filters. *Bioinformatics* **30**: 3402–3404.

Cosgrove DJ. **2024**. Structure and growth of plant cell walls. *Nature Reviews Molecular Cell Biology*
25: 340–358.

Creff A, Ali O, Bied C, Bayle V, Ingram G, Landrein B. **2023**. Evidence that endosperm turgor
pressure both promotes and restricts seed growth and size. *Nature Communications* **14**.

Creff A, Brocard L, Ingram G. **2015**. A mechanically sensitive cell layer regulates the physical
properties of the Arabidopsis seed coat. *Nature Communications* **6**.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.

Ewels P, Magnusson M, Lundin S, Käller M. **2016**. MultiQC: summarize analysis results for multiple
tools and samples in a single report. *Bioinformatics* **32**: 3047–3048.

Finkelstein RR, Gampala SSL, Rock CD. **2002**. Abscisic Acid Signaling in Seeds and Seedlings. *The*
Plant Cell **14**: S15–S45.

Heino M, Kinnunen P, Anderson W, Ray DK, Puma MJ, Varis O, Siebert S, Kummu M. **2023**.
Increased probability of hot and dry weather extremes during the growing season threatens global
crop yields. *Scientific Reports* **13**: 3583.

Holdsworth MJ, Bentsink L, Soppe WJJ. **2008**. Molecular networks regulating Arabidopsis seed
maturation, after-ripening, dormancy and germination. *New Phytologist* **179**: 33–54.

Jägermeyr J, Müller C, Ruane AC, Elliott J, Balkovic J, Castillo O, Faye B, Foster I, Folberth C, Franke
JA, *et al.* 2021. Climate impacts on global agriculture emerge earlier in new generation of climate
and crop models. *Nature Food* **2**: 873–885.

Jiang Y, Deyholos MK. **2010**. Transcriptome analysis of secondary-wall-enriched seed coat tissues of
canola (Brassica napus L.). *Plant Cell Reports* **29**: 327–342.

Kassambara A. **2018**. 'ggpubr:"ggplot2" based publication ready plots.' R package version 0.17.

Kim W, Lee Y, Park J, Lee N, Choi G. **2013**. HONSU, a Protein Phosphatase 2C, Regulates Seed
Dormancy by Inhibiting ABA Signaling in Arabidopsis. *Plant and Cell Physiology* **54**: 555–572.

Kolde R. **2015**. Package 'pheatmap'. R Package 1.7 .

Leprince O, Pellizzaro A, Berriri S, Buitink J. **2016**. Late seed maturation: drying without dying.
Journal of Experimental Botany: erw363.

Li N, Xu R, Li Y. **2019**. Molecular Networks of Seed Size Control in Plants. *Annual Review of Plant*
Biology **70**: 435–463.

Liao Y, Smyth GK, Shi W. **2014**. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30**: 923–930.

Lima RB, Pankaj R, Ehlert ST, Finger P, Fröhlich A, Bayle V, Landrein B, Sampathkumar A,
Figueiredo DD. **2024**. Seed coat-derived brassinosteroid signaling regulates endosperm
development. *Nature Communications* **15**: 9352.

Liu X, Zhang H, Zhao Y, Feng Z, Li Q, Yang H-Q, Luan S, Li J, He Z-H. **2013**. Auxin controls seed
dormancy through stimulation of abscisic acid signaling by inducing ARF-mediated *ABI3* activation in
Arabidopsis. *Proceedings of the National Academy of Sciences* **110**: 15485–15490.

Love MI, Huber W, Anders S. **2014**. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biology* **15**: 550.

Lu W, Tang X, Huo Y, Xu R, Qi S, Huang J, Zheng C, Wu C. **2012**. Identification and characterization of
fructose 1,6-bisphosphate aldolase genes in Arabidopsis reveal a gene family with diverse responses
to abiotic stresses. *Gene* **503**: 65–74.

Mácová K, Prabhullachandran U, Štefková M, Spyroglou I, Pěnčík A, Endlová L, Novák O, Robert
HS. **2022**. Long-Term High-Temperature Stress Impacts on Embryo and Seed Development in
Brassica napus. *Frontiers in Plant Science* **13**.

Marin-Rodriguez MC. **2002**. Pectate lyases, cell wall degradation and fruit softening. *Journal of*
Experimental Botany **53**: 2115–2119.

Mendu V, Griffiths JS, Persson S, Stork J, Downie AB, Voiniciuc C, Haughn GW, DeBolt S. **2011**.
Subfunctionalization of Cellulose Synthases in Seed Coat Epidermal Cells Mediates Secondary Radial
Wall Synthesis and Mucilage Attachment   . *Plant Physiology* **157**: 441–453.

Mohnen D. **2008**. Pectin structure and biosynthesis. *Current Opinion in Plant Biology* **11**: 266–277.

Nagayama T, Nakamura A, Yamaji N, Satoh S, Furukawa J, Iwai H. **2019**. Changes in the Distribution
of Pectin in Root Border Cells Under Aluminum Stress. *Frontiers in Plant Science* **10**.

National Aeronautics and Space Administration. **2024**. Global temperature. *Goddard Institute for*
Space Studies.

Oh E, Yamaguchi S, Hu J, Yusuke J, Jung B, Paik I, Lee H-S, Sun T, Kamiya Y, Choi G. **2007**. PIL5, a
Phytochrome-Interacting bHLH Protein, Regulates Gibberellin Responsiveness by Binding Directly to
the *GAI* and *RGA* Promoters in *Arabidopsis* Seeds. *The Plant Cell* **19**: 1192–1208.

Okonechnikov K, Conesa A, García-Alcalde F. **2016**. Qualimap 2: advanced multi-sample quality
control for high-throughput sequencing data. *Bioinformatics* **32**: 292–294.

Park K-C, Kwon S-J, Kim N-S. **2010**. Intron loss mediated structural dynamics and functional
differentiation of the polygalacturonase gene family in land plants. *Genes & Genomics* **32**: 570–577.

Peaucelle A, Braybrook SA, Le Guillou L, Bron E, Kuhlemeier C, Höfte H. **2011**. Pectin-Induced
Changes in Cell Wall Mechanics Underlie Organ Initiation in Arabidopsis. *Current Biology* **21**: 1720–
1726.

Pěnčík A, Casanova-Sáez R, Pilařová V, Žukauskaitė A, Pinto R, Micol JL, Ljung K, Novák O. **2018**.
Ultra-rapid auxin metabolite profiling for high-throughput mutant screening in Arabidopsis. *Journal*
of Experimental Botany **69**: 2569–2579.

Rae GM, David K, Wood M. **2013**. The Dormancy Marker DRM1/ARP Associated with Dormancy but
a Broader Role In Planta. *Developmental Biology Journal* **2013**: 1–12.

Rolletschek H, Muszynska A, Borisjuk L. **2021**. The process of seed maturation is influenced by
mechanical constraints. *New Phytologist* **229**: 19–23.

Rolletschek H, Muszynska A, Schwender J, Radchuk V, Heinemann B, Hilo A, Plutenko I, Keil P,
Ortleb S, Wagner S, *et al.* **2024**. Mechanical forces orchestrate the metabolism of the developing
oilseed rape embryo. *New Phytologist* **244**: 1328–1344.

Safran J, Tabi W, Ung V, Lemaire A, Habrylo O, Bouckaert J, Rouffle M, Voxeur A, Pongrac P,
Bassard S, *et al.* **2023**. Plant polygalacturonase structures specify enzyme dynamics and
processivities to fine-tune cell wall pectins. *The Plant Cell* **35**: 3073–3091.

Secchi MA, Fernandez JA, Stamm MJ, Durrett T, Prasad PVV, Messina CD, Ciampitti IA. **2023**.
Effects of heat and drought on canola (Brassica napus L.) yield, oil, and protein: A meta-analysis.
Field Crops Research **293**: 108848.

Sechet J, Marion-Poll A, North HM. **2018**. Emerging Functions for Cell Wall Polysaccharides
Accumulated during Eudicot Seed Development. *Plants* **7**: 81.

Šimura J, Antoniadi I, Široká J, Tarkowská D, Strnad M, Ljung K, Novák O. **2018**. Plant
Hormonomics: Multiple Phytohormone Profiling by Targeted Metabolomics. *Plant Physiology* **177**:
476–489.

Tai L, Wang HJ, Xu XJ, Sun WH, Ju L, Liu WT, Li WQ, Sun J, Chen KM. **2021**. Pre-harvest sprouting in
cereals: Genetic and biochemical mechanisms. *Journal of Experimental Botany* **72**: 2857–2876.

Turbant A, Fournet F, Lequart M, Zabijak L, Pageau K, Bouton S, Van Wuytswinkel O. **2016**. PME58
plays a role in pectin distribution during seed coat mucilage extrusion through homogalacturonan
modification. *Journal of Experimental Botany* **67**: 2177–2190.

Vaistij FE, Gan Y, Penfield S, Gilday AD, Dave A, He Z, Josse E-M, Choi G, Halliday KJ, Graham IA.
2013. Differential control of seed primary dormancy in *Arabidopsis* ecotypes by the transcription
factor SPATULA. *Proceedings of the National Academy of Sciences* **110**: 10866–10871.

**Voxeur A, Habrylo O, Guénin S, Miart F, Soulié M-C, Rihouey C, Pau-Roblot C, Domon J-M,
Gutierrez L, Pelloux J, *et al.* 2019**. Oligogalacturonide production upon *Arabidopsis thaliana* –
Botrytis cinerea interaction. *Proceedings of the National Academy of Sciences* **116**: 19743–19752.

Wang L, Wang S, Li W. **2012**. RSeQC: quality control of RNA-seq experiments. *Bioinformatics* **28**:
2184–2185.

Wickham H. **2011**. ggplot2. *WIREs Computational Statistics* **3**: 180–185.

Wormit A, Usadel B. **2018**. The Multifaceted Role of Pectin Methylesterase Inhibitors (PMEIs).
International Journal of Molecular Sciences **19**: 2878.

Wu H-C, Bulgakov VP, Jinn T-L. **2018**. Pectin Methylesterases: Cell Wall Remodeling Proteins Are
Required for Plant Response to Heat Stress. *Frontiers in Plant Science* **9**: 1612.

Wu W, Cao S, Shi L, Chen W, Yin X, Yang Z. **2023**. Abscisic acid biosynthesis, metabolism and
signaling in ripening fruit. *Frontiers in Plant Science* **14**: 1279031.

Xiang Y, Nakabayashi K, Ding J, He F, Bentsink L, Soppe WJJ. **2014**. REDUCED DORMANCY5 Encodes
a Protein Phosphatase 2C That Is Required for Seed Dormancy in *Arabidopsis*. *The Plant Cell* **26**:
4362–4375.

Xu Z, Yang M, Li Z, Xiao J, Yang X, Wang H, Wang X. **2022**. Tissue-specific pectin methylesterification
and pectin methylesterase activities play a role in lettuce seed germination. *Scientia Horticulturae*
301: 111134.

**Yamagishi K, Tatematsu K, Yano R, Preston J, Kitamura S, Takahashi H, McCourt P, Kamiya Y,
Nambara E**. **2009**. CHOTTO1, a Double AP2 Domain Protein of Arabidopsis thaliana, Regulates
Germination and Seedling Growth Under Excess Supply of Glucose and Nitrate. *Plant and Cell
Physiology* **50**: 330–340.

Yano R, Kanno Y, Jikumaru Y, Nakabayashi K, Kamiya Y, Nambara E. **2009**. CHOTTO1, a Putative
Double APETALA2 Repeat Transcription Factor, Is Involved in Abscisic Acid-Mediated Repression of
Gibberellin Biosynthesis during Seed Germination in Arabidopsis   . *Plant Physiology* **151**: 641–654.

Young LW, Wilen RW, Bonham-Smith PC. **2004**. High temperature stress of Brassica napus during
flowering reduces micro- and megagametophyte fertility, induces fruit abortion, and disrupts seed
production. *Journal of Experimental Botany* **55**: 485–495.

Zhao C, Liu B, Piao S, Wang X, Lobell DB, Huang Y, Huang M, Yao Y, Bassu S, Ciais P, *et al.* 2017.
Temperature increase reduces global yields of major crops in four independent estimates.
Proceedings of the National Academy of Sciences of the United States of America **114**: 9326–9331.

    FIGURE LEGENDS

Fig. 1. SCR phenotype observed in Topas seeds developed under HT. (a, b) 26 DAP seeds (a)

and their isolated embryos (b) grown under CT. (c, d) 26 DAP seeds (c) and their isolated

embryos (d) grown under HT. SCR was observed (*, Δ, +, ●). * Indicates seeds with testa

rupture with intact embryo, Δ indicates testa rupture with radicle protrusion, + indicates testa

rupture with cotyledon protrusion, and ● indicates embryos with abnormal development. (e)

Quantification of the number of dry seeds per silique from CT (blue) and HT (red) showing the

number of total, viable, SCR and dead seeds. Asterisks indicate statistically significant

differences in HT in a multiple t-test using the Holm-Sidak method (**** corresponds to p-

values of $p < 0.0001$. (f) Embryo stage of seeds developed on CT (top) and HT (bottom) from

14 to 20 DAP. The isolated embryo shown on the right is from the seed on the left. Scale bars:

2 mm (a-d) and 3mm (f).

Fig. 2. ABA levels and expression of genes involved in ABA metabolism in *Brassica napus* Topas

seeds. (a) Seeds were from 14 DAP to 36 DAP of CT (blue) and HT (red, dark red). From 20

DAP, HT seeds were separated into intact seeds (red) and SCR seeds (dark red). Quantification

is presented as pmol per g fresh weight. (b) Measurements were performed for ABA, 7-

hydroxy-ABA, phaseic acid (PA), and dihydrophaseic acid (DPA) from embryos and seed coats

from 14 DAP and 18 DAP seeds grown in CT (blue) and HT (red). Asterisks indicate statistically

significant differences in HT and between 14 DAP HT and 18 DAP CT in a multiple t-test using

the Holm-Sidak method (*, **, *** and **** correspond to p-values of $0.05 > p > 0.01$, 0.01

$> p > 0.001$, $0.001 > p > 0.0001$ and $p < 0.0001$, respectively); n.s., not significant. (c) 15 siliques

of 14 DAP and 18 DAP from HT and CT, respectively, were cultured in the presence or absence

of 10 μM ABA (Fig. S1). Dry seeds were quantified for the number of total viable, SCR and

dead seeds per silique. Samples with the same letter were not statistically significantly

different. (d) Relative expression of selected genes representing DORMANCY ASSOCIATED

gene 1 (DRM1) and FRUCTOSE 1,6-BIPHOSPHATE ALDOLASE 6 (FBA6) in 14 DAP (blue), 20 DAP

intact (yellow) and 20 DAP SCR (grey) seeds. Data were extracted from the transcriptome

dataset (Tables S1 and S2). (e) Heatmap showing the differential expression of selected genes

involved in ABA metabolism in 14 DAP, 20 DAP intact, and 20 DAP SCR seeds. The full dataset

is provided in Tables S1 and S2.

Fig 3. Embryo and seed size measurements of *Brassica napus* Westar and Topas seeds developed at CT and HT. (a, b) Seed (a) and embryo (b) area measurements in mm^2 of 14, 18 and 20 DAP Topas seeds developed at CT (blue) and HT (red). (c, d) Seed (c) and embryo (d) area measurements in mm^2 of 14, 18 and 20 DAP Westar seeds developed at CT (blue) and HT (red). Asterisks indicate statistically significant differences in HT and between 14 DAP HT and 18 DAP CT in a two-way ANOVA test with Sidak's multiple comparisons (*, **, *** and **** correspond to *p*-values of $0.05 > p > 0.005$, $0.005 > p > 0.0005$, $0.0005 > p > 0.0005$ and $p < 0.00005$, respectively); n.s., not significant. (e) Embryo-to-seed size ratio in 14, 18 and 20 DAP Topas and Westar seeds developed at CT and HT. Samples with the same letter were not statistically significantly different. Representative pictures of the seeds and embryos are shown in Fig. S3.

Fig. 4 The thickness of the seed coat cells is reduced in HT. (a-e) Histological sections of Topas seeds focused on the seed coat, stained with toluidine blue. (a) The seed coat is composed of an epidermal layer, a palisade layer and an endodermal layer. (b-e) Seed coat of 14 DAP seeds from CT (b) and HT (d) and 18 DAP seeds from CT (c) and HT (e). Scale bars represent 200 μm. (f-h) Box-whisker plots show the size mean, median, outliers, and SD in μm of the thickness of the epidermal layer (f), palisade layer (g) and first endothelial cell layer (h) in 10, 14, 18, and 20 DAP seeds from CT (blue) and HT (red). Samples with the same letter were not statistically significantly different. (i) Percentage of reduction in thickness as the seed aged from 14 to 20 DAP.

Fig. 5 Transcriptional changes in Topas seeds associated with growth at HT indicate changes in cell wall composition. (a) Principal component analysis using gene expression data from the samples tested for differences between samples and biological replicates. (b) Heatmap showing differential transcription of selected genes involved in seed dormancy in 14 DAP, 18 DAP, 20 DAP intact and 20 DAP SCR seeds. Expression in 14 DAP HT seeds is compared with expression in 18 DAP CT seeds. (c) GO terms related to cell wall associated with the transcriptional response of Topas seeds to HT are shown. (d) Heatmap showing the differential transcription of selected genes involved in pectin metabolism in 14 and 18 DAP seeds grown at CT and HT. The expression in 14 DAP HT seeds is compared with the expression in 18 DAP CT seeds. The full dataset is presented in Tables S1, S2 and S3.

Fig. 6 The composition and properties of the seed coat cell wall are altered in HT-grown seeds. (a, b) Histological sections of Topas seeds focused on the seed coat, stained with Ruthenium Red for pectin (a) and Safranin-O for lignin (b). Sections of 14, 18, and 22 DAP CT seeds and 14 and 18 DAP HT seeds are shown. Scale bars represent 200 μm. (d-g) Immunolocalization of pectin and xyloglucans using LM19, JIM5, LM20 and LM25 antibodies on the testa of 14 and 18 DAP HT and CT seeds. The signal is visible in green, the cell structure is marked in magenta with calcofluor white. Scale bars represent 100 μm. Fluorescence intensity was measured in epidermal and palisade cells in 10, 14 and 18 DAP HT and CT seeds. Asterisks indicate statistically significant differences in HT and between seed ages in a one-way ANOVA test with Tukey multiple comparisons (*, **, *** and **** correspond to p-values of $0.05 > p > 0.01$, $0.01 > p > 0.001$, $0.001 > p > 0.0001$ and $p < 0.0001$, respectively); n.s., not significant. (h) Pectin composition of the seed coat cell wall of 10, 14, and 18 DAP seeds from HT and CT. The graph shows the measurements expressed as peak area from the MS analysis for GalA3, GalA3ox, GalA3Xyl, GalA3Me, GalA3MeAc, GalA4Ac, GalA4Me, GalA4Meox, GalA4Me-H_2O and DPx>4Mey (methylesterified OGs with degree of polymerisation greater than 4). (i) Graph showing the total intensity signal for pectin from (d-f). (j) Graph showing the amount grouped as methylesterified versus non-methylesterified OGs from (h). (k) Graph showing the measurements presented as peak area from the MS analysis for xyloglucans. (l) Graph showing the total intensity signal for xyloglucan from (g). Samples with the same letter were not statistically significantly different. (m) A 3-mm silicon tube was applied to siliques from 10 DAP CT and 12 DAP HT. Dry seeds were quantified for the number of total viable, SCR and dead seeds per silique. Asterisks indicate statistically significant differences in HT and between seed ages in a two-way ANOVA test with Tukey multiple comparisons (*, **, *** and **** correspond to p-values of $0.05 > p > 0.01$, $0.01 > p > 0.001$, and $p < 0.001$, respectively); n.s., not significant.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Fig. S1 Representative picture of the silique culture system for ABA treatment.

Fig. S2 Profiles of auxin and its metabolites in *Brassica napus* Topas seeds, isolated embryos

and seed coats.

Fig. S3 Representative pictures of the seeds and embryos measured in Fig. 3.

Fig. S4 Venn diagrams summarising the transcriptional response of Topas seeds to HT.

Fig. S5 GO terms associated with the transcriptional response of Topas seeds to HT.

Fig. S6 Expression clusters of the transcriptional response of Topas seeds to HT.

Table S1 Full dataset for the DEGs for all the comparisons.

Table S2 DEGs for dormancy, ABA and pectin metabolism related.

Table S3 Misregulated GO terms associated with plant cell wall, hormone, cell cycle,

metabolic, heat response and developmental pathways.

Table S3 DEGs and GO terms associated with each of the clusters.

Methods S1 Co-expression cluster mapping of genes.