The swarming behaviors of Vorticella

Tetsuya Tabata[*] and Yuko Maeyama

Institute for Quantitative Biosciences, The university of Tokyo,1-1-1 Yayoi, Bunkyo-ku, Tokyo

113-0032, JAPAN

*Corresponding author

Present address:The Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kamikitazawa,

Setagaya-ku, Tokyo, 156-8506, Japan, tabata-tt@igakuken.or.jp

Abstract Two Vorticella species undergo a synchronous transition from sessile zooids to

motile telotrochs, which swarm enveloped in secreted mucus, subsequently forming dense

aggregations on substrates and reverting to the zooid form. This cyclical process recurs on a

daily basis. Each species exhibits a unique mode of swarming behavior. We hypothesize that

these behaviors may serve to facilitate efficient feeding while concurrently acting as a

mechanism for predator avoidance.

Introduction

"It has long been known that animals may aggregate into groups or clusters, more or less

closely associated, in which physical contact may, or may not, occur. Normally actual physical

contact is found as a part of the aggregation phenomenon in many Protozoa, Paramecium for

example; in flatworms such as the planarians; in earthworms; in arthropods and molluscs; and

among many vertebrates, including fish, frogs, reptiles, birds and mammals" (Allee, 1927) In

this study, we present further examples within the ciliate protozoan species, Vorticella spp.,

demonstrating an alternation between two distinct life cycle phases: the sessile, stalked-

feeding zooid and the motile swarmer, known as the telotroch. This transition is typically

attributed to environmental stressors (Barlow and Finley, 1976). However, in this study, we

demonstrate that in these particular species, the transition occurs cyclically on a daily basis,

wherein zooids synchronously transform into telotrochs, which subsequently swarm and

collectively settle or congregate on solid substrates. We had recognized this behavior has only

been publicly documented in conference abstracts by the research group of Horikami and Ishii,

one of which was in the international meeting where they reported that four Vorticella species

exhibit this type of behavior (Horikami and Ishii, 1981). During manuscript preparation, we also

uncovered earlier reports describing the gregarious Vorticella species by Fox and Newth (Fox

and Newth, 1936; Fox and NEWTH, 1935). Our current study confirms their findings and

presents the gregarious behaviors using videos, a method not yet feasible in earlier pioneering

research.

Results and Discussion

Swarming behaviors of Vorticella

We identified two species of the genus *Vorticella* from Sanshirou-ike, the pond, situated within

the University of Tokyo campus. The pond's name originates from a famous novel by Souseki

Natsume, written during the Meiji era. We primarily focus on the first species, the provisionally

named *Vorticella sp11* (Sanshirou) (Figure 1) as its ITS1-5.8S-ITS2 DNA sequence is identical

to that of *Vorticella sp11* (Figure 1-supplementary figure1), a species originating from China

that has been solely characterized by its genetic sequence (Sun et al., 2013). *Vorticella sp11*

(Sanshirou) appeared seasonally, from late April to early July. Despite efforts, we were unable

to establish stable culture conditions and could only maintain the organisms for a few weeks

post-sampling. Periodic sequencing of samples was performed to verify the genotype of the

observed individuals. Although not all samples were sequenced, those displaying the

characteristic behaviors of V. sp11 (Sanshirou) consistently exhibited identical ITS1-5.8S-ITS2

DNA sequences, without exception, despite the sequence being relatively short at

approximately 400 base pairs. We acknowledge that species identification based solely on

morphology was not possible, and thus we cannot exclude the possibility that the individuals

presented as *V. sp11* (Sanshirou) in this study may encompass heterogenous species.

We collected submerged fallen leaves from the pond's bottom and examined them

under a compound microscope to locate stalked *V. sp11* (Sanshirou) aggregating densely on

the leaves. Leaf fragments were then transferred to an aquarium with aged water containing

aquatic plants, such as *Cabomba* sp., or artificial plastic aquatic plants to which the organisms

could attach (Figure 1A). It is noteworthy that while they were densely clustered, each

organism remained independent. This aggregation on plant fragments was video-recorded at

low magnification, and a series of snapshots is provided (Figure 1B-I and figure 1-video 1). The

entire population of zooids transitioned to telotrochs almost simultaneously, detaching from

their stalks (Figure 1C) and forming a spherical cloud resembling a miniature bee swarm

(Figure 1D), as documented by Fox and Newth (1936) for *Vorticella chlorostigma*. The swarm

moved about one and a half laps along the inner wall of the well before settling into a dense

cluster (Figure 1D-I). Additional examples are presented (Figure 1-videos 2-6). The ITS1-5.8S-

ITS2 DNA sequence of *V. chlorostigma* exhibits significant divergence from that of V. sp11

(Sanshirou)(Figure 1-figure supplement 1).

It is noteworthy that the telotroch excretes a gelatinous material, suggesting that the

globular cloud is sustained by the mucus as described previously (Fox and Newth, 1936;

Horikami and Ishii, 1981). Fox and Newth further identified that a mucous filament originating

at the center of the aboral ciliary wreath (described below), which became visible with the

addition of Indian ink. This discovery is crucial in deepening our comprehension of the

mechanisms driving swarming and/or group settlement behaviors. While we were not aware of

this filament during the experiments, we observed turbid material being dragged by the moving

globular cloud (arrow in Figure 1F), which became particularly conspicuous under dark-field

illumination, potentially arising from the subsequent solidification of the mucus fibers. At times,

this material was incorporated into the settled cluster (arrow in Figure 1H). They additionally

observed: 'At the end of swarming they are so solid that sometimes groups of vorticellids settle

down on them as they would on pondweed.' (Fox and Newth, 1936). A video illustrating the

formation of telotroch clusters settling without adherence to the substrate (Figure 1-video 7)

may exemplify the phenomenon described. The formation of the mucus fibers will be

addressed in more detail later.

Life cycle of Vorticella

The transition into telotrochs via binary fission was video-recorded at higher magnification

(Figure 1J-P and Figure 1-video 8). The zooid, equipped with a prominent wreath of cilia at its

oral end (oral cilia) used for feeding (Figure 1J), becomes more rounded (Figure 1K), initiates

division (Figure 1L), and forms two daughter cells (Figure 1M). The oral cilia retract, while the

aboral end becomes encircled by a wreath of cilia (aboral cilia) that emerge from aborally

positioned kinetosomes (Buhse, 1998) (Figure 1M). The cell elongates, transforms into a

telotroch, detaches from the stalk, and swims with the aboral end leading, propelled by the

aboral cilia (Figure 1M-P). After swarming, the telotroch settles on a solid surface, secretes a

stalk, and reverts into a zooid (Figure 1Q-S and Figure 1-video 9). Not all cells undergo division

prior to transformation. In certain samples, the majority undergo transit without cell division

(Figure 1-video 10), whereas others exhibit a combination of both types [Figure 1-video 11 and

Figure 1T, some individuals transform through cell division (..) while others proceed without it

(.)]. The genotypes of both populations are identical. As described later, this transformation

appears to occur daily, with cell division taking place on the first day but not on the following

day in the observed case, indicating that the presence or absence of division is driven by

physiological conditions, such as nutrition, rather than genetic inheritance. Despite the

presence of both modes within the population, the timing of the transformation post-division is

not significantly delayed compared to that of the population without division (Figure 1-video

11), suggesting a highly regulated transformation mechanism across individuals.

It is interesting to explore how and when telotrochs decide on a site for settlement. A

few individuals appear to select the settlement location first, with the remaining ones following

suit and positioning themselves adjacent to one another (Figure 1U and Figure 1-video 12,

additional examples Figure 1-video 13-15). The decision-making process is not straightforward;

they continue roaming, searching for a site, and may even communicate with each other via

physical contact. In another instance, when numerous telotrochs were involved, two focal

points of settlement were established in rapid succession, and the remaining individuals

arranged themselves between the two foci as if attempting to avoid gaps (Figure 1V-X and

Figure 1-video 16). An additional example underscores another facet of the behavior: later-

arriving individuals persist in seeking proximity to those that preceded them, even when a

substantial number have already commenced their transformation into zooids (Figure 1-video

17). This suggests a strong preference for settling in as densely packed a manner as possible.

We conducted prolonged observations exceeding 60 hours and determined that the

transformation cycle from zooid to telotroch occurs at intervals of approximately 21 hours. The

telotrochs congregated at location 1 (Figure 2 and Figure 2-video 1) and swarmed toward

location 2, and so forth. The intervals between the initiation of each swarming event were

approximately 20 hours (from swarming away from 1 to swarming away from 2), 20.5 hours

(from 2 to 3), and 23 hours (from 3 to 4). Hence, the average interval between successive

swarms was roughly 21 hours. This observation aligns with the previous report indicating an

interval of around 24 hours (Fox and Newth, 1936).

Mucous fiber formation

The existence of mucus is discernible upon the mechanical displacement of the swarm (Figure

3A-B and Figure 3-video 1), as articulated by Fox and Newth, who noted that 'the presence of

mucus would not be suspected but for the fact that the mass, which has a gelatinous

consistency, can be pulled aside bodily with a needle'(Fox and Newth, 1936). Under dark-field

illumination (Figure 3C-K), an indistinct thread—presumably composed of mucous fibers, as

inferred from their findings—emerges shortly after the swarm commences (Figure 3E). The

thread becomes increasingly defined, thickens, and ultimately condenses into a globular

structure (Figure 3F-J), consistent with their observation that 'the movements of the animals

themselves may cause the progressive solidification of the mucus which occurs during

swarming.' (Fox and Newth, 1936). It follows the swarm until settlement on the substrate but

does not necessarily participate in the settled population, as observed in this particular

instance. The condensed mucous fiber was attached above the plant fragment, while the

telotrochs settled beneath, out of focus (Figure 3K). We lacked the opportunity to stain the

mucous fiber for closer examination and thus cannot describe its characteristics in detail,

leaving it unclear whether the telotrochs retain the mucous fibers post-condensation at the end

of swarming.

What drives their propensity to swarm and congregate in groups upon settlement?

What is benefit of this behavior? In unicellular organisms, this phenomenon can be interpreted

from an evolutionary standpoint as the origin of multicellularity. A notable example is the slime

mold Dictyostelium discoideum, which proliferates as free-living unicellular amoebae through

cell division and, upon starvation, aggregates to construct a multicellular fruiting body with

differentiated cysts or spores. Organisms displaying such life cycles have independently

evolved at least six times across both eukaryotic and prokaryotic lineages, including fungi and

ciliates (Du et al., 2015). However, in the case of *Vorticella*, they remain independent,

exhibiting swarming and spatial proximity without coalescing into a multicellular structure akin

to a "slime mold." One hypothesis is that their close assemblage may facilitate sexual

conjugation, though we have yet to observe such events as previously noted (Fox and Newth,

1936). Recent investigations into *Stentor coeruleus*, a suspension-feeding unicellular ciliate like

Vorticella, which exists both solitarily and colonially, suggest that hydrodynamic interactions

between neighboring individuals enhance feeding efficiency through accelerated flow,

contingent on inter-individual spacing (Shekhar et al., 2023). This could potentially account for

the pronounced clustering observed in these ciliates. However, if this hypothesis is valid, then

why does *Vorticella* repeatedly undergo cycles of aggregation and dispersion daily, despite the

significant energy expenditure required for the transition between zooid and telotroch forms? A

potential explanation for this behavior may lie in their predator evasion strategy, as outlined

below.

We sporadically observed the predatory ciliate, *Amphileptus sp*., feeding on *Vorticella*

(Figure 4 A-D and Figure 4-video 1). Individuals within a group can be consumed sequentially if

they remain in their stalked sessile form (Figure 4 E-I and Figure4-video 2). The periodic

transition to the telotroch stage and relocation may serve as a strategy to evade predators,

while swarming enables them to regroup for more efficient feeding.

The case of *V. campanula*

We also identified a distinct species of *Vorticella*, tentatively designated as *V. cf. campanula*

(Sanshirou), due to its morphological resemblance to *V. campanula* (notably, the shape of the

cell body and the presence of refractive oil droplets within the cytoplasm, which give the cells a

darkened appearance) (Figure 4), as well as its genomic sequence. The ITS1-5.8S-ITS2

region of *V. cf. campanula* (Sanshirou) exhibits the highest sequence homology to that of *V.*

campanula, despite some degree of heterogeneity observed among individual specimens

(Figure 4-supplementary figure1). The behaviors of *V. cf. campanula* (Sanshirou) have not

been extensively studied in comparison to *V. sp11* (Sanshirou), yet they exhibit significant

similarities with previous research on *V. campanula* (Fox and Newth, 1936; Fox and NEWTH,

1935). *V. cf. campanula* (Sanshirou) transitions from the zooid to telotroch stage (Figure 5A-D)

and settle in groups (Figure 5E-G), much like *V. sp11* (Sanshirou). The most pronounced

difference lies in their swarming behavior. *V. sp11* (Sanshirou) forms a compact, ball-like

cluster resembling a cloud of bees, whereas *V. cf. campanula* (Sanshirou) moves back and

forth along mucous threads (Figure 5H), as shown by the line from the upper left corner to the

middle bottom, along which individuals slide (Figure 5H). In more natural conditions, these

threads can extend up to 10 cm in length (Fox and Newth, 1936; Fox and NEWTH, 1935). The

swarming behaviors of the two species were compared (Figure 5I-J). *Vorticella sp11*

(Sanshirou) exhibited a shorter swarming duration, forming a compact and dense cloud and

ultimately settling into a tightly clustered group (Figure 5I) as described. In contrast, *Vorticella*

cf. campanula (Sanshirou) moved along a trajectory for an extended period before settling in

smaller, more dispersed groups (Figure 5J). When *V. cf. campanula* (Sanshirou) transitioned to

telotrochs in a larger container, swarming units initially appeared locally, gradually merging and

morphing over time, and eventually coalescing into a unified, larger stream (Figure 5K-N). This

example illustrates the swarming behavior across a more extensive area, paralleling previous

observations (Fox and Newth, 1936; Fox and NEWTH, 1935), despite the inherent differences

between our two-dimensional experimental conditions and their more natural, three-

dimensional environment, which was described as follows: 'Numerous individuals swim close

to the mucous thread-some moving round it, some swimming upwards, others downwards-

while all the time individuals leave the thread and return to it after short horizontal

excursions.'(Fox and Newth, 1936)

The analogous swarming behaviors of the two types of *Vorticella* have been

documented, nearly a century apart, in Great Britain (Fox and Newth, 1936; Fox and NEWTH,

1935) and Japan (this study). Certain *Vorticella* species do not aggregate but instead settle

solitarily on surfaces. Horikami and Ishii, in their conference abstract, reported that *V.*

campanula, *V. convallaria*, *V. monilata*, and *V. vestita* exhibited group settlement behaviors.

Additionally, they described that a viscous substance could attract telotrochs in a species-

specific manner (Horikami and Ishii, 1981). The role of mucus in swarm formation remains

uncertain. Does it merely encapsulate the telotrochs to restrict their mobility, or does it possess

a physiological function that modulates telotroch behavior, as indicated by Horikami and Ishii?

Is the strand observed by Fox and Newth derived from the mucus, contributing to the formation

of the putative condensed mucus-like material we detected? Regrettably, we lack evidence to

address this question, as we only discovered the relevant publication on mucus fibers after

completing our experiments. We are also intrigued by the duration for which the mucus exerts

its influence during the settlement process. In certain instances (e.g., Figure 3-video 2), the

condensed mucus-like material does not participate in the settlement cluster. Conversely, upon

closer examination of the settlement process (e.g., Figure 1-video 12), a few telotrochs initially

rotate around the condensed mucus-like material, appearing to be constrained by a central

tension, suggesting a persistent physical linkage. Elucidating the molecular function of this

mucus, along with any potential swarming behaviors exhibited by other *Vorticella* species,

would present an intriguing direction for future research.

If the dense clustering observed in this *Vorticella* settlement is not considered the

precursor to multicellularity, then it may represent an evolutionary origin of coordinated group

actions in multicellular organisms. Eduardo Orias posited that ciliates could be perceived as

multicellular organisms. He proposed that ciliates with multiple cilia evolved from flagellates

bearing a single flagellum through a mutation that delayed cytokinesis, resulting in chains of

flagellates (Orias, 1976). In this context, ciliates can already be viewed as multicellular

organisms, and the *Vorticella* species described here could be characterized as "gregarious

species", as previously noted (Fox and Newth, 1936).

Materials and Methods

Identification of the *Vorticella* species

Genomic DNA was isolated from 5-10 zooids and the ITS1–5.8S–ITS2 region was amplified by

polymerase chain reaction (PCR) using the SB2 (5′-

GTAGGTGAACCTGCGGAAGGATCATTA-3′) primer (Goggin and Murphy, 2000) and ITS-R

(5′-TACTGATATGCTTAAGTTCAGCGG-3′) primer (Sun et al., 2010). The thermal cycling

conditions were as follows: initial denaturation at 94°C for 2 minutes, followed by 35 cycles of

denaturation at 98°C for 10 seconds, annealing at 60°C for 30 seconds, and extension at 72°C

for 60 seconds, with a final extension step at 72°C for 7 minutes. The DNA database was

queried for sequence homology utilizing the Basic Local Alignment Search Tool (BLAST)

available at https://blast.ncbi.nlm.nih.gov/Blast.cgi. The DNA sequences were aligned utilizing

the ApE Plasmid Editor v3.1.4 (Wayne Davis) and edited using Illustrator (Adobe).

Microscopy and video recordings

The organisms were examined using the Axio Observer, Axiovert, and Axio Zoom V.16

microscopes (Carl Zeiss). Video recordings were captured with the URSA Mini Pro 12K

[Blackmagic Design (BMD)], Blackmagic Pocket Cinema Camera 6K (BMD), ILCE-1 (Sony),

and S1H (Panasonic) cameras, and subsequently edited using DaVinci Resolve 19 (BMD).

The manuscript underwent partial grammatical refinement using ChatGPT.

Acknowledgements

We thank Toshinobu Suzaki for providing a conference abstract and Ryu Ueda for helpful

comments. We are grateful to Minoru Saitoe and Takashi-Adachi Yamada for their generous

supports. This work was supported by grants from the Ministry of Education, Culture, Sports,

Science and Technology (21K19288) to T.T.

Author contributions: T.T. designed the research, performed imaging and wrote the manuscript.

Y.M. performed the PCR analysis.

Competing interest

The authors declare that no competing interests exist.

Additional files

Supplementary files

Figure 1-supplementary figure1

Figure 4-supplementary figure1

References

Allee WC. 1927. Animal Aggregations. Q Rev Biol 2:367–398. doi:10.1086/394281

Barlow BM, Finley, HE. 1976. Comparative Studies on Four Species of Vorticella by

Conventional Microscopy. Trans. Amer. Micros. Soc., 95: 346–351.

doi.org/10.2307/3225125.

Buhse HE. 1998. Vorticella: "A Cell For All Seasons"1. J Eukaryot Microbiol 45:469–474.

doi:10.1111/j.1550-7408.1998.tb05102.x

Buonanno F, Ortenzi C. 2021. Extremophilic Microbes and Metabolites - Diversity,

Bioprospecting and Biotechnological Applications. doi:10.5772/intechopen.78622

Du Q, Kawabe Y, Schilde C, Chen Z, Schaap P. 2015. The Evolution of Aggregative

Multicellularity and Cell–Cell Communication in the Dictyostelia. J Mol Biol 427:3722–3733.

doi:10.1016/j.jmb.2015.08.008

Fox HM, Newth HG. 1936. On the Swarming of Vorticella. Proc Zoöl Soc Lond 106:309–313.

doi:10.1111/j.1096-3642.1936.tb02291.x

Fox HM, NEWTH HG. 1935. Swarming of Vorticella. Nature 136:435–435.

doi:10.1038/136435b0

Goggin C, Murphy N. 2000. Conservation of sequence in the internal transcribed spacers and

5.8S ribosomal RNA among geographically separated isolates of parasitic scuticociliates

(Ciliophora, Orchitophryidae). *Dis Aquat Org* **40**:79–83. doi:10.3354/dao040079

Horikami H, Ishii K. 1981. Cluster inducing factor secreted by Vorticella telotrochProgress in

Protozoology. Presented at the the 6th International Congress on Protozoololgy. Warsaw,

Porland. p. 145.

Shekhar S, Guo H, Colin SP, Marshall W, Kanso E, Costello JH. 2023. Cooperative

hydrodynamics accompany multicellular-like colonial organization in the unicellular ciliate

Stentor. bioRxiv 2023.01.10.523506. doi:10.1101/2023.01.10.523506

Sun P, Clamp JC, Xu D. 2010. Analysis of the secondary structure of ITS transcripts in peritrich

ciliates (Ciliophora, Oligohymenophorea): Implications for structural evolution and

phylogenetic reconstruction. Mol Phylogenetics Evol 56:242–251.

doi:10.1016/j.ympev.2010.02.030

Sun P, Clamp JC, Xu D, Huang B, Shin MK, Turner F. 2013. An ITS-based phylogenetic

framework for the genus Vorticella: finding the molecular and morphological gaps in a

taxonomically difficult group. Proceedings Biological sciences / The Royal Society

280:20131177. doi:10.1098/rspb.2013.1177

Figure 1. Life cycle of *V. sp11* (Sanshirou). (A) Zooids affixed to a *Cabomba sp.* leaf. (B-I)

The transition from zooids (B) to telotrochs (C) and their subsequent group settlement (I).

Arrows denote the condensed mucus (F, H). (J-P) Cellular division and the ensuing transition

from zooids (J) to telotrochs (O). (J) *oc* denotes oral cilia, while (N) *ac* denotes aboral cilia. (Q-

S) Telotroch settlement and subsequent transition to a zooid form. (T) Telotroch formation via

cellular division (..) and in its absence (.). The video content is identical to that in (A) but

presented in a vertically inverted orientation. (U) The settlement process is shown in high-

resolution video recordings. (V-X) An instance of the settlement process involving a substantial

number of telotrochs. Scale bars represent 100 μm in all instances, except for (B-I), where the

scale bar denotes 1 mm. All figures represent frames from video recordings, with online access

to corresponding video images provided below and hereafter.

Figure 1-video 1 (related to Figure1 B-I) The transition from zooids to telotrochs and their

subsequent group settlement. The playback rate is accelerated by a factor of 100 in select

segments. https://youtu.be/iCczJqXLKbA

Figure 1-video 2 (related to Figure1 B-I) The transition from zooids to telotrochs and their

subsequent group settlement. Accelerated by a factor of 100. https://youtu.be/ezEXs3lvFgo

Figure 1-video 3 (related to Figure1 B-I) The transition from zooids to telotrochs and their

subsequent group settlement. Accelerated by a factor of 100. https://youtu.be/eQwjoO6vllM

Figure 1-video 4 (related to Figure1 B-I) The transition from zooids to telotrochs and their

subsequent group settlement. Select segments are accelerated 100-fold.

https://youtu.be/EJKMewtThXo

Figure 1-video 5 (related to Figure1 B-I) The transition from zooids to telotrochs and their

subsequent group settlement. Select segments are accelerated 100-fold.

https://youtu.be/LKlsDLsH1g0

Figure 1-video 6 (related to Figure1 B-I) The transition from zooids to telotrochs and their

subsequent group settlement. Select segments are accelerated 100-fold.

https://youtu.be/bAN8v2Xz-WM

Figure 1-video 7 (related to Figure1 B-I) The transition from zooids to telotrochs and their

subsequent group settlement without adherence to the substrate. Select segments are

accelerated 100-fold. https://youtu.be/zO9UL59LDG8

Figure 1-video 8 (related to Figure1 J-P) The transition into telotrochs via binary fission.

Select segments are accelerated 100-fold. https://youtu.be/az9qcpWltYk

Figure 1-video 9 (related to Figure1 Q-S) After swarming, the telotroch settles on a solid

surface, secretes a stalk, and reverts into a zooid. Select segments are accelerated 8-fold.

https://youtu.be/-C86QUN5fyA

Figure 1-video 10 (related to Figure1) The majority of the zooids undergo transit without cell

division. Select segments are accelerated 100-fold. https://youtu.be/hb9d2x7elTQ

Figure 1-video 11 (related to Figure1 T) Some individuals transform through cell division

while others proceed without it. Select segments are accelerated 100- or 8-fold.

https://youtu.be/fD6hb5axIn0

Figure 1-video 12 (related to Figure1 U) A few individuals appear to select the settlement

location first, with the remaining ones following suit and positioning themselves adjacent to one

another. Select segments are accelerated 20.03-fold. https://youtu.be/JetWGHHWKYc

Figure 1-video 13 (related to Figure1 U) An example of the settlement process. Select

segments are accelerated 20.1-fold. https://youtu.be/8dOEmC7t2Zk

Figure 1-video 14 (related to Figure1 U) An example of the settlement process. Select

segments are accelerated 100.06-fold. https://youtu.be/XDPElb5KTqw

Figure 1-video 15 (related to Figure1U) A representative instance of the settlement process

involving a substantial number of telotrochs. Select segments are accelerated 100-fold.

https://youtu.be/z-5ND7Cm4JA

Figure 1-video 16 (related to Figure1 V-X) Two focal points of settlement are established in

rapid succession, and the remaining individuals arrange themselves between the two foci as if

attempting to avoid gaps. Select segments are accelerated 10-fold.

https://youtu.be/1EWbjksuKsQ

Figure 1-video 17 (related to Figure1) Individuals that arrive later still attempt to settle near

the earlier ones, even if those have already begun transforming into zooids. The inset presents

an expanded rendition of the upper-right corner. Select segments are accelerated 100- or 7.86-

fold. REC/SRC TC: 00:00:00:00/00:00:00:00 indicates the time line time code and source time

code in the format hours:minutes:seconds:frames, here and hereafter, unless otherwise

explicitly noted. https://youtu.be/knojt56FYc4

Figure 2. The swarming behaviors documented through video recordings spanning 60

hours. The settlement location transitioned sequentially from positions 1 to 5, with each shift

occurring at an interval of approximately 21 hours. Scale indicates 1 mm.

Figure 2-video 1 (related to Figure2) Behaviors were recorded at 30 frames per second for

the initial 2 hours and at 1 frame per second for the remaining duration. The recording length is

approximate, as SSD storage swaps were conducted for extended recording periods. All

source files were concatenated into a single compound file for editing purposes. The source

time code format 01:01:01:01 represents 30 hours:0.5 hours:0.5 minutes:1 second, with each

30-second increment advancing the format by one unit. Select segments are accelerated 120-,

240- or 3000-fold. https://youtu.be/1BrYJ43Uo88

Figure 3. Telotrochs secrete mucus. (A-B) A spherical telotroch swarm being drawn aside

bodily with a needle. (C-K) The progression of mucous fiber development accompanying

telotroch swarming. An arrow denotes the faint, early-stage mucous fiber (E), while an arrow

indicates the condensed mucous fiber positioned apart from the settled telotrochs (K). Scales

indicate 1 mm.

Figure 3-video 1 (related to Figure3A-B) Select segments are accelerated 100.06-fold.

https://youtu.be/TTx5ickofxs

Figure 3-video 2 (related to Figure3C-K) Accelerated by a factor of 100.

https://youtu.be/iLhBgE3D0kk

Figure 4. A predatory ciliate, *Amphileptus sp.*, preying upon Vorticella. (A-E) The intricate

process by which *Amphileptus sp.* preys upon *Vorticella sp11 (Sanshirou)* was observed. Upon

mechanical contact with *Vorticella* using its proboscis (B, arrow), *Amphileptus* appears to

discharge prexicysts or analogous organelles (C, arrow), akin to the behavior exhibited by

Didinium (Buonanno and Ortenzi, 2021) facilitating the capture of *Vorticella*. (F-I) A cluster of

Vorticella sp11 (Sanshirou) was sequentially consumed by a group of *Amphileptus sp* (an

asterisk).

Figure 4-video 1 (related to Figure 4A-E) Select segments are accelerated 8-fold.

https://youtu.be/nzoriPLGvcI

Figure 4-video 2 (related to Figure 4A-D) The instance of Amphileptus sp. preying upon

Vorticella is indicated by an asterisk. Select segments are accelerated 100-, 20.1-, 19.97- or 8-

fold. https://youtu.be/vZD5u9RqMDg

Figure 5. Swarming behaviors of *V. cf. campanula (Sanshirou).* (A-D) Transition from

zooids to telotrochs. (A) *oc* denotes oral cilia, while (B) *ac* indicates aboral cilia. (E-G)

Settlement of telotrochs. (H) Telotroch swarming. (I-J) Side-by-side comparison of swarming

behaviors between *V. sp11 (Sanshirou)* (The upper-left specimen is previously exhibited in

Figure 3C-K) (I) and *V. cf. campanula (Sanshirou)* (J). Scales indicate 1 mm. (K-N) Telotroch

swarming coalescing and forming a single, unified stream. Scales are not available.

Figure 5-video 1 (related to Figure 5A-D) Select segments are accelerated 50.18- or 50.01-

fold. https://youtu.be/DKpSIFoM1YU

Figure 5-video 2 (related to Figure 5E-G) Select segments are accelerated 20.09-fold.

https://youtu.be/9zkQYLnVqk8

Figure 5-video 3 (related to Figure 5H) https://youtu.be/9gwk3RgWn4M

Figure 5-video 4 (related to Figure 5I-J) The two video recordings (I and J) were conducted

independently and integrated during the editing process. Accelerated 20.09-fold.

https://youtu.be/0wlM_yDZvzY

Figure 5-video 5 (related to Figure 5K-N) The recording field expanded progressively as the

swarm's area increased. Select segments are accelerated 100.11-fold.

https://youtu.be/ZkqdtnE0GDA