Comparative genomic analysis of a metagenome-assembled genome reveals distinctive

symbiotic traits in a Mucoromycotina fine root endophyte arbuscular mycorrhizal

fungus

Joshua Cole[1*], Sébastien Raguideau[2], Payman Abbaszadeh-Dahaji[3,4], Sally Hilton[1], George

Muscatt[1], Ryan M. Mushinski[1], R. Henrik Nilsson[5], Megan H. Ryan[3], Christopher Quince[2,6,7]

and Gary D. Bending[1*]

[1]School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK

[2]Earlham Institute, Norwich Research Park, Colney Lane, Norwich NR4 7UZ, UK

[3]UWA School of Agriculture and Environment, University of Western Australia, Crawley,

WA, 6009, Australia

[4]Department of Soil Sciences, Faculty of Agricultural Science, Vali-e-Asr University of

Rafsanjan, Rafsanjan, Iran

[5]University of Gothenburg, Department of Biological and Environmental Sciences,

Gothenburg Global Biodiversity Centre, University of Gothenburg, Box 461, 405 30,

Göteborg, Sweden

[6]Gut Microbes and Health, Quadram Institute, Norwich NR4 7UQ, UK

[7]School of Biological Sciences, University of East Anglia, Norwich NR4 7T, UK

*Corresponding authors: Joshua.M.Cole@warwick.ac.uk; gary.bending@warwick.ac.uk

ORCID

Joshua Cole - 0009-0004-1563-8469

Sébastien Raguideau - 0000-0001-8727-2604

Payman Abbaszadeh-Dahaji - 0000-0003-3055-4744

Sally Hilton - 0000-0002-3786-1096

George Muscatt - 0000-0002-9218-1210

Ryan M Mushinski - 0000-0003-3572-3500

R. Henrik Nilsson – 0000-0002-8052-0107

Megan H. Ryan – 0000-0003-0749-0199

Christopher Quince – 0000-0003-1884-8440

Gary D. Bending - 0000-0002-9740-5244

Summary

• Recent evidence shows that arbuscular mycorrhizal (AM) symbiosis is established by
two distinct fungal groups, with the distinctive 'fine root endophyte' morphotype
formed by fungi from the sub-phylum Mucoromycotina rather than the sub-phylum
Glomeromycotina. While Mucoromycotina AM fungi are globally distributed, their
genomic basis for the symbiosis or how this compares to other mycorrhizal symbionts
is unknown.

• We used culture-independent metagenome sequencing to assemble and characterise
the metagenome-assembled genome (MAG) of a putative fine root endophyte, which
we show belonged to the family Planticonsortiaceae within the order Densosporales.

• The MAG shares key traits with Glomeromycotina fungi, which indicate obligate
biotrophy, including absence of fatty acid and thiamine biosynthesis pathways,
limited enzymatic abilities to degrade plant cell walls, and a high abundance of
calcium transporters. In contrast to Glomeromycotina fungi, it exhibited a higher
capacity for degradation of microbial cell walls, a complete cellulose degradation
pathway, low abundances of copper, nitrate and ammonium transporters, and a
complete pathway for vitamin B6 biosynthesis.

• These differences highlight the potential for contrasting interactions between
Mucoromycotina and Glomeromycotina AM fungi with their host plant and the
environment which could support niche differentiation and complementary ecological
functions.

60
61

Introduction

Mutualistic symbioses with microbes are exhibited by the overwhelming majority of plant species (Begum, 2019) with mycorrhizas by far the most widespread and well-described association (Hawkins et al., 2023). While mycorrhizal symbioses have evolved multiple times across plant and fungal taxa, they share the common characteristic of the fungus trading nutrients assimilated from the soil, particularly phosphorus and nitrogen, in exchange for carbon from the plant (Luginbuehl & Oldroyd, 2017). Additionally, mycorrhizal associations can confer a range of secondary benefits to the plant host, including disease resistance and stress tolerance (Chen, 2018). Because of these profound impacts on plants, mycorrhizal associations are considered to be major determinants of ecosystem productivity and diversity and to play key roles in directing terrestrial biogeochemical cycling processes. There are four main groups of mycorrhizal associations: arbuscular (AM), ecto- (ECM), ericoid (ERM), and orchid (ORM) (Martin & van der Heijden, 2024) mycorrhizas, which are distinguished by a number of features including the specific plant and fungal taxa involved, the ecosystems they inhabit, the way in which the fungus colonises and grows within root tissues, and the functional traits of the fungi involved (Brundrett & Tedersoo, 2018).

As a function of time and niche specificity, mycorrhizal symbionts appear to have evolved along a saprotrophy to symbiosis continuum (Strullu-Derrien et al., 2018; Martin & van der Heijden, 2024). This directional transition from free-living saprotrophs to obligate symbionts is believed to be driven by the provision of carbohydrates and vitamins from plants in nutrient-poor environments and the subsequent evolutionary success of a symbiotic lifestyle. As a result, key genomic signatures associated with this transition can be identified across this continuum (Martin & van der Heijden, 2024). As fungal symbionts shift away from primarily saprotrophic lifestyles to mutualisms, their genomes typically possess a reduced repertoire of the enzymatic machinery involved in the degradation of soil organic matter (SOM) and plant cell walls (Martino et al., 2018; Gong et al., 2023; Martin & van der Heijden, 2024). The two youngest mycorrhizal symbioses, ERM and ORM, with origins c 75-90 and 112 mya, respectively, (Martino et al., 2018; Brundrett & Tedersoo, 2018) involve symbionts with genomic repertoires capable of simultaneously facilitating a saprotrophic and a symbiotic lifestyle. However, fungi forming the older AM (origin 450 mya, Brundrett & Tedersoo, 2018) and ECM symbioses (origin c 200 mya) possess a much more fine-tuned enzymatic toolkit, noted for a reduced suite of plant cell wall degrading enzymes (PCWDE)

along with the presence of specific genes involved in regulating nutrient transport and host
plant signalling and metabolism (Martino et al., 2018; Martin & van der Heijden, 2024).

AM are considered the most common and widespread plant-microbe symbiosis on Earth
(Read et al., 2000; Parniske, 2008; Spatafora et al., 2017), and their presence in fossil roots of
the earliest land plants suggests that they were a significant factor in the evolution of land
plants. Fungi forming AM have traditionally been placed into the Mucoromycota sub-phylum
Glomeromycotina. These fungi are obligate symbionts; as fatty acid auxotrophs they depend
on the exchange of plant derived fatty acids (Schüßler et al., 2007; Luginbuehl et al., 2017).
As a result of long-term host dependence, arbuscular mycorrhizal fungi (AMF) present
hallmark traits such as limited ability to break down cellulose, biosynthesise thiamine, or
metabolise vitamin B6 derivatives. The genomic inability to express these hallmark traits
represent Missing Glomeromycotan Core Genes (MGCGs) (Tang et al., 2016; Kobayashi et
al., 2018; Rosling et al., 2024).

Until recently, all AMF were believed to belong to the Glomeromycotina (G-AMF)
(Spatafora et al., 2017). It has long been recognised that there are two broad groups of
arbuscular mycorrhiza based on morphology, with the 'coarse endophyte' G-AMF being the
focus of the vast majority of research on the biology and ecology of AMF (Orchard et al.,
2017b). Glomeromycotina comprises four orders, the Glomerales, Diversisporales,
Paraglomerales, and Archaeosporales, which likely diversified when plants evolved onto land
(Schüßle 2001; Chen et al., 2018; Tedersoo et al., 2018). The second morphological group of
'fine root endophytes (FRE)' arbuscular fungi are distinguished from coarse endophytes by
their small spores, fan-shaped or palmate entry points, diverging thin hyphae (<2 µm
diameter) which in intercellular spaces often form hyphal ropes, as well as intercalary and
terminal 'vesicles'', approximately 5–10 µm in diameter (Thippayarugs et al. 1999; Orchard et
al., 2017a, b; Walker et al. 2018). Interestingly, the arbuscules of FRE appear very similar in
structure to those of coarse endophytes (Albornoz et al. 2020). Recent evidence has placed
FRE within the sub-phylum Mucoromycotina (M-AMF) (Orchard et al., 2017b), indicating
that AM symbioses may have evolved independently several times, as is the case for
ectomycorrhizal symbioses (Miyauchi et al., 2020).

To date only one species of M-AMF, *Planticonsortium tenue*, has been described (Walker et
al., 2018). However, similarly to G-AMF, root systems are colonised by diverse
Mucoromycotina taxa, communities of which are typically comprised of a small number of
abundant and common taxa and a high diversity of rare low-abundance taxa (Albornoz et al.,

2021, 2022; Seeliger et al., 2024). Furthermore, distinctive morphologies of FRE have been
recognised (Thippayarugs et al., 1999) based on a variety of characteristics, including the
roughness, diameter, and branching of hyphae and the size and shape of vesicles. While fungi
within the Glomeromycotina are all assumed to be plant symbionts, fungi residing within the
Mucoromycotina also include saprotrophs and putative ectomycorrhizal symbionts. Those
Mucoromycotina sequences found associated with roots and rhizoids of vascular and non-
vascular plants appear to have similar phylogenetic ranges and occur within clades alongside
taxa with saprotrophic and putative ectomycorrhizal lifestyles (Desirò et al., 2017; Seeliger et
al., 2024).

M-AMF have been detected across multiple continents, spanning a wide range of ecosystems,
environmental conditions, and hosts, although in contrast to G-AMF they may be missing
from tropical and sub-tropical biomes, and may be favoured by agricultural management
practices (Orchard et al., 2017a; Albornoz et al., 2022). While the benefits of G-AMF to the
host, particularly for phosphorus nutrition, are well documented, the functional significance
of M-AMF within ecosystems is poorly understood (Orchard et al., 2017b; Hoysted et al.,
2019; Albornoz et al., 2020). G-AMF and M-AMF typically co-occur and intermingle within
root systems (Orchard et al., 2017a,b; Jeffery et al., 2018), but they appear to occupy distinct
but overlapping ecological niches (e.g. Orchard et al. 2016), with their distribution at the
landscape scale differentially determined by temperature, pH, and plant richness (Albornoz et
al., 2022). Unlike G-AMF, nutrient availability was found to be an important determinant of
M-AMF abundance, which increased with fertility (Albornoz et al., 2022). Furthermore, in
contrast to G-AMF, experimental evidence suggests that M-AMF may provide the host with
access to soil nitrogen, indicating a potential complementary functionality between the two
arbuscular symbioses and an explanation for the parallel evolution of symbiotic traits
(Hoysted et al., 2022; Howard et al., 2024).

Comparative genome analysis has emerged as a powerful tool to understand microbial
evolution, function, and ecology. However, these approaches largely rely on the availability
of cultured microbes or voucher specimens for whole genome sequencing (e.g. Martino et al.,
2017; Miyauchi et al., 2020; Rosling et al., 2024). Importantly, culturing microbes can prove
time-consuming and problematic as microbial populations have great complexity and many
microbes have specific nutritional requirements which cannot be met in culture. This is
particularly the case for obligate biotrophs such as symbionts which rely wholly or in-part on
the host for their nutritional requirements (Tanaka et al., 2022). As obligate symbionts, G-

AMF require a host plant to complete their lifecycle. Despite this, pure cultures of G-AMF

have been obtained using root organ cultures, and these have been used to produce inoculum

for genome sequencing (Tisserant et al., 2013). The challenges in obtaining and maintaining

root cultures have meant that direct genome sequencing of G-AMF cultures has been limited

to date. Recently, asymbiotic mycelium and spore production of the G-AMF *Rhizophagus*

clarus has been achieved by supplementing its growth medium with palmitoleic acid, and this

could provide a boost to rates of culture-based sequencing (Tanaka et al., 2022). An

alternative approach to obtain fungal genomes is sequencing of nuclei extracted from spores

(Montoliu-Nerin et al., 2021), although most G-AMF taxa appear to sporulate infrequently, if

at all (Öpik et al., 2014).

M-AMF are also putatively obligate symbionts, but because they have only recently been

recognised as a distinct group of symbionts, culturing efforts lag behind those of G-AMF.

Recently, monoxenic cultures of a Mucoromycotina fungus associated with the clubmoss

Lycopodiella inundata have been obtained and shown to form FRE colonisation in a vascular

plant (Hoystead et al., 2022). However, while genome sequencing and comparative genomic

analysis of a number of saprotrophic and putative ectomycorrhizal Mucoromycotina has been

achieved (Chang et al., 2019; Rosling et al., 2024), there is no understanding of the genome

of M-AMF fungi and how their functional repertoire and mutualistic signatures compare to

those of G-AMF, other mycorrhizal symbioses, and saprotrophic and putatively

ectomycorrhizal Mucoromycotina. To further complicate the matter, initial comparative

analysis of the sister clades indicate that some widely considered hallmark traits of

mycorrhizal symbioses may instead represent ancestral markers as opposed to adaptation

(Rosling et al., 2024).

Recently there has been great progress in the development of computational tools for the

culture-independent assembly of 'dark' microbial genomes from metagenome sequencing

datasets (Benoit et al., 2024). Such 'Metagenome Assembled Genomes' (MAGs) have been

widely used to identify the metabolic capabilities and ecological roles of specific prokaryotes

in diverse habitats (Yang et al., 2021). MAG construction of eukaryotic microbes is more

challenging because of a range of issues including their large sizes, high genome

heterozygosity, low abundance in metagenome datasets, and a lack of well annotated

reference genomes for gene annotation (Peng et al., 2021; Setubal 2021). As a result, few

studies have constructed eukaryotic MAGs from metagenomes, and these have typically been

performed in low-diversity communities (e.g. West et al. 2018; Peng et al., 2021). A further

particular issue with analysis of plant symbionts *in situ* is their intimate association with plant

tissue. This results in co-extraction of host DNA, increasing the sequencing depth required to

assemble symbiont genomes.

The aim of the current work was to use culture-independent metagenome sequencing to

assemble and characterise the genome of an FRE-forming M-AMF fungus for the first time

and compare its genomic repertoire to glomeromycotan fungi and other fungal symbionts.

Roots of *Trifolium subterraneum* were first enriched with FRE using a selective soil sieving

technique. Root samples were used for metagenome sequencing, from which a MAG of the

FRE was constructed. We compiled a database of 274 fungal genomes across the

Ascomycota, Basidiomycota and Mucoromycota, and assigned these to G-AMF, ECM, ERM,

OCM, saprotroph, pathogen, parasite, endophyte, lichenic, and mixed lifestyles. Comparative

analysis showed that the MAG shares key distinctive characteristics with G-AMF. This

includes overlaps in some MCGCs, presence of abundant calcium transporters, and an overall

reduced set of plant cell wall degrading enzymes. However, we found a relatively high

abundance of microbial cell wall degrading enzymes and evidence of complete cellulose

degradation capability, as well as low numbers of nitrogen transporters, suggesting a

distinctive niche relative to G-AMF and the other mycorrhizal lifestyles.

Materials and Methods

Soil collection and sample preparation

For the metagenome sequencing, plants with the root AMF community enriched in M-AMF

were produced following the methods of Orchard et al. (2017a). In brief, soil was collected

from the field studied by Orchard et al. (2016) and Orchard et al. (2017a) in the Peel Harvey

region in Western Australia, Australia: site coordinates were 32° 48 826'S, 115° 53346'E and

the site was ~8 m above sea level. The field contained winter-active annual grasses and

legumes and supported a low-intensity free-range beef cattle grazing enterprise. Soil

collection targeted a sandy loam soil with a pH in $CaCl_2$ of 5.4. Soil was wet-sieved through a

stack of four sieves with pore sizes ranging from 2 mm down to 20 µm. The material

collected on each sieve was retained and added to pasteurised, dried, washed coarse river

sand at a ratio of either 1:27 or 1:81 w/w. Free-draining pots were filled with 1 kg of each

treatment (n=4) and placed in a temperature-controlled glasshouse. Four plants of *T.*

subterraneum L. with appropriate rhizobia added were grown in each pot which were

maintained at 80% field capacity by watering to weight, and received phosphorus-free, but

otherwise complete, nutrient solution once a week. After 8 weeks, plants were harvested and

roots gently washed clean. The four root systems from each pot were allocated to one of two

future activities: staining for assessment of AMF colonisation (stored in 70% ethanol) or

DNA extraction (frozen at -20ºC). Roots were cleared and stained using the ink-vinegar

method of Vierheilig et al. (1998) and assessed using a modified version of the magnified

intercept method described by McGonigle et al. (1990) where the root at each grid-line root

intercept was recorded as either free of all colonisation, colonised by G-AMF only, colonised

by M-AMF only, or colonised by both M-AMF and G-AMF.

DNA extraction, 18S rRNA gene amplicon sequencing and metagenome sequencing

Twelve samples were selected for 18S rRNA amplicon and metagenome sequencing on the

basis of predominance and abundance of M-AMF colonisation as well as quality and quantity

of DNA obtained (Table S1). DNA was extracted from root samples using the DNeasy

PowerSoilPro Kit (Qiagen) following the manufacturer's protocol. DNA concentrations were

measured using a Qubit 3.0 fluorimeter using the Qubit dsDNA high sensitivity assay kit

(Invitrogen) according to the manufacturer's protocol.

To determine whether the dilution technique had enriched M-AMF we preformed community

18S rRNA amplicon analysis. Glomeromycotina and Mucoromycotina 18S rRNA genes in

root samples were amplified using AM-Sal-F and AMDGR primers which amplify a 220-

base fragment (Seeliger et al., 2024). PCR reactions, Illumina MiSeq sequencing, and

bioinformatic analysis were performed using the approaches outlined in Seeliger et al. (2024).

Long read nanopore amplicon sequencing of the nuclear ribosomal 18S rRNA, 5.8S, and

28S/LSU region of Mucoromycotina was performed on root samples to support construction

of the FRE MAG and to determine its phylogenetic relationships. PCR reactions were

conducted using AM-Sal-F in combination with RCA95m (Wurzbacher et al., 2018) using

the recommended Oxford Nanopore protocol, amplifying a 4.5 kb region. Briefly, an initial

PCR was performed with adapter-tagged oligos, and the PCR products were cleaned using

Ampure SPRI beads. Barcodes were added using a second PCR with an initial denaturation at

95ºC for 3 minutes followed by 30 cycles of denaturation at 98ºC for 15 seconds, 60ºC

annealing for 15 seconds, and 65ºC extension for 150 seconds, with a final extension at 65ºC

for 6 minutes. Following a clean-up with SPRI beads, the samples were used for sequencing

on a SpotON Nanopore flow cell following the manufacturer's instructions.

Fast5 files were base called and demultiplexed using the Guppy basecaller provided by Oxford Nanopore Technologies. The sample was then processed using amplicon_sorter v2022-03-28 (Vierstraete & Braeckman, 2022) which aims to group amplicons at species or genus level (~95%), output consensus sequences, and quantify the copy number of the representative consensus sequences in the sample. This resulted in 42 consensus sequences, 5 of which were placed into the Mucoromycotina.

To aid detection of M-AMF sequences in the metagenome, a reference 18S rRNA sequence was obtained from DNA extracted from *T. subterraneum* root samples which were enriched in M-AMF and contained minimal G-AMF. These samples were produced in a similar manner to that described above and are presented in Orchard et al. (2017a) as "Experiment 1". On average, 90% of the total AMF colonisation in these roots was contributed by M-AMF, presenting the first evidence linking FRE morphology to Mucoromycotina. The AM-Sal-F and RCA95m primers were used to amplify a 4.5 kb region, as described above, which was cloned using the TA cloning kit (ThermoFisher). Partial 18S rRNA analysis using AM-Sal-F and AMDGR primers revealed a clone, FRE4583, whose sequence showed a 100 % match both to OTU4 which was associated with FRE morphology in the earlier Orchard et al. (2017) study and the BEG249 *Planticonsortium tenue* FRE culture described by Walker et al (2018) (Albornoz et al., 2022). Furthermore, this and closely related sequences have been detected in multiple studies reporting FRE morphology in plant roots (Albornoz et al 2020, Seeliger et al., 2024). Primer walking was used to sequence overlapping regions of clone FRE4583 until the full sequence was covered, generating a 4.5 kb fragment.

Metagenome sequencing and assembly

The twelve root samples were used for Shotgun metagenomic sequencing using the HiSeq 150 paired end protocol to generate between 15 and 47 Gbp sequencing depth. Library construction and sequencing was conducted using Illumina protocols by Novogene UK Ltd.

Samples were first filtered for *T. subterraneum* DNA using bwa mem v0.7.18-r1243-dirty (Li, 2013) and samtools v1.10 (Danecek et al., 2021). To determine FRE depth and breadth of coverage the filtered metagenome reads were mapped, again using bwa mem and samtools, against the FRE4583 ribosomal operon and the resulting alignment files processed with custom Python scripts. To be sure that a high mean coverage was not an artefact of conserved regions, breadth of coverage was visually inspected. Sample P72 was predicted to have the

highest Mucoromycotina abundance, possessing the majority of the operon with a coverage

depth of over 100 (Fig. S1); the missing regions are likely due to repeats as we filtered

ambiguous mappings. We therefore focussed on this sample in our efforts to reconstruct an M-

AMF genome, assembling it using the SPAdes assembler v3.14.1 (Bankevich et al., 2012).

This produced a catalogue of contigs with an N50 of 446 and total length of 1.3 Gbp. We then

computed coverage depth profiles for these contigs across all samples by mapping reads with

bwa mem, samtools, and bedtools v2.30.0 (Quinlan & Hall., 2010). This improves the

discriminatory power of genome binning through multi-sample coverages whilst keeping the

best quality single sample assemblies. The contigs were then binned using these coverage

profiles and composition with the binning software CONCOCT v1.1.0 (Alneberg et al., 2014)

and a minimum contig length of 1000bp.

The putative M-AMF genome was identified based on a consensus of multiple lines of

evidence. As a first step, our reference M-AMF ribosomal sequence FRE4583 was mapped to

the assembly. A total of 12 unitigs were found with high similarity (>98%) and spanning 75%

of the operon without overlap, of which only four had a context in the assembly graph. Using

custom scripts, co-localisation of contigs binned by CONCOCT within the direct vicinity of

the aforementioned 4 untigs were visualised directly on the SPAdes assembly graph (Fig. S2)

using Bandage v0.8.1 (Wick et al., 2015). This figure encompasses all paths of 10 Kbp

starting from these 4 unitigs (red) but correspond to only 0.05% of the whole assembly,

showing a clear proximity of multiple numbered bins. Ignoring any miss-assemblies, all

genomes have a distinct path in the assembly graph and in particular the M-AMF genome

should include a path which passes through those unitigs labelled as ribosomal sequences

allowing us to identify the M-AMF, even though we could not assemble a complete FRE4583

operon in a single contig, likely due to strain diversity.

A total of 12 bins were found in close proximity to M-AMF unitigs. However, on assessing

quality, only bin 17 was found to have any marker genes, giving it a completion of 83.7%

using the BUSCO (Manni et al., 2021) general fungi_odb10 scheme with a size of 37.37

MBp (Table S2). In particular, while contigs from bin 70 were directly co-located with FRE

ribosomal sequences, this bin did not possess any marker genes and was distinctly different in

terms of coverage from bin 17. It may be that bin 70's location in the graph is due to a miss-

assembly, or it could be that this is due to strain diversity and that bin 70 is strain-specific

part of the pan-genome of the FRE which would result in it having a different coverage

profile to the core genome represented by bin 17. To further confirm that bin 17 is related to

M-AMF, we used phyloFlash v3.4.2 (Gruber-Vodicka et al., 2020) to quantify ribosomal

operon abundance from the 12 Illumina short read samples. It is indeed possible to use

phyloFlash to derive abundance, not only of the SSU rRNAs of the SILVA database (Quast et

al., 2013) but also of the 42 nanopore amplicon 18S rRNA consensus sequences. Filtering

any taxa with less than 10 reads mapping across all samples yielded a total of 251 entries

including the 42 consensus sequences (Fig. S3). These were normalised by the total

sequencing depth of each sample. To compare to these, the coverage depth of bin 17 was

calculated for each sample by taking the sum of all nucleotides mapping to all contigs of bin

17 and dividing by the total length of the contigs, followed by normalisation as above. Using

Pearson's correlation, the coverage of bin 17 was compared to the coverage of all 251 taxa

detected with phyloFlash. The strongest correlation was with one of the FRE consensus

sequences (consensus_all_seqs_0_16). This was found to be correlated with r = 0.97 and an

adjusted p value of 4.5e-11 (Fig. S4). The regression coefficient of 63 between coverage

profiles (consensus_all_seqs_0_16 and bin 17) would imply that there are that many

ribosomal operons in the genome, which is consistent with other fungi.

338   Reference genome access and ecological lifestyle annotation

To compare the MAG to known fungi, a database of reference fungal genomes was

constructed from JGI MycoCosm (Table S3). The fungal species, representing the three

phyla—Ascomycota, Basidiomycota, and Mucoromycota —were chosen based on two

criteria. First, fungal genomes which had been included in previous comparative genomics

analyses of mycorrhizal fungi (Martino et al., 2018; Miyauchi et al., 2020) were included

since the ecology of these fungi had already been curated. Secondly, the database was

supplemented with Mucoromycota genomes available in MycoCosm

(https://mycocosm.jgi.doe.gov/mucoromycota/mucoromycota.info.html) in May 2023. The

associated predicted fungal proteomes were accessed and downloaded in May 2023. We

acknowledge that this was not an exhaustive list of fungal genomes accessible on JGI

MycoCosm in that there were genomes representing further species within Ascomycota and

Basidiomycota that were not included in the analyses. Additionally, with our primary focus

being the provision of comparable context for the MAG, we aimed to obtain a broadly

representative database and not an all-encompassing list.

Fungal genomes were categorised according to their ecological lifestyle (Table S3). To

achieve this, fungal species were coded as follows. First, ecological classifications were

accessed from functional groups recorded on JGI MycoCosm (e.g.,

https://mycocosm.jgi.doe.gov/Ectomycorrhizal_fungi/ Ectomycorrhizal_fungi.info.html).

Secondly, for genomes which were included in previous comparative genomics analyses, the

classifications reported in those studies were used (Chang et al., 2019, 2022; Martino et al.,

2018; Miyauchi et al., 2020). Any remaining genomes without classifications were then

manually curated, as informed by published evidence. Similarly to previous studies, we

allowed a fungal species to occupy multiple ecological lifestyles when there was evidence of

this. We assigned 146 genomes as saprotrophic, 42 as ECM, 29 as pathogenic, 23 as mixed

lifestyle, 12 as G-AMF, 9 as endophytic, 5 as parasitic, 4 as ERM, 3 as OCM, and 1 as

lichenic. Within this total we included the following five Mucoromycotina genomes.

Endogone sp. 'Endsp1' is a putative saprotroph found on fallen oak logs in Florida, USA

(Chang et al., 2019). *Jimgerdemannia lactiflua,* 'Jimlac1' is a putative ectomycorrhizal

fungus isolated from under *Pseudotsuga menziesii* in Oregon, USA (Chang et al., 2019). Two

genomes of the putative ectomycorrhizal fungus *Jimgerdemannia flammicorona*: isolate

'Jimfl_GMNB39_1' from under *Picea abies* and *Picea pungens* in Michigan, USA (Chang et

al., 2019), and isolate 'Jimfl_AD_1' from under *Pinus strobus* in Biella, Italy (Chang et al.,

2019). *Bifiguratus adelaidae* 'Bifad1', a soil-dwelling putative saprotroph isolated from a

Pinus taeda plantation in North Carolina, USA (Torres-Cruz et al., 2017).

Phylogenetic assessment

Phylogenetic assessment of FRE4583 and those consensus sequences from root samples

which were assigned to the order Endogonales (including all_seqs_0_16) was carried out

through 18S rRNA gene alignment. These sequences were added to the Endogonomycetes

18S rRNA alignment from Tedersoo et al. (2024) using MAFFT v7.525 (Katoh and Standley,

2013). The resulting alignment was manually checked and refined in MEGA11 v11.0.13

(Tamura et al., 2021). The best fitting model for our alignment was selected using IQ-TREE

v2.3.6 and the option "-MFP" (Kalyaanamoorthy et al., 2017; Minh et al., 2020). A

phylogenetic tree was generated using the model "TIM2+F+R5" with 1000 approximate

likelihood ratio tests and 1000 ultrafast bootstraps (Hoang et al., 2018; Minh et al., 2020).

The tree was subsequently visualised in FigTree v1.4.4 (Rambaut, 2009).

In addition to the 18S rRNA alignment, fungal phylogeny was inferred from genome

sequences using a concatenated protein marker alignment. A set of 434 protein markers

identified by the JGI as pan-fungal markers for inferring high-level fungal phylogeny was

used to compare the phylogenetic placement of the MAG and all the reference fungal

genomes within our curated database. To identify these marker proteins, an HMM set was

downloaded from https://github.com/1KFG/Phylogenomics_HMMs/tree/master/

HMM/JGI_1086. Phylogenomic markers were extracted from fungal proteomes using the

above HMM, and a subsequent phylogenetic tree was built using PHYling v0.9.0 (available

at: https://github.com/stajichlab/PHYling) under default settings. The tree was then visualised

in R using ggtree v3.6.2 (Yu et al., 2017).

Annotation of MAG and reference genomes

Our comparative analyses focused on Carbohydrate-Active enZymes (CAZymes), lipases,

proteases, transporters, and small-secreted proteins. To identify CAZymes, fungal proteomes

were queried against an HMM profile database built from CAZy sequences downloaded from

the dbCAN2 database (accessed from:

https://bcb.unl.edu/dbCAN2/download/Databases/V11/dbCAN-HMMdb-V11.txt) (Zhang et

al., 2018) using HMMer v 3.3.2 (Eddy, 2011). Protein hits were subsequently filtered with an

e-value threshold of 1e-15 and an alignment threshold of 50% of the domain query. To

identify proteases, fungal proteomes were queried against the MEROPS peptide database

(accessed in July 2023 from:

https://ftp.ebi.ac.uk/pub/databases/merops/current_release/merops_scan.lib) using BLASTp.

Protein hits were subsequently filtered with an e-value threshold of 1e-10. To identify lipases,

fungal proteomes were queried against the Lipase Engineering Database v4.1.0 using

DIAMOND v2.1.8 (Buchfink et al., 2021) and the option "-sensitive". Protein hits were

subsequently filtered with an e-value threshold of 1e-5. To identify transporters, fungal

proteomes were queried against the TCDB transporter database (accessed in November 2023)

(Saier et al., 2021) queried using BLASTp. Protein hits were subsequently filtered with an e-

value threshold of 1e-10. Secreted proteins were identified using a custom pipeline as

reported previously (Pellegrin et al., 2015). Briefly, signal peptides were identified with

SignalP v4.1 (Petersen et al., 2011), extracellular localisation was predicted with WoLFPSort

v0.2 (Horton et al., 2007), transmembrane helices were identified with TMHMM v2.0

(Möller et al., 2001), secretory pathway association was inferred with TargetP v2.0 (Almagro

Armenteros et al., 2019), and the absence of a KDEL motif in the C-terminal region was

confirmed with PS-SCAN revision 1.86 (De Castro et al., 2006). Small-secreted proteins

were identified as secreted proteins < 300 amino acids with no CAZyme, lipase, protease, or

transporter annotation. Functional annotation of putative small-secreted proteins was

achieved with PfamScan v1.6 (Mistry et al., 2007), an e-value threshold of 1e-15, and an

alignment threshold of 50% of the domain query. For each functional protein category being

identified, a single best hit per locus was retained by selecting the hit with the smallest e-

value and greatest alignment of the domain query.

Comparative genomics analyses

All comparative genomics analyses were conducted using R v4.2.2 (R Core Team, 2021), and

all visualisations were generated with ggplot2 v3.4.0 (Wickham, 2016). Functional

annotation of EC numbers was assigned using DeepECtransformer v1.0 (Kim et al., 2023)

and the ECRECer web platform (Shi et al., 2023).

Results

Description of FRE in the roots

The appearance of the FRE in the roots of the twelve samples chosen for metagenomic

sequencing was consistent with the description of the genus *Planticonsortium* by Walker et

al. (2018) into which they placed *Glomus tenue* (previously generically referred to in the

literature as fine endophyte or fine root endophyte; Orchard et al. 2017b) (see photographs in:

Fig. 1 herein; Figure 2 and Figure S1 in Orchard et al., 2017a and Thippayarugs et al.

(1999)). In particular, entry points were characterised by structures with "feather-like, fan-

shaped, palmate or multilobed with finger-like projections" (Walker et al. 2018) which then

led to fine, diverging hyphae in cortex cells and, often, plentiful finely-branched arbuscules.

Small intercalary vesicles and terminal vesicles were also common. Droplets of lipids were

often evident within hyphae and arbuscules. In some instance, colonisation in the cortex was

too dense to clearly distinguish the structure of individual arbuscules.

In the samples used for amplicon and metagenome sequencing the percentage of total root

length colonised AM ranging between 26.8 and 72.5% for AMF with FRE morphology and 0

to 16.3% for G-AMF, producing a proportion of AMF colonisation by FRE between 62.2 and

100%. 18S rRNA Miseq sequencing of Glomeromycotina and Mucoromycotina communities

in root samples revealed that the proportion of Endogonales amplicons ranged between 25

and 82% (Table S1).

Phylogeny and Features of Fungal Genomes

In this study we constructed a putative FRE genome, referred to as the 'MAG'. This MAG

was identified as a putative M-AMF through two lines of evidence. First, the contig bin was

colocated on the assembly graph with unitigs that we could assign to the 18S rRNA gene of a

known M-AMF reference (FRE4583). Secondly, its abundance correlated across the 12

samples in the study with an 18S rRNA sequence 'consensus_all_seqs_0_16', which we

showed to be very closely related to FRE4583 (Fig. 2). The bin itself did not contain a

complete 18S rRNA sequence. This is unsurprising as this gene is well known for failing to

assemble, largely due to it being comprised of strongly conserved regions that repeat

throughout the genome, failing to assemble into substantial contigs. The MAG measured

37.37 Mbp with a genome coverage varying between 1x and 568 depending on sample. The

N50 was found to be 4928 with a GC content of 41.1%. Finally, its completion was assessed

to be 83.7% with only 1.5% contamination (Table S2). We identified a total of 10872

proteins. Of these proteins, we were able to assign 3358 proteins with a functional annotation,

comprising 218 CAZymes, 121 lipases, 280 proteases, 13 small secreted proteins, and 2726

transporters.

While we acknowledge that the revised phylogeny of the Mucoromycotina subphylum

produced in Tedersoo et al. (2024) lacks type specimens and is grounded in part on dark taxa,

as a poorly captured branch of the fungal tree of life we deem this an appropriate basis to

apply the phylogeny of our MAG. 18S rRNA analysis placed consensus_all_seqs_0_16, and

by proxy the MAG, within the order Densosporales of the Mucoromycotina, corresponding

with the Densosporaceae sensu Desirò et al. (2017) (Fig. 2). This order is associated with

FRE morphology (Orchard et al., 2017a; Chang et al., 2019). Further to this, strong support

values allow us to confidently place the MAG in the newly proposed family

Planticonsortiaceae (Tedersoo et al., 2024). Within this family, the MAG was placed with

very high similarity to the known M-AMF reference sequence, FRE4583 (Fig. 2). Further to

this, the MAG was closely related with three other Mucoromycotina 18S rRNA sequences

obtained from the root samples (consensus_all_seqs 0105, 0122 and 0260) and to

environmental sequences obtained from the thalli of early diverging plant sequences, e.g.,

MH174585 and MH174471. The wider *Planticonsortiaceae* clade included further sequences

from the thalli of non-vascular plants, sequences not annotated beyond the phylum level

associated with soil from across Europe (EUK sequences), and one sequence associated with

the rhizosphere of *Populus tremuloides* (EF23703), a known arbuscular mycorrhizal host

(Lee et al., 2020). Phylogenetic analysis therefore placed the MAG apart from other members

of the Mucoromycotina with putative ecology from the order Endogonales, including the

putative ectomycorrhizal Mucoromycotina member *Jimgerdemannia* spp. (Desirò et al.,

2017) and the saprotrophic *Endogone* sp. and *Vinositunica* sp. which have a putative

saprotrophic or ectomycorrhizal lifestyle (Yamamoto et al., 2020).

Similarly, the protein marker alignment placed the MAG within the Mucoromycota (Fig. 3).

The closest relatives are within the order Endogonales, including *Endogone* sp. and

Jimgerdamannia spp, since no Densosporales genomes are available to date. The MAG forms

a sister clade relative to *Endogone* and *Jimgerdamannia*, suggesting it diverged early from

the other Mucoromycotina.

Genomic Profile of the MAG

To contextualise the gene content of the MAG, we compared the CAZyme, protease, lipase,

and transporter profile to genomes within the reference dataset that represented the

saprotrophy-symbiont continuum. We also compared the MAG with the five available

genomes from the Endogonales, which the 18S rRNA and genome analysis confirmed as the

closest phylogenetic relatives to the MAG. The ecological lifestyle had a significant effect on

genome content across all four categories (p < 0.01) (Fig. 4). For the CAZymes, the

ecological lifestyles fell into two groups—G-AMF and ECM, ERM and OCM —with SAP

bridging the two (Fig. 4a). The MAG contained a low number of CAZymes (218), which was

similar to the mean of the ECM (240), but slightly higher than that of the G-AMF (175). A

similar number of CAZymes were found in the MAG compared to related Endogonales

genomes (149 to 208). The ERM genomes contained by far the largest median number of

CAZymes across the lifestyles (629). Only the ERM possessed a significantly (p < 0.01)

larger number of lipases than the other ecological lifestyles (Fig. 4b). The MAG possessed

121 lipases, lower than the means of the G-AMF (189), ECM (238), and SAP (263), OCM

(268). Similarly, the related Endogonales genomes all contained low numbers of lipases (72

to 132). The ERM genomes had significantly (p < 0.01) higher mean numbers of proteases

per genome (457) than the other lifestyles (Fig. 4c). The MAG had similar numbers of

proteases (280) compared to G-AMF (305), ECM (286), SAP (271), and to the related

Endogonales genomes (229 to 298). The G-AMF and ERM lifestyles had significantly higher

(p < 0.05) numbers of transporters per genome relative to the other genomes. The MAG

(2726) had a substantially lower number of transporters compared to G-AMF (3900);

however, it contained more than related Endogonales genomes (1932 to 2388) that sat in the

lower end of the dataset (Fig. 4d). Across the dataset, the saprotrophs contained an especially

high degree of intra-group variability in numbers of all four profiles compared to the other

ecological lifestyles (Fig. 4). The number of CAZymes, lipases, proteases, and transporters in

the MAG were also comparable to the endophyte, parasite pathogen, and mixed lifestyles

(Fig. S5). The MAG contained 13 small secreted proteins, a middling amount for the dataset

(Fig. S5).

CAZyme Profile

CAZyme proteins associated with the degradation of plant and microbial cell walls were

identified to investigate the saprotrophic repertoire of the MAG, in the context of the

mycorrhizal saprotrophy-symbiont continuum. The ERM (252) and OCM (215) genomes

possessed a significantly higher (p < 0.01) mean number of PCWDEs than the G-AMF (32),

ECM (59), and SAP (77) genomes (Fig. 5a). The MAG contained a similar number of

PCWDE (37) as the G-AMF genomes and the genomes of related Endogonales (33 to 44),

which was notably lower than the endophyte, parasite, pathogen, and mixed lifestyles (Fig.

S6). ERM genomes contained a significantly (p < 0.01) higher mean number of MCWDE

(109) than the saprotrophs and other mycorrhizal lifestyles (24 to 59) (Fig. 5b). The G-AMF

genomes had a significantly (p < 0.05) lower number of MWCDE (24) compared to the mean

ECM (45), OCM (59), and SAP (48) genomes. The MAG had 53 MCWDE, higher than the

median number of all other lifestyles except the ERM (Fig. 5b, Fig. S6), and similarly higher

than numbers in the parasite, pathogen and mixed lifestyles, but similar to numbers in

endophytes (Fig. S6). A similar number of MCWDE was detected across the Endogonales

genomes (36 to 53).

Patterns of PCWDE distribution across the mycorrhizal associations were similar (Fig. 6).

ERM and OCM genomes had significantly (p < 0.01) higher mean numbers of cellulases,

hemicellulases, and pectinases than the G-AMF and ECM genomes. Relative to the mean G-

AMF, the MAG had notably higher numbers of cellulases (18 compared to 8), but similar

numbers of hemicellulases (11 compared to 7) and pectinases (4 compared 2). The

Endogonales genomes had similar numbers of cellulases (11 to 13), hemicellulases (10 to

16), and pectinases (4 to 12) as the MAG. The numbers of lignin degrading enzymes were

significantly higher in the ERM (31, p < 0.01) relative to the other mycorrhizal lifestyles,

which all possessed similar numbers: ECM (13), G-AMF (15), OCM (9), and SAP (12).

Notably the MAG had a very low number of lignin degrading enzymes (4) compared to all

other lifestyles, this trend was consistent across the other Endogonales genomes (2 to 8).

Numbers of cellulose, hemicellulose, and pectin degrading enzymes in the MAG were also

typically lower than the mean in endophyte genomes (Fig. S7), and numbers of lignin

degrading genes were similarly notably lower than the endophyte, parasite, pathogen and

mixed lifestyles (Fig. S7).

Cellulases in the MAG genome comprised 8 families, with 6 copies of AA9 being the most

abundant (Table S4). The other Endogonales genomes contained an average of 6 cellulase

families, with GH9 being the most abundant in all four genomes, of which the MAG

contained 3. However, not a single copy of AA9 was identified within the other Endogonales

genomes (Table S4). The G-AMF genomes only contained an average of 3 families, with the

Glomerales containing a mean of just 6 cellulases. However, the Diversisporales contained a

much higher number of cellulases (17 and 12), including 7 and 6 of AA9 and 6 and 3 copies

of AA3, respectively (Table S4). The MAG contained a full suite of cellulose degradation

enzymes: Endoglucanase, β-glucosidase, and exoglucanase (Table S5). However, with the

absence of β-glucosidases in all G-AMF genomes and only two containing a single copy of

exoglucanase, none of the G-AMF genomes possessed the full cellulose degradation pathway

(Table S5). Hemicellulases in the MAG comprised the majority of the remaining PCWDEs,

with 7 families and 5 copies of GH31 being the most abundant (Table S4). Similarly, the

other Endogonales genomes contained an average 8 families, with GH31 being the most

abundant. The G-AMF genomes contained an average 3 families, with GH31 and GH5_7

being the most abundant (Table S4). No copies of GH29 or GH95 were found within the G-

AMF genomes. Similarly, no copies of GH95 were found within the MAG. However, a

single copy of GH29 was present (Table S4).

Pectinases in the MAG were limited, with only 4 copies across 3 families: 2 GH35s, 1 GH2,

and 1 PL1_7 (Table S4). The other Endogonales genomes contained an average of 4 families,

with the most abundant being GH28 and GH35 (Table S4). Within the G-AMF genomes, no

evidence for pectinases was found within 'Gigmar1'. Apart from that, all other genomes

contained only 2 or 3 copies of GH35 (Table S4).

The MAG contained 3 families of lignin degrading enzyme: 2 AA1, 1 AA1_2, and 1 AA2

(Table S4). This trend was conserved across the Endogonales genomes, in which lignin

degrading enzymes comprised of 3 families, with AA1 being slightly more abundant (Table

S4). The G-AMF genomes, while only comprising an average 3 families, contained a much

larger number of lignin degrading enzymes. The Diversisporales contained 37 and 25 lignin

degrading enzymes, including 35 and 23 copies of AA1 respectively. The Glomerales

contained an average of 12 lignin degrading enzymes per genome, with an average of 9

counts being attributed to AA1 (Table S4).

The G-AMF genomes had a smaller mean chitinase repertoire (16) than ERM (35, $p < 0.01$),

OCM (25, $p < 0.05$), and SAP (26, $p < 0.01$) (Fig. 7a). However, the MAG had one of the

highest numbers of chitinases across the genome database (38), roughly double the mean

number found in both the G-AMF and ECM (21), and notably higher than the mean found in

endophytes, parasite, pathogen, and mixed lifestyles (Fig. S8). However, there was

considerable variation in the numbers of chitinases across the G-AMF genomes, with the two

Diversisporales genomes containing 38 and 29 counts respectively, more comparable to the

MAG. The Endogonales genomes similarly contained high mean numbers of chitinases (29 to

42) compared to other genomes in the dataset. Over half of the MAG's chitinase repertoire

was attributed to just two families, CE4 and GH18, with 16 and 13 copies respectively, a

trend also seen across the other Endogonales genomes (Table S4). The G-AMF genomes

largely comprised of CE4 chitinases, however, the Diversisporales genomes also contained a

large number of GH18 (13 and 10), very few of which (0 to 2) were found in Glomerales

(Table S4). The MAG contained a fully comprehensive set of chitin degrading enzymes

(Table S5). Similarly, all but one of the G-AMF genomes contained a full chitin degradation

suite (Table S5).

G-AMF genomes contained significantly ($p < 0.01$) lower mean numbers of glycan degrading

genes (1) compared to the mean of the saprotroph (14) and other mycorrhizal lifestyles (15 to

42), and in fact were only detected in the two Diversisporales genomes (Fig. 7b). In contrast,

the MAG contained 3 glycanases, and this was consistent with a low number of glycanases in

the Endogonales genomes (2 to 8). The MAG only contained 2 copies of GH17 and 1 copy of

GH5_9 (Table S4). The only two other genomes to contain single copies of these were

'Jimfl_AD_1' and 'Jimfl_GMNB39_1'. All the other Endogonales genomes contained 1 or 2

copies of both GH72 and GH81 (Table S4). The Diversisporales genomes only contained low

counts of GH5_9 and GH81 (Table S4).

ERM genomes contained a higher mean number of mannanases (24) than the saprotrophic

and other mycorrhizal lifestyles (3 to 6) (Fig. 7c). However, the MAG contained a

substantially higher mean number of mannanases (9) than the G-AMF (3) and ECM (5), and

also the Endogonales genomes (2 to 4). Similarly, there was a range of mannanases across the

G-AMF genomes with the two Diversisporales genomes containing 11 and 10 counts

respectively, more comparable to the MAG (Table S4). The MAG again contained a

comparatively large set of genes for mannan degradation, with 5 copies of GH92, 3 copies of

GH125, and 1 copy of GH76. However, this represents the one substrate-specific group that

was not as conserved across the phylogenetically closest relatives. None of the Endogonales

genomes contained copies of GH76, while all contained 1 or 2 copies of CE17 (Table S4).

Additionally, while three of the four contained GH92, it was only in single-copy instances.

The G-AMF genomes showed a range of mannanases within their genomes, and particularly

the Diversisporales genomes alone contained many copies of GH92. All but one Glomerales

genome only contained 1 or 2 copies of GH125, with no mannaanase identified in the

'RhiirC2' genome (Table S4).

Low numbers of peptidoglycanases were found in the genomes of all lifestyles, with

significantly larger mean numbers in ERM (8, $p = <0.01$) compared to the ECM (3) and

saprotroph (2) genomes (Fig. 7d). The G-AMF genomes had a comparable mean number of

peptodoglycanases (4) to the MAG (3), while the corresponding numbers in the Endogonales

genomes were also low (0 to 3). The MAG only contained 2 copies of GH24 and a single

copy of GH89 (Table S4). Similarly, the other Endogonales genomes were sparse in

peptidoglycanases. Specifically, the 'Jimfl_GMNB39_1' genome contained no

peptidoglycanases, while 'Jimfl_AD_1' and 'Jimlac1' only possessed a single copy of GH89

and CMB50, respectively. 'Endsp1' also contained 2 copies of CBM50 and a single copy of

GH24. The G-AMF genomes also contained a small number of peptidoglycanases, mostly

consisting of GH24 (Table S4).

The MAG contained no genome-specific cell wall degrading enzymes families. However, a

poorly described GH151 family, "α-L-fucosidase enzyme encoding protein", was only

identified within 3 of the closest phylogenetic relatives (Jimfl_AD_1, Jimfl_GMNB39_1, and

Endsp1) and a far-removed saprotrophic basidiomycete, Dacsp1 (Table S4).

Transporters

Since the ability to transport nutrients and assimilate with the host plant is core to the lifestyle

of a fungal symbiont, we compared substrate-specific transporters between the ecological

lifestyles. Only the ERM genomes contained a statistically higher ($p = 0.044$) number of

phosphate transporters than the other genomes (Fig. 8a). The MAG contained a relatively low

number of phosphate transporters (14), although this was not too dissimilar from the mean of

the G-AMF (18) and the same as the mean ECM genome (14). Similarly, the Endogonales

genomes possessed low numbers of phosphate transporters (13 to 15) (Fig. 8a). The G-AMF

genomes had a significantly higher ($p < 0.01$) number of calcium transporters (55) compared

to the other mycorrhizal and saprotrophic lifestyles (Fig. 8b, which ranged between 26 and

39), and also the endophyte, parasite, pathogen, and mixed lifestyles (Fig. S9). The MAG

showed a similar number of calcium transporters (50) to the G-AMF, while the number of

calcium transporters in the Endogonales genomes varied considerably (25 to 47) (Fig. 8b).

There was no significant difference ($p = 0.48$) in the number of copper transporters across the

mycorrhizal and saprotrophic genomes, with the mean varying between 13 and 17 (Fig. 8c).

However, the MAG contained only 6 copper transporters, one of the lowest of all genomes

(Fig. 8c, Fig. S9). Of the Endogonales genomes, 'Jimlac1' had the second smallest number of

copper transporters (7) while the rest ranged from 8 to 13 (Fig. 8c). The G-AMF had a

significantly higher ($p < 0.01$) number of nitrate transporters (7) compared to the other

mycorrhizal and saprotrophic lifestyles (Fig. 8d). The MAG contained a low number (2) of

nitrate transporters relative to the mean G-AMF, although this was the same as the mean

ECM (2) and similar to the ERM (3), OCM (1), and SAP (1). There was considerable

variation in the mean number of nitrate transporters within the Endogonales genomes (0-5)

(Fig. 8d). The ERM genomes had a significantly higher ($p < 0.01$) mean number of

ammonium transporters (11) than all but the G-AMF genomes (9) (Fig. 8e). The MAG

contained 5 ammonium transporters, slightly higher than the median ECM (4) and OCM (4).

The MAG contained an especially low number of glycine transporters (1), the same as the

average ECM (1) and OCM (1) genome (Fig. 8f). The ERM contained the highest mean

number of glycine transporters (5), whereas the G-AMF (3) and SAP (4) were not

significantly different to any of the other ecological lifestyles (Fig. 8f).

Hallmark Missing Glomeromycotan Core Genes

We analysed both the MAG and G-AMF genomes for enzymes linked to pathways that are

missing from typical G-AMFs, often referred to as 'Missing Glomeromycotan Core Genes'

(MGCG). As fatty acid auxotrophs, G-AMF have been shown to lack the presence of

cytosolic fatty acid synthases required for fatty acid biosynthesis. We found no evidence of

proteins directly associated with the fatty acid synthase required to facilitate fatty acid

biosynthesis within either the G-AMF genomes or the MAG. We did identify 3-oxoacyl-ACP

reductase in all genomes, including 10 within the MAG. (Table S5). Additionally, neither the

MAG nor the G-AMF genomes possessed the ability to biosynthesis thiamine, a hallmark of

G-AMF genomes. Specifically, we found no evidence of the presence of thiamine phosphate

synthase or hydroxymethylpyrimidine kinase (Table S5).

The ability to synthesis all derivatives of the vitamin B6 pathway was present within the

MAG. This included a copy of pyridoxine kinase (2.7.1.35), the most prolific enzyme within

the vitamin B6 metabolism pathway. Only one enzyme, pyridoxine 4-dehydrogenase, was

missing from the pathway, while an indirect three-step metabolism for the conversion of

pyridoxal to pyridoxine was present. Conversely, no G-AMF genome possessed a full

vitamin B6 metabolism pathway, with no evidence of pyridoxine kinase being found across

the group (Table S5).

Discussion

In the current work we assembled the genome of a putative FRE-forming AMF from a root

metagenome. Methodologically this is a relatively unique achievement: fungal MAGs have

been obtained previously from host associated environments and niche environments, but not

from complex environments, such as the rhizosphere (Tagirdzhanova et al., 2021; Shen et al.,

2023; Singh et al., 2023), or from similar fungal symbionts. To achieve this, we had to

augment standard binning methods with assembly graph visualisations and coverage

correlations to known M-AMF 18S rRNA genes. We believe that this method has many

advantages over isolation and culturing, in particular it is less time consuming and in

principle may be more reflective of the actual community present particularly if multiple

species are present.

Phylogenetic analysis of the correlative 18S rRNA marker, consensus_all_seqs_0_16,

showed that the MAG sits within the order Densosporales of the Mucoromycotina. The MAG

represents the first published genome of both the order Densosporales and the family

Planticonsortiaceae. The most closely related taxa with sequenced genomes are putative

ectomycorrhizal and saprotrophic fungi from the order Endogonales, although a reasonable

degree of dissimilarity is evident. The MAG was closely related to environmental sequences

from the rhizoids of liverworts, which form arbuscular-like symbioses involving

Glomeromycota and Mucoromycotina (Rimmington, 2018). To date, understanding of the

phylogeny of fungi which form mycorrhizas with FRE morphology in vascular plants has

been restricted to short 18S rRNA fragments. Similarly, these studies have confirmed close phylogenetic relatedness with sequences from liverwort thalli (Orchard et al., 2017; Albornoz et al., 2021, 2022; Seeliger et al., 2024), suggesting these fungi form associations with both vascular and non-vascular plants, with a role in N and P nutrition of the plant host (Orchard et al, 2017a; Hoystead, 2023). Mucoromycotina sequences associated with plant roots may span both the Densosporales (Orchard et al., 2017b; Desirò et al., 2017; Chang et al., 2018) and Endogonales (Seeliger et al., 2024), and furthermore diverse assemblages of fungi from these orders may be present in roots (Seeliger et al., 2024), although fungi associated with the FRE morphology appear to be phylogenetically distinct from putative ectomycorrhizal and saprotrophic Endogonales (Seeliger et al., 2024).

Comparative genome analysis showed that the MAG's genome had many features in common with G-AMF. Some of these genomic features, including the low abundance of PCWDE and the absence of thiamine biosynthesis genes, were also seen in related Endogonales genomes and therefore aren't likely to be specific, hallmark features of an adaptation to a symbiotic lifestyle. However other signatures the MAG shares with G-AMF, including a high abundance of calcium transporters and the absence of fatty acid synthases, may point to physiological convergence in the way in which the symbioses operate. There were some notable differences between the MAG and G-AMF, with the MAG possessing a high abundance of chitinases, a complete vitamin B6 biosynthesis pathway, and low abundances of nitrate, ammonium and copper transporters. These features may point to differences between G-AMF and M-AMF in the way they interact with the host plant and the wider environment.

It has been suggested that the transition from saprotrophic to mycorrhizal lifestyles involved loss of PCWDE, enabling fungi to exist within roots without digesting them (Miyauchi et al., 2020). The G-AMF symbiosis is estimated to have evolved 450 mya, and Glomeromycotan genomes possess substantially lower numbers of these genes than the fungi forming the more recently evolved ectomycorrhizal, ericoid mycorrhizal, and orchid mycorrhizal symbioses. Furthermore, G-AMF show signatures of dependence on host metabolism that are absent from more recently evolved symbioses, including fatty acid auxotrophy and an inability to synthesise thiamine and metabolise vitamin B6 (Miyauchi et al., 2020). These represent the 'Missing Glomeromycota Core Genes' (MGCG, Tang et al., 2016; Kobayashi et al., 2018; Rosling et al., 2024).

Although genomic evidence indicates that G-AMF and M-AMF have independent origins
(Rosling et al., 2024), these symbioses co-occurred in the earliest land plants (Strullu-Derrien
et al., 2014), and they continue to co-occur in roots of extant plants (Orchard et al., 2017a).
Additionally, while G-AMF and M-AMF exhibit some niche differentiation, evidence of
overlapping ecological presence and similar responses to overarching edaphic and
environmental variables have been noted (Albornoz et al., 2022; Seeliger et al., 2024).
Furthermore, ultrastructural evidence points to similarities in the way in which G-AMF and
M-AMF function at a cellular level (Albornoz et al., 2020). Therefore, similar evolutionary
pressures may have acted on these symbioses, which could have resulted in convergence of
genomic traits. Indeed, many of the symbiotic genomic features shown by G-AMF were also
displayed by the MAG. This included a reduced repertoire of PCWDE and the absence of
several MGCG, including those related to thiamine and fatty acid biosynthesis.

Our analysis confirms the findings of Rosling et al. (2024) that Endogonales have a low
genomic PCWDE count and an incomplete thiamine biosynthesis pathway, suggesting that
these may be features common to both the Densosporales and Endogonales, rather than an
adaptation to biotrophy. However, the Endogonales genomes possess fatty acid biosynthesis
genes (Rosling et al., 2024), so this signature of biotrophy does likely represent the same
adaptation to biotrophy in G-AMF and M-AMF, which suggests that host assimilate is
provided as fatty acids in both symbioses.

The MAG had a number of features which distinguished it from G-AMF genomes. In
particular, while there was a shared low abundance of CAZymes, the MAG possessed a
greater repertoire of enzymes for degradation of organic substrates than typically seen in G-
AMF genomes. This included the full complement of enzymes required for degradation of
cellulose, which was absent in all G-AMF genomes. In notable contrast to G-AMF, the MAG
had a high count of both chitin and mannan fungal cell wall degrading enzymes. The high
chitinase count doesn't appear to be specific to the M-AMF, with the putative
ectomycorrhizal and saprotrophic *Endogonales* genomes showing a similar pattern, attributed
to elevated numbers of CE4 and GH18, as seen previously in Rosling et al. (2024). However,
the full complement of chitinases was present in the G-AMF genomes as well as in the MAG,
so this feature may not have any functional significance. Relative to other mycorrhizal fungi
(except ericoid mycorrhizal fungi), the MAG had a notably high count of genes for mannan
degradation. These were absent or present in low numbers in most G-AMF genomes and,
significantly, were also found in low abundance in the *Endogonales* genomes, therefore high

abundance of mannan degradation genes may represent a distinguishing feature of FRE

genomes. However, even within the G-AMF there appears to be some variation, with the

Diversisporales genomes containing a higher number of mannan and chitin degrading

enzymes than the *Glomerales* genomes, indicating a substantial range of genomic repertoires.

It is important to understand whether these contrasting enzymatic profiles contribute to niche

partitioning between G-AMF and M-AMF in nutrient mobilisation from contrasting mineral

and organic substrates. For example, high abundance of chitin and mannan degrading genes

could indicate a role for M-AMF in the degradation of fungal necromass, while a complete

cellulose degradation pathway and high abundance of cellulose and hemicellulose degrading

enzymes could suggest involvement in saprotrophic breakdown of plant-derived materials,

similar to some ECM fungi (Janusz et al., 2017; Martin & van der Heijden., 2024).

Alternatively, it could reflect different strategies for enzymatic degradation of plant cell walls

as they colonise and grow through host plant tissues and penetrate root cells.

Currently there is no indication that soil organic matter has any influence on the distribution

of G-AMF or M-AMF. In contrast to G-AMF, evidence indicates that M-AMF are more

abundant in agricultural than natural habitats, indicating a preference for low organic matter

soils (Albornoz et al., 2022). However, these functional differences could support niche

partitioning within these habitats.

A further distinguishing feature between the MAG and the G-AMF genomes is the presence

of a full B6 biosynthesis pathway in the MAG, including pyroxidine kinase, which was

missing in all the G-AMF genomes, as demonstrated in earlier studies (Tang et al., 2016;

Kobayashi et al., 2018). B6 is a cofactor required for amino acid biosynthesis, and it also

plays a role as an antioxidant. Reactive oxygen species (ROS) scavenging systems such as B6

may be required by AMF to overcome host defences, allowing intracellular colonisation and

the development of arbuscules (Belmondo et al., 2016; Lanfranco et al., 2018). Components

of the B6 biosynthesis pathway that G-AMF do possess are expressed throughout the fungal

life cycle, but they are at the highest expression levels within the root (Benabdellah et al.,

2009). Similarly, B6 plays a role in supporting the infection of root tissues by pathogens

(Wang et al., 2023). Differences in capacity for independent production of B6 between the

MAG and G-AMF could point to differences in the way the fungi scavenge ROS, and

potentially also differences in the way root colonisation and arbuscule development are

regulated by the plant. There could also be implications for the specific mechanisms

underlying M-AMF and G-AMF responses to external stressors which generate ROS such as

temperature, osmotic stress, and nutrient limitation, and the independence of these responses
to the plant's own response (Sahu et al., 2022).

Key physiological traits associated with G-AMF symbioses are the uptake of P from the soil,
its translocation through hyphae to the root, and its transfer to the plant in exchange for
carbon. Ultrastructural (Albornoz et al., 2020) and experimental evidence (Orchard et al.,
2017b) suggest a similar role for M-AMF. A number of P transporters have been identified in
G-AMF involved in P assimilation from soil, storage of P, and exchange of P with the host
(Javot et al., 2006; Harrison et al., 2010; Plassard et al., 2019). G-AMF genomes generally
showed higher mean numbers of P transporters than ECM, while numbers in the MAG were
low relative to other mycorrhizal genomes studied. While the significance of transporters to
the functioning of symbionts will depend on a range of factors including their efficiency, our
data doesn't provide genomic support for particular specialisation in P nutrition by M-AMF.

Evidence also points to a potential role for G-AMF in N nutrition (Gosling et al., 2013).
Indeed, G-AMF genomes contained higher numbers of nitrate transporters than the other
mycorrhizal symbionts, while also sitting towards the higher end of both ammonium and
glycine transporters within the genomes we analysed. However, further evidence of inter-
group differentiation can be seen across the Diversisporales and Glomerales genomes, with
the former containing especially high numbers of ammonium and glycine transporters. The
MAG contained a lower number of all three categories of N transporters than all G-AMF
genomes. While the MAG contained a typical number of ammonium transporters compared
to the other mycorrhizal lifestyles, it contained a low number of nitrate and glycine
transporters. Hoystead et al. (2023) suggested that M-AMF and G-AMF have complementary
roles in host nutrition, with M-AMF predominantly involved in supply of N to the host and
G-AMF in supplying P. Furthermore, Howard et al. (2024) showed a putative M-AMF
displaying both a preference for, and increased transfer of, glycine and ammonium derived
nitrogen sources. While there is evidence for the ability to transport varying forms of N, our
MAG does not provide exceptional evidence for this relative specialism.

The very low abundance of Cu transporters in the MAG - the lowest across the entire dataset
- is particularly striking. The three ectomycorrhizal Endogonales genomes also contain low
numbers of Cu transporters, however the saprotrophic Endogonales sit near the mean. G-
AMF may promote Cu uptake by the host plant (Taylor et al., 2015) and arbuscules have high
demands for Cu (Gomez-Gallego et al., 2019). Furthermore, Cu transporters in G-AMF show
differential expression profiles across intra- and extra- radical mycelium (Gomez-Gallego et

al., 2019). The extent to which M-AMF are involved in the supply of micronutrients to the host remains to be determined, but the evidence presented here could indicate that in contrast to G-AMF, this may not be an important function for M-AMF.

The high number of calcium transporters in both the MAG and G-AMF is a clear point of similarity between them, and of differentiation to all other groups. AMF have been implicated in transport of Ca to host plants (e.g. Liu et al., 2002), and our data could therefore indicate a role for both G-AMF and M-AMF in this function. However, Ca may also play a key role in the arbuscule lifecycle of both FRE and G-AMF. A clear link between a dramatic increase in calcium accumulation and arbuscule collapse has been demonstrated within G-AMF by both Ryan et al. (2003) and Sujkowska-Rybkowska and Znojek (2018). A similar trend within FRE was documented by Gianinazzi-Pearson et al. (1981) and Albornoz et al. (2020). While it is unknown whether the Ca accumulated originates from the host plant or the fungi, the high number of Ca transporters within both G-AMF and the MAG may indicate that Ca accumulation is driven by the fungal symbionts, perhaps depositing Ca to cease the function of the arbuscule once it reaches maturity.

Genome analyses, including those reported here, have identified differences across G-AMF which suggest contrasting interactions with the host and the wider environment. For both G-AMF and M-AMF, comprehensive analysis of genomic functional gene repertoires across the broad range of taxa found within roots is key to understanding the way that AMF communities support plant health, and the potential for engineering communities to promote beneficial traits. With the development of third generation long read sequencing technologies, genome assembly from metagenomes represents a promising route for understanding functions and traits associated with dark taxa in complex microbial communities, especially when combined with transcriptome sequencing (Benoit et al., 2024) to understand the factors which regulate gene expression within plant tissues and the soil.

Acknowledgements

We thank the Natural Environment Research Council (Grant NE/S010270/1), the Biotechnology and Biological Sciences Research Council (BBSRC) Midlands Integrative Biosciences Training Partnership (Grant BB/T00746X/1) and the Australian Research Council (Discovery Project DP180103157) for funding.

Competing interests

None declared.

Author contributions

GB, MHR, and CQ conceived the work and with RM obtained funding. MHR and PAD
conducted the M-AMF enrichment work and the assessment of M-AMF colonisation of plant
roots. SH, SR, and GB conducted sequencing and analysis of 18S rRNA genes. SR and CQ
assembled the MAG. RHN, GM, and JC conducted the 18S rRNA gene phylogenetic
analysis. JC, GM, SR, and GB performed the comparative genomic analyses. JC and GB
wrote the manuscript. All authors edited the manuscript.

Data Availability

Raw 18S rRNA Miseq sequences are available at NCBI (BioProject ID PRJNA1179892), and
Nanopore sequences at ENA (Experiment Accession ERX13330708). The FRE4583
sequence is available at NCBI (SUB14894546). The 18S rRNA phylogeny sequence
alignment FASTA file and Newick tree file are available on PlutoF Biodiversity Platform
(https://dx.doi.org/10.15156/BIO/3301225). The sequence of the MAG has been deposited on
the Zenodo repository (https://doi.org/10.5281/zenodo.14144746).

References
Albornoz FE, Hayes PE, Orchard S, Clode PL, Nazeri NK, Standish RJ, Bending GD, Hilton
S, Ryan MH. 2020. First cryo-scanning electron microscopy images and X-ray microanalyses
of mucoromycotinian fine root endophytes in vascular plants. *Frontiers in Microbiology* 11:
2018.

Albornoz FE, Orchard S, Standish RJ, Dickie IA, Bending GD, Hilton S, Lardner T, Foster
KJ, Gleeson DB, Bougoure J et al. 2021. Evidence for niche differentiation in the
environmental responses of co-occurring mucoromycotinian fine root endophytes and
glomeromycotinian arbuscular mycorrhizal fungi. *Microbial Ecology* 81(4): 864-873.

Albornoz FE, Ryan MH, Bending GD, Hilton S, Dickie IA, Gleeson DB, Standish RJ. 2022.

Agricultural land-use favours Mucoromycotinian, but not Glomeromycotinian, arbuscular

mycorrhizal fungi across ten biomes. *New Phytologist* 233(3): 1369-1382.

Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, ljaz UZ, Loman NJ, Andersson

AF, Quince C. 2014. Binning metagenomic contigs by coverage and composition. *Nature*

Methods 11: 1144–1146.

Almagro Armenteros JJ, Salvatore M, Emanuelsson O, Winther O, von Heijne G, Elofsson

A, Nielsen H. 2019. Detecting sequence signals in targeting peptides using deep learning.

Life Science Alliance 2(5): e201900429.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD et al 2012. SPAdes: a new genome assembly algorithm

and its applications to single-cell sequencing. *Journal of Computational Biology* 19(5): 455-

477.

Begum N, Qin C, Ahanger MA, Raza S, Khan MI, Ashraf M, Ahmed N, Zhang L. 2019. Role

of arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress

tolerance. *Frontiers in Plant Science* 10: 1068.

Belmondo S, Calcagno C, Genre A, Puppo A, Pauly N, Lanfranco L. 2016. NADPH oxidases

in the arbuscular mycorrhizal symbiosis. *Plant Signaling & Behavior* 11(4): e1165379.

Benabdellah L, Azcón-Aguilar C, Valderas A, Speziga D, Fitzpatrick TB, Ferrol N. 2009.

GintPDX1 encodes a protein involved in vitamin B6 biosynthesis that is up-regulated by

oxidative stress in the arbuscular mycorrhizal fungus *Glomus intraradices*. *New Phytologist*

184(3): 682-693.

Benoit G, Raguideau S, James R, Phillippy AM, Chikhi R, Quince C. 2024. High-quality

metagenome assembly from long accurate reads with metaMDBG. *Nature Biotechnology* 42:

1378–1383.

Brundrett MC, Tedersoo L. 2018. Evolutionary history of mycorrhizal symbioses and global

host plant diversity. *New Phytologist* 220(4):1108-1115.

Brundrett MC. 2006. Understanding the roles of multifunctional mycorrhizal and endophytic

fungi. In: BJE Schulz, CJC Boyle, TN Sieber, eds. *Microbial Root Endophytes*. Berlin,

Germany: Springer-Verlag, 281–293.

Buchfink B, Reuter K, Drost HG. 2021. Sensitive protein alignments at tree-of-life scale

using DIAMOND. *Nature Methods* 18: 366-368.

Chang Y, Desirò A, Na H, Sandor L, Lipzen A, Clum A, Barry K, Grigoriev IV, Martin FM,

Stajich JE et al. 2019. Phylogenomics of Endogonales and evolution of mycorrhizas within

Mucoromycota. *New Phytologist* 222(1): 511-525.

Chang Y, Wang Y, Mondo S, Ahrendt S, Andreopoulos W, Barry K, Beard J, Benny GL,

Blankenship S, Bonito G et al. 2022. Evolution of zygomycete secretomes and the origins of

terrestrial fungal ecologies. *iScience* 25(8): 104840.

Chen J, Tang Y, Kohler A, Lebreton A, Xing YM, Zhou D, Li Y, Martin FM, Guo S. 2022.

Comparative transcriptomics analysis of the symbiotic germination of *D. officinale*

(Orchidaceae) with emphasis on plant cell wall modification and cell wall-degrading

enzymes. *Frontiers in Plant Science* 13:880600.

Chen M, Arato M, Borghi L, Nouri E, Reinhardt D. 2018. Beneficial services of arbuscular

mycorrhizal fungi – from ecology to application. *Frontiers in Plant Science* 9: 1270.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,

McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and BCFtools.

GigaScience 10(2): giab008.

De Castro E, Sigrist CJ, Gattiker A, Bulliard V, Langendijk-Genevaux PS, Gasteiger E,

Bairoch A, Hulo N. 2006. ScanProsite: detection of PROSITE signature matches and

ProRule-associated functional and structural residues in proteins. *Nucleic Acids Research* 34:

362-5.

Desirò A, Rimington WR, Jacob AS, Pol NV, Smith MR, Trappe JM, Bidartondo MI,

Bonito G. 2017. Multigene phylogeny of Endogonales, an early diverging lineage of fungi

associated with plants. *IMA Fungus* 8(2): 245–257.

Eddy SR. 2011. Accelerated profile HMM searches. *PLOS Computational Biology* 7(10):

e1002195.

Frąc M, Hannula SE, Bełka M, Jędryczka M. 2018. Fungal biodiversity and their role in soil

health. *Frontiers in Microbiology* 9: 707.

Gianinazzi-Pearson V, Morandi D, Dexheimer J, Gianinazzi S. 1981. Ultrastructural and

ultracytochemical features of a *Glomus tenuis* mycorrhiza. *New Phytologist* 88(4): 633-639.

Gómez-Gallego T, Benabdellah K, Merlos MA, Jiménez-Jiménez AM, Alcon C, Berthomieu

P, Ferrol N. 2019. The *Rhizophagus irregularis* genome encodes two CTR copper

transporters that mediate Cu import into the cytosol and a CTR-like protein likely involved in

copper tolerance. *Frontiers in Plant Science* 10: 604.

Gong Y, Lebreton A, Zhang F, Martin FM. 2023. Role of carbohydrate-active enzymes in

mycorrhizal symbioses. *Essays in Biochemistry* 67(3): 471-478.

Gosling P, Mead A, Proctor M, Hammond JP, Bending GD. 2013. Contrasting arbuscular

mycorrhizal communities colonizing different host plants show a similar response to a soil

phosphorus concentration gradient. *New Phytologist* 198(2): 546-556.

Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao X,

Korzeniewski F et al. 2013. MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic

Acids Research* 42: 699-704.

1000

1001 Gruber-Vodicka HR, Seah BKB, Pruesse E. 2020. phyloFlash: Rapid small-subunit rRNA

1002 profiling and targeted assembly from metagenomes. m*Systems* 5(5): e00920-20.

1003

1004 Harrison MJ, Pumplin N, Breuillin FJ, Noar RD, Park HJ. 2010. Phosphate transporters in

1005 arbuscular mycorrhizal symbiosis. In: Koltai H, Kapulnik Y, eds. *Arbuscular Mycorrhizas:*

1006 *Physiology and Function.* Dordrecht, Netherlands: Springer, 117–135.

1007

1008 Hawkins HJ, Cargill RIM, Nuland MEV, Hagen SC, Field KJ, Sheldrake M, Soudzilovskaia

1009 NA, Kiers TE. 2023. Mycorrhizal mycelium as a global carbon pool. *Current Biology* 33(11):

1010 560-573.

1011

1012 Hirose D, Degawa Y, Yamamoto K, Yamada A. 2014. Sphaerocreas pubescens is a member

1013 of the Mucoromycotina closely related to fungi associated with liverworts and hornworts.

1014 *Mycoscience* 55(3): 221-226.

1015

1016 Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving

1017 the ultrafast bootstrap approximation. *Molecular Biology and Evolution*. 35: 518–522.

1018

1019 Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai N. 2007.

1020 WoLF PSORT: protein localization predictor. *Nucleic Acids Research* 5(2): 585–587.

1021

1022 Howard NAO, Williams A, Durant E, Pressel S, Daniell TJ, Field KJ. 2024. Preferential

1023 nitrogen and carbon exchange dynamics in Mucoromycotina "fine root endophyte"-plant

1024 symbiosis. *Current Biology*.

1025

1026 Hoysted GA, Field KJ, Sinanaj B, Bell CA, Bidartondo MI, Pressel S. 2022. Direct nitrogen,

1027 phosphorus and carbon exchanges between Mucoromycotina 'fine root endophyte' fungi and

1028 a flowering plant in novel monoxenic cultures. *New Phytologist* 238(1): 70-79.

1029

1030 Hoysted GA, Jacob AS, Kowal J, Giesemann P, Bidartondo MI, Duckett JG, Gebauer G,

1031 Rimington WR, Schornack S, Pressel S et al. 2019. Mucoromycotina fine root endophyte

1032 fungi form nutritional mutualisms with vascular plants. *Plant Physiology* 181(2): 565-577.

1033

1034 Huey CJ, Gopinath SCB, Uda MNA, Zulhaimi HI, Jaafar MN, Kasim FH, Yaakub ARW.

1035 2020. Mycorrhiza: a natural resource assists plant growth under varied soil conditions.

1036 *3Biotech* 10(5): 204.

1037

1038 Janusz G, Pawlik A, Sulej J, Świderska-Burek U, Jarosz-Wilkołazka A, Paszczyński A. 2017.

1039 Lignin degradation: microorganisms, enzymes involved, genomes analysis and evolution.

1040 *FEMS Microbiology Reviews* 41(6): 941-962.

1041

1042 Javot H, Pumplin N, Harrison MJ. 2006. Phosphate in the arbuscular mycorrhizal symbiosis:

1043 Transport properties and regulatory roles. *Plant, Cell & Environment* 30: 310–322.

1044

1045 Jeffery RP, Simpson RJ, Lambers H, Orchard S, Kidd DR, Haling RE, Ryan MH. 2018.

1046 Contrasting communities of arbuscule-forming root symbionts change external critical

1047 phosphorus requirements of some annual pasture legumes. *Applied Soil Ecology* 126: 88-97.

1048

1049 Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:

1050 Fast model selection for accurate phylogenetic estimates. *Nature Methods.* 14: 587-589.

1051

1052 Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

1053 improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–

1054 780.

1055

1056 Kim GB, Kim JY, Lee JA, Norsigian CJ, Palsson BO, Lee SY. 2023. Functional annotation

1057 of enzyme-encoding genes using deep learning with transformer layers. *Nature*

1058 *Communications* 14: 7370.

1059

1060 Kobayashi Y, Maeda T, Yamaguchi K, Kameoka H, Tanaka S, Ezawa T, Shigenobu S,

1061 Kawaguchi M. 2018. The genome of Rhizophagus clarus HR1 reveals a common genetic

1062 basis for auxotrophy among arbuscular mycorrhizal fungi. *BMC Genomics* 19(1): 465.

1063

1064 Lanfranco L, Fiorilli V, Gutjahr C. 2018. Partner communication and role of nutrients in the

1065 arbuscular mycorrhizal symbiosis. *New Phytologist* 220(4): 1031-1046.

1066

1067 Lebreton A, Zeng Q, Miyauchi S, Kohler A, Dai YC, Martin FM. 2021. Evolution of the

1068 mode of nutrition in symbiotic and saprotrophic fungi in forest ecosystems. *Annual Review of*

1069 *Ecology, Evolution, and Systematics* 52(1): 385-404.

1070

1071 Lee S-J, Kong M, St-Arnaud M, Hijri M. 2020. Arbuscular mycorrhizal fungal communities

1072 of native plant species under high petroleum hydrocarbon contamination highlights

1073 Rhizophagus as a key tolerant genus. *Microorganisms* 8(6): 872.

1074

1075 Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

1076 MEM. *ArXiv*. doi: 10.48550/arXiv.1303.3997

1077

1078 Liu A, Hamel C, Elmi A, Costa C, Ma B, Smith DL. 2002. Concentrations of K, Ca and Mg

1079 in maize colonized by arbuscular mycorrhizal fungi under field conditions. *Canadian Journal*

1080 *of Soil Science* 82(3): 271-279.

1081

1082 Luginbuehl LH, Menard GN, Kurup S, Van Erp H, Radhakrishnan GV, Breakspear A,

1083 Oldroyd GED, Eastmond PJ. 2017. Fatty acids in arbuscular mycorrhizal fungi are

1084 synthesized by the host plant. *Science* 365(6343): 1175-1178.

1085

1086 Luginbuehl LH, Oldroyd GED. 2017. Understanding the arbuscule at the heart of

1087 endomycorrhizal symbioses in plants. *Current Biology* 27(17): 952-963.

1088

1089 Maillard F, Schilling J, Andrews E, Schreiner KM, Kennedy P. 2020. Functional

1090 convergence in the decomposition of fungal necromass in soil and wood. *FEMS*

1091 *Microbiology Ecology* 96(2): fiz209.

1092

1093 Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel

1094 and streamlined workflows along with broader and deeper phylogenetic coverage for scoring

1095 of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution* 38(10):

1096 4647-4654.

1097

1098 Mansfield TM, Albornoz FE, Ryan MH, Bending GD, Standish RJ. 2023. Niche

1099 differentiation of Mucoromycotinian and Glomeromycotinian arbuscular mycorrhizal fungi

1100 along a 2-million-year soil chronosequence. *Mycorrhiza* 33(3): 139-152.

1101

1102 Martin FM, van der Heijden MGA. 2024. The mycorrhizal symbiosis: research frontiers in

1103 genomics, ecology, and agricultural application. *New Phytologist* 242(4): 1486-1506.

1104

1105   Martino E, Morin E, Grelet G-A, Kuo A, Kohler A, Daghino S, Barry KW, Cichocki N,

1106   Clum A, Dockter RB et al. 2018. Comparative genomics and transcriptomics depict ericoid

1107   mycorrhizal fungi as versatile saprotrophs and plant mutualists. *New Phytologist* 217(3):

1108   1213-1229.

1109

1110   McGonigle TP, Miller MH, Evans DG, Fairchild GL, Swan JA. 1990. A new method which

1111   gives an objective measure of colonization of roots by vesicular—arbuscular mycorrhizal

1112   fungi. *New Phytologist* 155(3): 495-501.

1113

1114   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

1115   Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference

1116   in the genomic era. *Molecular Biology and Evolution*. 37(5): 1530-1534.

1117

1118   Mistry J, Bateman A, Finn RD. 2007. Predicting active site residue annotations in the Pfam

1119   database. *BMC Bioinformatics* 8: 298.

1120

1121   Miyauchi S, Kiss E, Kuo A, Drula E, Kohler A, Sánchez-García M, Morin E, Andreopoulos

1122   B, Barry KW, Bonito G et al. 2020. Large-scale genome sequencing of mycorrhizal fungi

1123   provides insights into the early evolution of symbiotic traits. *Nature Communications* 11:

1124   5125.

1125

1126   Möller S, Croning MDR, Apweiler R. 2001. Evaluation of methods for the prediction of

1127   membrane spanning regions. *Bioinformatics* 17(7): 646–653.

1128

1129   Montoliu-Nerin M, Sánchez-García M, Bergin C, Kutschera VE, Johannesson H, Bever JD,

1130   Rosling A. 2021. In-depth phylogenomic analysis of arbuscular mycorrhizal fungi based on a

1131   comprehensive set of de novo genome assemblies. *Frontiers in Fungal Biology* 29(2):

1132   716385.

1133

1134   Morin E, Miyauchi S, Clemente HS, Chen ECH, Pelin A, Providencia I, Ndikumana S,

1135   Beaudet D, Hainaut M, Drula E et al. 2019. Comparative genomics of *Rhizophagus*

1136   *irregularis* , *R. cerebriforme* , *R. diaphanus* and *Gigaspora rosea* highlights specific genetic

1137   features in Glomeromycotina. *New Phytologist* 222(3): 1584-1598.

1138

1139 Morris JL, Puttick MN, Clark JW, Edwards D, Kenrick P, Pressel S, Wellman CH, Yang Z,

1140 Schneider H, Donoghue PCJ. 2018. The timescale of early land plant evolution. *Proceedings*

1141 *of the National Academy of Sciences* 115(10): 2274-2283.

1142

1143 Naranjo-Ortiz MA, Gabaldón T. 2019. Fungal evolution: major ecological adaptations and

1144 evolutionary transitions. *Biological Reviews* 94(4): 1443–1476.

1145

1146 Öpik M, Davison J, Moora M, Zobel M. 2014. DNA-based detection and identification of

1147 Glomeromycota: the virtual taxonomy of environmental sequences. *Botany-Botanique* 92:

1148 135–147.

1149

1150 Orchard S, Hilton S, Bending GD, Dickie IA, Standish RJ, Gleeson DB, Jeffery RP, Walker

1151 C, Bass D, Monk J et al. 2017a. Fine endophytes (Glomus tenue) are related to

1152 Mucoromycotina, not Glomeromycota. *New Phytologist* 213(2): 481-486.

1153

1154 Orchard S, Standish RJ, Dickie IA, Renton M, Walker C, Moot D, Ryan MH. 2017b. Fine

1155 root endophytes under scrutiny: a review of the literature on arbuscule-producing fungi

1156 recently suggested to belong to the Mucoromycotina. *Mycorrhiza* 27(7): 619-638.

1157

1158 Orchard S, Standish RJ, Nicol D, Gupta VVSR, Ryan MH. 2016. The response of fine root

1159 endophyte (Glomus tenue) to waterlogging is dependent on host plant species and soil type.

1160 *Plant and Soil* 403: 305-315.

1161

1162 Palos-Fernández R, Aguilar-Pontes MV, Puebla-Planas G, Berger H, Studt-Reinhold L,

1163 Strauss J, Di Pietro A, López-Berges MS. 2024. Copper acquisition is essential for plant

1164 colonization and virulence in a root-infecting vascular wilt fungus. *bioRxiv*. doi:

1165 10.1101/2024.04.23.590778

1166

1167 Parniske, M. 2008. Arbuscular mycorrhiza: the mother of plant root endosymbiosis. *Nature*

1168 *Reviews Microbiology* 6: 763–775.

1169

1170 Peay KG, Kennedy PG, Talbot JM. 2016. Dimensions of biodiversity in the Earth

1171 mycobiome. *Nature Reviews Microbiology* 14: 434–447.

1172

1173   Pellegrin C, Morin E, Martin FM, Veneault-Fourrey C. 2015. Comparative analysis of

1174   secretomes from ectomycorrhizal fungi with an emphasis on small-secreted proteins.

1175   *Frontiers in Microbiology* 6: 1278.

1176

1177   Peng X, Wilken SE, Lankiewicz TS, Gilmore SP, Brown JL, Henske JK, Swift CL, Salamov

1178   A, Barry K, Grigoriev IV et al. 2021. Genomic and functional analyses of fungal and

1179   bacterial consortia that enable lignocellulose breakdown in goat gut microbiomes. *Nature*

1180   *Microbiology* 6: 499–511.

1181

1182   Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal

1183   peptides from transmembrane regions. *Nature Methods* 8(10): 785-6.

1184

1185   Plassard C, Becquer A, Garcia, K. 2019. Phosphorus transport in mycorrhiza: How far are

1186   we?. *Trends in Plant Science* 24: 794–801.

1187

1188   Prout JN, Williams A, Wanke A, Schornack S, Ton J, Field KJ. 2024. Mucoromycotina 'fine

1189   root endophytes': a new molecular model for plant–fungal mutualisms?. *Trends in Plant*

1190   *Science* 29(6): 650-661.

1191

1192   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013.

1193   The SILVA ribosomal RNA gene database project: improved data processing and web-based

1194   tools. *Nucleic Acids Research* 41: 590–596.

1195

1196   Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

1197   features. *Bioinformatics* 26(6): 841-2.

1198

1199   R Core Team. 2023. R: A language and environment for statistical computing. R Foundation

1200   for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

1201

1202   Rambaut, A. 2009. Figtree. Tree figure drawing tool, v1.4.4. Edinburgh, Scotland.

1203   http://tree.bio.ed.ac.uk/software/figtree.

1204

1205   Rasmussen HN, Rasmussen FN. 2014. Seedling mycorrhiza: a discussion of origin and

1206   evolution in Orchidaceae. *Botanical Journal of the Linnean Society* 175(3): 313-327.

1207

1208 Read DJ, Duckett JG, Francis R, Ligron R, Russell A. 2000. Symbiotic fungal associations in

1209 'lower' land plants. *Philosophical Transactions of the Royal Society of London* 355(1398):

1210 815–831.

1211

1212 Rimington WR, Pressel S, Duckett JG, Field KJ, Read DJ, Bidartondo MI. 2018. Ancient

1213 plants with ancient fungi: liverworts associate with early-diverging arbuscular mycorrhizal

1214 fungi. *Proceedings of the Royal Society B: Biological Sciences* 285(1888): 20181600.

1215

1216 Rosling A, Sahraei SE, Khan FK, Desirò A, Bryson AE, Mondo SJ, Grigoriev IV, Bonito G.

1217 Sánchez-García M. 2024. Evolutionary history of arbuscular mycorrhizal fungi and genomic

1218 signatures of obligate symbiosis. *BMC Genomics* 25: 529.

1219

1220 Ryan MH, McCully ME, Huang CX. 2003. Location and quantification of phosphorus and

1221 other elements in fully hydrated, soil-grown arbuscular mycorrhizas: a cryo-analytical

1222 scanning electron microscopy study. *New Phytologist* 160(2):429-441.

1223

1224 Sahu PK, Jayalakshmi K, Tilgam J, Gupta A, Nagaraju Y, Kumar A, Hamid S, Singh HV,

1225 Minkina T et al. 2022. ROS generated from biotic stress: Effects on plants and alleviation by

1226 endophytic microbes. *Frontiers in Plant Science* 13: 1042936.

1227

1228 Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V, Lam

1229 KJK, Tian N, Russum S, Wang J, Medrano-Soto A. 2021. The transporter classification

1230 database (TCDB): 2021 update. *Nucleic Acids Research* 49(1): 461-467.

1231

1232 Schüßler A, Martin H, Cohen D, Fitz M, Wipf D. 2007. Arbuscular mycorrhiza: studies on

1233 the geosiphon symbiosis lead to the characterization of the first glomeromycotan sugar

1234 transporter. *Plant Signaling & Behavior* 2(5), 431–434.

1235

1236 Schüßler A, Schwarzott D, Walker C. 2001. A new fungal phylum, the Glomeromycota:

1237 phylogeny and evolution. *Mycological Research* 105(12): 1413-1421.

1238

1239 Seeliger M, Hilton S, Muscatt G, Walker C, Bass D, Albornoz F, Standish RJ, Gray ND,

1240 Mercy L, Rempelos L et al. 2024. New fungal primers reveal the diversity of

1241 Mucoromycotinian arbuscular mycorrhizal fungi and their response to nitrogen application.
1242 *Environmental Microbiome* 19: 71.

1243

1244 Setubal JC. 2021. Metagenome-assembled genomes: concepts, analogies, and challenges.
1245 *Biophysical Reviews* 13(6): 905–909.

1246

1247 Shen Z, Robert L, Stolpman M, Che Y, Allen KJ, Saffery R, Walsh A, Young A, Eckert J,
1248 Deming C et al. 2023. A genome catalog of the early-life human skin microbiome. *Genome*
1249 *Biology* 24(252).

1250

1251 Shi Z, Deng R, Yuan Q, Mao Z, Wang R, Li H, Liao X, Ma H. 2023. Enzyme commission
1252 number prediction and benchmarking with hierarchical dual-core multitask learning
1253 framework. *Research* 31(6): 0153

1254

1255 Singh NK, Wood JM, Patane J, Moura LMS, Lombardino J, Setubal JC, Venkateswaran K.
1256 2023. Characterization of metagenome-assembled genomes from the International Space
1257 Station. Microbiome 11(1): 125.

1258

1259 Smith AD, Logeman BL, Thiele DJ. 2017. Copper acquisition and utilization in fungi.
1260 *Annual Review of Microbiology* 8(71): 597-623.

1261

1262 Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G, Corradi N,
1263 Grigoriev I, Gryganskyi A et al. 2016. A phylum-level phylogenetic classification of
1264 zygomycete fungi based on genome-scale data. *Mycologia* 108(5): 1028–1046.

1265

1266 Strullu-Derrien C, Selosse M-A, Kenrick P, Martin FM. 2018. The origin and evolution of
1267 mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytologist* 220(4):
1268 1012-1030.

1269

1270 Sujkowska-Rybkowska M, Znojek E. 2018. Localization of calreticulin and calcium ions in
1271 mycorrhizal roots of Medicago truncatula in response to aluminum stress. *Journal of Plant*
1272 *Physiology* 229: 22-31.

1273

1274   Tagirdzhanova G, Saary P, Tingley JP, Díaz-Escandón D, Abbott DW, Finn RD, Spribille T.
1275   2021. Predicted input of uncultured fungal symbionts to a lichen symbiosis from
1276   metagenome-assembled genomes. *Genome Biology and Evolution* 13(4).

1277

1278   Talbot JM, Allison SD, Treseder KK. 2008. Decomposers in disguise: mycorrhizal fungi as
1279   regulators of soil C dynamics in ecosystems under global change. *Functional Ecology* 22(6):
1280   955-963.

1281

1282   Tamura K, Stecher G, Kumar S. 2021. MEGA11: molecular evolutionary genetics analysis
1283   version 11. *Molecular Biology and Evolution* 38(7): 3022–3027.

1284

1285   Tanaka S, Hashimoto K, Kobayashi Y, Yano K, Maeda T, Kameoka H, Ezawa T, Saito K,
1286   Akiyama K, Kawaguchi M. 2022. Asymbiotic mass production of the arbuscular mycorrhizal
1287   fungus Rhizophagus clarus. *Communications Biology* 5: 43.

1288

1289   Tang N, Clemente HS, Roy S, Bécard G, Zhao B, Roux C. 2016. A survey of the gene
1290   repertoire of *Gigaspora rosea* unravels conserved features among Glomeromycota for
1291   obligate biotrophy. *Frontiers in Microbiology* 1(7):233.

1292

1293   Taylor A, Pereira N, Thomas B, Pink DAC, Jones JE, Bending GD. 2015. Growth and
1294   nutritional responses to arbuscular mycorrhizal fungi are dependent on onion genotype and
1295   fungal species. *Biology and Fertility of Soils* 51: 801–813.

1296

1297   Tedersoo L, Magurno F, Alkahtani S, Mikryukov V. 2024. Phylogenetic classification of
1298   arbuscular mycorrhizal fungi: new species and higher-ranking taxa in Glomeromycota and
1299   Mucoromycota (class Endogonomycetes). *MycoKeys* 107: 249-271.

1300

1301   Tedersoo L, Sánchez-Ramírez S, Kõljalg U, Bahram M, Döring M, Schigel D, May T,
1302   Ryberg M, Abarenkov K. 2018. High-level classification of the Fungi and a tool for
1303   evolutionary ecological analyses. *Fungal Diversity* 90: 135–159.

1304

1305   Thippayarugs S, Bansal M, Abbott LK. 1999. Morphology and infectivity of fine endophyte
1306   in a Mediterranean environment. *Mycological Research* 103(11): 1369-1379.

1307

1308 Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, Charron P, Duensing
1309 N, Frei dit Frey N, Gianinazzi-Pearson V et al. 2013. Genome of an arbuscular mycorrhizal
1310 fungus provides insight into the oldest plant symbiosis. *Proceedings of the National Academy*
1311 *of Sciences* 110(50):20117-20122.

1312

1313 Torres-Cruz TJ, Billingsley-Tobias TL, Almatruk M, Hesse CN, Kuske CR, Desirò A,
1314 Niccolò-Benucci GM, Bonito G, Stajich JE, Dunlap C et al. 2017. *Bifiguratus adelaidae*, gen.
1315 et sp. nov., a new member of Mucoromycotina in endophytic and soil-dwelling habitats.
1316 *Mycologia* 222(1): 511-525.

1317

1318 Vierheilig H, Coughlan AP, Wyss U, Piche Y. 1998. Ink and vinegar, a simple staining
1319 technique for arbuscular mycorrhizal fungi. *Applied and Environmental Microbiology*
1320 64(12):5004-5007.

1321

1322 Vierstraete AR, Braeckman BP. 2022. Amplicon_sorter: A tool for reference-free amplicon
1323 sorting based on sequence similarity and for building consensus sequences. *Ecology and*
1324 *Evolution* 12(3): e8603.

1325

1326 Walker C, Gollotte A, Redecker D. 2018. A new genus, Planticonsortium (Mucoromycotina),
1327 and new combination (P. tenue), for the fine root endophyte, *Glomus tenue* (basionym
1328 *Rhizophagus tenuis*). *Mycorrhiza* 12(3): e8603.

1329

1330 Wang S, Zhang X, Zhang Z, Chen Y, Tian Q, Zeng D, Xu M, Wang Y, Dong S, Ma Z et al.
1331 2023. Fusarium-produced vitamin B6 promotes the evasion of soybean resistance by
1332 *Phytophthora sojae*. *Journal of Integrative Plant Biology* 65(9): 2204-2217.

1333

1334 West PT, Probst AJ, Grigoriev IV, Thomas BC, Banfield JF. 2018. Genome-reconstruction
1335 for eukaryotes from complex natural microbial communities. *Genome Research* 28(4):569-
1336 580.

1337

1338 Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualisation of de novo
1339 genome assemblies. *Bioinformatics* 31(20): 3350-3352.

1340

1341   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

1342   ISBN 978-3-319-24277-4

1343

1344   Wurzbacher C, Larsson E, Johan Bengtsson-Palme, Van den Wyngaert S, Svantesson S,

1345   Kristiansson E, Kagami M, Nilsson HR. 2018. Introducing ribosomal tandem repeat

1346   barcoding for fungi. *Molecular Ecology Resources* 19(1):118-127.

1347

1348   Yamamoto K, Degawa Y, Yamada A. 2020. Taxonomic study of Endogonaceae in the

1349   Japanese islands: New species of *Endogone*, *Jimgerdemannia*, and *Vinositunica*, gen. nov..

1350   *Mycologia* 112(2): 309-328.

1351

1352   Yang C, Chowdhury D, Zhang Z, Cheung WK, Lu A, Bian Z, Zhang L. 2021. A review of

1353   computational tools for generating metagenome-assembled genomes from metagenomic

1354   sequencing data. *Computational and Structural Biotechnology Journal* 19: 6301-6314.

1355

1356   Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. 2017. ggtree: an r package for visualization and

1357   annotation of phylogenetic trees with their covariates and other associated data. *Methods in*

1358   *Ecology and Evolution* 8(1): 28-36.

1359

1360   Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018.

1361   dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids*

1362   *Research* 46: 95-101.

1363

1364

1376

1377

Figure 1. Colonisation by arbuscular mycorrhizal fungi (AMF) with fine root endophyte

(FRE) morphology in the roots of *Trifolium subterraneum* from: (A, B, C) the experiment

from which roots were chosen for metagenomic sequencing and (D) roots enriched in FRE

which were used as the reference 18S rRNA sequence (FRE 4583) of Mucoromycotina-AMF

(Orchard et al. (2017); experiment 1). Evident are the characteristic thin hyphae of FRE

morphology, as well as entry points (e), several cells filled by an arbuscule (a) which also

contained lipid droplets (l), terminal vesicles (tv) and intercalary vesicles (icv). (Photo credit:

Payman Abbaszadeh-Dahaji A, B, C; Suzanne Orchard D)

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

Figure 2. Phylogenetic assessment of the Mucoromycotina-AMF reference sequence FRE

4583 and 'consensus_all_seqs_0_16', an 18S rRNA gene sequence associated with the MAG,

and related 18S rRNA sequences amplified from root samples (consensus_all_seqs_0122,

0105 and 0122), in the context of the Tedersoo et al. (2024) *Endogonomycetes* SSU

alignment. The phylogenetic tree was generated using IQ-TREE, the model "TIM2+F+R5",

1000 ultrafast bootstrap replicates, and approximate Likelihood Ratio Tests. Branch support

is presented as a percentage (aLTR support/bootstrap support).

1407

1408

1409

1410

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

Phylum: Ascomycota (79) Basidiomycota (94) Mucoromycota (101) Unknown (1)

Ecology:
Ectomycorrhiza (42) Endophyte (9) Ericoid Mycorrhiza (4) G-AMF (12)
Lichenic symbiont (1) Mixed (23) Orchid Mycorrhiza (3) Parasite (5)
Pathogen (29) Saprotroph (146) Unknown (1)

Protein marker hits:
340 360 380 400 420

BUSCO hits:
50 60 70 80 90 100

1448

1449 Figure 3. Phylogenetic placement of the MAG (Shown here as Candidate FRE) based upon a

1450 set of 434 protein markers identified by the JGI as pan-fungal markers for inferring high-level

1451 fungal phylogeny. Associated ecological group, protein marker hits and BUSCO hits are

1452 shown.

1453

1454

1455

1456

Figure 4. Number of CAZyme, lipase, protease, and transporter predicted proteins per
genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points
represent the phylogenetically closest relatives to the MAG: yellow for 'Endsp1', red for
'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.
Data point shape within the G-AMF distinguishes between the Diversisporales (star) and
Glomerales (triangle). Different letters indicate significant differences between lifestyles as
determined by a Kruskal-Wallis test (p < 0.01) followed by a post-hoc Dunn's test (p < 0.05).

1457

1458

1459

1460

1461

1462

1463

1464

1465

1466

1467

1468

1469

Figure 5. Number of plant cell wall degrading enzyme (PCWDE) and microbial cell wall

degrading enzyme (MCWDE) predicted proteins per genome distributed by ecological

lifestyle. Lifestyle is colour coded. Coloured data points represent the phylogenetically

closest relatives to the MAG: yellow for 'Endsp1', red for 'Jimlac1', green for 'Bifad1', and

purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'. Data point shape within the G-AMF

distinguishes between the Diversisporales (star) and Glomerales (triangle). Different letters

indicate significant differences between lifestyles as determined by a Kruskal-Wallis test (p <

0.01) followed by a post-hoc Dunn's test (p < 0.05).

1478

1479

1480

1481

Figure 6. Cellulose, hemicellulose, lignin and pectin degrading predicted proteins per genome

distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points represent

the phylogenetically closest relatives to the FRE genome: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shape within the G-AMF distinguishes between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test (p < 0.01) followed by a post-hoc Dunn's test (p < 0.05).

1489

1490

none
1491

1492

Figure 7. Chitin, glycan, mannan, and peptidoglycan degrading predicted proteins per
genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points
represent the phylogenetically closest relatives to the MAG: yellow for 'Endsp1', red for
'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.
Data point shape within the G-AMF distinguishes between the Diversisporales (star) and
Glomerales (triangle). Different letters indicate significant differences between lifestyles as
determined by a Kruskal-Wallis test (p = 0.012 for chitin, p = 0.024 for peptidoglycan, and p
< 0.01 for the remainder) followed by a post-hoc Dunn's test (p < 0.05).

1502

1503

1504

1505

Figure 8: Phosphate, calcium, copper, nitrate, ammonium and glycine transporters per

genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points

represent the phylogenetically closest relatives to the MAG: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shape within the G-AMF distinguishes between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test ($p = 0.028$ for phosphate, $p = 0.62$ for copper, $p = 0.041$

for nitrate, and $p < 0.01$ for the remainder) followed by a post-hoc Dunn's test ($p < 0.05$).