Comparative genomic analysis of a metagenome-assembled genome reveals distinctive

symbiotic traits in a Mucoromycotina fine root endophyte arbuscular mycorrhizal

fungus

Joshua Cole[1][*], Sébastien Raguideau[2], Payman Abbaszadeh-Dahaji[3,4], Sally Hilton[1], George

Muscatt[1], Ryan M Mushinski[1], R. Henrik Nilsson[5], Megan H. Ryan[3], Christopher Quince[2,6,7]

and Gary D. Bending[1][*]

[1]School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK

[2]Earlham Institute, Norwich Research Park, Colney Lane, Norwich NR4 7UZ, UK

[3]UWA School of Agriculture and Environment, University of Western Australia, Crawley,

WA, 6009, Australia

[4]Department of Soil Sciences, Faculty of Agricultural Science, Vali-e-Asr University of

Rafsanjan, Rafsanjan, Iran

[5]University of Gothenburg, Department of Biological and Environmental Sciences,

Gothenburg Global Biodiversity Centre, University of Gothenburg, Box 461, 405 30,

Göteborg, Sweden

[6]Gut Microbes and Health, Quadram Institute, Norwich NR4 7UQ, UK

[7]School of Biological Sciences, University of East Anglia, Norwich NR4 7T, UK

*Corresponding authors

ORCID

Joshua Cole - 0009-0004-1563-8469

Sébastien Raguideau - 0000-0001-8727-2604

Payman Abbaszadeh-Dahaji - 0000-0003-3055-4744

Sally Hilton - 0000-0002-3786-1096

George Muscatt - 0000-0002-9218-1210

Ryan M Mushinski - 0000-0003-3572-3500

R. Henrik Nilsson – 0000-0002-8052-0107

Megan H. Ryan – 0000-0003-0749-0199

Christopher Quince – 0000-0003-1884-8440

Gary D. Bending - 0000-0002-9740-5244

Summary

• Recent evidence shows that the arbuscular mycorrhizal (AM) symbiosis is established
by two distinct groups of fungi, with the distinctive 'fine root endophyte' morphotype
formed by fungi from the sub-phylum Mucoromycotina rather than the sub-phylum
Glomeromycotina. While Mucoromycotina AM are abundant and globally distributed,
there is currently no understanding of the genomic basis for the symbiosis or how this
compares to other mycorrhizal symbionts.

• We used culture-independent metagenome sequencing to assemble and characterise
the metagenome-assembled genome (MAG) of a putative fine root endophyte, which
we show belonged to the family Planticonsortiaceae within the order Densosporales.

• The MAG shares key traits with Glomeromycotina fungi, which indicate obligate
biotrophy, including absence of fatty acid and thiamine biosynthesis pathways,
limited enzymatic abilities to degrade plant cell walls, and a high abundance of
calcium transporters, chitin and mannan degrading genes. In contrast to
Glomeromycotina fungi, it exhibited a higher capacity for degradation of microbial
cell walls, a complete cellulose degradation pathway, low abundances of copper,
nitrate and ammonium transporters, and a complete pathway for vitamin B6
biosynthesis.

• These differences highlight the potential for contrasting interactions between
Mucoromycotina and Glomeromycotina AM fungi with their host plant and the
environment which could support niche differentiation and complementary ecological
functions.

56
57

Introduction

Mutualistic symbioses with microbes are exhibited by the overwhelming majority of plant species (Begum, 2019) with mycorrhizas by far the most widespread and well-described association (Hawkins et al., 2023). While mycorrhizal symbioses have evolved multiple times across plant and fungal taxa, they share the common characteristic of the fungus trading nutrients assimilated from the soil, particularly phosphorus and nitrogen, in exchange for carbon from the plant (Luginbuehl & Oldroyd, 2017). Additionally, mycorrhizal associations can confer a range of secondary benefits to the plant host, including disease resistance and stress tolerance (Chen, 2018). Because of these profound impacts on plants, mycorrhizal associations are considered to be major determinants of ecosystem productivity and diversity and to play key roles in directing terrestrial biogeochemical cycling processes. There are four main groups of mycorrhizal associations: arbuscular (AM), ecto- (ECM), ericoid (ERM), and orchid (ORM) (Martin & van der Heijden, 2024) mycorrhizas, which are distinguished by a number of features including the specific plant and fungal taxa involved, the ecosystems they inhabit, the way in which the fungus colonises and grows within root tissues, and the functional traits of the fungi involved (Brundrett & Tedersoo, 2018).

As a function of time and niche specificity, mycorrhizal symbionts appear to have evolved along a saprotrophy to symbiosis continuum (Strullu-Derrien et al., 2018; Martin & van der Heijden, 2024). This directional transition from free-living saprotrophs to obligate symbionts is believed to be driven by the provision of carbohydrates and vitamins from plants in nutrient-poor environments and the subsequent evolutionary success of a symbiotic lifestyle. As a result, key genomic signatures associated with this transition can be identified across this continuum (Martin & van der Heijden, 2024). As fungal symbionts shift away from primarily saprotrophic lifestyles to mutualisms, their genomes typically possess a reduced repertoire of the enzymatic machinery involved in the degradation of soil organic matter (SOM) and plant cell walls (Martino et al., 2018; Gong et al., 2023; Martin & van der Heijden, 2024). The two youngest mycorrhizal symbioses, ERM and ORM, with origins c 75-90 and 112 mya, respectively, (Martino et al., 2018; Brundrett & Tedersoo, 2018) involve symbionts with genomic repertoires capable of simultaneously facilitating a saprotrophic and a symbiotic lifestyle. However, fungi forming the older AM (origin 450 mya, Brundrett & Tedersoo, 2018) and ECM symbioses (origin c 200 mya) possess a much more fine-tuned enzymatic toolkit, noted for a reduced suite of plant cell wall degrading enzymes (PCWDE)

along with the presence of specific genes involved in regulating nutrient transport and host
plant signalling and metabolism (Martino et al., 2018; Martin & van der Heijden, 2024).

AM are considered the most common and widespread plant-microbe symbiosis on Earth
(Read et al., 2000; Parniske, 2008; Spatafora et al., 2017), and their presence in fossil roots of
the earliest land plants suggests that they were a significant factor in the evolution of land
plants. Fungi forming AM have traditionally been placed into the Mucoromycota sub-phylum
Glomeromycotina. These fungi are obligate symbionts; as fatty acid auxotrophs they depend
on the exchange of plant derived fatty acids (Schüßler et al., 2007; Luginbuehl et al., 2017).
As a result of long-term host dependence, arbuscular mycorrhizal fungi (AMF) present
hallmark traits such as limited ability to break down cellulose, biosynthesise thiamine, or
metabolise vitamin B6 derivatives. The genomic inability to express these hallmark traits
represent Missing Glomeromycotan Core Genes (MGCGs) (Tang et al., 2016; Kobayashi et
al., 2018; Rosling et al., 2024).

Until recently, all AMF were believed to belong to the Glomeromycotina (G-AMF)
(Spatafora et al., 2017). It has long been recognised that there are two broad groups of
arbuscular mycorrhiza based on morphology, with the 'coarse endophyte' G-AMF being the
focus of the vast majority of research on the biology and ecology of AMF (Orchard et al.,
2017b). Glomeromycotina comprises four orders, the Glomerales, Diversisporales,
Paraglomerales, and Archaeosporales, which likely diversified when plants evolved onto land
(Schüßle 2001; Chen et al., 2018; Tedersoo et al., 2018). The second morphological group of
'fine root endophytes (FRE)' arbuscular fungi are distinguished from coarse endophytes by
their small spores, fan-shaped or palmate entry points, diverging thin hyphae (<2 µm
diameter) which in intercellular spaces often form hyphal ropes, as well as intercalary and
terminal 'vesicles", approximately 5–10 µm in diameter (Thippayarugs et al. 1999; Orchard et
al., 2017a, b; Walker et al. 2018). Interestingly, the arbuscules of FRE appear very similar in
structure to those of coarse endophytes (Albornoz et al. 2020). Recent evidence has placed
FRE within the sub-phylum Mucoromycotina (M-AMF) (Orchard et al., 2017b), indicating
that AM symbioses may have evolved independently several times, as is the case for
ectomycorrhizal symbioses (Miyauchi et al., 2020).

To date only one species of M-AMF, *Planticonsortium tenue*, has been described (Walker et
al., 2018). However, similarly to G-AMF, root systems are colonised by diverse
Mucoromycotina taxa, communities of which are typically comprised of a small number of
abundant and common taxa and a high diversity of rare low-abundance taxa (Albornoz et al.,

2021, 2022; Seeliger et al., 2024). Furthermore, distinctive morphologies of FRE have been

recognised (Thippayarugs et al., 1999) based on a variety of characteristics, including the

roughness, diameter, and branching of hyphae and the size and shape of vesicles. While fungi

within the Glomeromycotina are all assumed to be plant symbionts, fungi residing within the

Mucoromycotina also include saprotrophs and putative ectomycorrhizal symbionts. Those

Mucoromycotina sequences found associated with roots and rhizoids of vascular and non-

vascular plants appear to have similar phylogenetic ranges and occur within clades alongside

taxa with saprotrophic and putative ectomycorrhizal lifestyles (Desirò et al., 2017; Seeliger et

al., 2024).

M-AMF have been detected across multiple continents, spanning a wide range of ecosystems,

environmental conditions, and hosts, although in contrast to G-AMF they may be missing

from tropical and sub-tropical biomes, and may be favoured by agricultural management

practices (Orchard et al., 2017a; Albornoz et al., 2022). While the benefits of G-AMF to the

host, particularly for phosphorus nutrition, are well documented, the functional significance

of M-AMF within ecosystems is poorly understood (Orchard et al., 2017b; Hoysted et al.,

2019; Albornoz et al., 2020). G-AMF and M-AMF typically co-occur and intermingle within

root systems (Orchard et al., 2017a,b; Jeffery et al., 2018), but they appear to occupy distinct

but overlapping ecological niches (e.g. Orchard et al. 2016), with their distribution at the

landscape scale differentially determined by temperature, pH, and plant richness (Albornoz et

al., 2022). Unlike G-AMF, nutrient availability was found to be an important determinant of

M-AMF abundance, which increased with fertility (Albornoz et al., 2022). Furthermore, in

contrast to G-AMF, experimental evidence suggests that M-AMF may provide the host with

access to soil nitrogen, indicating a potential complementary functionality between the two

arbuscular symbioses and an explanation for the parallel evolution of symbiotic traits

(Hoysted et al., 2022; Howard et al., 2024).

Comparative genome analysis has emerged as a powerful tool to understand microbial

evolution, function, and ecology. However, these approaches largely rely on the availability

of cultured microbes or voucher specimens for whole genome sequencing (e.g. Martino et al.,

2017; Miyauchi et al., 2020; Rosling et al., 2024). Importantly, culturing microbes can prove

time-consuming and problematic as microbial populations have great complexity and many

microbes have specific nutritional requirements which cannot be met in culture. This is

particularly the case for obligate biotrophs such as symbionts which rely wholly or in-part on

the host for their nutritional requirements (Tanaka et al., 2022). As obligate symbionts, G-

AMF require a host plant to complete their lifecycle. Despite this, pure cultures of G-AMF

have been obtained using root organ cultures, and these have been used to produce inoculum

for genome sequencing (Tisserant et al., 2013). The challenges in obtaining and maintaining

root cultures have meant that direct genome sequencing of G-AMF cultures has been limited

to date. Recently, asymbiotic mycelium and spore production of the G-AMF *Rhizophagus*

clarus has been achieved by supplementing its growth medium with palmitoleic acid, and this

could provide a boost to rates of culture-based sequencing (Tanaka et al., 2022). An

alternative approach to obtain fungal genomes is sequencing of nuclei extracted from spores

(Montoliu-Nerin et al., 2021), although most G-AMF taxa appear to sporulate infrequently, if

at all (Öpik et al., 2014).

M-AMF are also putatively obligate symbionts, but because they have only recently been

recognised as a distinct group of symbionts, culturing efforts lag behind those of G-AMF.

Recently, monoxenic cultures of a Mucoromycotina fungus associated with the clubmoss

Lycopodiella inundata have been obtained and shown to form FRE colonisation in a vascular

plant (Hoystead et al., 2022). However, while genome sequencing and comparative genomic

analysis of a number of saprotrophic and putative ectomycorrhizal Mucoromycotina has been

achieved (Chang et al., 2019; Rosling et al., 2024), there is no understanding of the genome

of M-AMF fungi and how their functional repertoire and mutualistic signatures compare to

those of G-AMF, other mycorrhizal symbioses, and saprotrophic and putatively

ectomycorrhizal Mucoromycotina. To further complicate the matter, initial comparative

analysis of the sister clades indicate that some widely considered hallmark traits of

mycorrhizal symbioses may instead represent ancestral markers as opposed to adaptation

(Rosling et al., 2024).

Recently there has been great progress in the development of computational tools for the

culture-independent assembly of 'dark' microbial genomes from metagenome sequencing

datasets (Benoit et al., 2024). Such 'Metagenome Assembled Genomes' (MAGs) have been

widely used to identify the metabolic capabilities and ecological roles of specific prokaryotes

in diverse habitats (Yang et al., 2021). MAG construction of eukaryotic microbes is more

challenging because of a range of issues including their large sizes, high genome

heterozygosity, low abundance in metagenome datasets, and a lack of well annotated

reference genomes for gene annotation (Peng et al., 2021; Setubal 2021). As a result, few

studies have constructed eukaryotic MAGs from metagenomes, and these have typically been

performed in low-diversity communities (e.g. West et al. 2018; Peng et al., 2021). A further

particular issue with analysis of plant symbionts *in situ* is their intimate association with plant
tissue. This results in co-extraction of host DNA, increasing the sequencing depth required to
assemble symbiont genomes.

The aim of the current work was to use culture-independent metagenome sequencing to
assemble and characterise the genome of an FRE-forming M-AMF fungus for the first time
and compare its genomic repertoire to glomeromycotan fungi and other fungal symbionts.
Roots of *Trifolium subterraneum* were first enriched with FRE using a selective soil sieving
technique. Root samples were used for metagenome sequencing, from which a MAG of the
FRE was constructed. We compiled a database of 274 fungal genomes across the
Ascomycota, Basidiomycota and Mucoromycota, and assigned these to G-AMF, ECM, ERM,
OCM, saprotroph, pathogen, parasite, endophyte, lichenic, and mixed lifestyles. Comparative
analysis showed that the MAG shares key distinctive characteristics with G-AMF. This
includes overlaps in some MCGCs, presence of abundant calcium transporters, and an overall
reduced set of plant cell wall degrading enzymes. However, we found a relatively high
abundance of microbial cell wall degrading enzymes and evidence of complete cellulose
degradation capability, as well as low numbers of nitrogen transporters, suggesting a
distinctive niche relative to G-AMF and the other mycorrhizal lifestyles.

Materials and Methods

Soil collection and sample preparation

For the metagenome sequencing, plants with the root AMF community enriched in M-AMF
were produced following the methods of Orchard et al. (2017a). In brief, soil was collected
from the field studied by Orchard et al. (2016) and Orchard et al. (2017a) in the Peel Harvey
region in Western Australia, Australia: site coordinates were 32° 48 826'S, 115° 53346'E and
the site was ~8 m above sea level. The field contained winter-active annual grasses and
legumes and supported a low-intensity free-range beef cattle grazing enterprise. Soil
collection targeted a sandy loam soil with a pH in $CaCl_2$ of 5.4. Soil was wet-sieved through a
stack of four sieves with pore sizes ranging from 2 mm down to 20 µm. The material
collected on each sieve was retained and added to pasteurised, dried, washed coarse river
sand at a ratio of either 1:27 or 1:81 w/w. Free-draining pots were filled with 1 kg of each
treatment (n=4) and placed in a temperature-controlled glasshouse. Four plants of *T.*
subterraneum L. with appropriate rhizobia added were grown in each pot which were

maintained at 80% field capacity by watering to weight, and received phosphorus-free, but

otherwise complete, nutrient solution once a week. After 8 weeks, plants were harvested and

roots gently washed clean. The four root systems from each pot were allocated to one of two

future activities: staining for assessment of AMF colonisation (stored in 70% ethanol) or

DNA extraction (frozen at -20ºC). Roots were cleared and stained using the ink-vinegar

method of Vierheilig et al. (1998) and assessed using a modified version of the magnified

intercept method described by McGonigle et al. (1990) where the root at each grid-line root

intercept was recorded as either free of all colonisation, colonised by G-AMF only, colonised

by M-AMF only, or colonised by both M-AMF and G-AMF.

233   DNA extraction, 18S rRNA gene amplicon sequencing and metagenome sequencing

Twelve samples where a high proportion of total AMF colonisation consisted of M-AMF

(Table S1) were selected for 18S rRNA amplicon and metagenome sequencing based on the

proportion of root length colonised by FRE and the quality and quantity of DNA obtained

(Table S1). DNA was extracted from root samples using the DNeasy PowerSoilPro Kit

(Qiagen) following the manufacturer's protocol. DNA concentrations were measured using a

Qubit 3.0 fluorimeter using the Qubit dsDNA high sensitivity assay kit (Invitrogen)

according to the manufacturer's protocol.

To determine whether the dilution technique had enriched M-AMF we preformed community

18S rRNA amplicon analysis. Glomeromycotina and Mucoromycotina 18S rRNA genes in

root samples were amplified using AM-Sal-F and AMDGR primers which amplify a 220-

base fragment (Seeliger et al., 2024). PCR reactions, Illumina MiSeq sequencing, and

bioinformatic analysis were performed using the approaches outlined in Seeliger et al. (2024).

Long read nanopore amplicon sequencing of the nuclear ribosomal 18S rRNA, 5.8S, and

28S/LSU region of Mucoromycotina was performed on root samples to support construction

of the FRE MAG and to determine its phylogenetic relationships. PCR reactions were

conducted using AM-Sal-F in combination with RCA95m (Wurzbacher et al., 2018) using

the recommended Oxford Nanopore protocol, amplifying a 4.5 kb region. Briefly, an initial

PCR was performed with adapter-tagged oligos, and the PCR products were cleaned using

Ampure SPRI beads. Barcodes were added using a second PCR with an initial denaturation at

95ºC for 3 minutes followed by 30 cycles of denaturation at 98ºC for 15 seconds, 60ºC

annealing for 15 seconds, and 65ºC extension for 150 seconds, with a final extension at 65ºC

for 6 minutes. Following a clean-up with SPRI beads, the samples were used for sequencing on a SpotON Nanopore flow cell following the manufacturer's instructions.

Fast5 files were base called and demultiplexed using the Guppy basecaller provided by Oxford Nanopore Technologies. The sample was then processed using amplicon_sorter v2022-03-28 (Vierstraete & Braeckman, 2022) which aims to group amplicons at species or genus level (~95%), output consensus sequences, and quantify the copy number of the representative consensus sequences in the sample. This resulted in 42 consensus sequences, 5 of which were placed into the Mucoromycotina.

To aid detection of M-AMF sequences in the metagenome, a reference 18S rRNA sequence was obtained from DNA extracted from *T. subterraneum* root samples which were enriched in M-AMF and contained minimal G-AMF. These samples were produced in a similar manner to that described above and are presented in Orchard et al. (2017a) as "Experiment 1". On average, 90% of the total AMF colonisation in these roots was contributed by M-AMF, presenting the first evidence linking FRE morphology to Mucoromycotina. The AM-Sal-F and RCA95m primers were used to amplify a 4.5 kb region, as described above, which was cloned using the TA coning kit (ThermoFisher). Partial 18S rRNA analysis using AM-Sal-F and AMDGR primers revealed a clone, FRE4583, which showed a 100 % match to both OTU4 which was associated with FRE morphology in the earlier Orchard et al. (2017) study and the BEG249 *Planticonsortium tenue* FRE culture described by Walker et al (2018) (Albornoz et al., 2022). Furthermore, this and closely related sequences have been detected in multiple studies reporting FRE morphology in plant roots (Albornoz et al 2020, Seeliger et al., 2024). Primer walking was used to sequence overlapping regions of clone FRE4583 until the full sequence was covered, generating a 4.5 kb fragment.

Metagenome sequencing and assembly

The twelve root samples were used for Shotgun metagenomic sequencing using the HiSeq 150 paired end protocol to generate between 15 and 47 Gbp sequencing depth. Library construction and sequencing was conducted using Illumina protocols by Novogene UK Ltd.

Samples were first filtered for *T. subterraneum* DNA using bwa mem v0.7.18-r1243-dirty (Li, 2013) and samtools v1.10 (Danecek et al., 2021). To determine FRE depth and breadth of coverage the filtered metagenome reads were mapped, again using bwa mem and samtools, against the FRE4583 ribosomal operon and the resulting alignment files processed with

custom Python scripts. To be sure that a high mean coverage was not an artefact of conserved

regions, breadth of coverage was visually inspected. Sample P72 was predicted to have the

highest Mucoromycotina abundance, possessing the majority of the operon with a coverage

depth of over 100 (Fig. S1); the missing regions are likely due to repeats as we filtered

ambiguous mappings. We therefore focussed on this sample in our efforts to reconstruct an M-

AMF genome, assembling it using the SPAdes assembler v3.14.1 (Bankevich et al., 2012).

This produced a catalogue of contigs with an N50 of 446 and total length of 1.3 Gbp. We then

computed coverage depth profiles for these contigs across all samples by mapping reads with

bwa mem, samtools, and bedtools v2.30.0 (Quinlan & Hall., 2010). This improves the

discriminatory power of genome binning through multi-sample coverages whilst keeping the

best quality single sample assemblies. The contigs were then binned using these coverage

profiles and composition with the binning software CONCOCT v1.1.0 (Alneberg et al., 2014)

and a minimum contig length of 1000bp.

The putative M-AMF genome was identified based on a consensus of multiple lines of

evidence. As a first step, our reference M-AMF ribosomal sequence FRE4583 was mapped to

the assembly. A total of 12 unitigs were found with high similarity (>98%) and spanning 75%

of the operon without overlap, of which only four had a context in the assembly graph. Using

custom scripts, co-localisation of contigs binned by CONCOCT within the direct vicinity of

the aforementioned 4 untigs were visualised directly on the SPAdes assembly graph (Fig. S2)

using Bandage v0.8.1 (Wick et al., 2015). This figure encompasses all paths of 10 Kbp

starting from these 4 unitigs (red) but correspond to only 0.05% of the whole assembly,

showing a clear proximity of multiple numbered bins. Ignoring any miss-assemblies, all

genomes have a distinct path in the assembly graph and in particular the M-AMF genome

should include a path which passes through those unitigs labelled as ribosomal sequences

allowing us to identify the M-AMF, even though we could not assemble a complete FRE4583

operon in a single contig, likely due to strain diversity.

A total of 12 bins were found in close proximity to M-AMF unitigs. However, on assessing

quality, only bin 17 was found to have any marker genes, giving it a completion of 83.7%

using the BUSCO (Manni et al., 2021) general fungi_odb10 scheme with a size of 37.37

MBp (Table S2). In particular, while contigs from bin 70 were directly co-located with FRE

ribosomal sequences, this bin did not possess any marker genes and was distinctly different in

terms of coverage from bin 17. It may be that bin 70's location in the graph is due to a miss-

assembly, or it could be that this is due to strain diversity and that bin 70 is strain-specific

part of the pan-genome of the FRE which would result in it having a different coverage

profile to the core genome represented by bin 17. To further confirm that bin 17 is related to

M-AMF, we used phyloFlash v3.4.2 (Gruber-Vodicka et al., 2020) to quantify ribosomal

operon abundance from the 12 Illumina short read samples. It is indeed possible to use

phyloFlash to derive abundance, not only of the SSU rRNAs of the SILVA database (Quast et

al., 2013) but also of the 42 nanopore amplicon 18S rRNA consensus sequences. Filtering

any taxa with less than 10 reads mapping across all samples yielded a total of 251 entries

including the 42 consensus sequences (Fig. S3). These were normalised by the total

sequencing depth of each sample. To compare to these, the coverage depth of bin 17 was

calculated for each sample by taking the sum of all nucleotides mapping to all contigs of bin

17 and dividing by the total length of the contigs, followed by normalisation as above. Using

Pearson's correlation, the coverage of bin 17 was compared to the coverage of all 251 taxa

detected with phyloFlash. The strongest correlation was with one of the FRE consensus

sequences (consensus_all_seqs_0_16). This was found to be correlated with r = 0.97 and an

adjusted p value of 4.5e-11 (Fig. S4). The regression coefficient of 63 between coverage

profiles (consensus_all_seqs_0_16 and bin 17) would imply that there are that many

ribosomal operons in the genome, which is consistent with other fungi.

338 Reference genome access and ecological lifestyle annotation

To compare the MAG to known fungi, a database of reference fungal genomes was

constructed from JGI MycoCosm (Table S3). The fungal species, representing the three

phyla—Ascomycota, Basidiomycota, and Mucoromycota —were chosen based on two

criteria. First, fungal genomes which had been included in previous comparative genomics

analyses of mycorrhizal fungi (Martino et al., 2018; Miyauchi et al., 2020) were included

since the ecology of these fungi had already been curated. Secondly, the database was

supplemented with Mucoromycota genomes available in MycoCosm

(https://mycocosm.jgi.doe.gov/mucoromycota/mucoromycota.info.html) in May 2023. The

associated predicted fungal proteomes were accessed and downloaded in May 2023. We

acknowledge that this was not an exhaustive list of fungal genomes accessible on JGI

MycoCosm in that there were genomes representing further species within Ascomycota and

Basidiomycota that were not included in the analyses. Additionally, with our primary focus

being the provision of comparable context for the MAG, we aimed to obtain a broadly

representative database and not an all-encompassing list.

Fungal genomes were categorised according to their ecological lifestyle (Table S3). To
achieve this, fungal species were coded as follows. First, ecological classifications were
accessed from functional groups recorded on JGI MycoCosm (e.g.,
https://mycocosm.jgi.doe.gov/Ectomycorrhizal_fungi/ Ectomycorrhizal_fungi.info.html).
Secondly, for genomes which were included in previous comparative genomics analyses, the
classifications reported in those studies were used (Chang et al., 2019, 2022; Martino et al.,
2018; Miyauchi et al., 2020). Any remaining genomes without classifications were then
manually curated, as informed by published evidence. Similarly to previous studies, we
allowed a fungal species to occupy multiple ecological lifestyles when there was evidence of
this. We assigned 146 genomes as saprotrophic, 42 as ECM, 29 as pathogenic, 23 as mixed
lifestyle, 12 as G-AMF, 9 as endophytic, 5 as parasitic, 4 as ERM, 3 as OCM, and 1 as
lichenic. Within this total we included the following five Mucoromycotina genomes.
Endogone sp. 'Endsp1' is a putative saprotroph found on fallen oak logs in Florida, USA
(Chang et al., 2019). *Jimgerdemannia lactiflua,* 'Jimlac1' is a putative ectomycorrhizal
fungus isolated from under *Pseudotsuga menziesii* in Oregon, USA (Chang et al., 2019). Two
genomes of the putative ectomycorrhizal fungus *Jimgerdemannia flammicorona*: isolate
'Jimfl_GMNB39_1' from under *Picea abies* and *Picea pungens* in Michigan, USA (Chang et
al., 2019), and isolate 'Jimfl_AD_1' from under *Pinus strobus* in Biella, Italy (Chang et al.,
2019). *Bifiguratus adelaidae* 'Bifad1', a soil-dwelling putative saprotroph isolated from a
Pinus taeda plantation in North Carolina, USA (Torres-Cruz et al., 2017).

Phylogenetic assessment

Phylogenetic assessment of FRE4583 and those consensus sequences from root samples
which were assigned to the order Endogonales (including all_seqs_0_16) was investigated
using 18S rRNA gene alignment. These sequences were added to the Endogonomycetes 18S
rRNA alignment from Tedersoo et al. (2024) using MAFFT v7.525 (Katoh and Standley,
2013). The resulting alignment was manually checked and refined in MEGA11 v11.0.13
(Tamura et al., 2021). The best fitting model for our alignment was selected using IQ-TREE
v2.3.6 and the option "-MFP" (Kalyaanamoorthy et al., 2017; Minh et al., 2020). A
phylogenetic tree was generated using the model "TIM2+F+R5" with 1000 approximate
likelihood ratio tests and 1000 ultrafast bootstraps (Hoang et al., 2018; Minh et al., 2020).
The tree was subsequently visualised in FigTree v1.4.4 (Rambaut, 2009).

In addition to the 18S rRNA alignment, fungal phylogeny was inferred from genome

sequences using a concatenated protein marker alignment. A set of 434 protein markers

identified by the JGI as pan-fungal markers for inferring high-level fungal phylogeny was

used to compare the phylogenetic placement of the MAG and all the reference fungal

genomes within our curated database. To identify these marker proteins, an HMM set was

downloaded from https://github.com/1KFG/Phylogenomics_HMMs/tree/master/

HMM/JGI_1086. Phylogenomic markers were extracted from fungal proteomes using the

above HMM, and a subsequent phylogenetic tree was built using PHYling v0.9.0 (available

at: https://github.com/stajichlab/PHYling) under default settings. The tree was then visualised

in R using ggtree v3.6.2 (Yu et al., 2017).

Annotation of MAG and reference genomes

Our comparative analyses focused on Carbohydrate-Active enZymes (CAZymes), lipases,

proteases, transporters, and small-secreted proteins. To identify CAZymes, fungal proteomes

were queried against an HMM profile database built from CAZy sequences downloaded from

the dbCAN2 database (accessed from:

https://bcb.unl.edu/dbCAN2/download/Databases/V11/dbCAN-HMMdb-V11.txt) (Zhang et

al., 2018) using HMMer v 3.3.2 (Eddy, 2011). Protein hits were subsequently filtered with an

e-value threshold of 1e-15 and an alignment threshold of 50% of the domain query. To

identify proteases, fungal proteomes were queried against the MEROPS peptide database

(accessed in July 2023 from:

https://ftp.ebi.ac.uk/pub/databases/merops/current_release/merops_scan.lib) using BLASTp.

Protein hits were subsequently filtered with an e-value threshold of 1e-10. To identify lipases,

fungal proteomes were queried against the Lipase Engineering Database v4.1.0 using

DIAMOND v2.1.8 (Buchfink et al., 2021) and the option "-sensitive". Protein hits were

subsequently filtered with an e-value threshold of 1e-5. To identify transporters, fungal

proteomes were queried against the TCDB transporter database (accessed in November 2023)

(Saier et al., 2021) queried using BLASTp. Protein hits were subsequently filtered with an e-

value threshold of 1e-10. Secreted proteins were identified using a custom pipeline as

reported previously (Pellegrin et al., 2015). Briefly, signal peptides were identified with

SignalP v4.1 (Petersen et al., 2011), extracellular localisation was predicted with WoLFPSort

v0.2 (Horton et al., 2007), transmembrane helices were identified with TMHMM v2.0

(Möller et al., 2001), secretory pathway association was inferred with TargetP v2.0 (Almagro

Armenteros et al., 2019), and the absence of a KDEL motif in the C-terminal region was

confirmed with PS-SCAN revision 1.86 (De Castro et al., 2006). Small-secreted proteins

were identified as secreted proteins < 300 amino acids with no CAZyme, lipase, protease, or

transporter annotation. Functional annotation of putative small-secreted proteins was

achieved with PfamScan v1.6 (Mistry et al., 2007), an e-value threshold of 1e-15, and an

alignment threshold of 50% of the domain query. For each functional protein category being

identified, a single best hit per locus was retained by selecting the hit with the smallest e-

value and greatest alignment of the domain query.

Comparative genomics analyses

428   All comparative genomics analyses were conducted using R v4.2.2 (R Core Team, 2021), and

429   all visualisations were generated with ggplot2 v3.4.0 (Wickham, 2016). Functional

430   annotation of EC numbers was assigned using DeepECtransformer v1.0 (Kim et al., 2023)

431   and the ECRECer web platform (Shi et al., 2023).

Results

Description of FRE in the roots

The appearance of the FRE in the roots of the twelve samples chosen for metagenomic

sequencing was consistent with the description of the genus *Planticonsortium* by Walker et

al. (2018) into which they placed *Glomus tenue* (previously generically referred to in the

literature as fine endophyte or fine root endophyte; Orchard et al. 2017b) (see photographs in:

Figure 1 herein; Figure 2 and Figure S1 in Orchard et al., 2017a and Thippayarugs et al.

(1999)). In particular, entry points were characterised by structures with "feather-like, fan-

shaped, palmate or multilobed with finger-like projections" (Walker et al. 2018) which then

led to fine, diverging hyphae in cortex cells and, often, plentiful finely-branched arbuscules.

Small intercalary vesicles and terminal vesicles were also common. Droplets of lipids were

often evident within hyphae and arbuscules. In some instance, colonisation in the cortex was

too dense to clearly distinguish the structure of individual arbuscules.

In the samples used for amplicon and metagenome sequencing the percentage of total root

length colonised AM ranging between 26.8 and 72.5% for AMF with FRE morphology and 0

to 16.3% for G-AMF, producing a proportion of AMF colonisation by FRE between 62.2 and

100%. 18S rRNA Miseq sequencing of Glomeromycotina and Mucoromycotina communities

in root samples revealed that the proportion of Endogonales amplicons ranged between 25

and 82% (Table S1).

Phylogeny and Features of Fungal Genomes

In this study we constructed a putative FRE genome, referred to as the 'MAG'. This MAG

was identified as a putative M-AMF through two lines of evidence. First, the contig bin was

colocated on the assembly graph with unitigs that we could assign to the 18S rRNA gene of a

known M-AMF reference (FRE4583). Secondly, its abundance correlated across the 12

samples in the study with an 18S rRNA sequence 'consensus_all_seqs_0_16', which we

showed to be very closely related to FRE4583 (Figure 2). The bin itself did not contain a

complete 18S rRNA sequence. This is unsurprising as this gene is well known for failing to

assemble, largely due to it being comprised of strongly conserved regions that repeat

throughout the genome, failing to assemble into substantial contigs. The MAG measured

37.37 Mbp with a genome coverage varying between 1x and 568 depending on sample. The

N50 was found to be 4928 with a GC content of 41.1%. Finally, its completion was assessed

to be 83.7% with only 1.5% contamination (Table S2). We identified a total of 10872

proteins. Of these proteins, we were able to assign 3358 proteins with a functional annotation,

comprising 218 CAZymes, 121 lipases, 280 proteases, 13 small secreted proteins, and 2726

transporters.

While we acknowledge that the revised phylogeny of the Mucoromycotina subphylum

produced in Tedersoo et al. (2024) lacks type specimens and is grounded in part on dark taxa,

as a poorly captured branch of the fungal tree of life we deem this an appropriate basis to

apply the phylogeny of our MAG. 18S rRNA analysis placed consensus_all_seqs_0_16, and

by proxy the MAG, within the order Densosporales of the Mucoromycotina, corresponding

with the Densosporaceae sensu Desirò et al. (2017) (Figure 2). This order is associated with

FRE morphology (Orchard et al., 2017a; Chang et al., 2019). Further to this, strong support

values allow us to confidently place the MAG in the newly proposed family

Planticonsortiaceae (Tedersoo et al., 2024). Within this family, the MAG was placed with

very high similarity to the known M-AMF reference sequence, FRE4583 (Figure 2). Further

to this, the MAG was closely related with three other Mucoromycotina 18S rRNA sequences

obtained from the root samples (consensus_all_seqs 0105, 0122 and 0260) and to

environmental sequences obtained from the thalli of early diverging plant sequences, e.g.,

MH174585 and MH174471. The wider *Planticonsortiaceae* clade included further sequences

from the thalli of non-vascular plants, sequences not annotated beyond the phylum level

associated with soil from across Europe (EUK sequences), and one sequence associated with

the rhizosphere of *Populus tremuloides* (EF23703), a known arbuscular mycorrhizal host

(Lee et al., 2020). Phylogenetic analysis therefore placed the MAG apart from other members

of the Mucoromycotina with putative ecology from the order Endogonales, including the

putative ectomycorrhizal Mucoromycotina member *Jimgerdemannia* spp. (Desirò et al.,

2017) and the saprotrophic *Endogone* sp. and *Vinositunica* sp. which have a putative

saprotrophic or ectomycorrhizal lifestyle (Yamamoto et al., 2020). Similarly, the protein

marker alignment placed the MAG within the Mucoromycota (Figure 3). The closest relatives

are within the order Endogonales, including *Endogone* sp. and *Jimgerdamannia* spp, since no

Densosporales genomes are available to date. The MAG forms a sister clade relative to

Endogone and *Jimgerdamannia*, suggesting it diverged early from the other

Mucoromycotina.

Genomic Profile of the MAG

To contextualise the gene content of the MAG, we compared the CAZyme, protease, lipase,

and transporter profile to genomes within the reference dataset that represented the

saprotrophy-symbiont continuum. We also compared the MAG with the five available

genomes from the Endogonales, which the 18S rRNA and genome analysis confirmed as the

closest phylogenetic relatives to the MAG. The ecological lifestyle had a significant effect on

genome content across all four categories ($p < 0.01$) (Figure 4). For the CAZymes, the

ecological lifestyles fell into two groups—G-AMF and ECM, ERM and OCM —with SAP

bridging the two (Figure 4a). The MAG contained a low number of CAZymes (218), which

was similar to the mean of the ECM (240), but slightly higher than that of the G-AMF (175).

A similar number of CAZymes were found in the MAG compared to related Endogonales

genomes (149 to 208). The ERM genomes contained by far the largest median number of

CAZymes across the lifestyles (629). Only the ERM possessed a significantly ($p < 0.01$)

larger number of lipases than the other ecological lifestyles (Figure 4b). The MAG possessed

121 lipases, lower than the means of the G-AMF (189), ECM (238), and SAP (263), OCM

(268). Similarly, the related Endogonales genomes all contained low numbers of lipases (72

to 132). The ERM genomes had significantly ($p < 0.01$) higher mean numbers of proteases

per genome (457) than the other lifestyles (Figure 4c). The MAG had similar numbers of

proteases (280) compared to G-AMF (305), ECM (286), SAP (271), and to the related

Endogonales genomes (229 to 298). The G-AMF and ERM lifestyles had significantly higher

($p < 0.05$) numbers of transporters per genome relative to the other genomes. The MAG

(2726) had a substantially lower number of transporters compared to G-AMF (3900);

however, it contained more than related Endogonales genomes (1932 to 2388) that sat in the

lower end of the dataset (Figure 4d). Across the dataset, the saprotrophs contained an

especially high degree of intra-group variability in numbers of all four profiles compared to

the other ecological lifestyles (Figure 4). The number of CAZymes, lipases, proteases, and

transporters in the MAG were also comparable to the endophyte, parasite pathogen, and

mixed lifestyles (Fig S5). The MAG contained 13 small secreted proteins, a middling amount

for the dataset (Fig S5).

CAZyme Profile

CAZyme proteins associated with the degradation of plant and microbial cell walls were

identified to investigate the saprotrophic repertoire of the MAG, in the context of the

mycorrhizal saprotrophy-symbiont continuum. The ERM (252) and OCM (215) genomes

possessed a significantly higher ($p < 0.01$) mean number of PCWDEs than the G-AMF (32),

ECM (59), and SAP (77) genomes (Figure 5a). The MAG contained a similar number of

PCWDE (37) as the G-AMF genomes and the genomes of related Endogonales (33 to 44),

which was notably lower than the endophyte, parasite, pathogen, and mixed lifestyles (Fig

S6). ERM genomes contained a significantly ($p < 0.01$) higher mean number of MCWDE

(109) than the saprotrophs and other mycorrhizal lifestyles (24 to 59) (Figure 5b). The G-

AMF genomes had a significantly ($p < 0.05$) lower number of MWCDE (24) compared to the

mean ECM (45), OCM (59), and SAP (48) genomes. The MAG had 53 MCWDE, higher

than the median number of all other lifestyles except the ERM (Figure 5b, Fig S6), and

similarly higher than numbers in the parasite, pathogen and mixed lifestyles, but similar to

numbers in endophytes (Fig S6). A similar number of MCWDE was detected across the

Endogonales genomes (36 to 53).

Patterns of PCWDE distribution across the mycorrhizal associations were similar (Figure 6).

ERM and OCM genomes had significantly ($p < 0.01$) higher mean numbers of cellulases,

hemicellulases, and pectinases than the G-AMF and ECM genomes. Relative to the mean G-

AMF, the MAG had notably higher numbers of cellulases (18 compared to 8), but similar

numbers of hemicellulases (11 compared to 7) and pectinases (4 compared 2). The

Endogonales genomes had similar numbers of cellulases (11 to 13), hemicellulases (10 to 16), and pectinases (4 to 12) as the MAG. The numbers of lignin degrading enzymes were significantly higher in the ERM (31, $p < 0.01$) relative to the other mycorrhizal lifestyles, which all possessed similar numbers: ECM (13), G-AMF (15), OCM (9), and SAP (12). Notably the MAG had a very low number of lignin degrading enzymes (4) compared to all other lifestyles, this trend was consistent across the other Endogonales genomes (2 to 8). Numbers of cellulose, hemicellulose, and pectin degrading enzymes in the MAG were also typically lower than the mean in endophyte genomes (Fig S7), and numbers of lignin degrading genes were similarly notably lower than the endophyte, parasite, pathogen and mixed lifestyles (Fig S7).

Cellulases in the MAG genome comprised 8 families, with 6 copies of AA9 being the most abundant (Table S4). The other Endogonales genomes contained an average of 6 cellulase families, with GH9 being the most abundant in all four genomes, of which the MAG contained 3. However, not a single copy of AA9 was identified within the other Endogonales genomes (Table S4). The G-AMF genomes only contained an average of 3 families, with the Glomerales containing a mean of just 6 cellulases. However, the Diversisporales contained a much higher number of cellulases (17 and 12), including 7 and 6 of AA9 and 6 and 3 copies of AA3, respectively (Table S4). The MAG contained a full suite of cellulose degradation enzymes: Endoglucanase, β-glucosidase, and exoglucanase (Table S5). However, with the absence of β-glucosidases in all G-AMF genomes and only two containing a single copy of exoglucanase, none of the G-AMF genomes possessed the full cellulose degradation pathway (Table S5). Hemicellulases in the MAG comprised the majority of the remaining PCWDEs, with 7 families and 5 copies of GH31 being the most abundant (Table S4). Similarly, the other Endogonales genomes contained an average 8 families, with GH31 being the most abundant. The G-AMF genomes contained an average 3 families, with GH31 and GH5_7 being the most abundant (Table S4). No copies of GH29 or GH95 were found within the G-AMF genomes. Similarly, no copies of GH95 were found within the MAG. However, a single copy of GH29 was present (Table S4).

Pectinases in the MAG were limited, with only 4 copies across 3 families: 2 GH35s, 1 GH2, and 1 PL1_7 (Table S4). The other Endogonales genomes contained an average of 4 families, with the most abundant being GH28 and GH35 (Table S4). Within the G-AMF genomes, no evidence for pectinases was found within 'Gigmar1'. Apart from that, all other genomes contained only 2 or 3 copies of GH35 (Table S4).

The MAG contained 3 families of lignin degrading enzyme: 2 AA1, 1 AA1_2, and 1 AA2

(Table S4). This trend was conserved across the Endogonales genomes, in which lignin

degrading enzymes comprised of 3 families, with AA1 being slightly more abundant (Table

S4). The G-AMF genomes, while only comprising an average 3 families, contained a much

larger number of lignin degrading enzymes. The Diversisporales contained 37 and 25 lignin

degrading enzymes, including 35 and 23 copies of AA1 respectively. The Glomerales

contained an average of 12 lignin degrading enzymes per genome, with an average of 9

counts being attributed to AA1 (Table S4).

The G-AMF genomes had a smaller mean chitinase repertoire (16) than ERM (35, $p < 0.01$),

OCM (25, $p < 0.05$), and SAP (26, $p < 0.01$) (Figure 7a). However, the MAG had one of the

highest numbers of chitinases across the genome database (38), roughly double the mean

number found in both the G-AMF and ECM (21), and notably higher than the mean found in

endophytes, parasite, pathogen, and mixed lifestyles (Fig S8). However, there was

considerable variation in the numbers of chitinases across the G-AMF genomes, with the two

Diversisporales genomes containing 38 and 29 counts respectively, more comparable to the

MAG. The Endogonales genomes similarly contained high mean numbers of chitinases (29 to

42) compared to other genomes in the dataset. Over half of the MAG's chitinase repertoire

was attributed to just two families, CE4 and GH18, with 16 and 13 copies respectively, a

trend also seen across the other Endogonales genomes (Table S4). The G-AMF genomes

largely comprised of CE4 chitinases, however, the Diversisporales genomes also contained a

large number of GH18 (13 and 10), very few of which (0 to 2) were found in Glomerales

(Table S4). The MAG contained a fully comprehensive set of chitin degrading enzymes

(Table S5). Similarly, all but one of the G-AMF genomes contained a full chitin degradation

suite (Table S5).

G-AMF genomes contained significantly ($p < 0.01$) lower mean numbers of glycan degrading

genes (1) compared to the mean of the saprotroph (14) and other mycorrhizal lifestyles (15 to

42), and in fact were only detected in the two Diversisporales genomes (Figure 7b). In

contrast, the MAG contained 3 glycanases, and this was consistent with a low number of

glycanases in the Endogonales genomes (2 to 8). The MAG only contained 2 copies of GH17

and 1 copy of GH5_9 (Table S4). The only two other genomes to contain single copies of

these were 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'. All the other Endogonales genomes

contained 1 or 2 copies of both GH72 and GH81 (Table S4). The Diversisporales genomes

only contained low counts of GH5_9 and GH81 (Table S4).

ERM genomes contained a higher mean number of mannanases (24) than the saprotrophic

and other mycorrhizal lifestyles (3 to 6) (Figure 7c). However, the MAG contained a

substantially higher mean number of mannanases (9) than the G-AMF (3) and ECM (5), and

also the Endogonales genomes (2 to 4). Similarly, there was a range of mannanases across the

G-AMF genomes with the two Diversisporales genomes containing 11 and 10 counts

respectively, more comparable to the MAG (Table S4). The MAG again contained a

comparatively large set of genes for mannan degradation, with 5 copies of GH92, 3 copies of

GH125, and 1 copy of GH76. However, this represents the one substrate-specific group that

was not as conserved across the phylogenetically closest relatives. None of the Endogonales

genomes contained copies of GH76, while all contained 1 or 2 copies of CE17 (Table S4).

Additionally, while three of the four contained GH92, it was only in single-copy instances.

The G-AMF genomes showed a range of mannanases within their genomes, and particularly

the Diversisporales genomes alone contained many copies of GH92. All but one Glomerales

genome only contained 1 or 2 copies of GH125, with no mannaanase identified in the

'RhiirC2' genome (Table S4).

Low numbers of peptidoglycanases were found in the genomes of all lifestyles, with

significantly larger mean numbers in ERM (8, p = <0.01) compared to the ECM (3) and

saprotroph (2) genomes (Figure 7d). The G-AMF genomes had a comparable mean number

of peptodoglycanases (4) to the MAG (3), while the corresponding numbers in the

Endogonales genomes were also low (0 to 3). The MAG only contained 2 copies of GH24

and a single copy of GH89 (Table S4). Similarly, the other Endogonales genomes were

sparse in peptidoglycanases. Specifically, the 'Jimfl_GMNB39_1' genome contained no

peptidoglycanases, while 'Jimfl_AD_1' and 'Jimlac1' only possessed a single copy of GH89

and CMB50, respectively. 'Endsp1' also contained 2 copies of CBM50 and a single copy of

GH24. The G-AMF genomes also contained a small number of peptidoglycanases, mostly

consisting of GH24 (Table S4).

The MAG contained no genome-specific cell wall degrading enzymes families. However, a

poorly described GH151 family, "α-L-fucosidase enzyme encoding protein", was only

identified within 3 of the closest phylogenetic relatives (Jimfl_AD_1, Jimfl_GMNB39_1, and

Endsp1) and a far-removed saprotrophic basidiomycete, Dacsp1 (Table S4).

Transporters

Since the ability to transport nutrients and assimilate with the host plant is core to the lifestyle of a fungal symbiont, we compared substrate-specific transporters between the ecological lifestyles. Only the ERM genomes contained a statistically higher ($p = 0.044$) number of phosphate transporters than the other genomes (Figure 8a). The MAG contained a relatively low number of phosphate transporters (14), although this was not too dissimilar from the mean of the G-AMF (18) and the same as the mean ECM genome (14). Similarly, the Endogonales genomes possessed low numbers of phosphate transporters (13 to 15) (Figure 8a). The G-AMF genomes had a significantly higher ($p < 0.01$) number of calcium transporters (55) compared to the other mycorrhizal and saprotrophic lifestyles (Figure 8b, which ranged between 26 and 39), and also the endophyte, parasite, pathogen, and mixed lifestyles (Fig S9). The MAG showed a similar number of calcium transporters (50) to the G-AMF, while the number of calcium transporters in the Endogonales genomes varied considerably (25 to 47) (Figure 8b). There was no significant difference ($p = 0.48$) in the number of copper transporters across the mycorrhizal and saprotrophic genomes, with the mean varying between 13 and 17 (Figure 8c). However, the MAG contained only 6 copper transporters, one of the lowest of all genomes (Figure 8c, Fig S9). Of the Endogonales genomes, 'Jimlac1' had the second smallest number of copper transporters (7) while the rest ranged from 8 to 13 (Figure 8c). The G-AMF had a significantly higher ($p < 0.01$) number of nitrate transporters (7) compared to the other mycorrhizal and saprotrophic lifestyles (Figure 8d). The MAG contained a low number (2) of nitrate transporters relative to the mean G-AMF, although this was the same as the mean ECM (2) and similar to the ERM (3), OCM (1), and SAP (1). There was considerable variation in the mean number of nitrate transporters within the Endogonales genomes (0-5) (Figure 8d). The ERM genomes had a significantly higher ($p < 0.01$) mean number of ammonium transporters (11) than all but the G-AMF genomes (9) (Figure 8e). The MAG contained 5 ammonium transporters, slightly higher than the median ECM (4) and OCM (4). The MAG contained an especially low number of glycine transporters (1), the same as the average ECM (1) and OCM (1) genome (Figure 8f). The ERM contained the highest mean number of glycine transporters (5), whereas the G-AMF (3) and SAP (4) were not significantly different to any of the other ecological lifestyles (Figure 8f).

Hallmark Missing Glomeromycotan Core Genes

We analysed both the MAG and G-AMF genomes for enzymes linked to pathways that are

missing from typical G-AMFs, often referred to as 'Missing Glomeromycotan Core Genes'

(MGCG). As fatty acid auxotrophs, G-AMF have been shown to lack the presence of

cytosolic fatty acid synthases required for fatty acid biosynthesis. We found no evidence of

proteins directly associated with the fatty acid synthase required to facilitate fatty acid

biosynthesis within either the G-AMF genomes or the MAG. We did identify 3-oxoacyl-ACP

reductase in all genomes, including 10 within the MAG. (Table S5). Additionally, neither the

MAG nor the G-AMF genomes possessed the ability to biosynthesis thiamine, a hallmark of

G-AMF genomes. Specifically, we found no evidence of the presence of thiamine phosphate

synthase or hydroxymethylpyrimidine kinase (Table S5).

The ability to synthesis all derivatives of the vitamin B6 pathway was present within the

MAG. This included a copy of pyridoxine kinase (2.7.1.35), the most prolific enzyme within

the vitamin B6 metabolism pathway. Only one enzyme, pyridoxine 4-dehydrogenase, was

missing from the pathway, while an indirect three-step metabolism for the conversion of

pyridoxal to pyridoxine was present. Conversely, no G-AMF genome possessed a full

vitamin B6 metabolism pathway, with no evidence of pyridoxine kinase being found across

the group (Table S5).

Discussion

In the current work we assembled the genome of a putative FRE-forming AMF from a root

metagenome. Methodologically this is a relatively unique achievement: fungal MAGs have

been obtained previously from host associated environments and niche environments, but not

from complex environments, such as the rhizosphere (Tagirdzhanova et al., 2021; Shen et al.,

2023; Singh et al., 2023), or from similar fungal symbionts. To achieve this, we had to

augment standard binning methods with assembly graph visualisations and coverage

correlations to known M-AMF 18S rRNA genes. We believe that this method has many

advantages over isolation and culturing, in particular it is less time consuming and in

principle may be more reflective of the actual community present particularly if multiple

species are present.

Phylogenetic analysis of the correlative 18S rRNA marker, consensus_all_seqs_0_16,

showed that the MAG sits within the order Densosporales of the Mucoromycotina. The MAG

represents the first published genome of both the order Densosporales and the family
Planticonsortiaceae. The most closely related taxa with sequenced genomes are putative
ectomycorrhizal and saprotrophic fungi from the order Endogonales, although a reasonable
degree of dissimilarity is evident. The MAG was closely related to environmental sequences
from the rhizoids of liverworts, which form arbuscular-like symbioses involving
Glomeromycota and Mucoromycotina (Rimmington, 2018). To date, understanding of the
phylogeny of fungi which form mycorrhizas with FRE morphology in vascular plants has
been restricted to short 18S rRNA fragments. Similarly, these studies have confirmed close
phylogenetic relatedness with sequences from liverwort thalli (Orchard et al., 2017; Albornoz
et al., 2021, 2022; Seeliger et al., 2024), suggesting these fungi form associations with both
vascular and non-vascular plants, with a role in N and P nutrition of the plant host (Orchard et
al, 2017a; Hoystead, 2023). Mucoromycotina sequences associated with plant roots may span
both the Densosporales (Orchard et al., 2017b; Desirò et al., 2017; Chang et al., 2018) and
Endogonales (Seeliger et al., 2024), and furthermore diverse assemblages of fungi from these
orders may be present in roots (Seeliger et al., 2024), although fungi associated with the FRE
morphology appear to be phylogenetically distinct from putative ectomycorrhizal and
saprotrophic Endogonales (Seeliger et al., 2024).

Comparative genome analysis showed that the MAG's genome had many features in
common with G-AMF. Some of these genomic features, including the low abundance of
PCWDE and the absence of thiamine biosynthesis genes, were also seen in related
Endogonales genomes and therefore aren't likely to be specific, hallmark features of an
adaptation to a symbiotic lifestyle. However other signatures the MAG shares with G-AMF,
including a high abundance of calcium transporters and the absence of fatty acid synthases,
may point to physiological convergence in the way in which the symbioses operate. There
were some notable differences between the MAG and G-AMF, with the MAG possessing a
high abundance of chitinases, a complete vitamin B6 biosynthesis pathway, and low
abundances of nitrate, ammonium and copper transporters. These features may point to
differences between G-AMF and M-AMF in the way they interact with the host plant and the
wider environment.

It has been suggested that the transition from saprotrophic to mycorrhizal lifestyles involved
loss of PCWDE, enabling fungi to exist within roots without digesting them (Miyauchi et al.,
2020). The G-AMF symbiosis is estimated to have evolved 450 mya, and Glomeromycotan
genomes possess substantially lower numbers of these genes than the fungi forming the more

recently evolved ectomycorrhizal, ericoid mycorrhizal, and orchid mycorrhizal symbioses.

Furthermore, G-AMF show signatures of dependence on host metabolism that are absent

from more recently evolved symbioses, including fatty acid auxotrophy and an inability to

synthesise thiamine and metabolise vitamin B6 (Miyauchi et al., 2020). These represent the

'Missing Glomeromycota Core Genes' (MGCG, Tang et al., 2016; Kobayashi et al., 2018;

Rosling et al., 2024).

Although genomic evidence indicates that G-AMF and M-AMF have independent origins

(Rosling et al., 2024), these symbioses co-occurred in the earliest land plants (Strullu-Derrien

et al., 2014), and they continue to co-occur in roots of extant plants (Orchard et al., 2017a).

Additionally, while G-AMF and M-AMF exhibit some niche differentiation, evidence of

overlapping ecological presence and similar responses to overarching edaphic and

environmental variables have been noted (Albornoz et al., 2022; Seeliger et al., 2024).

Furthermore, ultrastructural evidence points to similarities in the way in which G-AMF and

M-AMF function at a cellular level (Albornoz et al., 2020). Therefore, similar evolutionary

pressures may have acted on these symbioses, which could have resulted in convergence of

genomic traits. Indeed, many of the symbiotic genomic features shown by G-AMF were also

displayed by the MAG. This included a reduced repertoire of PCWDE and the absence of

several MGCG, including those related to thiamine and fatty acid biosynthesis.

Our analysis confirms the findings of Rosling et al. (2024) that Endogonales have a low

genomic PCWDE count and an incomplete thiamine biosynthesis pathway, suggesting that

these may be features common to both the Densosporales and Endogonales, rather than an

adaptation to biotrophy. However, the Endogonales genomes possess fatty acid biosynthesis

genes (Rosling et al., 2024), so this signature of biotrophy does likely represent the same

adaptation to biotrophy in G-AMF and M-AMF, which suggests that host assimilate is

provided as fatty acids in both symbioses.

The MAG had a number of features which distinguished it from G-AMF genomes. In

particular, while there was a shared low abundance of CAZymes, the MAG possessed a

greater repertoire of enzymes for degradation of organic substrates than typically seen in G-

AMF genomes. This included the full complement of enzymes required for degradation of

cellulose, which was absent in all G-AMF genomes. In notable contrast to G-AMF, the MAG

had a high count of both chitin and mannan fungal cell wall degrading enzymes. The high

chitinase count doesn't appear to be specific to the M-AMF, with the putative

ectomycorrhizal and saprotrophic *Endogonales* genomes showing a similar pattern, attributed

to elevated numbers of CE4 and GH18, as seen previously in Rosling et al. (2024). However, the full complement of chitinases was present in the G-AMF genomes as well as in the MAG, so this feature may not have any functional significance. Relative to other mycorrhizal fungi (except ericoid mycorrhizal fungi), the MAG had a notably high count of genes for mannan degradation. These were absent or present in low numbers in most G-AMF genomes and, significantly, were also found in low abundance in the *Endogonales* genomes, therefore high abundance of mannan degradation genes may represent a distinguishing feature of FRE genomes. However, even within the G-AMF there appears to be some variation, with the *Diversisporales* genomes containing a higher number of mannan and chitin degrading enzymes than the *Glomerales* genomes, indicating a substantial range of genomic repertoires.

It is important to understand whether these contrasting enzymatic profiles contribute to niche partitioning between G-AMF and M-AMF in nutrient mobilisation from contrasting mineral and organic substrates. For example, high abundance of chitin and mannan degrading genes could indicate a role for M-AMF in the degradation of fungal necromass, while a complete cellulose degradation pathway and high abundance of cellulose and hemicellulose degrading enzymes could suggest involvement in saprotrophic breakdown of plant-derived materials, similar to some ECM fungi (Janusz et al., 2017; Martin & van der Heijden., 2024). Alternatively, it could reflect different strategies for enzymatic degradation of plant cell walls as they colonise and grow through host plant tissues and penetrate root cells.

Currently there is no indication that soil organic matter has any influence on the distribution of G-AMF or M-AMF. In contrast to G-AMF, evidence indicates that M-AMF are more abundant in agricultural than natural habitats, indicating a preference for low organic matter soils (Albornoz et al., 2022). However, these functional differences could support niche partitioning within these habitats.

A further distinguishing feature between the MAG and the G-AMF genomes is the presence of a full B6 biosynthesis pathway in the MAG, including pyroxidine kinase, which was missing in all the G-AMF genomes, as demonstrated in earlier studies (Tang et al., 2016; Kobayashi et al., 2018). B6 is a cofactor required for amino acid biosynthesis, and it also plays a role as an antioxidant. Reactive oxygen species (ROS) scavenging systems such as B6 may be required by AMF to overcome host defences, allowing intracellular colonisation and the development of arbuscules (Belmondo et al., 2016; Lanfranco et al., 2018). Components of the B6 biosynthesis pathway that G-AMF do possess are expressed throughout the fungal life cycle, but they are at the highest expression levels within the root (Benabdellah et al.,

2009). Similarly, B6 plays a role in supporting the infection of root tissues by pathogens
(Wang et al., 2023). Differences in capacity for independent production of B6 between the
MAG and G-AMF could point to differences in the way the fungi scavenge ROS, and
potentially also differences in the way root colonisation and arbuscule development are
regulated by the plant. There could also be implications for the specific mechanisms
underlying M-AMF and G-AMF responses to external stressors which generate ROS such as
temperature, osmotic stress, and nutrient limitation, and the independence of these responses
to the plant's own response (Sahu et al., 2022).

Key physiological traits associated with G-AMF symbioses are the uptake of P from the soil,
its translocation through hyphae to the root, and its transfer to the plant in exchange for
carbon. Ultrastructural (Albornoz et al., 2020) and experimental evidence (Orchard et al.,
2017b) suggest a similar role for M-AMF. A number of P transporters have been identified in
G-AMF involved in P assimilation from soil, storage of P, and exchange of P with the host
(Javot et al., 2006; Harrison et al., 2010; Plassard et al., 2019). G-AMF genomes generally
showed higher mean numbers of P transporters than ECM, while numbers in the MAG were
low relative to other mycorrhizal genomes studied. While the significance of transporters to
the functioning of symbionts will depend on a range of factors including their efficiency, our
data doesn't provide genomic support for particular specialisation in P nutrition by M-AMF.

Evidence also points to a potential role for G-AMF in N nutrition (Gosling et al., 2013).
Indeed, G-AMF genomes contained higher numbers of nitrate transporters than the other
mycorrhizal symbionts, while also sitting towards the higher end of both ammonium and
glycine transporters within the genomes we analysed. However, further evidence of inter-
group differentiation can be seen across the Diversisporales and Glomerales genomes, with
the former containing especially high numbers of ammonium and glycine transporters. The
MAG contained a lower number of all three categories of N transporters than all G-AMF
genomes. While the MAG contained a typical number of ammonium transporters compared
to the other mycorrhizal lifestyles, it contained a low number of nitrate and glycine
transporters. Hoystead et al. (2023) suggested that M-AMF and G-AMF have complementary
roles in host nutrition, with M-AMF predominantly involved in supply of N to the host and
G-AMF in supplying P. Furthermore, Howard et al. (2024) showed a putative M-AMF
displaying both a preference for, and increased transfer of, glycine and ammonium derived
nitrogen sources. While there is evidence for the ability to transport varying forms of N, our
MAG does not provide exceptional evidence for this relative specialism.

The very low abundance of Cu transporters in the MAG - the lowest across the entire dataset - is particularly striking. The three ectomycorrhizal Endogonales genomes also contain low numbers of Cu transporters, however the saprotrophic Endogonales sit near the mean. G-AMF may promote Cu uptake by the host plant (Taylor et al., 2015) and arbuscules have high demands for Cu (Gomez-Gallego et al., 2019). Furthermore, Cu transporters in G-AMF show differential expression profiles across intra- and extra- radical mycelium (Gomez-Gallego et al., 2019). The extent to which M-AMF are involved in the supply of micronutrients to the host remains to be determined, but the evidence presented here could indicate that in contrast to G-AMF, this may not be an important function for M-AMF.

The high number of calcium transporters in both the MAG and G-AMF is a clear point of similarity between them, and of differentiation to all other groups. AMF have been implicated in transport of Ca to host plants (e.g. Liu et al., 2002), and our data could therefore indicate a role for both G-AMF and M-AMF in this function. However, Ca may also play a key role in the arbuscule lifecycle of both FRE and G-AMF. A clear link between a dramatic increase in calcium accumulation and arbuscule collapse has been demonstrated within G-AMF by both Ryan et al. (2003) and Sujkowska-Rybkowska and Znojek (2018). A similar trend within FRE was documented by Gianinazzi-Pearson et al. (1981) and Albornoz et al. (2020). While it is unknown whether the Ca accumulated originates from the host plant or the fungi, the high number of Ca transporters within both G-AMF and the MAG may indicate that Ca accumulation is driven by the fungal symbionts, perhaps depositing Ca to cease the function of the arbuscule once it reaches maturity.

Genome analyses, including those reported here, have identified differences across G-AMF which suggest contrasting interactions with the host and the wider environment. For both G-AMF and M-AMF, comprehensive analysis of genomic functional gene repertoires across the broad range of taxa found within roots is key to understanding the way that AMF communities support plant health, and the potential for engineering communities to promote beneficial traits. With the development of third generation long read sequencing technologies, genome assembly from metagenomes represents a promising route for understanding functions and traits associated with dark taxa in complex microbial communities, especially when combined with transcriptome sequencing (Benoit et al., 2024) to understand the factors which regulate gene expression within plant tissues and the soil.

Acknowledgements

We thank the Natural Environment Research Council (Grant NE/S010270/1), the

Biotechnology and Biological Sciences Research Council (BBSRC) Midlands Integrative

Biosciences Training Partnership (Grant BB/T00746X/1) and the Australian Research

Council (Discovery Project DP180103157) for funding.

Author contributions

GB, MHR, and CQ conceived the work and with RM obtained funding. MHR and PAD

conducted the M-AMF enrichment work and the assessment of M-AMF colonisation of plant

roots. SH, SR, and GB conducted sequencing and analysis of 18S rRNA genes. SR and CQ

assembled the MAG. RHN, GM, and JC conducted the 18S rRNA gene phylogenetic

analysis. JC, GM, SR, and GB performed the comparative genomic analyses. JC and GB

wrote the manuscript. All authors edited the manuscript.

Data Availability

Raw 18S rRNA Miseq sequences are available at NCBI (BioProject ID PRJNA1179892), and

Nanopore sequences at ENA (Experiment Accession ERX13330708). The FRE4583

sequence is available at NCBI (SUB14894546). The 18S rRNA phylogeny sequence

alignment FASTA file and Newick tree file are available on PlutoF Biodiversity Platform

(https://dx.doi.org/10.15156/BIO/3301225). The sequence of the MAG has been deposited on

the Zenodo repository (https://doi.org/10.5281/zenodo.14144746).

References

Albornoz FE, Hayes PE, Orchard S, Clode PL, Nazeri NK, Standish RJ, Bending GD, Hilton

S, Ryan MH. 2020. First cryo-scanning electron microscopy images and X-ray microanalyses

of mucoromycotinian fine root endophytes in vascular plants. *Frontiers in Microbiology* 11:

2018.

Albornoz FE, Orchard S, Standish RJ, Dickie IA, Bending GD, Hilton S, Lardner T, Foster

KJ, Gleeson DB, Bougoure J et al. 2021. Evidence for niche differentiation in the

environmental responses of co-occurring mucoromycotinian fine root endophytes and

glomeromycotinian arbuscular mycorrhizal fungi. *Microbial Ecology* 81(4): 864-873.

Albornoz FE, Ryan MH, Bending GD, Hilton S, Dickie IA, Gleeson DB, Standish RJ. 2022.

Agricultural land-use favours Mucoromycotinian, but not Glomeromycotinian, arbuscular

mycorrhizal fungi across ten biomes. *New Phytologist* 233(3): 1369-1382.

Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, ljaz UZ, Loman NJ, Andersson

AF, Quince C. 2014. Binning metagenomic contigs by coverage and composition. *Nature*

Methods 11: 1144–1146.

Almagro Armenteros JJ, Salvatore M, Emanuelsson O, Winther O, von Heijne G, Elofsson

A, Nielsen H. 2019. Detecting sequence signals in targeting peptides using deep learning.

Life Science Alliance 2(5): e201900429.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD et al 2012. SPAdes: a new genome assembly algorithm

and its applications to single-cell sequencing. *Journal of Computational Biology* 19(5): 455-

477.

Begum N, Qin C, Ahanger MA, Raza S, Khan MI, Ashraf M, Ahmed N, Zhang L. 2019. Role

of arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress

tolerance. *Frontiers in Plant Science* 10: 1068.

Belmondo S, Calcagno C, Genre A, Puppo A, Pauly N, Lanfranco L. 2016. NADPH oxidases

in the arbuscular mycorrhizal symbiosis. *Plant Signaling & Behavior* 11(4): e1165379.

Benabdellah L, Azcón-Aguilar C, Valderas A, Speziga D, Fitzpatrick TB, Ferrol N. 2009.

GintPDX1 encodes a protein involved in vitamin B6 biosynthesis that is up-regulated by

oxidative stress in the arbuscular mycorrhizal fungus *Glomus intraradices*. *New Phytologist*

184(3): 682-693.

Benoit G, Raguideau S, James R, Phillippy AM, Chikhi R, Quince C. 2024. High-quality
metagenome assembly from long accurate reads with metaMDBG. *Nature Biotechnology* 42:
1378–1383.
939
940 Brundrett MC, Tedersoo L. 2018. Evolutionary history of mycorrhizal symbioses and global
941 host plant diversity. *New Phytologist* 220(4):1108-1115.
942
943 Brundrett MC. 2006. Understanding the roles of multifunctional mycorrhizal and endophytic
944 fungi. In: BJE Schulz, CJC Boyle, TN Sieber, eds. *Microbial Root Endophytes*. Berlin,
945 Germany: Springer-Verlag, 281–293.
946
947 Buchfink B, Reuter K, Drost HG. 2021. Sensitive protein alignments at tree-of-life scale
948 using DIAMOND. *Nature Methods* 18: 366-368.
949
950 Chang Y, Desirò A, Na H, Sandor L, Lipzen A, Clum A, Barry K, Grigoriev IV, Martin FM,
951 Stajich JE et al. 2019. Phylogenomics of Endogonales and evolution of mycorrhizas within
952 Mucoromycota. *New Phytologist* 222(1): 511-525.
953
954 Chang Y, Wang Y, Mondo S, Ahrendt S, Andreopoulos W, Barry K, Beard J, Benny GL,
955 Blankenship S, Bonito G et al. 2022. Evolution of zygomycete secretomes and the origins of
956 terrestrial fungal ecologies. *iScience* 25(8): 104840.
957
958 Chen J, Tang Y, Kohler A, Lebreton A, Xing YM, Zhou D, Li Y, Martin FM, Guo S. 2022.
959 Comparative transcriptomics analysis of the symbiotic germination of *D. officinale*
960 (Orchidaceae) with emphasis on plant cell wall modification and cell wall-degrading
961 enzymes. *Frontiers in Plant Science* 13:880600.
962
963 Chen M, Arato M, Borghi L, Nouri E, Reinhardt D. 2018. Beneficial services of arbuscular
964 mycorrhizal fungi – from ecology to application. *Frontiers in Plant Science* 9: 1270.
965
966 Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
967 McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and BCFtools.
968 *GigaScience* 10(2): giab008.
969

De Castro E, Sigrist CJ, Gattiker A, Bulliard V, Langendijk-Genevaux PS, Gasteiger E,
Bairoch A, Hulo N. 2006. ScanProsite: detection of PROSITE signature matches and
ProRule-associated functional and structural residues in proteins. *Nucleic Acids Research* 34:
362-5.

Desirò A, Rimington WR, Jacob AS, Pol NV, Smith MR, Trappe JM, Bidartondo MI,
Bonito G. 2017. Multigene phylogeny of Endogonales, an early diverging lineage of fungi
associated with plants. *IMA Fungus* 8(2): 245–257.

Eddy SR. 2011. Accelerated profile HMM searches. *PLOS Computational Biology* 7(10):
e1002195.

Frąc M, Hannula SE, Bełka M, Jędryczka M. 2018. Fungal biodiversity and their role in soil
health. *Frontiers in Microbiology* 9: 707.

Gianinazzi-Pearson V, Morandi D, Dexheimer J, Gianinazzi S. 1981. Ultrastructural and
ultracytochemical features of a *Glomus tenuis* mycorrhiza. *New Phytologist* 88(4): 633-639.

Gómez-Gallego T, Benabdellah K, Merlos MA, Jiménez-Jiménez AM, Alcon C, Berthomieu
P, Ferrol N. 2019. The *Rhizophagus irregularis* genome encodes two CTR copper
transporters that mediate Cu import into the cytosol and a CTR-like protein likely involved in
copper tolerance. *Frontiers in Plant Science* 10: 604.

Gong Y, Lebreton A, Zhang F, Martin FM. 2023. Role of carbohydrate-active enzymes in
mycorrhizal symbioses. *Essays in Biochemistry* 67(3): 471-478.

Gosling P, Mead A, Proctor M, Hammond JP, Bending GD. 2013. Contrasting arbuscular
mycorrhizal communities colonizing different host plants show a similar response to a soil
phosphorus concentration gradient. *New Phytologist* 198(2): 546-556.

Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao X,
Korzeniewski F et al. 2013. MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic*
Acids Research 42: 699-704.

1003

1004 Gruber-Vodicka HR, Seah BKB, Pruesse E. 2020. phyloFlash: Rapid small-subunit rRNA
1005 profiling and targeted assembly from metagenomes. m*Systems* 5(5): e00920-20.

1006

1007 Harrison MJ, Pumplin N, Breuillin FJ, Noar RD, Park HJ. 2010. Phosphate transporters in
1008 arbuscular mycorrhizal symbiosis. In: Koltai H, Kapulnik Y, eds. *Arbuscular Mycorrhizas:*
1009 *Physiology and Function.* Dordrecht, Netherlands: Springer, 117–135.

1010

1011 Hawkins HJ, Cargill RIM, Nuland MEV, Hagen SC, Field KJ, Sheldrake M, Soudzilovskaia
1012 NA, Kiers TE. 2023. Mycorrhizal mycelium as a global carbon pool. *Current Biology* 33(11):
1013 560-573.

1014

1015 Hirose D, Degawa Y, Yamamoto K, Yamada A. 2014. Sphaerocreas pubescens is a member
1016 of the Mucoromycotina closely related to fungi associated with liverworts and hornworts.
1017 *Mycoscience* 55(3): 221-226.

1018

1019 Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving
1020 the ultrafast bootstrap approximation. *Molecular Biology and Evolution*. 35: 518–522.

1021

1022 Horton P, Park KJ, Obayashi T, Fujita N, Harada H, Adams-Collier CJ, Nakai N. 2007.
1023 WoLF PSORT: protein localization predictor. *Nucleic Acids Research* 5(2): 585–587.

1024

1025 Howard NAO, Williams A, Durant E, Pressel S, Daniell TJ, Field KJ. 2024. Preferential
1026 nitrogen and carbon exchange dynamics in Mucoromycotina "fine root endophyte"-plant
1027 symbiosis. *Current Biology*.

1028

1029 Hoysted GA, Field KJ, Sinanaj B, Bell CA, Bidartondo MI, Pressel S. 2022. Direct nitrogen,
1030 phosphorus and carbon exchanges between Mucoromycotina 'fine root endophyte' fungi and
1031 a flowering plant in novel monoxenic cultures. *New Phytologist* 238(1): 70-79.

1032

1033 Hoysted GA, Jacob AS, Kowal J, Giesemann P, Bidartondo MI, Duckett JG, Gebauer G,
1034 Rimington WR, Schornack S, Pressel S et al. 2019. Mucoromycotina fine root endophyte
1035 fungi form nutritional mutualisms with vascular plants. *Plant Physiology* 181(2): 565-577.

1036

1037   Huey CJ, Gopinath SCB, Uda MNA, Zulhaimi HI, Jaafar MN, Kasim FH, Yaakub ARW.

1038   2020. Mycorrhiza: a natural resource assists plant growth under varied soil conditions.

1039   *3Biotech* 10(5): 204.

1040

1041   Janusz G, Pawlik A, Sulej J, Świderska-Burek U, Jarosz-Wilkołazka A, Paszczyński A. 2017.

1042   Lignin degradation: microorganisms, enzymes involved, genomes analysis and evolution.

1043   *FEMS Microbiology Reviews* 41(6): 941-962.

1044

1045   Javot H, Pumplin N, Harrison MJ. 2006. Phosphate in the arbuscular mycorrhizal symbiosis:

1046   Transport properties and regulatory roles. *Plant, Cell & Environment* 30: 310–322.

1047

1048   Jeffery RP, Simpson RJ, Lambers H, Orchard S, Kidd DR, Haling RE, Ryan MH. 2018.

1049   Contrasting communities of arbuscule-forming root symbionts change external critical

1050   phosphorus requirements of some annual pasture legumes. *Applied Soil Ecology* 126: 88-97.

1051

1052   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder:

1053   Fast model selection for accurate phylogenetic estimates. *Nature Methods.* 14: 587-589.

1054

1055   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

1056   improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–

1057   780.

1058

1059   Kim GB, Kim JY, Lee JA, Norsigian CJ, Palsson BO, Lee SY. 2023. Functional annotation

1060   of enzyme-encoding genes using deep learning with transformer layers. *Nature*

1061   *Communications* 14: 7370.

1062

1063   Kobayashi Y, Maeda T, Yamaguchi K, Kameoka H, Tanaka S, Ezawa T, Shigenobu S,

1064   Kawaguchi M. 2018. The genome of Rhizophagus clarus HR1 reveals a common genetic

1065   basis for auxotrophy among arbuscular mycorrhizal fungi. *BMC Genomics* 19(1): 465.

1066

1067   Lanfranco L, Fiorilli V, Gutjahr C. 2018. Partner communication and role of nutrients in the

1068   arbuscular mycorrhizal symbiosis. *New Phytologist* 220(4): 1031-1046.

1069

1070   Lebreton A, Zeng Q, Miyauchi S, Kohler A, Dai YC, Martin FM. 2021. Evolution of the

1071   mode of nutrition in symbiotic and saprotrophic fungi in forest ecosystems. *Annual Review of*

1072   *Ecology, Evolution, and Systematics* 52(1): 385-404.

1073

1074   Lee S-J, Kong M, St-Arnaud M, Hijri M. 2020. Arbuscular mycorrhizal fungal communities

1075   of native plant species under high petroleum hydrocarbon contamination highlights

1076   Rhizophagus as a key tolerant genus. *Microorganisms* 8(6): 872.

1077

1078   Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

1079   MEM. *ArXiv*. doi: 10.48550/arXiv.1303.3997

1080

1081   Liu A, Hamel C, Elmi A, Costa C, Ma B, Smith DL. 2002. Concentrations of K, Ca and Mg

1082   in maize colonized by arbuscular mycorrhizal fungi under field conditions. *Canadian Journal*

1083   *of Soil Science* 82(3): 271-279.

1084

1085   Luginbuehl LH, Menard GN, Kurup S, Van Erp H, Radhakrishnan GV, Breakspear A,

1086   Oldroyd GED, Eastmond PJ. 2017. Fatty acids in arbuscular mycorrhizal fungi are

1087   synthesized by the host plant. *Science* 365(6343): 1175-1178.

1088

1089   Luginbuehl LH, Oldroyd GED. 2017. Understanding the arbuscule at the heart of

1090   endomycorrhizal symbioses in plants. *Current Biology* 27(17): 952-963.

1091

1092   Maillard F, Schilling J, Andrews E, Schreiner KM, Kennedy P. 2020. Functional

1093   convergence in the decomposition of fungal necromass in soil and wood. *FEMS*

1094   *Microbiology Ecology* 96(2): fiz209.

1095

1096   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel

1097   and streamlined workflows along with broader and deeper phylogenetic coverage for scoring

1098   of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution* 38(10):

1099   4647-4654.

1100

1101   Mansfield TM, Albornoz FE, Ryan MH, Bending GD, Standish RJ. 2023. Niche

1102   differentiation of Mucoromycotinian and Glomeromycotinian arbuscular mycorrhizal fungi

1103   along a 2-million-year soil chronosequence. *Mycorrhiza* 33(3): 139-152.

1104

1105 Martin FM, van der Heijden MGA. 2024. The mycorrhizal symbiosis: research frontiers in

1106 genomics, ecology, and agricultural application. *New Phytologist* 242(4): 1486-1506.

1107

1108 Martino E, Morin E, Grelet G-A, Kuo A, Kohler A, Daghino S, Barry KW, Cichocki N,

1109 Clum A, Dockter RB et al. 2018. Comparative genomics and transcriptomics depict ericoid

1110 mycorrhizal fungi as versatile saprotrophs and plant mutualists. *New Phytologist* 217(3):

1111 1213-1229.

1112

1113 McGonigle TP, Miller MH, Evans DG, Fairchild GL, Swan JA. 1990. A new method which

1114 gives an objective measure of colonization of roots by vesicular—arbuscular mycorrhizal

1115 fungi. *New Phytologist* 155(3): 495-501.

1116

1117 Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

1118 Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference

1119 in the genomic era. *Molecular Biology and Evolution*. 37(5): 1530-1534.

1120

1121 Mistry J, Bateman A, Finn RD. 2007. Predicting active site residue annotations in the Pfam

1122 database. *BMC Bioinformatics* 8: 298.

1123

1124 Miyauchi S, Kiss E, Kuo A, Drula E, Kohler A, Sánchez-García M, Morin E, Andreopoulos

1125 B, Barry KW, Bonito G et al. 2020. Large-scale genome sequencing of mycorrhizal fungi

1126 provides insights into the early evolution of symbiotic traits. *Nature Communications* 11:

1127 5125.

1128

1129 Möller S, Croning MDR, Apweiler R. 2001. Evaluation of methods for the prediction of

1130 membrane spanning regions. *Bioinformatics* 17(7): 646–653.

1131

1132 Montoliu-Nerin M, Sánchez-García M, Bergin C, Kutschera VE, Johannesson H, Bever JD,

1133 Rosling A. 2021. In-depth phylogenomic analysis of arbuscular mycorrhizal fungi based on a

1134 comprehensive set of de novo genome assemblies. *Frontiers in Fungal Biology* 29(2):

1135 716385.

1136

1137 Morin E, Miyauchi S, Clemente HS, Chen ECH, Pelin A, Providencia I, Ndikumana S,

1138 Beaudet D, Hainaut M, Drula E et al. 2019. Comparative genomics of *Rhizophagus*

1139 *irregularis* , *R. cerebriforme* , *R. diaphanus* and *Gigaspora rosea* highlights specific genetic

1140 features in Glomeromycotina. *New Phytologist* 222(3): 1584-1598.

1141

1142 Morris JL, Puttick MN, Clark JW, Edwards D, Kenrick P, Pressel S, Wellman CH, Yang Z,

1143 Schneider H, Donoghue PCJ. 2018. The timescale of early land plant evolution. *Proceedings*

1144 *of the National Academy of Sciences* 115(10): 2274-2283.

1145

1146 Naranjo-Ortiz MA, Gabaldón T. 2019. Fungal evolution: major ecological adaptations and

1147 evolutionary transitions. *Biological Reviews* 94(4): 1443–1476.

1148

1149 Öpik M, Davison J, Moora M, Zobel M. 2014. DNA-based detection and identification of

1150 Glomeromycota: the virtual taxonomy of environmental sequences. *Botany-Botanique* 92:

1151 135–147.

1152

1153 Orchard S, Hilton S, Bending GD, Dickie IA, Standish RJ, Gleeson DB, Jeffery RP, Walker

1154 C, Bass D, Monk J et al. 2017a. Fine endophytes (Glomus tenue) are related to

1155 Mucoromycotina, not Glomeromycota. *New Phytologist* 213(2): 481-486.

1156

1157 Orchard S, Standish RJ, Dickie IA, Renton M, Walker C, Moot D, Ryan MH. 2017b. Fine

1158 root endophytes under scrutiny: a review of the literature on arbuscule-producing fungi

1159 recently suggested to belong to the Mucoromycotina. *Mycorrhiza* 27(7): 619-638.

1160

1161 Orchard S, Standish RJ, Nicol D, Gupta VVSR, Ryan MH. 2016. The response of fine root

1162 endophyte (Glomus tenue) to waterlogging is dependent on host plant species and soil type.

1163 *Plant and Soil* 403: 305-315.

1164

1165 Palos-Fernández R, Aguilar-Pontes MV, Puebla-Planas G, Berger H, Studt-Reinhold L,

1166 Strauss J, Di Pietro A, López-Berges MS. 2024. Copper acquisition is essential for plant

1167 colonization and virulence in a root-infecting vascular wilt fungus. *bioRxiv*. doi:

1168 10.1101/2024.04.23.590778

1169

1170 Parniske, M. 2008. Arbuscular mycorrhiza: the mother of plant root endosymbiosis. *Nature*

1171 *Reviews Microbiology* 6: 763–775.

1172

1173 Peay KG, Kennedy PG, Talbot JM. 2016. Dimensions of biodiversity in the Earth

1174 mycobiome. *Nature Reviews Microbiology* 14: 434–447.

1175

1176 Pellegrin C, Morin E, Martin FM, Veneault-Fourrey C. 2015. Comparative analysis of

1177 secretomes from ectomycorrhizal fungi with an emphasis on small-secreted proteins.

1178 *Frontiers in Microbiology* 6: 1278.

1179

1180 Peng X, Wilken SE, Lankiewicz TS, Gilmore SP, Brown JL, Henske JK, Swift CL, Salamov

1181 A, Barry K, Grigoriev IV et al. 2021. Genomic and functional analyses of fungal and

1182 bacterial consortia that enable lignocellulose breakdown in goat gut microbiomes. *Nature*

1183 *Microbiology* 6: 499–511.

1184

1185 Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal

1186 peptides from transmembrane regions. *Nature Methods* 8(10): 785-6.

1187

1188 Plassard C, Becquer A, Garcia, K. 2019. Phosphorus transport in mycorrhiza: How far are

1189 we?. *Trends in Plant Science* 24: 794–801.

1190

1191 Prout JN, Williams A, Wanke A, Schornack S, Ton J, Field KJ. 2024. Mucoromycotina 'fine

1192 root endophytes': a new molecular model for plant–fungal mutualisms?. *Trends in Plant*

1193 *Science* 29(6): 650-661.

1194

1195 Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013.

1196 The SILVA ribosomal RNA gene database project: improved data processing and web-based

1197 tools. *Nucleic Acids Research* 41: 590–596.

1198

1199 Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

1200 features. *Bioinformatics* 26(6): 841-2.

1201

1202 R Core Team. 2023. R: A language and environment for statistical computing. R Foundation

1203 for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

1204

1205   Rambaut, A. 2009. Figtree. Tree figure drawing tool, v1.4.4. Edinburgh, Scotland.

1206   http://tree.bio.ed.ac.uk/software/figtree.

1207

1208   Rasmussen HN, Rasmussen FN. 2014. Seedling mycorrhiza: a discussion of origin and

1209   evolution in Orchidaceae. *Botanical Journal of the Linnean Society* 175(3): 313-327.

1210

1211   Read DJ, Duckett JG, Francis R, Ligron R, Russell A. 2000. Symbiotic fungal associations in

1212   'lower' land plants. *Philosophical Transactions of the Royal Society of London* 355(1398):

1213   815–831.

1214

1215   Rimington WR, Pressel S, Duckett JG, Field KJ, Read DJ, Bidartondo MI. 2018. Ancient

1216   plants with ancient fungi: liverworts associate with early-diverging arbuscular mycorrhizal

1217   fungi. *Proceedings of the Royal Society B: Biological Sciences* 285(1888): 20181600.

1218

1219   Rosling A, Sahraei SE, Khan FK, Desirò A, Bryson AE, Mondo SJ, Grigoriev IV, Bonito G.

1220   Sánchez-García M. 2024. Evolutionary history of arbuscular mycorrhizal fungi and genomic

1221   signatures of obligate symbiosis. *BMC Genomics* 25: 529.

1222

1223   Ryan MH, McCully ME, Huang CX. 2003. Location and quantification of phosphorus and

1224   other elements in fully hydrated, soil-grown arbuscular mycorrhizas: a cryo-analytical

1225   scanning electron microscopy study. *New Phytologist* 160(2):429-441.

1226

1227   Sahu PK, Jayalakshmi K, Tilgam J, Gupta A, Nagaraju Y, Kumar A, Hamid S, Singh HV,

1228   Minkina T et al. 2022. ROS generated from biotic stress: Effects on plants and alleviation by

1229   endophytic microbes. *Frontiers in Plant Science* 13: 1042936.

1230

1231   Saier MH, Reddy VS, Moreno-Hagelsieb G, Hendargo KJ, Zhang Y, Iddamsetty V, Lam

1232   KJK, Tian N, Russum S, Wang J, Medrano-Soto A. 2021. The transporter classification

1233   database (TCDB): 2021 update. *Nucleic Acids Research* 49(1): 461-467.

1234

1235   Schüßler A, Martin H, Cohen D, Fitz M, Wipf D. 2007. Arbuscular mycorrhiza: studies on

1236   the geosiphon symbiosis lead to the characterization of the first glomeromycotan sugar

1237   transporter. *Plant Signaling & Behavior* 2(5), 431–434.

1238

1239 Schüßler A, Schwarzott D, Walker C. 2001. A new fungal phylum, the Glomeromycota:

1240 phylogeny and evolution. *Mycological Research* 105(12): 1413-1421.

1241

1242 Seeliger M, Hilton S, Muscatt G, Walker C, Bass D, Albornoz F, Standish RJ, Gray ND,

1243 Mercy L, Rempelos L et al. 2024. New fungal primers reveal the diversity of

1244 Mucoromycotinian arbuscular mycorrhizal fungi and their response to nitrogen application.

1245 *Environmental Microbiome* 19: 71.

1246

1247 Setubal JC. 2021. Metagenome-assembled genomes: concepts, analogies, and challenges.

1248 *Biophysical Reviews* 13(6): 905–909.

1249

1250 Shen Z, Robert L, Stolpman M, Che Y, Allen KJ, Saffery R, Walsh A, Young A, Eckert J,

1251 Deming C et al. 2023. A genome catalog of the early-life human skin microbiome. *Genome*

1252 *Biology* 24(252).

1253

1254 Shi Z, Deng R, Yuan Q, Mao Z, Wang R, Li H, Liao X, Ma H. 2023. Enzyme commission

1255 number prediction and benchmarking with hierarchical dual-core multitask learning

1256 framework. *Research* 31(6): 0153

1257

1258 Singh NK, Wood JM, Patane J, Moura LMS, Lombardino J, Setubal JC, Venkateswaran K.

1259 2023. Characterization of metagenome-assembled genomes from the International Space

1260 Station. Microbiome 11(1): 125.

1261

1262 Smith AD, Logeman BL, Thiele DJ. 2017. Copper acquisition and utilization in fungi.

1263 *Annual Review of Microbiology* 8(71): 597-623.

1264

1265 Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G, Corradi N,

1266 Grigoriev I, Gryganskyi A et al. 2016. A phylum-level phylogenetic classification of

1267 zygomycete fungi based on genome-scale data. *Mycologia* 108(5): 1028–1046.

1268

1269 Strullu-Derrien C, Selosse M-A, Kenrick P, Martin FM. 2018. The origin and evolution of

1270 mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytologist* 220(4):

1271 1012-1030.

1272

1273   Sujkowska-Rybkowska M, Znojek E. 2018. Localization of calreticulin and calcium ions in

1274   mycorrhizal roots of Medicago truncatula in response to aluminum stress. *Journal of Plant*

1275   *Physiology* 229: 22-31.

1276

1277   Tagirdzhanova G, Saary P, Tingley JP, Díaz-Escandón D, Abbott DW, Finn RD, Spribille T.

1278   2021. Predicted input of uncultured fungal symbionts to a lichen symbiosis from

1279   metagenome-assembled genomes. *Genome Biology and Evolution* 13(4).

1280

1281   Talbot JM, Allison SD, Treseder KK. 2008. Decomposers in disguise: mycorrhizal fungi as

1282   regulators of soil C dynamics in ecosystems under global change. *Functional Ecology* 22(6):

1283   955-963.

1284

1285   Tamura K, Stecher G, Kumar S. 2021. MEGA11: molecular evolutionary genetics analysis

1286   version 11. *Molecular Biology and Evolution* 38(7): 3022–3027.

1287

1288   Tanaka S, Hashimoto K, Kobayashi Y, Yano K, Maeda T, Kameoka H, Ezawa T, Saito K,

1289   Akiyama K, Kawaguchi M. 2022. Asymbiotic mass production of the arbuscular mycorrhizal

1290   fungus Rhizophagus clarus. *Communications Biology* 5: 43.

1291

1292   Tang N, Clemente HS, Roy S, Bécard G, Zhao B, Roux C. 2016. A survey of the gene

1293   repertoire of *Gigaspora rosea* unravels conserved features among Glomeromycota for

1294   obligate biotrophy. *Frontiers in Microbiology* 1(7):233.

1295

1296   Taylor A, Pereira N, Thomas B, Pink DAC, Jones JE, Bending GD. 2015. Growth and

1297   nutritional responses to arbuscular mycorrhizal fungi are dependent on onion genotype and

1298   fungal species. *Biology and Fertility of Soils* 51: 801–813.

1299

1300   Tedersoo L, Magurno F, Alkahtani S, Mikryukov V. 2024. Phylogenetic classification of

1301   arbuscular mycorrhizal fungi: new species and higher-ranking taxa in Glomeromycota and

1302   Mucoromycota (class Endogonomycetes). *MycoKeys* 107: 249-271.

1303

1304 Tedersoo L, Sánchez-Ramírez S, Kõljalg U, Bahram M, Döring M, Schigel D, May T,

1305 Ryberg M, Abarenkov K. 2018. High-level classification of the Fungi and a tool for

1306 evolutionary ecological analyses. *Fungal Diversity* 90: 135–159.

1307

1308 Thippayarugs S, Bansal M, Abbott LK. 1999. Morphology and infectivity of fine endophyte

1309 in a Mediterranean environment. *Mycological Research* 103(11): 1369-1379.

1310

1311 Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, Charron P, Duensing

1312 N, Frei dit Frey N, Gianinazzi-Pearson V et al. 2013. Genome of an arbuscular mycorrhizal

1313 fungus provides insight into the oldest plant symbiosis. *Proceedings of the National Academy*

1314 *of Sciences* 110(50):20117-20122.

1315

1316 Torres-Cruz TJ, Billingsley-Tobias TL, Almatruk M, Hesse CN, Kuske CR, Desirò A,

1317 Niccolò-Benucci GM, Bonito G, Stajich JE, Dunlap C et al. 2017. *Bifiguratus adelaidae*, gen.

1318 et sp. nov., a new member of Mucoromycotina in endophytic and soil-dwelling habitats.

1319 *Mycologia* 222(1): 511-525.

1320

1321 Vierheilig H, Coughlan AP, Wyss U, Piche Y. 1998. Ink and vinegar, a simple staining

1322 technique for arbuscular mycorrhizal fungi. *Applied and Environmental Microbiology*

1323 64(12):5004-5007.

1324

1325 Vierstraete AR, Braeckman BP. 2022. Amplicon_sorter: A tool for reference-free amplicon

1326 sorting based on sequence similarity and for building consensus sequences. *Ecology and*

1327 *Evolution* 12(3): e8603.

1328

1329 Walker C, Gollotte A, Redecker D. 2018. A new genus, Planticonsortium (Mucoromycotina),

1330 and new combination (P. tenue), for the fine root endophyte, *Glomus tenue* (basionym

1331 *Rhizophagus tenuis*). *Mycorrhiza* 12(3): e8603.

1332

1333 Wang S, Zhang X, Zhang Z, Chen Y, Tian Q, Zeng D, Xu M, Wang Y, Dong S, Ma Z et al.

1334 2023. Fusarium-produced vitamin B6 promotes the evasion of soybean resistance by

1335 *Phytophthora sojae*. *Journal of Integrative Plant Biology* 65(9): 2204-2217.

1336

1337   West PT, Probst AJ, Grigoriev IV, Thomas BC, Banfield JF. 2018. Genome-reconstruction

1338   for eukaryotes from complex natural microbial communities. *Genome Research* 28(4):569-

1339   580.

1340

1341   Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualisation of de novo

1342   genome assemblies. *Bioinformatics* 31(20): 3350-3352.

1343

1344   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

1345   ISBN 978-3-319-24277-4

1346

1347   Wurzbacher C, Larsson E, Johan Bengtsson-Palme, Van den Wyngaert S, Svantesson S,

1348   Kristiansson E, Kagami M, Nilsson HR. 2018. Introducing ribosomal tandem repeat

1349   barcoding for fungi. *Molecular Ecology Resources* 19(1):118-127.

1350

1351   Yamamoto K, Degawa Y, Yamada A. 2020. Taxonomic study of Endogonaceae in the

1352   Japanese islands: New species of *Endogone*, *Jimgerdemannia*, and *Vinositunica*, gen. nov..

1353   *Mycologia* 112(2): 309-328.

1354

1355   Yang C, Chowdhury D, Zhang Z, Cheung WK, Lu A, Bian Z, Zhang L. 2021. A review of

1356   computational tools for generating metagenome-assembled genomes from metagenomic

1357   sequencing data. *Computational and Structural Biotechnology Journal* 19: 6301-6314.

1358

1359   Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. 2017. ggtree: an r package for visualization and

1360   annotation of phylogenetic trees with their covariates and other associated data. *Methods in*

1361   *Ecology and Evolution* 8(1): 28-36.

1362

1363   Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018.

1364   dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids*

1365   *Research* 46: 95-101.

1366

1367 Figure 1.

1379

1380

Figure 1. Colonisation by arbuscular mycorrhizal fungi (AMF) with fine root endophyte

(FRE) morphology in the roots of *Trifolium subterraneum* from: (A, B, C) the experiment

from which roots were chosen for metagenomic sequencing and (D) roots enriched in FRE

which were used as the reference 18S rRNA sequence (FRE 4583) of Mucoromycotina-AMF

(Orchard et al. (2017); experiment 1). Evident are the characteristic thin hyphae of FRE

morphology, as well as entry points (e), several cells filled by an arbuscule (a) which also

contained lipid droplets (l), terminal vesicles (tv) and intercalary vesicles (icv). (Photo credit:

Payman Abbaszadeh-Dahaji A, B, C; Suzanne Orchard D)

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401   Figure 2.

1402

1403

Figure 2. Phylogenetic assessment of the Mucoromycotina-AMF reference sequence FRE

4583 and 'consensus_all_seqs_0_16', an 18S rRNA gene sequence associated with the MAG,

and related 18S rRNA sequences amplified from root samples (consensus_all_seqs_0122,

0105 and 0122), in the context of the Tedersoo et al. (2024) *Endogonomycetes* SSU

alignment. The phylogenetic tree was generated using IQ-TREE, the model "TIM2+F+R5",

1000 ultrafast bootstrap replicates, and approximate Likelihood Ratio Tests. Branch support

is presented as a percentage (aLTR support/bootstrap support).

1411

1412

1413

1414

1415

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425 Figure 3.

1426

Phylum: Ascomycota (79) Basidiomycota (94) Mucoromycota (101) Unknown (1)

Ecology:
- Ectomycorrhiza (42)
- Endophyte (9)
- Ericoid Mycorrhiza (4)
- G-AMF (12)
- Lichenic symbiont (1)
- Mixed (23)
- Orchid Mycorrhiza (3)
- Parasite (5)
- Pathogen (29)
- Saprotroph (146)
- Unknown (1)

Protein marker hits: 340 360 380 400 420

BUSCO hits: 50 60 70 80 90 100

1453

1454 Figure 3. Phylogenetic placement of the MAG based upon a set of 434 protein markers

1455 identified by the JGI as pan-fungal markers for inferring high-level fungal phylogeny.

1456 Associated ecological group, protein marker hits and BUSCO hits are shown.

1457

1458

1459   Figure 4.

1460

1461

Figure 4. Number of CAZyme, lipase, protease, and transporter predicted proteins per

genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points

represent the phylogenetically closest relatives to the MAG: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shape within the G-AMF distinguishes between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test (p < 0.01) followed by a post-hoc Dunn's test (p < 0.05).

1469

1470

1471   Figure 5.

1472

1473

Figure 5. Number of plant cell wall degrading enzyme (PCWDE) and microbial cell wall

degrading enzyme (MCWDE) predicted proteins per genome distributed by ecological

lifestyle. Lifestyle is colour coded. Coloured data points represent the phylogenetically

closest relatives to the MAG: yellow for 'Endsp1', red for 'Jimlac1', green for 'Bifad1', and

purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'. Data point shape within the G-AMF

distinguishes between the Diversisporales (star) and Glomerales (triangle). Different letters

indicate significant differences between lifestyles as determined by a Kruskal-Wallis test (p <

0.01) followed by a post-hoc Dunn's test (p < 0.05).

1482

Figure 6. Cellulose, hemicellulose, lignin and pectin degrading predicted proteins per genome

distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points represent

the phylogenetically closest relatives to the FRE genome: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shape within the G-AMF distinguishes between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test ($p < 0.01$) followed by a post-hoc Dunn's test ($p < 0.05$).

1493

1494

1495   Figure 7.

1496

1497

Figure 7. Chitin, glycan, mannan, and peptidoglycan degrading predicted proteins per

genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points

represent the phylogenetically closest relatives to the MAG: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shape within the G-AMF distinguishes between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test (p = 0.012 for chitin, p = 0.024 for peptidoglycan, and p

< 0.01 for the remainder) followed by a post-hoc Dunn's test (p < 0.05).

1506

1507   Figure 8.

1508

1509

Figure 8: Phosphate, calcium, copper, nitrate, ammonium and glycine transporters per

genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points

represent the phylogenetically closest relatives to the MAG: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shape within the G-AMF distinguishes between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test (p = 0.028 for phosphate, p = 0.62 for copper, p = 0.041

for nitrate, and p < 0.01 for the remainder) followed by a post-hoc Dunn's test (p < 0.05).