Comparative genomic analysis of a metagenome-assembled genome reveals distinctive

symbiotic traits in a Mucoromycotina fine root endophyte arbuscular mycorrhizal

fungus

Joshua Cole[1*], Sébastien Raguideau[2], Payman Abbaszadeh-Dahaji[3,4], Sally Hilton[1], George

Muscatt[1], Ryan M. Mushinski[1], R. Henrik Nilsson[5], Megan H. Ryan[3], Christopher Quince[2,6,7]

and Gary D. Bending[1*]

[1]School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK

[2]Earlham Institute, Norwich Research Park, Colney Lane, Norwich NR4 7UZ, UK

[3]UWA School of Agriculture and Environment, University of Western Australia, Crawley,

WA, 6009, Australia

[4]Department of Soil Sciences, Faculty of Agricultural Science, Vali-e-Asr University of

Rafsanjan, Rafsanjan, Iran

[5]University of Gothenburg, Department of Biological and Environmental Sciences,

Gothenburg Global Biodiversity Centre, University of Gothenburg, Box 461, 405 30,

Göteborg, Sweden

[6]Gut Microbes and Health, Quadram Institute, Norwich NR4 7UQ, UK

[7]School of Biological Sciences, University of East Anglia, Norwich NR4 7T, UK

*Corresponding authors: Joshua.M.Cole@warwick.ac.uk; gary.bending@warwick.ac.uk

Abstract

Background

Recent evidence shows that arbuscular mycorrhizal (AM) symbiosis is established by two

distinct fungal groups, with the distinctive 'fine root endophyte' morphotype formed by fungi

from the sub-phylum Mucoromycotina rather than the sub-phylum Glomeromycotina. While

Mucoromycotina AM fungi are globally distributed, there is currently no understanding of

the genomic basis for their symbiosis or how this symbiosis compares to that of other

mycorrhizal symbionts.

Results

We used culture-independent metagenome sequencing to assemble and characterise the

metagenome-assembled genome (MAG) of a putative fine root endophyte, which we show

belonged to the family Planticonsortiaceae within the order Densosporales. The MAG shares

key traits with Glomeromycotina fungi, which indicate obligate biotrophy, including the

absence of fatty acid and thiamine biosynthesis pathways, limited enzymatic abilities to

degrade plant cell walls, and a high abundance of calcium transporters. In contrast to

Glomeromycotina fungi, it exhibits a higher capacity for degradation of microbial cell walls,

a complete cellulose degradation pathway, low abundances of copper, nitrate and ammonium

transporters, and a complete pathway for vitamin B6 biosynthesis.

Conclusion

These differences highlight the potential for contrasting interactions between

Mucoromycotina and Glomeromycotina AM fungi with their host plant and the environment

which could support niche differentiation and complementary ecological functions.

Key Words

Mucoromycotina, Fine Root Endophyte, Metagenome-Assembled Genome, Arbuscular

Mycorrhizal Fungi, Planticonsortiaceae

Background

Mutualistic symbioses with microbes are exhibited by the overwhelming majority of plant species [1] with mycorrhizas by far the most widespread and well-described association [2]. While mycorrhizal symbioses have evolved multiple times across plant and fungal taxa, they share the common characteristic of the fungus trading nutrients assimilated from the soil, particularly phosphorus and nitrogen, in exchange for carbon from the plant [3]. Additionally, mycorrhizal associations can confer a range of secondary benefits to the plant host, including disease resistance and stress tolerance [4]. Because of these profound impacts on plants, mycorrhizal associations are considered to be major determinants of ecosystem productivity and diversity and to play key roles in directing terrestrial biogeochemical cycling processes. There are four main groups of mycorrhizal associations: arbuscular (AM), ecto- (ECM), ericoid (ERM), and orchid (ORM) mycorrhizas [5], which are distinguished by a number of features including the specific plant and fungal taxa involved, the ecosystems they inhabit, the way in which the fungus colonises and grows within root tissues, and the functional traits of the fungi involved [6].

As a function of time and niche specificity, mycorrhizal symbionts appear to have evolved along a saprotrophy to symbiosis continuum [5, 7]. This directional transition from free-living saprotrophs to obligate symbionts is believed to be driven by the provision of carbohydrates and vitamins from plants in nutrient-poor environments and the subsequent evolutionary success of a symbiotic lifestyle. As a result, key genomic signatures associated with this transition can be identified across this continuum [5]. As fungal symbionts shift away from primarily saprotrophic lifestyles to mutualisms, their genomes typically possess a reduced repertoire of the enzymatic machinery involved in the degradation of soil organic matter (SOM) and plant cell walls [5, 8, 9]. The two youngest mycorrhizal symbioses, ERM and ORM, with origins c 75-90 and 112 mya, respectively [6, 8], involve symbionts with genomic repertoires capable of simultaneously facilitating a saprotrophic and symbiotic lifestyle. However, fungi forming the older AM (origin 450 mya, [6]) and ECM symbioses (origin c 200 mya) possess a much more fine-tuned enzymatic toolkit, noted for a reduced suite of plant cell wall degrading enzymes (PCWDE) along with the presence of specific genes involved in regulating nutrient transport and host plant signalling and metabolism [5, 8].

AM are considered the most common and widespread plant-microbe symbiosis on Earth [10-12], and their presence in fossil roots of the earliest land plants suggests that they were a

significant factor in the evolution of land plants. Fungi forming AM have traditionally been
placed into the Mucoromycota sub-phylum Glomeromycotina. These fungi are obligate
symbionts; as fatty acid auxotrophs they depend on the exchange of plant-derived fatty acids
[13, 14]. As a result of long-term host dependence, arbuscular mycorrhizal fungi (AMF)
present hallmark traits such as limited ability to break down cellulose, biosynthesise
thiamine, or metabolise vitamin B6 derivatives. The genomic inability to express these
hallmark traits represent Missing Glomeromycotan Core Genes (MGCGs) [15-17].

Until recently, all AMF were believed to belong to the Glomeromycotina (G-AMF) [12]. It
has long been recognised that there are two broad groups of arbuscular mycorrhiza based on
morphology, with the 'coarse endophyte' G-AMF being the focus of the vast majority of
research on the biology and ecology of AMF [18]. Glomeromycotina comprises four orders,
the Glomerales, Diversisporales, Paraglomerales, and Archaeosporales, which likely
diversified when plants evolved onto land [4, 19, 20]. The second morphological group of
'fine root endophytes (FRE)' arbuscular fungi are distinguished from coarse endophytes by
their small spores, fan-shaped or palmate entry points, diverging thin hyphae (<2 µm
diameter) which in intercellular spaces often form hyphal ropes, as well as intercalary and
terminal 'vesicles'', approximately 5–10 µm in diameter [18, 21-23]. Interestingly, the
arbuscules of FRE appear very similar in structure to those of coarse endophytes [24]. Recent
evidence has placed FRE within the sub-phylum Mucoromycotina (M-AMF) [18], indicating
that AM symbioses may have evolved independently several times, as is the case for
ectomycorrhizal symbioses [25].

To date only one species of M-AMF, Planticonsortium tenue, has been described [23].
However, similarly to G-AMF, root systems are colonised by diverse Mucoromycotina taxa,
communities of which are typically comprised of a small number of abundant and common
taxa and a high diversity of rare low-abundance taxa [26-28]. Furthermore, distinctive
morphologies of FRE have been recognised [21] based on a variety of characteristics,
including the roughness, diameter, and branching of hyphae and the size and shape of
vesicles. While fungi within the Glomeromycotina are all assumed to be plant symbionts,
fungi residing within the Mucoromycotina also include saprotrophs and putative
ectomycorrhizal symbionts. Those Mucoromycotina sequences found associated with roots
and rhizoids of vascular and non-vascular plants appear to have similar phylogenetic ranges
and occur within clades alongside taxa with saprotrophic and putative ectomycorrhizal
lifestyles [28, 29].

M-AMF have been detected across multiple continents, spanning a wide range of ecosystems,

environmental conditions, and hosts, although in contrast to G-AMF they may be missing

from tropical and sub-tropical biomes, and may be favoured by agricultural management

practices [22, 27]. While the benefits of G-AMF to the host, particularly for phosphorus

nutrition, are well documented, the functional significance of M-AMF within ecosystems is

poorly understood [18, 24, 30]. G-AMF and M-AMF typically co-occur and intermingle

within root systems [18, 22, 31], but they appear to occupy distinct but overlapping

ecological niches [32], with their distribution at the landscape scale differentially determined

by temperature, pH, and plant richness [27]. Unlike G-AMF, nutrient availability was found

to be an important determinant of M-AMF abundance, which increased with fertility [27].

Furthermore, in contrast to G-AMF, experimental evidence suggests that M-AMF may favour

providing the host with access to soil nitrogen, indicating a potential complementary

functionality between the two arbuscular symbioses and an explanation for the parallel

evolution of symbiotic traits [33, 34].

Comparative genome analysis has emerged as a powerful tool to understand microbial

evolution, function, and ecology. However, these approaches largely rely on the availability

of cultured microbes or voucher specimens for whole genome sequencing (e.g. [8, 17, 25]).

Importantly, culturing microbes can prove time-consuming and problematic as microbial

populations have great complexity and many microbes have specific nutritional requirements

which cannot be met in culture. This is particularly the case for obligate biotrophs such as

symbionts which rely wholly or in-part on the host for their nutritional requirements [35]. As

obligate symbionts, G-AMF require a host plant to complete their lifecycle. Despite this, pure

cultures of G-AMF have been obtained using root organ cultures, and these have been used to

produce inoculum for genome sequencing [36]. The challenges in obtaining and maintaining

root cultures have meant that direct genome sequencing of G-AMF cultures has been limited

to date. Recently, asymbiotic mycelium and spore production of the G-AMF Rhizophagus

clarus has been achieved by supplementing its growth medium with palmitoleic acid, and this

could provide a boost to rates of culture-based sequencing [35]. An alternative approach to

obtain fungal genomes is sequencing of nuclei extracted from spores [37], although most G-

AMF taxa appear to sporulate infrequently, if at all [38].

M-AMF are also putatively obligate symbionts, but because they have only recently been

recognised as a distinct group of symbionts, culturing efforts lag behind those of G-AMF.

Recently, monoxenic cultures of a Mucoromycotina fungus associated with the clubmoss

Lycopodiella inundata have been obtained and shown to form FRE colonisation in a vascular

plant [33]. However, while genome sequencing and comparative genomic analysis of a

number of saprotrophic and putative ectomycorrhizal Mucoromycotina has been achieved

[17, 39], there is no understanding of the genome of M-AMF fungi and how their functional

repertoire and mutualistic signatures compare to those of G-AMF, other mycorrhizal

symbioses, and saprotrophic and putatively ectomycorrhizal Mucoromycotina. To further

complicate the matter, initial comparative analysis of the sister clades indicate that some

widely considered hallmark traits of mycorrhizal symbioses may instead represent ancestral

markers as opposed to adaptation [17].

Recently there has been great progress in the development of computational tools for the

culture-independent assembly of 'dark' microbial genomes from metagenome sequencing

datasets [40]. Such 'Metagenome Assembled Genomes' (MAGs) have been widely used to

identify the metabolic capabilities and ecological roles of specific prokaryotes in diverse

habitats [41]. MAG construction of eukaryotic microbes is more challenging because of a

range of issues including their large sizes, high genome heterozygosity, low abundance in

metagenome datasets, and a lack of well-annotated reference genomes for gene annotation

[42, 43]. As a result, few studies have constructed eukaryotic MAGs from metagenomes, and

these have typically been performed in low-diversity communities (e.g. [42, 44]). A further

particular issue with analysis of plant symbionts in situ is their intimate association with plant

tissue. This results in co-extraction of host DNA, increasing the sequencing depth required to

assemble symbiont genomes.

The aim of the current work was to use culture-independent metagenome sequencing to

assemble and characterise the genome of an FRE-forming M-AMF fungus for the first time

and compare its genomic repertoire to glomeromycotan fungi and other fungal symbionts.

Roots of Trifolium subterraneum were first enriched with FRE using a selective soil sieving

technique. Root samples were used for metagenome sequencing, from which a MAG of the

FRE was constructed. We compiled a database of 274 fungal genomes across the

Ascomycota, Basidiomycota and Mucoromycota, and assigned these to G-AMF, ECM, ERM,

OCM, saprotroph, pathogen, parasite, endophyte, lichenic, and mixed lifestyles. Comparative

analysis showed that the MAG shares key distinctive characteristics with G-AMF. This

includes overlaps in some MCGCs, presence of abundant calcium transporters, and an overall

reduced set of plant cell wall degrading enzymes. However, we found a relatively high

abundance of microbial cell wall degrading enzymes and evidence of complete cellulose

degradation capability, as well as low numbers of nitrogen transporters, suggesting a

distinctive niche relative to G-AMF and the other mycorrhizal lifestyles.

Materials and Methods

Soil collection and sample preparation

For the metagenome sequencing, plants with the root AMF community enriched in M-AMF

were produced following the methods of Orchard et al. (2017a). In brief, soil was collected

from the field studied by Orchard et al. (2016) and Orchard et al. (2017a) in the Peel Harvey

region in Western Australia, Australia: site coordinates were 32° 48 826'S, 115° 53346'E.

Further detail on the location, soil properties, enrichment process and mycorrhizal staining

protocol are available in the supplementary information.

DNA extraction, 18S rRNA gene amplicon sequencing and metagenome sequencing

Twelve samples were selected for 18S rRNA amplicon and metagenome sequencing on the

basis of predominance and abundance of M-AMF colonisation as well as quality and quantity

of DNA obtained (Table S1). DNA was extracted from root samples using the DNeasy

PowerSoilPro Kit (Qiagen) following the manufacturer's protocol. DNA concentrations were

measured using a Qubit 3.0 fluorimeter using the Qubit dsDNA high sensitivity assay kit

(Invitrogen) according to the manufacturer's protocol.

To determine whether the dilution technique had enriched M-AMF, we performed

community 18S rRNA amplicon analysis. Glomeromycotina and Mucoromycotina 18S rRNA

genes in root samples were amplified using AM-Sal-F and AMDGR primers which amplify a

220-base fragment [28]. PCR reactions, Illumina MiSeq sequencing, and bioinformatic

analysis were performed using the approaches outlined in Seeliger et al. (2024).

Long read nanopore amplicon sequencing of the nuclear ribosomal 18S rRNA, 5.8S, and

28S/LSU region of Mucoromycotina was performed on root samples to support construction

of the FRE MAG and to determine its phylogenetic relationships. PCR reactions were

conducted using AM-Sal-F in combination with RCA95m [45] using the recommended

Oxford Nanopore protocol, amplifying a 4.5 kb region. Further detail is available in the

supplementary information.

Fast5 files were base called and demultiplexed using the Guppy (2022) basecaller provided

by Oxford Nanopore Technologies. The sample was then processed using amplicon_sorter

v2022-03-28 [46] which aims to group amplicons at species or genus level (~95%), output

consensus sequences, and quantify the copy number of the representative consensus

sequences in the sample. This resulted in 42 consensus sequences, 5 of which were placed

into the Mucoromycotina.

To aid detection of M-AMF sequences in the metagenome, a reference 18S rRNA sequence

was obtained from DNA extracted from T. subterraneum root samples which were enriched

in M-AMF and contained minimal G-AMF. These samples were produced in a similar

manner to that described above and are presented in Orchard et al. (2017a) as "Experiment

1". On average, 90% of the total AMF colonisation in these roots was contributed by M-

AMF, presenting the first evidence linking FRE morphology to Mucoromycotina. The AM-

Sal-F and RCA95m primers were used to amplify a 4.5 kb region, as described above, which

was cloned using the TA cloning kit (ThermoFisher). Partial 18S rRNA analysis using AM-

Sal-F and AMDGR primers revealed a clone, FRE4583, whose sequence showed a 100%

match both to OTU4 which was associated with FRE morphology in the earlier Orchard et al.

(2017a) study and the BEG249 Planticonsortium tenue FRE culture described by Walker et

al. (2018) [27]. Furthermore, this and closely related sequences have been detected in

multiple studies reporting FRE morphology in plant roots [24, 28]. Primer walking was used

to sequence overlapping regions of clone FRE4583 until the full sequence was covered,

generating a 4.5 kb fragment.

Metagenome sequencing and assembly

The twelve root samples were used for Shotgun metagenomic sequencing using the HiSeq

150 paired end protocol to generate between 15 and 47 Gbp sequencing depth. Library

construction and sequencing was conducted using Illumina protocols by Novogene UK Ltd.

A collection of reference genomes for T. subterraneum was built from downloading all

available sequences from the NCBI at the date of the work (2022). This collection was then

used to filter samples DNA using bwa mem v0.7.18-r1243-dirty [47] and samtools v1.10

[48]. To determine FRE depth and breadth of coverage, the filtered metagenome reads were

mapped, again using bwa mem and samtools, against the FRE4583 ribosomal operon and the

resulting alignment files processed with custom Python scripts. To be sure that a high mean

coverage was not an artefact of conserved regions, breadth of coverage was visually

inspected. Sample P72 was predicted to have the highest Mucoromycotina abundance,

possessing the majority of the operon with a coverage depth of over 100 (Fig. S1); the

missing regions are likely due to repeats as we filtered ambiguous mappings. We therefore

focused on this sample in our efforts to reconstruct an M-AMF genome, assembling it using

the SPAdes assembler v3.14.1 [49] with option -meta. This produced a catalogue of contigs

with an N50 of 446 and total length of 1.3 Gbp. We then computed coverage depth profiles

for these contigs across all samples by mapping reads with bwa mem, samtools, and bedtools

v2.30.0 [50]. This improves the discriminatory power of genome binning through multi-

sample coverages whilst keeping the best quality single sample assemblies. The contigs were

then binned using these coverage profiles and composition with the binning software

CONCOCT v1.1.0 [51] and a minimum contig length of 1000 bp.

The putative M-AMF genome was identified based on a consensus of multiple lines of

evidence, see Fig. S2. Initially, we searched for the reference M-AMF ribosomal sequence

FRE4583 in our dataset by mapping it to the assembly. A total of 12 unitigs were found with

high similarity (>98%) and spanning 75% of the operon without overlap. Since the assembly

is fragmented, we decided to explore the assembly graph and more particularly the context of

these small unitigs. Out of the 12, only 4 had a context, i.e., they were linked to other unitigs

in the assembly graph. The idea behind this approach is that ignoring any miss-assemblies, all

genomes have a distinct path in the assembly graph and in particular the M-AMF genome

should include a path which passes through those unitigs labelled as ribosomal sequences

thus allowing identification of M-AMF. We could not assemble a complete FRE4583 operon

in a single contig, which is likely due to strain diversity. Using custom scripts available on

GitHub (https://github.com/Sebastien-Raguideau/Graph-tools) we extracted a subgraph

encompassing all 10 Kbp around these 4 unitigs, and then visualised all contigs binned by

CONCOCT on this subgraph using Bandage v0.8.1 [52], see Fig. S3. In this figure, the 4

unitigs are in red, un-binned unitigs are in gray, and all other colored unitigs are related to

different CONCOCT bins. While this figure looks complex, it only corresponds to 0.05% of

the whole assembly and refers to a limited radius of 10 kb nucleotides around the ribosomal

unitigs, showing a clear proximity of multiple numbered bins.

A total of 12 bins were found in close proximity to M-AMF unitigs. However, on assessing

quality, only bin 17 was found to have any marker genes, giving it a completion of 83.7%

using the BUSCO v5.2.2 [53] general fungi_odb10 scheme with a size of 37.37 MBp (Table

S2). In particular, while contigs from bin 70 were directly co-located with FRE ribosomal

sequences, this bin did not possess any marker genes and was distinctly different in terms of

coverage from bin 17. It may be that bin 70's location in the graph is due to a miss-assembly,

or it could be that this is due to strain diversity and that bin 70 is strain-specific part of the

pan-genome of the FRE which would result in it having a different coverage profile to the

core genome represented by bin 17.

To provide additional QC for this putative FRE MAG (bin 17) we used varScan [54] to detect

SNVs on all the contigs assigned to it. In Fig. S4 we show the frequencies of these SNVs, and

in Fig. S5 their density on contigs. There are a large number of SNVs throughout the genome

(Fig. S5) but these are distributed evenly in frequency suggesting that they are likely

generated by strain diversity rather than multiple haplotypes. We also computed the

distribution of coverage depths of all ORFs predicted by prodigal (Fig. S6). While not

sufficient to correctly detect all ORF from a eukaryotic genome, it still suggests that most

genes are in a single copy despite the strain diversity.

We used phyloFlash v3.4.2 [55] to quantify abundances of all the SSU rRNAs in the SILVA

database [56] and also our 42 target nanopore amplicon 18S rRNA consensus sequences.

Filtering any taxa with less than 10 reads mapping across all samples yielded a total of 251

entries including the 42 consensus sequences (Fig. S7). These were normalised by the total

sequencing depth of each sample. To compare to these, the coverage depth of bin 17 was

calculated for each sample by taking the sum of all nucleotides mapping to all contigs of bin

17 and dividing by the total length of the contigs, followed by normalisation as above. Using

Pearson's correlation, the coverage of bin 17 was compared to the coverage of all 251 taxa

detected with phyloFlash. The strongest correlation (r = 0.97 and an adjusted p value of 4.5e-

11, Fig. S8) was with one of the FRE consensus sequences (consensus_all_seqs_0_16), thus

enabling us to provide finer scale phylogenetic information for bin 17, unambiguously

linking it to this M-AMF sequence. The regression coefficient of 63 between coverage

profiles (consensus_all_seqs_0_16 and bin 17) provides an indication of the number of

ribosomal operons in the genome, a quantity consistent with other fungi [57].

Reference genome access and ecological lifestyle annotation

To compare the MAG to known fungi, a database of reference fungal genomes was

constructed from JGI MycoCosm (Table S3). The fungal species, representing the three

phyla—Ascomycota, Basidiomycota, and Mucoromycota—were chosen based on two

criteria. First, fungal genomes which had been included in previous comparative genomics

analyses of mycorrhizal fungi [8, 25] were included since the ecology of these fungi had

already been curated. Secondly, the database was supplemented with Mucoromycota

genomes available in MycoCosm

(https://mycocosm.jgi.doe.gov/mucoromycota/mucoromycota.info.html) in May 2023. The

associated predicted fungal proteomes were accessed and downloaded in May 2023. Fungal

genomes were categorised according to their ecological lifestyle (Table S3). We assigned 146

genomes as saprotrophic, 42 as ECM, 29 as pathogenic, 23 as mixed lifestyle, 12 as G-AMF,

9 as endophytic, 5 as parasitic, 4 as ERM, 3 as OCM, and 1 as lichenic. Within this total, we

included five Mucoromycotina genomes. Full descriptions of the ecological assignment

process and the Mucoromycotina genomes are available in the supplementary materials.

Phylogenetic assessment

Phylogenetic assessment of FRE4583 and those consensus sequences from root samples

which were assigned to the order Endogonales (including all_seqs_0_16) was carried out

through 18S rRNA gene alignment. These sequences were added to the Endogonomycetes

18S rRNA alignment from Tedersoo et al. (2024) using MAFFT v7.525 [58]. The resulting

alignment was manually checked and refined in MEGA11 v11.0.13 [59]. The best fitting

model for our alignment was selected using IQ-TREE v2.3.6 and the option "-MFP" [60, 61].

A phylogenetic tree was generated using the model "TIM2+F+R5" with 1000 approximate

likelihood ratio tests and 1000 ultrafast bootstraps [62, 61]. The tree was subsequently

visualised in FigTree v1.4.4 [63].

In addition to the 18S rRNA alignment, fungal phylogeny was inferred from genome

sequences using a concatenated protein marker alignment. A set of 434 protein markers

identified by the JGI as pan-fungal markers for inferring high-level fungal phylogeny was

used to compare the phylogenetic placement of the MAG and all the reference fungal

genomes within our curated database. To identify these marker proteins, an HMM set was

downloaded from

https://github.com/1KFG/Phylogenomics_HMMs/tree/master/HMM/JGI_1086.

Phylogenomic markers were extracted from fungal proteomes using the above HMM, and a

subsequent phylogenetic tree was built using PHYling v0.9.0 (available at:

https://github.com/stajichlab/PHYling) under default settings. The tree was then visualised in

R using ggtree v3.6.2 [64].

Annotation of MAG and reference genomes

Our comparative analyses focused on Carbohydrate-Active enZymes (CAZymes), lipases,

proteases, transporters, and small-secreted proteins. Full details on the methodology used to

identify these groups within the genomes is available in the supplementary information.

Comparative genomics analyses

All comparative genomics analyses were conducted using R v4.2.2 [65], and all

visualisations were generated with ggplot2 v3.4.0 [66]. Functional annotation of EC numbers

was assigned using DeepECtransformer v1.0 [67] and the ECRECer web platform [68].

Results

Description of FRE in the roots

The appearance of the FRE in the roots of the twelve samples chosen for metagenomic

sequencing was consistent with the description of the genus *Planticonsortium* by Walker et

al. (2018) into which they placed *Glomus tenue* (previously generically referred to in the

literature as fine endophyte or fine root endophyte [17] (see photographs in: Fig. 1 herein;

Figure 2 and Figure S1 in Orchard et al., 2017a and Thippayarugs et al. (1999)). In particular,

entry points were characterised by structures with "feather-like, fan-shaped, palmate or

multilobed with finger-like projections" [22] which then led to fine, diverging hyphae in

cortex cells and, often, plentiful finely-branched arbuscules. Small intercalary vesicles and

terminal vesicles were also common. Droplets of lipids were often evident within hyphae and

arbuscules. In some instance, colonisation in the cortex was too dense to clearly distinguish

the structure of individual arbuscules.

Figure 1. Colonisation by arbuscular mycorrhizal fungi (AMF) with fine root endophyte

(FRE) morphology in the roots of Trifolium subterraneum from: (A, B, C) the experiment

from which roots were chosen for metagenomic sequencing and (D) roots enriched in FRE

which were used as the reference 18S rRNA sequence (FRE 4583) of Mucoromycotina-AMF

(Orchard et al. (2017); experiment 1). Evident are the characteristic thin hyphae of FRE

morphology, as well as entry points (e), several cells filled by an arbuscule (a) which also

contained lipid droplets (l), terminal vesicles (tv) and intercalary vesicles (icv). (Photo credit:

Payman Abbaszadeh-Dahaji A, B, C; Suzanne Orchard D)

In the samples used for amplicon and metagenome sequencing, the percentage of total root

length colonised by AM ranged between 26.8 and 72.5% for AMF with FRE morphology and

0 to 16.3% for G-AMF, producing a proportion of AMF colonisation by FRE between 62.2

and 100%. 18S rRNA Miseq sequencing of Glomeromycotina and Mucoromycotina

communities in root samples revealed that the proportion of Endogonales amplicons ranged

between 25 and 82% (Table S1).

Phylogeny and Features of Fungal Genomes

In this study, we constructed a putative FRE genome, referred to as the 'MAG'. This MAG

was identified as a putative M-AMF through two lines of evidence. First, the contig bin was

colocated on the assembly graph with unitigs that we could assign to the 18S rRNA gene of a

known M-AMF reference (FRE4583). Secondly, its abundance correlated across the 12

samples in the study with an 18S rRNA sequence 'consensus_all_seqs_0_16', which we

showed to be very closely related to FRE4583 (Fig. 2). The bin itself did not contain a

complete 18S rRNA sequence. This is unsurprising as this gene is well known for failing to

assemble, largely due to it being comprised of strongly conserved regions that repeat

throughout the genome, failing to assemble into substantial contigs. The MAG measured

37.37 Mbp with a genome coverage varying between 1x and 568 depending on the sample.

The N50 was found to be 4928 with a GC content of 41.1%. Finally, its completion was

assessed to be 83.7% with only 1.5% contamination (Table S2). We identified a total of

10,872 proteins. Of these proteins, we were able to assign 3,358 proteins with a functional

annotation, comprising 218 CAZymes, 121 lipases, 280 proteases, 13 small secreted proteins,

and 2,726 transporters.

While we acknowledge that the revised phylogeny of the Mucoromycotina subphylum

produced in Tedersoo et al. (2024) lacks type specimens and is grounded in part on dark taxa,

as a poorly captured branch of the fungal tree of life, we deem this an appropriate basis to

apply the phylogeny of our MAG. 18S rRNA analysis placed consensus_all_seqs_0_16, and

by proxy the MAG, within the order Densosporales of the Mucoromycotina, corresponding

with the Densosporaceae sensu Desirò et al. (2017) (Fig. 2). This order is associated with

FRE morphology [22, 39]. Further to this, strong support values allow us to confidently place

the MAG in the newly proposed family Planticonsortiaceae [69]. Within this family, the

MAG was placed with very high similarity to the known M-AMF reference sequence,

FRE4583 (Fig. 2). The MAG was closely related to three other Mucoromycotina 18S rRNA

sequences obtained from the root samples (consensus_all_seqs 0105, 0122, and 0260) and to

environmental sequences obtained from the thalli of early diverging plant sequences, e.g.,

MH174585 and MH174471. The wider Planticonsortiaceae clade included further sequences

from the thalli of non-vascular plants, sequences not annotated beyond the phylum level

associated with soil from across Europe (EUK sequences), and one sequence associated with

the rhizosphere of Populus tremuloides (EF23703), a known arbuscular mycorrhizal host

[70]. Phylogenetic analysis therefore placed the MAG apart from other members of the

Mucoromycotina with putative ecology from the order Endogonales, including the putative

ectomycorrhizal Mucoromycotina member Jimgerdemannia spp. [29] and the saprotrophic

Endogone sp. and Vinositunica sp. which have a putative saprotrophic or ectomycorrhizal

lifestyle [71].

Figure 2: Phylogenetic assessment of the Mucoromycotina-AMF reference sequence FRE

4583 and 'consensus_all_seqs_0_16', an 18S rRNA gene sequence associated with the MAG,

and related 18S rRNA sequences amplified from root samples (consensus_all_seqs_0122,

0105 and 0122, highlighted red), in the context of the Tedersoo et al. (2024)

Endogonomycetes SSU alignment, showing the family Planticonsortiaceae. The phylogenetic

tree was generated using IQ-TREE, the model "TIM2+F+R5", 1000 ultrafast bootstrap

replicates, and approximate Likelihood Ratio Tests. Branch support is presented as a

percentage (aLTR support/bootstrap support).

Similarly, the protein marker alignment placed the MAG within the Mucoromycota (Fig. 3).

In our phylogenetic analysis, the MAG forms a sister group to all other Mucoromycotina

species except for Calcarisporiella. This placement suggests that the MAG represents an

early-diverging lineage within the Mucoromycotina. Due to the lack of available genomes

from the order Densosporales, we cannot definitively assign the MAG to this order, but its

basal position indicates it may be related to early diverging Mucoromycotina lineages,

possibly including the Densosporales.

Figure 3: Phylogenetic placement of the MAG (shown here as Candidate FRE) based upon a set of 434 protein markers identified by the JGI as pan-fungal markers for inferring high-level fungal phylogeny. Associated ecological group, protein marker hits and BUSCO hits are shown.

Genomic Profile of the MAG

To contextualize the gene content of the MAG, we compared the CAZyme, protease, lipase, and transporter profiles to genomes within the reference dataset that represented the saprotrophy-symbiont continuum. Due to the MAG's early-diverging position within the Mucoromycotina (Fig. 3), and the absence of available genomes from some lineages such as the Densosporales, we highlighted the inclusion of five available genomes from the Endogonales for comparative purposes (Fig. 4-8). While the 18S rRNA and genome-wide analyses indicate that the MAG is a sister group to other Mucoromycotina species (excluding Calcarisporiella), the Endogonales genomes provide the closest available genomic context within this subphylum. This comparison allows us to explore potential functional attributes and evolutionary relationships within the Mucoromycotina. The ecological lifestyle had a significant effect on genome content across all four categories ($p < 0.01$) (Fig. 4). For the CAZymes, the ecological lifestyles fell into two groups—G-AMF and ECM, ERM and OCM —with SAP bridging the two (Fig. 4a). The MAG contained a low number of CAZymes (218), which was similar to the mean of the ECM (240), but slightly higher than that of the G-AMF (175). A similar number of CAZymes were found in the MAG compared to Endogonales genomes (149 to 208). The ERM genomes contained by far the largest median number of CAZymes across the lifestyles (629). Only the ERM possessed a significantly ($p < 0.01$) larger number of lipases than the other ecological lifestyles (Fig. 4b). The MAG possessed 121 lipases, lower than the means of the G-AMF (189), ECM (238), and SAP (263), OCM (268). Similarly, the Endogonales genomes all contained low numbers of lipases (72 to 132). The ERM genomes had significantly ($p < 0.01$) higher mean numbers of proteases per genome (457) than the other lifestyles (Fig. 4c). The MAG had similar numbers of proteases (280) compared to G-AMF (305), ECM (286), SAP (271), and to the Endogonales genomes (229 to 298). The G-AMF and ERM lifestyles had significantly higher ($p < 0.05$) numbers of transporters per genome relative to the other genomes. The MAG (2726) had a substantially lower number of transporters compared to G-AMF (3900); however, it contained more than the Endogonales genomes (1932 to 2388) that sat in the lower end of the dataset (Fig. 4d). Across the dataset, the saprotrophs contained an especially high degree of intra-group variability in numbers of all four profiles compared to the other ecological lifestyles (Fig. 4). The number of CAZymes, lipases, proteases, and transporters in the MAG were also comparable to the endophyte, parasite pathogen, and mixed lifestyles

(Fig. S9). The MAG contained 13 small secreted proteins, a middling amount for the dataset

(Fig. S9).

Figure 4: Number of CAZyme, lipase, protease, and transporter predicted proteins per

genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points

represent selected Mucoromycotina species for comparison: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shape within the G-AMF distinguishes between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test ($p < 0.01$) followed by a post-hoc Dunn's test ($p < 0.05$).

CAZyme Profile

CAZyme proteins associated with the degradation of plant and microbial cell walls were

identified to investigate the saprotrophic repertoire of the MAG, in the context of the

mycorrhizal saprotrophy-symbiont continuum. The ERM (252) and OCM (215) genomes

possessed a significantly higher (p < 0.01) mean number of PCWDEs than the G-AMF (32),

ECM (59), and SAP (77) genomes (Fig. 5a). The MAG contained a similar number of

PCWDE (37) as the G-AMF genomes and the genomes of the Endogonales (33 to 44). ERM

genomes contained a significantly (p < 0.01) higher mean number of MCWDE (109) than the

saprotrophs and other mycorrhizal lifestyles (24 to 59) (Fig. 5b). The G-AMF genomes had a

significantly (p < 0.05) lower number of MWCDE (24) compared to the mean ECM (45),

OCM (59), and SAP (48) genomes. The MAG had 53 MCWDE, higher than the median

number of all other lifestyles except the ERM (Fig. 5b, Fig. S10). A similar number of

MCWDE was detected across the Endogonales genomes (36 to 53).

a)

b)

Figure 5: Number of plant cell wall degrading enzyme (PCWDE) and microbial cell wall
degrading enzyme (MCWDE) predicted proteins per genome distributed by ecological
lifestyle. Lifestyle is colour coded. Coloured data points represent selected Mucoromycotina
species for comparison: yellow for 'Endsp1', red for 'Jimlac1', green for 'Bifad1', and purple
for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'. Data point shapes within the G-AMF
distinguish between the Diversisporales (star) and Glomerales (triangle). Different letters
indicate significant differences between lifestyles as determined by a Kruskal-Wallis test
(p < 0.01) followed by a post-hoc Dunn's test (p < 0.05).

Patterns of PCWDE distribution across the mycorrhizal associations were similar (Fig. 6).
ERM and OCM genomes had significantly (p < 0.01) higher mean numbers of cellulases,

hemicellulases, and pectinases than the G-AMF and ECM genomes. Relative to the mean G-

AMF, the MAG had notably higher numbers of cellulases (18 compared to 8), but similar

numbers of hemicellulases (11 compared to 7) and pectinases (4 compared 2). The

Endogonales genomes had similar numbers of cellulases (11 to 13), hemicellulases (10 to

16), and pectinases (4 to 12) as the MAG. The numbers of lignin degrading enzymes were

significantly higher in the ERM (31, $p < 0.01$) relative to the other mycorrhizal lifestyles,

which all possessed similar numbers: ECM (13), G-AMF (15), OCM (9), and SAP (12).

Notably the MAG had a very low number of lignin degrading enzymes (4) compared to all

other lifestyles, this trend was consistent across the Endogonales genomes (2 to 8).

Figure 6: Number of cellulose, hemicellulose, lignin, and pectin degrading predicted proteins

per genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points

represent selected Mucoromycotina species for comparison: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shapes within the G-AMF distinguish between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test ($p < 0.01$) followed by a post-hoc Dunn's test

($p < 0.05$).

Cellulases in the MAG genome comprised 8 families, with 6 copies of AA9 being the most

abundant (Table S4). The Endogonales genomes contained an average of 6 cellulase families,

with GH9 being the most abundant in all four genomes, of which the MAG contained 3.

However, not a single copy of AA9 was identified within the Endogonales genomes (Table

S4). The G-AMF genomes only contained an average of 3 families, with the Glomerales

containing a mean of just 6 cellulases. However, the Diversisporales contained a much higher

number of cellulases (17 and 12), including 7 and 6 of AA9 and 6 and 3 copies of AA3,

respectively (Table S4). The MAG contained a full suite of cellulose degradation enzymes:

Endoglucanase, β-glucosidase, and exoglucanase (Table S5). However, with the absence of

β-glucosidases in all G-AMF genomes and only two containing a single copy of

exoglucanase, none of the G-AMF genomes possessed the full cellulose degradation pathway

(Table S5). Hemicellulases in the MAG comprised the majority of the remaining PCWDEs,

with 7 families and 5 copies of GH31 being the most abundant (Table S4). Similarly, the

Endogonales genomes contained an average 8 families, with GH31 being the most abundant.

The G-AMF genomes contained an average 3 families, with GH31 and GH5_7 being the

most abundant (Table S4). No copies of GH29 or GH95 were found within the G-AMF

genomes. Similarly, no copies of GH95 were found within the MAG. However, a single copy

of GH29 was present (Table S4).

Pectinases in the MAG were limited, with only 4 copies across 3 families: 2 GH35s, 1 GH2,

and 1 PL1_7 (Table S4). The Endogonales genomes contained an average of 4 families, with

the most abundant being GH28 and GH35 (Table S4). Within the G-AMF genomes, no

evidence for pectinases was found within 'Gigmar1'. Apart from that, all other genomes

contained only 2 or 3 copies of GH35 (Table S4).

The MAG contained 3 families of lignin degrading enzyme: 2 AA1, 1 AA1_2, and 1 AA2

(Table S4). This trend was conserved across the Endogonales genomes, in which lignin

degrading enzymes comprised of 3 families, with AA1 being slightly more abundant (Table

degrading enzymes comprised of 3 families, with AA1 being slightly more abundant (Table

S4). The G-AMF genomes, while only comprising an average 3 families, contained a much

larger number of lignin degrading enzymes. The Diversisporales contained 37 and 25 lignin

degrading enzymes, including 35 and 23 copies of AA1 respectively. The Glomerales

contained an average of 12 lignin degrading enzymes per genome, with an average of 9

counts being attributed to AA1 (Table S4).

The G-AMF genomes had a smaller mean chitinase repertoire (16) than ERM (35, p < 0.01),

OCM (25, p < 0.05), and SAP (26, p < 0.01) (Fig. 7a). However, the MAG had one of the

highest numbers of chitinases across the genome database (38), roughly double the mean

number found in both the G-AMF and ECM (21). However, there was considerable variation

in the numbers of chitinases across the G-AMF genomes, with the two Diversisporales

genomes containing 38 and 29 counts respectively, more comparable to the MAG. The

Endogonales genomes similarly contained high mean numbers of chitinases (29 to 42)

compared to other genomes in the dataset. Over half of the MAG's chitinase repertoire was

attributed to just two families, CE4 and GH18, with 16 and 13 copies respectively, a trend

also seen across the Endogonales genomes (Table S4). The G-AMF genomes largely

comprised of CE4 chitinases, however, the Diversisporales genomes also contained a large

number of GH18 (13 and 10), very few of which (0 to 2) were found in Glomerales (Table

S4). The MAG contained a fully comprehensive set of chitin degrading enzymes (Table S5).

Similarly, all but one of the G-AMF genomes contained a full chitin degradation suite (Table

S5).

Figure 7: Number of chitin, glycan, mannan, and peptidoglycan degrading predicted proteins

per genome distributed by ecological lifestyle. Lifestyle is colour coded. Coloured data points

represent selected Mucoromycotina species for comparison: yellow for 'Endsp1', red for

'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and 'Jimfl_GMNB39_1'.

Data point shapes within the G-AMF distinguish between the Diversisporales (star) and

Glomerales (triangle). Different letters indicate significant differences between lifestyles as

determined by a Kruskal-Wallis test (p = 0.012 for chitin, p = 0.024 for peptidoglycan,

and p < 0.01 for the remainder) followed by a post-hoc Dunn's test (p < 0.05).

ERM genomes contained a higher mean number of mannanases (24) than the saprotrophic

and other mycorrhizal lifestyles (3 to 6) (Fig. 7c). However, the MAG contained a

substantially higher mean number of mannanases (9) than the G-AMF (3) and ECM (5), and

also the Endogonales genomes (2 to 4). Similarly, there was a range of mannanases across the

G-AMF genomes with the two Diversisporales genomes containing 11 and 10 counts

respectively, more comparable to the MAG (Table S4). The MAG again contained a

comparatively large set of genes for mannan degradation, with 5 copies of GH92, 3 copies of

GH125, and 1 copy of GH76. However, this represents the one substrate-specific group that

was not as conserved across the chosen Endogonales genomes. None of the Endogonales

genomes contained copies of GH76, while all contained 1 or 2 copies of CE17 (Table S4).

Additionally, while three of the four contained GH92, it was only in single-copy instances.

The G-AMF genomes showed a range of mannanases within their genomes, and particularly

the Diversisporales genomes alone contained many copies of GH92. All but one Glomerales

genome only contained 1 or 2 copies of GH125, with no mannanase identified in the

'RhiirC2' genome (Table S4).

The MAG contained no genome-specific cell wall degrading enzymes families. However, a

poorly described GH151 family, "α-L-fucosidase enzyme encoding protein", was only

identified within 3 of the Endogonales genomes (Jimfl_AD_1, Jimfl_GMNB39_1, and

Endsp1) and a far-removed saprotrophic basidiomycete, Dacsp1 (Table S4). The MAG

possessed similar numbers of glycanases and peptidoglycanases to G-AMF and Endogonales

genomes (Fig 7b, d and supplementary information).

Transporters

Since the ability to transport nutrients and assimilate with the host plant is core to the lifestyle

of a fungal symbiont, we compared substrate-specific transporters between the ecological

lifestyles. Only the ERM genomes contained a statistically higher (p = 0.044) number of

phosphate transporters than the other genomes (Fig. 8a). The MAG contained a relatively low

number of phosphate transporters (14), although this was not too dissimilar from the mean of

the G-AMF (18) and the same as the mean ECM genome (14). Similarly, the Endogonales

genomes possessed low numbers of phosphate transporters (13 to 15) (Fig. 8a). The G-AMF

genomes had a significantly higher (p < 0.01) number of calcium transporters (55) compared

to the other mycorrhizal and saprotrophic lifestyles (Fig. 8b, which ranged between 26 and

39), and also the endophyte, parasite, pathogen, and mixed lifestyles (Fig. S13). The MAG

showed a similar number of calcium transporters (50) to the G-AMF, while the number of

calcium transporters in the Endogonales genomes varied considerably (25 to 47) (Fig. 8b).

There was no significant difference (p = 0.48) in the number of copper transporters across the

mycorrhizal and saprotrophic genomes, with the mean varying between 13 and 17 (Fig. 8c).

However, the MAG contained only 6 copper transporters, one of the lowest of all genomes

(Fig. 8c, Fig. S13). Of the Endogonales genomes, 'Jimlac1' had the second smallest number

of copper transporters (7) while the rest ranged from 8 to 13 (Fig. 8c). The G-AMF had a

significantly higher ($p < 0.01$) number of nitrate transporters (7) compared to the other

mycorrhizal and saprotrophic lifestyles (Fig. 8d). The MAG contained a low number (2) of

nitrate transporters relative to the mean G-AMF, although this was the same as the mean

ECM (2) and similar to the ERM (3), OCM (1), and SAP (1). There was considerable

variation in the mean number of nitrate transporters within the Endogonales genomes (0-5)

(Fig. 8d). The ERM genomes had a significantly higher ($p < 0.01$) mean number of

ammonium transporters (11) than all but the G-AMF genomes (9) (Fig. 8e). The MAG

contained 5 ammonium transporters, slightly higher than the median ECM (4) and OCM (4).

The MAG contained an especially low number of glycine transporters (1), the same as the

average ECM (1) and OCM (1) genome (Fig. 8f). The ERM contained the highest mean

number of glycine transporters (5), whereas the G-AMF (3) and SAP (4) were not

significantly different to any of the other ecological lifestyles (Fig. 8f).

Figure 8: Number of phosphate, calcium, copper, nitrate, ammonium, and glycine

transporters per genome distributed by ecological lifestyle. Lifestyle is colour coded.

Coloured data points represent selected Mucoromycotina species for comparison: yellow for

'Endsp1', red for 'Jimlac1', green for 'Bifad1', and purple for both 'Jimfl_AD_1' and

'Jimfl_GMNB39_1'. Data point shapes within the G-AMF distinguish between the

Diversisporales (star) and Glomerales (triangle). Different letters indicate significant

differences between lifestyles as determined by a Kruskal-Wallis test ($p = 0.028$ for

phosphate, $p = 0.62$ for copper, $p = 0.041$ for nitrate, and $p < 0.01$ for the remainder)

followed by a post-hoc Dunn's test ($p < 0.05$).

Hallmark Missing Glomeromycotan Core Genes

We analysed both the MAG and G-AMF genomes for enzymes linked to pathways that are

missing from typical G-AMFs, often referred to as 'Missing Glomeromycotan Core Genes'

(MGCG). As fatty acid auxotrophs, G-AMF have been shown to lack the presence of

cytosolic fatty acid synthases required for fatty acid biosynthesis. We found no evidence of

proteins directly associated with the fatty acid synthase required to facilitate fatty acid

biosynthesis within either the G-AMF genomes or the MAG. We did identify 3-oxoacyl-ACP

reductase in all genomes, including 10 within the MAG. (Table S5). Additionally, neither the

MAG nor the G-AMF genomes possessed the ability to biosynthesis thiamine, a hallmark of

G-AMF genomes. Specifically, we found no evidence of the presence of thiamine phosphate

synthase or hydroxymethylpyrimidine kinase (Table S5).

The ability to synthesis all derivatives of the vitamin B6 pathway was present within the

MAG. This included a copy of pyridoxine kinase (2.7.1.35), the most prolific enzyme within

the vitamin B6 metabolism pathway. Only one enzyme, pyridoxine 4-dehydrogenase, was

missing from the pathway, while an indirect three-step metabolism for the conversion of

pyridoxal to pyridoxine was present. Conversely, no G-AMF genome possessed a full

vitamin B6 metabolism pathway, with no evidence of pyridoxine kinase being found across

the group (Table S5).

Discussion

In the current work, we assembled the genome of a putative FRE-forming AMF from a root

metagenome. Methodologically, this is a relatively unique achievement: fungal MAGs have

been obtained previously from host-associated environments and niche environments, but not

from complex environments, such as the rhizosphere [72-74], or from similar fungal

symbionts. To achieve this, we had to augment standard binning methods with assembly

graph visualisations and coverage correlations to known M-AMF 18S rRNA genes. We

believe that this method has many advantages over isolation and culturing, in particular, it is

less time-consuming and in principle may be more reflective of the actual community

present, particularly if multiple species are present.

We acknowledge that our assembled genome size of 37.37 Mb may underestimate the actual

genome size. Short reads can, at times, have inherent difficulties in resolving repetitive

regions and complex structures, especially when the repeats exceed the length of the reads.

Consequently, repetitive sequences or duplications may collapse, presenting a smaller

genome [75]. The early diverging G-AMF Paraglomus occultum genome exhibited a very

similar sized (39.6 Mb) genome with very small numbers of repeats, highlighting that a large

genome is not a prerequisite of an AM lifestyle [76, 77]. Furthermore, many

Mucoromycotina genomes of varying ecological lifestyles are represented by genomes of

similar size and limited complexity [78]. There is conflicting evidence as to whether

arbuscular mycorrhizal genomes are diploid or simply 'diploid-looking' examples of

heterokaryosis [79]. Since short-read assemblers cannot phase chromosomes, in the event that

M-AMF is a diploid organism, while the genome contents will be assembled correctly, it will

be as a haploid chromosome.

Phylogenetic analysis of the correlative 18S rRNA marker, consensus_all_seqs_0_16,

showed that the MAG sits within the order Densosporales of the Mucoromycotina. The MAG

represents the first published genome of both the order Densosporales and the family

Planticonsortiaceae. Due to the absence of sequenced genomes from the Densosporales—the

order to which the MAG likely belongs—the only available genomes from the

Mucoromycotina for comparison are those of putative ectomycorrhizal and saprotrophic

fungi from the order Endogonales. Although these genomes provide valuable comparative

insights, the MAG is phylogenetically distinct from these taxa, and a reasonable degree of

dissimilarity is evident. The MAG was closely related to environmental sequences from the

rhizoids of liverworts, which form arbuscular-like symbioses involving Glomeromycota and

Mucoromycotina [80]. To date, understanding of the phylogeny of fungi which form

mycorrhizas with FRE morphology in vascular plants has been restricted to short 18S rRNA

fragments. Similarly, these studies have confirmed close phylogenetic relatedness with

sequences from liverwort thalli [22, 26-28], suggesting these fungi form associations with

both vascular and non-vascular plants, with a role in N and P nutrition of the plant host [22,

33]. Mucoromycotina sequences associated with plant roots may span both the Densosporales

[18, 29, 39] and Endogonales [28], and furthermore diverse assemblages of fungi from these

orders may be present in roots [28], although fungi associated with the FRE morphology

appear to be phylogenetically distinct from putative ectomycorrhizal and saprotrophic

Endogonales [28].

Comparative genome analysis showed that the MAG's genome had many features in

common with G-AMF. Some of these genomic features, including the low abundance of

PCWDE and the absence of thiamine biosynthesis genes, were also seen in the Endogonales

genomes and therefore aren't likely to be specific, hallmark features of an adaptation to a

symbiotic lifestyle. However, other signatures the MAG shares with G-AMF, including a

high abundance of calcium transporters and the absence of fatty acid synthases, may point to

physiological convergence in the way in which the symbioses operate. There were some

notable differences between the MAG and G-AMF, with the MAG possessing a high

abundance of chitinases, a complete vitamin B6 biosynthesis pathway, and low abundances

of nitrate, ammonium, and copper transporters. These features may point to differences

between G-AMF and M-AMF in the way they interact with the host plant and the wider

environment.

It has been suggested that the transition from saprotrophic to mycorrhizal lifestyles involved

loss of PCWDE, enabling fungi to exist within roots without digesting them [25]. The G-

AMF symbiosis is estimated to have evolved 450 mya, and Glomeromycotan genomes

possess substantially lower numbers of these genes than the fungi forming the more recently

evolved ectomycorrhizal, ericoid mycorrhizal, and orchid mycorrhizal symbioses.

Furthermore, G-AMF show signatures of dependence on host metabolism that are absent

from more recently evolved symbioses, including fatty acid auxotrophy and an inability to

synthesise thiamine and metabolise vitamin B6 [25]. These represent the 'Missing

Glomeromycota Core Genes' (MGCG) [15-17].

Although genomic evidence indicates that G-AMF and M-AMF have independent origins

[17], these symbioses co-occurred in the earliest land plants [7], and they continue to co-

occur in roots of extant plants [22]. Additionally, while G-AMF and M-AMF exhibit some

niche differentiation, evidence of overlapping ecological presence and similar responses to

overarching edaphic and environmental variables have been noted [27, 28]. Furthermore,

ultrastructural evidence points to similarities in the way in which G-AMF and M-AMF

function at a cellular level [24]. Therefore, similar evolutionary pressures may have acted on

these symbioses, which could have resulted in convergence of genomic traits. Indeed, many

of the symbiotic genomic features shown by G-AMF were also displayed by the MAG. This

included a reduced repertoire of PCWDE and the absence of several MGCG, including those

related to thiamine and fatty acid biosynthesis.

Our analysis confirms the findings of Rosling et al. (2024) that Endogonales genomes have a

low genomic PCWDE count and an incomplete thiamine biosynthesis pathway. Similarly,

our MAG, which represents an early-diverging lineage within the Mucoromycotina and is

possibly related to the Densosporales, also displays these features. This suggests that these

traits may be present in multiple lineages within the Mucoromycotina and may not be

exclusive adaptations to biotrophy. However, the Endogonales genomes possess fatty acid

biosynthesis genes [17], whereas both the G-AMF genomes and the MAG lack these genes.

This absence in both G-AMF and the MAG could indicate a convergent adaptation to

biotrophy, potentially suggesting that host assimilate is provided as fatty acids in both

symbioses.

The MAG had a number of features which distinguished it from G-AMF genomes. In

particular, while there was a shared low abundance of CAZymes, the MAG possessed a

greater repertoire of enzymes for degradation of organic substrates than typically seen in G-

AMF genomes. This included the full complement of enzymes required for degradation of

cellulose, which was absent in all G-AMF genomes. In notable contrast to G-AMF, the MAG

had a high count of both chitin and mannan fungal cell wall degrading enzymes. The high

chitinase count doesn't appear to be specific to the M-AMF, with the putative

ectomycorrhizal and saprotrophic Endogonales genomes showing a similar pattern, attributed

to elevated numbers of CE4 and GH18, as seen previously in Rosling et al. (2024). However,

the full complement of chitinases was present in the G-AMF genomes as well as in the MAG,

so this feature may not have any functional significance. Relative to other mycorrhizal fungi

(except ericoid mycorrhizal fungi), the MAG had a notably high count of genes for mannan

degradation. These were absent or present in low numbers in most G-AMF genomes and,

significantly, were also found in low abundance in the Endogonales genomes, therefore high

abundance of mannan degradation genes may represent a distinguishing feature of FRE

genomes. However, even within the G-AMF there appears to be some variation, with the

Diversisporales genomes containing a higher number of mannan and chitin degrading

enzymes than the Glomerales genomes, indicating a substantial range of genomic repertoires.

It is important to understand whether these contrasting enzymatic profiles contribute to niche

partitioning between G-AMF and M-AMF in nutrient mobilisation from contrasting mineral

and organic substrates. For example, high abundance of chitin and mannan degrading genes

could indicate a role for M-AMF in the degradation of fungal necromass, while a complete

cellulose degradation pathway and high abundance of cellulose and hemicellulose degrading

enzymes could suggest involvement in saprotrophic breakdown of plant-derived materials,

similar to some ECM fungi [81, 5]. Alternatively, it could reflect different strategies for

enzymatic degradation of plant cell walls as they colonise and grow through host plant tissues

and penetrate root cells.

Currently, there is no indication that soil organic matter has any influence on the distribution

of G-AMF or M-AMF. In contrast to G-AMF, evidence indicates that M-AMF are more

abundant in agricultural than natural habitats, indicating a preference for low organic matter

soils [27]. However, these functional differences could support niche partitioning within

these habitats.

A further distinguishing feature between the MAG and the G-AMF genomes is the presence

of a full B6 biosynthesis pathway in the MAG, including pyridoxine kinase, which was

missing in all the G-AMF genomes, as demonstrated in earlier studies [15, 16]. B6 is a

cofactor required for amino acid biosynthesis, and it also plays a role as an antioxidant.

Reactive oxygen species (ROS) scavenging systems such as B6 may be required by AMF to

overcome host defences, allowing intracellular colonisation and the development of

arbuscules [82, 83]. Components of the B6 biosynthesis pathway that G-AMF do possess are

expressed throughout the fungal life cycle, but they are at the highest expression levels within

the root [84]. Similarly, B6 plays a role in supporting the infection of root tissues by

pathogens [85]. Differences in capacity for independent production of B6 between the MAG

and G-AMF could point to differences in the way the fungi scavenge ROS, and potentially

also differences in the way root colonisation and arbuscule development are regulated by the

plant. There could also be implications for the specific mechanisms underlying M-AMF and

G-AMF responses to external stressors which generate ROS such as temperature, osmotic

stress, and nutrient limitation, and the independence of these responses to the plant's own

response [86].

Key physiological traits associated with G-AMF symbioses are the uptake of P from the soil,

its translocation through hyphae to the root, and its transfer to the plant in exchange for

carbon. Ultrastructural [24] and experimental evidence [18] suggest a similar role for M-

AMF. A number of P transporters have been identified in G-AMF involved in P assimilation

from soil, storage of P, and exchange of P with the host [87-89]. G-AMF genomes generally

showed higher mean numbers of P transporters than ECM, while numbers in the MAG were

low relative to other mycorrhizal genomes studied. While the significance of transporters to

the functioning of symbionts will depend on a range of factors including their efficiency, our

data doesn't provide genomic support for particular specialisation in P nutrition by M-AMF.

Evidence also points to a potential role for G-AMF in N nutrition [90]. Indeed, G-AMF

genomes contained higher numbers of nitrate transporters than the other mycorrhizal

symbionts, while also sitting towards the higher end of both ammonium and glycine

transporters within the genomes we analysed. However, further evidence of inter-group

differentiation can be seen across the Diversisporales and Glomerales genomes, with the

former containing especially high numbers of ammonium and glycine transporters. The MAG

contained a lower number of all three categories of N transporters than all G-AMF genomes.

While the MAG contained a typical number of ammonium transporters compared to the other

mycorrhizal lifestyles, it contained a low number of nitrate and glycine transporters.

Hoystead et al. (2023) suggested that M-AMF and G-AMF have complementary roles in host

nutrition, with M-AMF predominantly involved in supply of N to the host and G-AMF in

supplying P. Furthermore, Howard et al. (2024) showed a putative M-AMF displaying both a

preference for, and increased transfer of, glycine and ammonium derived nitrogen sources.

While there is evidence for the ability to transport varying forms of N, our MAG does not

provide exceptional evidence for this relative specialism.

The very low abundance of Cu transporters in the MAG - the lowest across the entire dataset

- is particularly striking. The three ectomycorrhizal Endogonales genomes also contain low

numbers of Cu transporters, however the saprotrophic Endogonales sit near the mean. G-

AMF may promote Cu uptake by the host plant [91] and arbuscules have high demands for

Cu [92]. Furthermore, Cu transporters in G-AMF show differential expression profiles across

intra- and extra-radical mycelium [92]. The extent to which M-AMF are involved in the

supply of micronutrients to the host remains to be determined, but the evidence presented

here could indicate that in contrast to G-AMF, this may not be an important function for M-

AMF.

The high number of calcium transporters in both the MAG and G-AMF is a clear point of

similarity between them, and of differentiation to all other groups. AMF have been

implicated in transport of Ca to host plants [93], and our data could therefore indicate a role

for both G-AMF and M-AMF in this function. However, Ca may also play a key role in the

arbuscule lifecycle of both FRE and G-AMF. A clear link between a dramatic increase in

calcium accumulation and arbuscule collapse has been demonstrated within G-AMF [94, 95].

A similar trend within FRE has also been documented [24, 96]. While it is unknown whether

the Ca accumulated originates from the host plant or the fungi, the high number of Ca

transporters within both G-AMF and the MAG may indicate that Ca accumulation is driven

by the fungal symbionts, perhaps depositing Ca to cease the function of the arbuscule once it

reaches maturity.

We must exercise some caution when interpreting these findings since we have constructed a

relatively fragmented genome from a metagenome. However, our predicted level of

contamination is only 1.5%, this suggests that we are unlikely to be assigning functions to the

MAG that are not really present. In contrast, completion is potentially low at 83.7% so it is a

definite possibility that we have missed a significant number of genes but most of the

analysis above is at the level of gene families, if the missing genes are randomly distributed

through these families we are unlikely to predict the absence of an entire family that is

actually possessed by our MAG.

Conclusion

In this study, we assembled and characterized a MAG of a putative fine root endophyte

forming Mucoromycotina AMF from samples collected in Australia. Comparative analyses

revealed key genomic features shared with related fungal taxa, shedding light on potential

functional adaptations to the root-associated lifestyle. Our findings contribute to the growing

understanding of Mucoromycotan-AMF diversity and their ecological roles, providing a

valuable genomic resource for future research on endophytic symbioses and nutrient

exchange in plant-fungal interactions. For both G-AMF and M-AMF, comprehensive analysis

of genomic functional gene repertoires across the broad range of taxa found within roots is

key to understanding the way that AMF communities support plant health, and the potential

for engineering communities to promote beneficial traits. With the development of third

generation long read sequencing technologies, genome assembly from metagenomes

represents a promising route for understanding functions and traits associated with dark taxa

in complex microbial communities, especially when combined with transcriptome sequencing

[40] to understand the factors which regulate gene expression within plant tissues and the

soil.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Consent for use of Figure 1d obtained from Suzanne Orchard. Credit given in the figure

legend.

Availability of data and materials

Raw 18S rRNA Miseq sequences are available at NCBI (BioProject ID PRJNA1179892), and

Nanopore sequences at ENA (Experiment Accession ERX13330708). The FRE4583

sequence is available at NCBI (SUB14894546). The 18S rRNA phylogeny sequence

alignment FASTA file and Newick tree file are available on PlutoF Biodiversity Platform

(https://dx.doi.org/10.15156/BIO/3301225). The sequence of the MAG has been deposited on

the Zenodo repository (https://doi.org/10.5281/zenodo.14144746).

Competing interests

900   The authors declare that they have no competing interests

Funding

902   Natural Environment Research Council (Grant NE/S010270/1), Biotechnology and

903   Biological Sciences Research Council (BBSRC) Midlands Integrative Biosciences Training

904   Partnership (Grant BB/T00746X/1) and Australian Research Council (Discovery Project

905   DP180103157)

Authors' contributions

GB, MHR, and CQ conceived the work and with RM obtained funding. MHR and PAD

conducted the M-AMF enrichment work and the assessment of M-AMF colonisation of plant

roots. SH, SR, and GB conducted sequencing and analysis of 18S rRNA genes. SR and CQ

assembled the MAG. RHN, GM, and JC conducted the 18S rRNA gene phylogenetic

analysis. JC, GM, SR, and GB performed the comparative genomic analyses. JC and GB

wrote the manuscript. All authors edited the manuscript.

Acknowledgements

914   We thank the Natural Environment Research Council (Grant NE/S010270/1), the

915   Biotechnology and Biological Sciences Research Council (BBSRC) Midlands Integrative

916   Biosciences Training Partnership (Grant BB/T00746X/1) and the Australian Research

917   Council (Discovery Project DP180103157) for funding.

Authors' information

ORCID:

JC- 0009-0004-1563-8469

SR - 0000-0001-8727-2604

PAD - 0000-0003-3055-4744

SH - 0000-0002-3786-1096

GM - 0000-0002-9218-1210

RMM - 0000-0003-3572-3500

RHN – 0000-0002-8052-0107

MHR – 0000-0003-0749-0199

CQ – 0000-0003-1884-8440

GDB - 0000-0002-9740-5244

References

[1] Begum N, Qin C, Ahanger MA, Raza S, Khan MI, Ashraf M, Ahmed N, Zhang L. Role of

arbuscular mycorrhizal fungi in plant growth regulation: implications in abiotic stress

tolerance. *Frontiers in Plant Science.* 2019;10: 1068.

[2] Hawkins HJ, Cargill RIM, Nuland MEV, Hagen SC, Field KJ, Sheldrake M,

Soudzilovskaia NA, Kiers TE. Mycorrhizal mycelium as a global carbon pool. *Current*

Biology. 2023;33(11): 560-573.

[3] Luginbuehl LH, Menard GN, Kurup S, Van Erp H, Radhakrishnan GV, Breakspear A,

Oldroyd GED, Eastmond PJ. Fatty acids in arbuscular mycorrhizal fungi are synthesized by

the host plant. *Science.* 2017;365(6343): 1175-1178.

[4] Chen M, Arato M, Borghi L, Nouri E, Reinhardt D. Beneficial services of arbuscular

mycorrhizal fungi – from ecology to application. *Frontiers in Plant Science.* 2018;9: 1270.

[5] Martin FM, van der Heijden MGA. The mycorrhizal symbiosis: research frontiers in

genomics, ecology, and agricultural application. *New Phytologist.* 2024;242(4): 1486-1506.

[6] Brundrett MC, Tedersoo L. Evolutionary history of mycorrhizal symbioses and global

host plant diversity. *New Phytologist*. 2018;220(4):1108-1115.

[7] Strullu-Derrien C, Selosse M-A, Kenrick P, Martin FM. The origin and evolution of

mycorrhizal symbioses: from palaeomycology to phylogenomics. *New Phytologist.*

2018;220(4): 1012-1030.

[8] Martino E, Morin E, Grelet G-A, Kuo A, Kohler A, Daghino S, Barry KW, Cichocki N,

Clum A, Dockter RB et al. Comparative genomics and transcriptomics depict ericoid

mycorrhizal fungi as versatile saprotrophs and plant mutualists. *New Phytologist.*

2018;217(3): 1213-1229.

[9] Gong Y, Lebreton A, Zhang F, Martin FM. Role of carbohydrate-active enzymes in

mycorrhizal symbioses. *Essays in Biochemistry.* 2023;67(3): 471-478.

[10] Read DJ, Duckett JG, Francis R, Ligron R, Russell A. Symbiotic fungal associations in

'lower' land plants. *Philosophical Transactions of the Royal Society of London.*

2000;355(1398): 815–831.

[11] Parniske, M. Arbuscular mycorrhiza: the mother of plant root endosymbiosis. *Nature*

Reviews Microbiology. 2008;6: 763–775.

[12] Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G,

Corradi N, Grigoriev I, Gryganskyi A et al. A phylum-level phylogenetic classification of

zygomycete fungi based on genome-scale data. *Mycologia.* 2016;108(5): 1028–1046.

[13] Schüßler A, Martin H, Cohen D, Fitz M, Wipf D. Arbuscular mycorrhiza: studies on the

geosiphon symbiosis lead to the characterization of the first glomeromycotan sugar

transporter. *Plant Signaling & Behavior.* 2007;2(5), 431–434.

[14] Luginbuehl LH, Oldroyd GED. Understanding the arbuscule at the heart of

endomycorrhizal symbioses in plants. *Current Biology.* 2017b; 27(17): 952-963.

[15] Tang N, Clemente HS, Roy S, Bécard G, Zhao B, Roux C. A survey of the gene

repertoire of *Gigaspora rosea* unravels conserved features among Glomeromycota for

obligate biotrophy. *Frontiers in Microbiology.* 2016*;*1(7):233.

[16] Kobayashi Y, Maeda T, Yamaguchi K, Kameoka H, Tanaka S, Ezawa T, Shigenobu S,

Kawaguchi M. The genome of Rhizophagus clarus HR1 reveals a common genetic basis for

auxotrophy among arbuscular mycorrhizal fungi. *BMC Genomics.* 2018;19(1): 465.

[17] Rosling A, Sahraei SE, Khan FK, Desirò A, Bryson AE, Mondo SJ, Grigoriev IV,

Bonito G. Sánchez-García M. Evolutionary history of arbuscular mycorrhizal fungi and

genomic signatures of obligate symbiosis. *BMC Genomics.* 2024;25: 529.

[18] Orchard S, Standish RJ, Dickie IA, Renton M, Walker C, Moot D, Ryan MH. Fine root

endophytes under scrutiny: a review of the literature on arbuscule-producing fungi recently

suggested to belong to the Mucoromycotina. *Mycorrhiza.* 2017b*;*27(7): 619-638.

[19] Schüßler A, Schwarzott D, Walker C. A new fungal phylum, the Glomeromycota:

phylogeny and evolution. *Mycological Research.* 2001;105(12): 1413-1421.

[20] Tedersoo L, Sánchez-Ramírez S, Kõljalg U, Bahram M, Döring M, Schigel D, May T,

Ryberg M, Abarenkov K. High-level classification of the Fungi and a tool for evolutionary

ecological analyses. *Fungal Diversity.* 2018;90: 135–159.

[21] Thippayarugs S, Bansal M, Abbott LK. Morphology and infectivity of fine endophyte in

a Mediterranean environment. *Mycological Research.* 1999;103(11): 1369-1379.

[22] Orchard S, Hilton S, Bending GD, Dickie IA, Standish RJ, Gleeson DB, Jeffery RP,

Walker C, Bass D, Monk J et al. Fine endophytes (Glomus tenue) are related to

Mucoromycotina, not Glomeromycota. *New Phytologist.* 2017a;213(2): 481-486.

[23] Walker C, Gollotte A, Redecker D. A new genus, Planticonsortium (Mucoromycotina),

and new combination (P. tenue), for the fine root endophyte, *Glomus tenue* (basionym

Rhizophagus tenuis). *Mycorrhiza.* 2018;12(3): e8603.

[24] Albornoz FE, Hayes PE, Orchard S, Clode PL, Nazeri NK, Standish RJ, Bending GD,

Hilton S, Ryan MH. First cryo-scanning electron microscopy images and X-ray

microanalyses of mucoromycotinian fine root endophytes in vascular plants. *Frontiers in*

Microbiology. 2020; 11: 2018.

[25] Miyauchi S, Kiss E, Kuo A, Drula E, Kohler A, Sánchez-García M, Morin E,

Andreopoulos B, Barry KW, Bonito G et al. Large-scale genome sequencing of mycorrhizal

fungi provides insights into the early evolution of symbiotic traits. *Nature Communications.*

2020;11: 5125.

[26] Albornoz FE, Orchard S, Standish RJ, Dickie IA, Bending GD, Hilton S, Lardner T,

Foster KJ, Gleeson DB, Bougoure J et al. Evidence for niche differentiation in the

environmental responses of co-occurring mucoromycotinian fine root endophytes and

glomeromycotinian arbuscular mycorrhizal fungi. *Microbial Ecology.* 2021;81(4): 864-873.

[27] Albornoz FE, Ryan MH, Bending GD, Hilton S, Dickie IA, Gleeson DB, Standish RJ.

Agricultural land-use favours Mucoromycotinian, but not Glomeromycotinian, arbuscular

mycorrhizal fungi across ten biomes. *New Phytologist.* 2022;233(3): 1369-1382.

[28] Seeliger M, Hilton S, Muscatt G, Walker C, Bass D, Albornoz F, Standish RJ, Gray ND,

Mercy L, Rempelos L et al. New fungal primers reveal the diversity of Mucoromycotinian

arbuscular mycorrhizal fungi and their response to nitrogen application. *Environmental*

Microbiome. 2024;19: 71.

[29] Desirò A, Rimington WR, Jacob AS, Pol NV, Smith MR, Trappe JM, Bidartondo MI,

Bonito G. Multigene phylogeny of Endogonales, an early diverging lineage of fungi

associated with plants. *IMA Fungus.* 2017;8(2): 245–257.

[30] Hoysted GA, Jacob AS, Kowal J, Giesemann P, Bidartondo MI, Duckett JG, Gebauer G,

Rimington WR, Schornack S, Pressel S et al. Mucoromycotina fine root endophyte fungi

form nutritional mutualisms with vascular plants. *Plant Physiology.* 2019;181(2): 565-577.

[31] Jeffery RP, Simpson RJ, Lambers H, Orchard S, Kidd DR, Haling RE, Ryan MH.

Contrasting communities of arbuscule-forming root symbionts change external critical

phosphorus requirements of some annual pasture legumes. *Applied Soil Ecology.* 2018;126:

88-97.

[32] Orchard S, Standish RJ, Nicol D, Gupta VVSR, Ryan MH. The response of fine root

endophyte (Glomus tenue) to waterlogging is dependent on host plant species and soil type.

Plant and Soil. 2016;403: 305-315.

[33] Hoysted GA, Field KJ, Sinanaj B, Bell CA, Bidartondo MI, Pressel S. Direct nitrogen,

phosphorus and carbon exchanges between Mucoromycotina 'fine root endophyte' fungi and

a flowering plant in novel monoxenic cultures. *New Phytologist.* 2022;238(1): 70-79.

[34] Howard NAO, Williams A, Durant E, Pressel S, Daniell TJ, Field KJ. Preferential

nitrogen and carbon exchange dynamics in Mucoromycotina "fine root endophyte"-plant

symbiosis. *Current Biology.* 2024;34(23): 5484-5493.

[35] Tanaka S, Hashimoto K, Kobayashi Y, Yano K, Maeda T, Kameoka H, Ezawa T, Saito

K, Akiyama K, Kawaguchi M. Asymbiotic mass production of the arbuscular mycorrhizal

fungus Rhizophagus clarus. *Communications Biology.* 2022;5: 43.

[36] Tisserant E, Malbreil M, Kuo A, Kohler A, Symeonidi A, Balestrini R, Charron P,

Duensing N, Frei dit Frey N, Gianinazzi-Pearson V et al. Genome of an arbuscular

mycorrhizal fungus provides insight into the oldest plant symbiosis. *Proceedings of the*

National Academy of Sciences. 2013;110(50):20117-20122.

[37] Montoliu-Nerin M, Sánchez-García M, Bergin C, Kutschera VE, Johannesson H, Bever

JD, Rosling A. In-depth phylogenomic analysis of arbuscular mycorrhizal fungi based on a

comprehensive set of de novo genome assemblies. *Frontiers in Fungal Biology.* 2021;29(2):

716385.

[38] Öpik M, Davison J, Moora M, Zobel M. DNA-based detection and identification of

Glomeromycota: the virtual taxonomy of environmental sequences. *Botany-Botanique.*

2014;92: 135–147.

[39] Chang Y, Desirò A, Na H, Sandor L, Lipzen A, Clum A, Barry K, Grigoriev IV, Martin

FM, Stajich JE et al. Phylogenomics of Endogonales and evolution of mycorrhizas within

Mucoromycota. *New Phytologist.* 2019;222(1): 511-525.

[40] Benoit G, Raguideau S, James R, Phillippy AM, Chikhi R, Quince C. High-quality

metagenome assembly from long accurate reads with metaMDBG. *Nature Biotechnology*.

2024;42: 1378–1383.

[41] Yang C, Chowdhury D, Zhang Z, Cheung WK, Lu A, Bian Z, Zhang L. A review of

computational tools for generating metagenome-assembled genomes from metagenomic

sequencing data. *Computational and Structural Biotechnology Journal*. 2021;19: 6301-6314.

[42] Peng X, Wilken SE, Lankiewicz TS, Gilmore SP, Brown JL, Henske JK, Swift CL,

Salamov A, Barry K, Grigoriev IV et al. Genomic and functional analyses of fungal and

bacterial consortia that enable lignocellulose breakdown in goat gut microbiomes. *Nature*

Microbiology. 2021;6: 499–511.

[43] Setubal JC. Metagenome-assembled genomes: concepts, analogies, and challenges.

Biophysical Reviews. 2021; 13(6): 905–909.

[44] West PT, Probst AJ, Grigoriev IV, Thomas BC, Banfield JF. Genome-reconstruction for

eukaryotes from complex natural microbial communities. *Genome Research*. 2018;28(4):569-

580.

[45] Wurzbacher C, Larsson E, Johan Bengtsson☐Palme, Van den Wyngaert S, Svantesson

S, Kristiansson E, Kagami M, Nilsson HR. Introducing ribosomal tandem repeat barcoding

for fungi. *Molecular Ecology Resources*. 2018;19(1):118-127.

[46] Vierstraete AR, Braeckman BP. Amplicon_sorter: A tool for reference-free amplicon

sorting based on sequence similarity and for building consensus sequences. *Ecology and*

Evolution. 2022;12(3): e8603.

[47] Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. *ArXiv*. doi: 10.48550/arXiv.1303.3997

[48] Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A,

Keane T, McCarthy SA, Davies RM, Li H. Twelve years of SAMtools and BCFtools.

GigaScience 2021;10(2): giab008.

[49] Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD et al. SPAdes: a new genome assembly algorithm and

its applications to single-cell sequencing. *Journal of Computational Biology*. 2012;19(5):

455-477.

[50] Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics*. 2010;26(6): 841-2.

[51] Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, ljaz UZ, Loman NJ,

Andersson AF, Quince C. Binning metagenomic contigs by coverage and composition.

Nature Methods. 2014;11: 1144–1146.

[52] Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualisation of de novo

genome assemblies. *Bioinformatics.* 2015;31(20): 3350-3352.

[53] Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel

and streamlined workflows along with broader and deeper phylogenetic coverage for scoring

of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution.*

2021;38(10): 4647-4654.

[54] Koboldt DC, Chen K, Wylie T, Larson DE, McLellan MD, Mardis ER, Weinstock GM,

Wilson RK, Ding L. VarScan: variant detection in massively parallel sequencing of

individual and pooled samples. *Bioinformatics*. 2009;25(17): 2283–2285.

[55] Gruber-Vodicka HR, Seah BKB, Pruesse E. phyloFlash: Rapid small-subunit rRNA

profiling and targeted assembly from metagenomes. m*Systems.* 2020;5(5): e00920-20.

[56] Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

The SILVA ribosomal RNA gene database project: improved data processing and web-based

tools. *Nucleic Acids Research.* 2013;41: 590–596.

[57] Antunes PM, Stürmer SL, Bever JD, Chagnon PC, Chaudhary VB, Deveautour C, Fahey

C, Kokkoris V, Lekberg Y, Powell JR et al. Enhancing consistency in arbuscular mycorrhizal

trait-based research to improve predictions of function. *Mycorrhiza*. 2025;53(14).

[58] Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular Biology and Evolution.* 2013;30(4):

772–780.

[59] Tamura K, Stecher G, Kumar S. MEGA11: molecular evolutionary genetics analysis

version 11. *Molecular Biology and Evolution.* 2021;38(7): 3022–3027.

[60] Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:

Fast model selection for accurate phylogenetic estimates. *Nature Methods.* 2017;14: 587-589.

[61] Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,

Lanfear R. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the

genomic era. *Molecular Biology and Evolution*. 2020;37(5): 1530-1534.

[62] Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the

ultrafast bootstrap approximation. *Molecular Biology and Evolution*. 2018;35: 518–522.

[63] Rambaut, A. 2009. Figtree. Tree figure drawing tool, v1.4.4. Edinburgh, Scotland.

http://tree.bio.ed.ac.uk/software/figtree.

[64] Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. ggtree: an r package for visualization and

annotation of phylogenetic trees with their covariates and other associated data. *Methods in*

Ecology and Evolution. 2017;8(1): 28-36.

[65] R Core Team. 2023. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

[66] Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New

York. ISBN 978-3-319-24277-4

[67] Kim GB, Kim JY, Lee JA, Norsigian CJ, Palsson BO, Lee SY. Functional annotation of

enzyme-encoding genes using deep learning with transformer layers. *Nature*

Communications. 2023;14: 7370.

[68] Shi Z, Deng R, Yuan Q, Mao Z, Wang R, Li H, Liao X, Ma H. Enzyme commission

number prediction and benchmarking with hierarchical dual-core multitask learning

framework. *Research.* 2023;31(6): 0153

[69] Tedersoo L, Magurno F, Alkahtani S, Mikryukov V. Phylogenetic classification of

arbuscular mycorrhizal fungi: new species and higher-ranking taxa in Glomeromycota and

Mucoromycota (class Endogonomycetes). *MycoKeys.* 2024;107: 249-271.

[70] Lee S-J, Kong M, St-Arnaud M, Hijri M. Arbuscular mycorrhizal fungal communities of

native plant species under high petroleum hydrocarbon contamination highlights

Rhizophagus as a key tolerant genus. *Microorganisms.* 2020;8(6): 872.

[71] Yamamoto K, Degawa Y, Yamada A. Taxonomic study of Endogonaceae in the

Japanese islands: New species of *Endogone*, *Jimgerdemannia*, and *Vinositunica*, gen. nov..

Mycologia. 2020;112(2): 309-328.

[72] Tagirdzhanova G, Saary P, Tingley JP, Díaz-Escandón D, Abbott DW, Finn RD,

Spribille T. Predicted input of uncultured fungal symbionts to a lichen symbiosis from

metagenome-assembled genomes. *Genome Biology and Evolution.* 2021;13(4).

[73] Shen Z, Robert L, Stolpman M, Che Y, Allen KJ, Saffery R, Walsh A, Young A, Eckert

J, Deming C et al. A genome catalog of the early-life human skin microbiome. *Genome*

Biology. 2023;24(252).

[74] Singh NK, Wood JM, Patane J, Moura LMS, Lombardino J, Setubal JC, Venkateswaran

K. Characterization of metagenome-assembled genomes from the International Space Station.

Microbiome. 2023;11(1): 125.

[75] Treangen TJ, Salzberg SL. Repetitive DNA and next-generation sequencing:

computational challenges and solutions. *Nature Reviews Genetics.* 2012;13(1): 36–46.

[76] Malar CM, Wang Y, Stajich JE, Kokkoris V, Villeneuve-Laroche M, Yildirir G, Corradi

N. Early branching arbuscular mycorrhizal fungus Paraglomus occultum carries a small and

repeat-poor genome compared to relatives in the Glomeromycotina. *Microbial Genomics.*

2022;8(4): 000810.

[77] Lanfranco, L, Bonfante P. Lessons from arbuscular mycorrhizal fungal genomes.

Current Opinion in Microbiology. 2023;75(1): 102357.

[78] Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao

X, Korzeniewski F et al. MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic*

Acids Research. 2013;42: 699-704.

[79] Oliveira J, Yildirir G, Corradi N. From Chaos Comes Order: Genetics and Genome

Biology of Arbuscular Mycorrhizal Fungi. *Annual Review of Microbiology.* 2024;78: 147-

168.

[80] Rimington WR, Pressel S, Duckett JG, Field KJ, Read DJ, Bidartondo MI. Ancient

plants with ancient fungi: liverworts associate with early-diverging arbuscular mycorrhizal

fungi. *Proceedings of the Royal Society B: Biological Sciences.* 2018;285(1888): 20181600.

[81] Janusz G, Pawlik A, Sulej J, Świderska-Burek U, Jarosz-Wilkołazka A, Paszczyński A.

Lignin degradation: microorganisms, enzymes involved, genomes analysis and evolution.

FEMS Microbiology Reviews. 2017;41(6): 941-962.

[82] Belmondo S, Calcagno C, Genre A, Puppo A, Pauly N, Lanfranco L NADPH oxidases

in the arbuscular mycorrhizal symbiosis. *Plant Signaling & Behavior.* 2016;11(4): e1165379.

[83] Lanfranco L, Fiorilli V, Gutjahr C. Partner communication and role of nutrients in the

arbuscular mycorrhizal symbiosis. *New Phytologist.* 2018;220(4): 1031-1046.

[84] Benabdellah L, Azcón-Aguilar C, Valderas A, Speziga D, Fitzpatrick TB, Ferrol N.

GintPDX1 encodes a protein involved in vitamin B6 biosynthesis that is up-regulated by

oxidative stress in the arbuscular mycorrhizal fungus *Glomus intraradices*. *New Phytologist.*

2009;184(3): 682-693.

[85] Wang S, Zhang X, Zhang Z, Chen Y, Tian Q, Zeng D, Xu M, Wang Y, Dong S, Ma Z et

al. Fusarium-produced vitamin B6 promotes the evasion of soybean resistance by

Phytophthora sojae. *Journal of Integrative Plant Biology.* 2023;65(9): 2204-2217.

[86] Sahu PK, Jayalakshmi K, Tilgam J, Gupta A, Nagaraju Y, Kumar A, Hamid S, Singh

HV, Minkina T et al. ROS generated from biotic stress: Effects on plants and alleviation by

endophytic microbes. *Frontiers in Plant Science.* 2022;13: 1042936.

[87] Javot H, Pumplin N, Harrison MJ. Phosphate in the arbuscular mycorrhizal symbiosis:

Transport properties and regulatory roles. *Plant, Cell & Environment.* 2006;30: 310–322.

[88] Harrison MJ, Pumplin N, Breuillin FJ, Noar RD, Park HJ. 2010. Phosphate transporters
in arbuscular mycorrhizal symbiosis. In: Koltai H, Kapulnik Y, eds. *Arbuscular Mycorrhizas:*
Physiology and Function. Dordrecht, Netherlands: Springer, 117–135.

[89] Plassard C, Becquer A, Garcia, K. Phosphorus transport in mycorrhiza: How far are we?.
Trends in Plant Science. 2019;24: 794–801.

[90] Gosling P, Mead A, Proctor M, Hammond JP, Bending GD. Contrasting arbuscular
mycorrhizal communities colonizing different host plants show a similar response to a soil
phosphorus concentration gradient. *New Phytologist.* 2013;198(2): 546-556.

[91] Taylor A, Pereira N, Thomas B, Pink DAC, Jones JE, Bending GD. Growth and
nutritional responses to arbuscular mycorrhizal fungi are dependent on onion genotype and
fungal species. *Biology and Fertility of Soils.* 2015;51: 801–813.

[92] Gómez-Gallego T, Benabdellah K, Merlos MA, Jiménez-Jiménez AM, Alcon C,
Berthomieu P, Ferrol N. The *Rhizophagus irregularis* genome encodes two CTR copper
transporters that mediate Cu import into the cytosol and a CTR-like protein likely involved in
copper tolerance. *Frontiers in Plant Science.* 2019;10: 604.

[93] Liu A, Hamel C, Elmi A, Costa C, Ma B, Smith DL. Concentrations of K, Ca and Mg in
maize colonized by arbuscular mycorrhizal fungi under field conditions. *Canadian Journal of*
Soil Science. 2002;82(3): 271-279.

[94] Ryan MH, McCully ME, Huang CX. Location and quantification of phosphorus and
other elements in fully hydrated, soil grown arbuscular mycorrhizas: a cryo analytical
scanning electron microscopy study. *New Phytologist.* 2003;160(2):429-441.

[95] Sujkowska-Rybkowska M, Znojek E. Localization of calreticulin and calcium ions in
mycorrhizal roots of Medicago truncatula in response to aluminum stress. *Journal of Plant*
Physiology. 2018;229: 22-31.

[96] Gianinazzi-Pearson V, Morandi D, Dexheimer J, Gianinazzi S. Ultrastructural and
ultracytochemical features of a *Glomus tenuis* mycorrhiza. *New Phytologist.* 1981;88(4): 633-
639.