Enalapril mitigates aging by targeting antioxidative genes via phosphorylated Smad1/5/9

Wencong Lyu[1†], Haochen Wang[1†], Zhehao Du[1], Ran Wei[1], Jianuo He[1], Fanju Meng[1], Jinlong Bi[1], Lijun Zhang[1], Chao Zhang[2], Yiting Guan[3*] and Wei Tao[1*]

[1]The MOE Key Laboratory of Cell Proliferation and Differentiation, School of Life Sciences, Peking University, Beijing 100871, P. R. China

[2]Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650201, P. R. China

[3]Zhanjiang Institute of Clinical Medicine, Central People's Hospital of Zhanjiang, Guangdong Medical University, Zhanjiang 524045, P. R. China

[†]These authors contributed equally to this work

[*]These authors jointly supervised this work

Correspondence:

Wei Tao (weitao@pku.edu.cn)

ABSTRACT

Aging increases the risk of a myriad of chronic diseases, which are expensive and difficult to treat owing to their various risk factors. Repurposing existing anti-aging medications has accelerated the development of therapies aimed at slowing aging. In this study, we revealed that enalapril, a drug widely prescribed for hypertension, can improve both cellular senescence and individual health. Mechanistically, phosphorylated Smad1/5/9 act as pivotal mediators of the anti-aging properties of enalapril. It stimulates downstream genes involved in cell cycle regulation and antioxidative defense, facilitating cell proliferation and diminishing the production of reactive oxygen species (ROS), thus increasing the antioxidative ability of enalapril. At the organismal level, enalapril has been shown to bolster the physiological performance of various organs; it notably enhances memory capacity and renal function and relieves lipid accumulation. Our work highlights the potential of enalapril to augment antioxidant defenses and combat the effects of aging, thereby indicating its promise as a treatment strategy for aging-associated diseases and its use for healthy aging.

INTRODUCTION

Aging is a multifaceted process influenced by numerous factors, and its intricate mechanisms are not yet fully understood. Geroscience is a burgeoning field of study in which researchers hypothesize that aging is a significant risk factor for many chronic diseases. These findings suggest that interventions targeting the mechanisms of aging might delay the onset or slow the progression of these diseases(Kirkland, 2016; Lopez-Otin, Blasco, Partridge, Serrano, & Kroemer, 2023). Cellular senescence, a pivotal indicator of aging, is characterized by irreversible cell cycle arrest triggered by cellular damage or stress(Khosla, Farr, Tchkonia, & Kirkland, 2020). During cellular senescence, cells exhibit increased activity of the lysosomal enzyme senescence-associated β-galactosidase (SA-β-gal), decreased proliferation marker Ki67 signals and increased senescence-associated secretory phenotype (SASP) secretion(Coppe et al., 2008; Dimri et al., 1995).

Senescence is the result of a complex interplay of endogenous and exogenous signaling pathways(Cohen et al., 2022). Currently, the activation of tumor suppressor pathways, such as $p16^{INK4a}$/RB, is deemed essential for senescence-related cessation of proliferation(Baker et al., 2011). The disruption of signaling pathways disrupts tissue homeostasis, leading to proinflammatory responses that are increasingly recognized as fundamental aspects of senescence(Coppe et al., 2008; Shah et al., 2013). Researchers have suggested that the expression of p16 is suppressed by inhibitor of DNA-binding (ID) proteins. The expression of the *ID1* gene is downregulated by transforming growth factor-β (TGF-β), while it is upregulated by BMP-mediated phosphorylated SMAD signaling in various cell types(Hayashi et al., 2016; Liang, Brunicardi, & Lin, 2009). SMADs are crucial transcriptional effector factors responsible for transmitting TGF-β/BMP signals(Urist, 1965). They convey signals from the TGF-β superfamily and homomeric serine/threonine kinases on the cell surface to the nucleus, directly regulating transcriptional programs critical for key events(Miyazono, Maeda, & Imamura, 2005). Numerous studies have shown that the BMP-SMAD signaling pathway can modulate genes involved in the cell cycle, which affects the balance between cell proliferation and senescence(Genander et al., 2014; Hayashi et al., 2016). Phosphorylated SMAD plays a crucial role in the induction and maintenance of

pluripotency, helping maintain tissue homeostasis during aging(Orlowski, 2017).

However, dysregulation of BMP signaling during aging can lead to fibrosis, impaired

wound healing, or excessive tissue calcification, which are detrimental to tissue

function(Lyu et al., 2018; Rahman, Akhtar, Jamil, Banik, & Asaduzzaman, 2015).

Owing to increasing insights into cellular and individual aging, novel aging

mechanisms and signaling pathways have been identified, leading to the development

of new therapeutic interventions. Senescent cells have recently been targeted in therapy

given their substantial impact on aging (Childs et al., 2017). Consequently, eliminating

senescent cells through pharmacological interventions, known as "senolytics", may

delay the onset or slow the progression of age-related diseases and enhance

physiological function(Childs et al., 2017). Research has proven that specific senolytic

drugs, such as dasatinib and quercetin, can   eliminate senescent cells, reverse aging

phenotypes and increase the survival rate of mice(Xu et al., 2018; Zhu et al., 2015).

However, the current senolytics are not universally effective across different cell types

and   display   inconsistent   sensitivities,   thereby   limiting   their   broad

application(Yousefzadeh et al., 2018; Zhu et al., 2016). In addition to senolytics,

researchers have explored pharmacological interventions that aim to ameliorate or

reverse aging phenotypes without the need to eliminate senescent cells. This included

both the repurposing of existing medications and the development of new ones(Bharath

et al., 2020; Harrison et al., 2014). However, repurposing is particularly attractive due

to its cost-effectiveness; these drugs have already undergone rigorous safety evaluations,

allowing for quicker implementation in new contexts. For example, drugs such as

metformin, which is typically used to treat type 2 diabetes, and rapamycin, an

immunosuppressant, have shown promise in reducing aging phenotypes and extending

the lifespan(Bharath et al., 2020; Harrison et al., 2009). This strategy is an innovative

pharmacological intervention in which existing compounds target the aging process.

In this study, we employed a drug repurposing strategy to evaluate the anti-aging

potential of several FDA-approved drugs and identified enalapril as a promising

candidate. Our investigation revealed that enalapril increases pSmad1/5/9 levels, which

in turn upregulates key downstream genes involved in cell cycle regulation and

antioxidative defense, thereby improving cell proliferation, reducing inflammation, and

decreasing reactive oxygen species (ROS) levels. Our findings confirmed the anti-aging mechanism of enalapril at both the cellular and organismal levels. Enalapril has the ability to delay the aging of various organs, with notable effects on the brain, kidney, and liver. Furthermore, treating aged mice with enalapril improved spatial memory and anxiety behavior, alleviated lipid accumulation, and slowed kidney aging. Taken together, the potential of enalapril to extend the lifespan and ameliorate age-related diseases makes it a compelling candidate for research and development in the field of anti-aging therapeutics.

MATERIALS AND METHODS

Cell culture

IMR90 cells were cultured in DMEM (Gibco, C11995500) supplemented with 10% fetal bovine serum (Cellmax, SA111) and 1% non-essential amino acids (Gibco, 11140050) at 37°C and 5% CO_2. When the cell density in the in vitro culture reached 80–90% (usually within 2 or 3 days), cell passage was performed.

Drug preparation and addition

Enalapril (MCE, HY-B0331) was dissolved in DMSO to prepare a 10 mM stock solution. When used, it was diluted to 10 μM in the culture medium. After the cells adhered to the surface, enalapril was added for treatment, and RNA and protein were collected for analysis. Recombinant BMP4 protein (MCE, HY-P7007) was dissolved in sterile ddH_2O to prepare a 100 μg/mL stock solution. When used, it was diluted to 50 ng/mL in the culture medium and added to the cells. The BMP receptor inhibitor (LDN193189, MCE, HY-12071) was dissolved in sterile ddH_2O to prepare a 1 mM stock solution. When used, it was diluted to 1 μM or 0.5 μM in the culture medium. The ID inhibitor (Pimozide, MCE, HY-12987) was dissolved in DMSO to prepare a 10 mM stock solution. When used, it was diluted to 5 μM in the culture medium. After the cells adhered to the surface, drugs were added for a 3-day treatment, and samples were collected for analysis.

Senescence-associated β-galactosidase (SA-β-gal) Staining

SA-β-gal staining was performed using a Senescence β-Galactosidase Staining Kit (Sigma–Aldrich, CAS0030). Briefly, 0.6×10^5 cells were seeded in a 6-well plate. The cells were fixed with fixation buffer and incubated at room temperature for 10 minutes. After the cells were washed three times with PBS, the prepared staining mixture was added to the 6-well plate, and the plate was incubated at 37°C overnight. The stained cells were observed and photographed under a microscope.

Western blot

Proteins were extracted using TRIzol reagent (Invitrogen, 15596018CN) and dissolved in 1% SDS, after which SDS–PAGE was performed. The samples were transferred to nitrocellulose membranes and incubated with 5% skimmed milk at room temperature for 30 minutes. The primary antibodies were incubated overnight at 4°C, followed by three washes with PBST. The membranes were then incubated with secondary antibody at room temperature for two hours. The band signals were visualized using an Odyssey Infrared Imaging System (Odyssey, LI-COR).

RT-qPCR

Total RNA from cultured cells was extracted using TRIzol reagent (Invitrogen, 15596018CN) according to the manufacturer's instructions. The DEPC-treated water-dissolved RNA was subsequently reverse transcribed into cDNA using the HiScript III All-in-one RT SuperMix Perfect for qPCR (Vazyme, R333) following the product manual. The cDNA levels of specific primers were analyzed using the LightCycler® 96 qPCR system (Roche), and the results were normalized to those of β-actin. The primers used in this analysis are listed in Table S1.

RNA-seq and data processing

Total RNA was extracted using TRIzol reagent (Invitrogen, 15596018CN) following the manufacturer's instructions. The extracted total RNA was then sent to Novogene for subsequent cDNA library construction. The library was sequenced with paired-end reads on the Illumina HiSeq PE150 platform.

For data processing, FastQC (version 0.11.5) software was first used for data

quality control. Raw paired-end sequencing reads were trimmed by Trim Galore

(version 0.6.10) software to remove adapter sequences and low-quality reads. The high-

quality RNA-seq reads were mapped to the hg38/mm10 genome using HISAT2 (version

2.2.1) software(Kim, Paggi, Park, Bennett, & Salzberg, 2019). SAMtools (version 1.17)

software was then used to convert the generated SAM files to bam files(Li et al., 2009).

To qualify gene expression, the number of reads mapped to each gene was counted

using featureCounts (version 2.0.1) software(Liao, Smyth, & Shi, 2014). Differentially

expressed genes (DEGs) were calculated by R package DESeq2 (version 1.34.0)(Love,

Huber, & Anders, 2014) with the cutoff values of Benjamini-Hochberg adjusted p-value

< 0.05 and $|\log_2(\text{fold change})| > 0.5$. The gene expression count matrix was converted

into the reads per kilobase per million mapped reads (RPKM) matrix, followed by Z

score normalization for visualization.

CUT&Tag and data processing

For the CUT&Tag experiments, 8×10^5 cells were processed using the CUT&Tag Kit

(Vazyme, TD903). The cells were collected and incubated with ConA beads and then

mixed with the primary antibody pSmad1/5/9 (CST, #13820) overnight at 4°C. After

incubation with the secondary antibody for one hour, the cells were further incubated

with pA/G-Tnp for one hour. The transposase was then activated for DNA

fragmentation. Following DNA extraction, library amplification and purification were

performed. The libraries were sequenced using paired-end reads on the Illumina HiSeq

PE150 platform.

For data processing, raw CUT&Tag data was trimmed using Trim Galore (version

0.6.10) software, and the clean reads were mapped to the human reference genome hg38

using Bowtie2 (version 2.5.1) software(Langmead & Salzberg, 2012) with the

following parameters: "--very-sensitive --no-mixed --no-discordant -I 10 -X 1000". The

SAM files were transformed into bam files via SAMtools (version 1.17) software(Li et

al., 2009). The CUT&Tag peaks were performed using the MACS2 (version 2.2.8) with

the default parameters(Zhang et al., 2008). Differential peaks were calculated using the

R package DiffBind (version 3.4.11) with the cutoff values of adjusted p-value < 0.05

and |log₂(fold change)| > 0.5 and annotated using the R package ChIPpeakAnno

(version 3.28.0)(Lihua, Claude, & Nathan, 2010). To visualize the CUT&Tag signal,

the reads per kilobase per million mapped reads (RPKM) were calculated in each 50-

bp bin size using DeepTools (version 3.5.2) software(Ramirez, Dundar, Diehl, Gruning,

& Manke, 2014)· and the RPKM matrix was normalized using the Z score algorithm.

ChIP-qPCR

At room temperature, approximately 2×10^6 IMR90 cells were cross-linked with 1%

paraformaldehyde for 10 minutes. The cross-linking reaction was stopped by adding

0.125 M glycine for 6 minutes. The cells were then harvested and lysed on ice in nuclear

lysis buffer (50 mM Tris-Cl, 10 mM EDTA, 1% SDS, and protease inhibitors).

Chromatin immunoprecipitation (ChIP) was performed using a Bioruptor (Diagenode)

to shear the chromatin. Fragmented chromatin (25 μg) was immunoprecipitated with 5

μg of anti-pSmad1/5/9 antibody (CST, #13820) or anti-rabbit IgG (Abcam, ab171870).

The immunoprecipitated DNA was quantified via RT-qPCR. The amount of DNA in the

immunoprecipitates was calculated as a percentage relative to 10% of the input

chromatin. All ChIP-qPCR-related primers are available in Table S2.

Immunofluorescence

The slides were washed twice in PBS, fixed with 4% PFA for 10 minutes, and washed

again three times with PBS. Subsequently, permeabilization was performed via

incubation with 0.5% Triton X-100 for 10 minutes, followed by three additional washes

with PBS. Next, blocking was performed with 1% BSA, and after blocking, the slides

were washed three times with PBS. The primary antibody was then diluted at a ratio of

1:500 in PBST (PBS + 0.1% Tween 20) and incubated overnight at 4°C. Afterward, the

slides were washed four times with PBST, with each wash lasting 5 minutes. The

secondary antibody was diluted at a ratio of 1:500 in PBST (1 × PBS + 0.1% Triton X-

100) and incubated at room temperature for 1–1.5 hours. After three washes with PBST,

a final wash with PBS was conducted. DAPI was diluted to a final concentration of 1–

2 ng/μL and incubated at room temperature for 3–5 minutes. Following three washes

with PBS and one wash with ddH₂O, the slides were sealed with Fluoromount-G®

(SouthernBiotech, 0100-010) and covered with a coverslip.

Lentiviral production and viral transduction

First, the pLKO.1 empty vector or vectors encoding shRNAs targeting *ID1* or *ID2* (Table S3), along with the assistant plasmids psPAX2 and pMD2.G, were cotransfected into HEK293T cells. After 18 h of transfection, the medium was replaced with DMEM supplemented with 30% complement-inactivated FBS, and viral supernatants were collected after 24 h and 48 h of transfection. When IMR90 cells reached 70% confluence, they were infected with the packaged virus with 5 μg/mL polybrene (MCE, HY-K1057). At 36–48 hours after infection, successfully transduced cells were selected via treatment with 1 μg/mL puromycin (M&C Gene Technology, MA009) for 24–36 hours.

Detection of Reactive Oxygen Species (ROS)

IMR90 cells were harvested from a 6-well plate. After the cells were washed, the ROS-sensitive fluorescent probe 2',7'-dichlorofluorescin diacetate (DCFH-DA, MCE, HY-D0940) was used for detection. A 10 mM stock solution of DCFH-DA was prepared by dissolving it in anhydrous DMSO. This stock solution was subsequently diluted in serum-free DMEM to obtain a 10 μM working solution of DCFH-DA. The samples were incubated with the DCFH-DA probe at 37°C for 30 minutes. After incubation, the samples were washed to remove excess probe. The fluorescence intensity was measured via a fluorescence microscope. The fluorescence intensity corresponds to the level of ROS present in the sample.

Mice feeding

All the mouse experiments were approved by the Institutional Animal Care and Use Committee of Peking University. Twelve-month-old C57BL/6 mice (purchased from Wukong Biotech Co., Ltd.) were housed in a specific pathogen-free (SPF) animal facility under controlled environmental conditions with a temperature range of 20–25°C, 55±10% humidity, and a 12-h light/dark cycle. The mice had ad libitum access to food and water. A daily dosage of 30 mg/kg of enalapril was dissolved into the mice's drinking water. Four to ten mice were included in each group.

Forelimb grip strength test

The mice were placed on a grip strength meter, and their tails were gently pulled to allow them to grasp the pull bar with their forelimbs. The grip strength meter readings were recorded at the point of maximum force exerted by each mouse. The measurements were repeated multiple times to ensure reliable and consistent results.

Rotarod fatigue test

The mouse was placed on the rotarod, and rotation was initiated. The device was set to an accelerating mode that increased from 4 to 40 revolutions per minute over 300 seconds. The time at which each mouse fell off or lost balance was recorded. The test was repeated multiple times to assess the endurance and motor performance of the mice.

Y-maze test

The Y-maze consists of three opaque arms arranged at 120° angles relative to each other. The mice were placed in the central position of the maze and allowed to explore the three arms freely for 10 minutes. Each consecutive entry into one of the three arms was defined as spontaneous alternation behavior. The total number of arm entries and the number of spontaneous alternations were recorded. The percentage of spontaneous alternation behavior was then calculated.

Open field test

The mice were removed from their cages and placed in an open field, with the central area designated. The mice were allowed to move freely within the open field for 10 minutes, and the number of times that they crossed the central area was recorded.

Mouse serum extraction and analysis

After the mice were anesthetized, blood (approximately 1 mL per mouse) was collected from the eyeballs. The blood was allowed to clot and settle at room temperature. The mixture was then centrifuged at 3,000 rpm for 20 minutes, and the upper clear layer, which was the serum, was collected. The serum was subsequently subjected to biochemical assays, including ALT, CREA, LDH, TG, LDL, and HDL tests.

Mouse tissue extraction

After the mice were euthanized, they were perfused with prechilled PBS until no blood

was expelled. The heart, liver, spleen, kidney, brain, and muscle tissues were harvested.

A portion of each tissue sample was used for RNA and protein extraction for sequencing

and Western blot analysis, respectively, while another portion was fixed in 4%

paraformaldehyde (PFA). The tissues were then subjected to either frozen or paraffin

embedding. Various staining techniques, including β-galactosidase (β-gal) staining,

periodic acid-Schiff (PAS) staining, Congo red staining, Sirius red staining, Oil Red O

staining, and hematoxylin and eosin (H&E) staining, were performed.

Pathway enrichment analysis

Gene Ontology (GO) enrichment analysis was performed with Metascape(Zhou et al.,

2019), and gene set enrichment analysis (GSEA) was performed with the R package

ClusterProfiler (version 4.2.2)(Wu et al., 2021). Pathways with p values < 0.05 were

considered significantly enriched. Representative terms were visualized with the

ggplot2 (version 3.3.6) R package and ComplexHeatmap (version 2.10.0) R

package(Gu, 2022).

Statistical analysis

The data are expressed as means ± SEMs. The data were analyzed via a two-tailed

unpaired Student's t test. A p value < 0.05 was considered statistically significant.

RESULTS

Enalapril restrains cellular replicative senescence.

To identify drugs that are capable of delaying cellular senescence, we established an in

vitro replicative senescence model using primary human embryonic lung fibroblasts

(IMR90). Compared with young cells, long-term-cultured IMR90 cells presented a

characteristic senescence phenotype, characterized by increased SA-β-gal staining,

increased expression of the cell cycle arrest factors p16 and p21, and reduced expression

of the proliferation marker Ki67 (**Fig. S1A-C**). Additionally, the release of SASP factors

that significantly alter the cellular microenvironment, including the well-described

factors *IL-1β*, *IL-6*, and *CXCL10*, has also been reported to increase the degree of

senescence (**Fig. S1D**).

Employing a drug repurposing strategy, we discovered that enalapril, an
antihypertensive medication, delayed senescence. Notably, enalapril treatment at
various concentrations reduced SA-β-gal staining and increased Ki67 positivity in
IMR90 cells (**Fig. 1A-B, D-E**). Western blot analysis confirmed a decrease in the aging
markers p16 and p21 with the optimal effect of 10 μM enalapril (**Fig. 1C**). To further
analyze the effects of enalapril on transcriptome dynamics upon senescence entry,
RNA-seq was performed across enalapril-treated cells at different concentrations. In
general, we found that the expression of SASP genes decreased significantly (**Fig. 1F**).
The expression of cell cycle arrest factors typically increased during senescence;
however, the expression of these factors decreased after exposure to enalapril, whereas
the expression of cell cycle genes related to cell proliferation increased upon treatment
with enalapril at the optimal concentration of 10 μM (**Fig. 1F**). We therefore treated the
cells with 10 μM enalapril via RT-qPCR and detected that many prominent SASP genes,
such as *IL-1β and IL-6*, were significantly reduced compared with those in the control
(**Fig. 1G**). Moreover, enalapril treatment increased the cell growth rate and delayed the
transition to the growth arrest phase (**Fig. 1H**). Our comprehensive analysis underscores
enalapril's potential as an anti-senescence agent, warranting further exploration for its
role in combating cellular senescence.

The function of enalapril depends on phosphorylated Smad1/5/9.

Given the potential of enalapril to decelerate senescence, we screened several important
targets of the aging-related pathway to identify the underlying mechanism of
senescence entry. We found that enalapril notably elevated the phosphorylation level of
Smad1/5/9 without affecting Smad2/3 phosphorylation, suggesting a possible role for
the BMP-pSmad1/5/9 pathway (**Fig. 2A, Fig. S2A**). To validate the pathway alterations
in the gene expression profile, GSEA confirmed that enalapril treatment increased BMP
signaling (**Fig. 2B**). Concurrently, the level of the Smad4 protein, which is required for
pSmad1/5/9 to enter the nucleus, also increased, suggesting that enalapril might
facilitate the nuclear entry of pSmad1/5/9 and modulate downstream genes (**Fig.
2A**)(Guglielmi et al., 2021; Senft, Bikoff, Robertson, & Costello, 2019). To further

assess the impact on gene regulation, we employed CUT&Tag analysis of pSmad1/5/9 and found that enalapril treatment increased the pSmad1/5/9 binding signals at the transcription start site (TSS) regions (**Fig. 2C**).

To explore the potential role of pSmad1/5/9 in anti-senescence mechanisms, we experimentally manipulated its protein levels by supplementing with BMP4 to increase pSmad1/5/9 levels or employing the BMP receptor inhibitor LDN193189 (LDN) to reduce it. Our findings revealed that BMP4 increased pSmad5 levels, which correlated with decreased p16 and p21, reduced β-gal staining and increased Ki67 positivity, indicating a mitigated senescent phenotype (**Fig. S2B-D**). In contrast, when cells were treated with the BMP receptor inhibitor, the changes in pSmad1/5/9 levels and senescence markers were completely reversed, highlighting the accelerated progression of senescence under the influence of the BMP receptor inhibitor (**Fig. S2E-G**).

Encouraged by the association between increased pSmad1/5/9, we tested whether pSmad1/5/9 serves as an intermediary in the anti-senescence effects of enalapril by combining LDN with enalapril. RNA-seq analysis revealed that LDN upregulated SASP genes and downregulated cell cycle genes. On the basis of these findings, the subsequent addition of enalapril did not reverse these changes, which suggests that the anti-senescence function of enalapril relies on pSmad1/5/9 (**Fig. 2D**). Furthermore, western blot results also revealed that in the presence of LDN, neither enalapril nor BMP4 restored Smad1/5/9 phosphorylation or the expression of p16 and p21 (**Fig. 2E**). Similarly, β-gal and Ki67 staining also revealed that enalapril failed to counteract LDN-induced deterioration (**Fig. 2F, G**). Collectively, these results indicate that elevated pSmad1/5/9 could increase the expression of critical genes, thereby exerting a potent anti-senescence effect.

pSmad1/5/9 modulates the cell cycle and SASP secretion by upregulating ID1.

The ID family, which is recognized as a key downstream target of pSmad1/5/9, serves as a biomarker for Smad1/5/9 phosphorylation levels(Genander et al., 2014; Hayashi et al., 2016; Ramachandran et al., 2018; Sun et al., 2022; Ying, Nichols, Chambers, & Smith, 2003). Coincidentally, we found that ID1 was the top-ranked transcription factor that pSmad1/5/9 bound to its promoter region via CUT&Tag analysis after enalapril

treatment (**Fig. 3A**). Moreover, IGV analysis confirmed that enalapril significantly

increased pSmad1/5/9 enrichment at the *ID1* promoter (**Fig. 3B**). Simultaneously, the

expression levels of both *ID1* and *ID2* were elevated (**Fig. 3C, D**). Conversely,

inhibiting pSmad1/5/9 with LDN led to decreased *ID1* and *ID2* levels, confirming the

regulatory link between pSmad1/5/9 and the ID family (**Fig. 3E**). This finding indicated

that the ID family was indeed a downstream gene regulated by pSmad1/5/9, and the

increase in ID1 induced by enalapril further confirmed that enalapril could upregulate

pSmad1/5/9 levels, thereby affecting the expression of downstream genes.

Prior studies have indicated that the ID family can suppress p16 and p21

expression, thus inhibiting cell cycle arrest and delaying cellular senescence(Hayashi

et al., 2016; Lyden et al., 1999; Ying et al., 2003). This prompted us to examine ID

protein expression during senescence, where we observed a decrease in ID1 and ID2

levels (**Fig. S3A**). The knockdown of *ID1* and *ID2* via shRNA or an ID1 inhibitor

resulted in a decreasing trend of ID1 or ID2 expression and a significant increase in p16

and p21 expression (**Fig. 3F, I**). In addition, while β-gal increased the proportion of

positively stained cells, the Ki67 proliferation signal decreased nearly fourfold after the

knockdown of *ID1* or *ID2*, resembling the changes observed during senescence (**Fig.**

S3B, C). Given that enalapril reduced SASP factors while LDN increased them, we

investigated whether ID1, which acts as a transcriptional repressor downstream of

pSmad1/5/9, could modulate SASP expression. Indeed, the expression of traditional

SASP factors, such as *IL1β* and *IL6*, which decrease with enalapril treatment,

significantly increased after *ID* knockdown (**Fig. 3G**). By overlapping the genes

downregulated by enalapril with those upregulated upon *ID* knockdown, several SASP

genes, as well as *CDKN2A* and *CDKN2B,* were identified (**Fig. 3H**). Consistent with

these findings, we found that SASP factor expression also increased in *ID* knockout

mouse bone marrow cells in previously reported data (**Fig. S3E**)(Fei et al., 2023).

Importantly, *ID* knockdown did not affect pSmad1/5/9 levels, confirming that ID is a

downstream effector of pSmad1/5/9, influencing the expression of genes such as p16,

p21, and SASP-related genes (**Fig. 3J**).

After validating the prosenescence effects of *ID1* or *ID2* through ID1 knockdown

or the use of small molecule inhibitors, we next examined whether the anti-senescence

effects of enalapril are mediated by ID proteins. The continued addition of enalapril to

these *ID*-knockdown cells did not significantly alter β-gal or Ki67 signals compared

with those in the *ID*-knockdown group without enalapril, implying that the benefits of

enalapril may be ID dependent (**Fig. 3K-L, Fig. S3D**). Taken together, these findings

indicate that enalapril elevates pSmad1/5/9 levels, which in turn upregulate *ID1*

expression. This dual action has two key implications: first, it confirms that enalapril

can modulate the expression of pSmad1/5/9 downstream genes. Second, by

upregulating ID1, enalapril can suppress the expression of p16, p21, and SASP-related

genes, thereby potentially restoring cell proliferation and reducing the expression of

inflammatory factors. These findings suggest that the anti-senescence mechanism of

enalapril involves the pSmad1/5/9-ID1 axis, highlighting a novel pathway for its

therapeutic effects on aging.

Enalapril has antioxidative effects through pSmad1/5/9.

Since an increase in pSmad1/5/9 after enalapril treatment was observed and

pSmad1/5/9 is a transcription factor, to understand the in-depth mechanism of enalapril

regulation, we further explored the downstream genes of pSmad1/5/9. Analysis of

CUT&Tag-upregulated peaks after enalapril treatment revealed that enalapril treatment

led to enrichment of pSmad1/5/9 in the GO pathway associated with DNA replication,

the cell cycle and the oxidative stress response, all of which are crucial for cell growth

(**Fig. 4A**). Moreover, transcriptome analysis revealed that the genes whose expression

increased after enalapril treatment were also enriched in pathways related to the

response to oxidative stress (**Fig. 4A**). Notably, genes upregulated by enalapril in

response to oxidative stress included antioxidative enzymes, electron transport chain

components and lipid oxidation inhibitors, with a significant increase in pSmad1/5/9 at

their TSS regions (**Fig. 4B, C**). Specifically, we also examined the promoter regions of

antioxidative genes, such as the antioxidative enzyme *PRDX5* and the lipid oxidation

inhibitor *TXN*, and found increased pSmad1/5/9 enrichment along with upregulated

gene expression (**Fig. 4D, F-H**). During cellular senescence, a decrease in these

antioxidative genes exacerbates oxidative stress, reduces energy metabolism, and

increases lipid oxidation, leading to the release of inflammatory cytokines (**Fig.**

4E)(Coppe et al., 2008).

It has been reported that changes in the expression of antioxidative genes can affect

the ROS level of cells; thus, we then used DCFH-DA probes to detect ROS levels in

cells(Coppe et al., 2008). The results revealed that cellular senescence increased ROS,

whereas enalapril slowed the increase in ROS, indicating its ability to upregulate

antioxidative genes and reduce ROS, thus affecting cellular senescence (**Fig. 4I**).

Importantly, the continued addition of BMP receptor inhibitors reversed the beneficial

effects of enalapril on ROS and decreased pSmad1/5/9 enrichment of these

antioxidative genes (**Fig. 4G, I**). Collectively, these results reinforce the notion that

enalapril promotes the upregulation of antioxidative genes through pSmad1/5/9,

reduces ROS levels and alleviates oxidative stress, thus playing a crucial role in

delaying senescence.

Enalapril upregulates pSmad1/5/9 and improves physiological function in mice.

Enalapril, which is known for its anti-senescence potential, was further investigated for

its impact at the organ and individual levels in 12-month-old mice (**Fig. 5A**). After

continuous feeding at a dosage of 30 mg/kg/day for 3 months, we conducted a

comprehensive RNA-seq analysis of six key organs—brain, kidney, liver, heart, muscle,

and spleen—to assess the impact of enalapril on the transcriptome. We found that

enalapril induced organ-specific gene expression changes, thereby affecting the

physiological functions of these genes. Moreover, GSEA revealed upregulated

pathways related to forebrain and kidney development, long-chain fatty acid

metabolism and other processes, indicating that enalapril has a beneficial effect on

organ functions (**Fig. S4A**). Additionally, enalapril reduced the expression of SASP

factors in multiple organs, particularly the CCL, CXCL, and MMP families (**Fig. 5B**).

Serum SASP array analysis also revealed a decrease in the secretion of certain SASP

proteins, such as CCL6, CCL11, CXCL13, IGFBP-1, and MMP3, suggesting that

enalapril has anti-inflammatory effects in mice (**Fig. 5C**).

Given the known upregulation of pSmad1/5/9 levels and antioxidative genes in

cellular studies by enalapril, we measured Smad1/5/9 phosphorylation and the

expression of the antioxidative genes *Txn1*, *Txn2*, and *Prdx5* in the tested organs. We

observed increased pSmad1/5/9 levels, especially in the brain, kidney and liver, along

with upregulated *Id1* and antioxidative genes, notably *Txn1* and *Txn2* (**Fig. 5D, Fig.**

S4B). Further analysis of BMP signaling and antioxidative stress-related pathways in

these organs revealed increased gene expression after enalapril treatment, indicating the

activation of this pathway (**Fig. 5E**). These findings align with our cellular findings,

implying that enalapril may enhance the pSmad1/5/9-antioxidative gene axis to

increase organ physiological function and delay individual aging.

Enalapril alleviates aging-related behavior in naturally aged mice.

Building on our finding of pSmad1/5/9 activation and antioxidative genes upregulation

in multiple organs, we further investigated the broader phenotypic effects of enalapril

in mice. After three months of enalapril feeding, behavioral experiments, serum tests,

and tissue section staining were performed. Notably, enalapril had no significant effect

on body weight in the mice, suggesting that the drug has no overt toxic side effects (**Fig.**

6A). However, enalapril improved the decrease in age-related mobility, as evidenced

by increased grip strength and extended time in the rotarod fatigue experiments,

suggesting increased muscle strength and endurance (**Fig. 6D, E**).

During the aging process, the brain undergoes varying degrees of functional

decline. To assess the impact of enalapril on brain function, we conducted Y-maze and

open field tests. The Y-maze, which is frequently employed to assess short-term

memory in mice by measuring spontaneous alternation as a marker of spatial working

memory, indicates notable improvements. Under normal circumstances, mice prefer to

explore new maze branches rather than return to previously visited branches. Several

brain regions, including the hippocampus, septum, basal forebrain, and prefrontal

cortex, contribute to this task, and the age-related decline in spontaneous alternation

ability in mice is associated with degeneration or lesions in brain regions such as the

hippocampus(Radulescu, Cerar, Haslehurst, Kopanitsa, & Barnes, 2021). Enalapril

promoted a substantial increase in the rate of spontaneous alternation in the mice,

indicating enhanced spatial memory (**Fig. 6B, C**). As aging progresses, anxiety-like

behaviors often increase with age in mice(Radulescu et al., 2021). The open-field test,

a valuable tool for assessing autonomous and exploratory behaviors in new

environments, demonstrated that enalapril increased the number of central area
crossings, suggesting a reduction in aging-related anxiety (**Fig. 6F, G**). In conclusion,
these results suggest that enalapril has the potential to mitigate the deterioration of
various behavioral aspects associated with aging, including endurance, spatial memory
and anxiety.

Enalapril ameliorates aging-related pathological changes in aged mice.

The results of these behavioral tests provided valuable insights. To further explore the
effects of enalapril on pathological changes, our investigation was extended to serum
analyses. This comprehensive analysis encompassed the evaluation of vital organ
function, including the liver, kidneys, and heart, as well as an assessment of lipid
content, specifically triglycerides (TGs), low-density lipoprotein cholesterol (LDL),
and high-density lipoprotein cholesterol (HDL). After enalapril administration, there
was no significant change in the serum alanine transaminase (ALT) level, indicating
that no liver toxicity from enalapril occurred (**Fig. 6H**). Moreover, the level of
creatinine (CREA), a key marker of kidney function, was significantly decreased in
enalapril-treated mice, indicating potential functional recovery and improvement of
renal function (**Fig. 6I**). A heart health indicator, lactate dehydrogenase (LDH), was
also reduced, indicating a positive cardiac effect of enalapril (**Fig. 6J**). These findings
underscore the multifaceted effects of enalapril on various physiological parameters,
emphasizing its potential as a therapeutic intervention in aging-related conditions.

The transcriptomic data from the enalapril treatment group revealed an increase in
long-chain fatty acid metabolism in the liver, suggesting that enalapril may play a
positive role in lipid metabolism (**Fig. S4A**). Accordingly, we analyzed serum lipid
levels in enalapril-treated mice. Typically, aging leads to a decline in lipid metabolism
in the serum, causing an accumulation of triglycerides (TGs) and low-density
lipoprotein (LDL) and a decrease in high-density lipoprotein (HDL), contributing to
oxidative stress and conditions such as hyperlipidemia and fatty liver(A. A. Johnson &
Stolzing, 2019). Following enalapril treatment, the mice presented significant
reductions in TG and LDL levels and a noticeable increase in HDL levels (**Fig. 6K-M**).
These results suggest that enalapril has the potential to increase lipid metabolism in

aged mice, thereby mitigating the increase in blood lipid levels.

To evaluate the improvements in the aging process induced by enalapril in various

organ systems, we conducted pathological examinations of mouse tissues. A decrease

in β-gal staining was observed in the brain, liver and spleen of enalapril-treated mice,

suggesting a delay in organ aging (**Fig. 6N**). Given the effects of enalapril on brain

function, we further examined pathological markers in the brain. Periodic acid-Schiff

(PAS) staining, which is indicative of glycogen deposition, revealed a significant

reduction in glycogen near the hippocampus (**Fig. 6O**). Congo red staining, which

detects amyloid protein deposition, also revealed reduced staining in the cerebral cortex

(**Fig. 6P**). In the kidney, Sirius red staining, which highlights abnormal collagen fibers

or fibrosis, revealed significant red staining around blood vessels in control mice.

However, this staining was noticeably reduced in the kidneys of the mice treated with

enalapril (**Fig. 6Q**). These findings suggest that enalapril has a beneficial effect on

fibrosis in the kidneys of aged mice, which aligns with the observed improvement in

plasma creatinine levels (**Fig. 6I**). Furthermore, analysis of spleen, kidney and liver

sections from enalapril-treated mice revealed a reduction in Oil Red O staining, with

the kidneys and liver showing the most significant changes (**Fig. 6R**). This finding

aligned with the consistent decrease in lipid-related molecular content in plasma,

indicating that enalapril has a role in reducing age-related lipid accumulation (**Fig. 6K-**

M). Hematoxylin and eosin (H&E) staining of the liver revealed a decrease in cavities

and increased tissue density in mice treated with enalapril, suggesting its role in

enhancing liver cell proliferation and regeneration (**Fig. 6S**).

In summary, enalapril significantly mitigated multiple pathological markers

associated with aging, including lipid droplets, amyloid, glycogen deposition, and

fibrosis, in the brain, kidneys, liver, heart and spleen of aged mice. These

comprehensive improvements contribute to an overall amelioration of their aging status.

DISCUSSION

Since the early 2000s, the National Institutes of Health (NIH) has been pioneering the

Interventions Testing Program (ITP), an initiative aimed at harnessing existing drugs or

small molecules to decelerate the aging process(Harrison et al., 2009; Nadon et al., 2008;

Strong et al., 2016). This approach repurposes established medications for novel uses,
maximizing the potential of known compounds. It has accelerated the transition of anti-
aging therapies from discovery to clinical application, ensures their safety and bolsters
a comprehensive anti-aging drug screening strategy.

In our study, we discovered that enalapril, a commonly prescribed ACE inhibitor
for hypertension, has significant anti-aging effects at both the cellular and organismal
levels (**Fig. 7**). Enalapril demonstrated a notable ability to attenuate aging-related
physiological decline across multiple organs, including the brain, kidneys, and liver.
This finding suggests its potential to reduce the risk of disease across multiple organs,
slow the overall aging process and extend both the healthspan and lifespan of mice.
This observation aligns with previous reports on the lifespan-extending effects of other
ACE inhibitors, such as captopril, which have been shown to increase longevity in
organisms such as *C. elegans* and mice(Egan et al., 2024; Kumar, Dietrich, & Kornfeld,
2016; Strong et al., 2022). However, our screening process revealed that enalapril
outperformed captopril and other antihypertensive drugs in ameliorating senescent
phenotypes, although captopril and other antihypertensive drugs also exhibited the
ability to delay senescence. These findings indicate that enalapril may have unique or
more potent mechanisms of action that contribute to its anti-aging effects, warranting
further mechanistic studies to better understand its molecular targets.

Mechanistically, our data revealed that enalapril enhances the phosphorylation of
Smad1/5/9, a critical signaling pathway that regulates various cellular processes,
including cell proliferation and the oxidative stress response. Phosphorylation of
Smad1/5/9 facilitates its nuclear translocation, where it acts as a transcriptional
regulator to upregulate downstream genes, such as those involved in cell cycle
regulation and antioxidative responses. These findings suggest that the ability of
enalapril to modulate the pSmad1/5/9 pathway contributes to its protective effects
against oxidative stress, a well-established hallmark of aging. By reducing ROS levels
and promoting cell proliferation, enalapril effectively delays the progression of cellular
senescence and organismal aging. This finding not only highlights the potential of
phosphorylated Smad1/5/9 as a therapeutic target for anti-aging interventions but also
reveals a novel antioxidative mechanism of enalapril action.

Several FDA-approved drugs have been repurposed for anti-aging applications, demonstrating a wide range of mechanisms and effects. These include rapamycin, metformin, and NAD+ precursors. For example, rapamycin, which was originally used as an immunosuppressant to prevent organ transplant rejection, inhibits the mTOR pathway, which is associated with cell growth and metabolism(S. C. Johnson, Rabinovitch, & Kaeberlein, 2013). It extends the lifespan of mice and shows potential anti-aging effects across various species(Harrison et al., 2009). Compared with other repurposed anti-aging drugs, enalapril offers distinct advantages because of its unique mechanism of action. As an ACE inhibitor, enalapril inherently targets the renin–angiotensin system, providing vascular protection(Schmieder, Hilgers, Schlaich, & Schmidt, 2007; Yamamoto & Rakugi, 2021). Compared with repurposed drugs such as rapamycin, which may have more pronounced side effects, such as immunosuppression, limiting their long-term use, enalapril has a well-established safety profile(Janes & Fruman, 2009; Saunders, Metcalfe, & Nicholson, 2001). This can facilitate faster clinical translation for anti-aging applications. Moreover, our study revealed that enalapril reduces ROS through the pSmad1/5/9-antioxidant axis, highlighting its potential for antioxidation, a hallmark of aging that is less directly targeted by drugs such as rapamycin or metformin. Additionally, enalapril enhances multi-organ function, particularly in the brain, kidneys, and liver. Its well-established safety profile further supports enalapril as a promising candidate for clinical anti-aging applications, with the potential to complement existing therapies by promoting both vascular health and systemic antioxidative and anti-inflammatory effects.

Aging is a highly complex and multifactorial process(Campisi, 2013; Jazwinski, 1996). It involves intricate interactions among various signaling pathways and crosstalk between tissues and organs, especially during individual aging(Jazwinski, 1996; Munoz-Espin & Serrano, 2014). Notably, enalapril is not uniform across all organs and tissues, as indicated by the differential improvement in aging phenotypes among various tissues. For example, enalapril significantly improved the physiological functions of the brain, muscle and other organs, but the specific pathways and mechanisms by which these effects are mediated might differ among organs. This observation suggests that the tissue-specific effects of enalapril may be mediated by

differential activation of signaling pathways, including those beyond the Smad1/5/9
axis, which are known to play crucial roles in aging. Future studies should focus on
delineating these tissue-specific mechanisms to fully understand the scope of the anti-
aging effects of enalapril. Moreover, improvements in metabolic function in the liver
could have downstream effects on brain function, as metabolic dysregulation is known
to exacerbate neurodegenerative diseases. Similarly, enhanced vascular health resulting
from the antihypertensive properties of enalapril may contribute to improved brain
perfusion, reducing the risk of age-related cognitive decline. Therefore, exploring the
systemic effects of enalapril and how improvements in one organ system may benefit
others should be a priority for future research. Furthermore, given the established role
of enalapril as an antihypertensive drug, assessing its potential side effects when it is
repurposed for anti-aging purposes is essential. Identifying strategies to minimize these
side effects while optimizing their anti-aging effects is an important area of
investigation. In conclusion, our study identified enalapril as a promising candidate for
repurposing as an anti-aging drug with antioxidative and promoting effects on cell
proliferation. These findings open avenues for the development of targeted therapies
aimed at mitigating aging and promoting healthy longevity in clinical settings,
providing a foundation for guiding future research in the realm of anti-aging drug
interventions.

DATA AVAILABILITY

All related sequencing data have been uploaded to the NCBI Gene Expression Omnibus and are accessible through accession numbers GSE277861 and GSE277862.

ACKNOWLEDGMENTS

We thank Dr. Xuehui Lyu (Peking University) and Shuo Chen (Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences) for suggestions regarding data organization. We thank the National Center for Protein Sciences at Peking University for assisting with the cDNA library and the Core Facilities of Life Sciences, Peking University, for assisting with the confocal microscopy. We are also grateful to the Behavioral Platform of the Animal Center at Peking University for assisting with the behavioral experiments involving the mice.

AUTHOR CONTRIBUTIONS

W.T. oversaw the study and obtained financial support. W.C.L. conducted the main bench experiments. H.C.W. analyzed the data. Z.H.D. and J.N.H. helped prepare the mice samples. R.W., F.J.M. and J.L.B. helped analyze the data. L.J.Z. helped guide the experiment. W.C.L., Y. T. G. and C. Z. prepared the original manuscripts. All the authors read and approved the final version.

FUNDING

This work was supported by the National Key Research and Development Program of China (2021YFA0909300).

COMPETING INTEREST DECLARATION

The authors have no conflicts of interest to declare.

SUPPLEMENTAL INFORMATION

Additional supplementary information includes supplementary figures and tables in the attachment.

REFERENCES

Baker, D. J., Wijshake, T., Tchkonia, T., LeBrasseur, N. K., Childs, B. G., van de Sluis, B., . . . van Deursen, J. M. (2011). Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. *Nature, 479*(7372), 232-236. doi:10.1038/nature10600

Bharath, L. P., Agrawal, M., McCambridge, G., Nicholas, D. A., Hasturk, H., Liu, J., . . . Nikolajczyk, B. S. (2020). Metformin Enhances Autophagy and Normalizes Mitochondrial Function to Alleviate Aging-Associated Inflammation. *Cell Metab, 32*(1), 44-55 e46. doi:10.1016/j.cmet.2020.04.015

Campisi, J. (2013). Aging, cellular senescence, and cancer. *Annu Rev Physiol, 75*, 685-705. doi:10.1146/annurev-physiol-030212-183653

Childs, B. G., Gluscevic, M., Baker, D. J., Laberge, R. M., Marquess, D., Dananberg, J., & van Deursen, J. M. (2017). Senescent cells: an emerging target for diseases of ageing. *Nat Rev Drug Discov, 16*(10), 718-735. doi:10.1038/nrd.2017.116

Cohen, A. A., Ferrucci, L., Fülöp, T., Gravel, D., Hao, N., Kriete, A., . . . Varadhan, R. (2022). A complex systems approach to aging biology. *Nature Aging, 2*(7), 580-591. doi:10.1038/s43587-022-00252-6

Coppe, J. P., Patil, C. K., Rodier, F., Sun, Y., Munoz, D. P., Goldstein, J., . . . Campisi, J. (2008). Senescence-associated secretory phenotypes reveal cell-nonautonomous functions of oncogenic RAS and the p53 tumor suppressor. *PLoS Biol, 6*(12), 2853-2868. doi:10.1371/journal.pbio.0060301

Dimri, G. P., Lee, X., Basile, G., Acosta, M., Scott, G., Roskelley, C., . . . et al. (1995). A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A, 92*(20), 9363-9367. doi:10.1073/pnas.92.20.9363

Egan, B. M., Pohl, F., Anderson, X., Williams, S. C., Gregory Adodo, I., Hunt, P., . . . Kornfeld, K. (2024). The ACE inhibitor captopril inhibits ACN-1 to control dauer formation and aging. *Development, 151*(3). doi:10.1242/dev.202146

Fei, M. Y., Wang, Y., Chang, B. H., Xue, K., Dong, F. Y., Huang, D., . . . Wang, L. (2023). The non-cell-autonomous function of ID1 promotes AML progression via ANGPTL7 from the microenvironment. *Blood, 142*(10), 903-917. doi:10.1182/blood.2022019537

Genander, M., Cook, P. J., Ramsköld, D., Keyes, B. E., Mertz, A. F., Sandberg, R., & Fuchs, E. (2014). BMP Signaling and Its pSMAD1/5 Target Genes Differentially Regulate Hair Follicle Stem Cell Lineages. *Cell Stem Cell, 15*(5), 619-633. doi:10.1016/j.stem.2014.09.009

Gu, Z. (2022). Complex heatmap visualization. *Imeta, 1*(3), e43. doi:10.1002/imt2.43

Guglielmi, L., Heliot, C., Kumar, S., Alexandrov, Y., Gori, I., Papaleonidopoulou, F., . . . Hill, C. S. (2021). Smad4 controls signaling robustness and morphogenesis by differentially contributing to the Nodal and BMP pathways. *Nature Communications, 12*(1). doi:ARTN 6374 10.1038/s41467-021-26486-3

Harrison, D. E., Strong, R., Allison, D. B., Ames, B. N., Astle, C. M., Atamna, H., . . . Miller, R. A.
(2014). Acarbose, 17-α-estradiol, and nordihydroguaiaretic acid extend mouse lifespan
preferentially in males. *Aging Cell, 13*(2), 273-282. doi:10.1111/acel.12170

Harrison, D. E., Strong, R., Sharp, Z. D., Nelson, J. F., Astle, C. M., Flurkey, K., . . . Miller, R. A.
(2009). Rapamycin fed late in life extends lifespan in genetically heterogeneous mice.
Nature, 460(7253), 392-U108. doi:10.1038/nature08221

Hayashi, Y., Hsiao, E. C., Sami, S., Lancero, M., Schlieve, C. R., Nguyen, T., . . . Yamanaka, S.
(2016). BMP-SMAD-ID promotes reprogramming to pluripotency by inhibiting
p16/INK4A-dependent senescence. *Proc Natl Acad Sci U S A, 113*(46), 13057-13062.
doi:10.1073/pnas.1603668113

Janes, M. R., & Fruman, D. A. (2009). Immune Regulation by Rapamycin: Moving Beyond T Cells.
Science Signaling, 2(67). doi:ARTN pe25 10.1126/scisignal.267pe25

Jazwinski, S. M. (1996). Longevity, genes, and aging. *Science, 273*(5271), 54-59.
doi:10.1126/science.273.5271.54

Johnson, A. A., & Stolzing, A. (2019). The role of lipid metabolism in aging, lifespan regulation,
and age-related disease. *Aging Cell, 18*(6). doi:ARTN e13048 10.1111/acel.13048

Johnson, S. C., Rabinovitch, P. S., & Kaeberlein, M. (2013). mTOR is a key modulator of ageing
and age-related disease. *Nature, 493*(7432), 338-345. doi:10.1038/nature11861

Khosla, S., Farr, J. N., Tchkonia, T., & Kirkland, J. L. (2020). The role of cellular senescence in
ageing and endocrine disease. *Nat Rev Endocrinol, 16*(5), 263-275. doi:10.1038/s41574-
020-0335-y

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol, 37*(8), 907-
915. doi:10.1038/s41587-019-0201-4

Kirkland, J. L. (2016). Translating the Science of Aging into Therapeutic Interventions. *Cold Spring
Harb Perspect Med, 6*(3), a025908. doi:10.1101/cshperspect.a025908

Kumar, S., Dietrich, N., & Kornfeld, K. (2016). Angiotensin Converting Enzyme (ACE) Inhibitor
Extends Caenorhabditis elegans Life Span. *PLoS Genet, 12*(2), e1005866.
doi:10.1371/journal.pgen.1005866

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods,
9*(4), 357-359. doi:10.1038/nmeth.1923

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., . . . Genome Project Data
Processing, S. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics,
25*(16), 2078-2079. doi:10.1093/bioinformatics/btp352

Liang, Y. Y., Brunicardi, F. C., & Lin, X. (2009). Smad3 mediates immediate early induction of Id1
by TGF-beta. *Cell Res, 19*(1), 140-148. doi:10.1038/cr.2008.321

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics, 30*(7), 923-930.

700   doi:10.1093/bioinformatics/btt656

701   Lihua, Z., Claude, G., & Nathan, L. (2010). ChIPpeakAnno: a Bioconductor package to annotate
702   ChIP-seq and ChIP-chip data. *BMC Bioinformatics, 11*, 237.

703   Lopez-Otin, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2023). Hallmarks of
704   aging: An expanding universe. *Cell, 186*(2), 243-278. doi:10.1016/j.cell.2022.11.001

705   Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion
706   for RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-014-0550-
707   8

708   Lyden, D., Young, A. Z., Zagzag, D., Yan, W., Gerald, W., O'Reilly, R., . . . Benezra, R. (1999). Id1
709   and Id3 are required for neurogenesis, angiogenesis and vascularization of tumour
710   xenografts. *Nature, 401*(6754), 670-677. doi:10.1038/44334

711   Lyu, G. L., Guan, Y. T., Zhang, C., Zong, L., Sun, L., Huang, X. K., . . . Tao, W. (2018). TGF-β
712   signaling alters H4K20me3 status via miR-29 and contributes to cellular senescence and
713   cardiac aging (vol 9, 2560, 2018). *Nature Communications, 9*. doi:ARTN 4134
714   10.1038/s41467-018-06710-3

715   Miyazono, K., Maeda, S., & Imamura, T. (2005). BMP receptor signaling: Transcriptional targets,
716   regulation of signals, and signaling cross-talk. *Cytokine & Growth Factor Reviews, 16*(3),
717   251-263. doi:10.1016/j.cytogfr.2005.01.009

718   Munoz-Espin, D., & Serrano, M. (2014). Cellular senescence: from physiology to pathology. *Nat
719   Rev Mol Cell Biol, 15*(7), 482-496. doi:10.1038/nrm3823

720   Nadon, N. L., Strong, R., Miller, R. A., Nelson, J., Javors, M., Sharp, Z. D., . . . Harrison, D. E.
721   (2008). Design of aging intervention studies: the NIA interventions testing program. *Age,
722   30*(4), 187-199. doi:10.1007/s11357-008-9048-1

723   Orlowski, J. (2017). SMAD5 signaling: more than meets the nuclei. *Cell Res, 27*(9), 1075-1076.
724   doi:10.1038/cr.2017.101

725   Radulescu, C. I., Cerar, V., Haslehurst, P., Kopanitsa, M., & Barnes, S. J. (2021). The aging mouse
726   brain: cognition, connectivity and calcium. *Cell Calcium, 94*, 102358.
727   doi:10.1016/j.ceca.2021.102358

728   Rahman, M. S., Akhtar, N., Jamil, H. M., Banik, R. S., & Asaduzzaman, S. M. (2015). TGF-β/BMP
729   signaling and other molecular events: regulation of osteoblastogenesis and bone formation.
730   *Bone Research, 3*. doi:ARTN 15005 10.1038/boneres.2015.5

731   Ramachandran, A., Vizán, P., Das, D., Chakravarty, P., Vogt, J., Rogers, K. W., . . . Hill, C. S. (2018).
732   TGF-β uses a novel mode of receptor activation to phosphorylate SMAD1/5 and induce
733   epithelial-to-mesenchymal transition. *Elife, 7*. doi:ARTN e31756 10.7554/eLife.31756

734   Ramirez, F., Dundar, F., Diehl, S., Gruning, B. A., & Manke, T. (2014). deepTools: a flexible
735   platform for exploring deep-sequencing data. *Nucleic Acids Res, 42*(Web Server issue),
736   W187-191. doi:10.1093/nar/gku365

Saunders, R. N., Metcalfe, M. S., & Nicholson, M. L. (2001). Rapamycin in transplantation: A

review of the evidence. *Kidney International, 59*(1), 3-16. doi:DOI 10.1046/j.1523-
1755.2001.00460.x

Schmieder, R. E., Hilgers, K. F., Schlaich, M. P., & Schmidt, B. M. W. (2007). Renin-angiotensin
system and cardiovascular risk. *Lancet, 369*(9568), 1208-1219. doi:10.1016/S0140-
6736(07)60242-6

Senft, A. D., Bikoff, E. K., Robertson, E. J., & Costello, I. (2019). Genetic dissection of Nodal and
Bmp signalling requirements during primordial germ cell development in mouse. *Nature
Communications, 10*. doi:ARTN 1089 10.1038/s41467-019-09052-w

Shah, P. P., Donahue, G., Otte, G. L., Capell, B. C., Nelson, D. M., Cao, K., . . . Berger, S. L. (2013).
Lamin B1 depletion in senescent cells triggers large-scale changes in gene expression and
the chromatin landscape. *Genes Dev, 27*(16), 1787-1799. doi:10.1101/gad.223834.113

Strong, R., Miller, R. A., Antebi, A., Astle, C. M., Bogue, M., Denzel, M. S., . . . Harrison, D. E.
(2016). Longer lifespan in male mice treated with a weakly estrogenic agonist, an
antioxidant, an -glucosidase inhibitor or a Nrf2-inducer. *Aging Cell, 15*(5), 872-884.
doi:10.1111/acel.12496

Strong, R., Miller, R. A., Cheng, C. J., Nelson, J. F., Gelfond, J., Allani, S. K., . . . Harrison, D. E.
(2022). Lifespan benefits for the combination of rapamycin plus acarbose and for captopril
in genetically heterogeneous mice. *Aging Cell, 21*(12). doi:10.1111/acel.13724

Sun, Y., Yan, K., Wang, Y., Xu, C., Wang, D., Zhou, W., . . . Xi, Q. R. (2022). Context-dependent
tumor-suppressive BMP signaling in diffuse intrinsic pontine glioma regulates stemness
through epigenetic regulation of CXXC5. *Nature Cancer, 3*(9), 1105-+.
doi:10.1038/s43018-022-00408-8

Urist, M. R. (1965). Bone - Formation by Autoinduction. *Science, 150*(3698), 893-+. doi:DOI
10.1126/science.150.3698.893

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., . . . Yu, G. (2021). clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. *Innovation (Camb), 2*(3), 100141.
doi:10.1016/j.xinn.2021.100141

Xu, M., Pirtskhalava, T., Farr, J. N., Weigand, B. M., Palmer, A. K., Weivoda, M. M., . . . Kirkland,
J. L. (2018). Senolytics improve physical function and increase lifespan in old age. *Nat
Med, 24*(8), 1246-1256. doi:10.1038/s41591-018-0092-9

Yamamoto, K., & Rakugi, H. (2021). Angiotensin receptor-neprilysin inhibitors: Comprehensive
review and implications in hypertension treatment. *Hypertension Research, 44*(10), 1239-
1250. doi:10.1038/s41440-021-00706-1

Ying, Q. L., Nichols, J., Chambers, I., & Smith, A. (2003). BMP induction of Id proteins suppresses
differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3.
Cell, 115(3), 281-292. doi:Doi 10.1016/S0092-8674(03)00847-X

Yousefzadeh, M. J., Zhu, Y., McGowan, S. J., Angelini, L., Fuhrmann-Stroissnigg, H., Xu, M., . . .
Niedernhofer, L. J. (2018). Fisetin is a senotherapeutic that extends health and lifespan.
Ebiomedicine, 36, 18-28. doi:10.1016/j.ebiom.2018.09.015

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., . . . Liu, X. S. (2008).
Model-based analysis of ChIP-Seq (MACS). *Genome Biol, 9*(9), R137. doi:10.1186/gb-
2008-9-9-r137

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., . . . Chanda, S. K.
(2019). Metascape provides a biologist-oriented resource for the analysis of systems-level
datasets. *Nat Commun, 10*(1), 1523. doi:10.1038/s41467-019-09234-6

Zhu, Y., Tchkonia, T., Fuhrmann-Stroissnigg, H., Dai, H. M. M., Ling, Y. Y. Y., Stout, M. B., . . .
Kirkland, J. L. (2016). Identification of a novel senolytic agent, navitoclax, targeting the
Bcl-2 family of anti-apoptotic factors. *Aging Cell, 15*(3), 428-435. doi:10.1111/acel.12445

Zhu, Y., Tchkonia, T., Pirtskhalava, T., Gower, A. C., Ding, H. S., Giorgadze, N., . . . Kirkland, J. L.
(2015). The Achilles' heel of senescent cells: from transcriptome to senolytic drugs. *Aging
Cell, 14*(4), 644-658. doi:10.1111/acel.12344

FIGURE AND FIGURE LEGENDS

Figure 1 Enalapril alleviates cellular senescence. (A, D) SA-β-Gal staining **(A)** and statistical analysis of SA-β-Gal ratios **(D)** in IMR90 cells after enalapril treatment. Scale

bars, 200 μm. Enlarged scale bars, 100 μm. (**B, E**) Ki67 immunofluorescence

experiment (**B**) and statistical analysis of Ki67 intensity (**E**) after enalapril treatment.

Scale bars, 80 μm. Enlarged scale bars, 40 μm. (**C**) Western blot analysis showing the

protein levels of p16 and p21 in IMR90 cells after enalapril treatment. (**F**) Heatmap

showing cell cycle-related gene (green) upregulation and SASP gene (purple)

downregulation after enalapril treatment. (**G**) RNA expression of SASP factors in

IMR90 cells after enalapril treatment, with blue bars representing the control group and

red bars representing the enalapril-treated group. (**H**) Growth curves of IMR90 cells

after enalapril treatment. The x-axis represents the passage time, while the y-axis

represents population doubling (PD). A two-tailed t test was employed; ns indicates no

significant difference, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure 2 Phosphorylated Smad1/5/9 mediate the anti-senescence effect of enalapril.
(**A**) Western blot results for the detection of pSmad5, pSmad1/5/9 and other proteins
after enalapril treatment. (**B**) GSEA showing the enrichment of the BMP signaling
pathway after enalapril treatment. NES, normalized enrichment score. ES, enrichment
score. (**C**) Profile of pSmad1/5/9 enrichment at the TSS regions in the control group
(Ctrl) and the enalapril-treated group (Enalapril). (**D**) Heatmap of cell cycle (green) and
SASP-related genes (purple) in the control group (Ctrl), BMP receptor inhibitor-treated
group (LDN193189, LDN), and enalapril and BMP receptor inhibitor-cotreated group

(EP+LDN). (**E**) Western blot analysis showing the protein levels of pSmad1/5/9, p16,
and p21 in response to different combinations of enalapril, a BMP receptor inhibitor
(LDN193189, LDN), and BMP4. (**F**) SA-β-Gal staining (left) and corresponding ratio
statistical chart (right) of IMR90 cells treated with different combinations of enalapril,
a BMP receptor inhibitor (LDN193189, LDN), and BMP4. Scale bars, 200 μm.
Enlarged scale bars, 100 μm. (**G**) Ki67 immunofluorescence experiment (left) and
intensity statistical chart (right) of IMR90 cells treated with different combinations of
enalapril, a BMP receptor inhibitor (LDN193189, LDN), and BMP4. Scale bars, 80 μm.
Enlarged scale bars, 40 μm. A two-tailed t test was employed; ns indicates no significant
difference, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Figure 3 pSmad1/5/9 regulate ID-induced inhibition of p16, p21 and SASP. (A)
Changes in the peaks of all transcription factors identified by CUT&Tag following
enalapril treatment, with red dots indicating transcription factors with increased peak
intensity (Upregulated TFs). (**B**) Integrative Genomics Viewer (IGV) showing
pSmad1/5/9 signals near the *ID1* region between the control group (Ctrl) and the
enalapril treatment group (Enalapril). The vertical yellow boxes indicate regions with
increased signal intensity. (**C**) Western blot analysis showing the changes in the protein
levels of ID1 and ID2 following enalapril treatment. (**D**) Changes in the RNA levels of
ID1 and ID2 following enalapril treatment. (**E**) Western blot analysis showing the
protein levels of ID1 and ID2 following treatment with a BMP receptor inhibitor

(LDN193189, LDN). (**F, I**) Western blot showing the changes in the protein levels of

p16 and p21 following the knockdown of *ID1* or *ID2* (**F**) or the inhibition of ID1 (**I**).

(**G**) RNA expression of SASP factors after *ID* knockdown, with pink representing *ID1*

knockdown and orange representing *ID2* knockdown. (**H**) Normalized average RNA

expression levels of selected SASP factors and cell cycle arrest factors after enalapril

treatment and *ID* knockdown. Positive values indicate upregulation, while negative

values indicate downregulation. The values represent the expression levels relative to

those of the Ctrl. (**J**) Western blot analysis showing pSmad1/5/9 levels following *ID1*

and *ID2* knockdown. (**K**) SA-β-Gal staining (left) and SA-β-Gal ratio quantification

(right) in the control group (Ctrl) and *ID*-knockdown groups (ID1_KD, ID2_KD) with

or without enalapril treatment. Scale bars, 200 μm. Enlarged scale bars, 100 μm. (L)

Ki67 immunofluorescence intensity in the control group (Ctrl) and *ID* knockdown

groups (ID1_KD, ID2_KD), with or without enalapril treatment. A two-tailed t test was

employed; ns indicates no significant difference, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.

Figure 4 pSmad1/5/9 increase the expression of antioxidative genes. (**A**) Bar plot showing the enriched GO terms and pathways associated with the CUT&Tag-upregulated peaks (left) and the upregulated RNA-seq genes (right) following enalapril treatment. (**B**) Heatmap showing the changes in the RNA expression of antioxidative

genes. (**C**) Profile plot showing the increase in the pSmad1/5/9 binding signal of

antioxidative genes in the TSS region following enalapril treatment. (**D**) Western blot

analysis showing the protein levels of representative antioxidative genes after enalapril

treatment. (**E**) Relative RNA levels of *TXN*, *GPX4*, and *PRDX5* during cellular

senescence. The blue bars represent young cells, and the red bars represent senescent

cells. (**F, H**) Integrative Genomics Viewer (IGV) showing pSmad1/5/9 signals near the

PRDX5 (**F**) and *TXN* (**H**) regions between the control group (Ctrl) and the enalapril

treatment group (Enalapril). The vertical yellow boxes indicate regions with increased

signal intensity. (**G**) ChIP-qPCR results showing pSmad1/5/9 levels at many peaks

following enalapril or combination treatment with enalapril and LDN193189. The y-

axis represents the normalized pSmad1/5/9 signals relative to 10% input. pSmad1/5/9

enrichment for *HPRT1* and *HBB* served as a negative control, and pSmad1/5/9

enrichment for *ID1* and *ID2* served as positive controls, as previously described. A two-

tailed t test was employed, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. (**I**)

Detection of ROS fluorescence levels via the DCFH-DA probe in young cells, senescent

cells (Ctrl), cells treated with enalapril, and cells treated with a combination of enalapril

and LDN193189. Scale bars, 200 μm.

Figure 5 Enalapril exerts its anti-aging effects on mouse organs via pSmad1/5/9-antioxidative genes. (**A**) A schematic workflow of the mouse feeding and experimental procedures. Created in BioRender. Li, D. (2024) BioRender.com/u88h339 (**B**) Heatmap of RNA levels of SASP factors in various organs following enalapril treatment (EP). (**C**) Cytokine array analysis of secreted proteins (below) and relative quantitation

877   (above) of SASP factors in control serum (Ctrl) and enalapril-treated serum (Enalapril).

878   (**D**) Western blot analysis showing the protein levels of pSmad1/5/9 and antioxidative

879   genes in various organs following enalapril treatment. (**E**) Heatmap of the RNA levels

880   of BMP signaling pathway-related genes and antioxidative genes in the brain, kidney,

881   and liver following enalapril (EP) treatment.

Figure 6 Enalapril mitigates age-related degenerative changes in aged mice. (**A**) Changes in the body weights of the mice after enalapril feeding. (**B, C**) Movement

trajectory (**B**) and spontaneous alternation rate (**C**) in the Y-maze of the mice after enalapril treatment. (**D, E**) Changes in the grip strength (**D**) and rotarod fall time (**E**) of the mice after enalapril treatment. (**F, G**) Heatmap of movement trajectories (**F**) and the number of entries into the central area (**G**) in the open-field test after enalapril treatment. (**H–M**) Alterations in the serum levels of aspartate transaminase (AST) (**H**), creatinine (CREA) (**I**), lactate dehydrogenase (LDH) (**J**), triglycerides (TGs) (**K**), low-density lipoprotein cholesterol (LDL) (**L**), and high-density lipoprotein cholesterol (HDL) (**M**) after enalapril treatment in mice. (**N**) SA-β-Gal staining of the brain, liver, and spleen after enalapril treatment. Scale bars, 20 μm. (**O, P**) Periodic Acid-Schiff (PAS) staining (**O**) and Congo red staining (**P**) of the brain after enalapril treatment. Scale bars, 50 μm. (**Q**) Sirius red staining of the kidney after enalapril treatment. Scale bars, 50 μm. (**R**) Oil Red O staining of the spleen, kidney and liver after enalapril treatment. Scale bars, 20 μm. (**S**) Hematoxylin and eosin (H&E) staining of the liver after enalapril treatment. Scale bars, 50 μm.

Figure 7 Summary schematic. It showing pSmad1/5/9-mediated combating cellular senescence of enalapril, and improvement of organ function in mice. Specifically, enalapril increases the level of pSmad1/5/9 and Smad4, promoting the expression of downstream genes, such as antioxidative and cell cycle related genes, thereby promoting cell proliferation and reducing ROS. Moreover, enalapril mitigates age-related degenerative changes, including enhanced memory, improved kidney function, increased liver metabolism and other organ functions. Created in BioRender. Li, D. (2024) BioRender.com/u88h339.

A SA-β-Gal Staining

Enalapril (μM) 0 5 10 20 40

B Immunofluorescence of IMR90 (tagged Ki67)

Ki67

DAPI

Enalapril (μM) 0 5 10 20 40

C

Enalapril (μM)
0 5 10 20 40

p16

p21

β-actin

D SA-β-Gal Staining

The percentage of positive cells (%)

0 5 10 20 40 (μM)

E Immunofluorescence of IMR90 (tagged Ki67)

The percentage of positive cells (%)

0 5 10 20 40 (μM)

F Heatmap of Cell Cycle and SASP genes

RNA-seq

CIT
MCM2
CDC7
FBXO5
CDK4
MCM7
CDT1
DDX11
MCM5

IL6
CDKN1B
CDKN2A
FOS
CXCL5
CDKN2B
CXCL8
ADAM17
STAT3
NFKB1

Z-score
3
2
1
-1
-2

Ctrl-1 Ctrl-2 5μM-1 5μM-2 10μM-1 10μM-2 20μM-1 20μM-2 40μM-1 40μM-2

G RT-qPCR for SASP factors

Relative expression

0μM 10μM

IL1β IL6 CXCL10 CXCL16 CCL2 MMP2

H The growth curve of IMR90

PD

0μM 10μM

Enalapril

Time (d)
0 3 6 9 12 15

A GO terms and pathways

CUT&Tag of pSmad1/5/9 signal upregulated peaks

- DNA replication — LgPvalue
- SMAD binding
- Positive regulation of cell adhesion
- Chromatin organization
- Cell growth
- Extracellular matrix assembly
- **Cellular response to oxidative stress**
- Cell cycle

RNA-seq upregulated genes

- DNA replication — LgPvalue
- Cell cycle
- SMAD binding
- Extracellular matrix assembly
- Positive regulation of cell adhesion
- Response to oxidative stress
- **Cellular response to oxidative stress**
- BMP signaling pathway

B Heatmap of anti-oxidative stress genes

C CUT&Tag of pSmad1/5/9 signal in antioxidative stress genes

Ctrl / Enalapril

D Enalapril (μM) 0 / 10

PRDX5, TXN, NDUFB10, GPX4, β-actin

E RT-qPCR for antioxidative genes in senescent IMR90

Young / Old

TXN, GPX4, PRDX5

F CUT&Tag: pSmad1/5/9 signal

Ctrl-1, Ctrl-2, Enalapril-1, Enalapril-2, Chromatin State, Refseq Genes

PRDX5

G ChIP-qPCR for pSmad1/5/9 binding genes

Ctrl / Enalapril / Enalapril + LDN

HPRT1, HBB, ID1, ID2, TXN, PRDX5, GPX4, SOD3

H CUT&Tag: pSmad1/5/9 signal

Ctrl-1, Ctrl-2, Enalapril-1, Enalapril-2, Chromatin State, Refseq Genes

TXN

I ROS fluorescence level (using DCFN-DA probe)

Young / Ctrl / Enalapril / Enalapril+LDN