Changes in interlimb coordination induced by within-stride changes in treadmill speed

Brooke L. Hall[1], Ryan T. Roemmich[1,2]*, Caitlin L. Banks[1,2]*

[1]Center for Movement Studies, Kennedy Krieger Institute, Baltimore, MD, USA

[2]Department of Physical Medicine and Rehabilitation, Johns Hopkins University School of Medicine,

Baltimore, MD, USA

*designates co-senior authors

X handles: @RyanRoemmich (RTR), @Biomch_LBanks (CLB)

ORCiDs: https://orcid.org/0000-0003-0797-6455 (RTR), https://orcid.org/0000-0003-1500-199X (CLB),

https://orcid.org/0009-0001-2611-7675 (BLH)

Contact information:

Room 240

Kennedy Krieger Institute

716 North Broadway

Baltimore, MD 21205

rroemmi1@jhmi.edu

Abstract word count: 246   Text word count: 7274

Number of figures: 5   Number of tables: 0

Supplemental figures: 4

Key words: interlimb coordination, gait analysis, walking, gait asymmetry

ABSTRACT

Human walking involves tightly coordinated movements of the right and left legs. We recently

developed and tested a 'dynamic treadmill walking' paradigm that changes the treadmill speed

within a single step to provide asymmetric training for persons with gait dysfunction. We

previously demonstrated that this approach could induce changes in human gait symmetry;

however, if this approach is to be used in rehabilitation, we also need to understand how

movements of the legs are coordinated to produce these asymmetric gait changes. The goal of

this study was to examine the temporal (phase shift) and spatial (center of oscillation difference)

aspects of interlimb coordination during dynamic treadmill walking in ten young adults without

gait impairment. We found that dynamic treadmill walking drove significant changes in phase

shift and center of oscillation difference that were dependent on the timing of the treadmill speed

change within the gait cycle. For example, slowing the treadmill during the stance phase

extended the double limb support period, and these changes were strongly correlated with a

phase shift between the two legs. Accelerating the treadmill late in stance led to extensions in the

trailing limb angle that were strongly correlated with changes in the center of oscillation

difference. Overall, dynamic treadmill walking can be configured to drive changes in many

spatiotemporal, kinematic, and interlimb coordination parameters, creating a variety of options

for restoring gait symmetry and targeting aspects of spatial and temporal interlimb coordination

in clinical populations with heterogenous patterns of gait asymmetry.

NEW & NOTEWORTHY

Dynamic treadmill walking has been previously shown to drive asymmetric gait changes, but

little is known about its effect on interlimb coordination. Here, we find that this approach

induces significant changes in interlimb coordination (relative to normal walking) in both spatial

and temporal domains. Altogether, dynamic treadmill walking offers a customizable gait

rehabilitation strategy for asymmetric gait training that can induce changes in interlimb

coordination using only a single-belt treadmill.

INTRODUCTION

Many clinical populations – e.g., persons with stroke[1], Parkinson's disease[2], or cerebral palsy[3] – walk with gait patterns characterized by disrupted interlimb coordination (i.e., poorly coordinated movements of the left and right legs). Unimpaired gait is characterized by oscillating, anti-phase movements between the two legs: one leg generally moves forward as the other extends backward and vice versa in a bilaterally coordinated manner[4,5]. Damage to the nervous system can affect these well-coordinated movements between the legs such that the oscillations deviate from their typical anti-phase relationship and become asymmetric[4,6]. Restoration of anti-phase interlimb coordination then becomes important because deviations from symmetric, coordinated gait can lead to increased fall risk[7,8] and elevated energy expenditure[9–11] that can decrease mobility and community engagement[12].

Interlimb coordination can be quantified in several ways. Locomotion literature often uses a combination of temporal (e.g., limb phasing) and spatial (e.g., difference in the axes around which each limb oscillates, or "center of oscillation difference") metrics to measure and understand how the legs are controlled in synchrony[5,13–15]. In the temporal domain, "phasing" can be used to capture the degree to which the leg oscillations are in anti-phase[16]. In the spatial domain, "center of oscillation" describes the axis around which each leg rotates in the sagittal plane as a person walks (i.e., a positive or negative center of oscillation would indicate that the leg rotates around a flexed or extended axis, respectively)[17]. The center of oscillation difference then captures the difference in the axes of rotation of the right and left legs. A difference value near zero is expected for symmetric walking, indicating that both legs rotate about a similar axis. However, both phasing and center of oscillation are often altered in persons with asymmetric gait patterns or when walking under asymmetrically perturbed conditions[16].

The ability to use gait perturbations to drive changes in interlimb coordination offers an

opportunity to leverage these perturbations to improve interlimb coordination in persons with

gait impairment. Our laboratory recently developed a 'dynamic treadmill walking' approach that

can drive changes in gait symmetry by changing the speed of a conventional treadmill within a

single stride[18]. Our previous study in young adults without gait impairment revealed that the

treadmill speed changes could be delivered at different timepoints within the gait cycle (via

pacing of participant heel-strikes using a metronome) to induce customizable asymmetries in

several clinically relevant gait parameters[18]. For example, slowing the treadmill immediately

following right heel-strike induced asymmetries in step time, peak braking, and propulsion

impulse whereas slowing the treadmill late in stance induced asymmetries in step time, leading

limb angle, and peak braking force[18].

It is likely that different patterns of changes in interlimb coordination underlie these

various spatiotemporal, kinematic, and kinetic gait asymmetries induced by dynamic treadmill

walking. However, given that only one limb is paced by a metronome and the other is

unconstrained during this task, it is not yet clear how the coordination patterns between the limbs

change with treadmill belt speed changes at different timings throughout the gait cycle. We

anticipate that a better understanding of how dynamic treadmill walking affects interlimb

coordination could be useful for providing customizable rehabilitation approaches aimed at

restoring proper interlimb coordination in persons with heterogeneous asymmetric gait deficits.

The goal of this study was to investigate interlimb coordination across a range of

dynamic treadmill walking conditions in young adults without gait impairment. We hypothesized

that interlimb coordination would be changed during dynamic treadmill walking (relative to

normal walking) as demonstrated by non-zero center of oscillation differences and deviations

from anti-phase coupling. More specifically, we hypothesized that these changes in interlimb

coordination would differ depending on the timing of the speed change within a gait cycle. In the

trials where the leg paced by the metronome (the right leg) stepped onto a fast or accelerating

treadmill, we hypothesized that leading limb angles on the right leg would increase and stance

times would decrease, resulting in the right leg showing a center of oscillation about a more

flexed axis when compared to that of the left leg (and the opposite to be true when the right leg

stepped onto a slow or decelerating treadmill). Similarly, we expected that the legs would be

more in-phase when participants stepped onto a fast or accelerating treadmill with their paced

(right) leg with the right leg leading the left in time (and the right leg to lag the left in time when

stepping onto a slow or decelerating treadmill).

METHODS

Ten young adults without gait impairment participated in this study (2 M, 8 F, age:

25.1±3.0 years). These participants composed one group from a larger, unpublished experiment

(16 M, 24 F, age: 25.6±3.5 years, randomly assigned to four groups). This group walked under

relatively slow dynamic treadmill conditions (as compared to our previous work[18]) for the

purpose of understanding potential translational implications to neurologic populations. All

participants reported no current neurological, musculoskeletal, or cardiovascular conditions and

provided written, informed consent in accordance with the Declaration of Helsinki. The Johns

Hopkins School of Medicine Institutional Review Board approved all study procedures, and

participants received monetary compensation for their participation (IRB00394009).

Data collection

Participants walked on an instrumented split-belt treadmill (Motek Medical BV, Houten,

The Netherlands), but we emphasize that the treadmill belt speeds were tied for all trials (i.e.,

participants did not walk under split-belt treadmill conditions with the belts moving at different

speeds simultaneously; the split-belt treadmill allowed us to collect ground reaction force data

from each individual leg). We collected kinematic data using an eight-camera motion capture

system (Vicon Motion Systems, Centennial, CO; 100 Hz; RRID: SCR_015001) with 30 passive

retroreflective markers on the seventh cervical vertebrae, jugular notch, tenth thoracic vertebrae,

xiphoid process, bilaterally over the iliac crest, anterior superior iliac spine, posterior superior

iliac spine, trochanter, thigh, lateral femoral epicondyle, medial femoral epicondyle, shank,

lateral malleolus, medial malleolus, calcaneus, second metatarsal head, and fifth metatarsal head.

We collected triaxial ground reaction forces using force plates embedded independently under

each treadmill belt (1000 Hz). We executed dynamic treadmill walking on our treadmill with a

custom D-Flow script (Motek Medical BV, Houten, The Netherlands). The dynamic treadmill

controller changed between a slow (0.5 m/s) and fast (1.0 m/s) speed within a single stride at an

acceleration rate of 6.0 m/s^2. For each dynamic treadmill trial, 50% of the gait cycle was spent at

the fast speed and 50% at the slow speed.

Experimental protocol

The experimental protocol is outlined in Figure 1. All participants completed three two-

minute baseline trials at the slow (0.5 m/s), intermediate (0.75 m/s), and fast speeds (1.0 m/s).

Four five-minute dynamic treadmill walking trials (Slow, Fast, Accelerate, and Decelerate)

followed the baseline trials in a randomized order. Our experimental design included a similar

set of conditions that we used in our previous work[18], but here we tested at slower treadmill

speeds (previous study: slow=0.75 m/s, intermediate=1.125 m/s, fast=1.5 m/s)[18]. We instructed

participants to synchronize each right heel-strike with a metronome tone that was aligned with

the treadmill speed changes. We determined the timing of the metronome tone by measuring

each participant's average stride time across the intermediate baseline trial, as calculated by

dividing the trial duration (120 seconds) by the total number of strides completed in the trial.

For each dynamic treadmill trial, the treadmill speed changed at different points within

the gait cycle (Figure 1). The metronome sounded to pace the participant's right heel-strike to

guide the desired timing pattern set by each dynamic treadmill trial:

• Slow: the right heel-strike was paced to occur with the treadmill at the slow speed (0.5 m/s).

This resulted in the treadmill moving slow over the first and last ~25% of the gait cycle and

fast over the middle ~50% of the gait cycle.

• Fast: the right heel-strike was paced to occur with the treadmill at the fast speed (1.0 m/s).

This resulted in the inverse of Slow: the treadmill moved fast over the first and last ~25% of

the gait cycle and slow over the middle ~50% of the gait cycle.

• Accelerate: the right heel-strike was paced to occur when the treadmill was moving at

approximately the intermediate speed (0.75 m/s) while accelerating to the fast speed. This

resulted in the treadmill moving at the fast speed during the first ~50% of the gait cycle and

then decelerating to the slow speed for the last ~50% of the gait cycle

• Decelerate: the right heel-strike was paced to occur when the treadmill moved at

approximately the intermediate speed (0.75 m/s) while decelerating to the slow speed. This

resulted in the inverse of Accelerate: the treadmill moved slow over the first ~50% of the gait

cycle and fast over the last ~50% of the gait cycle.

Data analysis

We used custom MATLAB software for data processing (R2023b, The Mathworks,

Natick, MA; RRID: SCR_001622). We filtered the marker and ground reaction force data using

fourth-order low-pass Butterworth filters with cutoff frequencies of 6 Hz and 28 Hz,

respectively. We determined heel-strikes as the local maxima and minima where the change in

limb angle (calculated as the sagittal vectors between the iliac crest and second metatarsal head

markers) shifted from positive to negative and toe-offs as the opposite[18]. We calculated step

length, step time, and leading and trailing limb angles as in our previous work[18] as well as stance

time (time between heel-strike and toe-off of the ipsilateral limb), swing time (time between toe-

off and heel-strike of the ipsilateral limb), and double support time (time between heel-strike of

the contralateral limb and toe-off of the ipsilateral limb).

We also recorded the treadmill belt speeds within each trial to assess how well

participants synchronized their right heel-strikes with the metronome tone to assess adherence to

the desired heel-strike/speed change pairings of the different dynamic treadmill trials. We

calculated cross-correlation coefficients (at time lag zero) between the prescribed and actual

treadmill speed time series for each gait cycle within each participant to understand how well

participants adhered to the prescribed conditions. We also calculated a general metric of how

accurately participants stepped to the metronome: for each gait cycle, we considered a

participant's step successful if their right heel-strike occurred within ±200 ms of the metronome

tone initiation.

We calculated phase shift and center of oscillation difference as temporal and spatial

measures of interlimb coordination, respectively. As a first step to calculating phase shift, we

calculated limb phasing as the stride-by-stride maximum cross-correlation coefficient between

right and left limb angle trajectories. Next, we subtracted limb phasing during dynamic treadmill

walking from limb phasing during intermediate baseline walking to calculate phase shift (i.e., a

phase shift of zero would indicate that a participant had identical phasing values to the baseline

trial). We calculated center of oscillation as the limb angle at midstance (or at the timepoint

midway between heel-strike and toe-off) for each leg. We then calculated the center of

oscillation difference as the difference between centers of oscillation measured for the right and

left centers of oscillation.

Statistical analysis

We used R version 4.3.3 (R Core Team, 2023: RRID: SCR_001905) and RStudio[19]

version 2023.12.1 (RRID: SCR_000432) for data analysis and visualization, including the readxl

(RRID: SCR_018083), dplyr (RRID: SCR_016708), rstatix (RRID: SCR_021240), and psych

(RRID: SCR_021744) packages[19–24]. Given that we paced all participants to their stride time

during the intermediate baseline trial, we first used one-way repeated measures ANOVA to

confirm that stride time was consistent across the four dynamic treadmill trials. Next, we used

one-way repeated measures ANOVA to assess whether order effects were present for our

primary interlimb coordination outcome variables (phase shift and center of oscillation

difference) across the different trial orders. To test how well participants matched the prescribed

belt speeds across the four dynamic treadmill trials, we then performed one-way repeated

measures ANOVAs to compare cross-correlation coefficients (averaged across all gait cycles for

each participant) of the prescribed versus actual treadmill speed time series across the four

dynamic treadmill trials.

To test our primary study hypotheses, we performed a 4 (Trial) x 2 (Leg) repeated

measures ANOVA to compare Trial and Leg differences for spatiotemporal and kinematic gait

parameters (i.e., stance time, swing time, double support time, leading limb angle, and trailing

limb angle). We then investigated interlimb coordination across the dynamic treadmill trials

using one-way repeated measures ANOVA to compare phase shift among four dynamic

treadmill trials and performed a similar analysis on center of oscillation difference. We analyzed

individual leg centers of oscillation using 4 (Trial) x 2 (Leg) repeated measures ANOVA to

understand whether any center of oscillation differences were driven by changes in center of

oscillation in the right versus left leg. All repeated measures ANOVAs were tested for sphericity

using Mauchly's Test, and Greenhouse-Geisser corrected values are provided when the

sphericity test failed $(p<0.05)$[25]. Finally, we performed Pearson's correlations to assess the

relationships between the interlimb coordination measures and kinematic and spatiotemporal gait

parameters. For all intralimb and interlimb measures, we performed statistical analyses on the

mean of the last 30 strides for each trial. We set $\alpha<0.05$ for all analyses and used Bonferroni

post-hoc corrections for pairwise comparisons where appropriate.

RESULTS

Our preliminary checks verified that 1) participants walked with similar stride times

across all four dynamic treadmill walking trials and 2) there was no effect of the dynamic

treadmill trial order on the outcomes. We verified that participants indeed walked with similar

stride times across all four dynamic treadmill walking conditions using one-way repeated

measures ANOVA, where there was no significant main effect of Trial $(F(3,67)=2.73, p=0.05)$.

Similarly, we did not observe a significant main effect of Block (or order) on phase shift

$(F(3,27)=0.92, p=0.45)$ or center of oscillation difference $(F(3,27)=0.41, p=0.75)$.

Participants stepped in synchrony with the metronome during dynamic treadmill walking

Participants synchronized their right heel-strikes to a metronome tone with strong cross-

correlation coefficients (at time lag zero) between the prescribed and actual treadmill speeds

across each of the trials (mean±standard error of the mean): Slow: $r=0.97\pm0.01$, Fast:

$r=0.98\pm0.01$, Accelerate: $r=0.98\pm0.01$, Decelerate: $r=0.97\pm0.01$(treadmill speed profiles shown

in Figure 2A). One-way repeated measures ANOVA revealed no significant main effect of Trial

(F(3,26)=1.59, p=0.21) on the cross-correlation coefficients, indicating that there were no

significant differences in how successfully participants stepped to the metronome tone across the

four dynamic treadmill trials. Moreover, participants were generally successful with

synchronizing their right heel-strikes to the metronome tone: group mean success rates ranged

from 89% to 94% across the four dynamic treadmill trials (i.e., participants stepped within ±200

ms of the metronome tone on 89% to 94% of strides).

Dynamic treadmill walking drives asymmetric changes in spatiotemporal and kinematic gait

parameters

Next, we aimed to understand the spatiotemporal and kinematic factors that contributed

to the different patterns of limb angle timing and amplitude observed across the four dynamic

treadmill trials (Figure 2B). For comparison to our previous work, we calculated step length in

addition to the step-by-step values of spatiotemporal and kinematic parameters below

(Supplemental Figures S1-S4).

<u>Stance time</u>

Stance time generally increased when the treadmill was slow during a longer portion of

the stance phase (e.g., right leg stance during the Fast and Decelerate conditions). A 4 (Trial) x 2

(Leg) repeated measures ANOVA revealed a significant main effect of Trial (F(3,27)=3.44,

p=0.03) on stance time and a significant Trial*Leg interaction (F(3,27)=14.3, p<0.01), but no

significant main effect of Leg (F(1,9)=2.09, p=0.18; Figure 2C). Post-hoc analyses showed

significant within-trial Leg differences for stance time in Slow (left longer than right, p<0.01),

Fast (right longer than left, p<0.01), Accelerate (left longer than right, p=0.04), and Decelerate

(right longer than left, p<0.01). We also observed that left stance time in Fast was significantly

shorter than left stance time in Accelerate (p=0.03) and Slow (p<0.01). Right stance time in Slow

was significantly shorter than in Decelerate (p=0.01) and Fast (p<0.01). Left stance time was

significantly longer in Slow than in Decelerate (p=0.046). Right stance time in Accelerate was

significantly shorter than in Decelerate (p=0.01) and Fast (p<0.01).

<u>Swing time</u>

238   Because swing time has a strong, inverse correlation with stance time across all trials

(r^2=-0.99, p<0.01), we expected similar findings as stance time but with opposite directionality.

Indeed, swing times generally decreased when the treadmill was slow during a large portion of

the stance phase and fast during the swing phase. A 4 (Trial) x 2 (Leg) repeated-measures

ANOVA revealed no significant main effect of Trial (F(3,27)=2.85, p=0.06) or Leg

(F(1,9)=2.07, p=0.18) on swing time but did reveal a significant Trial*Leg interaction

(F(3,27)=14.55, p<0.01, Figure 2C). Post-hoc analyses revealed significant within-trial Leg

differences in swing time for Slow (right longer than left, p<0.01), Fast (left longer than right,

p<0.01), Accelerate (right longer than left, p=0.04), and Decelerate (left longer than right,

p<0.01). Furthermore, post-hoc analyses revealed that within-leg swing times were significantly

different between Fast and Accelerate, with Fast producing a longer swing time on the left leg

than Accelerate (p=0.02) and Accelerate driving a longer swing time on the right leg (p=0.01).

Slow and Fast exhibited significantly different swing times on the left and right legs, with Fast

producing a longer left swing time than Slow (p=0.01) and Slow producing a longer right swing

time than Fast (p<0.01). Decelerate produced a significantly shorter right swing time than

Accelerate (p=0.02) and Slow (p<0.01).

<u>Double support time</u>

255   Double support times were generally longer when the treadmill was slow during double

support (e.g., double support period from right heel-strike to left toe-off – or left double support

time using our naming convention – during Slow or Decelerate). A 4 (Trial) x 2 (Leg) repeated

measures ANOVA revealed a significant main effect of Trial $(F(3,27)=3.31, p=0.04)$ and Leg

$(F(1,9)=8.62, p=0.02)$ on double support time as well as a significant Trial*Leg interaction

$(F(3,27)=13.17, p<0.01)$. Post-hoc analyses revealed significant Leg differences within Slow

(left longer than right, $p<0.01$), Fast (right longer than left, $p=0.047$), and Decelerate (left longer

than right, $p<0.01$). Left double support time in Fast was significantly shorter than left double

support time in Decelerate $(p<0.01)$ and Slow $(p<0.01)$. Left double support time in Slow was

significantly longer than that in Accelerate $(p<0.01)$. Slow produced significantly shorter right

double support times than Accelerate $(p<0.01)$ and Fast $(p=0.01)$. Right double support time was

significantly longer in Fast than in Decelerate $(p<0.01)$.

<u>Leading limb angle</u>

Leading limb angles were largely similar across trials. A 4 (Trial) x 2 (Leg) repeated measures

ANOVA revealed no significant main effect of Trial $(F(3,27) =0.81, p=0.50)$ or Leg

$(F(1,9)=1.61, p=0.24)$ on leading limb angle. We also did not observe a significant Trial*Leg

interaction $(F(3,27)=2.27, p=0.10)$.

<u>Trailing limb angle</u>

Trailing limb angles were generally larger (i.e., more negative) when the treadmill moved

fast during late stance (e.g., right leg in Slow). A 4 (Trial) x 2 (Leg) repeated measures ANOVA

revealed no significant main effects of Trial $(F(3,27)=1.56, p=0.22)$ or Leg $(F(1,9)=0.19, p=0.67)$

on trailing limb angle, but we did observe a significant Trial*Leg interaction $(F(3,27)=12.66,$

$p<0.01)$. Post-hoc analyses revealed that Slow, Fast, and Decelerate showed significant

differences between the left and right legs, with right trailing limb angle being larger (i.e., more

negative) than left trailing limb angle in Slow $(p<0.01)$, the left trailing limb angle being larger

than the right trailing limb angle in Fast (p<0.01), and right trailing limb angle being larger than

left trailing limb angle in Decelerate (p=0.049). Left trailing limb angle in Fast was significantly

larger than in Decelerate (p=0.03) and Slow (p<0.01). Right trailing limb angle in Fast was

significantly smaller than in Decelerate (p<0.01) and Slow (p<0.01).

Dynamic treadmill walking alters spatial and temporal components of interlimb coordination

Temporal interlimb coordination: phase shift

After understanding how the different dynamic treadmill trials affected spatiotemporal

and kinematic gait parameters, we next studied how the different trials affected interlimb

coordination. A phase shift value of zero indicates that the limbs are in anti-phase (similar to

baseline walking). Positive phase shift indicates that the right leg oscillations are advanced in

time relative to those of the left and vice versa. In Figure 3A, we now show the limb angles with

0% of the gait cycle representing right heel-strike to visualize the coordination between the two

limbs.

Dynamic treadmill walking conditions showed the most pronounced changes in phase

shift when the treadmill speed changes occurred largely during double support periods (e.g.,

Slow and Fast, see Figures 3B-C). One-way repeated measures ANOVA revealed a significant

main effect of Trial ($F(2.2, 19.76) = 9.7$, $p < 0.01$) on phase shift (Figure 3C). Post-hoc analyses

revealed that phase shift in Fast was significantly more positive (i.e., right leg oscillates

advanced in time relative to the left) than in Slow (p<0.01) and Decelerate (p=0.02). Phase shift

in Slow was significantly more negative (i.e., left leg oscillates advanced in time relative to the

right) than in Accelerate (p<0.01).

Spatial interlimb coordination: center of oscillation and center of oscillation difference

A positive center of oscillation value indicates that the right leg is rotating about a flexed

axis, whereas a negative value indicates rotation about an extended axis. For center of oscillation

difference, a positive value indicates that the right leg rotates about a more flexed axis than the

left and vice versa.

Similar to our findings in the temporal domain, dynamic treadmill walking conditions

where the treadmill speed changes occurred largely during double support periods showed the

most pronounced changes in center of oscillation (Figure 4). A 4 (Trial) x 2 (Leg) repeated

measures ANOVA revealed no significant main effects of Trial ($F(3,27)=0.85$, $p=0.45$) or Leg

($F(1,9)=0.96$, $p=0.35$) on center of oscillation. However, the ANOVA did reveal a significant

Trial*Leg interaction ($F(3,27)=7.24$, $p<0.01$; Figure 4C). Post-hoc analyses revealed that center

of oscillation in Fast was significantly more positive on the right leg than the left ($p=0.02$). Slow

produced a significantly more positive center of oscillation on the left leg than the right ($p<0.01$).

Similarly, Decelerate drove a significantly more positive center of oscillation on the left leg than

the right ($p=0.03$). In Fast, the left leg oscillated about a more extended axis when compared to

the left leg in Decelerate ($p<0.01$). In Slow, the left leg oscillated about a more flexed axis when

compared to the left leg in Accelerate ($p=0.049$). In Fast, the left leg oscillated about a more

extended axis than the left leg in Slow ($p<0.01$). Conversely, the right leg in Fast oscillated about

a significantly more flexed axis when compared to the right leg in Slow ($p=0.03$).

For center of oscillation difference (Figure 4D), a one-way repeated measures ANOVA

revealed a significant main effect of Trial ($F(3,27)=7.24$, $p<0.01$). Post-hoc analyses revealed

that Fast displayed a significantly larger center of oscillation difference than Decelerate ($p<0.01$).

Additionally, Fast displayed a significantly larger center of oscillation difference than Slow

(p<0.01).

Correlational relationships between interlimb gait parameters and spatiotemporal and kinematic

gait parameters

Phase shift

We observed several statistically significant correlations between phase shift and

spatiotemporal and kinematic gait parameters (Figure 5A). We observed a significant positive

correlation between phase shift and stance time asymmetry (r=0.41, p<0.01) and a significant

negative correlation between phase shift and swing time asymmetry (r=-0.42, p<0.01). We also

observed a significant positive correlation between phase shift and double support time (r=0.93,

p<0.01). Furthermore, we observed significant negative correlations between phase shift and

leading and trailing limb angles (r=-0.33, p=0.03; r=-0.47, p<0.01).

Center of oscillation difference

Center of oscillation difference was also significantly associated with spatiotemporal and

kinematic gait parameters (Figure 5B). Similar to phasing, we observed significant positive

correlations between center of oscillation difference and stance time asymmetry (r=0.36, p=0.02)

and double support time asymmetry (r=0.47, p<0.01) as well as a significant negative correlation

between center of oscillation difference and swing time asymmetry (r=-0.37, p=0.02). We

observed a significant positive correlation between center of oscillation difference and leading

limb angle (r=0.47, p<0.01) and a significant negative correlation between center of oscillation

difference and trailing limb angle (r=-0.88, p<0.01).

DISCUSSION

In this study, we showed that dynamic treadmill walking drove significant interlimb

coordination changes that were dependent on the timing of the speed change within the gait cycle

(i.e., trial type) and were related to a variety of changes in spatiotemporal and kinematic gait

parameters. Overall, our hypotheses were supported in Slow and Fast, which were the trials

showing the most strongly perturbed interlimb coordination in both the spatial and temporal

domains. During Slow and Fast, the legs were more in-phase and showed larger asymmetries in

the axes of rotation for each leg when compared to the other dynamic treadmill trials and to

normal walking. Here, we will discuss potential implications of the different dynamic treadmill

walking trials in driving gait changes in populations with gait asymmetry, and future

implications for the potential clinical translation of dynamic treadmill walking.

Spatiotemporal and kinematic changes depended on the timing of treadmill speed changes

Dynamic treadmill walking drove significant changes in spatiotemporal and kinematic

gait parameters. Despite the changing nature of the treadmill speed during dynamic treadmill

walking, the spatiotemporal and kinematic changes were in accordance with those that may be

expected based on the speeds of the treadmill isolated at specific points in the gait cycle. For

example, the treadmill moved at the slow speed for a majority of the stance phase of the right leg

during Fast; accordingly, the right stance time was significantly longer than the left stance time

during this trial. Similarly, trailing limb angle was influenced by treadmill speed during late

stance. A fast treadmill speed during late stance drove a larger (more negative) trailing limb

angle. Surprisingly, leading limb angle did not follow our expectations, as we expected the leg

that stepped onto a fast treadmill to show a larger leading limb angle than the other leg (i.e., we

expected larger leading limb angles in the right leg during Fast as compared to Slow[18]).

However, there were no significant differences between the legs within any or between the trials.

Our leading limb angle findings do not agree with our previous work[18]; we hypothesize that this

may be due to the slower belt speeds being less likely to alter foot placement than the speeds

used previously[18].

Changes in interlimb coordination were related to changes in spatiotemporal and kinematic gait

parameters

Dynamic treadmill walking induced significant temporal and spatial interlimb

coordination changes as quantified by phase shift and center of oscillation difference,

respectively. When the treadmill was fast at heel-strike but then decelerated during double

support (i.e., Fast), the leading leg rotated about a more flexed axis and progressed through an

extended stance phase. This also drove the leading leg to lag in phase relative to the contralateral

leg. As expected, opposite interlimb coordination changes were driven between dynamic

treadmill walking trials with opposite behavior (e.g., Slow and Fast). Overall, the legs were more

in-phase and participants walked about a more forward axis of oscillation than in normal

walking.

We also observed that these changes in interlimb coordination were strongly associated

with a variety of spatiotemporal and kinematic gait parameters. For instance, we found that

double support time asymmetry was strongly correlated with phase shift, indicating that double

support periods of the gait cycle were sensitive to treadmill speed change and related to similar

shifts in phasing. In the spatial domain, we observed a strong correlation between center of

oscillation difference and trailing limb angle asymmetry. Unsurprisingly, when the treadmill

accelerated or moved at the faster speed during late stance, the treadmill extended the leg in

stance further behind the body than the other, ultimately contributing to the right leg oscillating

about a more extended axis of rotation.

Comparison between dynamic treadmill walking and split-belt walking

Dynamic treadmill walking could be intuited as a split-belt walking-like environment (in

split-belt walking, there are separate treadmill belts under each foot that drive each leg to walk at

a different speed simultaneously[5]) with a conventional, single-belt treadmill. However, these two

approaches to asymmetric gait training have clear differences. During split-belt walking, the belt

speeds experienced by the left and right legs are independent; in dynamic treadmill walking,

each leg experiences the same belt speed during double support periods. Therefore, in dynamic

treadmill walking, the bilateral mismatch in the belt speed predictions that occurs during split-

belt treadmill walking is absent.

The interlimb coordination demands of split-belt treadmill walking are also significantly

different than those observed here in dynamic treadmill walking. During split-belt walking, the

slow leg lags the fast leg in timing, creating a more in-phase gait pattern than normal walking[16].

Additionally, the slow leg oscillates about a more forward axis than the fast leg during split-belt

walking[16]. Here, Slow and Fast dynamic treadmill walking conditions drove similar patterns of

temporal and spatial changes, but Accelerate and Decelerate produced changes in only one, or

neither, domain of interlimb coordination due to the timing of the speed changes. The magnitude

of interlimb coordination change created in Slow and Fast was similar to that of gradually

introduced split-belt walking but about half the magnitude of abruptly introduced split-belt

walking[16]. We hypothesize that this lower magnitude of interlimb coordination change in Slow

and Fast – and smaller deviations in the remaining trials – play a role in dynamic treadmill

walking not following the characteristics of gait adaptation that are commonly observed in split-

belt walking. For example, our previous work showed that there were minimal aftereffects after

dynamic treadmill walking[18], certainly lower than the robust aftereffects observed in split-belt

walking[18]. We suspect that these phenomena are driven by the lack of bilateral mismatch in belt

speed predictions that occur in split-belt walking. Additionally, the explicit task of pacing the

right limb to a metronome provides a different kind of stimulus for gait pattern change than the

implicit task of responding to asymmetric belt speed changes alone[26].

Clinical translation of dynamic treadmill walking

The ability to selectively target asymmetric gait using dynamic treadmill walking, which

can be executed on a conventional treadmill, offers new opportunities for use in traditional

rehabilitation settings. To understand how people with a neurologic impairment or injury will

respond to dynamic treadmill walking, it is important to first understand the interlimb

coordination demands of this pattern. We have shown that this pattern can be leveraged to drive

significant interlimb coordination changes in Slow or Fast but does not require the same

magnitude of changes as other methods (i.e., split-belt walking[27]). For people with gait

impairment, the field is still gaining insights into the necessary ingredients for creating a long-

term change in gait symmetry[28] and perturbing interlimb coordination is one of the potential

candidates. Additionally, we cannot assume that people with asymmetric gait patterns, and

especially those with neurologic conditions, respond to dynamic treadmill walking in the same

way as unimpaired young adults. Therefore, ongoing work is assessing the feasibility of using

dynamic treadmill walking to normalize asymmetry patterns after stroke, as well as the potential

impact of repeated exposure on changing overground gait asymmetry.

Limitations

We note several limitations to our study. Young adults with intact neurologic systems

tend to walk at faster speeds than studied here (approximately 1.5 m/s[29]); however, in our study,

participants walked at 0.5 m/s and 1.0 m/s on the dynamic treadmill. These slower speeds are

advantageous from a clinical perspective because clinical populations tend to walk more slowly,

though there is potential that these slower speeds may have simultaneously disrupted the normal

walking patterns of our participants[30]. Another limitation of the current study is that we only

used two measures of interlimb coordination (center of oscillation difference and phase shift).

There are other measures that we could have used to gain a deeper understanding of interlimb

coordination changes driven by dynamic treadmill walking (e.g., limb orientation at weight

shift[5]). Similarly, we operationally defined interlimb coordination as the coordination between

the left and right legs. We therefore did not assess movement of the upper limbs or how they

interact with the legs in these analyses. Finally, while our long-term goals of this line of research

are focused on clinical applications, we did not study clinical populations directly.

CONCLUSION

In this study, we investigated the interlimb coordination changes that dynamic treadmill

walking may require. We found that dynamic treadmill walking drove significant interlimb

coordination changes, both in the spatial and temporal domains. These changes were dependent

on the timing of the treadmill speed change. These findings show that dynamic treadmill walking

is a tunable tool that can drive changes in spatiotemporal, kinematic, and interlimb coordination

parameters. As a potential clinical tool, this finding is encouraging because it offers additional

flexibility to remediate gait asymmetry while choosing whether to challenge interlimb

coordination and the direction in which it is challenged. Given that we tested interlimb

coordination changes driven by this tool in slower-than-average walking for control participants,

there is potential for implementation into clinical populations that present with gait asymmetry.

DATA AVAILABILITY

The data and analysis code that supports the findings of this study are archived on

Zenodo: https://doi.org/10.5281/zenodo.15705993.

ACKNOWLEDGEMENTS

The authors would like to thank the members of the Kennedy Krieger Institute Center for

Movement Studies for their valuable contributions and thoughtful discussion to the findings

presented here. This article was posted to BioRxiv as a preprint, available at

https://doi.org/10.1101/2024.11.27.625740.

FUNDING SOURCES

This work was supported by an American Heart Association Career Development Award

#935556 (RTR). Brooke Hall was supported by a Research Supplement to Promote Diversity in

Health-Related Research (#3R21HD110686-02S1, PI RTR). Caitlin Banks is supported by an

American Heart Association Postdoctoral Fellowship

(https://doi.org/10.58275/AHA.25POST1366391.pc.gr.227425), and previously a training grant

from the Eunice Kennedy Shriver National Institute of Child Health and Human Development

(#5T32HD007414-29, PI Bastian). BLH and CLB implemented data science and visualization

training in this work from the ReproRehab program (NIH NICHD/NCMRR R25HD105583, PI

Liew).

DISCLOSURES

The authors have declared no conflicts of interest or disclosures.

AUTHOR CONTRIBUTIONS

Conceived and designed research: BH, RTR

Performed experiments: BH

Analyzed data: CLB, BH

Interpreted results of experiments: CLB, BH, RTR

Prepared figures: CLB, BH, RTR

Drafted manuscript: BH

Edited and revised manuscript: CLB, BH, RTR

Approved final version of the manuscript: CLB, BH, RTR

REFERENCES

1. 1. Olney SJ, Richards C. Hemiparetic gait following stroke. Part I: Characteristics. *Gait & Posture*. 1996;4(2):136-148. doi:10.1016/0966-6362(96)01063-6

2. Knutsson E. An analysis of Parkinsonian gait. *Brain*. 1972;95(3):475-486. doi:10.1093/brain/95.3.475

3. White R, Agouris I, Selbie RD, Kirkpatrick M. The variability of force platform data in normal and cerebral palsy gait. *Clinical Biomechanics*. 1999;14(3):185-192. doi:10.1016/S0268-0033(99)80003-5

4. Dec. JB, Saunders M, Inman VT, Eberhart HD. The Major Determinants in Normal and Pathological Gait. *J Bone Joint Surg Am*. 1953;35:543-558.

5. Reisman DS, Block HJ, Bastian AJ. Interlimb Coordination During Locomotion: What Can be Adapted and Stored? *Journal of Neurophysiology*. 2005;94(4):2403-2415. doi:10.1152/jn.00089.2005

6. Sousa ASP, Silva A, Santos R, Sousa F, Tavares JMRS. Interlimb Coordination During the Stance Phase of Gait in Subjects With Stroke. *Archives of Physical Medicine and Rehabilitation*. 2013;94(12):2515-2522. doi:10.1016/j.apmr.2013.06.032

7. Bloem BR, Hausdorff JM, Visser JE, Giladi N. Falls and freezing of gait in Parkinson's disease: A review of two interconnected, episodic phenomena. *Movement Disorders*. 2004;19(8):871-884. doi:10.1002/mds.20115

8. Wei TS, Liu PT, Chang LW, Liu SY. Gait asymmetry, ankle spasticity, and depression as independent predictors of falls in ambulatory stroke patients. 2017;12(5). doi:10.1371/journal.pone.0177136

9. Finley J. Associations between foot placement asymmetries and metabolic cost of transport in hemiparetic gait. 2016;31(2). doi:https://doi.org/10.1177/1545968316675428

10. Awad L, Palmer J, Reisman D, Binder-Macleod S, Pohlig R. Walking speed and step length asymmetry modify the energy cost of walking after stroke. 2015;29(5):416-423. doi:10.1177/1545968314552528

11. Gill P. Causal modelling demonstrates metabolic power is largely affected by gait kinematics and motor control in children with cerebral palsy. Published online May 24, 2023. doi:https://doi.org/10.1371/journal.prone.0285667

12. Lord S, McPherson K, McNaughton H, Rochester L, Weatherall M. Community ambulation after stroke: how important and obtainable is it and what measures appear predictive? 2004;85(2):234-239. doi:https://doi.org/10.1016/j.apmr.2003.05.002

13. Reisman DS, Wityk R, Silver K, Bastian AJ. Locomotor adaptation on a split-belt treadmill
can improve walking symmetry post-stroke. *Brain*. 2007;130(7):1861-1872.
doi:10.1093/brain/awm035

14. Roemmich RT, Bastian AJ. Two ways to save a newly learned motor pattern. *J
Neurophysiol*. 2015;113(10):3519-3530. doi:10.1152/jn.00965.2014

15. Malone LA, Bastian AJ, Torres-Oviedo G. How does the motor system correct for errors in
time and space during locomotor adaptation? *Journal of Neurophysiology*. 2012;108(2):672-
683. doi:10.1152/jn.00391.2011

16. Torres-Oviedo G, Bastian AJ. Natural error patterns enable transfer of motor learning to
novel contexts. *J Neurophysiol*. 2012;107(1):346-356. doi:10.1152/jn.00570.2011

17. Malone LA, Bastian AJ. Spatial and Temporal Asymmetries in Gait Predict Split-Belt
Adaptation Behavior in Stroke. *Neurorehabil Neural Repair*. 2014;28(3):230-240.
doi:10.1177/1545968313505912

18. Browne MG, Stenum J, Padmanabhan P, Roemmich RT. Simple within-stride changes in
treadmill speed can drive selective changes in human gait symmetry. *PLOS ONE*.
2023;18(10):e0287568. doi:10.1371/journal.pone.0287568

19. Posit Team. RStudio: Integrated Development for R. Published online 2024.
http://www.posit.co/ Version 2023.12.1

20. Kassambara A. rstatix: Pipe-Friendly Framework for Basic Statistical Tests. Published online
2021. https://CRAN.R-project.org/package=rstatix Version 1.4.3

21. Wickham H, Bryan J. readxl: Read Excel Files. Published online 2019. https://CRAN.R-
project.org/package=readxl Version 1.4.3

22. Wickham H, Francois R, Henry L, Muller K, Vaughan D. dplyr: A grammar of data
manipulation. Published online 2023. https://CRAN.R-project.org/package=dplyr Version
1.1.4

23. R Core Team. R: A language and environment for statistical computing [Internet]. Published
online 2023. Available from: https://www.r-project.org/

24. Revelle W. psych: Procedures for Personality and Psychological Research. Published online
2021. https://CRAN.R-project.org/package=psych Version = 2..4.6.26,

25. Field A. *Discovering Statistics Using IBM SPSS Statistics*. Sage; 2013.

26. Roemmich RT, Long AW, Bastian AJ. Seeing the Errors You Feel Enhances Locomotor
Performance but Not Learning. *Current Biology*. 2016;26(20):2707-2716.
doi:10.1016/j.cub.2016.08.012

27. Reisman DS, Block HJ, Bastian AJ. Interlimb Coordination During Locomotion: What Can
be Adapted and Stored? *Journal of Neurophysiology*. 2005;94(4):2403-2415.
doi:10.1152/jn.00089.2005

28. Reisman DS, Bastian AJ, Morton SM. Neurophysiologic and Rehabilitation Insights From
the Split-Belt and Other Locomotor Adaptation Paradigms. *Phys Ther*. 2010;90(2):187-195.
doi:10.2522/ptj.20090073

29. Herssens N, Verbecque E, Hallemans A, Vereeck L, Van Rompaey V, Saeys W. Do
spatiotemporal parameters and gait variability differ across the lifespan of healthy adults? A
systematic review. *Gait & Posture*. 2018;64:181-190. doi:10.1016/j.gaitpost.2018.06.012

30. Little VL, McGuirk TE, Patten C. Slower than normal walking speeds involve a pattern shift
in joint and temporal coordination contributions. *Experimental Brain Research*.
2019;237(11):2973-2982. doi:10.1007/s00221-019-05648-2

FIGURE CAPTIONS

Figure 1. Experimental paradigm: All participants completed three baseline trials at slow (0.5

m/s), intermediate (*inter*, 0.75 m/s), and fast (1.0 m/s) treadmill speeds for two minutes each.

Participants then completed the four five-minute, metronome-paced dynamic treadmill walking

trials (*Slow, Fast, Accelerate, Decelerate)* in a randomized order. The shaded, gray box depicts

the target speed profiles of each trial over a gait cycle. Each trial is named according to treadmill

speed at heel-strike (i.e., Slow is at slow treadmill speed at right heel-strike, Fast is at the fast

speed, Accelerate is accelerating between slow and fast, and Decelerate is decelerating between

fast and slow).

Figure 2. Group speed profiles, limb angles, and spatiotemporal gait parameters: **A)** Group

average treadmill speed traces are shown as solid black lines with standard error of the mean

represented with gray fill. Individual participant speed profiles are layered behind the group

traces in light gray. The group average gait event timings (LTO, LHS, RTO) are represented by

vertical dashed lines shown in each profile. *RHS*: right heel-strike; *LTO:* left toe-off; *LHS:* left

heel-strike; *RTO:* right toe-off. The four speed conditions include *Slow, Fast, Accelerate* (*Accel*),

and *Decelerate* (*Decel*). **B)** Limb angle plots with the right (dark) and left (light) gait cycles

anchored to their own heel-strike (i.e., 0% of the gait cycle), respectively. Line conventions and

abbreviations follow those in Figure 2A. **C and D)** Box and whisker plots for spatiotemporal (**C**)

and kinematic (**D**) parameters. The black scatter points show individual average values over the

last 30 strides for each participant. The horizontal lines indicate statistical significance as

denoted with asterisks (*: $p<0.05$, **: $p<0.01$), ***: $p<0.001$). Shading conventions follow

Figure 2B.

Figure 3. Phase shift: **A)** Limb angle plots anchored to the right heel-strike. Lines and shading

conventions follow Figure 2B. Individual participant traces for each leg are displayed by the thin

lines. Group averages of each gait event for each gait cycle are represented by vertical solid

lines. **B)** Time-series plots of left and right limb angles through the duration of each dynamic

treadmill walking condition. Plots depict group mean (thick line) ± standard error of the mean

(lighter colored fill). On the far-right side of each plot, group mean ± standard error over the last

30 strides is depicted by a solid dot with error bar. **C)** Box and whisker plots for phase shift

follow the same conventions as Figures 2C and 2D. The black scatter points show individual

average values over the last 30 strides for each participant. The horizontal lines indicate

statistical significance as denoted with asterisks (*: $p<0.05$, **: $p<0.01$), ***: $p<0.001$).

Figure 4. Center of oscillation: **A)** Time-series plots of the spatial shift between the left (lighter

color) and right (darker color) limb angles through each dynamic treadmill trial. Plots depict

group mean (thick line) ± standard error of the mean (lighter fill). On the far-right side of each

plot, group mean ± standard error over the last 30 strides is depicted by a solid dot with error bar.

B) Time series plots of center of oscillation difference for each walking trial. **C)** Boxplots of

center of oscillation from each trial. Boxes display the group median ± interquartile range of

phase shift values. Individual participant data are represented as black scatter points. **D)** Boxplots

of center of oscillation difference for each trial. Asterisks indicate statistical significance within a

repeated measures analysis of variance: *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$.

Figure 5. Correlational relationships between spatiotemporal and kinematic asymmetry and

intralimb parameters: **A)** Scatter plots of center of oscillation difference versus asymmetry values

for stance time (top left), swing time (top center), double support time (top right), leading limb

angle (lower left), and trailing limb angle (lower center). **B)** Scatter plots of phase shift versus

spatiotemporal (upper) and kinematic (lower) asymmetry values. Pearson correlation coefficients

and statistical significance are listed on each plot.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5