1

Tobacco smoke alters the trajectory of lung adenocarcinoma evolution

via effects on somatic selection and epistasis

Krishna Dasari,[1,2] Jorge A. Alfaro-Murillo,[1] Jeffrey P. Townsend[1,3,4,5]*

[1]*Department of Biostatistics, Yale School of Public Health, New Haven, CT.*

[2]*Medical Scientist Training Program, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA.*

[3]*Yale Cancer Center, Yale University, New Haven, CT.*

[4]*Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT.*

[5]*Program in Computational Biology and Bioinformatics, Yale University, New Haven, CT.*

*Corresponding author. Email: jeffrey.townsend@yale.edu

Abstract

The mutagenic effects of tobacco smoke are well-reported. However, tobacco smoke also likely alters the somatic selective pressures that drive lung cancer evolution, and these altered pressures have never been quantified nor compared to the oncogenic effects of tobacco-induced mutagenesis. This comparison is necessary for prediction of targeted therapy efficacy given exposure history. Here, we estimated background mutation rates, somatic selection, and selective epistasis for 21 driver genes in ever- and never-smoker lung adenocarcinoma (ES- and NS-LUAD).

As expected, mutation rates were gene-specifically elevated in ES-LUAD. However, variant prevalence differences between ES- and NS- LUAD could not be explained by differential mutagenesis. *KRAS*, *KEAP1*, and *STK11* mutations were more strongly selected in ES-LUAD. *EGFR*, *PIK3CA*, and *SMAD4* mutations were more strongly selected in NS-LUAD, wherein *EGFR* mutations were associated with upregulation of epithelial-mesenchymal transition—a functional consequence specific to exposure history.

Epistasis was pervasive and context-dependent: ES-LUAD exhibited more synergy and less antagonism, implying a broader, more navigable adaptive landscape than NS-LUAD. We also performed the first systematic quantification of higher-order-than-pairwise epistasis, revealing sub-additive and emergent synergies that shape cancer evolution. Our results disambiguate the mutagenic and selective effects of tobacco smoke, demonstrating how environmental insults reshape cancer evolution and enabling prediction of therapeutic efficacy based on smoking history and somatic genotype.

Author Summary

Smoking is a well-known cause of lung cancer because it increases the number of mutations in lung cells. But tobacco smoke also causes inflammation and other physiological changes that might influence how tumors evolve. Using DNA from lung tumors, we asked whether these changes make certain mutations more helpful to cancer growth and thereby better targets for therapy. We also asked whether some mutations were more helpful to cancer growth in those who haven't smoked. To do so, we estimated not only how often mutations occurred, but also how strongly they were favored during cancer development. Focusing on 21 key lung cancer genes, including *KRAS*, *EGFR,* and *TP53*, we found that mutations in some, such as *KRAS* and *KEAP1* were more strongly favored in smokers. Mutations in genes such as *EGFR* and *PIK3CA* were more strongly favored in non-smokers, where they were also linked to non-smoker-specific changes in gene activity. We also found that mutations in tumors interact differently depending on smoking history: smokers' tumors showed more cooperation between mutations, which may help them evolve in more flexible ways. This study shows how smoking reshapes the genetic paths lung cancers take—and why a patient's smoking history matters when choosing targeted treatments. Understanding these evolutionary patterns can help improve treatment decisions and lead to more personalized care for lung cancer patients.

Introduction

Tobacco smoke is well known to cause cancer, in part through its mutagenic effects [1]. Carcinogen exposure from tobacco generates distinct patterns of mutations [2], a subset of which then contribute to oncogenesis [3–9]. In addition to this mutagenic effect, tobacco smoke induces physiological changes in the lung—inflammation, oxidative stress, immune dysregulation, and tissue damage [1,10–12]—that can alter the lung microenvironment and thereby the selective pressures acting on specific somatic mutations. These microenvironmental

disruptions could modulate the fitness advantages conferred by specific mutations, thereby

influencing their likelihood of contributing to cancer development. Indeed, several studies have

suggested that, in principle, endogenous or exogenous alterations to the microenvironment

caused by factors such as aging or tobacco exposure likely shape the trajectories of cancer

evolution [13–17]. However, the extent to which such physiological effects alter the adaptive

landscape of cancer has not previously been quantified. Therefore, quantification of the

adaptive impacts of a highly disruptive exogenous insult such as tobacco smoke is crucial to

understanding how environmental exposures shape the evolution of individual cancer

evolutionary trajectories.

Among lung cancer subtypes, LUAD is the most common and accounts for the highest

proportion of cases in never-smokers [18]. As of 2002, nearly 23% of LUADs in the United

States occurred in never-smokers [19], a proportion that has been rising as cigarette usage

declines [20]. Numerous studies have reported that LUAD exhibits distinct somatic variant

profiles in never-smokers compared to ever-smokers, which has been taken as evidence of their

divergent genetic vulnerabilities [21–25]. However, differences in variant prevalence alone are

confounded as to whether they are a consequence of effects of tobacco-induced mutagenesis

or a consequence of tobacco-altered selection dynamics. Therefore, quantification of the

relative oncogenic roles of the mutagenic and physiological effects of smoking in LUAD will be

increasingly crucial to determination of optimal use of targeted therapies for patients with and

without a prior smoking history.

Computational models now enable inference of baseline somatic mutation rates from

synonymous mutations observed in tumor genomes—implicitly [6] or explicitly [26] accounting

for carcinogen-induced mutagenesis, such as that caused by tobacco smoke. Comparing these

baseline mutation rates to the observed prevalence of nonsynonymous mutations enables

estimation of somatic selection acting on specific substitutions [8,27,28]. This framework

provides a foundation for disentangling the oncogenic contributions of the mutagenic and

selective effects of smoking. Moreover, selection on each mutation is likely influenced not only

by the microenvironmental effects of tobacco smoke, but also by pairwise [28–30] and higher-

order [28] selective epistasis between mutations. In selective epistasis, mutations affect the

selection operating on other mutations. These interactions between the selective effects of

mutations shape the adaptive landscape of cancer evolution [5,28,31,32]. Understanding how

tobacco smoke influences not just mutation and selection, but also epistasis, is essential for a

complete picture of LUAD evolution.

Because knowledge of epistasis is crucial to inference of cancer-driving mutation modules

and synthetic vulnerabilities, a number of algorithms have been developed that attempt to detect

epistasis between mutations or sets of mutations by quantifying over- and under-

representation [33,34] in the context of heterogeneous mutation rates [30,35–37]. However, with

the exception of Coselens [30], these methods do not enable meaningful quantitative

comparisons of the effects of mutations in distinct somatic genotypes. This quantification is a

crucial step for prioritization and translation of knowledge of epistasis into effective therapeutic

strategies. Moreover, all of these approaches are restricted to pairwise inferences [30,34] or to

inference of discrete modules [33,35–37], and are not well suited to the quantitative inference of

higher-order epistasis (involving more than two mutations). When present, higher-order

epistasis substantially influences the adaptive landscape and shapes evolutionary trajectories

[38–43], including the degree to which selective epistasis diminishes along the adaptive

trajectory [44–47]. Systematic characterization of the presence and biological significance of

higher-order epistasis to cancer evolution has been challenging. Recent theory facilitates

understanding of higher-order epistasis in cancer by providing the selection strength of

mutations for each somatic genotype [28]. This novel approach has not yet been applied in the

context of environmental risk factors that could also affect selection, such as tobacco smoke.

Accordingly, analysis of the effect of smoke on the adaptive landscape of LUAD will benefit from

consideration of pairwise and higher-order epistatic effects and potential differences in epistasis

between microenvironments.

To address these gaps, we quantified the mutagenic, selective, and epistatic effects of

tobacco smoke on single-nucleotide variants (SNVs) in LUAD. We assembled a large dataset of

lung adenocarcinoma tumor genomes and classified them by smoking status into ever-smoker

(ES) and never-smoker (NS) groups. We estimated background mutation rates in each group

and inferred the strength of somatic selection acting on driver mutations in each gene. We then

investigated how selection pressures differed between ES- and NS-LUAD and associated these

differences with differential gene expression. Finally, we extended our analysis to inference of

selection in specific somatic genotypes, detecting and quantifying the extent of pairwise and

higher-order epistasis. The degree of divergence in the mutational, selective, epistatic, and

transcriptional landscapes of LUAD between ever-smokers and never-smokers enables novel

insight into how environmental exposure shapes cancer evolution—identifying genotype- and

environment-specific vulnerabilities that can inform personalized therapeutic strategies.

Results

Mutation rates and somatic selection pressures jointly shape the prevalence of

driver variants in lung adenocarcinoma

To disentangle the relative contributions of mutation and selection to observed mutation

frequencies in LUAD, we deconvolved the prevalence of somatic variants in our curated set of

21 well-supported driver genes into estimates of the background oncogenic mutation rates and

scaled selection coefficients. Across all smoking-status-classified LUAD samples in our dataset

(n = 1,722), *TP53* and *KRAS* were the most frequently mutated genes, followed distantly by

EGFR, *KEAP1*, and *STK11* (**Fig. 1A**). Background oncogenic mutation rates for the majority of

these genes spanned slightly more than an order of magnitude—between 7.6×10^{-8} and

8.7 × 10⁻⁷ (**Fig. 1B**). When compared to the observed variant prevalences, this variation in

background oncogenic mutation rate frequently implies substantial and variable impacts of

somatic selection on the prevalence of each variant.

Figure 1. Fixation rates and scaled somatic selection coefficients for 21 known and potential driver mutations among 1,722 lung adenocarcinomas. (**A**) Poisson-corrected fixation rates (rates at which each mutation occurs and becomes fixed within the population). (**B**) Background rate at which oncogenic gene mutations occur. (**C**) Strength of somatic selection (log-scaled) experienced by oncogenic gene mutations. Inset: Correlation between the prevalence of an oncogenic mutation and the strength of selection acting on it ($P < 0.001$).

Scaled selection coefficients for oncogenic mutations in each gene exhibited a long-tailed

distribution (**Fig. 1C**), with mutations in *KRAS* experiencing the strongest selection, followed

distantly by mutations of *TP53* and *CTNNB1*, then mutations in *ATM*, *KEAP1*, and *PIK3CA*, and

then other genes that experienced weaker but still substantial selection (**Fig. 1C**). Notably,

CTNNB1 mutations were present in only 3.7% of tumors. Nevertheless, knowledge of the low

underlying oncogenic mutation rate of *CTNNB1* indicates that these infrequent mutations

experienced stronger selection than did the more frequent mutations in *STK11, RBM10*, and

SMARCA4. Generally, selection and mutation prevalence were correlated ($r = 0.68$, $P < 0.001$),

but several discrepancies were present between the prevalence of a mutation and the relative

intensity of selection it experienced (**Fig. 1C,** inset). These discrepancies highlight the

importance of reporting selection as the effect size of mutations, rather than reporting solely

prevalence, which is only moderately associated with strength of selection.

The oncogenic mutation rates and selection coefficients presented above were quantified

based on all tumors, regardless of smoking status. However, both the mutagenic processes and

selective advantages of mutations were plausibly influenced by the genotoxic and physiological

effects of tobacco smoke. To assess these distinct effects, we stratified tumors by smoking

status and performed separate analyses of somatic selection in ES- and NS-LUAD tumors.

Mutation and selection differ substantially between ever-smoker and never-

smoker LUAD

Stratification of LUAD tumors by smoking status based on mutational signature attributions

and clinical annotation yielded 1,066 ES-LUADs and 656 NS-LUADs. In each group, we

independently estimated oncogenic mutation rates and selection coefficients for the 21 well-

established driver genes (**Fig. 2**). Both mutation rates and selection pressures differed markedly

between ES- and NS-LUAD. Oncogenic mutation rates were elevated in ES-LUAD across all 21

genes. Increases ranged from modest (39%–43% for *RB1* and *CTNNB1*) to extreme (571–

1474% for *EGFR*, *SMAD4*, *GNAS*, and *ALK*; **Fig. 2A inset**). These elevated mutation rates

demonstrate the intense mutagenic effect of tobacco smoking on the peripheral tissue of the

lung.

Figure 2. Mutational and adaptive landscapes of ever-smoker lung adenocarcinoma (ES-LUAD) and never-smoker lung adenocarcinoma (NS-LUAD). (**A**) Strengths of somatic selection (*y* axis is square-root scaled; *: non-overlapping 95% confidence intervals) for mutations of 21 known and potential driver genes in NS-LUAD (teal bars; *n* = 656) and ES-LUAD (salmon bars; *n* = 1,066), averaged across somatic genotypes. Inset: Background oncogenic mutation rates for these genes in ES-LUAD and NS-LUAD (dashed line: equality). (**B**) Differential expression of genes between *EGFR* wild-type (WT) and *EGFR* mutant samples in ES-LUAD (*n* = 273) and NS-LUAD (*n* = 59; log2FC: log₂ of fold change in expression) for genes deemed to have a significant gene expression interaction between *EGFR* allelic status and smoking status. Color of each gene name denotes whether the change in expression was more positive (red) or more negative (blue) in NS-LUAD compared to ES-LUAD (*: *P* < 0.05, •: *P* < 0.1, *SOX21*: *P* = 0.116; background color: change in gene expression from normal (*n* = 33) to LUAD (*n* = 332) is increased [pink], decreased [light blue], or insignificant [gray]). Leftward squares designate whether the gene produces a transmembrane (TM) or extracellular matrix (ECM) protein as well as whether it facilitates or interacts with the Wnt, and TGF-β and the epithelial-mesenchymal transition (EMT) signaling pathways. (**C**) Interactions between proteins (red: implicated in EMT) whose *EGFR*-mutation-associated expression changes differ between ES-LUAD and NS-LUAD. (**D**) Sample-specific DNA replication expression scores for *KEAP1* WT and mutant ES- and NS-LUAD (**: *P* < 0.01).

We found that reported differences between ES- and NS-LUAD in the prevalences of key

driver mutations [21,22,48] were often driven by differential selection, rather than solely by

differential mutagenesis (**Fig. 2A**). Among the 21 driver genes analyzed, *TP53*, *EGFR*, *PIK3CA*,

SMAD4, *MET*, *MGA*, and *GNAS* mutations experienced stronger positive selection in NS-LUAD.

STK11, *KEAP1*, and *KRAS* mutations experienced stronger positive selection in ES-LUAD.

RB1, *BRAF*, and *BRCA2* mutations experienced similar selection in both cancer types.

Most mutations that experienced significantly different selection between ES- and NS-

LUAD were strongly selected in one context and weakly selected in the other, demonstrating

pronounced context-dependence in LUAD evolution. In ES-LUAD, *KEAP1* mutations

experienced the third strongest selection—after *TP53* and *KRAS*—and experienced over three

times stronger selection than in NS-LUAD. Likewise, *STK11* mutations experienced over twice

as much selection in ES-LUAD. Indeed, both *KEAP1* and *STK11* mutations are enriched in

ever-smoker populations [23,49,50]. *EGFR* mutations exhibited the most dramatic difference:

they were more than twenty times more strongly selected in NS- than ES-LUAD, and were also

the third most strongly selected mutations in NS-LUAD after *TP53* and *KRAS*. *EGFR* mutations

are much more common in NS-LUAD than in ES-LUAD [21,51]. Similarly, *PIK3CA*, *SMAD4*, and

MET mutations experienced over three times stronger selection in NS- relative to ES-LUAD.

To investigate mechanisms underlying the differential selection of *EGFR* mutations, we

assessed their differential transcriptional effects in ES- and NS-LUAD using TCGA expression

data. In fewer than one percent of genes did the expression fold change from *EGFR* wild type

samples compared to *EGFR* mutant samples differ significantly between ES- and NS-LUAD

(**Fig. 2B**). However, several of these genes exhibited markedly stronger upregulation or

downregulation in EGFR-mutant NS-LUAD compared to ES-LUAD. Genes that were

upregulated were frequently components of Wnt, TGF-β, and Hedgehog signaling pathways or

of the extracellular matrix (**Fig. 2B–C**). Indeed, in an analysis of gene sets upregulated in NS-

versus ES-LUAD, the Hallmark gene set associated with epithelial-mesenchymal transition was

the most enriched. On the other hand, *KEAP1* mutations were associated with a substantially

higher DNA replication enrichment score in ES-LUAD but not in NS-LUAD (**Fig. 2D**). These

examples relate microenvironmental context to modifications of the downstream effects of driver

mutations and their accompanying selective advantage.

Extending our analysis from 21 to 1,200 LUAD-associated genes revealed a striking

asymmetry in selection by smoking status: only six genes—*KRAS*, *KEAP1*, *STK11*, *PBRM1*,

DNMT3B, and *CDH1*—were significantly more strongly selected in ES- than NS-LUAD, whereas

87 experienced stronger selection in NS-LUAD (**Fig. S1**). Among the mutations most strongly

favored in NS-LUAD were those in *EGFR*, *SMAD4*, *SMAD2*, *PTEN*, *PIK3R1, PAK1*, *FGF3*,

GATA3, *JUN*, *CTNNA2*, and *TLR4*. Mutations often experienced weaker selection in ES-LUAD,

which is more heavily mutated and frequently possesses multiple known drivers, than in NS-

LUAD, which often features a single known driver (**Fig. S1**).

Overall, somatic mutations often experienced substantially altered selection in the

tobacco-smoke-altered ES-LUAD compared to NS-LUAD, contributing to differential evolution of

the two cancers. Along with the somatic environment, the somatic genotype can also induce

differential selection on specific mutations: effects of each mutation can be analyzed across

somatic genotypes to reveal selective epistasis that shapes the evolutionary trajectories of

tumors in ever-smokers and never-smokers. Therefore, we assessed the influence of pairwise

selective epistasis.

Pairwise epistasis shapes the adaptive landscape of ES-LUAD and NS-LUAD

To assess how frequently and how strongly the evolutionary trajectory of LUAD is affected

by pairwise epistasis, we quantified the effects of driver mutations both in isolation (in tumors

that were wild type at a paired gene) and in the presence of each paired gene mutation. For ES-

and NS-LUAD, we evaluated all 420 possible gene pairs among the 21 driver genes: pairwise

interactions were identified that modulate the selective advantage of mutations, and did so in a

smoking-dependent manner.

Mutation of TP53 *followed by* RB1 *is the most likely trajectory for ES-LUAD and NS-*

LUAD

To investigate directional epistasis between two common LUAD drivers, *TP53* and *RB1*,

we first stratified tumors by smoking status and somatic genotype at these two loci, and then

estimated the fixation rates of each mutation from each somatic genotype (**Fig. 3A**). In ES-

LUAD, *TP53* mutations exhibited a high fixation rate when *RB1* was wild-type but were inferred

to exhibit a negligible fixation rate when *RB1* was mutated (λ, **Fig. 3B**). Conversely, *RB1*

mutations exhibited a low fixation rate when *TP53* was wild-type but exhibited a high fixation

rate when *TP53* was mutated. These asymmetries indicate the most likely evolutionary

sequence in ES-LUAD: mutation of *TP53* followed by mutation of *RB1*. In wild-type tumors, the

mutation rate of TP53 was over double the mutation rate of *RB1* ($\mu_{WT \rightarrow TP53} = 7.4 \times 10^{-7}$,

$\mu_{WT \rightarrow RB1} = 2.0 \times 10^{-7}$); the same was true of mutant tumors ($\mu_{RB1 \rightarrow RB1+TP53} = 8.9 \times 10^{-7}$,

$\mu_{TP53 \rightarrow RB1+TP53} = 3.6 \times 10^{-7}$; **Fig. 3B**). Accordingly, *RB1* variants in a *TP53*-mutant genotype must

have experienced comparable selection to *TP53*-variants in a wild-type genotype (γ, **Fig. 3B**),

despite the substantially lower fixation rate of the *RB1* variants. This pattern reflects synergistic

epistasis: prior mutation of *TP53* increases selection for *RB1*. In contrast, the reverse order

exhibited antagonistic epistasis—*TP53* was less strongly selected in the presence of *RB1*

mutation—though this last effect did not achieve statistical significance.

Figure 3. Pairwise selective epistasis inferred within the evolutionary trajectories of lung adenocarcinoma (LUAD), depicted for *TP53* and *RB1*. (**A**) Genomic sequence data from lung adenocarcinomas can be stratified by clinically- and genetically-determined patient smoking status (blue: never-smokers; red: ever-smokers) and somatic genotype. Analysis of (**B**) ever-smoker LUAD and (**C**) never-smoker LUAD deconvolves somatic genotype prevalences (gray circle radius) into genotype-specific fixation rates (λ; arrow width), oncogenic mutation rates (μ; arrow width), and strengths of somatic selection (γ; arrow width) on mutations of *TP53* (dark blue arrows) and *RB1* (brown arrows).

In NS-LUAD, the fixation rate for *TP53* mutations from a wild-type genotype was lower

than in ES-LUAD, and the fixation rate for *RB1* mutations in a *TP53*-mutant background was

higher (λ, **Fig. 3C**). In single-mutant tumors, the mutation rate of *TP53* is nearly double that of

RB1 ($\mu_{RB1 \to RB1+TP53}$ = 4.0 × 10^{-7}, $\mu_{TP53 \to RB1+TP53}$ = 2.2 × 10^{-7}), but in wild-type NS-LUAD tumors,

the mutation rate of *TP53* was similar to the mutation rate of *RB1* ($\mu_{WT \to TP53}$ = 1.48 × 10^{-7},

$\mu_{WT \to RB1}$ = 1.8 × 10^{-7}). This similarity of mutation rates contrasts with the disparity in ES-LUAD.

Despite these differences, the inferred evolutionary trajectory mirrored that of ES-LUAD: the

most likely (λ) and most adaptive (γ) path involved mutation of *TP53* followed by mutation of

RB1. As in ES-LUAD, prior *TP53* mutation significantly synergistically enhanced selection for

RB1 mutation, and prior *RB1* mutation antagonized selection for *TP53* mutation, without

reaching statistical significance (γ, **Fig. 3C**).

Synergistic epistasis is widespread and often substantial in ES-LUAD

Analysis of all pairwise combinations of the 21 reputed LUAD driver genes revealed

frequent synergistic and antagonistic epistasis. In ES-LUAD, synergistic interactions were

especially pronounced: for genes whose mutations were under strong positive selection (with

the notable exceptions of *TP53* and *KRAS*) synergistic epistasis led to two- to eight-fold

increases in selection. For example, selection on mutations in *KEAP1, ATM*, *BRCA2*, *PIK3CA*,

and *ARID1A* was markedly enhanced in the presence of prior *TP53* or *KRAS* mutations

(**Fig. 4A**). Additional strong synergistic interactions included those affecting mutants of *ARID1A*

in the context of somatic genotypes with *ATM* or *RBM10* mutations; mutants of *STK11* in the

context of mutated *KEAP1*; and mutants of *ATM* in the context of mutated *STK11*. Mutations in

the non-*TP53* tumor suppressors *BRCA2*, *SETD2*, *RBM10*, *RB1*, *MGA*, and *APC*, as well as in

the chromatin-remodeling genes *ARID1A* and *SMARCA4*, experienced moderate baseline

selection that was frequently enhanced by the presence of diverse prior mutations. These genes

never experienced antagonistic epistasis with each other and rarely with other mutations.

Figure 4. Pairwise epistasis in ever-smoker (ES) and never-smoker (NS) lung adenocarcinoma (LUAD). Strengths of somatic selection for mutations in wild-type tissue (green) and in previously mutated tissue (gray, labeled with previous mutation in brackets) in (**A**) ES-LUAD and (**B**) NS-LUAD. Significant pairwise epistatic interactions depicted in heatmap form for (**C**) ES-LUAD and (**D**) NS-LUAD based on non-overlapping confidence intervals for epistatic strengths of selection. Heatmap values are ratios of selection coefficients for mutations (in each row) from a somatic genotype containing a mutation in another gene (in each column) to the selection coefficient from a somatic genotype lacking the mutation. (**E**) Strengths of somatic selection for mutation of *KRAS* in *TP53*-mutant and *TP53*-wild-type somatic genotypes in ES-LUAD (red) and NS-LUAD (blue).

277   Patterns of epistasis involving oncogenes were heterogeneous. Notably, *KRAS* and

278 *CTNNB1* mutations showed no elevation of selection in the presence of any prior mutations in

279 the 21-gene set. Conversely, selection for mutations of *GNAS* and *ALK* was enhanced by three

280 prior mutations (**Fig. 4A**). For many oncogenes, selection was enhanced by prior *TP53*

281 mutation, but was antagonized by mutations in other oncogenes such as *MET*. Across all tested

282 genes, the median effect size of statistically significant synergistic epistasis was a 3.6× increase

283 in selection, and the magnitude of this synergy did not correlate with the baseline strength of

284 selection for either the first or the second mutation.

286 *Antagonistic effects can reduce positive selection or induce negative selection on*

287 *drivers*

288   In ES-LUAD, we identified eight gene pairs exhibiting statistically significant antagonistic

289 sign epistasis, in which selection for a driver mutation shifted from positive to negative in the

290 presence of another mutated gene. Mutations in *CTNNB1* and *GNAS* became selectively

291 disadvantageous in the presence of *MET* and *APC* mutations. Similarly, selection for *SMAD4*

292 mutations was antagonized by prior *BRCA2* and *PIK3CA* mutations, and *MET* mutations were

293 antagonized by mutations of *RB1*, *EGFR*, *CTNNB1*, and *GNAS*. In each of these eight cases,

294 the reciprocal mutation order also induced negative selection (e.g., mutated *CTNNB1* induced

negative selection for *MET* mutations). For over a hundred other ordered pairs, antagonistic

sign epistasis was estimated, without reaching statistical significance. Antagonistic shifts in the

magnitude of selection, in which selection for a mutation remained positive but was substantially

reduced, were indicated in 33 cases, of which three were significant: the presence of *KRAS*

mutations caused a 64% reduction in selection for *TP53* mutations and an 81% reduction in

selection for *EGFR* mutations, and *TP53* mutations caused a 58% reduction in selection for

KRAS mutations (**Fig. 4A**). Across all analyses of ES- and NS-LUAD, new mutations in *TP53*

and *KRAS* were never inferred to experience significantly stronger selection in the context of

another driver mutation.

Antagonistic epistasis is more frequent in NS-LUAD

Compared to ES-LUAD, NS-LUAD exhibited a higher frequency of antagonistic epistasis

and fewer instances of synergistic epistasis (**Fig. 4B**). Of the 65 gene pairs exhibiting significant

antagonistic epistasis in ES-LUAD, 25 also reached significance in NS-LUAD, and an additional

36 exhibited antagonistic trends without achieving significance (**Fig. 4C–D**). Thus, instances of

selective antagonism inferred in ES-LUAD were nearly always present in NS-LUAD.

This consistency contrasted with a substantial inconsistency in selective synergies

between ES- and NS-LUAD. Of the 42 gene pairs whose mutations exhibited significant

synergistic epistasis in ES-LUAD, only five were significantly synergistic in NS-LUAD. An

additional 25 trended synergistic without reaching significance. Strikingly, 16 of these 42

instances were inferred to be significantly antagonistic in NS-LUAD, including many involving

BRCA2. For example, *KEAP1* mutation enhanced selection for *BRCA2* mutation in ES-LUAD

but antagonized it in NS-LUAD (**Fig. 4C–D**). These reversals reveal distinct adaptive constraints

imposed by the smoking-altered tumor microenvironment. Overall, antagonistic interactions

were relatively stable across smoking status, whereas synergistic epistasis was highly

contingent on smoking status.

Mutations in these 21 driver genes were subject to a broad range of epistatic and

environmental interactions, as well as combinations thereof (**Fig. S2**). For example, *TP53*

mutations antagonized *KRAS* mutations in ES-LUAD but had no substantial effect on *KRAS*

mutations in NS-LUAD (**Fig. 4E**). These context-dependent epistatic effects substantially

contribute to the divergent trajectories of ES- and NS-LUAD.

Fewer examples of synergistic epistasis were observed in NS-LUAD compared to ES-

LUAD. Nevertheless, synergistic epistasis contributed to a median twelve-fold increase in

selection on secondary mutations. Notable examples include enhanced selection for *CTNNB1*,

RB1, and *RBM10* mutations after *EGFR* mutation, for *PIK3CA*, *BRCA2*, *SETD2*, and *ARID1A*

mutations after *ATM* mutation, and for *SETD2* mutation after *ARID1A* mutation (**Fig. 4B**). Even

in relatively less mutated NS-LUAD, cooperative genetic interactions dramatically reshape the

fitness landscape of evolving tumors.

Pairwise epistasis can be asymmetric in both sign and magnitude

In both ES- and NS-LUAD, several gene pairs exhibited asymmetric epistasis, in which the

direction or strength of interaction depended on the order of mutation acquisition. For instance,

in ES-LUAD, mutation of either *KEAP1* or *STK11* enhanced selection on mutations of the other,

but with markedly different magnitudes: primary *STK11* mutation was estimated to increase

selection for a secondary *KEAP1* mutation by 140%, whereas primary *KEAP1* mutation

significantly increased selection for a secondary mutation of *STK11* by 360%. In other cases,

the sign of epistasis reversed depending on mutational order. Most notably in ES-LUAD,

primary *KRAS* mutation synergistically increased selection for a secondary *KEAP1* mutation, but

primary *KEAP1* mutation strongly antagonized selection for a secondary *KRAS* mutation

(**Fig. 4A**). This asymmetry means that a mutation in *KRAS* followed by a mutation in *KEAP1*

was advantageous, but a mutation in *KRAS* was disadvantageous or lethal when preceded by a

mutation in *KEAP1*. This dramatic directional asymmetry and eighteen other such instances

exemplify how somatic evolutionary trajectories can be constrained or potentiated by the

specific sequence in which mutations occur.

Higher-order epistasis shapes evolutionary trajectories of ES-and NS-LUAD

KRAS*,* KEAP1*, and* STK11 *constitute an epistatically interacting triad in ES-LUAD*

Having identified *KRAS*, *KEAP1*, and *STK11* as experiencing strong, ever-smoker-specific

selection, we next investigated whether their interactions extended beyond pairwise epistasis to

form a higher-order epistatic network. Mutations in all three genes exhibited substantial fixation

rates from nearly any initial somatic genotype, particularly en route to the triple-mutant

genotype. Fixation rates for *KRAS* mutations were estimated to be very low when arising after

KEAP1 or *STK11* mutations alone compared to when arising as the initial mutation or in the

KEAP1-STK11 double-mutant (**Fig. 5A**). Nevertheless, these three genes form an epistatic triad

in which mutations in any one tend to increase selection for the others.

Figure 5. Evolutionary trajectory for *KRAS*, *KEAP1*, and *STK11* mutations in ever-smoker lung adenocarcinoma. (**A**) Fixation rates, (**B**) oncogenic mutation rates, and (**C**) scaled selection coefficients for mutations of *KRAS* (orange arrows; arrows width proportional to rate or coefficient), *KEAP1* (light blue arrows), and *STK11* (yellow arrows) conditioned on somatic genotype (gray circles; radius proportional to genotype prevalence).

Among the three genes, *STK11* manifested the highest oncogenic mutation rate at each

progressive step, followed by *KEAP1*, and then *KRAS* (**Fig. 5B**). Given these differential

underlying rates of mutation, we computed the consequent scaled selection coefficients for

mutations in each genotype in the three-gene model, and compared selection between

genotypes to quantify pairwise and higher-order selective epistasis. Many tumors possessed

mutations in at least one of these genes (**Fig. 5C**, second column of nodes), providing high

power to detect pairwise epistasis. Indeed, our three-gene model recapitulated the pairwise

epistatic relationships inferred from the two-gene models (**Fig. 4C**). However, concurrent

mutations of two or more of these genes were relatively infrequent (**Fig. 5C**, third and fourth

columns of nodes), limiting power to detect higher-order epistasis with statistical confidence. For

example, substantial uncertainty in the selection estimate for *KRAS* mutations in tumors

harboring *STK11* and *KEAP1* mutations precluded a definitive inference of higher-order

interactions. In contrast, because of the sufficiently high prevalence of *KRAS-KEAP1-* and

KRAS-KEAP1-STK11-mutant genotypes, we were able to infer higher-order epistatic effects on

STK11 with greater confidence.

STK11 mutations experienced significantly stronger selection in the presence of either

KRAS or *KEAP1* mutations. If the synergistic effects of *KRAS* and *KEAP1* mutations on *STK11*

compounded additively, then *STK11* mutations would experience selection equivalent to the

three-way product of the individual synergistic effects and the baseline selection for *STK11*

mutations. However, the upper bound for selection for *STK11* mutations in a *KRAS-KEAP1*

double-mutant genotype was less than this product (**Fig. 5C**). Accordingly, the compound

synergistic effect of concurrent *KEAP1* and *KRAS* mutations represents an example of sub-

additive higher-order epistasis.

Most pairwise epistatic effects compound without strong higher-order interaction

The sub-additive epistasis affecting mutations of *STK11* demonstrates that assessments

of higher-order epistasis will be necessary to thoroughly understand cancer evolution. However,

most effects of driver mutations compounded in a manner consistent with minimal higher-order

epistasis. Examining all 1330 triads out of the 21 genes, we compared selection for a new

mutation in single- versus double-mutant backgrounds. In the vast majority of cases, a

combination of pairwise epistatic effects was sufficient to explain significant differences in

selection for a new mutation between single- and double-mutant genotypes. Generally, when

selection for a secondary mutation in one gene was epistatically altered by presence of a

primary mutation in a second gene and was unaffected by a primary mutation in a third gene,

selection on the secondary mutation remained commensurately epistatically altered when both

other variants of the triad were concurrently present. For example, in ES-LUAD, selection for

SMARCA4 mutations was synergistically enhanced by *STK11* mutations but unaffected by

TP53 mutations, and correspondingly, selection for mutations in *SMARCA4* was synergistically

increased in a *STK11-TP53*-mutant genotype (**Fig. 6A,B**). Similarly, in NS-LUAD, selection for

RBM10 mutations was synergistically increased in both a *TP53*-mutant and *TP53-EGFR*-mutant

genotype (**Fig. 6C**). Overall, pairwise interactions typically dominate the adaptive landscape of

LUAD, but higher-order epistasis can have substantial influence on specific evolutionary

trajectories.

Figure 6. Higher-order selective epistasis in lung adenocarcinoma (LUAD). (**A**) Selection within gene triads (*y* axis is square-root scaled), for a mutation (bold font) in a context of no co-occurring mutations (circle), one co-occurring mutation (square; labeled with first letter(s) of the gene(s) indicated in the tick label that are mutated), and two co-occurring mutations (diamond) in ever-smoker (ES) LUAD [pink: visualized in panels B and F; purple: visualized in panel D; green: visualized in panel E]. (**B**) In ES-LUAD, the synergistic effect of *SMARCA4* mutations (light blue arrows) on selection for *APC* mutations (pink arrows; arrow width proportional to scaled selection coefficient; circle radius proportional to somatic genotype prevalence) does not diminish in the presence of a neutrally-interacting *KRAS* mutation (orange arrows). (**C**) Selection within gene triads in never-smoker (NS) LUAD; *y* axis is square-root scaled, symbols are as in panel A]. (**D**) Pairwise and higher-order epistasis in ES-LUAD: co-mutation of *TP53* (blue) and *KRAS* (orange), each of which independently increase selection for *ARID1A* mutations (purple), produces an even stronger synergy for *ARID1A* mutations; and (**E**) antagonistic effect of *KRAS* mutations (orange) on selection for *EGFR* mutations (green) outweighs the synergistic effect of *TP53* mutations (blue). (**F**) In ES-LUAD, co-mutation of *KEAP1* (light blue) and *STK11* (yellow) synergistically increases selection for *APC* mutations (pink) despite neutral pairwise effects. (**G**) Comparison of the epistatic effect of double-mutant-genotype (label: bottom row, two-letter abbreviation) on selection for new mutation (label: top row) to the "expected" product of pairwise epistatic effects (blue points: both pairwise epistatic effects are statistically significant, but not the higher order effect; green: the higher-order effect is significant, but neither pairwise effect is; red: all effects are significant)

When two co-occurring mutations each independently enhanced selection for a third

mutation, the synergistic effects frequently compounded. For instance, *ARID1A* mutations

experienced increased selection in both KRAS- and *TP53*-mutant genotype. These synergistic

effects compounded, resulting in mutations of *ARID1A* experiencing three times stronger

selection in a *KRAS-TP53*-mutant genotype than in a *TP53*-mutant genotype and nine times

stronger selection than in a wild-type genotype (**Fig. 6D**). Similar patterns of compounded

synergy were evident for *ATM* mutations in *KRAS-STK11*-mutant genotypes, as well as for

SMARCA4 mutations in a *KEAP1-STK11*-mutant background. Across cases, compounded

synergy frequently drove selection to levels up to an order of magnitude higher than in

genotypes lacking either partner mutation. This substantial compounding constitutes a crucial

element of how driver mutations accumulate in cancer genomes, positively reinforcing

progression of cancer toward more aggressive forms.

Higher-order epistasis often manifests as sub-additive or emergent synergistic

interactions

In the case of compounding synergistic effects, it is also possible to observe super-additive higher-order epistasis, in which selection for a mutation in the presence of two independently synergistic mutations is even larger than what would be expected from the product of their independent synergistic effects. Indeed, some cases are consistent with either additive synergy or super-additive epistasis, such as mutations of *SMARCA4* in a *KRAS-KEAP1*-mutant genotype in ES-LUAD (**Fig. 6A**). More frequently, cases are consistent with either additive synergy or sub-additive epistasis, in which selection for a mutation in the presence of two independently synergistic mutations is smaller than expected from the sum of synergistic effects. Indeed, in 43 of 45 cases wherein a new mutation arose in a genotype of two significantly synergistic mutations, estimates of selection for the new mutation were lower than would be expected from purely additive synergistic epistasis. Such widespread sub-additivity pervades the adaptive landscape, tempering rather than amplifying selective pressures, even in the context of strong pairwise synergy.

We also identified cases of higher-order epistasis involving the co-occurrence of synergistic and antagonistic partner mutations, as well as the emergence of synergy in the co-occurrence of neutrally-interacting partner mutations. In ES-LUAD, for example, selection for *EGFR* mutations was synergistically enhanced by prior *TP53* mutations but antagonized by *KRAS* mutations. In a *TP53-KRAS*-mutant genotype, *EGFR* mutations experienced significantly less selection than in a *TP53*-mutant *KRAS*-wildtype genotype (**Fig. 6A,E**). In fact, these *EGFR* mutations experienced levels of selection similar to those they would experience in a genotype lacking either mutation, indicating a neutralizing result of the combination of synergistic and antagonistic epistasis.

Another form of higher-order epistasis is the emergence of synergy or antagonism from a

combination of mutations, each of which does not individually exhibit pairwise epistasis with the

affected gene. We observed several instances in which selection for a mutation was significantly

increased by the co-occurrence of two mutations that did not individually have significant

synergistic effects: in ES-LUAD, selection for mutations of *APC* increased by an order of

magnitude in a *KEAP1-STK11*-mutant genotype despite not being significantly altered when

only one of *KEAP1* or *STK11* was mutated (**Fig. 6A**,**F**). In this instance, significance was

reached for synergy when compared to wild-type or a *KEAP1*-mutant genotype, but not when

compared to the *STK11*-mutant genotypes, which was itself epistatically neutral. We detected

multiple similar instances of emergent synergy, such as for *ARID1A* mutation in an *EGFR-*

PIK3CA-mutant genotype in NS-LUAD (**Fig. 6G**). We also observed one instance of emergent

antagonism, in which *STK11* mutations were antagonized in a *TP53-EGFR*-mutant genotype

but not in either single-mutant genotype alone.

Overall, we observed two major classes of higher-order epistasis: sub-additive synergy

and emergent—and consequently super-additive—synergy. In particular, when both component

epistatic effects were strongly synergistic, their combination was almost always sub-additively

synergistic (**Fig. 6G**). On the other hand, when both component epistatic effects were neutral or

insignificant, a super-additive synergistic effect occasionally emerged from their combination

(**Fig. 6G**). Interestingly, as the strength of the component epistatic effects increases, the inferred

epistatic effect of the double-mutant combination rapidly plateaus at an approximately ten to

twenty-fold increase in selection.

In this survey of the adaptive landscape of LUAD, we discovered several cases of higher-

order epistasis that substantially influence the selective advantage of oncogenic mutations.

Over a third of LUAD samples in ever-smokers possessed mutations in three or more of 21

tested genes—a proportion that would likely rise in an analysis of a greater number of driver

genes. In these patients, higher-order epistasis may play a substantial role in the evolutionary

trajectories of their tumors and their responses to targeted therapies. Indeed, we found that a

wide variety of adaptive trajectories ensued from the combined epistatic effects of multiple

mutations. Some of these trajectories cannot be predicted from pairwise analyses. However, our

large-scale analysis of the adaptive landscape of 21 mutations has significantly supported only

a subset of the many potential instances of higher-order epistasis between oncogenic mutations

in LUAD. Indeed, multiple other instances of higher-order epistasis were estimated to be

present. Their effect would be clarified with greater sample sizes.

Discussion

479
Here we have characterized the distinct adaptive landscapes of lung adenocarcinoma in

ever-smokers and never-smokers by first estimating gene-specific background mutation rates in

ES- and NS-LUAD, then comparing the intensity of somatic selection and selective epistasis

acting on those mutations between the two groups. Consistent with the established mutagenic

effects of tobacco carcinogens [4], we detected elevated oncogenic mutation rates in all tested

genes in ES- compared to NS-LUAD. Accounting for these background rates revealed that

mutations in genes such as *EGFR*, *PIK3CA*, and *MET* provide a stronger selective advantage in

NS-LUAD, while mutations in *KRAS*, *KEAP1*, and *STK11* provide a stronger selective

advantage in ES-LUAD. Focusing on *EGFR* and *KEAP1*, we found that these adaptive

differences were at least partially explained by the differential effects of these mutations on

epithelial-mesenchymal transition and DNA replication transcriptional program activity in ES-

and NS-LUAD.

By quantifying the selective advantage of these mutations within specific somatic

genotypes, our analysis recovered known epistatic interactions—such as the enhancement of

selection on *RB1* and *EGFR* mutations by prior *TP53* mutations—and revealed context-specific

interactions, such as the synergistic effect of *EGFR* mutations on *CTNNB1* mutations uniquely

in NS-LUAD. We also detected substantial asymmetries: *CTNNB1* mutations did not impact

EGFR mutations in ES- or NS-LUAD. Notably, we found more frequent synergy in ES-LUAD

and substantially more pervasive antagonism in NS-LUAD, constituting fundamental differences

in the evolutionary navigability of their adaptive landscapes. Finally, we showed that higher-

order epistasis—particularly sub-additive synergistic and emergent synergistic epistasis—is

frequent in and substantially guides the evolutionary trajectories of cancer. These findings of

differential selection and epistasis reveal that exposure to tobacco smoke reshapes the adaptive

landscape of LUAD not only by elevating mutation burden, but also by altering the selective

value of mutations and their epistatic interactions—highlighting opportunities to leverage these

adaptive differences for biomarker development and therapeutic stratification.

The prevalences of driver mutations have been reported to substantially differ between

smokers and never-smokers in LUAD [21,22] as well as in non-small cell lung cancer [NSCLC;

23,24]. However, previous studies have either refrained from explaining these differences or

attributed them primarily to tobacco-induced mutagenesis [22,23]. Our results show that they

often cannot be explained by differential mutagenesis alone. For example, *KRAS* variants are

over five times more frequent in ES- than NS-LUAD tumors, but the background, neutral rate at

which *KRAS* mutations occur is only about three times higher in ES-LUAD. To resolve this

discrepancy, *KRAS* mutations must be nearly twice as oncogenic in ES- compared to NS-

LUAD. In general, the differential prevalence of these driver mutations is often not solely

attributable to tobacco-induced mutagenesis, but also due to their altered selective advantage in

the tobacco-altered lung environment. This physiological-adaptive effect of smoking is crucial to

account for: changes in the selective advantage of a mutation suggest altered functional impact

within a given tissue environment, with direct implications for the efficacy of targeted therapies

and the interpretation of biomarkers in smokers versus non-smokers.

Multiple perspectives have proposed that the adaptive landscape of oncogenesis is

altered by physiological insults from endogenous and exogenous factors [14,17,52,53].

However, data-driven evidence for this hypothesis has remained limited [cf. 54]. Here, we

provide compelling evidence that canonical driver mutations confer altered selective advantages

in the smoking-affected lung microenvironment of ever-smokers with LUAD. These shifts in

selection are linked to context-dependent phenotypic effects of mutations. For example, *EGFR*

mutations are associated with greater activity of the epithelial-mesenchymal transition (EMT)

transcriptional program in NS-LUAD, but not in ES-LUAD—likely explaining their stronger

selective advantage in never-smokers. Such divergent phenotypic consequences can be

attributed to smoking-induced alterations to transcriptional program activity [55–57], tumor

microenvironment composition [58], and cell fitness. These findings provide a mechanistic basis

for how exogenous exposures can reshape the evolutionary trajectory of cancer by altering the

adaptive value of oncogenic mutations.

In the case of *TP53*, Rodin and Rodin [13] reported comparable background mutation

rates between smokers and nonsmokers with lung cancer, and attributed the higher frequency

of *TP53* mutations in smokers to increased selection in the smoking-altered microenvironment.

By contrast, our analysis—leveraging a substantially larger dataset, incorporating tissue- and

context-specific covariates, and applying advanced statistical methodologies to estimate

background mutation rates [6]—identified a markedly elevated *TP53* mutation rate in ES-LUAD.

With this substantial differential mutagenesis accounted for, our analysis revealed that *TP53*

mutations are actually under weaker positive selection in ever-smokers than in never-smokers.

This finding challenges earlier conclusions and underscores the importance of accurately

resolving both mutational processes and selection when interpreting variant prevalence across

environmental exposures.

Despite mounting evidence of the substantial biological impact of tobacco smoke, clinical

trials for lung cancer therapies often do not assess the influence of smoking status—and

frequently do not even collect usage history [59], perhaps due to a conception that tobacco

smoke has a negligible effect on outcomes [60]. However, our findings reveal that tobacco

smoke profoundly reshapes the evolutionary trajectory of lung adenocarcinoma, necessitating

more systematic incorporation of smoking history into trial design and treatment decision-

making, particularly for targeted therapies. For example, the substantially stronger selection for

EGFR mutations in NS-LUAD implies that EGFR tyrosine kinase inhibitors (TKIs) should yield

greater clinical benefit in never-smokers. Indeed, a retrospective analysis of patients with

EGFR-mutant LUAD found that the objective response rate (ORR) of patients receiving EGFR

TKIs declined from 73% among never-smokers to 46% among individuals with over 10 pack-

years of smoking history [61]. Similar patterns have been reported in additional retrospective

studies of LUAD [62] and across NSCLC more broadly [63,64]. This concordance between

selection-based predictions and observed outcomes highlights the translational value of

selection coefficients as predictors of targeted therapy efficacy. By identifying likely responders

based on environmental exposures that shape the tumor microenvironment, exposure-informed

selection coefficients can improve clinical trial stratification, enhance statistical power, and refine

treatment personalization in precision oncology.

Prior studies have questioned whether computational analyses can reliably detect

epistasis, citing concerns about confounding by subtype-specific mutation profiles, variation in

mutation burden, and limited statistical power [31,65]. To address these concerns, we analyzed

ES- and NS-LUAD independently and comparatively, modeled subtype- and gene-specific

background mutation rates and effect sizes of mutations in specific somatic genotypes instead

of suboptimal measures of co-occurrence or mutual exclusivity, and integrated data from

multiple cohorts to ensure sufficient statistical power. Consequently, we were able to make

robust inferences of both pairwise and higher-order selective epistasis, many of which align with

experimental and clinical evidence. For example, consistent with our finding that *KRAS*

mutations synergistically enhance selection for *KEAP1* and *STK11* mutations, prior studies

show that *KEAP1* loss accelerates LUAD progression in *KRAS*-mutant mouse tumors [66,67]

and compensates for the deleterious effects of *KRAS*-*STK11*-co-mutation [68], and mutations of

KEAP1 and/or *STK11* worsen survival and treatment response in *KRAS*-mutant LUAD [69,70]

and NSCLC [71–76]. Furthermore, in accordance with the inferred mutually synergistic effects of

mutations of *KEAP1* and *STK11* in ES-LUAD, co-mutation of *KEAP1* and *STK11* promote cell

proliferation [77] and worsen survival in NSCLC independently of *KRAS* [78,79].

Our analyses also identified synergies between *EGFR* and other drivers: *RBM10* and

CTNNB1 in NS-LUAD, and *RB1* in both NS- and ES-LUAD. These interactions are corroborated

by experimental and clinical studies showing that *RBM10* mutations promote *EGFR*-mutant

LUAD [80] and confer resistance to EGFR TKIs [81], that *CTNNB1* mutations are enriched in

advanced *EGFR*-mutant NSCLC and associated with metastasis and TKI resistance [82], and

that *RB1* mutations in *EGFR*-mutant LUAD mediate histologic transformation to small-cell lung

cancer [83,84]. We also revealed that mutations of *TP53* and *KRAS* were widely antagonized by

other driver mutations, consistent with their early, clonal emergence during LUAD evolution

[85,86] and supporting their frequently primary role in tumor initiation. This panoply of

alignments between our results and experimental and clinical evidence encourages deeper

investigation into the many epistatic relationships we have revealed that remain untested in the

laboratory or clinic. They represent fertile ground for mechanistic investigation and promise new

therapeutic targets and biomarker candidates.

Previous studies have shown that epistatic interaction networks vary across cancer types

in humans [87] and across environmental contexts in other organisms [88–93]. However,

extensive differences in selective epistasis within a single cancer type exposed to distinct tissue

environments have not previously been demonstrated. Here, we have shown that tobacco

smoke exposure markedly reshapes the selective epistatic interactions underlying LUAD

evolution. Broadly, synergistic epistasis was more frequent in ES-LUAD, whereas antagonistic

epistasis was more frequent in NS-LUAD. Indeed, in some cases, epistasis between driver

genes was synergistic in ES-LUAD, but antagonistic in NS-LUAD. In other cases, the magnitude

of synergy differed across exposures: synergy for mutation of *TP53* in *EGFR*-mutant tumors

was stronger in ES- than in NS-LUAD. The higher frequency of synergy and lower frequency of

antagonism in ES-LUAD suggests a more navigable adaptive landscape—characterized by a

higher frequency of beneficial mutational combinations and fewer evolutionary barriers—than in

NS-LUAD. This increased accessibility enables ES-LUADs to develop along a greater diversity

of evolutionary trajectories, contributing to increased interpatient heterogeneity. Moreover, it

may also help to explain why tobacco smoke elevates the risk of lung cancer to such an

exceptional degree. In NS-LUAD, widespread antagonistic epistasis may reflect a greater

fitness cost of mutation accumulation, possibly due to a more immunologically reactive tumor

microenvironment [12,48,58,94] or due to a lower tolerance for mutation accumulation in the

non-dysregulated lung tissue cells of never-smokers. Taken together, these findings reveal that

environmental exposures modulate selective epistasis within a cancer type, suggesting the

importance of exposure history to predicting tumor evolutionary trajectories and to guiding

epistasis-informed treatment strategies.

It has often been assumed that the adaptive landscape of cancer can be adequately

described without epistasis [6,95–99] or with only pairwise epistasis [31,32,100]. However, our

results demonstrate that not only pairwise but also higher-order epistasis—particularly sub-

additive and emergent epistasis—play a crucial role in modulating the adaptive value of cancer

drivers. For instance, in ES-LUAD, selection for *APC* mutations increased by an order of

magnitude in an *KEAP1-STK11*-mutant genotype, despite showing no significant pairwise

synergy with either mutation alone. This emergent epistasis, arising only in the triadic context,

would be missed by analyses restricted to pairwise comparisons, necessitating consideration of

multi-mutant genotypes in both computational and experimental studies. Moreover, almost all

significant pairwise synergistic effects combined sub-additively in triads, rapidly reaching a

plateau of synergistic benefit. This pattern aligns with the diminishing-returns epistasis observed

in asexual populations [44–47]. Overall, higher-order synergistic epistasis biases tumors toward

some evolutionary trajectories, while higher-order antagonism prunes most trajectories. Both

make tumor evolution more predictable. Thus, incorporation of higher-order epistasis into

models of tumorigenesis is essential to understanding and guiding the evolutionary paths that

tumors take under therapy.

Current NSCLC treatment guidelines for EGFR TKIs do not account for co-occurring

mutations. However, our findings suggest that concurrent mutations could serve as biomarkers

of treatment response. Antagonistic epistasis between *EGFR* mutations and mutations in *KRAS*

or *KEAP1* implies that the efficacy of EGFR TKIs may be reduced in tumors harboring these

concurrent alterations. This prediction aligns with clinical observations: *KRAS* mutations

decrease survival in patients with *EGFR*-mutant LUAD patients treated with EGFR TKIs [101],

and *KEAP1* mutations reduced EGFR TKI efficacy in *EGFR*-mutant lung cancer [102,103]. More

broadly, antagonistic epistasis is a signal of synthetic lethality [32]. Our results also demonstrate

mutual negative selection between *BRCA2* and *SMAD4* loss-of-function mutations—suggesting

that pharmacological inhibition of *SMAD4* could be therapeutically beneficial in *BRCA2*-mutant

tumors, in analogy to usage of PARP inhibitors in *BRCA1/2* deficient tumors. In general,

selective epistasis can clarify the frequent heterogeneity of response to targeted therapies

[37,104], and can improve stratification of responders and nonresponders to targeted therapy in

clinical trials [105,106], enhancing trial power and therapeutic precision.

Because *TP53* mutations synergize with mutations of *EGFR*, we would expect that in

tumors with concurrent *EGFR* and *TP53* mutations, EGFR TKIs would be especially effective

because they would both eliminate the baseline effect of the *EGFR* mutation and diminish the

effect of the concurrent *TP53* mutation that is mediated through the *EGFR* mutation. However,

TP53 mutations are consistently associated with resistance to EGFR TKIs in *EGFR*-mutant

LUAD [107–109] and NSCLC [110–114]. This contradiction may be explained by the pleiotropic

effects of *TP53* loss, including increased somatic nucleotide mutation rates [109], widespread

copy-number alterations [107,115], and dysregulation of gene expression programs linked to

resistance [116] and progression [115]. These pleiotropic effects provide greater genetic

variation that facilitates the evolution of resistance independently of the selective synergy

between *TP53* loss and *EGFR* mutation.

Previous computational studies of pairwise interactions have demonstrated asymmetries

in the magnitude of pairwise epistasis [cf. 30]. Our analysis has revealed not only epistasis of

asymmetric magnitudes, but also asymmetric signs of selection, in which one ordering of the

mutations is advantageous while the reverse is disadvantageous. These directional

asymmetries likely arise from cellular or stable transcriptional transitions initiated by the primary

mutation that unidirectionally alter the selective context for subsequent alterations—an effect not

captured by conventional analyses of mutual exclusivity or co-occurrence. Indeed, the presence

of such asymmetries of epistasis are evident in established genetic models of tumor evolution,

such as the *APC → KRAS → TP53* mutation sequence in colorectal cancer [117,118], whose

strongly preferential ordering requires directional epistasis. Asymmetric epistasis has also been

supported by genetic and computational genomic studies of *TP53* and *CCNE1* mutations [32],

TP53 and *RB1* mutations [30], and *JAK2* and *TET2* mutations [119]. The presence of directional

epistasis implies preferential orders of mutation acquisition that offer new opportunities to

predict and potentially redirect cancer evolution.

This study was subject to several limitations. First, its retrospective nature introduces the

possibility that covariates of smoking behavior—such as comorbid health states, environmental

exposures, inherited genetics, and sex biases—may contribute to the observed differences in

the adaptive landscapes of ES-LUAD and NS-LUAD. However, none of these factors are known

to exert a physiological impact on the lung that is comparable to smoking tobacco. Accordingly,

the selective differences we report are most parsimoniously attributed to the direct and

downstream effects of smoking. Second, smoking status in our study was primarily inferred from

mutational signatures in tumor sequences, which can be susceptible to misclassification.

However, this method has been validated for classification of smoking status in NSCLC and is

believed to outperform self-reported smoking histories [21,25,120]; self-reported status has

been found to be unreliable [121,122] and often does not account for the risk conferred by

secondhand smoke [123]. Furthermore, because our approach explicitly quantifies the smoking-

associated mutational burden in the lung periphery—where LUAD develops—it also implicitly

identifies patients who have experienced significant injury from smoking in the regions where

LUAD originates. Finally, our analysis was restricted to somatic SNVs and did not consider

indels, copy number alterations (CNAs), or other structural variants (SVs). CNAs are common in

cancer [124] and pairwise selective epistasis has been indicated between SNVs and CNAs and

other SVs [125,126] or between SVs [37,87,127]. Unfortunately, accurate background mutation

rates are currently unavailable for SVs. As such rates become calculable, integrated analysis of

SNVs and SVs will reveal new epistatic relationships and clarify the somatic adaptive landscape

of cancer.

Overall, we have demonstrated that the adaptive landscapes of ES- and NS-LUAD diverge

substantially, implicating tobacco smoking not only as a mutagen but also as a modulator of

somatic selection. Our analysis also shows that symmetric and asymmetric selective epistasis

among driver mutations is both substantial and pervasive in lung adenocarcinoma and

generates the distinct evolutionary trajectories of ES-LUAD and NS-LUAD. Furthermore, we

provide a systematic characterization of higher-order selective epistasis in LUAD, revealing its

underrecognized role in shaping tumor evolution. Together, these results show that the

contribution of mutations to tumor growth is profoundly context-dependent—modulated by both

tumor somatic genetics and the environmental exposures it has experienced. This insight

illuminates new opportunities for precision oncology through the integration of somatic genotype

and exposure history into treatment strategies. Continued investigation into epistatic,

environmental, and even gene-by-gene-by-environment interactions in lung adenocarcinoma

and other cancers will be essential for the advancement of our understanding of cancer

evolution and the improvement of patient-specific therapeutic strategies.

Methods

Design overview

The aim of this study was to test for differences in background mutation rates, subsequent

somatic selection, and gene-by-gene epistatic effects between ES- and NS-LUAD. To do so, we

aggregated mutation data of 2,097 LUAD tumor samples from multiple institutional sources. We

classified tumors as either ES- or NS-LUAD either by detection of the smoking-associated

mutational signature SBS4 in exome or genome sequences or, in the case of targeted panel

sequence data, by relying on the patient's self-reported smoking history. We applied established

approaches to estimate background mutation rates, scaled selection coefficients, and pairwise

and higher-order selective epistasis within the ES- and NS-LUAD cohorts. To assess the

differential transcriptional effects of driver mutations, we compared ES- and NS-LUAD tumors

with and without specific mutations. Differences in selection were conservatively evaluated

based on non-overlapping 95% confidence intervals.

Data sources

We sourced clinicogenomic data from (i) eight published datasets hosted by cBioPortal

[85,128–135], (ii) The Cancer Genome Atlas dataset hosted on Genomic Data Commons [136],

and (iii) 108 samples sequenced at Yale University [137]. All datasets reported sample sizes,

sequencing methods, and clinical covariates, and were incorporated into our analysis

accordingly (**Supplementary Table 1**). We only included samples from primary LUAD tumors,

and we did not include any datasets from studies that preselected patients based on their

mutational profiles. When multiple samples from a single patient were available, one was

selected at random. In cases where patients appeared in more than one dataset, we retained

the sample with the highest-quality sequencing and most complete clinical annotations.

For consistency across datasets, we standardized all variant genomic coordinates to the

hg19 reference genome using the liftOver function of rtracklayer v1.62.0 [138]. We used

cancereffectsizeR v2.10.1 to remove variants that failed liftover, were incorrectly classified as

single-nucleotide variants (SNVs), or did not meet quality criteria for downstream analysis. For

panel-sequenced samples, we inferred the genomic regions covered by each panel from the

gene list provided for each panel using the corresponding hg19 coordinates.

Mutational signature attributions and smoking-status classification

We performed mutational signature deconvolution for tumor samples with whole-exome

or whole-genome sequence data, but not for panel-sequenced samples, due to the low reliability

of signature attribution when few silent somatic variants are present. We conducted signature

attribution using cancereffectsizeR with MutationalPatterns v3.12.0 [139], applying COSMIC

v3.2 single-base substitution signatures [26,140]. We excluded COSMIC signatures that were

unlikely to have been observed in lung adenocarcinomas [140], and we excluded treatment-

associated signatures when deconvolving samples known to be treatment-naive. Samples in

which the smoking-associated SBS4 signature contributed more than the MutationalPatterns

default threshold of 5% of the total signature burden were classified as ever-smoker ES-LUAD,

and otherwise were classified as never-smoker NS-LUAD. Zhang et al. detected no signature of

SBS4 in the samples used in their study; therefore we directly classified these samples as NS-

LUAD. As in Rosenthal et al. [141], we excluded samples with ≤50 single-nucleotide variants in

whole-exome data from classification due to insufficient mutation burden for reliable signature

deconvolution. We compared signature-based smoking assignments to available clinical

annotation of smoking status across datasets, validating that there was 67–90% concordance.

Discrepancies were likely due to underreporting of smoking, second-hand smoke exposure,

minimal mutational consequences in light smokers, or misclassification in clinical records. The

selection and epistasis results produced from either classification method were 99.4%

consistent with each other. We classified panel-sequenced samples according to clinical

annotations of smoking status when available; we excluded samples annotated as exposed to

"former light" smoking from all smoking-stratified analyses due to ambiguity in exposure and

effect.

Estimation of oncogenic mutation rates

We estimated gene-specific oncogenic mutation rates separately for the ES- and NS-

LUAD cohorts following the approach of Alfaro-Murillo and Townsend [28]. The oncogenic

mutation rate for a gene was defined as the neutral rate at which oncogenic variants occur

within the gene. We defined oncogenic variants as variants that were observed at least once in

any tumor sample in the dataset. The background oncogenic mutation rate of each gene was

computed as the sum of the background rates of all oncogenic variants within the gene. For a

specific variant derived from a trinucleotide mutation type m (*e.g.* CCT → CAT) at a site i within

gene g in a tumor sample cohort c, the somatic background mutation rate

768
$$\mu_{i,g,c}^m = \mu_{g,c} \cdot \mathrm{Prob}\left\{T_m|c\right\} \cdot \mathrm{Prob}\left\{S_{i,m_{\mathrm{WT}},g}\right\},$$

where $\mu_{g,c}$ is the background mutation rate of the entire gene g (the rate at which variants occur

in g) within cohort c, T_m is the event that an arising variant is of trinucleotide mutation type m,

and $S_{i,m_{\mathrm{WT}},g}$ is the event that the variant occurs at site i among all other trinucleotide sequences

that match the wild-type (WT) allele of m (m_{WT}, *e.g.* CCT), within g.

The probability of the variant being of type m given a cohort c, $\mathrm{Prob}\left\{T_m|c\right\}$, is computed

as the proportional frequency of m within cohort c, equivalent to the ratio of the number of

variants of type m in cohort c, n_m^c, to the total number of variants in c, N_c. Assuming an equal

probability of such a variant occurring at all possible sites within g, the probability of such a

variant occurring at site i, $\mathrm{Prob}\left\{S_{i,m_{\mathrm{WT}},g}\right\}$, is the inverse of the number of sites within g with

the same trinucleotide sequence as m_{WT}, $S_{m_{\mathrm{WT}}}^g$. Therefore

779
$$\mu_{i,g,c}^m = \frac{\mu_{g,c} \cdot n_m^c}{S_{m_{\mathrm{WT}}}^g \cdot N_c}.$$

780   The background oncogenic mutation rate of gene g is the sum of the background rates of all
781   oncogenic variants within g:

782
$$\hat{\mu}_{g,c} = \sum_{(i,m) \in \mathcal{S}_g} \mu_{i,g,c}^m$$

where $\mathcal{S}_g$ denotes the set of pairs of site indices and mutation types identifying the oncogenic

variants in g. This approach enables estimation of mutation rates that are both gene-specific

and context-dependent, accounting for trinucleotide mutational biases across the genome. The

oncogenic mutation rates of the genes were consistently lower than the mutation rates of the

genes.

Finally, the oncogenic mutation rates per somatic genotypes were computed by rescaling

the per-gene background rates using the tumor mutation burden (TMB) observed in each

genotype. Specifically, for a given cohort c, the mutation rate for a jump between the somatic

genotype **x** the somatic genotype **y** is given by

792
$$\mu_{\mathbf{x}\to\mathbf{y},c} = \frac{\mathrm{TMB}_{\{\mathbf{x},\mathbf{y}\}}}{\mathrm{TMB}_{\mathrm{all}}}\hat{\mu}_{g,c},$$

where $\mathrm{TMB}_{\{\mathbf{x},\mathbf{y}\},c}$ is the average TMB in tumors with the somatic genotypes **x** or **y**, $\mathrm{TMB}_{\mathrm{all}}$ is

average TMB for all tumors, and g is the gene that is mutated in the somatic genotype **y** but not

in **x** (the mutation gained by the jump). In cases where one genotype has no samples, the

average is taken only from the observed partner. If both endpoints have zero samples, the rate

defaults to the baseline per-gene rate $\mu_{\mathbf{x}\to\mathbf{y},c} = \hat{\mu}_{g,c}$, avoiding spurious rescaling in unsupported

genotype classes. We considered TMB only from our largest dataset (TCGA) because after

comparing the datasets with the Kolmogorov–Smirnov test, all of them proved to come from

different distributions.

Estimation of scaled selection coefficients and selective epistasis

To quantify the growth and survival advantages conferred by somatic SNVs in LUAD, we

calculated scaled selection coefficients for recurrent mutations in a curated set of 1,200 genes

[136,142–145]. These coefficients roughly equate to the ratio of the fixation rate for the

oncogenic mutant (the rate at which the mutant is generated and subsequently selected to a

high enough frequency to be observed as a somatic variant within the cancer cell population) to

its background mutation rate. Precisely, we computed the fixation rate by analytical

maximization of the binomial likelihood of observing at least one non-silent SNV in the gene

given the number of samples with and without a non-silent SNV in the gene. Scaled selection

coefficients for each mutated gene were estimated separately for ES- and NS-LUAD, as well as

for a combined cohort. Within each group, we included samples with whole-exome, whole-

genome, or targeted gene panel sequences (TGS). Results were consistent when excluding

TGS samples, indicating robustness to sequencing strategy. Inclusion of TGS samples provided

greater statistical power to detect significant differences in selection between cohorts. Results

were also robust to cohort-specific normalization: applying median-scaling of coefficients prior to

between-cohort comparison yielded consistent outcomes.

From the curated set of 1,200 genes, we selected 21 for detailed analysis of selective

epistasis based on strong prior evidence as LUAD drivers [136,143] and their elevated average

strength of selection across somatic genotypes. We quantified epistasis by calculating scaled

selection coefficients conditional on the somatic genotype as in Alfaro-Murillo and Townsend

[28]: for each pair or triplet of genes, we calculated the fixation rate of mutations in one gene

conditional on the presence or absence of somatic variants in the others. We performed this

calculation by maximizing the multinomial likelihood for the number of samples with each

somatic genotype using PyMC [146]. The probability of a tumor being in a specific somatic

genotype during the somatic evolutionary trajectory from organogenesis to biopsy was

computed via a continuous-time Markov chain model [28]. We computed asymmetric 95%

confidence intervals using Wilks' theorem [147]. To obtain scaled selection coefficients for each

mutation occurring within each somatic genotype, we divided the respective Poisson-corrected

fixation rate [148] by the background oncogenic mutation rate of the gene.

To systematically assess selective epistasis in both the ever-smoker and the never-

smoker groups, we evaluated 3,080 models representing all possible two- and three-gene

combinations from the 21 LUAD driver genes. For each model, somatic genotypes were defined

exclusively by the variant status (mutant or wildtype) of the genes included in each combination.

To ensure complete genotype classification, we only included panel-sequenced samples in a

given model if the sequencing panel covered all of the genes under analysis.

Classification of synergistic and antagonistic selective epistasis

To classify a mutant somatic genotype as interacting synergistically or antagonistically with

a certain mutation, we compared the scaled selection coefficients for a mutation when it

occurred in the mutant somatic genotype to the scaled selection coefficients obtained for the

same mutation in a somatic genotype lacking mutations in the genes included in the model. If

the scaled selection coefficient for a mutation in gene A when it occurred in a wild-type

genotype ($\emptyset \rightarrow A$) was less than the scaled selection coefficient for a mutation in gene A when it

occurred in a somatic genotype with a mutation in gene B ($B \rightarrow A + B$), then we deemed the

epistatic effect of mutant gene B on mutant gene A to be synergistic. If the scaled selection

coefficient for $\emptyset \rightarrow A$ was greater than the scaled selection coefficient for $B \rightarrow A + B$, then we

deemed the epistatic effect of mutant gene B on mutant gene A to be antagonistic. We further

distinguished between two scales of antagonism: antagonistic sign epistasis when there was a

reversal of the direction of selection—from positive (coefficient $\gamma > 1$) to negative selection

(coefficient $\gamma < 1$)—in the antagonistic genotype, and antagonistic magnitude epistasis when

there was a reduction in strength of selection but selection remained positive in the antagonistic

genotype.

Differential expression analysis

To evaluate the transcriptional impact of key driver mutations across smoking-stratified

LUAD tumors, we performed differential expression analysis using RNA-seq data from the

TCGA-LUAD project. We obtained unstranded raw read counts for 59 normal lung and 539

primary LUAD tumor samples using TCGAbiolinks v2.30.0. We excluded formalin-fixed paraffin-

embedded samples and tumors which received neoadjuvant treatment to minimize technical

and biological confounders. We retained solid-tissue normal samples as controls. We excluded

genes with fewer than 10 total counts across all samples. For each sample, we assigned

smoking status based on SBS4 signature weight from the matched MAF. The genes *EGFR*,

KRAS, and *KEAP1* were considered mutated if at least one non-silent SNV, insertion, or

deletion was present in the gene (excluding intronic regions). After this filtering and annotation,

we validated that a principal-component analysis revealed no evidence of batch effects,

confirming the suitability of the dataset for differential expression analysis.

We performed differential expression analysis with DESeq2 v1.42.1 [149] using two

designs. Differential transcriptional effects of *EGFR* mutations in ES- and NS-LUAD were

quantified as

$$K_i \; = \; \beta_0 + \beta_1 x_t + \beta_2 x_s + \beta_3 x_e + \beta_4 x_r + \beta_5 (x_s * x_e) + \beta_6 (x_s * x_r),$$

where K_i is the count for gene i, and x_t, x_s, x_e, x_r are indicator variables for the sample type,

smoking status, *EGFR* mutation status, and *KRAS* mutation status. The differential effect in

question was measured by the interaction term between smoking status and *EGFR* allele

status. The *EGFR*-focused-design accounts for the fact that most *EGFR*-wildtype LUADs are

KRAS-mutant. Differential transcriptional effects of *KEAP1* mutations in ES- and NS-LUAD,

were quantified as

$$K_i \; = \; \beta_0 + \beta_1 x_t + \beta_2 x_s + \beta_3 x_k + \beta_4 (x_s * x_k),$$

where x_k is the *KEAP1* mutation status.

We performed gene-set enrichment analysis with fgsea v1.28.0 using the MSigDB

Hallmark gene sets. We ranked transcripts by signed *t*-statistics to capture both degree of

significance and magnitude of fold change. We performed gene set variation analysis with

GSVA v1.50.5 using the MSigDB canonical gene sets.

Statistical Analysis

Differences in selection coefficients were assessed conservatively by non-overlapping

95% confidence intervals for each estimate. Differential gene expression was analyzed using

two-tailed Wald tests, and statistical significance was determined based on Benjamini-

Hochberg–adjusted $P < 0.05$. Gene set enrichment analyses were likewise considered

significant at a Benjamini-Hochberg–adjusted $P < 0.05$. Analysis was performed in Python

(3.9.5) and R (4.3.0).

Data and code availability

All code, data, and instructions necessary to reproduce the results and figures in the current
study is available in a public Zenodo repository, https://zenodo.org/records/16380037 [150] with
no access restrictions.

Acknowledgements

We thank Jeff Mandell for his advice on mutational signature attribution and mutation rate

estimation, and Elizabeth Perry for assistance with gene expression analysis.

Authors' contributions

JPT, JAA-M, and KD conceptualized the project. KD curated the data and performed the formal

analysis. KD and JAA-M wrote the software. KD, JAA-M, and JPT contributed to the

methodology and investigation. KD drafted the original manuscript and JPT and JAA-M

reviewed and edited the manuscript. All authors read and approved the final manuscripts.

Competing interests

The authors declare that they have no competing interests.

910 **Materials and correspondence**

911 Correspondence and material requests should be addressed to Jeffrey Townsend.

912 **Funding**

References

1. HHS. How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-attributable Disease. U.S. Department of Health and Human Services; 2010.

2. Alexandrov LB, Ju YS, Haase K, Van Loo P, Martincorena I, Nik-Zainal S, et al. Mutational signatures associated with tobacco smoking in human cancer. Science. 2016;354: 618–622.

3. Pfeifer GP, Denissenko MF, Olivier M, Tretyakova N, Hecht SS, Hainaut P. Tobacco smoke carcinogens, DNA damage and p53 mutations in smoking-associated cancers. Oncogene. 2002;21: 7435–7451.

4. Yoshida K, Gowers KHC, Lee-Six H, Chandrasekharan DP, Coorens T, Maughan EF, et al. Tobacco exposure and somatic mutations in normal human bronchial epithelium. Nature. 2020;578: 266.

5. Dasari K, Somarelli JA, Kumar S, Townsend JP. The somatic molecular evolution of cancer: Mutation, selection, and epistasis. Prog Biophys Mol Biol. 2021;165: 56–65.

6. Martincorena I, Raine KM, Gerstung M, Dawson KJ, Haase K, Van Loo P, et al. Universal patterns of selection in cancer and somatic tissues. Cell. 2017;171: 1029–1041.e21.

7. Cannataro VL, Kudalkar S, Dasari K, Gaffney SG, Lazowski HM, Jackson LK, et al. APOBEC mutagenesis and selection for NFE2L2 contribute to the origin of lung squamous-cell carcinoma. Lung Cancer. 2022;171: 34–41.

8. Tan C, Mandell JD, Dasari K, Cannataro VL, Alfaro-Murillo JA, Townsend JP. Heavy mutagenesis by tobacco leads to lung adenocarcinoma tumors with KRAS G12 mutations other than G12D, leading KRAS G12D tumors-on average-to exhibit a lower mutation burden. Lung Cancer. 2022;166: 265–269.

9. Mochizuki A, Shiraishi K, Honda T, Higashiyama RI, Sunami K, Matsuda M, et al. Passive smoking-induced Mutagenesis as a promoter of lung carcinogenesis. J Thorac Oncol. 2024;19: 984–994.

10. Lee J, Taneja V, Vassallo R. Cigarette Smoking and Inflammation: Cellular and Molecular Mechanisms. J Dent Res. 2012;91: 142.

11. van der Vaart H, Timens W, Ten Hacken NHT. Acute effects of cigarette smoke on inflammation and oxidative stress: a review. Thorax. 2004;59: 713–721.

12. Li X, Li J, Wu P, Zhou L, Lu B, Ying K, et al. Smoker and non-smoker lung adenocarcinoma is characterized by distinct tumor immune microenvironments. Oncoimmunology. 2018;7: e1494677.

13. Rodin SN, Rodin AS. Human lung cancer and p53: the interplay between mutagenesis and selection. Proc Natl Acad Sci U S A. 2000;97: 12244–12249.

14. Laconi E, Marongiu F, DeGregori J. Cancer as a disease of old age: changing mutational and microenvironmental landscapes. Br J Cancer. 2020;122: 943–952.

15. Marusyk A, DeGregori J. Declining cellular fitness with age promotes cancer initiation by selecting for adaptive oncogenic mutations. Biochim Biophys Acta. 2008;1785: 1–11.

16. Lopez-Bigas N, Gonzalez-Perez A. Are carcinogens direct mutagens? Nat Genet. 2020;52: 1137–1138.

17. Stratton MR, Humphreys L, Alexandrov LB, Balmain A, Brennan P, Campbell PJ, et al. Implementing mutational epidemiology on a global scale: Lessons from mutographs. Cancer Discov. 2025;15: 22–27.

18. Okamoto T, Suzuki Y, Fujishita T, Kitahara H, Shimamatsu S, Kohno M, et al. The prognostic impact of the amount of tobacco smoking in non-small cell lung cancer--differences between adenocarcinoma and squamous cell carcinoma. Lung Cancer. 2014;85: 125–130.

19. Kenfield SA, Wei EK, Stampfer MJ, Rosner BA, Colditz GA. Comparison of Aspects of Smoking Among Four Histologic Types of Lung Cancer. Tob Control. 2008;17: 198.

20. National Center for Health Statistics. Table 17, Current cigarette smoking among adults aged 18 and over, by sex, race, and age: United States, selected years 1965–2018. National Center for Health Statistics (US); 2021.

21. Devarakonda S, Li Y, Rodrigues FM, Sankararaman S, Kadara H, Goparaju C, et al. Genomic Profiling of Lung Adenocarcinoma in Never-Smokers. J Clin Oncol. 2021 [cited 25 Apr 2024]. doi:10.1200/JCO.21.01691

22. Moorthi S, Paguirigan A, Itagi P, Ko M, Pettinger M, Hoge AC, et al. The genomic landscape of lung cancer in never-smokers from the Women's Health Initiative. JCI Insight. 2024;9. doi:10.1172/jci.insight.174643

23. Wang X, Ricciuti B, Nguyen T, Li X, Rabin MS, Awad MM, et al. Association between smoking history and tumor mutation burden in advanced non-small cell lung cancer. Cancer Res. 2021;81: 2566–2573.

24. Govindan R, Ding L, Griffith M, Subramanian J, Dees ND, Kanchi KL, et al. Genomic landscape of non-small cell lung cancer in smokers and never-smokers. Cell. 2012;150: 1121–1134.

25. Díaz-Gay M, Zhang T, Hoang PH, Leduc C, Baine MK, Travis WD, et al. The mutagenic forces shaping the genomes of lung cancer in never smokers. Nature. 2025; 1–12.

26. Blokzijl F, Janssen R, van Boxtel R, Cuppen E. MutationalPatterns: comprehensive genome-wide analysis of mutational processes. Genome Med. 2018;10: 1–11.

27. Mandell JD, Cannataro VL, Townsend JP. Estimation of Neutral Mutation Rates and Quantification of Somatic Variant Selection Using cancereffectsizeR. Cancer Res. 2023;83: 500–505.

28. Alfaro-Murillo JA, Townsend JP. Pairwise and higher-order epistatic effects among somatic

cancer mutations across oncogenesis. Math Biosci. 2023;366: 109091.

29. Wang X, Fu AQ, McNerney ME, White KP. Widespread genetic epistasis among cancer genes. Nat Commun. 2014;5: 1–10.

30. Iranzo J, Gruenhagen G, Calle-Espinosa J, Koonin EV. Pervasive conditional selection of driver mutations and modular epistasis networks in cancer. Cell Rep. 2022;40: 111272.

31. Blair LM, Juan JM, Sebastian L, Tran VB, Nie W, Wall GD, et al. Oncogenic context shapes the fitness landscape of tumor suppression. Nat Commun. 2023;14: 1–19.

32. Mina M, Iyer A, Ciriello G. Epistasis and evolutionary dependencies in human cancers. Curr Opin Genet Dev. 2022;77: 101989.

33. Zhang Z, Yang Y, Zhou Y, Fang H, Yuan M, Sasser K, et al. A forward selection algorithm to identify mutually exclusive alterations in cancer studies. J Hum Genet. 2021;66: 509–518.

34. Fedrizzi T, Ciani Y, Lorenzin F, Cantore T, Gasperini P, Demichelis F. Fast mutual exclusivity algorithm nominates potential synthetic lethal gene pairs through brute force matrix product computations. Comput Struct Biotechnol J. 2021;19: 4394–4403.

35. Liu S, Liu J, Xie Y, Zhai T, Hinderer EW, Stromberg AJ, et al. MEScan: a powerful statistical framework for genome-scale mutual exclusivity analysis of cancer mutations. Bioinformatics. 2021;37: 1189–1197.

36. Klein MI, Cannataro VL, Townsend JP, Newman S, Stern DF, Zhao H. Identifying modules of cooperating cancer drivers. Mol Syst Biol. 2021 [cited 2 Jul 2024]. doi:10.15252/msb.20209810

37. Mina M, Raynaud F, Tavernari D, Battistello E, Sungalee S, Saghafinia S, et al. Conditional Selection of Genomic Alterations Dictates Cancer Evolution and Oncogenic Dependencies. Cancer Cell. 2017;32: 155–168.e6.

38. Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Global epistasis makes adaptation predictable despite sequence-level stochasticity. Science. 2014 [cited 24 Apr 2024]. doi:10.1126/science.1250939

39. Weinreich DM, Lan Y, Wylie CS, Heckendorn RB. Should evolutionary geneticists worry about higher-order epistasis? Curr Opin Genet Dev. 2013;23: 700–707.

40. Sailer ZR, Harms MJ. High-order epistasis shapes evolutionary trajectories. PLoS Comput Biol. 2017;13: e1005541.

41. Weinreich DM, Lan Y, Jaffe J, Heckendorn RB. The influence of higher-order epistasis on biological fitness landscape topography. J Stat Phys. 2018;172: 208–225.

42. Domingo J, Diss G, Lehner B. Pairwise and higher-order genetic interactions during the evolution of a tRNA. Nature. 2018;558: 117–121.

43. Papkou A, Garcia-Pastor L, Escudero JA, Wagner A. A rugged yet easily navigable fitness landscape. Science. 2023;382: eadh3860.

44. Wei X, Zhang J. Patterns and Mechanisms of Diminishing Returns from Beneficial Mutations. Mol Biol Evol. 2019;36: 1008–1021.

45. Aggeli D, Li Y, Sherlock G. Changes in the distribution of fitness effects and adaptive mutational spectra following a single first step towards adaptation. Nat Commun. 2021;12: 1–14.

46. Chou H-H, Chiu H-C, Delaney NF, Segrè D, Marx CJ. Diminishing returns epistasis among beneficial mutations decelerates adaptation. Science. 2011;332: 1190–1192.

47. Wünsche A, Dinh DM, Satterwhite RS, Arenas CD, Stoebel DM, Cooper TF. Diminishing-returns epistasis decreases adaptability along an evolutionary trajectory. Nature Ecology & Evolution. 2017;1: 1–6.

48. Sui Q, Liang J, Hu Z, Chen Z, Bi G, Huang Y, et al. Genetic and microenvironmental differences in non-smoking lung adenocarcinoma patients compared with smoking patients. Transl Lung Cancer Res. 2020;9: 1407–1421.

49. Frank R, Scheffler M, Merkelbach-Bruse S, Ihle MA, Kron A, Rauer M, et al. Clinical and Pathological Characteristics of KEAP1- and NFE2L2-Mutated Non-Small Cell Lung Carcinoma (NSCLC). Clin Cancer Res. 2018;24. doi:10.1158/1078-0432.CCR-17-3416

50. Nadal E, Palmero R, Muñoz-Pinedo C. Mutations in the Antioxidant KEAP1/NRF2 Pathway Define an Aggressive Subset of NSCLC Resistant to Conventional Treatments. J Thorac Oncol. 2019;14. doi:10.1016/j.jtho.2019.08.005

51. Subramanian J, Govindan R. Lung Cancer in Never Smokers: A Review. J Clin Oncol. 2016 [cited 25 Apr 2024]. doi:10.1200/JCO.2006.06.8015

52. Scott J, Marusyk A. Somatic clonal evolution: A selection-centric perspective. Biochim Biophys Acta. 2017;1867: 139–150.

53. Fleenor CJ, Marusyk A, DeGregori J. Ionizing radiation and hematopoietic malignancies: altering the adaptive landscape. Cell Cycle. 2010;9: 3005–3011.

54. Hill W, Lim EL, Weeden CE, Lee C, Augustine M, Chen K, et al. Lung adenocarcinoma promotion by air pollutants. Nature. 2023;616: 159–167.

55. Bossé Y, Postma DS, Sin DD, Lamontagne M, Couture C, Gaudreault N, et al. Molecular signature of smoking in human lung tissues. Cancer Res. 2012;72: 3753–3763.

56. Pintarelli G, Noci S, Maspero D, Pettinicchio A, Dugo M, De Cecco L, et al. Cigarette smoke alters the transcriptome of non-involved lung tissue in lung adenocarcinoma patients. Sci Rep. 2019;9: 13039.

57. Ramirez JM, Ribeiro R, Soldatkina O, Moraes A, García-Pérez R, Oliveros W, et al. The molecular impact of cigarette smoking resembles aging across tissues. Genome Medicine. 2025;17: 1–23.

58. Luo W, Zeng Z, Jin Y, Yang L, Fan T, Wang Z, et al. Distinct immune microenvironment of lung adenocarcinoma in never-smokers from smokers. Cell Rep Med. 2023;4: 101078.

59. Price SN, Land SR, Pebley K, Fahey MC, Palmer AM, McCall MH, et al. Tobacco assessment in actively accruing National Cancer Institute clinical trials network trials. Nicotine Tob Res. 2025; ntaf071.

60. Peters EN, Warren GW, Sloan JA, Marshall JR. Tobacco assessment in completed lung cancer treatment trials: Pooling Project: Smoking and Lung Cancer. Cancer. 2016;122: 3260–3262.

61. Kim IA, Lee JS, Kim HJ, Kim WS, Lee KY. Cumulative smoking dose affects the clinical outcomes of EGFR-mutated lung adenocarcinoma patients treated with EGFR-TKIs: a retrospective study. BMC Cancer. 2018;18: 768.

62. Kim MH, Kim HR, Cho BC, Bae MK, Kim EY, Lee CY, et al. Impact of cigarette smoking on response to epidermal growth factor receptor (EGFR)-tyrosine kinase inhibitors in lung adenocarcinoma with activating EGFR mutations. Lung Cancer. 2014;84: 196–202.

63. Zhang P, Nie X, Bie Z, Li L. Impact of heavy smoking on the benefits from first-line EGFR-TKI therapy in patients with advanced lung adenocarcinoma. Medicine . 2018;97: e0006.

64. Igawa S, Sasaki J, Otani S, Ishihara M, Takakura A, Katagiri M, et al. Impact of smoking history on the efficacy of gefitinib in patients with non-small cell lung cancer harboring activating epidermal growth factor receptor mutations. Oncology. 2015;89: 275–280.

65. van de Haar J, Canisius S, Yu MK, Voest EE, Wessels LFA, Ideker T. Identifying epistasis in cancer genomes: A delicate affair. Cell. 2019;177: 1375–1383.

66. Romero R, Sayin VI, Davidson SM, Bauer MR, Singh SX, LeBoeuf SE, et al. Keap1 loss promotes Kras-driven lung cancer and results in dependence on glutaminolysis. Nat Med. 2017;23: 1362–1368.

67. Best SA, Ding S, Kersbergen A, Dong X, Song J-Y, Xie Y, et al. Distinct initiating events underpin the immune and metabolic heterogeneity of KRAS-mutant lung adenocarcinoma. Nat Commun. 2019;10: 4190.

68. Galan-Cobo A, Sitthideatphaiboon P, Qu X, Poteete A, Pisegna MA, Tong P, et al. LKB1 and KEAP1/NRF2 pathways cooperatively promote metabolic reprogramming with enhanced glutamine dependence in KRAS-mutant lung adenocarcinoma. Cancer Res. 2019;79: 3251–3267.

69. Ricciuti B, Arbour KC, Lin JJ, Vajdi A, Vokes N, Hong L, et al. Diminished Efficacy of Programmed Death-(Ligand)1 Inhibition in STK11- and KEAP1-Mutant Lung Adenocarcinoma Is Affected by KRAS Mutation Status. J Thorac Oncol. 2022;17: 399–410.

70. Kwack WG, Shin SY, Lee SH. Primary resistance to immune checkpoint blockade in an STK11/TP53/KRAS-mutant lung adenocarcinoma with high PD-L1 expression. Onco Targets Ther. 2020;13: 8901–8905.

71. Arbour KC, Jordan E, Kim HR, Dienstag J, Yu HA, Sanchez-Vega F, et al. Effects of Co-occurring Genomic Alterations on Outcomes in Patients with KRAS-Mutant Non–Small Cell Lung Cancer. Clin Cancer Res. 2018;24: 334–340.

72. Proulx-Rocray F, Routy B, Nassabein R, Belkaid W, Tran-Thanh D, Malo J, et al. The

prognostic impact of KRAS, TP53, STK11 and KEAP1 mutations and their influence on the NLR in NSCLC patients treated with immunotherapy. Cancer Treat Res Commun. 2023;37: 100767.

73. Baptiste Oudart J, Garinet S, Leger C, Barlesi F, Mazières J, Jeannin G, et al. STK11/LKB1 alterations worsen the poor prognosis of KRAS mutated early-stage non-squamous non-small cell lung carcinoma, results based on the phase 2 IFCT TASTE trial. Lung Cancer. 2024;190: 107508.

74. Bange E, Marmarelis ME, Hwang W-T, Yang Y-X, Thompson JC, Rosenbaum J, et al. Impact of KRAS and TP53 co-mutations on outcomes after first-line systemic therapy among patients with STK11-mutated advanced non-small-cell lung cancer. JCO Precis Oncol. 2019;3: 1–11.

75. West HJ, McCleland M, Cappuzzo F, Reck M, Mok TS, Jotte RM, et al. Clinical efficacy of atezolizumab plus bevacizumab and chemotherapy in KRAS-mutated non-small cell lung cancer with STK11, KEAP1, or TP53 comutations: subgroup results from the phase III IMpower150 trial. J Immunother Cancer. 2022;10: e003027.

76. Aredo JV, Padda SK, Kunder CA, Han SS, Neal JW, Shrager JB, et al. Impact of KRAS mutation subtype and concurrent pathogenic mutations on non-small cell lung cancer outcomes. Lung Cancer. 2019;133: 144–150.

77. Wohlhieter CA, Richards AL, Uddin F, Hulton CH, Quintanal-Villalonga À, Martin A, et al. Concurrent mutations in STK11 and KEAP1 promote ferroptosis protection and SCD1 dependence in lung cancer. Cell Rep. 2020;33: 108444.

78. Julian C, Pal N, Gershon A, Evangelista M, Purkey H, Lambert P, et al. Overall survival in patients with advanced non-small cell lung cancer with KRAS G12C mutation with or without STK11 and/or KEAP1 mutations in a real-world setting. BMC Cancer. 2023;23. doi:10.1186/s12885-023-10778-6

79. Boeschen M, Kuhn CK, Wirtz H, Seyfarth H-J, Frille A, Lordick F, et al. Comparative bioinformatic analysis of KRAS, STK11 and KEAP1 (co-)mutations in non-small cell lung cancer with a special focus on KRAS G12C. Lung Cancer. 2023;184: 107361.

80. Bao Y, Zhang S, Zhang X, Pan Y, Yan Y, Wang N, et al. RBM10 Loss Promotes EGFR-Driven Lung Cancer and Confers Sensitivity to Spliceosome Inhibition. Cancer Res. 2023;83: 1490–1502.

81. Nanjo S, Wu W, Karachaliou N, Blakely CM, Suzuki J, Chou YT, et al. Deficiency of the splicing factor RBM10 limits EGFR inhibitor response in EGFR-mutant lung cancer. The Journal of clinical investigation. 2022;132. doi:10.1172/JCI145099

82. Gini B, Thomas N, Blakely CM. Impact of concurrent genomic alterations in epidermal growth factor receptor (EGFR)-mutated lung cancer. J Thorac Dis. 2020;12: 2883–2895.

83. Offin M, Chan JM, Tenet M, Rizvi HA, Shen R, Riely GJ, et al. Concurrent RB1 and TP53 alterations define a subset of EGFR-mutant lung cancers at risk for histologic transformation and inferior clinical outcomes. J Thorac Oncol. 2019;14: 1784–1793.

84. Pros E, Saigi M, Alameda D, Gomez-Mariano G, Martinez-Delgado B, Alburquerque-Bejar

JJ, et al. Genome-wide profiling of non-smoking-related lung cancer cells reveals common RB1 rearrangements associated with histopathologic transformation in EGFR-mutant tumors. Ann Oncol. 2020;31: 274–282.

85. Jamal-Hanjani M, Wilson GA, McGranahan N, Birkbak NJ, Watkins TBK, Veeriah S, et al. Tracking the Evolution of Non-Small-Cell Lung Cancer. N Engl J Med. 2017;376: 2109–2121.

86. Frankell AM, Dietzen M, Al Bakir M, Lim EL, Karasaki T, Ward S, et al. The evolution of lung cancer and impact of subclonal selection in TRACERx. Nature. 2023;616: 525–533.

87. Park S, Lehner B. Cancer type-dependent genetic interactions between cancer driver alterations indicate plasticity of epistasis across cell types. Mol Syst Biol. 2015;11: 824.

88. Guénolé A, Srivas R, Vreeken K, Wang ZZ, Wang S, Krogan NJ, et al. Dissection of DNA damage responses using multiconditional genetic interaction maps. Mol Cell. 2013;49: 346–358.

89. Harrison R, Papp B, Pál C, Oliver SG, Delneri D. Plasticity of genetic interactions in metabolic networks of yeast. Proc Natl Acad Sci U S A. 2007;104: 2307–2312.

90. Zhu C-T, Ingelmo P, Rand DM. G×G×E for lifespan in Drosophila: mitochondrial, nuclear, and dietary interactions that modify longevity. PLoS Genet. 2014;10: e1004354.

91. Shi X, Reinstadler B, Shah H, To T-L, Byrne K, Summer L, et al. Combinatorial GxGxE CRISPR screen identifies SLC25A39 in mitochondrial glutathione transport linking iron homeostasis to OXPHOS. Nat Commun. 2022;13: 2483.

92. Flynn KM, Cooper TF, Moore FB-G, Cooper VS. The environment affects epistatic interactions to alter the topology of an empirical fitness landscape. PLoS Genet. 2013;9: e1003426.

93. Costanzo M, Hou J, Messier V, Nelson J, Rahman M, VanderSluis B, et al. Environmental robustness of the global yeast genetic interaction network. Science. 2021;372. doi:10.1126/science.abf8424

94. Sun Y, Yang Q, Shen J, Wei T, Shen W, Zhang N, et al. The Effect of Smoking on the Immune Microenvironment and Immunogenicity and Its Relationship With the Prognosis of Immune Checkpoint Inhibitors in Non-small Cell Lung Cancer. Front Cell Dev Biol. 2021;9: 745859.

95. Boström M, Larsson E. Somatic mutation distribution across tumour cohorts provides a signal for positive selection in cancer. Nat Commun. 2022;13: 7023.

96. Bolton KL, Ptashkin RN, Gao T, Braunstein L, Devlin SM, Kelly D, et al. Cancer therapy shapes the fitness landscape of clonal hematopoiesis. Nat Genet. 2020;52: 1219–1226.

97. Salehi S, Kabeer F, Ceglia N, Andronescu M, Williams MJ, Campbell KR, et al. Clonal fitness inferred from time-series modelling of single-cell cancer genomes. Nature. 2021;595: 585–590.

98. Liggett LA, DeGregori J. Changing mutational and adaptive landscapes and the genesis of

cancer. Biochim Biophys Acta Rev Cancer. 2017;1867: 84–94.

99. Hsu T-K, Asmussen J, Koire A, Choi B-K, Gadhikar MA, Huh E, et al. A general calculus of fitness landscapes finds genes under selection in cancers. Genome Res. 2022;32: 916–929.

100. Misra N, Szczurek E, Vingron M. Inferring the paths of somatic evolution in cancer. Bioinformatics. 2014;30: 2456–2463.

101. Marchetti A, Milella M, Felicioni L, Cappuzzo F, Irtelli L, Del Grammastro M, et al. Clinical implications of KRAS mutations in lung cancer patients treated with tyrosine kinase inhibitors: an important role for mutations in minor clones. Neoplasia. 2009;11: 1084–1092.

102. Hellyer JA, Stehr H, Das M, Padda SK, Ramchandran K, Neal JW, et al. Impact of KEAP1/NFE2L2/CUL3 mutations on duration of response to EGFR tyrosine kinase inhibitors in EGFR mutated non-small cell lung cancer. Lung Cancer. 2019;134: 42–45.

103. Krall EB, Wang B, Munoz DM, Ilic N, Raghavan S, Niederst MJ, et al. KEAP1 loss modulates sensitivity to kinase targeted therapy in lung cancer. Elife. 2017;6. doi:10.7554/eLife.18970

104. Iyer G, Hanrahan AJ, Milowsky MI, Al-Ahmadie H, Scott SN, Janakiraman M, et al. Genome sequencing identifies a basis for everolimus sensitivity. Science. 2012;338: 221.

105. Weigelt B, Reis-Filho JS. Epistatic interactions and drug response. J Pathol. 2014;232: 255–263.

106. Wilkins JF, Cannataro VL, Shuch B, Townsend JP. Analysis of mutation, selection, and epistasis: an informed approach to cancer clinical trials. Oncotarget. 2018;9: 22243–22253.

107. Hobor S, Al Bakir M, Hiley CT, Skrzypski M, Frankell AM, Bakker B, et al. Mixed responses to targeted therapy driven by chromosomal instability through p53 dysfunction and genome doubling. Nat Commun. 2024;15: 4871.

108. Bria E, Pilotto S, Amato E, Fassan M, Novello S, Peretti U, et al. Molecular heterogeneity assessment by next-generation sequencing and response to gefitinib of EGFR mutant advanced lung adenocarcinoma. Oncotarget. 2015;6: 12783–12795.

109. Vokes NI, Chambers E, Nguyen T, Coolidge A, Lydon CA, Le X, et al. Concurrent TP53 Mutations Facilitate Resistance Evolution in EGFR-Mutant Lung Adenocarcinoma. J Thorac Oncol. 2022;17: 779–792.

110. Canale M, Petracci E, Delmonte A, Chiadini E, Dazzi C, Papi M, et al. Impact of TP53 Mutations on Outcome in EGFR-Mutated Patients Treated with First-Line Tyrosine Kinase Inhibitors. Clin Cancer Res. 2017;23: 2195–2202.

111. Ibusuki R, Iwama E, Shimauchi A, Tsutsumi H, Yoneshima Y, Tanaka K, et al. TP53 gain-of-function mutations promote osimertinib resistance via TNF-α-NF-κB signaling in EGFR-mutated lung cancer. NPJ Precis Oncol. 2024;8: 60.

112. Canale M, Petracci E, Delmonte A, Bronte G, Chiadini E, Ludovini V, et al. Concomitant TP53 Mutation Confers Worse Prognosis in EGFR-Mutated Non-Small Cell Lung Cancer

Patients Treated with TKIs. J Clin Med Res. 2020;9. doi:10.3390/jcm9041047

113.   Choudhury NJ, Lavery JA, Brown S, de Bruijn I, Jee J, Tran TN, et al. The GENIE BPC NSCLC cohort: A real-world repository integrating standardized clinical and genomic data for 1,846 patients with non-small cell lung cancer. Clin Cancer Res. 2023;29: 3418–3428.

114.   Fu J, Tong Y, Xu Z, Li Y, Zhao Y, Wang T, et al. Impact of TP53 mutations on EGFR-tyrosine kinase inhibitor efficacy and potential treatment strategy. Clin Lung Cancer. 2023;24: 29–39.

115.   Donehower LA, Soussi T, Korkut A, Liu Y, Schultz A, Cardenas M, et al. Integrated analysis of TP53 gene and pathway alterations in The Cancer Genome Atlas. Cell Rep. 2019;28: 1370–1384.e5.

116.   Rho JK, Choi YJ, Ryoo B-Y, Na III, Yang SH, Kim CH, et al. p53 enhances gefitinib-induced growth inhibition and apoptosis by regulation of Fas in non-small cell lung cancer. Cancer Res. 2007;67: 1163–1169.

117.   Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. Cell. 1990;61: 759–767.

118.   Paterson C, Clevers H, Bozic I. Mathematical model of colorectal cancer initiation. Proc Natl Acad Sci U S A. 2020;117: 20681–20688.

119.   Ortmann CA, Kent DG, Nangalia J, Silber Y, Wedge DC, Grinfeld J, et al. Effect of mutation order on myeloproliferative neoplasms. N Engl J Med. 2015;372: 601–612.

120.   Ernst SM, Mankor JM, van Riet J, von der Thüsen JH, Dubbink HJ, Jgjv A, et al. Tobacco Smoking-Related Mutational Signatures in Classifying Smoking-Associated and Nonsmoking-Associated NSCLC. J Thorac Oncol. 2023;18. doi:10.1016/j.jtho.2022.11.030

121.   Hobbs SD, Wilmink AB, Adam DJ, Bradbury AW. Assessment of smoking status in patients with peripheral arterial disease. J Vasc Surg. 2005;41. doi:10.1016/j.jvs.2004.12.039

122.   Hald J, Overgaard J, Grau C. Evaluation of objective measures of smoking status--a prospective clinical study in a group of head and neck cancer patients treated with radiotherapy. Acta Oncol. 2003;42. doi:10.1080/02841860310005020

123.   Brennan P, Buffler PA, Reynolds P, Wu AH, Wichmann HE, Agudo A, et al. Secondhand smoke exposure in adulthood and risk of lung cancer among never smokers: a pooled analysis of two large studies. International journal of cancer. 2004;109. doi:10.1002/ijc.11682

124.   Beroukhim R, Mermel CH, Porter D, Wei G, Raychaudhuri S, Donovan J, et al. The landscape of somatic copy-number alteration across human cancers. Nature. 2010;463: 899–905.

125.   Besedina E, Supek F. Copy number losses of oncogenes and gains of tumor suppressor genes generate common driver mutations. Nat Commun. 2024;15: 6139.

126.   Tomanek I, Guet CC. Adaptation dynamics between copy-number and point mutations.

Elife. 2022;11. doi:10.7554/eLife.82240

127.   Ciriello G, Cerami E, Sander C, Schultz N. Mutual exclusivity analysis identifies oncogenic network modules. Genome Res. 2012;22: 398–406.

128.   Chen J, Yang H, Teo ASM, Amer LB, Sherbaf FG, Tan CQ, et al. Genomic landscape of lung adenocarcinoma in East Asians. Nat Genet. 2020;52: 177–186.

129.   Imielinski M, Berger AH, Hammerman PS, Hernandez B, Pugh TJ, Hodis E, et al. Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. Cell. 2012;150: 1107–1120.

130.   Jordan EJ, Kim HR, Arcila ME, Barron D, Chakravarty D, Gao J, et al. Prospective Comprehensive Molecular Characterization of Lung Adenocarcinomas for Efficient Patient Matching to Approved and Emerging Therapies. Cancer Discov. 2017;7: 596–609.

131.   Rizvi NA, Hellmann MD, Snyder A, Kvistborg P, Makarov V, Havel JJ, et al. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. Science. 2015;348: 124–128.

132.   Rizvi H, Sanchez-Vega F, La K, Chatila W, Jonsson P, Halpenny D, et al. Molecular Determinants of Response to Anti-Programmed Cell Death (PD)-1 and Anti-Programmed Death-Ligand 1 (PD-L1) Blockade in Patients With Non-Small-Cell Lung Cancer Profiled With Targeted Next-Generation Sequencing. J Clin Oncol. 2018;36: 633–641.

133.   Zhang T, Joubert P, Ansari-Pour N, Zhao W, Hoang PH, Lokanga R, et al. Genomic and evolutionary classification of lung cancer in never smokers. Nat Genet. 2021;53: 1348.

134.   Gillette MA, Satpathy S, Cao S, Dhanasekaran SM, Vasaikar SV, Krug K, et al. Proteogenomic characterization reveals therapeutic vulnerabilities in lung adenocarcinoma. Cell. 2020;182: 200–225.e35.

135.   Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, et al. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. Sci Signal. 2013;6: l1.

136.   Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. Nature. 2014;511: 543–550.

137.   Kadara H, Choi M, Zhang J, Parra ER, Rodriguez-Canales J, Gaffney SG, et al. Whole-exome sequencing and immune profiling of early-stage lung adenocarcinoma with fully annotated clinical follow-up. Ann Oncol. 2017;28: 75–82.

138.   Lawrence M, Gentleman R, Carey V. rtracklayer: an R package for interfacing with genome browsers. Bioinformatics. 2009;25: 1841–1842.

139.   Manders F, Brandsma AM, de Kanter J, Verheul M, Oka R, van Roosmalen MJ, et al. MutationalPatterns: the one stop shop for the analysis of mutational processes. BMC Genomics. 2022;23: 134.

140.   Alexandrov LB, Kim J, Haradhvala NJ, Huang MN, Tian Ng AW, Wu Y, et al. The repertoire of mutational signatures in human cancer. Nature. 2020;578: 94–101.

141.   Rosenthal R, McGranahan N, Herrero J, Taylor BS, Swanton C. deconstructSigs: delineating mutational processes in single tumors distinguishes DNA repair deficiencies and patterns of carcinoma evolution. Genome Biol. 2016;17: 1–11.

142.   Devarakonda S, Morgensztern D, Govindan R. Genomic alterations in lung adenocarcinoma. Lancet Oncol. 2015;16. doi:10.1016/S1470-2045(15)00077-7

143.   Greulich H. The Genomics of Lung Adenocarcinoma: Opportunities for Targeted Therapies. Genes Cancer. 2010;1: 1200.

144.   Seo JS, Ju YS, Lee WC, Shin JY, Lee JK, Bleazard T, et al. The transcriptional landscape and mutational profile of lung adenocarcinoma. Genome Res. 2012;22. doi:10.1101/gr.145144.112

145.   Li S, Choi YL, Gong Z, Liu X, Lira M, Kan Z, et al. Comprehensive Characterization of Oncogenic Drivers in Asian Lung Adenocarcinoma. J Thorac Oncol. 2016;11. doi:10.1016/j.jtho.2016.08.142

146.   Abril-Pla O, Andreani V, Carroll C, Dong L, Fonnesbeck CJ, Kochurov M, et al. PyMC: a modern, and comprehensive probabilistic programming framework in Python. PeerJ Comput Sci. 2023;9: e1516.

147.   Wilks SS. The Large-Sample Distribution of the Likelihood Ratio for Testing Composite Hypotheses. aoms. 1938;9: 60–62.

148.   Cannataro VL, Gaffney SG, Townsend JP. Effect sizes of somatic mutations in cancer. J Natl Cancer Inst. 2018;110: 1171–1177.

149.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550.

150.   Dasari K. Townsend-Lab-Yale/lung-smoking: Software for "Tobacco smoke alters the trajectory of lung adenocarcinoma evolution via effects on somatic selection and epistasis." [cited 23 Jul 2025]. doi:10.5281/zenodo.16379015

Supporting information

Supplementary Figure 1. Strengths of somatic selection on 1,200 oncogenic gene mutations between lung adenocarcinoma (LUAD) in ever-smokers (n = 1,066) and in never-smokers (n = 656; significant: yellow circle, insignificant: no circle, red gene name: stronger selection in ES-LUAD, blue gene name: stronger selection in NS-LUAD).

Supplementary Figure 2. Four patterns of the influences of genetic and environmental interactions in ever-smoker and never-smoker lung adenocarcinoma. (**A**) *RB1* mutations with wild-type (WT) or mutant *KRAS*, (**B**) *RB1* mutations with wild-type or mutant *EGFR*, © MET mutations with wild-type or mutant *TP53*, and (**D**) *STk11* mutations with wild-type or mutant *KEAP1*.

Table S1. Characteristics of datasets included in study.

Study	Institution/project	Sequencing methods	Treatment prior to sample retrieval	Number of samples included in final analysis
The Cancer Genome Atlas Research Network., 2014	The Cancer Genome Atlas	Whole-exome sequencing	3 / 451[*]	451
Jordan et al., 2017	Memorial Sloan Kettering Cancer Center (MSKCC)	Targeted sequencing (MSK-IMPACT 341, MSK-IMPACT 410)	101 / 305	305
Chen et al., 2020	OncoSG	Whole-exome sequencing	≥ 70 / 247[‡,§]	247
Zhang et al., 2021	National Cancer Institute	Whole-genome sequencing	None	187
Rizvi et al., 2018	MSKCC	Targeted sequencing (MSK-IMPACT 341, MSK-IMPACT 410, MSK-IMPACT 468)	164 / 164[‖]	164

		Whole-exome sequencing, whole-genome sequencing, or both	None	153
Imielinski et al., 2012.	Broad Institute			
Gillette et al., 2020	Clinical Proteomic Tumor Analysis Consortium	Whole-exome sequencing	None	74
Kadara et al., 2017	Yale University	Whole-exome sequencing	Not available	73
Jamal-Hanjani et al., 2017	TracerX	Whole-exome sequencing	None	61
Rizvi et al., 2015	MSKCC	Whole-exome sequencing	7 / 7[¶]	7

[*] Treatment not indicated
[†] Combination of targeted therapy, immunotherapy, and/or chemotherapy
[‡] Treatment information is incomplete for 75 samples
[§] Tyrosine-kinase inhibitor and/or chemotherapy
[‖] Anti–PD-(L)1 therapy, sometimes combined with anti–CTLA-4 therapy
[¶] Pembrolizumab