A meta-analysis of age-dependent changes in extracellular vesicle proteins in *C. elegans*

Prasun Kumar Bhunia, Prasad Kasturi [*]

School of Biosciences and Bioengineering, Indian Institute of Technology Mandi, Kamand, 175005, Himachal Pradesh, India

*Corresponding author: prasadkasturi@iitmandi.ac.in

Prasad Kasturi   ORCID ID: 0000-0002-3611-5489

Prasun Kumar Bhunia   ORCID ID: 0000-0003-4201-143X

Abstract

Extracellular vesicles (EVs) contribute to the maintenance of organism-wide proteostasis by mediating intercellular communication. Loss of proteostasis and altered intercellular communication are associated with aging and age-related diseases, suggesting key roles for EVs. However, it is unclear how the proteome of the EVs changes with age. To identify EV-associated proteins and their fate with age, we curated publicly available proteome data from C.elegans model organism. Our analysis reveals that EVs carry proteins that involve protein quality control. We found that abundance of the EV proteins changes significantly with age. Many of these EV proteins also aggregate with age and overlap with Aβ driven protein aggregates. Further, we find that a subset of proteins that alter their abundance in response to heat stress and pathogen infections are also associated with the EVs. This meta-analysis highlights EVs proteome composition, their abundance changes, and aggregation during aging and stress conditions. Overall, this study provides new insights into the dynamics of EV proteins during and aging and may possibly help in identifying potential biomarkers for age-related diseases.

Keywords Aging, proteostasis, extracellular vesicles, *C.elegans*

1. Introduction

Aging is a complex, multifactorial process characterized by the gradual decline in the cellular functions, leading to increased susceptibility to age-related diseases and ultimately, death. One of the hallmarks of aging is loss of protein homeostasis or proteostasis (López-Otín et al., 2023), the state of balanced protein synthesis, folding, and degradation (Hipp et al., 2019). While the molecular chaperones assist in protein folding and prevent aggregation, the proteasomes and autophagy systems degrade misfolded or damaged proteins. As organisms age, the ability to maintain proteostasis deteriorates, leading to the accumulation of damaged or misfolded proteins, which is a hallmark of many age-associated diseases, including neurodegenerative disorders and cancer (Wilson et al., 2023). In an organism-wide context, proteostasis is not limited to individual cells or tissues, but requires coordination across the entire organism. This coordination is facilitated by inter-tissue communication, a process through which tissues and organs exchange signals to maintain systemic balance (Sala et al., 2017; Morimoto 2020). Notably, extracellular vesicles (EVs) have emerged as critical mediators of inter-tissue communication.

Extracellular vesicles (EVs) are small membrane-bound particles secreted by cells into the extracellular space (van Niel et al., 2018; Wang et al., 2024). They carry a wide range of biomolecules, including proteins, lipids, nucleic acids, and metabolites, that can be transferred between cells and influence various physiological processes. The content of EVs can vary depending on the cell type, the environmental conditions, and the state of the cell. As an organism ages, changes in the content, function, and fate of EVs can have significant implications for aging-related processes, including stress responses, inflammation, and tissue maintenance (Guix 2020; Manni et al., 2023). They play a key role in cell-to-cell communication and in maintaining proteostasis across different tissues (Beer and Wehman 2017; Wang and Barr 2018; Sarkar and Patranabis 2024). EVs can facilitate the exchange of protein quality control related molecules such as molecular chaperones, protein aggregates, signalling proteins between tissues, helping coordinate responses to cellular stress, damage, or misfolded proteins across the organism (Takeuchi et al., 2015). In the context of aging or diseases like neurodegeneration, EVs facilitate the transmission of both beneficial and harmful signals such as misfolded proteins (e.g., tau, amyloid-β) between cells, influencing the systemic response to stress, and potentially spreading pathological proteins across tissues (Wallis et al., 2020).

The content of the extracellular proteins and how do they change with age is of particular interest in the context of age-related diseases. Further how proteins involved in proteostasis and stress responses change with age help in understanding organism-wide proteostasis. Here we report a meta-analysis of EV proteins at organism level using *C.elegans* proteome data during conditions of aging, disease, stress and infection. We extended our analysis to protein abundance changes and aggregation with age in wildtype and also in lifespan mutant *C.elegans*. Our analysis suggests that the EV associated proteins

undergo significant changes during aging and their aggregation might affect EVs biogenesis, transport and their function and offer an opportunity to identify potential biomarkers of age-related diseases.

2. Methods

2.1. Data collection and analysis

The EV proteome of *C. elegans* were downloaded from two published datasets (Russell et al., 2020; Nikonorova et al., 2022). The aging proteome of *C. elegans* was downloaded from published dataset (Walther et al., (2015). The aggregation or insoluble proteome datasets of *C. elegans* were downloaded from three published studies (David et al., 2010; Rodrigues et al., 2011; Walther et al., 2015; Anderton et a., 2024). The *C. elegans* heat shock datasets were downloaded from Larance et al., (2011) and Liang et al., 2014 and proteome change after pathogen infection dataset was downloaded from Treitz et al., (2015). The data was analysed in MS-Excel and using Perseus_2.0.6.0 program (Tyanova et al., 2016).

From the Nikonorova dataset, they covered a large number of EV proteins by isolating and separated according to heavy and light subunits. There are 7 replicates for light fractionation and 6 for the heavy one. We summed all the replicates and get rid of the proteins which have zero values (didn't found in any replicate). After that, we got 2363 proteins in light subunit and 2277 in the heavy subunit, in which 1714 proteins to be common in both fractionations. In Russell dataset, we got 617 proteins which considered for further analysis. We merged the both datasets and got 3207 proteins after removing the duplicate proteins and used this dataset and our working EV protein set.

Gene Ontology (GO) term enrichment analysis of biological processes (BP) and cellular components (CC) was performed using Shiny GO 0.81 webserver (Ge et al., 2020) and R Studio.Protein-protein interaction (PPI) network analysis was performed using Metascape (Zhou et al., 2019). Protein abundance changes in wildtype and lifespan mutant worms (*daf-2* and *daf-16* mutant worms) were calculated as fold changes at indicated time points (days) and comparisons were made relative to day1 or as mentioned in the results section.

2.2. Statistical analysis

Origin and GraphPad software were used for graphing and statistical analysis. Mann Whitney U test, these statistical tests and p-values are mentioned in the relevant figure legends.

3. Results

3.1. Analysis of extracellular vesicles associated proteins

Extracellular vesicles (EVs) have been isolated and their content identified from various systems and tissues. They carry a spectrum of biomolecules such as proteins, lipids, RNA and metabolites and implicated in various cellular functions including as signalling molecules. Identification of EV proteome from a complete organism help in finding their role and possible molecular mechanisms of inter-tissue communication and organism wide proteostasis (Wang and Barr 2018; Guix 2020; Manni et al., 2023). To do this, we selected *C. elegans* model organism which is a good system to study aging, proteostasis and inter-tissue communication. There were two studies (Russel et al., 2020; Nikonorova et al., 2022) that reported isolation and identification of EV proteins from *C.elegans*. While Russel et al reported 513 proteins, Nikonorova et al reported 2841 proteins in the EVs (at least in one replicate). There were 348 common proteins (Figure 1A). We combined these two data sets to make a comprehensive list of 3207 EV proteins. We call these proteins as EVAPs (extracellular vesicles associated proteins). The EVs with their content leave the originating cells and travel in the extracellular fluid to reach their target cells. How do the EVAPs maintain their folded structure and function? To answer this, we checked whether the EVs also contain proteins related to proteostasis network (PN). We compared the known 821 PN components from *C.elegans* (Walther et al., 2015) with the EVAPs. We found that 250 proteins related to PN are associated with the EVs (Figure 1B). Among these 95 belong to protein synthesis, 89 belong to protein folding and 66 belong to protein degradation. Based on these numbers, we assume that at least one fourth of the all PN components might present in the EVs. The GO term analysis for cellular components (CC) revealed that these EVAPs are part of vesicles, endomembrane systems and ribosomes (top 20 CC terms) and involve in localization, transport and metabolism related biological processes (BP) (Figure 1 D-E). Nikonorova et al also reported proteins from two types of EVs (heavy and light). Our GO term analysis revealed that there is some difference in the content of these two types of EVs (Suppl Figure 1). These finding confirm earlier observation that PN components can transmit via EVs to maintain organism-wide proteostasis (Takeuchi et al., 2015). In summary we made a comprehensive list of EVAPs from *C.elegans* model organism and found that they carry PN components in addition to many other classes of proteins.

3.2. Identification of age-related changes in the EV proteins

Extracellular vesicles are implicated in aging and proposed to consider as hallmarks of aging (Yin et al., 2021; Manni et al., 2023). EV proteins were analysed from a few senescence cells or plasma from young and old animals. It has been reported that cellular senescence or aging alter EVs cargo (Alibhai et al., 2020). Further it has been observed that the EVs from young animals can improve health span of

old animals (Sanz-Ros et al., 2022; Chen et al., 2024). However, many of these studies mainly focused on the miRNAs of the EVs and studies on proteins were limited.

To found how the EVAPs change with age in an organism level, we compared the EVAPs with an age-related proteome data by Walther et al 2015. In this study Walther et al., identified and quantified proteome wide changes during aging of wildtype and lifespan mutants (*daf-2* and *daf-16*) of *C.elegans*. We analysed abundance changes of the EVAPs in wildtype and lifespan mutants relative to day1 adult worms. We found a small change in the mean abundance levels and this trend is similar in the wildtype as well as in the lifespan mutants (Figure 2A-C). These results suggest that the abundance of the EVAPs change in both directions. Then we specifically focused on proteins that increase or decrease more than 4-fold. We found that the EVAPs that increase more than 4-fold continue to increase with age. In wildtype and lifespan mutant worms, the abundance increase is more than the decrease (Figure 2 D-F). To gain more insight on the EVAPs that altered during aging, we selected proteins that either increase or decrease 2-fold at day 12 and performed GO term analysis (Suppl Figure 2 A-B). We only found minor changes in the biological process and cellular components among the strains. A more detailed analysis is required find changes that vary in the lifespan mutants. It is noteworthy that these protein abundance changes reflect at organismal level, not necessarily in the EVAPs. However, one can assume similar changes in the EVAPs as well because the proteomes were analysed by lysing whole worms.

3.3. Aggregation of extracellular vesicles associated proteins with age

It has been reported that the EVs also carry misfolded, disease related proteins and implicated in age-related neurodegeneration (Hill 2019; Takeuchi and Nagai 2022). For example, Aβ peptides related to Alzheimer's disease has been shown to interact with the EVs (Rajendran et al., 2006; Coughlan et al., 2024). Further it has been observed that the EVs can spread a disease as well as can cause diseases in healthy animals (Aulston et al., 2019; Shippey et al., 2022). Here we wanted to identify whether the EVAPs become insoluble or aggregated during aging. For this we analysed age-related proteome aggregation data from 3 different studies (David et al., 2010; Reis-Rudrigues et al., 2011, Walther et al., 2015). These studies reported aggregation of proteins during aging of *C.elegans* (day10 to day12 of adulthood) by different isolation methods. Walther et al study also reported aggregation data for the lifespan mutants (*daf-2* and *daf-16* mutant worms).

Reis-Rudrigues et al identified 203, David et al identified 711 and Walther et al identified 1414 proteins that aggregated more than 1-fold in aged worms compared to day1 adult worms (Figure 3A). We merged these proteins and identified 1694 proteins that represent age-related insoluble or aggregated proteins. We found that 859 proteins that aggregate during aging overlap with the EVAPs which is corresponding to ~65%. Recently, it has been reported that Amyloid β promotes protein aggregation similar to aging

(Anderton et al., 2024). We wanted to check whether the Aβ induced protein aggregates are part of the EVAPs. We compared the Anderton et al data with the EVAPs and found that 452 of 595 proteins (~76%) overlap with the EVAPs.

Next, we looked at the protein aggregation during aging of the wildtype and lifespan mutant worms. Compared to the day1 worms, there is strong increase in the aggregation of the EVAPs with age in all the strains. This is in line with the suggestion that the EVs can carry protein aggregates (Lim and Lee 2017). This suggest that either the proteins in the EVs aggregate during aging or the aggregated proteins pack into the EVs. We did not detect significant changes in the GO-terms among the worm strains (Suppl Figure 3). However, the long-lived *daf-2* mutant worms may properly maintain their proteins in the EVs. For example, the dauers, a specific larval stage shares many features with the *daf-2* long-lived mutant worms. Recently it has been reported that the the EVs derived from the dauers extend lifespan of the wildtype worms (Ma et al., 2023).

In conclusion, we assume that the EVAPs aggregate during aging in *C.elegans*. Since the analysed proteome data were isolated by lysing whole animals, it is likely that the aggregated protein might represent both the intracellular and extracellular proteome including the EVAPs. Whether the aggregation of the EVAPs affect packing, formation or deliver of the EVs remain to be explored.

3.4. Association of proteins that alter during stress and infection with EV proteins

Extracellular vesicles are also implicated in protection from stresses such as heat stress and pathogen infections. EVs from heat stressed cells can promote survival of non-heat stressed cells by adaptive response, suggesting they carry protective molecules (Bewicke-Copley et al., 2017; Gebremedhn et al., 2020). During infections, EVs can also spread host as well as pathogen derived molecules to promote defence mechanisms and immunity (Rodrigues et al., 2018; Spencer et al., 2021). To identify whether the proteins that are altered during heat stress or pathogen infections are associated with the EVAPs, we analysed proteome data from *C.elegans* under heat stress (Larance et al., 2011; Liang et al., 2014) and pathogen infection conditions (Treitz et al., 2015).

First, we identified proteins that change their abundance during heat stress from both the data sets, combine them and sorted according to their fold change (more or less than 1.2-fold). We found 109 proteins upregulated and 349 proteins down regulated with heat stress. We found that 58 of the 109 upregulated proteins and 212 of the 349 downregulated proteins are associated with the EVAPs. Next, we looked at the proteins that change their abundance with infection to the pathogen, Bacillus thuringiensis. We found 171 proteins upregulated and 117 proteins down regulated in response to the pathogenic bacteria. We found that 110 of the 171 upregulated proteins and 48 of the 117 downregulated

proteins are associated with the EVAPs. This analysis revealed that the EVs contain proteins that are required to respond to stress conditions such as heat stress or pathogen infections.

4. Discussion

Inter-tissue communication has been observed across multiple tissues in a wide range of experimental systems and cell nonautonomous regulation of aging and stress response (O'Brien and van Oosten-Hawle 2016; Morimoto 2020). Organism-wide proteostasis involves complex cellular mechanisms that ensure protein quality control across different tissues (Sala et al., 2017). Inter-tissue communication, particularly via extracellular vesicles, plays a key role in maintaining organism-wide proteostasis by allowing for the exchange of signals and materials between cells and tissues. EVs are important players in this process, mediating both the spread of pathological proteins and potential protective responses that help maintain proteostasis at the organismal level (Takeuchi 2021; Wallis et al., 2020).

The fate of EV-associated proteins is of particular interest, as it remains unclear whether these proteins contribute to beneficial or harmful processes in recipient cells. Using the *C. elegans* model, we identified age-related changes in the composition of EV-associated proteins, including the aggregation of specific proteins and the accumulation of stress-induced factors, such as those triggered by heat stress and pathogen infection. These findings suggest that EVs may serve as a mechanism for the intercellular transfer of damaged or aggregated proteins, potentially contributing to the spread of cellular stress and the progressive decline in proteostasis observed during aging.

A key finding from this study is the identification of protein quality control (PQC) components as EVAPs, including chaperones, ribosome and degradation-related factors. The association of these proteins with EVs emphasizes their potential role in managing cellular stress and maintaining proteostasis through the intercellular transfer of quality-controlled proteins. However, with aging, we observed a shift in the nature of EVAPs, with a higher proportion of aggregated or misfolded proteins becoming associated with EVs. This suggests that as proteostasis deteriorates with age, EVs may become vehicles for the misdirected transfer of dysfunctional proteins.

Furthermore, the aggregation of EVAPs during aging has important implications for the biogenesis and delivery of EVs. The insolubility of aggregated proteins within EVs could interfere with their biogenesis, cargo sorting, or release, thus impairing the overall function of EVs. This disruption in EV function may further contribute to the decline in proteostasis and the progression of age-related diseases.

One of the most striking aspects of our analysis was the overlap between EVAPs and proteins that are known to be altered during heat stress and pathogen infection. We found that association of heat stress and pathogen infection-altered proteins with the EVs, which suggests that EVs play an important role in responding to environmental stressors and propagating stress responses across tissues. This association further underscores the dual role of EVs in both maintaining proteostasis and mediating cellular communication during stress conditions. The incorporation of misfolded or stress-related proteins into EVs may act as a protective mechanism to clear damaged proteins from the cells, but with aging, this process may become dysregulated, facilitating the spread of protein aggregation and exacerbating the aging phenotype.

Limitation of this study is that the data represents proteins from both intracellular and extracellular because of the use of whole worm lysate for proteomic identifications. Future studies focused on isolation and comparison of total as well as insoluble proteomes of EVs from young and aged samples will be essential for further unraveling the complex role of EVs in aging and proteostasis.

Acknowledgements
The authors acknowledge BioX centre, SBB at IIT Mandi for providing facilities.

CRediT authorship contribution statement
Prasun Kumar: Data curation, Formal analysis, Investigation, Methodology, Validation, Visualization, Writing – review & editing. Prasad Kasturi: Conceptualization, Data curation, Formal analysis, Funding acquisition, Methodology, Supervision, Visualization, Writing – original draft, Writing – review & editing.

Funding
This work was supported by funding from Department of Biotechnology, Government of India, as DBT-Ramalingaswami Re-entry grant (BT/RLF/Re-entry/31/2018 to P.K).

Declaration of Competing Interest
The authors have no conflicts of interest to report.

Appendix A. Supporting information
The supporting files containing datasets used in this work will be provided upon request.

References

Alibhai, F.J., Lim, F., Yeganeh, A., DiStefano, P.V., Binesh-Marvasti, T., Belfiore, A., Wlodarek, L., Gustafson, D., Millar, S., Li, S.H., Weisel, R.D., Fish, J.E., Li, R.K., 2020. Cellular senescence contributes to age-dependent changes in circulating extracellular vesicle cargo and function. Aging Cell. 19(3):e13103.

Anderton, E., Chamoli, M., Bhaumik, D., King, C.D., Xie, X., Foulger, A., Andersen, J.K., Schilling, B., Lithgow, G.J., 2024. Amyloid β accelerates age-related proteome-wide protein insolubility. GeroScience 46, 4585–4602.

Beer, K.B., Wehman, A.M., 2017. Mechanisms and functions of extracellular vesicle release in vivo-What we can learn from flies and worms. Cell Adh Migr. 11 (2), 135-150.

Bewicke-Copley, F., Mulcahy, L.A., Jacobs, L.A., Samuel, P., Akbar, N., Pink, R.C., Carter, D.R.F., 2017. Extracellular vesicles released following heat stress induce bystander effect in unstressed populations. J Extracell Vesicles. 6(1):1340746.

Chen, X., Luo, Y., Zhu, Q., Zhang, J., Huang, H., Kan, Y., Li, D., Xu, M., Liu, S., Li, J., Pan, J., Zhang, L., Guo, Y., Wang, B., Qi, G., Zhou, Z., Zhang, C.Y., Fang, L., Wang, Y., Chen, X., 2024. Small extracellular vesicles from young plasma reverse age-related functional declines by improving mitochondrial energy metabolism. Nat Aging. 4(6):814-838.

Coughlan, C., Lindenberger, J., Jacot, J.G., Johnson, N.R., Anton, P., Bevers, S., Welty, R., Graner, M.W., Potter, H., 2024. Specific Binding of Alzheimer's Aβ Peptides to Extracellular Vesicles. Int. J. Mol. Sci. 25 (7), 3703.

David, D.C., Ollikainen, N., Trinidad, J.C., Cary, M.P., Burlingame, A.L., Kenyon, C., 2010. Widespread protein aggregation as an inherent part of aging in C. elegans. PLoS Biol. 2010 8(8):e1000450. doi: 10.1371/journal.pbio.1000450.

Ge, S.X., Jung, D., Yao, R., 2020. ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics 36 (8), 2628-29.

Gebremedhn, S., Gad, A., Aglan, H.S., Laurincik, J., Prochazka, R., Salilew-Wondim, D., Hoelker, M., Schellander, K., Tesfaye, D., 2020. Extracellular vesicles shuttle protective messages against heat stress in bovine granulosa cells. Sci Rep. 10(1):15824. doi: 10.1038/s41598-020-72706-z.

Guix, F.X., 2020. The interplay between aging-associated loss of protein homeostasis and extracellular vesicles in neurodegeneration. J Neurosci Res. 98(2):262-283. doi: 10.1002/jnr.24526.

Hill, A.F., 2019. Extracellular Vesicles and Neurodegenerative Diseases. J Neurosci. 39(47):9269-9273.

Hipp, M.S., Kasturi, P., Hartl, F.U., 2019. The proteostasis network and its decline in ageing. Nat. Rev. Mol. Cell Biol. 20 (7), 421-35.

Larance, M., Bailly, A.P., Pourkarimi, E., Hay, R.T., Buchanan, G., Coulthurst, S., Xirodimas, D.P., Gartner, A., Lamond, A.I., 2011. Stable-isotope labeling with amino acids in nematodes. Nat Methods. 8(10):849-51.

Liang, V., Ullrich, M., Lam, H., Chew, Y.L., Banister, S., Song, X., et al., 2014. Altered proteostasis in aging and heat shock response in C. elegans revealed by analysis of the global and de novo synthesized proteome. Cell Mol Life Sci. 71 (17), 3339-61.

Lim, Y.J., Lee, S.J., 2017. Are exosomes the vehicle for protein aggregate propagation in neurodegenerative diseases? Acta Neuropathol Commun. 5(1):64.

López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., Kroemer, G., 2023. Hallmarks of aging: An expanding universe. Cell. 186(2):243-278.

Ma, J., Wang, Y.T., Chen, L.H., Yang, B.Y., Jiang, Y.Z., Wang, L.X., Chen, Z.Q., Ma, G.R., Fang, L.Q., Wang, Z.B., 2023. Dauer larva-derived extracellular vesicles extend the life of Caenorhabditis elegans. Biogerontology.24(4):581-592. doi: 10.1007/s10522-023-10030-5.

Manni, G., Buratta, S., Pallotta, M.T., Chiasserini, D., Di Michele, A., Emiliani, C., Giovagnoli, S., Pascucci, L., Romani, R., Bellezza, I., Urbanelli, L., Fallarino, F., 2023. Extracellular Vesicles in Aging: An Emerging Hallmark? Cells 12(4), 527.

Nikonorova, I.A., Wang, J., Cope, A.L., Tilton, P.E., Power, K.M., Walsh, J.D., Akella, J.S., Krauchunas, A.R., Shah, P., Barr, M.M., 2022. Isolation, profiling, and tracking of extracellular vesicle cargo in Caenorhabditis elegans. Curr Biol. 32 (9),1924-1936.e6.

O'Brien, D., van Oosten-Hawle, P., 2016. Regulation of cell-non-autonomous proteostasis in metazoans. Essays Biochem. 60 (2), 133-142.

Rajendran, L., Honsho, M., Zahn, T.R., Keller, P., Geiger, K.D., Verkade, P., Simons, K., 2006. Alzheimer's disease beta-amyloid peptides are released in association with exosomes. Proc Natl Acad Sci U S A. 103(30):11172-7.

Reis-Rodrigues, P., Czerwieniec, G., Peters, T.W., Evani, U.S., Alavez, S., Gaman, E.A., Vantipalli, M., Mooney, S.D., Gibson, B.W., Lithgow, G.J., Hughes, R.E., 2012. Proteomic analysis of age-dependent changes in protein solubility identifies genes that modulate lifespan. Aging Cell. 11(1):120-7.

Rodrigues, M., Fan, J., Lyon, C., Wan, M., Hu, Y., 2018. Role of Extracellular Vesicles in Viral and Bacterial Infections: Pathogenesis, Diagnostics, and Therapeutics. Theranostics. 8(10):2709-2721.

Russell, J.C., Kim, T.K., Noori, A., Merrihew, G.E., Robbins, J.E, Golubeva, A., Wang, K., MacCoss, M.J., Kaeberlein, M., 2020. Composition of Caenorhabditis elegans extracellular vesicles suggests roles in metabolism, immunity, and aging. Geroscience 42 (4), 1133-1145.

Sala, A.J., Bott, L.C., Morimoto, R.I., 2017. Shaping proteostasis at the cellular, tissue, and organismal level. J. Cell Biol. 216 (5), 1231-41.

Sanz-Ros, J., Romero-García, N., Mas-Bargues, C., Monleón, D., Gordevicius, J., Brooke, R.T., Dromant, M., Díaz, A., Derevyanko, A., Guío-Carrión, A., Román-Domínguez, A., Inglés, M., Blasco, M.A., Horvath, S., Viña, J., Borrás, C., 2022. Small extracellular vesicles from young adipose-derived stem cells prevent frailty, improve health span, and decrease epigenetic age in old mice. Sci Adv. 8(42):eabq2226.

Sarkar, S., Patranabis, S., 2024. Emerging Role of Extracellular Vesicles in Intercellular Communication in the Brain: Implications for Neurodegenerative Diseases and Therapeutics. Cell Biochem Biophys. 2024 Feb 1. doi: 10.1007/s12013-024-01221-z.

Spencer, N., Yeruva, L. 2021. Role of bacterial infections in extracellular vesicles release and impact on immune response. Biomed J. 44(2):157-164. doi: 10.1016/j.bj.2020.05.006.

Takeuchi, T., 2021. Pathogenic and protective roles of extracellular vesicles in neurodegenerative diseases. J. Biochem. 169 (2): 181-186.

Takeuchi, T., Suzuki, M., Fujikake, N., Popiel, H.A., Kikuchi, H., Futaki, S., et al., 2015. Intercellular chaperone transmission via exosomes contributes to maintenance of protein homeostasis at the organismal level. Proc. Natl. Acad. Sci. USA 112 (19), e2497-506.

Takeuchi, T., Nagai, Y., 2022. Emerging roles of extracellular vesicles in polyglutamine diseases: Mutant protein transmission, therapeutic potential, and diagnostics. Neurochem Int. 157:105357.

Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gíslason, M.H., Pihl, S.I., Tsirigos, K.D., Winther, O., Brunak, S., von Heijne, G., Nielsen H., 2022. SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat. Biotechnol. 40, 1023-1025.

Treitz, C., Cassidy, L., Höckendorf, A., Leippe, M., Tholey, A., 2015. Quantitative proteome analysis of Caenorhabditis elegans upon exposure to nematicidal Bacillus thuringiensis. J Proteomics.113:337-50.

Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., et al., 2016. The Perseus computational platform for comprehensive analysis of proteomics data. Nat. Methods 13 (9), 731-40.

van Niel, G., D'Angelo, G., Raposo, G., 2018. Shedding light on the cell biology of extracellular vesicles. Nat. Rev. Mol. Cell Biol. 19 (4), 213-228.

van Oosten-Hawle, P., Morimoto, R.I, 2014. Organismal proteostasis: role of cell-nonautonomous regulation and transcellular chaperone signaling. Genes Dev. 28(14):1533-43. doi: 10.1101/gad.241125.114.

Walther, D.M., Kasturi, P., Zheng, M., Pinkert, S., Vecchi, G., Ciryam, P., et al., 2015. Widespread Proteome Remodeling and Aggregation in Aging C. elegans. Cell 161 (4), 919-932.

Wallis, R., Mizen, H., Bishop, C.L., 2020. The bright and dark side of extracellular vesicles in the senescence-associated secretory phenotype. Mech Ageing Dev.189:111263.

Wang, J., Barr, M.M., 2018. Cell-cell communication via ciliary extracellular vesicles: clues from model systems. Essays Biochem. 62 (2), 205-213.

Wang, J., Barr, M.M., Wehman, A.M., 2024. Extracellular vesicles. Genetics. Jun 17:iyae088.

Wilson, D.M. 3rd., Cookson, M.R., Van Den Bosch, L., Zetterberg, H., Holtzman, D.M., Dewachter, I., 2023. Hallmarks of neurodegenerative diseases. Cell.186(4):693-714.

Yin, Y., Chen, H., Wang, Y., Zhang, L., Wang, X., 2021. Roles of extracellular vesicles in the aging microenvironment and age-related diseases. J. Extracell. Vesicles 10 (12), e12154.

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A.H., Tanaseichuk, O., Benner, C., Chanda, S.K., 2019. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. Nat. Commun. 10 (1). 1523.

Figure Legends

Fig. 1 Identification of extracellular vesicles associated proteins (EVAP).

A Venn diagram showing the overlap between EVAP found in light and heavy subunit of Nikonorova study with Russell study. There are 2841 and 513 proteins are found in Nikonorova both submit and Russell dataset respectively, in which 348 proteins are common in them.

B Identification of the PN components in the EVAPs. There are 3207 unique EVAPs found from the two different EV studies and 821 PN component proteins found from the Walther study. And there 250 proteins are common between them.

C The PN components associated with EVAPs. Out 250 EVAPs which were common with the PN components, 95 proteins are involved in synthesis, 89 in folding and 66in degradation, which are the three major arms of proteostasis.

D-E Enriched gene ontology (GO) cluster of biological processes (BP) and Cellular components (CC) of EVs found in both subunits respectively.

Fig. 2 Abundance analysis of the EV proteins during aging.

A-C Boxplots showing the relative abundance of the EV associated proteins in Walther et al total proteome data in wildtype and lifespan mutant worms. The relative abundance level is calculated with respect to day1 values (plotted on the y-axis). Time points in days and strains are mentioned on the x-axis. (P values: **wildtype** D1-D6: <0.0001, D1-D12: <0.0001, D1-D17: <0.0001, **daf-2** D1-D6: <0.0001, D1-D12: <0.0001, D1-D17: <0.0001, **daf-16** D1-D6: <0.0001, D1-D12: <0.0001, D1-D17: <0.0001). (* <0.05, **<0.01, ***<0.001, ****<0.0001).

D-F Bar chart representing the number of up and downregulated EV proteins (greater than 4-fold) that change with age in the Walther dataset. The different color codes for different days in WT, *daf-2* and *daf-16* mutant worms respectively. Number of proteins are plotted on the y-axis and time points in days are mentioned on the x-axis.

Fig. 3 Insolubility of EVAPs during aging and disease condition

A Bar chart representing the number of EV proteins found in aggregation datasets of Reis-Rodrigues, Della and Walther datasets respectively. Number of proteins are plotted on the y-axis and different datasets are mentioned on the x-axis. The grey color denotes the number of proteins found in the respective studies and the red color denotes the number of proteins overlapping with EV dataset.

B Venn diagram illustrating the number of identified EV proteins in three different age-related aggregation datasets. There are 3207 EVAPs and 1694 age- related aggregated proteins, in which 859 proteins are common.

C Venn diagrams showing the overlap between EVAPs with Aβ aggregation proteome dataset. In which 3207 are total EVAPs and 595 proteins are found in Aβ aggregation dataset and 452 proteins are between them.

D Boxplots showing the relative abundance of the EV associated proteins in Walther et al aggregation data in wildtype and lifespan mutant worms. The relative abundance level is calculated with respect to day1 values (plotted on the y-axis). Time points in days and strains are mentioned on the x-axis. (P values: **wildtype** D1-D6: <0.0001, D1-D12: <0.0001, *daf-2* D1-D6: <0.0001, D1-D12: <0.0001, *daf-16* D1-D6: <0.0001, D1-D12: <0.0001). (* <0.05, **<0.01, ***<0.001, ****<0.0001).

Fig. 4 Overlap of EVAPs with the proteins that alter during heat stress and pathogen infection.

A-B Bar chart representing the number of up and downregulated proteins during heat stress and the number of proteins among them are EVAPs respectively.

C-D Bar chart representing the number of up and downregulated proteins during pathogen infection and the number of proteins among them are EVAPs respectively.

Supplementary Figure Legends

Suppl Fig 1.

A-B Enriched gene ontology (GO) cluster of biological processes (BP) and Cellular components (CC) of EVs found in only heavy subunit respectively.

C-D Enriched gene ontology (GO) cluster of biological processes (BP) and Cellular components (CC) of EVs found in only light subunit respectively.

Suppl Fig 2.

A Enriched gene ontology (GO) cluster of biological processes (BP) of the EVAPs upregulated and downregulated by 2 fold in day 12 in WT, *daf-2* and *daf-16* lifespan mutant worms.

B Enriched gene ontology (GO) cluster of cellular components (CC) of the EVAPs upregulated and downregulated by 2-fold in day 12 in WT, *daf-2* and *daf-16* lifespan mutant worms.

Suppl Fig 3.

A Enriched gene ontology (GO) cluster of biological processes (BP) of the EVAPs that aggregate at day 12 in in wildtype worms.

B Enriched gene ontology (GO) cluster of cellular components (CC) of the EVAPs that aggregate at day 12 in in wildtype worms.

Figure 1

Bhunia et al

A

Nikonorova et al
(2841)

Russell et al
(513)

2493   348   165

B

EV proteins
(3207)

PN components
(821)

2957   250   571

C

EV associated
PN components

Folding
89

Synthesis
95

Degradation
66

D

Biological Process

Organonitrogen compound metabolic.
Localization
Macromolecule localization
Cellular localization
Cellular organization or biogenesis
Cellular amide metabolic proc.
Establishment of localization
Protein metabolic proc.
Small molecule metabolic proc.
Cellular component organization
Nitrogen compound metabolic proc.
Establishment of localization in cell
Transport
Organonitrogen compound biosynthetic.
Peptide metabolic proc.
Protein localization
Amide biosynthetic proc.
Peptide biosynthetic proc.
Translation
Carboxylic acid metabolic proc.

N. of Genes
● 200
● 400
● 600

−log10(FDR)
60
80
100
120

−log10(FDR)

E

Cellular Components

Cytoplasm
Cytosol
Protein−containing complex
Endomembrane system
Intracellular membrane−bounded organelle
Organelle membrane
Cytosolic ribosome
Endoplasmic reticulum
Cytoplasmic vesicle
Intracellular vesicle
Vesicle
Non−membrane bounded organelle
Intracellular non−membrane bounded organelle
Bounding membrane of organelle
Membrane protein complex
Ribonucleoprotein complex
Organelle subcompartment
Ribosomal subunit
Ribosome
Cytosolic large ribosomal subunit

N. of Genes
● 300
● 600
● 900

−log10(FDR)
100
200
300

100   200   300
−log10(FDR)

Figure 2

Bhunia et al

A

B — EV proteins (3207) / Age-related aggregated proteins (1694): 2348, 859, 835

C — EV proteins (3207) / Aβ driven aggregated proteins (595): 2755, 452, 143

D

A

B

C

D

GO Biological Processes

Suppl Figure 2B

Bhunia et al

GO Cellular components

A

B