Trade-off between sample volume passaged and number of passages during serial dilution for bacterial enumeration

Monika Jain[a], Shuhada Begum[a], Shuvam Bhuyan[a], Chayanika Nath[a], Uchakankhi Kashyap[a], Lukapriya Dutta[a], Shubhra Jyoti Giri[a], Nishita Deka[a], Manabendra Mandal[a], Aditya Kumar[a], Suvendra Kumar Ray[a,*]

[a]Department of Molecular Biology & Biotechnology, Tezpur University, Tezpur–784028, Assam, India.

[*]Corresponding author: Suvendra Kumar Ray: suven@tezu.ernet.in

Abstract

Accurate enumeration of bacteria in a culture is the first step in both fundamental as well as applied research in microbiology. Serial dilution is an age old method used widely by researchers for enumerating viable bacteria in a culture where a specific sample volume is passaged successively to a specific diluent volume. Here, we demonstrated that a higher sample volume is a better representation of bacterial population than a lower sample volume, which was in concordance with the random nature of bacterial distribution in culture. Therefore, a bigger sample to diluent ratio during serial dilution appears more favorable for an accurate bacterial enumeration than a smaller ratio. But surprisingly, enumeration using the different dilution ratios such as 1:9, 1:99 and 1:999 in 1.0 mL final volume yielded similar results with the exception of 1:999, where 1 μL sample was passaged. However, in 10.0 mL final volume of dilution, the above three dilution ratios exhibited similar bacterial enumeration. The experiment was performed using two different bacterial cultures such as *Escherichia coli* and *Ralstonia pseudosolanacearum*. Our results indicated that the advantage gained due to lesser number of passages in case of a lower sample volume could overcome the disadvantage associated with it, thereby co-aligning the different dilution ratios with regards to enumeration. Hence, although in laboratory, 1:9 dilution ratio is usually performed during serial dilution, our results suggest that dilution ratios such as 1:99 in 1 mL dilution volume and ratios such as 1:99 and 1:999 in 10 mL dilution volume are equally effective, which also reduces time, cost and labor.

Keywords

31 Dilution plating, CFU (Colony Forming Units), serial dilution, microliter spotting, diluent

32 volume

1. Introduction

Estimating viable microbial units in a sample serves the fundamental pillar in

microbiology and has been practiced in laboratory for more than one-hundred years after

Robert Koch first attempted to enumerate bacterial count from water in 1883 (Ben-David and

Davidson, 2014). With bacteriological samples, estimation requires sampling of bacterial

population over nutrient agar but their abundance in any sample makes it necessary to begin

with diluting it many folds before performing the sampling (Reynold, 2005). The sample is

diluted sufficiently so that when plated over nutrient agar, each viable unit forms a distinct

colony (cellular aggregate) which can then be counted and estimated in the form of colony

forming units (CFU) (Thomas *et al.*, 2015). In this method of dilution plating, serial dilution

remains the classical technique being employed (Reynold, 2005), where the desirable fold

diluted suspension is attained by a series of successive dilutions with the passaging of a

specific sample volume in a specific diluent volume in order to produce the respective fold

dilution series. Likewise, different sample volumes can be passaged. For example, a 10^6 fold

dilution can be attained by stepwise passaging 100 µL or 10 µL or 1 µL sample volume to a

final 1 mL dilution volume leading to the respective 10-, 100- and 1000-fold dilution series

and requiring six, three and two stepwise dilutions respectively in the process. Considering

the random nature of bacterial distribution in a suspension, anomalies are expected in cell

counts passaged in the successive dilution steps, with lesser anomaly associated with

passaging a higher sample volume. This is attributed to the better tendency of a larger sample

size to contain the randomness prevalent in suspension. However apart from sample volume,

another variable factor that comes into picture here is the complexity of the process (defined

by number of steps involved). In biological experiments, there is a trade-off between

complexity of the process involved and accuracy of the result, influenced by potential errors

associated with each independent step. Evident from our previous study (Bhuyan *et al*. 2023),

CFU estimation by a 10 µL spotting of bacterial suspension was found to be more accurate

than 100 µL spreading of bacterial suspension on plate. The inaccuracy in the 100 µL

spreading was attributed to the complex procedures involving the use of a spreader, ethanol

and heat in the spreading process. So although in serial dilution, theoretically, using a higher

sample to diluent ratio (as 1:9 in 10-fold series) is better aligned with the random distribution

of bacteria; yet using a lower sample to diluent ratio (as 1:999 in 1000-fold series) appears

more accurate considering the reduced complexity of the process. Given the inverse relationship between sample volume and step complexity, it is crucial to assess the influence of varying dilution ratio in bacterial CFU estimation. Besides this, a given fold dilution series can be performed using different diluent volumes in order to adjust to a certain final volume such as 1 mL or 10 mL. Although the sample to diluent ratio remains constant while scaling up the final volume of dilution, the overall force exerted by the diluent while invert mixing might be different in the two cases. This might pose an impact on the degree of random distribution of bacteria in the diluent, overall affecting the CFU estimation upon sampling. Considering this, it was also pertinent to observe the impact of dilution volume on CFU estimation.

Therefore, the current study was carried out in order to understand the influence of two variables of serial dilution: (i) sample to diluent ratio (dilution ratio) and (ii) dilution volume on CFU estimation of bacteria by carrying out a critical examination of 10-, 100- and 1000-fold dilution series. Here, sample to dilution ratio was highest at 1:9 in a 10-fold series, followed by 1:99 in a 100-fold series, and lowest at 1:999 in a 1000-fold series. The studies were conducted using two different bacterial cell suspensions: *E. coli* DH5α and *R. pseudosolanacearum* F1C1 exhibiting two different colony morphologies, with colonies of the former being non-mucoid and latter being mucoid due to the secretion of exopolysaccharides. Microliter spotting approach was employed for the purpose of plating as our previous study (Bhuyan *et al*., 2023) demonstrates a clear advantage of using it over the traditional spread plate method. The present study serves a continuation of this study.

2. Material & Methods

2.1 Bacterial strains and growth conditions

E. coli DH5α (Lab collection) was cultured in Luria Bertani (LB) (Himedia) medium (that contains 1% casein enzyme hydrolysate, 1% NaCl, 0.5% yeast extract) and incubated at 37 ☐C with 200 rpm shaking condition. *R. pseudosolanacearum* F1C1 (Lab collection) (Kumar *et al*., 2013) was cultured in BG medium (Boucher *et al*., 1985) that contains 1% peptone (Himedia), 0.1% yeast extract (Himedia) and 0.1% casamino acid (SRL), supplemented with 0.5% glucose (Himedia) and incubated at 28 ☐C with 150 rpm shaking condition. For preparing solid medium, 1.5% agar (Himedia) was added.

2.2 Culturing of strains and serial dilutions

Saturated cultures of the mentioned strains were obtained after incubating for ⬜16 h
in their respective growth conditions as mentioned above; and then serial dilutions were
performed in saline (0.9% NaCl) as diluent. Three different approaches were employed for
serial dilution *viz*. (i) a 10-fold, (ii) a 100-fold and (iii) a 1000-fold dilution series. The 10-,
100- and 1000-fold dilution series involves the respective series of diluting 1-part sample in
9-parts diluent, 1-part sample in 99-parts diluent and 1-part sample in 999-parts diluent.
Thereby, producing the corresponding dilution ratios of 1:9, 1:99 and 1:999 respectively.
These dilutions were performed for two different final volumes: 1 mL and 10 mL. The
dilutions for 1 mL were performed in 1.5 mL micro-centrifuge tubes (MCT) (from Abdos) as
illustrated in Fig. 1 and in test tubes (from Borosil) for 10 mL dilution volume as illustrated
in S.Fig. 1. Tips were changed between different dilutions of the series. For a more reliable
dilution, precaution was taken during taking in the sample, dispensing it on the walls of next
tube in series and mixing the new suspension obtained as follows: Sample was immediately
taken from a well-mixed suspension (to reduce the chances of cellular sedimentation on
standby). While taking in the sample, the micro-tip was dipped just below the meniscus, not
too deep to reduce bacterial adhesion on the outer surface of tip. The sample was then
dispensed carefully on the walls of the next tube, without touching the diluent so that the
population adhered on the outer surface of the tip would not have a chance to contact the
diluent. Lastly the tip was discarded after dispensing the sample, and the sample on wall was
invert-mixed after closing the MCT or covering the mouth with sterile parafilm in case of
test-tube. From this suspension, next dilution of the series was similarly performed from a
fresh tip.

2.2.1 Dilutions to a final volume of 1 mL

As illustrated in Fig. 1, for a 1 mL final volume, a 10-fold dilution series was
performed by adding 100 µL of culture to 900 µL of saline in the first step to obtain a 10^1-
fold dilution, subsequently 100 µL of this suspension was added to the next tube in the series
containing 900 µL of saline to obtain a 10^2-fold dilution cell suspension. This was repeated
until a cell suspension of 10^7-fold dilution was obtained. Similarly, a 100-fold dilution series
was performed by adding 10 µL of culture to 990 µL of saline in the first step to obtain a 10^2-
fold dilution. In subsequent steps, 10^4- and 10^6-fold dilutions were similarly obtained. Lastly,
to proceed with a 1000-fold dilution series, 1 µL of culture was added to 999 µL of saline to
obtain 10^3-fold dilution in first step and 10^6-fold dilution in second step. In both the latter
dilution series (100-fold and 1000-fold series), the final 10^7-fold dilution was obtained by

performing a 10-fold dilution from their respective 10^6-fold diluted suspensions.

Fig. 1: Three different ways of attaining a 10^6 fold dilution has been shown *viz* a 10-, 100-

and 1000-fold dilution series. (i) A 10-fold series (1:9 dilution ratio) involves passaging 100

µL suspension to 900 µL saline in six successive dilutions. (ii) A 100-fold series (1:99

dilution ratio) involves passaging 10 µL suspension to 990 µL saline in three successive

dilutions. (iii) A 1000-fold series (1:999 dilution ratio) involves passaging 1 µL suspension to

999 µL saline in two successive dilutions. Bacterial culture is shown in yellow in the screw

caped bottle. Diluent (saline) is shown in blue in micro-centrifuge tubes. Created with

Biorender.com.

2.2.2 Dilutions to a final volume of 10 mL

Dilutions were performed in a similar way as mentioned above, only the volumes of

sample and diluent taken were both up scaled (although same in proportion) for adjusting it to

10 mL volume as illustrated in S.Fig. 1. In a 10-fold dilution series 1 mL of culture was

added to 9 mL of saline to obtain a 10^1-fold dilution, which was repeated with subsequent

diluents in series to obtain 10^7-fold dilution. In a 100-fold dilution series, 100 µL of culture

was added to 9.900 mL of saline to obtain a 10^2-fold dilution. In subsequent steps, 10^4- and

10^6-fold dilutions were similarly obtained. Lastly, in a 1000-fold dilution series, 10 µL of

culture was added to 9.990 mL of saline to obtain 10^3-fold dilution and then 10^6-fold dilution

in second step. In both the latter dilution series, a 10^7-fold dilution was obtained by

performing a 10-fold dilution from their respective 10^6-fold diluted suspensions.

2.3 Plating and colony observation

Plating was done from a 10^6- and 10^7-fold diluted suspensions obtained from each of

the series for *E. coli* DH5α and at 10^7- and 10^8-fold diluted suspensions obtained from each

of the series for *R. pseudosolanacearum* F1C1. A higher fold dilution was chosen for *R.*

pseudosolanacearum F1C1 as (i) with the saturated cultures of the two, the number of

colonies observed in this case were always higher than that of *E. coli* DH5α by 2-6 times at

same fold dilution (this can also be observed in Table 1) and (ii) *R. pseudosolanacearum*

forms typical mucoidal colonies once they appear completely in plate; also they show

twitching motility over solid media unlike *E. coli* (Bhuyan *et al*., 2023, 2024b); the latter one

forming non-mucoidal colonies. So apparently, colonies of the former one had greater

tendency to merge with other such colonies.

For plating, microliter spotting approach was followed (Bhuyan *et al*., 2023).

Colonies were observed and counted after 12-14 h of incubation in case of *E. coli* DH5α and

after 48 h of incubation in case of *R. pseudosolanacearum* F1C1. Although with the latter

one, colonies appear fully after 48 h of incubation (due to a slower generation time), but the

simultaneous production of copious amounts of EPS by the bacteria makes it fluidal and

difficult to count at this point. So in this case, a prior observation of the colonies was also

made at 28 h post incubation, where tiny colonies although not so prominent can be lightly

visualized against light and accordingly counted. These were tallied after the colonies

completely appear.

2.4 Study on the influence of plating volume on bacterial enumeration

To study the influence of plating/sampling volumes on bacterial enumeration, ten
different spots of each of the different volumes: 5 µL, 10 µL, 15 µL and 20 µL were made in
separate nutrient agar plates. These were sampled from a specific suspension which was at
10^6 fold dilution for *E. coli* DH5α and at 10^7 fold dilution for *R. pseudosolanacearum* F1C1,
obtained simply from a 10-fold series performed for 1 mL final volume. Plates were then kept
under incubation for colony observation. After incubation, average number of colonies per
spot was calculated in each case.

2.5 Study on the influence of dilution ratio on bacterial enumeration

To study the influence of dilution ratio used in serial dilution on bacterial
enumeration, all the three dilution series (10-fold, 100-fold and 1000-fold series) were
performed for the two final volumes: 1 mL and 10 mL in the independent experiments
generated from different cultures. For plating, ten spots of 10 µL each were made from a 10^6-
and 10^7- fold diluted suspensions of *E. coli* DH5α and a 10^7- and 10^8- fold diluted
suspensions of *R. pseudosolanacearum* F1C1. Plates were then kept under incubation for
colony observation. After incubation, average number of colonies per spot was calculated in
each case.

2.6 Study on the influence of dilution volume on bacterial enumeration

For this study, all the dilution series were performed for both 1 mL and 10 mL
dilution volume and were generated from same culture, so that except the factor of dilution
volume chosen, all other factors remain constant. This study was performed with *E. coli*
DH5α as representative. Plating was done from a 10^6 and 10^7 fold diluted suspension. For
each of the three dilution series, ten spots of 10 µL each were made for it being performed at
1 mL and 10 mL final volume in the different halves of same plate to enable clear
comparison. The plates were then kept under incubation and later average number of colonies
per spot was calculated for all the cases.

2.7 Statistical analyses

2.7.1 Coefficient of Variation (CoV)

197   For comparing different sample volumes used to estimate bacterial cell density, CoV
198   (which is a measure of variability in a population) was calculated among the colony numbers
199   observed in the ten replicates of each sampling volumes used. First, average number of

colonies and standard deviation were calculated for each of the different sample volumes, then CoV was calculated by dividing standard deviation by average.

$$CoV = \frac{\text{Standard deviation}}{\text{Average}}$$

2.7.2 Correlation analysis

To study the correlation between different sampling volumes used for bacterial cell density estimation and CoV obtained in the colony count data corresponding to each sampling volume, Pearson's correlation analysis was performed. The Pearson's correlation coefficient (r) was calculated as:

$$r = \frac{\sum(x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum(x_i - \bar{x})^2 \sum(y_i - \bar{y})^2}}$$

where x_i = values of x-variable (here, x takes the values of different sample volumes); $\bar{x}$ = mean of the values of x-variable; y_i = values of y-variable (here, y takes the values of CoV corresponding to each values of x); $\bar{y}$ = mean of the values of y-variable

2.7.3 One-way ANOVA

For comparing the mean of sample population of the suspensions obtained from different dilution series, one-way ANOVA test was performed in MS Excel at α=0.05 with H_0 stating that there is no significance difference between means of sample population of suspensions obtained from these different dilution series and H_1 stating that the means of sample population of suspensions obtained from these dilution series are significantly different.

2.7.4 Bonferroni post-hoc test

In cases where one-way ANOVA proved significant difference between the groups, a post-hoc test was performed using Bonferroni corrected p-value to identify which group differ significantly. This was calculated as:

Bonferroni-corrected p-value = α / number of comparative tests performed

When p-value of comparative tests < Bonferroni-corrected p-value, the test is significant.

2.7.5 Mann-Whitney U test

A two-tailed Mann-Whitney U test was performed to compare the mean difference of

colony counts obtained with dilutions performed for 1 mL and 10 mL volumes. Test is

significant when $U_{Stat} < U_{Crit}$.

U_{Stat} = minimum of the two values, U_1 and U_2.

$$U_1 = n_1 n_2 + \frac{n_1(n_1+1)}{2} - R_1 \qquad \text{and} \qquad U_2 = n_1 n_2 + \frac{n_2(n_2+1)}{2} - R_2$$

Where, n_1 and n_2 represents sample size of the two groups under comparison; R_1 and R_2

denotes rank sum of the two groups respectively when all of them are arranged in ascending

order.

3. Result

3.1 A higher sample volume of a bacterial suspension is a better representation of the

population size than a lower sample volume

During bacterial enumeration by dilution method, a specific volume of sample is

stepwise passaged to a specific volume of the diluent. As distribution of bacteria is random in

a culture, there would be anomalies associated with the colony numbers being passaged. In

consideration of the random distribution, the sample volume being passaged is likely to play

an important role in the enumeration. To get an insight into the influence of sample volume

on consistency of bacterial count, we studied the variation prevalent in colony count in the

plating of different sample volumes: 5 μL, 10 μL, 15 μL and 20 μL of a diluted bacterial

suspension. The colony numbers obtained in 10 spots of each of these sample volumes is

represented in Table 1. To enable clear distinction of colonies in spotted area, the suspension

used was at 10^6 fold dilution of the saturated culture in case of *E. coli* and at 10^7 fold dilution

in case of *R. pseudosolanacearum*. In case of *E. coli*, the CoV observed in colony numbers

obtained with 5 μL, 10 μL, 15 μL and 20 μL spotting were 0.18, 0.12, 0.10 and 0.07

respectively. When the same study was conducted with *R. pseudosolanacearum*, the CoV

was found to be 0.42, 0.37, 0.26 and 0.17 respectively for 5 μL, 10 μL, 15 μL and 20 μL

spotting. Clearly, for both the independently tested bacterial strains, the coefficient of

variation (CoV) of bacterial count in the different spots made of a given sampling volume

was observed to be highest with the lowest sampling volume used (5 μL) and lowest with the

highest sampling volume used (20 μL). Similar observations were recorded in four repetitive

experiments (S.Table1). This is in concordance with the view of random distribution of bacteria in a culture; as with a randomly distributed population, a larger sampling size is supposed to give more consistency in density estimation. As expected, a strong negative correlation was found between sampling volumes of the suspension used and coefficient of variation (CoV) of bacteria count, with the same trend being observed for both the bacterial species ($r = -0.97$ for *E. coli* and $r = -0.99$ for *R. pseudosolanacearum*) (Fig. 2). This suggests that higher the sampling volume of the suspension, greater the consistency in bacterial cell count and lesser the variability. Hence, higher sampling volume is a better representation of the bacterial population.

Table 1: Colony numbers obtained in different plating volumes of the two different bacterial cell suspensions.

	Colony count data of 10 spots							
Bacterial samples	***E. coli* DH5α**				***R. pseudosolanacearum* F1C1**			
Dilution factors	10^6				10^7			
Spotting volume	**5 µL**	**10 µL**	**15 µL**	**20 µL**	**5 µL**	**10 µL**	**15 µL**	**20 µL**
	10	27	47	44	7	6	11	14
	11	24	42	40	4	6	10	16
	12	21	36	49	1	10	17	15
	15	28	39	45	4	8	11	12
	12	26	36	42	5	6	9	17
	13	24	34	44	2	6	11	15
	12	28	42	45	4	8	9	19
	18	19	39	49	5	14	12	11
	15	25	38	41	4	4	6	15
	14	24	36	42	6	8	10	19
Mean	13.2	24.6	38.9	44.1	4.20	7.60	10.60	15.30
SD	2.35	2.91	3.87	3.07	1.75	2.80	2.80	2.63
CoV	**0.18**	**0.12**	**0.10**	**0.07**	**0.42**	**0.37**	**0.26**	**0.17**
CFU estimate	2.6×10^9	2.5×10^9	2.6×10^9	2.2×10^9	8.4×10^9	7.6×10^9	7.1×10^9	7.7×10^9

Standard deviation (SD), Coefficient of Variation (CoV) and CFU estimate calculated for parent culture in each of these cases are represented.

Fig. 2: Correlation between suspension volume spotted (X- axis) and coefficient of variation (CoV) obtained among the colony numbers (Y-axis) of (a) *E. coli* and (b) *R. pseudosolanacearum*. Pearson's correlation coefficient (r) is represented for each case. A strong negative correlation has been observed in each case.

3.2 Stepwise passaging of 1 mL, 100 µL and 10 µL (but not 1 µL) are similar with regards to CFU estimation of E. coli DH5α

In theory, to make a 10^6 fold dilution of an original culture different dilution series such as ten or hundred or thousand fold can be used which has been shown in Fig. 1. As evident from the Fig, in case of a higher fold dilution series such as thousand, only 1 µL of the bacterial suspension is passaged twice to achieve 10^6 fold dilution whereas in case of a lower fold dilution series such as ten, 100 µL of bacterial suspension is passaged six times. In the intermediary hundred fold dilution series, 10 µL of bacterial suspension is passaged thrice. In this way, the three different approaches of serial dilution are different with respect to (i) sample to diluent ratio and (ii) complexity of the series (number of passaged involved). As mentioned earlier, in view of the random distribution of bacteria, a higher dilution ratio (as 1:9 in 10-fold series) seems a better approach for CFU estimation. However, considering the complexity involved, a lower dilution ratio (as 1:999 in 1000-fold series) seems better. To find out the more accurate dilution series considering this trade-off, we estimated the CFU of *E. coli* culture using these three dilution series *viz* a 10-fold, a 100-fold and a 1000-fold dilution series as discussed below.

The CFU of bacterial suspension obtained from the three different dilution series (a 10-, 100-, and 1000-fold dilution series) is represented in Table 2a and a representative result

is shown in Fig. 3a. The 10 μL spotting of 10^6 fold dilution obtained from a 10-fold and a

100-fold dilution series produced average 11.9 ± 4.1 and 11.3 ± 4.5 numbers of colonies per

spot respectively; however that obtained from a 1000-fold dilution series produced average

28.6 ± 1.8 numbers of colonies per spot. Similarly with 10^7 fold dilution average number of

colonies per spot obtained from a 10- and 100-fold dilution series were 1.4 ± 1.0 and 1.1 ±

1.2 respectively; however a 1000-fold dilution series produced average 3.5 ± 1.4 number of

colonies per spot. A p-value < 0.05 was indicative of significant differences among the

groups, and thus a post-hoc test was followed in each case to identify the group being

significantly different. This was observed to be the 1000-fold series in almost all of these

cases. The results indicated that a 10-fold and 100-fold dilution series are producing similar

CFU count whereas the 1000-fold dilution series is giving a higher CFU count than the other

two. This trend was further observed in all the three repetitive experiments (Table 2a). In

addition the CFU count by the 1000-fold dilution series was inconsistent. This was

perplexing for us to find out if a 1000-fold dilution series is more accurate than the other two

dilution series. As the results by 1000-fold dilution series were inconsistent, we thought

pipetting a smaller volume like 1 μL might be a more complex procedure while taking in as

well as dispensing out the suspension. To avoid any bias related to pipette, we repeated the

experiments using pipettes from three neighboring labs for pipetting 1 μL volume. In this

case also, an inconsistently higher CFU count was observed with the 1000-fold series. This

led us to assume that 1 μL pipetting indeed is more complex and prone to inconsistencies

than 10 μL and 100 μL pipetting. This might be because a minor deviation in either taking in

and/or dispensing out the 1 μL suspension would contribute a larger percentage of change in

the total volume.

To compare the accuracies of these dilution series after eliminating the complex factor

of 1 μL pipetting, all the three series were re-performed this time to a final 10 mL volume by

passaging stepwise 1 mL, 100 μL and 10 μL respectively in the 10-, 100- and 1000-fold

dilution series. The CFU now obtained with each of these dilution series are represented in

Table 2b and a representative result is shown in Fig. 3b. The 10 μL spotting of 10^6 fold

dilution obtained from a 10-, 100- and 1000-fold dilution series produced average 13.7 ± 4.2,

14.7 ± 2.1 and 16.7 ± 3.9 numbers of colonies per spot respectively. Similarly with the 10^7

fold dilution, average number of colonies obtained from the 10-, 100- and 1000-fold dilution

series were 1.6 ± 1.4, 2.6 ± 1.8 and 2.0 ± 0.9 respectively. This trend was further observed in

all the three repetitive experiments. It was interesting to observe this time a similar number of

colonies with all the three dilution series. A p-value > 0.05 as obtained in one-way ANOVA
was indicative of no significant differences between the colony numbers obtained from the
different dilution series. This proves a 1000-fold dilution series to be equally accurate as the
10-fold and 100-fold dilution series provided that pipetting of 1 μL is not carried out.

Table 2(a): Average colony numbers (along with standard deviations) of *E. coli* DH5α
obtained from different dilution series performed to a final 1 mL volume.

Dilutions adjusted to a total 1 mL volume (*E. coli* DH5α)						
Dilution factors	**10^6**			**10^7**		
Dilution series	**10 fold**	**100 fold**	**1000 fold**	**10 fold**	**100 fold**	**1000 fold**
Dilution ratios	**1:9** (A)	**1:99** (B)	**1:999** (C)	**1:9** (A)	**1:99** (B)	**1:999** (C)
Experiment 1 — Colony Count	11.9 ± 4.1	11.3 ± 4.5	28.6 ± 1.8	1.4 ± 1.0	1.1 ± 1.2	3.5 ± 1.4
Experiment 1 — One-way ANOVA	$F(2, 27) = 71.64, p = 1.6 \times 10^{-11}$			$F(2, 27) = 11.60, p = 2.3 \times 10^{-4}$		
Experiment 1 — Bonferroni post-hoc test	(C) > (A), (B); (A) ☐ (B)			(C) > (A), (B); (A) ☐ (B)		
Experiment 2 — Colony Count	9.5 ± 1.6	8.3 ± 3.4	28.8 ± 5.2	1.8 ± 1.2	1.8 ± 1.3	3.5 ± 2.0
Experiment 2 — One-way ANOVA	$F(2, 27) = 108.21, p = 1.3 \times 10^{-13}$			$F(2, 27) = 4.08, p = 2.8 \times 10^{-2}$		
Experiment 2 — Bonferroni post-hoc test	(C) > (A), (B); (A) ☐ (B)			(A) ☐ (B) ☐ (C)		
Experiment 3 — Colony Count	22.5 ± 3.3	19.9 ± 3.5	45.4 ± 5.1	3.4 ± 1.8	2.9 ± 1.6	7.5 ± 2.0
Experiment 3 — One-way ANOVA	$F(2, 27) = 120.33, p = 3.5 \times 10^{-14}$			$F(2, 27) = 19.15, p = 6.6 \times 10^{-6}$		
Experiment 3 — Bonferroni post-hoc test	(C) > (A), (B); (A) ☐ (B)			(C) > (A), (B); (A) ☐ (B)		

329      $F_{statistic}$ and p-value calculated from One-way ANOVA and interpretation of Bonferroni post-

330   hoc tests for the different comparable groups are represented.

331   **Table 2(b):** Average colony numbers (along with standard deviations) of *E. coli* DH5α

332   obtained from different dilution series performed to a final 10 mL volume.

Dilutions adjusted to a total 10 mL volume (*E. coli* DH5α)						
Dilution factors	**10^6**			**10^7**		
Dilution series	**10 fold**	**100 fold**	**1000 fold**	**10 fold**	**100 fold**	**1000 fold**
Dilution ratios	**1:9** **(A)**	**1:99** **(B)**	**1:999** **(C)**	**1:9** **(A)**	**1:99** **(B)**	**1:999** **(C)**
Experiment 1 — **Colony Count**	13.7 ± 4.2	14.7 ± 2.1	16.7 ± 3.9	1.6 ± 1.4	2.6 ± 1.8	2.0 ± 0.9
Experiment 1 — **One-way ANOVA**	F (2, 27) = 1.87, p = 0.17			F (2, 27) = 1.20, p = 0.32		
Experiment 2 — **Colony Count**	17.5 ± 5.0	18.3 ± 4.5	21.5 ± 2.7	1.7 ± 1.5	2.3 ± 1.2	1.6 ± 1.0
Experiment 2 — **One-way ANOVA**	F (2, 27) = 2.57, p = 0.10			F (2, 27) = 0.95, p = 0.40		
Experiment 3 — **Colony Count**	16.0 ± 3.9	17.6 ± 5.2	18.8 ± 2.3	1.8 ± 0.9	1.7 ± 1.2	2.0 ± 1.2
Experiment 3 — **One-way ANOVA**	F (2, 27) = 1.23, p = 0.31			F (2, 27) = 0.20, p = 0.82		

333      $F_{statistic}$ and p-value calculated from One-way ANOVA for the different comparable groups

334   are represented.

336
Fig. 3: *E. coli* colonies at selected fold dilutions (10^6 and 10^7) obtained with the different
dilution series. These series were performed first to a final volume of (a) 1 mL and later to a
final volume of (b) 10 mL from independent cultures. At 1 mL dilution volume, a clearly
greater number of colonies were observed with the 1000-fold series; however, at 10 mL
dilution volume, similar numbers of colonies were observed with all the three dilution series.

3.3 Stepwise passaging of 1 mL, 100 μL and 10 μL (but not 1 μL) are similar with regards to
CFU estimation of R. pseudosolanacearum F1C1

We further performed this CFU estimation study for a different bacterium: *R.*
pseudosolanacearum, a phytopathogen, which in contrast to *E. coli*, forms mucoidal colonies
on plate due to the production of exopolysaccharides (Bhuyan *et al.*, 2024a; 2024c). This
study was undertaken to know whether the accuracies observed previously with the different
dilution series holds true for the other bacterial suspension too. Unlike *E. coli*, a higher fold
dilution (10^7 and 10^8) was chosen for plating this time as the mucoidal nature of the colonies

and twitching motility exhibited by *R. pseudosolanacearum* over solid media (Bhuyan *et al.*, 2024b) leads towards a greater tendency for the colonies to merge among each other. Also the CFU/mL of the saturated culture of *R. pseudosolanacearum* was always observed to be higher than that of *E. coli* by 2-6 times (this can also be observed in Table1). So plating a higher fold dilution in this case will enable clear distinction of the colonies. When the dilutions were performed to a final volume of 1 mL, average number of colonies observed per 10 µL spot of 10^7 fold dilution were 6.0 ± 2.5, 5.4 ± 1.0 and 14.1 ± 3.5 respectively with a 10-, 100- and 1000-fold dilution series (Table 3a). Similarly with 10^8 fold dilution, average number of colonies per spot obtained from a 10-, 100-, and 1000-fold dilution series were 0.6 ± 0.8, 0.4 ± 0.7 and 2.5 ± 1.3 respectively. A p-value < 0.05 was indicative of significant differences among the groups, and thus a post-hoc test was followed in each case to identify the group being significantly different. This was observed to be the 1000-fold series in almost all of these cases. A representative result is shown in Fig. 4(a). Aligning with our previous observation, CFU recorded with the 1000-fold series were again observed to be significantly higher and were inconsistent than that recorded with the other two dilution series. This further highlight the potential complexities associated with 1 µL pipetting in the 1000-fold dilution series and so to eliminate it, the dilutions were re-performed to a final volume of 10 mL as done previously with *E. coli* suspension. In this case, average number of colonies observed per 10 µL spot of 10^7 fold dilution were 9.8 ± 1.9, 8.4 ± 4.3 and 7.2 ± 2.0 respectively with a 10-, 100- and 1000-fold dilution series (Table 3b). Similarly with 10^8 fold dilution, average number of colonies per spot obtained from a 10-, 100-, and 1000-fold dilution series were 0.3 ± 0.7, 0.5 ± 0.7 and 0.2 ± 0.4 respectively. It was interesting to note that the colony numbers observed were similar in this case and this trend was repetitive in all the three experiments. A representative result is shown in Fig. 4(b). A p-value > 0.05 as obtained in one-way ANOVA was indicative of no significant differences between the colony numbers obtained from the different dilution series. This was again in concordance with our previous observation of *E. coli* CFU estimation. This establishes 1000-fold dilution series to be equally accurate as the 10-fold and 100-fold dilution series when 1 µL pipetting is not carried out and validates our understanding on the inconsistencies and complexities associated with 1 µL pipetting. In summary, when performing the dilutions to a final volume of 1 mL, a 10- and 100-fold series proved to be effective for CFU estimation and when performing them to a final volume of 10 mL, all the 10-, 100- and 1000-fold dilution series proved to be equally effective for CFU estimation. But considering the time, cost and labor efficiency of the approach with no compromise on accuracy of CFU estimation, a 1000-fold

 dilution series shall be preferred when performing the dilutions for 10 mL final volume and a

100-fold dilution series shall be preferred when performing the dilutions for 1 mL final

volume. This is because a relatively lower number of passages are involved in these

processes compared to the 10-fold series.

Table 3(a): Average colony numbers (along with standard deviations) of *R.*

pseudosolanacearum F1C1 obtained from different dilution series performed to a final 1 mL

volume.

Dilutions adjusted to a total 1 mL volume (*R. pseudosolanacearum* F1C1)						
Dilution factors	10^7			10^8		
Dilution series	**10 fold**	**100 fold**	**1000 fold**	**10 fold**	**100 fold**	**1000 fold**
Dilution ratios	**1:9** **(A)**	**1:99** **(B)**	**1:999** **(C)**	**1:9** **(A)**	**1:99** **(B)**	**1:999** **(C)**
Experiment 1 — **Colony Count**	6.0 ± 2.5	5.4 ± 1.0	14.1 ± 3.5	0.6 ± 0.8	0.4 ± 0.7	2.5 ± 1.3
Experiment 1 — **One-way ANOVA**	$F (2, 27) = 35.95$, $p = 2.4 \times 10^{-8}$			$F (2, 27) = 14.34$, $p = 5.7 \times 10^{-5}$		
Experiment 1 — **Bonferroni post-hoc test**	(C) > (A), (B); (A) �□ (B)			(C) > (A), (B); (A) �□ (B)		
Experiment 2 — **Colony Count**	7.1 ± 2.5	4.9 ± 1.4	21.9 ± 3.3	0.8 ± 0.6	0.2 ± 0.4	2.0 ± 1.6
Experiment 2 — **One-way ANOVA**	$F (2, 27) = 135.21$, $p = 8.6 \times 10^{-15}$			$F (2, 27) = 8.34$, $p = 1.5 \times 10^{-3}$		
Experiment 2 — **Bonferroni post-hoc test**	(C) > (A), (B); (A) �□ (B)			(C) �□ (A); (A) �□ (B); (C) > (B)		
Experiment 3 — **Colony Count**	3.7 ± 2.1	1.8 ± 0.9	11.9 ± 3.3	0.6 ± 0.7	0.3 ± 0.5	1.7 ± 0.9
Experiment 3 — **One-way ANOVA**	$F (2, 27) = 53.79$, $p = 3.8 \times 10^{-10}$			$F (2, 27) = 10.05$, $p = 5.5 \times 10^{-4}$		
Experiment 3 — **Bonferroni post-hoc test**	(C) > (A), (B); (A) �□ (B)			(C) > (A), (B); (A) �□ (B)		

391   $F_{statistic}$ and p-value calculated from One-way ANOVA and interpretation of Bonferroni post-
392   hoc tests for the different comparable groups are represented.

393   **Table 3(b):** Average colony numbers (along with standard deviations) of *R.*
394   *pseudosolanacearum* F1C1 obtained from different dilution series performed to a final 10 mL
395   volume.

Dilutions adjusted to a total 10 mL volume (*R. pseudosolanacearum* F1C1)							
Dilution factors		**10^7**			**10^8**		
Dilution series		**10 fold**	**100 fold**	**1000 fold**	**10 fold**	**100 fold**	**1000 fold**
Dilution ratios		**1:9** **(A)**	**1:99** **(B)**	**1:999** **(C)**	**1:9** **(A)**	**1:99** **(B)**	**1:999** **(C)**
Experiment 1	**Colony Count**	9.8 ± 1.9	8.4 ± 4.3	7.2 ± 2.0	0.3 ± 0.7	0.5 ± 0.7	0.2 ± 0.4
	One-way ANOVA	F (2, 27) = 1.97, p = 0.16			F (2, 27) = 0.62, p = 0.55		
Experiment 2	**Colony Count**	4.3 ± 1.7	4.8 ± 1.8	4.1 ± 1.3	0.5 ± 0.7	0.5 ± 0.7	0.3 ± 0.5
	One-way ANOVA	F (2, 27) = 0.51, p = 0.61			F (2, 27) = 0.32, p = 0.73		
Experiment 3	**Colony Count**	3.3 ± 2.5	3.2 ± 1.7	3.9 ± 1.5	0.5 ± 0.8	0.3 ± 0.5	0.5 ± 0.5
	One-way ANOVA	F (2, 27) = 0.37, p = 0.70			F (2, 27) = 0.32, p = 0.73		

396   $F_{statistic}$ and p-value calculated from One-way ANOVA for the different comparable groups
397   are represented.

Fig. 4: *R. pseudosolanacearum* colonies at selected fold dilutions (10^7 and 10^8) obtained with the different dilution series. These series were performed first to a final volume of (a) 1 mL and later to a final volume of (b) 10 mL from independent cultures. At 1 mL final volume, a clearly greater number of colonies were observed with the 1000-fold series; however, at 10 mL final volume, similar numbers of colonies were observed with all the three dilution series.

3.4 Bacterial enumeration is indifferent to the dilution volume used in serial dilution

To observe if dilution volume used in serial dilution has an influence on bacterial enumeration, each of the 10-, 100- and 1000-fold dilution series were performed from a saturated *E. coli* culture using different diluent volumes to reach the final volume of 1 mL and 10 mL respectively. CFU observed with the plating of 10^6 and 10^7 fold dilutions obtained from each of the three dilution series adjusted to the different dilution volumes are represented in Table 4 and a representative result is shown in Fig. 5. With 10^6 fold dilution, a

 10-fold series yielded 21.9 ± 3.0 and 18.1 ± 3.4, a 100-fold series yielded 19.7 ± 3.3 and 19.1

± 3.4, and a 1000-fold series yielded 46.9 ± 5.3 and 19.9 ± 2.6 average numbers of colonies

for the final 1 mL and 10 mL dilution volumes respectively. Similarly, with 10^7 fold dilution,

a 10-fold series yielded 2.6 ± 1.1 and 2.4 ± 1.3, a 100-fold series yielded 2.5 ± 1.3 and $1.9 \pm$

1.0, and a 1000-fold series yielded 5.6 ± 2.1 and 1.9 ± 1.3 average numbers of colonies for the

final 1 mL and 10 mL dilution volumes respectively. Clearly, the colony numbers observed

with the 1 mL and 10 mL dilution volumes were similar for most of the times for a 10-fold

and 100-fold series; however they were greatly different in all the cases for the 1000-fold

series. This suggests that the use of two dilution volumes brought no contrasting difference in

the bacterial density being passaged in the 10-fold and 100-fold dilution series. However, the

passaging in 1000-fold series was contrastingly different in the two diluent volumes used.

Here, the 1 µL passaging (in 1 mL dilution volume) produced significantly higher number of

colonies than the passaging of 10 µL (in 10 mL dilution volume); the latter yielding

comparable colonies to that of the 10-fold and 100-fold series performed in either dilution

volumes. It correlates to our previous understanding of the complexities associated with 1 µL

pipetting. Overall, this suggests that the dilution volume used in serial dilution doesn't affect

the accuracy of bacterial enumeration, given that 1 µL pipetting is avoided.

Table 4: Comparison of the colony numbers (along with standard deviations) of *E. coli*

obtained from each of the dilution series performed for 1 mL and 10 mL dilution volume,

generated from same culture.

Average colonies per spot											
Dilution factors	\multicolumn 10⁶						**10⁷**					
Dilution series	**10 fold**		**100 fold**		**1000 fold**		**10 fold**		**100 fold**		**1000 fold**	
Dilution ratios	**1:9 (A)**		**1:99 (B)**		**1:999 (C)**		**1:9 (A)**		**1:99 (B)**		**1:999 (C)**	
Final volume of dilutions	**1 mL**	**10 mL**	**1 mL**	**10 mL**	**1 mL**	**10 mL**	**1 mL**	**10 mL**	**1 mL**	**10 mL**	**1 mL**	**10 mL**
Experiment 1	$21.9^* \pm 3.0$	$18.1^* \pm 3.4$	19.7 ± 3.3	19.1 ± 3.4	$46.9^* \pm 5.3$	$19.9^* \pm 2.6$	2.6 ± 1.1	2.4 ± 1.3	2.5 ± 1.3	1.9 ± 1.0	$5.6^* \pm 2.1$	1.9 ± 1.3

Experiment 2	21.8 ± 5.1	20.6 ± 4.1	28.3* ± 5.4	20.5* ± 4.6	62.7* ± 8.9	27.7* ± 2.8	2.4 ± 1.9	2.6 ± 1.7	2.7 ± 2.2	2.0 ± 1.3	6.6* ± 2.4	2.8 ± 1.7
Experiment 3	27.6 ± 3.0	26.3 ± 3.9	36.5 ± 5.1	33.5 ± 4.4	86.9* ± 7.4	36.8* ± 7.9	3.0 ± 1.2	2.9 ± 1.3	5.7* ± 3.0	3.2* ± 1.1	15.7 ± 3.4	4.3* ± 1.3

Asterisk (*) denotes a significant difference ($U_{statistic} < U_{critical}$) between the colony numbers obtained with dilutions for 1 mL and 10 mL volumes, as calculated from a two-tailed Mann-Whitney U-Test at 95% confidence interval.

(a) 10^6-fold dilution plating

(b) 10^7-fold dilution plating

Fig. 5: Comparison of colony numbers of *E. coli* generated from same culture but attained using two different diluent volumes of serial dilution in order to adjust to 1 mL and 10 mL volumes. Plating at (a) 10^6- and (b) 10^7-fold dilution was done. Each petri-plate corresponds

439 to a specific dilution series where comparison is made between upper and lower halves

440 containing colonies from 1 mL and 10 mL dilution volumes respectively. In each of the case,

441 colony numbers observed with these dilution volumes appears similar for 10-fold and 100-

442 fold series. However a contrasting difference was observed with 1000-fold series, where a

443 clearly greater number of colonies were observed in the case of 1 mL dilution volume.

4. Discussion

In serial dilution process to reach a certain dilution fold, specific sample volume is

passaged stepwise to specific diluent volume. Our study first demonstrated that a higher

sample volume is a better representation of bacterial population size than a lower sample

volume, evident from the values of coefficient of variation (CoV) recorded (Table 1). Adding

to it, a strong negative correlation was also observed between sample volumes plated and

CoV observed. This was in concordance with the view of random nature of bacterial

distribution in culture. In this regard, a higher sample volume during serial dilution is

expected to better contain the randomness of population and give lesser anomalies in cell

density being passaged in the successive steps. But it also leads to a lower fold dilution in a

single passage, thereby requiring more number of passages to reach the desired dilution fold.

This increases the complexity of the process. Alternately, passaging a lower sample volume

leads to a higher fold dilution in a single step and requires fewer numbers of passages to

reach the desired dilution fold. This reduces the complexity of the process. So, although

passaging of a higher sample volume seems advantageous considering the random

distribution of bacteria but is disadvantageous considering the complexity of the process and

alternately, passaging of a lower suspension volume seems disadvantageous considering the

random distribution of bacteria but is advantageous considering the complexity of the

process. In order to know whether passaging a higher or lower sample volume is more

favourable for an accurate bacterial enumeration, a critical examination of 10-, 100- and

1000-fold dilution series was carried out in the current study. Accordingly, sample to diluent

ratio (dilution ratio) was highest at 1:9 in a 10-fold series, followed by 1:99 in a 100-fold

series, and lowest at 1:999 in a 1000-fold series. The studies were performed with two

different bacterial species: *E. coli* DH5α and *R. pseudosolanacearum* F1C1, exhibiting two

different colony morphologies, with colonies of the former one being non-mucoidal, and that

of the latter being mucoidal, exhibiting twitching motility and possessing a greater tendency

to merge with other such colonies. The study revealed that all the three dilution ratios (1:9,

1:99 and 1:999) were similar with regards to bacterial enumeration and in practice all of them

are reliable. However, it is recommended to use higher diluent volume to perform a higher fold dilution. Example, in our study, a 1000-fold dilution series proved good when performed to a final 10 mL volume, but inappropriate when performed to the final 1 mL volume. The inaccuracies observed in this case were perhaps attributed to passaging a lower sample volume, *i.e.*, 1 µL, which is replaced by 10 µL upon scaling up the final volume 10-times. Passaging 1 µL was found to result in significantly higher and inconsistent colony counts compared to passaging 100 µL and 10 µL. It might be attributed to the requirement of a more sophisticated handling. Furthermore, similar enumerations observed with the three dilution series upon eliminating 1 µL pipetting might be because the advantage of a using a higher sample volume in 10-fold series was balanced by the disadvantage of a more number of passages involved; similarly the advantage of a lesser number of passages in 1000-fold series was balanced by the disadvantage of using a lower sample volume; and the 100-fold series being moderate in both respect. But considering the time, cost and labor efficiency of the approach with no compromise on accuracy of enumeration, a 1000-fold dilution series shall be preferred when performing the dilutions for 10 mL final volume and a 100-fold dilution series shall be preferred when performing the dilutions for 1 mL final volume. This is because a relatively lower number of passages are involved in these processes compared to the 10-fold series. The study was carried forward to understand if diluent volumes used for serial dilution affects bacterial enumeration. For this, each of the 10-, 100- and 1000-fold dilution series were performed using two different diluent volumes to maintain the final volume of 1 mL and 10 mL respectively. Results revealed that the diluent volume used for dilutions doesn't affect CFU estimation. It is only the passaging of a lower sample volume (such as 1 µL) that bring in-congruencies and yield a higher than expected CFU count.

Besides highlighting the accuracies and complexities of various dilution series/ratios to make a conclusive comment on the practice that shall be followed or avoided, and understanding the impact of diluent volume on bacterial enumeration, our study offers a clear demonstration on the random distribution pattern of bacteria. Had it been uniform, all the sample volumes plated would have been equally consistent in terms of colony count. Whereas, the findings of our study that proves a higher sample volume to be more consistent (due to low CoV observed) in terms of bacterial count, is a clear indicative of randomness prevalent in the population that can't be contained in a smaller sampling size. On this note, more the sampling size required for bringing consistency in colony count, more the degree of randomness prevalent in the bacterial distribution. Our study serves a good and easy model

(in terms of reproducibility and cost efficiency) for students who want to observe the random nature of bacterial distribution. The advantage here is that a greater number of replicates (10 or even more) can be obtained in just a single solid agar plate, and a greater number of sampling is always better for addressing the randomness prevalent in the population. Many-a-times, researchers performing microbiological experiments evaluate the average number of colonies in a sampling population from triplicates, generally obtained from spread-plate method, which requires three separate solid agar plates. Even sometimes, due to less understanding of the mathematics of bacterial distribution, students performing the experiment consider singular plating for estimating cell density. The randomness in bacterial distribution can't be contained in just a single replicate. Hence, taking one replicate or three proves insufficient for getting precise cell density estimate of bacteria. In this context, our study by virtue of a greater number of sampling in a single plate becomes advantageous for depicting the randomness if any is prevalent in a population. Further, the precision of pipettes or the sample-adherence ability of tips is some of the additional aspects that could be examined by this reproducible study.

Funding

The authors would like to acknowledge DBT for the MSc project grants for the consumables.

Conflict of interest

The authors declare no conflict of interest in the subject matter discussed in this manuscript.

Acknowledgement

MJ is thankful to Tezpur University for the institutional fellowship. S Begum and SJG is thankful for the JRF fellowship from the DBT, GoI New Delhi grant (BT/PR41637/NER/95/1753/2021) awarded to SKR and also to Tezpur University for the institutional fellowship. S Bhuyan is thankful for the JRF/SRF fellowship from the UGC-NFSC, GoI, New Delhi. LD is thankful for the DST-INSPIRE JRF/SRF fellowship. SKR is thankful to Prof. Ramesh V. Sonti, Director, ICGEB New Delhi and Prof. Anil Tripathi, Director, IISER Mohali for teaching the fundaments of serial dilution.

References

Ben-David, A., Davidson, C. E., 2014. Estimation method for serial dilution experiments. J. Microbiol. Methods. 107, 214-221. https://doi.org/10.1016/j.mimet.2014.08.023

Bhuyan, S., Boruah, L., Jain, M., Begum, S., Giri, S. J., Dutta, L., Mandal, M., Ray, S. K., 2024a. Morphology of softening and translucency in the stems of tomato and eggplant seedlings due to infection by *Ralstonia pseudosolanacearum* F1C1. bioRxiv, 2024-09. https://doi.org/10.1101/2024.09.10.612199

Bhuyan, S., Dutta, L., Begum, S., Giri, S. J., Jain, M., Mandal, M., Ray, S. K., 2024b. A study on twitching motility dynamics in *Ralstonia solanacearum* microcolonies by live imaging. J. Basic Microbiol. 64, 42-49. https://doi.org/10.1002/jobm.202300272

Bhuyan, S., Jain, M., Boruah, L., Tara, T. S., Begum, S., Dutta, L., Giri, S. J., Phukan, T., Kabyashree, K., Mandal, M., Ray, S. K., 2024c. Tissue-dependent nature of plant susceptibility: a comparative pathogenicity study of the systemic phytopathogen *Ralstonia pseudosolanacearum* in eggplant and tomato seedlings through root and leaf. bioRxiv, 2024-11. https://doi.org/10.1101/2024.11.25.625158

Bhuyan, S., Yadav, M., Giri, S. J., Begum, S., Das, S., Phukan, A., Priyadarshani, P., Sarkar, S., Jayswal, A., Kabyashree, K., Kumar, A., Mandal, M., Ray, S. K., 2023. Microliter spotting and micro-colony observation: A rapid and simple approach for counting bacterial colony forming units. J. Microbiol. Methods. 207, 106707. https://doi.org/10.1016/j.mimet.2023.106707

Boucher, C. A., Barberis, P. A., Trigalet, A. P., Demery, D. A., 1985. Transposon mutagenesis of *Pseudomonas solanacearum*: isolation of Tn 5-induced avirulent mutants. Microbiology. 131, 2449-2457. https://doi.org/10.1099/00221287-131-9-2449

Hollinger, F.B. (Ed.), 1993. AIDS Clinical Trials Group Virology Manual for HIV Laboratories. NIH (NIAID Division of AIDS), Rockville MD, pp. MIC-1–MIC-5.

Kumar, R., Barman, A., Jha, G., Ray, S. K., 2013. Identification and establishment of genomic identity of *Ralstonia solanacearum* isolated from a wilted chilli plant at Tezpur, North East India. Curr. Sci. 1571-1578. https://www.jstor.org/stable/24098856

MacLowry, J. D., Jaqua, M. J., Selepak, S. T., 1970. Detailed methodology and implementation of a semiautomated serial dilution microtechnique for antimicrobial susceptibility testing. Appl. microbiol. 20, 46-53. https://doi.org/10.1128/am.20.1.46-53.1970

Reynolds, J., 2005. Serial dilution protocols. American Society for Microbiology: Washington, DC, USA. 1-7.

Taswell, C., 1984. Limiting dilution assays for the determination of immunocompetent cell frequencies.III. Validity tests for the single-hit Poisson model. J. Immunol. Methods. 72, 29–40. https://doi.org/10.1016/0022-1759(84)90430-7

Thomas, P., Sekhar, A. C., Upreti, R., Mujawar, M. M., Pasha, S. S., 2015. Optimization of single plate-serial dilution spotting (SP-SDS) with sample anchoring as an assured method for bacterial and yeast cfu enumeration and single colony isolation from diverse samples. Biotechnol. Rep. 8, 45-55. https://doi.org/10.1016/j.btre.2015.08.003

Figure Legends

Fig. 1: Three different ways of attaining a 10^6 fold dilution has been shown *viz* a 10-, 100- and 1000-fold dilution series. (i) A 10-fold series (1:9 dilution ratio) involves passaging 100 μL suspension to 900 μL saline in six successive dilutions. (ii) A 100-fold series (1:99 dilution ratio) involves passaging 10 μL suspension to 990 μL saline in three successive dilutions. (iii) A 1000-fold series (1:999 dilution ratio) involves passaging 1 μL suspension to 999 μL saline in two successive dilutions. Bacterial culture is shown in yellow in the screw caped bottle. Diluent (saline) is shown in blue in micro-centrifuge tubes. Created with Biorender.com.

Fig. 2: Correlation between suspension volume spotted (X- axis) and coefficient of variation (CoV) obtained among the colony numbers (Y-axis) of (a) *E. coli* and (b) *R. pseudosolanacearum*. Pearson's correlation coefficient (r) is represented for each case. A strong negative correlation has been observed in each case.

Fig. 3: *E. coli* colonies at selected fold dilutions (10^6 and 10^7) obtained with the different dilution series. These series were performed first to a final volume of (a) 1 mL and later to a final volume of (b) 10 mL from independent cultures. At 1 mL dilution volume, a clearly

greater number of colonies were observed with the 1000-fold series; however, at 10 mL dilution volume, similar numbers of colonies were observed with all the three dilution series.

Fig. 4: *R. pseudosolanacearum* colonies at selected fold dilutions (10^7 and 10^8) obtained with the different dilution series. These series were performed first to a final volume of (a) 1 mL and later to a final volume of (b) 10 mL from independent cultures. At 1 mL final volume, a clearly greater number of colonies were observed with the 1000-fold series; however, at 10 mL final volume, similar numbers of colonies were observed with all the three dilution series.

Fig. 5: Comparison of colony numbers of *E. coli* generated from same culture but attained using two different diluent volumes of serial dilution in order to adjust to 1 mL and 10 mL volumes. Plating at (a) 10^6- and (b) 10^7-fold dilution was done. Each petri-plate corresponds to a specific dilution series where comparison is made between upper and lower halves containing colonies from 1 mL and 10 mL dilution volumes respectively. In each of the case, colony numbers observed with these dilution volumes appears similar for 10-fold and 100-fold series. However a contrasting difference was observed with 1000-fold series, where a clearly greater number of colonies were observed in the case of 1 mL dilution volume.

Dilutions adjusted to 1 mL

(a)

Sampling of *E. coli* DH5α population

y = -0.007x + 0.205
R² = 0.9459

Correlation coefficient (r) = -0.97

(b)

Sampling of *R. pseudosolanacearum* F1C1 population

y = -0.0172x + 0.52
R² = 0.9809

Correlation coefficient (r) = -0.99

(a)

10^6 DF		10^7 DF
I	10-fold series	I
II	100-fold series	II
III	1000-fold series	III

1 mL

1 mL

(b)

10^6 DF		10^7 DF
I	10-fold series	I
II	100-fold series	II
III	1000-fold series	III

10 mL

10 mL

(a)

10⁷ DF — 10⁸ DF

10-fold series
100-fold series
1000-fold series

1 mL — 1 mL

(b)

10⁷ DF — 10⁸ DF

10-fold series
100-fold series
1000-fold series

10 mL — 10 mL

(a) 10^6-fold dilution plating

(b) 10^7-fold dilution plating