Constructing a holistic map of cell fate decision by hyper solution landscape

Xiaoyi Zhang,[1] Zhiyuan Li,[1,2,*] and Lei Zhang[1,3,4,5**]

[1] Center for Quantitative Biology, Peking University, Beijing, 100871, China

[2] Peking-Tsinghua Center for Life Sciences, Peking University, Beijing 100871, China

[3] Beijing International Center for Mathematical Research, Peking University, Beijing 100871, China

[4] Center for Machine Learning Research, Peking University, Beijing 100871, China

[5] Lead contact

[*]Correspondence: zhiyuanli@pku.edu.cn

[**]Correspondence: zhangl@math.pku.edu.cn

Summary

The Waddington landscape metaphor has inspired extensive quantitative studies of cell fate decisions using dynamical systems. While these approaches provide valuable insights, the intrinsic nonlinear complexity and the parameter dependence limits systematic analysis of fate transitions. Here, we introduce the Hyper Solution Landscape (HSL), a parameter-independent methodology showing a comprehensive structure of all possible landscape configurations for gene regulatory networks. Building on the concept of solution landscape that primarily captures the complete stationary points in static landscape, HSL connects different solution landscapes to reflect dynamic changes of the landscapes associated with bifurcations. Applied to the Cross-Inhibition with Self-activation motif, HSL analysis identifies key hyperparameters driving distinct directional changes in cell fate propensity. Importantly, different routes through the HSL between the same initial and final states can produce markedly different fate distributions. This enables rational design of transition strategies. We validate HSL's utility in the seesaw model of cellular reprogramming, establishing a powerful framework for understanding and engineering cell fate decisions.

Key words:

cell fate decision, Waddington landscape, bifurcation, gene regulatory network, solution landscape, hyper solution landscape

Introduction:

The Waddington landscape has served as a powerful metaphor for understanding cell fate decisions since its inception[1], catalyzing extensive research in developmental biology[2,3]. Following Kauffman's pioneering work using Boolean networks to characterize the valleys of this landscape[4,5], researchers have made significant strides in quantifying cell fate decisions through dynamical systems, particularly gene regulatory networks (GRNs) [6–9]. This quantitative framework has established a crucial link between mathematical attractors and biological cell

40 fates, enabling computational studies of cellular decision-making through nonlinear
41 dynamics[10–12].
42
Among the various regulatory motifs studied, the Cross-Inhibition with Self-activation (CIS)
pattern has emerged as particularly significant[13–17], appearing in numerous biological contexts
including GATA1-PU.1[18], OCT4-CDX2[15], and NANOG-GATA6 interactions[19]. Cell state
transitions within these networks can occur through two primary mechanisms: stochastic
fluctuations (noise-driven)[20–23] or changes in external signals (signal-driven)[24–28]. These
biological processes correspond to static and dynamic landscapes, respectively[29].
49
While nonlinear dynamics models have proven valuable in characterizing cell fate decisions,
they face a fundamental limitation: parameter dependency. Recent advances in nonequilibrium
statistical physics have enabled the construction of energy landscapes from dynamical
systems[30–34]. However, experimental determination of all relevant parameters in a gene
regulatory network remains challenging, if not impossible, and different parameter sets can lead
to distinct landscapes with varying predictions on cell fate behaviors[35]. Furthermore, the
complexity of energy landscapes, which contain numerous details beyond the primary
attractors, makes systematic analysis of global changes particularly challenging. While recent
geometric approaches to cell fate decisions have provided valuable insights[36–39], their abstract,
gene-free nature has limited their direct application to real genetic networks[40].
60
The solution landscape concept offers a promising alternative approach[41]. This framework
creates a pathway map comprising all stationary points and their connections[42], having
demonstrated utility across various fields including gene regulatory networks[43], liquid crystals[44],
and quasicrystals[45]. Unlike traditional energy landscapes that expend computational resources
on fine details, solution landscapes focus on the essential elements of noise-driven cell fate
decisions: the attractors (stable cell fates), saddle points (unstable transition states) and their
connections (transition pathways). Their simplified graph structure of nodes and edges provides
an efficient framework for analyzing static landscape and transitions driven by stochastic
fluctuations. However, experimental manipulation of cell fates typically falls into the signal-
driven category, where external signals alter the landscape geometry[43,46]. In these cases, the
solution landscape itself undergoes transformation as external conditions change[47]. This
dynamic nature calls for a more systematic approach to connect and analyze different solution
landscapes, enabling better understanding and control of directed cell fate decisions[48].
74
Building on this foundation, we propose a novel framework: Hyper Solution Landscape (HSL),
which is the ensemble of all solution landscapes and their bifurcation relationships for a
dynamic system. HSL is a non-parameter-dependent methodology that shows a
comprehensive structure of all possible landscape configurations from a given GRN topology.
By connecting different solution landscapes through bifurcation relationships, HSL provides a
unified analysis for both noise-driven and signal-driven cell fate decisions. We apply this
approach to both the CIS motif of differentiation and the seesaw model of reprogramming,
establishing clear connections between solution landscape alterations and the Waddington
landscape. Our analysis reveals intrinsic nonlinear complexity of cell fate decisions, including

the most probable fate distributions and their responses to varying hyperparameters. Furthermore, we demonstrate how different bifurcation routes lead to distinct cell fate behaviors, enabling the rational design of cell fate decision routes through HSL.

Results

Solution Landscape Depicts the Geometry of Cell Fate Decisions Under a Given Signal Level

The dynamics of cell fate decisions are often studied through mathematical models of gene regulatory networks. Among these, the Cross-Inhibition with Self-activation (CIS) motif has emerged as a particularly important regulatory structure[13] (Figure 1A). In this motif, two lineage-specifying transcription factors interact: factor X promotes its own expression while inhibiting factor Y, and vice versa. This regulatory architecture can be mathematically described by a system of ordinary differential equations (Method):

$$\frac{dx}{dt} = F_1(x, y) = \frac{r_x + r_1 k_1 x^{n_1} + r_2 k_2 y^{n_2} + r_3 k_3 x^{n_1} y^{n_2}}{1 + k_1 x^{n_1} + k_2 y^{n_2} + k_3 x^{n_1} y^{n_2}} - d_1 x$$
$$\frac{dy}{dt} = F_2(x, y) = \frac{r_y + r_4 k_4 y^{n_2} + r_5 k_5 x^{n_1} + r_6 k_6 x^{n_1} y^{n_2}}{1 + k_4 y^{n_2} + k_5 x^{n_1} + k_6 x^{n_1} y^{n_2}} - d_2 y$$

(1)

n_1 and n_2 represent transcription factor cooperation efficiencies, d_1 and d_2 are degradation rates, $k_1 - k_6$ denote relative binding strengths to cis-regulatory elements, and $r_x, r_y, r_1 - r_6$, represent transcription rates for different binding configurations. For analytical convenience, we can express this system in vector form:

$$\frac{d\boldsymbol{x}}{dt} = \boldsymbol{F}(\boldsymbol{x})$$

(2)

where the $\boldsymbol{x} = (x, y)$ and $\boldsymbol{F}(\boldsymbol{x}) = [F_1(x, y), F_2(x, y)]$.

In the CIS motif, four distinct cell fates emerge as stable gene expression patterns, corresponding to mathematical attractors (Figure 1B). These states mirror the 'valleys' in Waddington's landscape metaphor: two stem cell states—S1 (low expression of both X and Y) and S2 (high expression of both)—and two lineage-committed states—LX (high X, low Y) and LY (low X, high Y). Between these stable states lie saddle points, analogous to the "peaks" in Waddington's landscape, which separate different attractor basins. Of particular importance are the index-2 saddle points, which connect multiple index-1 saddle points and help define attractor basin boundaries.

By connecting all stationary points—both attractors and saddles—the solution landscape provides a complete geometric representation of all possible cell fate decisions under specific conditions[41]. The solution landscape is a hierarchical structure with all stationary points in different layers (Figure 1C). The General High-index Saddle Dynamics (GHiSD) method allows us to efficiently compute the index-k saddle points [42]:

$$\begin{cases} \dot{\boldsymbol{x}} = \left(I - 2 \sum_{j=1}^{k} \boldsymbol{v}_j \boldsymbol{v}_j^{\mathrm{T}} \right) \boldsymbol{F}(\boldsymbol{x}) \\ \dot{\boldsymbol{v}}_i = \left(I - \boldsymbol{v}_i \boldsymbol{v}_i^{\mathrm{T}} \right) J(\boldsymbol{x}) \boldsymbol{v}_i - \sum_{j=1}^{i-1} \boldsymbol{v}_j \boldsymbol{v}_j^{\mathrm{T}} \left(J(\boldsymbol{x}) + J^{\mathrm{T}}(\boldsymbol{x}) \right) \boldsymbol{v}_i, i = 1, \dots, k \end{cases}$$

Where $J(x) = \nabla F(x)$ is the Jacobian matrix of $F(x)$ and the v is the eigenvector of $J(x)$
corresponding to the positive eigenvalues.
The solution landscape reveals a static landscape under a given signal. For our two-
dimensional (2D) system, the highest index for saddles is 2, and the solution landscape can be
constructed hierarchically from the index-2 saddle point (repeller) down to the attractors using
GHiSD combined with downward/upward searching algorithms[42]. The resulting landscape for
the CIS motif reveals a single index-2 saddle point, four index-1 saddle points, and four stable
attractors, arranged in a hierarchical structure that corresponds to the intersections of nullclines
in state space (Figure 1D). By calculating the energy barrier (i.e. quasi-potential) between
different attractors by the GMAM[49,50] (Figure 1E and Method), the solution landscape offers a
novel perspective of noise-driven cell fate decision.
130
The Alteration of Solution Landscapes Corresponds to Distinct Geometry of Cell Fate
Decision
A solution landscape captures the complete set of stationary points—both saddles and
attractors—under specific parameter conditions of a system, representing all possible cell fates
and their transition relationships in a static landscape. In biological systems, varying signaling
strengths manifest as parameter changes in mathematical models. These parameter shifts can
alter the solution landscape, leading to changes in both the number of stable cell fates and the
likelihood of transitions between them.
Consider the signal-induced differentiation process from pluripotent stem cells (S2) to lineage-
specified cells (LX and LY) as an illustrative example. This process demonstrates how
parameter-induced changes in the solution landscape govern the geometry of cell fate
decisions: Initially, the solution landscape exhibits three stable attractors connected by two
index-1 saddle points (Figure 1E, top panel). This configuration represents a stem cell fate (S2)
positioned between two differentiated states (LX and LY). As the parameter k decreases from
5 to 2 (Figure 1E, middle panel), the landscape's topology remains unchanged, but the relative
stabilities shift significantly: the stem cell state S2 becomes less stable, while the differentiated
states LX and LY gain more stability. Further reduction of k to 1 triggers a subcritical fork
bifurcation, causing the stable attractor corresponding to the stem cell state to vanish, leaving
only the LX and LY attractors connected by a saddle point (Figure 1E, bottom panel).
These progressive changes in the solution landscape directly correspond to cellular behavior.
In the initial configuration, stem cells predominantly maintain their undifferentiated state due to
the higher stability of the S2 attractor (Figure 1G, top panel). As k decreases, the reduced
stability of the stem cell fate increases the probability of spontaneous differentiation (Figure 1G,
middle panel). Following the bifurcation, the topological transformation of the landscape drives
all stem cells toward differentiated fates (Figure 1G, bottom panel).
This analysis reveals how signal-induced changes in cell fate decisions manifest as geometric
transformations in solution landscapes. Notably, bifurcations create qualitative changes in cell
fate geometries, highlighting the need for tools capable of characterizing sequential changes in

solution landscapes. Such tools would enable systematic analysis of complex cell fate transition processes.

Figure 1. Solution landscape characterizes the geometry of cell fate decisions.

(A) The cross-inhibition and self-activation (CIS) gene regulatory motif in cell fate decisions. Gene X and Y represent two antagonizing lineage-specifying genes.

(B) Relationship between cell fates (i.e. attractors) and gene expression levels for the model in (A). Symbols S2, S1, LX, and LY represent stem cell fates with high and low lineage specifying genes, lineage X, and lineage Y cell fates, respectively. These symbols are used for all following figures.

(C) A solution landscape of the CIS motif in (A). The lowest layer of the solution landscape is composed of stable attractors represented by colored circles, which are labeled as LX, S2, S1, and LY to represent corresponding cell fates. The middle layer is composed of four index-1 saddle points represented by red crosses, and the top layer contains an index-2 saddle point represented by yellow triangle. The line in the figure indicates the connection between these points. These annotations are used for the following figures.

(D) The phase diagram of the CIS motif in (A). The coordinate axis represents the expression of X and Y genes, respectively. The black line and the red line represent nullclines of $\frac{dx}{dt} = 0$,

$\frac{dy}{dt} = 0$, respectively. The intersections of nullclines are the stationary points in the system,

with the same symbol as that in (C). Basins of each attractor is filled by their corresponding colors. These annotations are used for the following figures.

(E) Three solution landscapes during the signal-driven differentiation from stem cell to Lineage X and Lineage Y, corresponding to the red lines in (F). The height of each point is

determined by the quasi-potential. Different parameter k $(k = k_1 = k_4)$ corresponds to each

solution landscape is shown in the figure.

(F) The schematic Waddington landscapes for the differentiation shown in (E). In each curve,

the valleys represent attractors, and the blue, green and purple balls represent cells with LX,

S2 and LY fates, respectively. The red dashed line indicates the boundary of the S2 attractor.

The situation corresponding to the red line is shown in (E) and (G).

(G) The schematic representation of cell fate decisions under three different situations, as

corresponds in (E).

Hyper Solution Landscape (HSL): Ensemble of Solution Landscapes and Bifurcations

We introduce the HSL, defined as the ensemble of solution landscapes and their bifurcation

relationships, for a given gene regulatory network across all parameter sets. For the CIS motif,

the HSL takes the form of a hierarchical graph (Figure 2A), with nodes representing topologies

of solution landscapes and edges representing bifurcations. The nodes are arranged

horizontally based on their number of stationary points and assigned unique identifiers based

on their position in the graph.

The HSL can be constructed through two complementary approaches: an analytical method

based on Morse inequality (Method) and a computational method using sampling and

bifurcation analysis (Figure S1 and Method).

The former provides theoretical validation for the HSL structure. Morse inequality establishes a

fundamental relationship between stationary points and manifolds, specifically that the

difference between the number of even-order and odd-order saddle points equals one for the

manifolds studied here. This relationship persists through local bifurcations, constraining the

possible transitions between different landscape topologies. Furthermore, we observed that

local bifurcations without limit cycles (such as saddle-node and pitchfork bifurcations) follow a

consistent pattern: they typically involve the simultaneous creation or annihilation of adjacent-

index saddle points: index$-k$ saddle point and index$-k \pm 1$ saddle point simultaneously. This

feature offers a generation rule of simplified HSL (method and Figure S2), instructed by which

we can generate it by a process of adding or deleting nodes and edges.

The latter approach offers a practical algorithm for constructing the HSL. Given that the HSL is

inherently a graph composed of nodes and edges, the algorithm is divided into two main

components: identifying the nodes and connecting them via edges. The node identification

process involves several steps: sampling parameter sets, constructing the solution landscape

for each set, and clustering these landscapes (Figure S1 and Method). Enhanced sampling

may be necessary to ensure node completeness. Notably, prior knowledge, such as the

simplified HSL derived from Morse inequality, can significantly aid in node identification. The

second component of the algorithm involves calculating bifurcations by examining solution

landscapes across a range of parameters. If new nodes are discovered during bifurcation

analysis, the process returns to enhanced sampling; otherwise, the HSL is finalized.

HSL Enables Systematic Analysis of Parameter-Dependent Bifurcations.

The HSL framework provides comprehensive insights into cell fate transition behaviors.

Through extensive bifurcation analysis across all 16 parameters using randomized parameter

sets, we found that bifurcations occur more frequently between landscapes with fewer

stationary points (Figure 2B). Notably, the solution landscape topology labeled ID 13 emerges

as the most highly connected node in the HSL. This topology, representing a bistable landscape

with two differentiated states (LX and LY) separated by a saddle point, appears to be the most

probable landscape of the CIS motif, as evidenced by its frequent connections to other

topologies (Figure S2).

Parameter changes can trigger sequential transformations between solution landscapes,

represented as directional routes in the HSL. For example, increasing the parameter

r_x produces a systematic progression through different landscape topologies. In this

progression, certain topologies act as "sources" (ID41 and ID23, with multiple outgoing edges)

or "sinks" (ID2, with multiple incoming edges), revealing how parameter changes can

systematically eliminate certain attractors and bias the system toward specific cell fates (Figure

2C).

This hyperparameter framework provides a simplified yet powerful way to understand how

different parameters influence cell fate decisions. Analysis of parameter-induced changes

revealed that the 16 parameters naturally cluster into four categories, characterized by their

effects on X and Y gene expression levels (Figure 2D and Figure S2). We formalized these

categories as hyperparameters: $p_{x\_up}, p_{y\_up}, p_{x\_down}, p_{y\_down}$ (Figure 2D-E). This categorization

emerges directly from equation (1) by analyzing how dx/dt responds to parameter changes.

For instance, parameters r_x and r_1, which represent the basal expression and self-activation

of gene X respectively, belong to the p_{x_up} category as they both increase X gene expression.

Figure 2. HSL connects different solution landscape topologies by bifurcations.

(A) The HSL for the CIS motif shown in Figure 1(A). Individual solution landscape topology in each dotted box represents the node of HSL, and their IDs are shown above each box. Bifurcations between solution landscape topologies are edges of HSL.

(B) The HSL with weighted edges. Each solution landscape topology is symbolled by a square with the corresponding ID. The widths of the edges are positively correlated with the

number of parameters capable of inducing transformations between the linked landscape

topologies, in 100 randomized parameter sets.

(C) Transformations induced by r_x. As r_x increases, the transformation from one solution

landscape topology to another is indicated by arrows, with the arrow widths positively

correlated with the count of transformation in 100 randomized parameter sets.

(D) Relationship between the hyper parameters and their directions of cell fate changes. For

example, with the increase of hyper parameter P_{x_up}, the steady state of the solution

landscape bias towards the direction of S2 and LX. So as the other hyper parameters.

(E) Categorization of parameters into the four classes of hyper-parameters.

Different routes in HSL induce distinct cell fate decision behaviors.

The HSL provides a global framework for understanding signal-driven cell fate decisions. Each

cell fate transition corresponds to a transformation between nodes in HSL, which can occur

either through direct parameter changes ("direct route," blue in Figure 3A) or through a series

of sequential bifurcations along HSL edges ("gradual route," red in Figure 3A). Using the

differentiation of stem cells (transition from landscape ID 21 to ID 13) as a model system, we

discovered that different routes between the same initial and final landscapes can produce

significantly different cell fate outcomes.

To investigate these route-dependent behaviors, we conducted simulations modulating four

key parameters: basal expression levels (r_x, r_y) and self-activation levels (r_1, r_4) of genes X and

Y through a signal Δr. To introduce a realistic fate bias, we set the parameter changes for the

X lineage 1.1 times stronger than those for the Y lineage. Specifically, the parameter set

$\{r_x, r_y, r_1, r_4\}$ increased by $\{0.55, 0.5, 0.11, 0.1\}$ from initial to final states (Method).

Our first key finding reveals that the gradual route promotes more efficient differentiation toward

the LX fate compared to the direct route (Figure 3B). In stochastic simulations starting with

1,000 cells in the S1 attraction basin, the gradual route produced 30% more LX cells than the

direct route under identical noise conditions. This difference appears to arise from distinct

decision-making mechanisms: in the direct route, a cell's final fate strongly correlates with its

initial gene expression state, analogous to direct lineage commitment. In contrast, the gradual

route allows fate decisions to evolve with the changing landscape, similar to passing through a

priming stage before lineage commitment.

Furthermore, the HSL framework reveals multiple gradual routes between the same

landscapes (Figure 3C): a short route (ID21→ID13) and a long route

(ID21→ID31→ID22→ID13). These routes can be implemented by alternating increases in

basal expression (Δr_0) and self-activation levels (Δr_1) of both genes (Figure 3D): the long route

prioritizes increasing Δr_1 followed by Δr_0, while the short route reverses this sequence.

Remarkably, despite sharing initial and final landscapes, these routes produce substantially

different outcomes: simulations with 1,000 stem cells showed that the long route generated 2.5

times more LY cells than the short route (Figure 3E). This route-dependent bias can be

explained through calculating energy barriers (i.e. quasi-potentials). By comparing the energy

barriers from S1 to LY (E_{SY}) and S1 to LX (E_{SX}) along both routes, we found that during the

critical decision period near the bifurcation point, the long route presents an E_{SY} approximately

three times lower than the short route (Figure 3F and Figure S3). This reduced barrier makes

the LY fate more accessible to stem cells following the long route, demonstrating how route

choice can significantly influence cell fate outcomes even when starting and ending points
remain the same.
301

Figure 3. Different routes in HSL induce different cell fate decision behaviors.

(A) The schematics of directly change and gradually changes in landscapes. The symbols are
the same as the Figure 1. The gray shadows indicate the boundary of the stem cell attractors
in the upper landscape. The black dash lines indicate the position of the saddle points in the
bottom landscape.

(B) The number of LX cells induced by direct (blue) and gradual routes(red), respectively. The
parameter changes from the initial to the final in these two routes were shown in the upper
box. In the gradual route, parameters changes by the following rules: $r_1(t) = r_1(0) + 5.5 *$
$\Delta r(t)$, $r_4(t) = r_4(0) + 5 * \Delta r(t)$, $r_x(t) = r_x(0) + 1.1 * \Delta r(t)$, $r_y(t) = r_y(0) + \Delta r(t)$. The noise

level in simulations is set to 0.02. Simulations in each route are repeated for 1000 times to
calculate the standard deviation.
(C) Long route and short route in HSL. In the subset of HSL, the short and the long routes
from ID 21 to ID 13 solution landscape topologies are indicated by yellow and blue arrows,
respectively.
(D) Changes in parameters over time for different cell fate decision routes. Blue and yellow
lines correspond to the long and the short routes, respectively. The routes correspond to
alternatively increase the basal expression level $\Delta r_0(t)$ and the self-activation level $\Delta r_1(t)$.
The parameters are changed as the following rules: $r_1(t) = r_1(0) + 1.1 * \Delta r_1(t)$, $r_4(t) =$
$r_4(0) + \Delta r_1(t), r_x(t) = r_x(0) + 1.1 * \Delta r_0(t), r_y(t) = r_y(0) + \Delta r_0(t)$.
(E) The number of LY cells induced by different routes. The correspondence between colors
and routes is the same as (B). The noise level in simulations is set to 0.02. Simulations in
each route were repeated for 100 times to calculate the standard deviation.
(F) The schematic of the landscape near bifurcation. Symbols are the same as the Figure1.
The energy barrier of different route is shown in the corresponding color in (E). The energy
barrier calculated by GMAM is shown in the table.
328
Utilizing HSL to Design the Route of Cell Fate Decision.
The HSL provides a comprehensive framework for designing signal-driven cell fate transitions.
Consider the differentiation from stem cells to lineage-X (transition from landscape ID 31 to ID
2), where HSL reveals two distinct routes: a "progression route" (red dashed line, Figure 4A)
and an "accuracy route" (blue dashed line, Figure 4A), each with unique fate decision
characteristics [43]. In the progression route, the stem cell attractor disappears first, forcing cells
to exit the stem cell state and differentiate (Figure 4B). In contrast, the accuracy route first
eliminates the antagonistic LY fate, restricting cell fate choices to either maintaining stemness
or differentiating specifically to LX, thus ensuring higher differentiation precision.
These theoretical routes can be implemented through specific parameter modifications. Based
on our analysis in Figure 2, the progression route can be achieved through linear-like increases
in basal expression rate r_1, while the accuracy route requires increases in self-activation level
r_x (Figure 4B, Figure S4). To compare these routes quantitatively, we conducted stochastic
simulations starting with 1,000 cells in the S2 state. After initial equilibration, we tracked the
population dynamics along both routes, recording the number of cells in S2, LX, and LY cell
type over time (cell type defined by expression bias $[X] - [Y]$ with a 0.5 threshold) (Figure 4C)
To characterize route efficiency, we defined three key temporal indicators (Figure 4C). The
lineage X arrival time measures when the cell type LX population exceeds 900 cells, providing
a metric for successful differentiation. The lineage Y leaving time indicates when the cell type
LY population decreases below 90% of its maximum, representing clearance of the alternative
fate. Finally, the stem cell leaving time marks when the stem cell (S2) population drops below
100 cells, indicating exit from the stem cell state.
These metrics revealed distinct rate-limiting steps for each route. In the progression route, rapid
stem cell exit precedes other transitions, with lineage X arrival time closely matching lineage Y
leaving time—indicating that exit from lineage Y is rate-limiting. Conversely, in the accuracy

route, lineage X arrival closely tracks stem cell leaving time, suggesting that stem cell exit is
the primary bottleneck.
Further analysis across varying noise levels and initial landscapes revealed environment-
dependent route efficiencies. Initial landscape geometry, controlled by parameter shift Δr_{ini},
particularly affects energy barriers between stem and differentiated states (Figure S4). Heat
map analysis of temporal indicators across these conditions (Figure 4D-F) revealed several key
insights into route behavior. The overall efficiency of each route, measured by lineage X arrival
time, strongly correlates with its characteristic rate-limiting step. Under conditions of low noise
and low energy barriers, the progression route shows diminished performance as cells have
difficulty exiting the LY state. In contrast, the accuracy route faces its greatest challenges at
low noise levels combined with high energy barriers, where cells exhibit delayed exit from the
stem cell state, resulting in extended differentiation times.
These findings demonstrate how HSL can guide the design of effective differentiation
strategies by identifying and addressing route-specific bottlenecks in cell fate transitions.

Figure 4. Targeted design of the cell fate transition routes by HSL.

(A) Designs of the progression route and accuracy route for differentiation in the HSL of
Figure 2(A). We set the initial landscape as ID 23 and the final landscape as ID 2, and there
are two routes connecting them. The red arrows mark the "progression route" and the blue
arrows mark the "accuracy route".

(B) Schematic changes of the landscapes for the two routes in (A). Along the progression
route, with the increasing of r_1, the stem cell fate disappears first, then the LY fate
disappears. Along the accuracy route, with the increasing of r_x, the LY fate disappears first,
then the stem cell fate. The green, blue, and purple balls stand for the cells in stem cell,
lineage X, and lineage Y fates, respectively.

(C) Changes of the percentage of cell states along the progression (left panel) and accuracy
(right panel) routes. The green, blue and purple lines stand for the cells in stem cell, lineage X
and lineage Y states respectively. The cell type at each time is defined by the fate bias ($[X]$ −
$[Y]$) with the cutoff setting to 0.5. The noise level and initial parameter is set as indicated by
the green box in (D) and (E).

(D-F) The heatmap of lineage X arrival time (D), Stem cell leaving time (E) and lineage Y
leaving time (F) in different noise levels and initial parameters for progression and accuracy
routes. The initial parameter is set as following: $r_x = r_x^0 + \Delta r_{ini}, r_y = r_y^0 + \Delta r_{ini}, r_1 = r_1^0 +$
$\Delta r_{ini}, r_4 = r_4^0 + \Delta r_{ini}$, where $\{r_x^0, r_y^0, r_1^0, r_4^0\} = \{0.15, 0.15, 0.95, 0.95\}$.

The Application of HSL in Seesaw Model

The HSL framework can be extended to various network structures. A particularly significant
application lies in the "seesaw" model of cellular reprogramming (Figure 5A), which
demonstrates how maintaining regulatory balance is crucial for restoring pluripotency[51]. In this
model, pluripotency factors OCT4 and SOX2 activate each other while promoting their
respective lineages, while lineage-specifying gene groups MEs and ECTs exhibit mutual
antagonism and suppress pluripotency (Figure 5A). We constructed the HSL using eight key
parameters from the original model: external signals that either induce (C_0, C_S, C_M, C_E) or inhibit
(I_O, I_S, I_M, I_E) the expression of OCT4, SOX2, MEs, and ECTs, respectively.
The resulting HSL reveals that cellular reprogramming primarily follows two routes: ID 8→ID
6→ID 2 or ID 8→ID 4→ID 2 (Figure 5B, Figure S5). Both paths begin from a three-attractor
landscape (ID 8 topology) containing the ME fate and culminate in a landscape containing only
the stem cell attractor. Even between the same initial (ID 8) and intermediate (ID 6) landscapes,
different parameter modifications can induce distinct reprogramming behaviors.
When ME fate suppression dominates (through increased I_M), the landscape transforms
directly from ID 8 to ID 6. However, when ECT fate activation prevails (through increased C_E),
the transition proceeds via an intermediate ID 9 landscape (Figure 5C). These distinct routes
produce markedly different fate outcomes: ME suppression forces direct reprogramming to
stem cells (Figure 5D and 5F), while ECT activation enables both reprogramming to stem cells
and transdifferentiation to ECT cells (Figure 5E and 5G).
These divergent fate distributions arise from different topological connections between
attractors. In the ID 9 landscape, ME cells connect to ECT cells through an index-1 saddle,
while in ID 8, ME cells connect directly to the stem cell state. As signaling increases, the ME
attractor gradually destabilizes, enhancing transition probability to adjacent states according to
the landscape's topology.
This application of HSL to the seesaw model not only demonstrates the framework's versatility
but also reveals how different routes through the landscape can fundamentally alter cell fate

outcomes. These insights suggest that careful consideration of transition routes, not just initial

and final states, is crucial for optimal control of cell fate engineering.

Figure 5. HSL of the seesaw model reveals how different routes influences fate distributions.

(A) The topology of the gene regulatory network underling the original seesaw model[51]. Color schemes and the symbols are the same as Figure1(A).

(B) The HSL for the seesaw model. The symbols are the same as Figure2(A). The pink lines between ID 8 and ID 9 (ID 7) represent heteroclinic flip bifurcations. The cell types are distinguished by the stemness and the bias (Method).

(C) Schematics on how different parameter changes induce different landscape transformations, forming different routes. The upper route corresponds to the change of solution landscape topologies from ID 8 to ID 6, induced by the I_M linear decreasing over time. The bottom route corresponds to the change from ID 8 to ID 9 and then to ID 6, induced by the C_E linear increasing over time.

(D-E) The distribution of cell types at different time for 8-6 route(D) and 8-9-6 route(E). The green, blue and purple bars stand for the cells in stem cell, lineage X and lineage Y states respectively. 1000 cells are set in ME state in ID 8 solution landscape at initial and reach the ID 6 solution landscape finally, with different parameters linear changing over time.

(F-G) The distribution of cell types at different parameter situation for 8-6 route(F) and 8-9-6 route(G). Dots with different colors represent cells at different parameter situation. The cells are clustered by Gaussian mixture method.

Discussion

The quest to understand cell fate decisions through landscape perspectives has evolved significantly since Waddington's initial metaphor. We construct a holistic map of cell fate decision by applying a HSL approach. HSL incorporates fundamental principles of landscape transitions, particularly how local bifurcations involve the simultaneous creation and annihilation of adjacent-order stationary points. Through systematic analysis of the CIS network, we identified hub nodes within the HSL and consolidated the numerous parameters into four key hyperparameters that characterize directional landscape changes. This framework demonstrates how different routes through the landscape can produce distinct cell fate distributions, providing a global perspective for designing cell fate transitions with specific properties, such as optimizing for either progression efficiency or fate accuracy.

The solution landscape approach offers several advantages over traditional methods. Unlike energy landscapes constructed through stochastic simulations[30], solution landscapes directly extract stationary points and their connections from gene regulatory networks using efficient saddle dynamics algorithms. This approach eliminates the need for extensive stochastic simulations, particularly beneficial for high-dimensional systems where reaction coordinates are difficult to obtain.

The HSL provides a unified framework for investigating both noise-driven and signal-driven cell fate decisions, each corresponding to distinct analytical scales. At the node level, each node (i.e. solution landscape) represents a static landscape, enabling the study of noise-driven effects through the extraction of transition states and transfer pathways. Conversely, at the edge level, the connections between different solution landscapes reflect dynamic landscape changes associated with bifurcations. This allows us to explore signal-driven behaviors and how cells adapt to varying environmental stimuli. Overall, the HSL enhances our understanding of the intricate interplay between noise and signals in cellular fate decision, positioning it as a valuable tool for advancing insights into complex biological processes.

The practical utility of HSL lies in its ability to guide the design of cell fate decision routes. Acting as a holistic map, it enables identification of optimal paths to desired cell states. Our analysis of progression and accuracy routes revealed distinct rate-limiting steps, providing crucial insights for route design in real systems. With advancing capabilities in synthetic biology[52–54],

these theoretically designed routes can be experimentally implemented. Future applications
could extend to optimizing pathways under multiple constraints to meet diverse functional
requirements.
While this work establishes HSL as a powerful framework, several challenges remain for future
investigation. The theory and algorithms of HSL need further refinement for application to more
complex systems, particularly those exhibiting dynamic behaviors like limit cycles[32].
Additionally, validation in more realistic biological systems is needed to bridge the gap between
theoretical predictions and experimental observations. The potential applications of HSL extend
beyond cellular systems to other dynamical systems, such as ecosystem stability and microbial
community dynamics, opening new avenues for future research.
This global perspective on landscape transitions not only advances our understanding of cell
fate decisions but also provides practical tools for cellular engineering. As experimental
techniques continue to evolve, the insights gained from HSL analysis will become increasingly
valuable for designing precise cellular programming strategies.
482

ACKNOWLEDGMENTS
We would like to thank G. Xue for his suggestions on the biological significance, J. Yin and H.
Su for their advice on mathematical methods, and N. Yang. and Y. Yu. for their support in
theoretical concepts. We also extend our gratitude to all members of Lei's group for their
support and assistance. L.Z. was supported by the National Natural Science Foundation of
China (No.12225102, T2321001, and 12288101) and the National Key Research and
Development Program of China (No. 2021YFF1200500). Z.L. was supported by the National
Natural Science Foundation of China (No. T2321001) and the National Key Research and
Development Program of China (No. 2021YFA0910700). This work supported in part by
Peking-Tsinghua Center for Life Sciences.
493

494   **DECLARATION OF INTERESTS**
495   The authors declare no competing interests.
496

497   **AUTHOR CONTRIBUTIONS**
498   L.Z. and X.Z. conceived the project; X.Z. performed the analysis and computational simulations;
499   Z.L. and L.Z. supervised the project; X.Z., Z.L. and L.Z. wrote the paper.
500

501   **STAR★METHODS**

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Software and algorithms		
MATLAB r2022b	The Mathworks Inc.	https://matlab.mathworks.com/

METHOD DETAILS
Gene Regulatory Network.
The GRN used in this work follows previous models[43,55]. The equation (1) derives from a
biochemical reaction model incorporating dimerization, reversible DNA binding, transcription,

translation, decay, and ectopic overexpression[55]. The original 14-dimensional system was
reduced to equation (1) using quasi-steady state assumptions.
509
510 **Generalized High-index Saddle Dynamics (GHiSD).**
511 k-GHiSD identifies index-k saddle points in dynamical systems[42,56], represented by dynamics
512 (3). This method can be interpreted as a reverting evolution along the unstable subspace
513 $\mathcal{W}^{\mathrm{u}}(x)$:

$$\dot{x} = -\mathcal{P}_{\mathcal{W}^{\mathrm{u}}(x)} F(x) + \left(F(x) - \mathcal{P}_{\mathcal{W}^{\mathrm{u}}(x)} F(x) \right)$$

515 Where $\mathcal{P}_{\mathcal{W}^{\mathrm{u}}(x)}$ stands for the orthogonal projection operator onto the unstable subspace
516 $\mathcal{W}^{\mathrm{u}}(x)$.
517
518 **Algorithm for constructions of the solution landscape.**
519 Generally speaking, we can use the downward-search and the upward-search algorithms to
520 construct the solution landscape[41,44], which starting from the k-index saddle points and the
521 stable points respectively.
522
Downward search:

1. Given an k-index saddle x, calculate the unstable directions $\{v_1, \ldots, v_k\}$.
2. Choose an unstable direction v_j and perturb the initial state along this direction.
3. Start from the point $x \pm \varepsilon v_j$, find the m-index saddle point $(m < k)$ using the m-GHiSD.
4. Record the new saddle point, and the connection relationship.
5. Repeat the process for all of the unstable directions and all of the indices until no new
saddles can be found.

Upward search:
1. Given an k-index saddle x, calculate the stable directions $\{v_1, \ldots, v_p\}$.(suppose the
highest index of saddle points is $k + p$)
2. Choose a stable direction v_j and perturb the initial state along this direction.
3. Start from the point $x \pm \varepsilon v_j$, find the m-index saddle point $(m > k)$ using the m-GHiSD.
4. Record the new saddle point, and the connection relationship.
5. Repeat the process for all of the stable directions and all of the indices until find the
highest index saddles.
In practice, we use the upward search and the downward search algorithm together.
539
Morse inequality
Morse theory is a geometrical theory that studies the topological relationship between stationary
points and manifolds. For the dynamical system we studied, it satisfies the constrains in
reference[57] and the Morse inequalities for the dynamical system can be shown by the following
expression:
Let α_i be the number of stationary points of index i and β_i be the i-th Betti number (i.e.,
the rank of the i-th homology group). We have the following inequalities:

$$\alpha_0 \geq \beta_0$$
$$\alpha_1 - \alpha_0 \geq \beta_1 - \beta_0$$

549	$$\alpha_2 - \alpha_1 + \alpha_0 \geq \beta_2 - \beta_1 + \beta_0$$
550	$$\alpha_3 - \alpha_2 + \alpha_1 - \alpha_0 \geq \beta_3 - \beta_2 + \beta_1 - \beta_0$$
551	... ...

$$\alpha_2 - \alpha_1 + \alpha_0 \geq \beta_2 - \beta_1 + \beta_0$$

$$\alpha_3 - \alpha_2 + \alpha_1 - \alpha_0 \geq \beta_3 - \beta_2 + \beta_1 - \beta_0$$

$$... ...$$

The Betti numbers for $\mathbb{R}^2$ are as follows: $\beta_0 = 1, \beta_i = 0$ for $i \geq 1$.

In the context of a two-dimensional system, the maximum index of saddles is 2, indicating that

there are no saddles of index 3 or higher: $\alpha_j = 0 \; for \; j \geq 3$.

Therefore, the inequalities can be simplified to the following:

$$\alpha_0 \geq 1,$$

$$\alpha_1 - \alpha_0 \geq -1,$$

$$\alpha_2 - \alpha_1 + \alpha_0 \geq 1,$$

$$-\alpha_2 + \alpha_1 - \alpha_0 \geq -1.$$

The last inequality can be rewritten as:

$$\alpha_2 - \alpha_1 + \alpha_0 \leq 1.$$

Considering the last two inequalities together, we find that the number of saddles satisfies:

$$\alpha_2 - \alpha_1 + \alpha_0 = 1.$$

Since bifurcations do not alter the topological features of the manifold, this equation holds

both before and after bifurcations. In our study, we observe that bifurcations affect the number

of stationary points according to the following rules: k-index and $(k \pm 1)$-index saddles appear

and disappear simultaneously.

A framework of constructing simplified HSL.

We present a theoretical framework for constructing simplified hyper solution landscape

(simplified HSL), which primarily focuses on the topological structures of the solution

landscape (Figure S2). The methodology encompasses the following steps: 1) fixing the

starting and ending solution landscapes; 2) applying generation rules derived from Morse

inequalities to create the solution landscapes between them; and 3) pruning based on prior

knowledge.

In the specific application to the CIS network, our algorithmic process is detailed as follows:

1. Fixing Starting and Ending Points: The simplest solution landscape is characterized by

the presence of a single attractor. According to our prior knowledge, the most complex

solution landscape for the CIS network consists of a 2-saddle, four 1-saddles, and four

attractors (Figure 1C).

2. Generation Rules: Within the CIS network, the methods for expansion include two types:

adding a 2-saddle and a 1-saddle to the solution landscape, or adding a 1-saddle and a

0-saddle (attractor) to the solution landscape.

3. Modifications: By integrating prior knowledge of the CIS network, we eliminate structures

that lack biological significance.

The resulting simplified hyper solution landscape (simplified HSL) can then be utilized in the

subsequent construction of the HSL.

HSL Construction Algorithm

The flowchart for constructing the HSL (Figure S1) is organized into the following steps:

1.   Input Data: Begin by inputting the GRNs (or ODEs), along with the corresponding
parameter space.
2.   Initial Parameter Sampling: Conduct initial sampling of the parameter set using Latin
Hypercube Sampling or leveraging prior knowledge, such as a simplified HSL constructed
based on Morse inequality.
3.   Solution Landscape Generation: For the parameter sets identified in the previous step,
generate solution landscapes through both upward and downward searches.
4.   Clustering: Cluster the pool of solution landscapes into distinct topologies using
isomorphic graph methods and other clustering techniques.
5.   Enhanced Sampling: Perform enhanced sampling for topologies that exhibit insufficient
sampling to ensure comprehensive representation.
6.   Bifurcation Search: Execute a bifurcation search by generating solution landscapes
across a range of parameters to identify critical transitions.
7.   Output: If a new topology of solution landscape is discovered, return to step 5 for further
enhanced sampling; otherwise, proceed to output the final HSL.

Geometric Minimum Action Method (GMAM) and quasi-potential.

The GMAM[49,50] is a numerical algorithm used to compute the Minimum Action Path (MAP)

connecting two metastable states in a dynamical system. The key idea behind the GMAM is

to reformulate the Freidlin-Wentzell action functional on the space of curves. We apply the

GMAM method to calculate the transition pathways between two attractors x_1 and x_2 and

the quasi-potential $V(x_1, x_2)$ representing the energy barrier.

613
$$V(x_1, x_2) = \inf_{\varphi \in \bar{C}_{x_1}^{x_2}(0,1)} \hat{S}(\varphi)$$

614 Where $\hat{S}(\varphi) = \int_0^1 \langle \varphi', \hat{\vartheta}(\varphi, \varphi') \rangle d\alpha$ is the action calculated by GMAM.

616 **Stochastic simulation.**

617 We introduce noise into Equation (1) to obtain the Langevin equation[14,30], expressed as

618 follows:

619
$$\frac{d\mathbf{x}}{dt} = \mathbf{F}(\mathbf{x}) + \xi(t),$$

620 where $F(x)$ represents the right-hand side of Equation (1), and $\xi(t)$ denotes the Gaussian

621 white noise. The properties of the noise are given by:

622
$$\langle \xi(t) \rangle = 0, \langle \xi(t)\xi(t') \rangle = 2D\delta(t - t'),$$

where D is the noise level.

Range of parameter variation in constructing HSL

The range for $\{r_x, r_y, r_1, r_2, r_3, r_4, r_5, r_6\}$: 0-1 with the interval of 0.05

The range for $\{k_1, k_2, k_3, k_4, k_5, k_6, d_1, d_2\}$: the following multiplier to the initial parameter:

{0.0001,0.001,0.01,0.1,0.5,0.9,0.95,0.99,0.995,0.999,1,1.001,1.005,1.01,1.05,1.1,1.5,2,2.5,5,10}

Parameter sets for solution landscape.

The following parameter sets for Figure 1-4 are shown by the format:

$$\{r_x, r_y, r_1, r_2, r_3, r_4, r_5, r_6, k_1, k_2, k_3, k_4, k_5, k_6, d_1, d_2\}.$$

Figure1C and 1D:

$\{0, 0, 0.95, 0, 0, 0.95, 0, 0, 2.9045, 0.3818, 0.0073, 2.9045, 0.3818, 0.0073, 0.55, 0.55\}.$

Figure1E: $\{0.15, 0.15, 0.95, 0, 0.95, 0.95, 0, 0.95, k, 1.7, 1.8, k, 1.7, 1.8, 0.55, 0.55\}.$ k is shown in

the Figure.

Figure3: the initial parameter set $\{0, 0, 0.9, 0, 0, 0.9, 0, 0, 1.5, 0.8, 1, 1.5, 0.8, 1, 0.55, 0.55\}$ and the

final parameter set $\{0.11, 0.1, 1.45, 0, 0, 1.4, 0, 0, 1.5, 0.8, 1, 1.5, 0.8, 1, 0.55, 0.55\}.$

Figure4: the initial parameter set:

$\{0.15, 0.15, 0.95, 0, 0.95, 0.95, 0, 0.95, 2.1, 1.7, 1.8, 2.1, 1.7, 1.8, 0.55, 0.5\}.$

Figure5: the initial parameter set: $co = cs = cm = ce = 0; io = is = im = ie = 1; do = ds =$

$0.2; dm = de = 1; kpact = 0.35; kpinh = 0.7; kdinh = 0.3; ka = 1.3; ki = 0.55; npact =$

$1; npinh = 8; ndinh = 8; na = 9; ni = 10; w = 0.8; KM = 0.4;$

Classification criteria of cell types

For the CIS motif:

The attractors are distinguished by the expression level of X and Y gene (Figure 1B).

In the simulations, we can use the following two methods to define the cell types:

1. By the distance between the cell state and attractors with the cutoff setting to 0.5. If the

distance of cell state and a certain attractor is less than 0.5, we define the state as the

corresponding cell type.

2. By the fate bias ($[X] - [Y]$) of the cell state with the cutoff setting to 0.5, where the $[X]$ and

$[Y]$ represent for the expression level of gene X and gene Y, respectively. The cell type of

stem cell, LX and LY is corresponding to $-0.5 < [X] - [Y] < 0.5, [X] - [Y] \geq 0.5 \, and \, [X] -$

$[Y] \leq -0.5.$

In our simulation, the two methods have the same result.

For the seesaw model:

The cell types are distinguished by the stemness ($stemness = [Oct4] + [Sox2]$) and the bias

($bias = ([Oct4] + [MEs]) - ([Sox2] + [ECTs])$). The stem cell state has $stemness \geq 1$. The ME

state and the ECT state corresponds to the case of $stemnes < 1, bias > 0$ and $stemnes <$

$1, bias > 0$, respectively.

Reference

1. Waddington, C.H. (2014). The Strategy of the Genes (Routledge).

2. Bhattacharya, S., Zhang, Q., and Andersen, M.E. (2011). A deterministic map of Waddington's epigenetic landscape for cell fate specification. BMC Syst Biol *5*, 85.

3. Demirkıran, G. (2024). Construction of a Waddington-like landscape model that can guide clinical exploration of p53-dynamics-activating parameters in the face of divergent p53 dynamics. Communications in Nonlinear Science and Numerical Simulation *132*, 107893.

4. Kauffman, S. (1969). Homeostasis and Differentiation in Random Genetic Control Networks. Nature *224*, 177–178.

5. Kauffman, S.A. (1993). The Origins of Order: Self-Organization and Selection in Evolution (Oxford University Press).

6. Ferrell, J.E. (2012). Bistability, Bifurcations, and Waddington's Epigenetic Landscape. Current Biology *22*, R458–R466.

7. Li, C., and Wang, J. (2013). Quantifying Waddington landscapes and paths of non-adiabatic cell fate decisions for differentiation, reprogramming and transdifferentiation. J R Soc Interface *10*, 20130787.

8. Huang, S., Eichler, G., Bar-Yam, Y., and Ingber, D.E. (2005). Cell fates as high-dimensional attractor states of a complex gene regulatory network. Phys Rev Lett *94*, 128701.

9. Qiao, L., Zhao, W., Tang, C., Nie, Q., and Zhang, L. (2019). Network Topologies That Can Achieve Dual Function of Adaptation and Noise Attenuation. Cell Systems *9*, 271-285.e7.

10. Hong, T., Watanabe, K., Ta, C.H., Villarreal-Ponce, A., Nie, Q., and Dai, X. (2015). An Ovol2-Zeb1 Mutual Inhibitory Circuit Governs Bidirectional and Multi-step Transition between Epithelial and Mesenchymal States. PLoS Comput Biol *11*, e1004569.

11. Huang, S. (2013). Hybrid T-Helper Cells: Stabilizing the Moderate Center in a Polarized System. PLoS Biol *11*, e1001632.

12. Graf, T., and Enver, T. (2009). Forcing cells to change lineages. Nature *462*, 587–594.

13. Huang, S., Guo, Y.-P., May, G., and Enver, T. (2007). Bifurcation dynamics in lineage-commitment in bipotent progenitor cells. Developmental Biology *305*, 695–713.

14. Wang, J., Xu, L., Wang, E., and Huang, S. (2010). The Potential Landscape of Genetic Circuits Imposes the Arrow of Time in Stem Cell Differentiation. Biophysical Journal *99*, 29–39.

15. Zhou, J.X., and Huang, S. (2011). Understanding gene circuits at cell-fate branch points for rational cell reprogramming. Trends in Genetics *27*, 55–62.

16. Semrau, S., Goldmann, J.E., Soumillon, M., Mikkelsen, T.S., Jaenisch, R., and van Oudenaarden, A. (2017). Dynamics of lineage commitment revealed by single-cell transcriptomics of differentiating embryonic stem cells. Nat Commun *8*, 1096.

17. Liu, Y., Zhou, Z., Su, H., Wu, S., Ni, G., Zhang, A., Tsimring, L.S., Hasty, J., and Hao, N. (2024). Enhanced cellular longevity arising from environmental fluctuations. Cell Systems *15*, 738-752.e5.

18. Hoppe, P.S., Schwarzfischer, M., Loeffler, D., Kokkaliaris, K.D., Hilsenbeck, O., Moritz, N., Endele, M., Filipczyk, A., Gambardella, A., Ahmed, N., et al. (2016). Early myeloid

lineage choice is not initiated by random PU.1 to GATA1 protein ratios. Nature *535*, 299–
302.

19.　Simon, C.S., Hadjantonakis, A., and Schröter, C. (2018). Making lineage decisions with
biological noise: Lessons from the early mouse embryo. WIREs Developmental Biology
7, e319.

20.　Guillemin, A., and Stumpf, M.P.H. (2020). Noise and the molecular processes underlying
cell fate decision-making. Phys. Biol. *18*, 011002.

21.　Qiao, L., Zhang, Z.-B., Zhao, W., Wei, P., and Zhang, L. (2022). Network design principle
for robust oscillatory behaviors with respect to biological noise. eLife *11*, e76188.

22.　Wells, D.K., Kath, W.L., and Motter, A.E. (2015). Control of Stochastic and Induced
Switching in Biophysical Networks. Phys. Rev. X *5*, 031036.

23.　Coomer, M.A., Ham, L., and Stumpf, M.P.H. (2022). Noise distorts the epigenetic
landscape and shapes cell-fate decisions. Cell Systems *13*, 83-102.e6.

24.　Li, Y., Jiang, Y., Paxman, J., O'Laughlin, R., Klepin, S., Zhu, Y., Pillus, L., Tsimring, L.S.,
Hasty, J., and Hao, N. (2020). A programmable fate decision landscape underlies single-
cell aging in yeast. Science *369*, 325–329.

25.　Zhu, R., Del Rio-Salgado, J.M., Garcia-Ojalvo, J., and Elowitz, M.B. (2022). Synthetic
multistability in mammalian cells. Science *375*, eabg9765.

26.　Driskill, J.H., and Pan, D. (2023). Control of stem cell renewal and fate by YAP and TAZ.
Nat Rev Mol Cell Biol *24*, 895–911.

27.　Smart, M., Moreno, D.F., and Acar, M. (2024). Rationally reprogramming single-cell aging
trajectories and lifespan through dynamic modulation of environmental inputs. Cell
Systems *15*, 676–678.

28.　Zhang, J., Tian, X.-J., Zhang, H., Teng, Y., Li, R., Bai, F., Elankumaran, S., and Xing, J.
(2014). TGF-β-induced epithelial-to-mesenchymal transition proceeds through stepwise
activation of multiple feedback loops. Sci Signal *7*, ra91.

29.　MacArthur, B.D. (2022). The geometry of cell fate. Cell Systems *13*, 1–3.

30.　Wang, J., Zhang, K., Xu, L., and Wang, E. (2011). Quantifying the Waddington landscape
and biological paths for development and differentiation. Proc. Natl. Acad. Sci. U.S.A.
108, 8257–8262.

31.　Zhou, P., and Li, T. (2016). Construction of the landscape for multi-stable systems:
Potential landscape, quasi-potential, A-type integral and beyond. The Journal of
Chemical Physics *144*, 094109.

32.　Li, C., and Wang, J. (2014). Landscape and flux reveal a new global view and physical
quantification of mammalian cell cycle. Proceedings of the National Academy of Sciences
111, 14130–14135.

33.　Yao, M., Su, Y., Xiong, R., Zhang, X., Zhu, X.-M., Chen, Y.-C., and Ao, P. (2023). Energy
Landscape near Glioma Decipher the Role of Glioma Stem Cells within Glioma
Hierarchy. Preprint at bioRxiv.
https://www.biorxiv.org/content/10.1101/2023.11.24.568616v2

34.　Shi, J., Aihara, K., Li, T., and Chen, L. (2022). Energy landscape decomposition for cell
differentiation with proliferation effect. Natl Sci Rev *9*, nwac116.

35.　Li, C. (2018). Landscape of gene networks for random parameter perturbation. Integr.
Biol. *10*, 92–99.

36. Rand, D.A., Raju, A., Sáez, M., Corson, F., and Siggia, E.D. (2021). Geometry of gene
regulatory dynamics. Proc. Natl. Acad. Sci. U.S.A. *118*, e2109729118.
37. Raju, A., and Siggia, E.D. (2024). A geometrical model of cell fate specification in the
mouse blastocyst. Development *151*, dev202467.
38. Raju, A., and Siggia, E.D. (2023). A geometrical perspective on development.
Development, Growth & Differentiation *65*, 245–254.
39. Sáez, M., Blassberg, R., Camacho-Aguilar, E., Siggia, E.D., Rand, D.A., and Briscoe, J.
(2022). Statistically derived geometrical landscapes capture principles of decision-making
dynamics during cell fate transitions. Cell Systems *13*, 12-28.e3.
40. Corson, F., and Siggia, E.D. (2017). Gene-free methodology for cell fate dynamics during
development. eLife *6*, e30743.
41. Yin, J., Wang, Y., Chen, J.Z.Y., Zhang, P., and Zhang, L. (2020). Construction of a
Pathway Map on a Complicated Energy Landscape. Phys. Rev. Lett. *124*, 090601.
42. Yin, J., Yu, B., and Zhang, L. (2021). Searching the solution landscape by generalized
high-index saddle dynamics. Sci. China Math. *64*, 1801–1816.
43. Xue, G., Zhang, X., Li, W., Zhang, L., Zhang, Z., Zhou, X., Zhang, D., Zhang, L., and Li,
Z. (2024). A logic-incorporated gene regulatory network deciphers principles in cell fate
decisions. eLife *12*, RP88742.
44. Yin, J., Zhang, L., and Zhang, P. (2022). Solution landscape of the Onsager model
identifies non-axisymmetric critical points. Physica D: Nonlinear Phenomena *430*,
133081.
45. Yin, J., Jiang, K., Shi, A.-C., Zhang, P., and Zhang, L. (2021). Transition pathways
connecting crystals and quasicrystals. Proc. Natl. Acad. Sci. U.S.A. *118*, e2106230118.
46. Li, L., Xu, Y., Yan, L., Li, X., Li, F., Liu, Z., Zhang, C., Lou, Y., Gao, D., Cheng, X., et al.
(2023). Dynamic network biomarker factors orchestrate cell-fate determination at tipping
points during hESC differentiation. Innovation *4*.
47. Moris, N., Pina, C., and Arias, A.M. (2016). Transition states and cell fate decisions in
epigenetic landscapes. Nat Rev Genet *17*, 693–703.
48. Wang, L.-Z., Su, R.-Q., Huang, Z.-G., Wang, X., Wang, W.-X., Grebogi, C., and Lai, Y.-C.
(2016). A geometrical approach to control and controllability of nonlinear dynamical
networks. Nat Commun *7*, 11323.
49. Heymann, M., and Vanden-Eijnden, E. (2008). The geometric minimum action method: A
least action principle on the space of curves. Comm Pure Appl Math *61*, 1052–1117.
50. Vanden-Eijnden, E., and Heymann, M. (2008). The geometric minimum action method for
computing minimum energy paths. The Journal of Chemical Physics *128*, 061103.
51. Shu, J., Wu, C., Wu, Y., Li, Z., Shao, S., Zhao, W., Tang, X., Yang, H., Shen, L., Zuo, X.,
et al. (2013). Induction of Pluripotency in Mouse Somatic Cells with Lineage Specifiers.
Cell *153*, 963–975.
52. Wu, F., Su, R.-Q., Lai, Y.-C., and Wang, X. (2017). Engineering of a synthetic
quadrastable gene network to approach Waddington landscape and cell fate
determination. eLife *6*, e23702.
53. Zhang, R., Goetz, H., Melendez-Alvarez, J., Li, J., Ding, T., Wang, X., and Tian, X.-J.
(2021). Winner-takes-all resource competition redirects cascading cell fate transitions.
Nat Commun *12*, 853.

54. Zhu, J., Chu, P., and Fu, X. (2023). Unbalanced response to growth variations reshapes
the cell fate decision landscape. Nat Chem Biol *19*, 1097–1104.
55. Cahan, P. ed. (2019). Computational Stem Cell Biology: Methods and Protocols
(Springer New York).
56. Yin, J., Zhang, L., and Zhang, P. (2019). High-Index Optimization-Based Shrinking Dimer
Method for Finding High-Index Saddle Points. SIAM J. Sci. Comput. *41*, A3576–A3595.
57. Smale, S. (1960). Morse inequalities for a dynamical system. Bull. Amer. Math. Soc. *66*,
43–49.

804
805

Supplement:

Figure S1: The algorithm of constructing HSL. Arrows show the information flow among the
various components described in this paper. The yellow rounded rectangles are
corresponding to the manipulations in Method. The symbols in solution landscapes are the
same as the Figure 1. The red ellipse marks the new topopogy of solution landscape. The
gray box indicates the part of identifying the nodes.

Figure S2. More details for HSL of CIS motif.

(A) The simplified HSL. The symbols are the same as those in Figure 2.

(B) The transformation matrix for solution landscapes. IDs in rows and columns denote the indices of solution landscapes in HSL, and the color represent the count of transformation induced by parameter changes. Transformation between solution landscapes of the adjacent columns in the HSL are highlighted by yellow borders.

(C) Transformations of solution landscapes induced by increasing different parameters. The
corresponding parameters are shown in the figure. The arrows in each panel exhibit the
transformations more than 100 times in our simulations.

Figure S3: More details for long route and short route.

(A) The alternation of solution landscapes for long route (blue arrows) and short route (yellow
arrows). The symbols are the same as those in Figure 3. The height of each point is
determined by the energy (i.e.quasi-potential) calculated by the GMAM. These annotations
are used in the Figure S4.

(B) The flow fields and the nullclines for long routes (blue arrows) and short route (yellow
arrows). More details of the region near S1 are enlarged and shown in the right panel.

(C) The schematic of landscape containing three attractors and the definition of E_{SY} (the
energy barrier between S1 and the LY) and E_{SX} (the energy barrier between S1 and the LX).

(D) The change of energy barriers along different routes. The route was normalized to [0,1].
The color is the same as former.

(E) The change of the $\Delta E = E_{SY} - E_{SX}$ along different routes. The route was normalized to
[0,1]. The color is the same as former.

Figure S4: More details for progression route and accuracy route.

(A) The schematic of landscape containing three attractors and the definition of E_{SY} (the energy barrier from S2 to LY), E_{SX} (the energy barrier from S2 to LX), E_{XS} (the energy barrier from LX to S2) and E_{YS} (the energy barrier from LY to S2).

(B) The relationship of Δr_{ini} and the energy barriers (E_{SX} and E_{XS}). As the symmetry of LX and LY, $E_{SX} = E_{SY}$ and $E_{XS} = E_{YS}$.

(C) The alternation of solution landscapes for progression route.

(D) The alternation of solution landscapes for accuracy route.

(E) The change of energy barriers along progression route.

(F) The change of energy barriers along accuracy route.

Figure S5: More details for seesaw model.

(A) The different routes for reprogramming. The red route corresponds to 8-4-2 and the blue route corresponds to 8-6-2.

(B) The 8-6 route and 8-9-6 route.

(C) The schematic of heteroclinic flip bifurcations. The arrows stand for the flow lines. The symbols are the same as those in Figure 5.

(D) The schematic of different traces from the saddle (near ME) to the attractor (S or ECT) near the heteroclinic flip bifurcations. Different colors correspond to different parameters.

(E) The distance between the critical saddle (between S and ECT) and all the points on the trace in (D) at a certain parameter. Different colors correspond to different parameters.