High threat intensity increases conditioned fear generalization and makes extinction of a generalization stimulus less effective

Alba López-Moraga[a,b], Zeynep Gültekin[a], Laura Luyten[a,b,c]*, Tom Beckers[a,b,c]*

   a. Centre for the Psychology of Learning and Experimental Psychopathology, Faculty of Psychology and Educational Sciences, KU Leuven, Leuven 3000, Belgium.

   b. Leuven Brain Institute, KU Leuven, Leuven 3000, Belgium

   c. Corresponding authors: tom.beckers@kuleuven.be , laura.luyten@kuleuven.be

*joint last authors

Abstract

Generalization of conditioned fear is adaptive for survival. However, overgeneralization of

fear from threat cues to loosely similar yet safe stimuli is a hallmark of anxiety-related

disorders. Such overgeneralization may impact other fear learning processes. In particular,

broad fear generalization might limit the effectiveness of extinction training, which may further

maintain anxiety.

Here, we examined if increased generalization of conditioned fear might reduce the later

generalization of fear extinction. To this end, we compared rats conditioned with moderate-

versus high-intensity footshocks, the latter showing stronger generalization of acquired fear

than the former. Within each shock intensity group, we then conducted extinction training

using the original conditioned stimulus (CS) for half of the animals and a generalization

stimulus (GS) for the others. We found stronger preserved responding to the CS after GS

extinction in rats that had initially been conditioned using the high-intensity footshock than in

rats conditioned using the moderate-intensity footshock, in line with the notion of reduced

generalization of extinction from the GS to the CS in the former group. However, we also

found indications for weaker retention of GS extinction in rats conditioned with the high-

intensity footshock, which may in part or in whole explain the apparent difference in

generalization of GS extinction to the CS.

Our results may be important to consider in extinction-based exposure therapy, where

patients often present with broadly generalized fears and exposure treatment is usually not

conducted using the exact cues or situations for which fear was initially acquired.

Keywords: generalization, extinction, fear, anxiety, rats, freezing, fear conditioning.

Highlights:

- We see greater CS-to-GS fear generalization using a strong (vs moderate) US in rats

- Paradoxically, we see more preserved CS fear upon GS extinction in the former
group

- This may reflect an effect on extinction generalization and/or extinction retention

- CS extinction reduced GS fear similarly in both groups

- Male rats showed higher freezing during acquisition and extinction than females

Introduction

Generalization of acquired fear is an adaptive phenomenon. In daily life, animals and humans may encounter threat-predictive stimuli in various forms and under various conditions, and learning to generalize from one cue for impending danger to another one can be vital for survival. However, overgeneralization, or the tendency to exhibit fear when confronted with essentially safe cues that merely resemble a genuine cue for danger, can interfere with adaptive functioning, and has been suggested to be a hallmark of anxiety and stress-related disorders (Cooper et al., 2022).

Generalization can be investigated experimentally through fear conditioning studies in humans, but thanks to a high degree of conservation across species in the laws and mechanisms that govern fear learning (Haaker et al., 2019), the fear conditioning paradigm can readily be applied in rodents as well, allowing to study generalization in a highly controlled environment and to investigate sex differences in the absence of sociocultural factors. Studies on fear generalization in rodents typically start with a fear acquisition phase, where a neutral stimulus (CS, e.g., a tone) is paired with an aversive footshock (US). Later, animals are presented with different cues that resemble the original CS (e.g., other tones). The degree of defensive responding that generalizes to those cues depends on how perceptually similar they are to the CS (Asok et al., 2019; Luyten et al., 2016; Zhang et al., 2017).

Prior research has shown that while fear acquired to a CS will generalize readily to a sufficiently similar generalization stimulus (GS), extinction training of such a GS, in which the GS is repeatedly presented without the US, will leave fear responding to the original CS or other GSs relatively intact in humans (Vervliet et al., 2004, 2005; but see Struyf et al., 2018) and rodents (Alfei et al., 2020, 2021; Boddez et al., 2012; Bouton and Bolles, 1979), whereas extinction training of the original CS will reduce fear responding to the CS as well as GSs. Individuals with high trait anxiety show such lack of generalization of GS extinction to the

original CS even more markedly (Wong and Lovibond, 2020). This observation has clinical

relevance, because exposure therapy is often conducted using generalization cues (e.g.,

cars and traffic situations in general) rather than with the original cues that a patient acquired

fear for (e.g., the car they had a crash in, the intersection they got run over at). In fact, as

fear generalizes more broadly, chances increase that a therapist will expose a patient to

generalized fear-eliciting situations – if they have become afraid of all sorts of traffic, it is all

the more likely that a therapist will just expose them to a traffic situation that elicits strong

fear, regardless of the exact nature of the initial incident.

It remains untested whether individuals who exhibit broader generalization of acquisition

would show more of a reduction of defensive responding to the CS following extinction

training with a GS. Formal models of associative learning suggest that the amount of

generalization depends on the degree of perceived similarity between the CS and GS. Thus,

strong generalization from the CS to a GS would imply that GS extinction should also

generalize well to the CS (Pearce, 1987). As such, individuals who overgeneralize threat

following fear acquisition should also show increased generalization of safety following

extinction learning. However, a better-safe-than-sorry strategy would suggest that strong

generalization of acquisition might in fact go hand-in-hand with *reduced* generalization of

extinction, because incorrectly identifying a dangerous stimulus as safe is more costly than

incorrectly identifying a safe stimulus as dangerous (Dunsmoor and Paz, 2015). This

asymmetry might prevent especially cautious individuals, who would show stronger

generalization from the CS to a GS, from generalizing extinction learning for that GS to the

original fear-eliciting CS or to other GSs (Beckers et al., 2023). Here, we studied whether

heightened generalization of fear after acquisition would indeed weaken the later

generalization of fear extinction.

In the present study, we performed fear conditioning using moderate- and high-intensity

footshock USs in male and female rats, in light of the evidence that high-intensity USs

promote stronger fear generalization than moderate-intensity USs (Laxmi et al., 2003; Xuan

et al., 2023). Subsequently, we conducted fear extinction using either the CS or a GS. We

hypothesized that enhanced generalization of fear from CS to GS after acquisition would

weaken the later generalization of fear extinction from GS to CS. Additionally, we expected

female rats to show overall slower extinction than male rats (Baran et al., 2009; Fenton et al.,

2016; Greiner et al., 2019) and more generalization of freezing from CS to GS (Greiner et al.,

2019; Keiser et al., 2017).

Methods

Preregistration and data availability

The design, procedures, sample size and analysis plan were preregistered on the Open

Science Framework (OSF) prior to the start of data collection

(http://doi.org/10.17605/osf.io/9yd83). All data and scripts are available through the same

OSF link as well.

Subjects

The experiment was performed in accordance with Belgian and European laws (Belgian

Royal Decree of 29/05/2013 and European Directive 2010/63/EU) and the ARRIVE 2.0

guidelines (Percie du Sert et al., 2020) and approved by the KU Leuven animal ethics

committee (project license number 011/2019). The study was conducted in 64 8-week-old

male (270-300 g at arrival) and female (200-230 g at arrival) rats (Janvier Labs, France).

Animals were housed in groups of 4, on a 12-hour light-dark cycle (lights on at 7 am);

experimental procedures were performed between 9 am and 5 pm. The cages had bedding

and cage enrichment in the form of a hanging red polycarbonate tunnel. Water and food were

available ad libitum for the entire experiment, except during behavioral testing. Animals were

acclimated for one week to our facility and habituated to handling for 2 days before the start

of the experiment.

Apparatus

Eight identical operant chambers (30.5 cm width, 25.4 cm depth and 30.5 cm height; Rat Test Cage, Coulbourn Instruments, Pennsylvania, USA) were used simultaneously and were enclosed in sound-attenuating boxes. The shock grid consisted of 18 stainless steal rods of 0.4 cm diameter, with 1.5 cm spacing between the center of the rods (Modular Shock Floor H10-11R-TC-SF, Coulbourn Instruments). This grid was connected to a scrambled shocker (Precision Animal Shocker H13-15, Coulbourn Instruments). The different tones were generated by a modified tone generator (Seven-tone generator H12-07, Coulbourn Instruments) and transmitted to the operant chamber through two speakers in opposite walls of the chamber. The experiment was conducted using Graphic State 4 (Coulbourn Instruments) software with the house light off. Behavior was continuously recorded during the experimental sessions using an IP camera (Foscam C2M, Shenzhen, China) attached to the ceiling of the sound-attenuating box. The operant chambers were cleaned between animals with soapy water (5.55% Instanet, Henkel, Vilvoorde, Belgium).

Experimental design

Rats (n = 64) were evenly divided into four groups: moderate shock during fear conditioning, extinction training with CS (MSCS), moderate shock during fear conditioning, extinction training with GS (MSGS), strong shock during fear conditioning, extinction training with CS (SSCS), strong shock during fear conditioning, extinction training with GS (SSGS) (see Figure 1). Rats of both sexes were divided equally over the four groups, the final group sizes were: MSCS (n = 16; 8 females), MSGS (n = 16; 8 females), SSCS (n = 16; 8 females) and SSGS (n = 16; 8 females). All sessions were scheduled 24h apart, except for the spontaneous recovery test, which was conducted 7 days after the reinstatement test.

Figure 1. Experimental design. MSCS: moderate shock acquisition, extinction training with CS (n = 16); MSGS: moderate shock acquisition, extinction training with GS (n = 16); SSCS: strong shock acquisition, extinction training with CS (n = 16); SSGS: strong shock acquisition, extinction training with GS (n = 16). All sessions were scheduled 24h apart, except for the spontaneous recovery test, which was conducted 7 days after the reinstatement test. At the end of extinction test, some rats were euthanized for tissue collection. Thus, in reinstatement and reinstatement test all groups had n = 12, except the SSCS group which had n = 13 (n = 7 females, n = 6 males). After reinstatement test, some rats were euthanized for tissue collection and sample sizes for spontaneous recovery test were n = 10, except the SSCS group which had n = 11 (females n = 6, males n = 5) and the MSCS group which had n = 9 (n = 4 females, n = 5 males).

First, all rats underwent a habituation session during which the CS and GS were each

presented twice. Half of the rats received the tone presentations in the sequence CS, GS,

CS, GS, while the other half received them in the sequence GS, CS, GS, CS. The CS and

GS consisted of a pure 3-kHz tone and a pure 7-kHz tone, counterbalanced between

animals. After 3 min of acclimation, each tone was presented for 30 s, with an average ITI of

300 s (270-330 s). After the last tone, the rats remained in the testing chamber for one

additional minute.

The next day, rats underwent an acquisition session. After 3 min of acclimation, 9 CS-US

pairings were presented, with an ITI averaging 180 s (150-210 s). Half of the rats (n = 32)

were trained using a moderate footshock US of 0.4 mA intensity and 1 s duration. The other

half (n = 32) were trained using a strong footshock US of 1 mA intensity and 2 s duration.

The 30-s CS co-terminated with the US in both conditions. After the last CS-US pairing, rats

remained in the testing chamber for an additional minute. One day later, an acquisition

retention test was performed, during which the CS and GS were each presented twice, with

the same parameters as in the habituation session.

The extinction session consisted of 18 CS or GS presentations, after 3 min of acclimation,

and with an ITI averaging 180 s (150-210 s). After the last tone presentation, rats remained in

the testing chamber for an additional minute. Four rats per group, 2 of each sex, were

perfused 90 min after the extinction test to dissect their brains for immunohistochemical

staining. The next day, an extinction retention test was performed, during which the CS and

GS were each presented twice, with the same parameters as in the habituation session.

A reinstatement manipulation took place one day later. After 3 min of acclimation, a single

footshock of 0.4 mA and 1 s duration was delivered, followed by 2 more minutes in the

testing chamber. The next day, a reinstatement test session was conducted, consisting of 2

CSs and 2 GSs, as in the habituation session. Two rats per group, one of each sex, were

perfused 90 min after the reinstatement test.

Seven days after the reinstatement test, a spontaneous recovery test took place with 2 CSs

and 2 GSs, using the same parameters as in the habituation session. Four rats per group,

two of each sex, were perfused 90 min after the spontaneous recovery test.

Due to problems with tissue quality after freezing the specimens, the planned

immunohistochemical stainings were not performed.

Behavior

Freezing during CS presentations was scored manually from the videos as a percentage of

CS duration. Freezing was defined as full immobility except for minimal movements

associated with breathing.

A freezing discrimination index was calculated for sessions where both CSs and GSs were

presented. It was calculated as the difference in freezing between the two different tones

(i.e., CS minus GS) divided by the sum of freezing to both tones (Xu and Südhof, 2013).

Automated analyses of behavior

To investigate darting behavior and contextual freezing, we used a combination of

DeepLabCut (version 2.3.9) (Mathis et al., 2018; Nath et al., 2019) and SimBA (version

1.99.5) (Goodwin et al., 2024).

First, we used DeepLabCut to obtain pose estimation. We labeled 200 frames taken from 10

videos, each video coming from a different animal; 95% of these frames were used for

training. Five body parts were selected: nose, left ear, right ear, centroid, and tail base. We

used a RestNet-50 neural network with default parameters for 300000 training iterations.

Given that our results were initially unsatisfactory (see OSF for further details), we extracted

outliers, relabeled 20 of those additional outlier frames for each video and ran an additional

training iteration with the same parameters as above. We validated with 1 shuffle and found

that the test error was 1.8 pixels and the training error was 5.78 pixels (image size: 456 x 256

pixels, downsampled from original 1280 x 720 pixels).

Second, we used SimBA to further analyze pose estimation data obtained with DeepLabCut.

We created a single animal project configuration with user-defined body parts (same body

parts as for DeepLabCut). We defined pixels per mm by considering the distance between

the width of the bottom of the fear conditioning chamber, which in our case was 291 mm. We

used the body-part: nearest interpolation method and a gaussian smoothing method of

100 ms, outlier correction was skipped.

To analyze darting, we obtained velocity traces per animal at 1-s time bins. We filtered

speeds under 23.5 cm/s. An animal was considered a darter when a darting bout occurred

during CS 3 to 7 of the acquisition session, excluding intertrial intervals and US presentations

(Gruene et al., 2015).

To analyze contextual freezing, we started from a model that we described and used

previously (López-Moraga et al., 2025). We annotated three additional videos; we added 972

frames (32.4 s) annotated with freezing present and 24491 frames (816.33 s) annotated with

freezing absent. Here, we specifically focused on labelling sniffing as a non-freezing bout.

We built the model with the following settings: n_estimators = 2000, RF_criterion = gini,

RF_max_features = sqrt, RF_min_sample_leaf = 1, under_sample_ratio = 3, class_weights =

custom: Freezing present = 2; Freezing Absent = 1, with an 80% training split. The evaluation

performance of the freezing classifier had an F1 score of 0.877, precision of 0.876, and recall

of 0.878 for freezing present and F1 score of 0.960, precision of 0.960, and recall of 0.959 for

freezing absent. We set the discrimination threshold at Pr = 0.75 and a minimum duration of

500 ms (see OSF for further details).

Data analyses

Statistical analyses were preregistered; any deviations from the preregistered analyses are

specified below. Data were analyzed using t-tests and one-way, two-way, repeated-measures

or mixed ANOVAs, as appropriate. Post-hoc tests with corrections for multiple testing

followed if significant main effects or interactions were found. If assumptions were violated,

nonparametric statistical analyses were performed. For the nonparametric mixed RM

ANOVAs, we applied an aligned rank transform ANOVA using the ARTool package (Elkin et

al., 2021) in R (R Core Team, 2022) and R Studio (RStudio Team, 2021). All other statistical

analyses were performed using performance (Lüdecke et al., 2021), afex (Singmann et al.,

2023), emmeans (Lenth et al., 2024) and effectsize (Ben-Shachar et al., 2024) R packages.

Data were processed and plotted using the tidyverse (Wickham and RStudio, 2022) R

packages.

Analyses for habituation and acquisition sessions were not preregistered. Habituation was

analyzed with a non-parametric mixed ANOVA with shock intensity (moderate, strong) as a

between-subjects variable and stimulus (CS, GS) as a within-subjects variable. Acquisition

performance was analyzed with a two-way ANOVA with shock intensity and sex as between-

subjects factors.

Data from the acquisition retention test, extinction retention test, reinstatement test and

spontaneous recovery test were analyzed with sex and either shock intensity (moderate,

strong) or experimental group (MSCS, MSGS, SSCS, SSGS) as between-subjects factors

and stimulus (CS, GS) as within-subjects factor. If effects of sex were not significant, both

sexes were collapsed in the mixed ANOVA as preregistered. As a deviation from the

preregistration, we did not directly compare retention test sessions (i.e., acquisition test

versus extinction test) within one mixed ANOVA. Two-tailed one-sample t-tests were

performed on the discrimination indices for the acquisition test and extinction test of each

shock intensity group, comparing the mean of each group to 0 (e.g., no discrimination). This

comparison was not preregistered. The analysis of contextual freezing during the extinction

retention test was exploratory and consisted of a mixed ANOVA with sex and group as

between-subject variables and minute as within-subjects variable.

Differences in freezing during extinction training were analyzed using a mixed ANOVA with

trial as a within-subjects factor and sex and experimental group as between-subjects factors.

In a separate mixed ANOVA, to assess changes in freezing, the first three tones (i.e., early

block) were compared to the last three tones (i.e., late block) using block as a within-subjects

factor and group and sex as between-subjects factors.

Criteria for classifying a rat as darter or non-darter were preregistered as described above.

The chi-square test to compare the ratio of darters between sexes was preregistered, but the

analyses comparing the ratio of darters between groups were exploratory. The analysis of

differences in freezing between darters and non-darters by experimental session and sex

were preregistered, except for the analysis of the extinction test session. The correlation

analysis between number of darting bouts and freezing was exploratory.

Results

The aim of this experiment was to investigate whether increased generalization of acquisition

from CS to GS would go hand in hand with reduced generalization of extinction from GS to

CS. To evaluate this, we needed to manipulate the generalization of acquisition. We therefore

divided our sample in two groups, one that received a moderate intensity footshock as US

during acquisition and one that received a strong intensity footshock as US, as higher US

intensity has been shown to promote increased conditioned fear generalization (Dunsmoor et

al., 2017; Laxmi et al., 2003; Xuan et al., 2023). After a test of generalization of acquisition,

we conducted extinction training with either the CS or a GS and performed an extinction

retention test 24 h later to test our central hypothesis. Finally, we performed exploratory tests

of reinstatement and spontaneous recovery.

Manipulation checks

In the habituation session, all rats exhibited uniformly low levels of freezing ($F_{(1, 62)} = 1.58$,

$p = 0.213$, $\omega_p^2 < -0.01$, see further analyses in Table 1), with no significant differences in

discrimination between CS and GS ($t(31) = 0.65$, $p = 0.520$, $d = 0.12$) (see Figure 2).

During acquisition, we did not observe differences in freezing between rats that experienced

moderate footshock USs or strong footshock USs ($F_{(1, 60)} = 1.58$, $p = 0.213$, $\omega_p^2 < -0.01$),

but males showed significantly higher freezing than females ($F_{(1, 60)}$ = 10.00, p = 0.002, ω_p^2

= 0.12).

We also investigated darting behavior during acquisition. Rats were classified as darters or

non-darters according to previous literature (Gruene et al., 2015). Forty-seven percent of

female rats (15/32 rats) were classified as darters, against 19% (6/32) of male rats. A chi-

square test revealed a significant association between darter status and sex ($\chi^2(1)$= 4.54, p =

0.033, φ = 0.3, see Figure S1A). This suggests that the distribution of darters was different

between sexes. Additionally, female darters performed more dart bouts (M = 3.47, SE = 0.68)

than male darters (M = 1.83, SE = 0.31) during acquisition. We did not find a significant

association between shock intensity group and darter status ($\chi^2(1)$= 0.28, p = 0.594, φ = 0.1).

Given the different distribution of darters between sexes, we analyzed by sex whether there

were differences in freezing during acquisition between darter and non-darter rats (Figure

S1B). Female darters showed significantly less freezing than female non-darters (t(30) = -

4.8, p < 0.001, d = -1.7). However, in male rats, darting did not make a difference (t(30) = -

0.34, p = 0.73, d = -0.17). We did not find differences in freezing between darters and non-

darters in later sessions, regardless of sex (see Supplementary Table 1 for further statistics).

In the acquisition test, there was a significant interaction between shock intensity group and

stimulus ($F_{(1, 62)}$ = 5.22, p = 0.026, ω_p^2 = 0.02, see further analyses in Table 1). Specifically,

the moderate shock group showed significantly higher freezing to the CS than the GS (t(124)

= 2.63, p = 0.01, d = 0.66), whereas the strong shock group showed no significant difference

in freezing between the stimuli (t(124) = 0.43, p = 0.669, d = 0.11), suggesting stronger

generalization in the latter group, although the freezing discrimination index did not differ

significantly between both groups (t(62) = 1.47, p = 0.148, d = 0.37). Additionally, we

performed an exploratory one-sample t-test to assess if the discrimination index was

significantly different from 0 in each group. For rats in the moderate shock group, the

discrimination index was significantly greater than 0 (t(31) = 3.34, p = 0.002, d = 0.59),

whereas for rats in the strong shock group it was not significantly different from 0 (t(31) =

1.32, p = 0.196, d = 0.32).

For the subsequent extinction training phase, both shock intensity groups were to be split in

two, with half of the rats receiving extinction training for the CS and the other half receiving

extinction training for the GS. We checked whether any differences between the four groups-

to-be were present already in the acquisition test (see Figure S2 and Table S2).

Figure 2. Freezing to the tones during habituation, acquisition, and the acquisition retention

test and discrimination indices during habituation and acquisition test. Freezing boxplots

represent average freezing to each stimulus, expressed as % of tone duration. Discrimination

boxplots represent the average freezing discrimination index per group. Discrimination

indices were calculated as the difference in freezing during the presentation of the CS versus

the GS, divided by the sum of freezing to both tones. **a.** Graphical representation and results

of the habituation session. **b.** Graphical representation and results of the acquisition session.

Males showed higher freezing than females during acquisition ($F(1, 60) = 10$, $p = 0.002$). In

the line plot bold lines represent the mean and the surrounding shaded area represents the

standard error of the mean. **c.** Graphical representation and results of the acquisition

retention test. An interaction between shock intensity group and freezing ($F(1, 62) = 5.22$, $p =$

0.026) reflects the fact that there were significant differences in freezing between CS and GS

in the moderate shock group ($t(124) = 2.63$, $p = 0.01$), but not in the strong shock group

($t(124) = 0.43$, $p = 0.669$). The discrimination index was significantly greater than 0 for rats in

the moderate shock group ($t(31) = 3.34$, $p = 0.002$), but not the strong shock group ($t(31) =$

1.32, $p = 0.196$).

In extinction training, we observed a reduction of freezing over trials ($F(17, 1020) = 16.06$, p

< 0.001, $\omega_p^2 = 0.13$), and overall differences in freezing levels between groups ($F(3, 60) =$

8.54, $p < 0.001$, $\omega_p^2 = 0.24$) (see Figure 3). In particular, the SSCS group exhibited

significantly higher levels of freezing than the SSGS ($t(60) = 3.39$, $p = 0.006$, $r = 0.32$) and

MSGS groups ($t(60) = 4.92$, $p < 0.001$, $r = 0.43$).

Comparing freezing levels between an early block and a late block of the extinction training

session, we found that males froze significantly more than females ($F(1, 56) = 4.66$, $p =$

0.035, $\omega_p^2 = 0.05$) and that freezing was lower in the late block than in the early block, as

expected ($F(1, 56) = 93.47$, $p < 0.001$, $\omega_p^2 = 0.37$). Additionally, significant differences in

freezing levels were detected between groups ($F(3, 56) = 8.43$, $p = 0.001$, $\omega_p^2 = 0.22$). In

particular, freezing levels differed between the two moderate shock groups, with rats

undergoing extinction with the CS (MSCS) exhibiting higher freezing than rats undergoing

extinction with the GS (MSGS) ($t(56) = 2.7$, $p = 0.044$, $r = 0.27$). Similarly, for the strong

shock groups, SSCS rats showed higher levels of freezing than SSGS rats ($t(56) = 3.02$, $p =$

0.019, $r = 0.33$). We evaluated differences in freezing depending on sex and whether animals

had been classified as darters during acquisition. We did not find significant differences

between female darters and non-darters ($t(30) = -1.37$, $p = 0.181$, $d = -0.49$, see Figure S1C)

nor between male darters and non-darters ($t(30) = -1.3$, $p = 0.202$, $d = -0.64$).

Figure 3. Freezing during extinction training, expressed in % of tone duration. **a.** Graphical representation of the extinction training session. **b.** Bold lines represent the mean and the surrounding shaded area the standard error of the mean. A reduction in freezing behavior was observed during extinction training ($F_{(17, 1020)} = 16.06$, $p < 0.001$), along with significant differences between groups ($F_{(3, 60)} = 8.54$, $p < 0.001$). **c.** Boxplots represent the average of the first three (i.e., early) and last three (i.e., late) tone presentations of the extinction session. While a general reduction in freezing was observed between the early and late block ($F_{(1, 56)} = 93.47$, $p < 0.001$), males had higher freezing levels than females ($F_{(1, 56)} = 4.66$, $p = 0.035$) and significant differences between groups were found ($F_{(3, 56)} = 8.43$, $p = 0.001$).

Effects of generalization of acquisition on generalization of extinction

In the extinction retention test session, we observed a group by sex interaction ($F_{(3, 56)} = 3.01$, $p = 0.038$, $\omega_p^2 = 0.09$) and a group by stimulus interaction ($F_{(3, 56)} = 4.13$, $p = 0.01$, $\omega_p^2 = 0.06$) (see Figure 4). Further investigation of the group by stimulus interaction, zooming in on the degree of generalization of extinction to the stimulus *not* presented during extinction training, revealed that rats in the SSGS group showed significantly higher freezing to the CS than rats in the MSGS group ($t_{(112)} = 3.41$, $p = 0.005$, $d = 1.21$), whereas rats from the

SSCS group and rats from the MSCS group did not display significant differences in freezing

to the GS ($t(112) = 1.38$, $p = 0.514$, $d = 0.49$). Rats in the SSGS group had not shown higher

freezing to the CS than rats in the MSGS groups in the acquisition test. Do note, however,

that rats in the MSGS group also tended to show better extinction retention for the GS than

rats in the SSGS group, even if not significantly so ($t(112) = -2.54$, $p = 0.06$, $d = -0.9$).

Importantly, rats in the MSCS group did show significantly less freezing to the CS than those

in the SSCS group ($t(112) = -3.20$, $p = 0.01$, $d = -1.13$). Group differences were detected in

the freezing discrimination index as well ($F(3, 60) = 10.89$, $p < 0.001$, $\omega2 = 0.32$). The MSCS

group showed a significantly lower discrimination index than the MSGS group ($t(60) = -5.47$,

$p < 0.001$, $d = -1.94$) and the SSGS group ($t(60) = -3.24$, $p = 0.01$, $d = -1.15$). In follow-up of

those results, we explored whether the discrimination index was significantly different from 0

for each group separately. Regardless of shock intensity, the discrimination index was not

significantly different from 0 for rats that underwent CS extinction, suggesting that extinction

generalized well from CS to GS (MSCS: $t(15) = -0.66$, $p = 0.516$, $d = -0.17$; SSCS: $t(15) =$

1.658, $p = 0.118$, $d = 0.41$). In contrast, the discrimination index was significantly different

from 0 for rats that underwent GS extinction (MSGS: $t(15) = 7.95$, $p = < 0.001$, $d = 1.99$;

SSGS: $t(15) = 3.665$, $p = 0.002$, $d = 0.92$), indicating that GS extinction did not fully

generalize to the CS. One might argue that the calculation of the discrimination index as a

relative measure might obscure differences in generalization between the MSGS and SSGS

groups, given the higher degree of freezing to the GS in the latter than in the former group.

However, an alternative (non-preregistered) measure of discrimination, calculated as the

absolute difference in freezing between CS and GS, yielded no difference in discrimination

between the MSGS and SSGS groups either ($t(60) = -0.695$, $p = 0.8987$, $d = -0.25$), with

discrimination again significantly different from 0 in both groups (MSGS: $t(15) = 4.7$, $p <$

0.001, $d = 1.17$; SSGS: $t(15) = 3.39$, $p = 0.004$, $d = 0.85$).

Further investigation of the significant group by sex interaction revealed differential freezing

between sexes in the SSCS group only ($t(112) = -3.49$, $p < 0.001$, $d = -1.23$), with males

showing higher freezing than females. Given the high levels of freezing by the male rats of

the SSCS group, we investigated if these differences were already present before tone

onset. While we observed that contextual freezing levels increased over time ($F(4, 224) =$

5.39, $p < 0.001$, $\omega_p^2 = 0.02$), overall contextual freezing levels were low (see Figure S3). We

observed differences between groups ($F(3, 56) = 2.84$, $p = 0.046$, $\omega_p^2 = 0.02$, see

Supplementary Table 3), but these differences did not withstand multiple testing correction.

To further evaluate differences in GS extinction retention as a potential source for the

differences in CS responding between the SSGS and MSGS group, we compared the

difference in freezing between the last two tones of the extinction session and the relevant

extinction retention test tones between groups that received the same extinction training (i.e.,

comparing SSCS to MSCS in their change in CS responding from the end of extinction

training to the extinction retention test and SSGS to MSGS in their change in GS responding

from the end of extinction training to the extinction retention test). We observed that the

difference between the SSGS and MSGS groups in the increase in freezing between both

sessions failed to reach significance ($F(1, 28) = 3.78$, $p = 0.062$, $\omega_p^2 = 0.08$), although it was

nominally larger in the SSGS group ($M = 12.2$, $SE = 4.59$) than in the MSGS ($M = 2.11$, $SE =$

2.43). There was no difference in the increase in freezing from the end of extinction to the

extinction retention test when comparing the SSCS and MSCS groups ($F(1, 28) = 2.58$, $p =$

0.119, $\omega_p^2 = 0.05$).

Figure 4. Results of the extinction retention test session. **a.** Graphical representation of the extinction retention test. **b.** The discrimination index boxplot represents the average of the freezing discrimination index. Discrimination indices were calculated as the difference in freezing to the different tones (CS versus GS), divided by the sum of freezing to both tones. Significant differences in discrimination were observed between groups ($F_{(3, 60)} = 10.89$, $p < 0.001$). **c.** The freezing boxplots represent the average percentage of freezing to each stimulus, expressed in % of tone duration. Interactions between group and sex ($F_{(3, 56)} = 3.01$, $p = 0.038$) and between group and stimulus ($F_{(3, 56)} = 4.13$, $p = 0.01$) were found in freezing behavior. Rats in the MSGS group showed significantly lower freezing to the CS than rats in the SSGS group ($t_{(112)} = -3.41$, $p = 0.005$). In the SSCS group, males showed higher freezing than females ($t_{(112)} = -3.49$, $p < 0.001$).

Return of fear tests

We presented unsignaled USs, followed by a reinstatement test session 24 hours later (see Figure 5). While we did not observe significant group differences ($F_{(3, 45)} = 1.48$, $p = 0.231$, $\omega_p^2 = 0.03$), we observed significantly higher freezing to the CS than to the GS ($F_{(1, 45)} =$

18.1, $p < 0.001$, $\omega_p^2 = 0.07$). No group differences were found in freezing discrimination

indices either ($F_{(3, 45)} = 1.62$, $p = 0.199$, $\omega_p^2 = 0.04$).

Seven days later, we tested differences in spontaneous recovery, and found higher freezing

to the CS than to the GS again ($F_{(1, 36)} = 10.54$, $p = 0.003$, $\omega_p^2 = 0.06$), in the absence of

group differences ($F_{(3, 36)} = 0.4$, $p = 0.757$, $\omega_p^2 = -0.05$) (see Figure 5). We did find group

differences in the freezing discrimination index ($F_{(3, 36)} = 3.77$, $p = 0.019$, $\omega_p^2 = 0.17$).

Specifically, rats in the MSGS group showed a significantly higher discrimination index than

those in the MSCS group ($t_{(36)} = -2.78$, $p = 0.041$, $d = -1.28$) and the SSCS group ($t_{(36)} = $

2.99, $p = 0.024$, $d = 1.31$).

Figure 5. Freezing during reinstatement test and spontaneous recovery test. The freezing

boxplots represent the average for each stimulus; results are expressed in % of tone

duration. The discrimination index boxplots represent the average freezing discrimination

index. Discrimination indices were established by calculating the difference in freezing to the

tones (CS minus GS) and dividing by the sum of freezing to both tones. **a.** Graphical

representation of the reinstatement test and the spontaneous recovery test. **b.** Results of the

reinstatement test session. Significant differences between CS and GS ($F_{(1, 45)} = 18.1$, $p < $

0.001) in freezing behavior, but no significant differences in discrimination index between

groups were found (F(3, 45) = 1.62, p = 0.199). **c.** Results of the spontaneous recovery test

session. Significant differences between CS and GS were found in freezing behavior (F(1,

36) = 10.54, p = 0.003), as well as significant differences in the discrimination index between

groups (F(3, 36) = 3.77, p = 0.019).

Discussion

We investigated how increased generalization of acquisition affects the generalization of

extinction in a Pavlovian fear conditioning task. To achieve differences in generalization of

acquisition, in line with previous literature (Laxmi et al., 2003; Xuan et al., 2023), we

subjected rats to cued fear conditioning with strong versus moderate intensity shocks as US.

While rats that received moderate shocks during acquisition discriminated between the CS

and a GS on a subsequent retention test, rats that received strong shocks did not and

showed similar levels of freezing to both tones.

In the extinction test session, we confirmed some of our preregistered hypotheses. In the

moderate shock groups, we observed the expected asymmetry: in the MSGS group, we

observed a high discrimination index, with higher freezing to the CS than to the GS, meaning

that extinction did not strongly generalize from GS to CS, while in the MSCS group, the

discrimination index was low, in other words, in the MSCS group extinction generalized

strongly from the CS to the GS, yielding similar freezing for both tones. In the strong shock

groups, we expected that the enhanced generalization of acquisition would further reduce the

generalization of extinction, yielding larger discrimination at test between CS and GS

particularly in the SSGS group. In support of our hypothesis, we found that animals in the

SSGS group showed significantly higher freezing to the CS than those in the MSGS group.

We did not observe higher freezing to the GS in the SSCS group than in the MSCS group,

which suggests that the enhanced generalization of acquisition from CS to GS as a result of

high US intensity went hand-in-hand with reduced generalization of extinction from an

extinguished GS back to the CS but not from the extinguished CS to a GS. However, we also

noted a tendency for higher responding to the GS in the SSGS group compared to the

MSGS group. While not significant after multiple-test correction, this suggests that there was

less retention of GS extinction in the SSGS group than in the MSGS group. Hence, it is

possible that the SSGS group shows higher responding to the CS than the MSGS group

after GS extinction not because of weaker generalization from the GS to the CS but because

of the weaker retention of GS extinction leaving less to generalize to the CS. Further

research should aim to disentangle these possibilities.

Regarding the return of fear tests, we did not find any group differences in the reinstatement

test. We did find that rats showed higher CS freezing than GS freezing, indicative of a

selective return of fear to the CS, regardless of initial US intensity or the nature of the

stimulus used for extinction training. One week later, we performed a spontaneous recovery

test and again observed higher CS-elicited than GS-elicited freezing. Not all our

preregistered differences were confirmed here. One reason could be a reduction of power

relative to the extinction test session, as some animals were euthanized in between to collect

their brain tissue.

It is important to note that the results for the discrimination indices should be interpreted with

caution. Rats that underwent GS extinction had mostly received reinforced CS presentations,

while the GS was never paired with shock. However, in the groups that underwent CS

extinction, the CS became effectively ambiguous, as the CS was paired with shock in

acquisition and then not followed by shock during extinction training. Thus, after extinction

training, the results of discrimination indices may be biased towards increased discrimination

in the MSGS and SSGS groups compared to the MSCS and SSCS groups.

Our data show that the greater generalization that is observed after acquisition with a strong-

intensity US is accompanied by stronger retention of CS-elicited fear upon GS extinction. Our

results do not allow to discern to what degree this is due to a reduction in generalization of

extinction from GS to CS proper, as hypothesized, or to an impaired retention of GS

extinction in rats trained with strong-intensity shocks. Future research is needed to dissect

this. Of note, the effect did not extend to rats that underwent CS extinction, whose freezing

towards the GS at test was the same whether trained with moderate- or high-intensity

shocks.

While future studies might seek to replicate and elaborate these results in humans, it would

also be worthwhile to investigate the neurobiological mechanisms involved in and

constraining the generalization of extinction or the retention of GS extinction in further animal

work. Whereas proactively limiting the generalization of fear after a threatening event might

be difficult to achieve, finding ways to improve the generalization of extinction might be more

feasible. This could prove beneficial then as an adjunct to exposure therapy. As one avenue

for further exploration, previous research has shown that vagus nerve stimulation during

extinction can increase generalization of extinction in male rats (Noble et al., 2019; Souza et

al., 2020).

The use of equal numbers of male and female animals allowed us to explore sex differences.

We detected such differences, especially in acquisition and extinction, although less

pronounced than expected. We compared freezing between an early extinction block (i.e.,

first three tones) and a late block (i.e., last three tones), which revealed that male rats

showed higher freezing in early and late trials than female rats during those same trials.

Such generally reduced freezing in female rats during extinction training has been reported

previously in other tasks (Shanazz et al., 2022). While we also expected slower extinction in

female rats based on prior literature (Baran et al., 2009; Fenton et al., 2016; Graham et al.,

2009; Greiner et al., 2019), we found no supporting evidence for this in our results.

Surprisingly, males in the SSCS group showed high freezing to the CS despite having

undergone extinction training with that same CS in the preceding session. This result is

puzzling given that their extinction training results look as expected, exhibiting a gradual

reduction in freezing over extinction trials. Thus, it seems that males in the SSCS group

showed poor extinction retention during the extinction test session. This result could not be

attributed to increased contextual fear, nor to technical issues, as rats from all experimental

groups were tested simultaneously.

The lack of sex differences during the extinction test was unexpected, as we anticipated

lower discrimination indices in female rats, in line with previous reports (Greiner et al., 2019;

Keiser et al., 2017). Note that Keiser et al. (2017) assessed generalization of contextual

rather than cued fear. While Greiner et al. (2019) found a lack of extinction to a conditioned

fear cue and a lack of discrimination between a conditioned fear cue and a conditioned

safety cue in females when assessing freezing behavior, females in that study did show

differential darting behavior in response to a safety versus a fear cue.

Darting behavior has previously been introduced as a conditioned behavior expressed

largely by female rats (Gruene et al., 2015; Mitchell et al., 2022), although the associative

properties of darting have been challenged (Trott et al., 2022). In line with previous research

(Gruene et al., 2015; Mitchell et al., 2022), we observed that females were more likely to be

classified as darters than males (47% compared to 19%). We also found that during

acquisition, female darters showed less freezing than female non-darters, but this difference

was not significant in their male counterparts nor in any of the other experimental sessions.

Two important sidenotes are that female darters darted more than male darters, and that,

whereas our female sample had a well-balanced distribution of darters versus non-darters

(15 darters vs 17 non-darters), the distribution was very unbalanced for the males (6 darters

vs 26 non-darters), which makes these statistical comparisons more unreliable in male rats.

In conclusion, we found higher responding to the CS after GS extinction in strong shock rats

compared to moderate shock rats. This could be either due to weaker retention of GS

extinction in strong shock rats (while CS extinction is not similarly affected) or due to weaker

generalization, as hypothesized. This may be important to consider in extinction-based

exposure therapy, as patients frequently exhibit broadly generalized fears, and exposure

treatment typically involves cues or situations different from those to which the fear was

originally acquired.

Research data statement

Acknowledgements

We thank Yana van der Heyden for assistance with behavioral scoring. Graphical

representations of the behavioral setups in the figures were made with Biorender.

Author contributions

A. L-M.: conceptualization, data curation, formal analysis, funding acquisition, investigation,

methodology, project administration, software, validation, visualization, writing – original draft.

Z. G.: data curation, investigation; writing – review & editing. L. L.: conceptualization, funding

acquisition, methodology, project administration, resources, supervision, writing – review &

editing. T. B.: conceptualization, methodology, funding acquisition, project administration,

resources, supervision, writing – review & editing.

Funding sources

This work was supported by the KU Leuven Research Grant 3H190245 and FWO PhD

fellowship 11K3821N.

Declaration of competing interests

The authors have no known competing financial interests or personal relationships that could

have appeared to influence the work reported in this paper.

References

Alfei, J.M., De Gruy, H., De Bundel, D., Luyten, L., Beckers, T., 2021. Apparent

     reconsolidation interference without generalized amnesia. Progress in Neuro-

Psychopharmacology and Biological Psychiatry 108, 110161.
https://doi.org/10.1016/j.pnpbp.2020.110161

Alfei, J.M., Ferrer Monti, R.I., Molina, V.A., De Bundel, D., Luyten, L., Beckers, T., 2020.
Generalization and recovery of post-retrieval amnesia. Journal of Experimental
Psychology: General 149, 2063–2083. https://doi.org/10.1037/xge0000765

Asok, A., Kandel, E.R., Rayman, J.B., 2019. The neurobiology of fear generalization.
Frontiers in Behavioral Neuroscience 12, 329.
https://doi.org/10.3389/fnbeh.2018.00329

Baran, S.E., Armstrong, C.E., Niren, D.C., Hanna, J.J., Conrad, C.D., 2009. Chronic stress
and sex differences on the recall of fear conditioning and extinction. Neurobiology of
Learning and Memory 91, 323–332. https://doi.org/10.1016/j.nlm.2008.11.005

Beckers, T., Hermans, D., Lange, I., Luyten, L., Scheveneels, S., Vervliet, B., 2023.
Understanding clinical fear and anxiety through the lens of human fear conditioning.
Nat Rev Psychol 2, 233–245. https://doi.org/10.1038/s44159-023-00156-1

Ben-Shachar, M.S., Makowski, D., Lüdecke, D., Patil, I., Wiernik, B.M., Thériault, R., Kelley,
K., Stanley, D., Caldwell, A., Burnett, J., Karreth, J., Waggoner, P., 2024. effectsize:
Indices of Effect Size.

Boddez, Y., Callaerts-Vegh, Z., Vervliet, B., Baeyens, F., D'Hooge, R., Hermans, D., Beckers,
T., 2012. Stimulus generalization and return of fear in C57BL/6J mice. Frontiers in
Behavioral Neuroscience 6. https://doi.org/10.3389/fnbeh.2012.00041

Bouton, M.E., Bolles, R.C., 1979. Contextual control of the extinction of conditioned fear.
Learning and Motivation 10, 445–466. https://doi.org/10.1016/0023-9690(79)90057-2

Cooper, S.E., van Dis, E.A.M., Hagenaars, M.A., Krypotos, A.-M., Nemeroff, C.B., Lissek, S.,
Engelhard, I.M., Dunsmoor, J.E., 2022. A meta-analysis of conditioned fear
generalization in anxiety-related disorders. Neuropsychopharmacol. 47, 1652–1661.
https://doi.org/10.1038/s41386-022-01332-2

Dunsmoor, J.E., Kroes, M.C.W., Braren, S.H., Phelps, E.A., 2017. Threat intensity widens

fear generalization gradients. Behavioral Neuroscience 131, 168–175.

https://doi.org/10.1037/bne0000186

Dunsmoor, J.E., Paz, R., 2015. Fear Generalization and Anxiety: Behavioral and Neural

Mechanisms. Biological Psychiatry, Understanding PTSD: From Mind to Molecules

78, 336–343. https://doi.org/10.1016/j.biopsych.2015.04.010

Elkin, L.A., Kay, M., Higgins, J.J., Wobbrock, J.O., 2021. An Aligned Rank Transform

Procedure for Multifactor Contrast Tests, in: The 34th Annual ACM Symposium on

User Interface Software and Technology. Presented at the UIST '21: The 34th Annual

ACM Symposium on User Interface Software and Technology, ACM, Virtual Event

USA, pp. 754–768. https://doi.org/10.1145/3472749.3474784

Fenton, G.E., Halliday, D.M., Mason, R., Bredy, T.W., Stevenson, C.W., 2016. Sex

differences in learned fear expression and extinction involve altered gamma

oscillations in medial prefrontal cortex. Neurobiology of Learning and Memory, MCCS

2016 135, 66–72. https://doi.org/10.1016/j.nlm.2016.06.019

Goodwin, N.L., Choong, J.J., Hwang, S., Pitts, K., Bloom, L., Islam, A., Zhang, Y.Y., Szelenyi,

E.R., Tong, X., Newman, E.L., Miczek, K., Wright, H.R., McLaughlin, R.J., Norville,

Z.C., Eshel, N., Heshmati, M., Nilsson, S.R.O., Golden, S.A., 2024. Simple

Behavioral Analysis (SimBA) as a platform for explainable machine learning in

behavioral neuroscience. Nat Neurosci 27, 1411–1424.

https://doi.org/10.1038/s41593-024-01649-9

Graham, L.K., Yoon, T., Lee, H.J., Kim, J.J., 2009. Strain and sex differences in fear

conditioning: 22 kHz ultrasonic vocalizations and freezing in rats. Psychology &

Neuroscience 2, 219–225. https://doi.org/10.3922/j.psns.2009.2.015

Greiner, E.M., Müller, I., Norris, M.R., Ng, K.H., Sangha, S., 2019. Sex differences in fear

regulation and reward-seeking behaviors in a fear-safety-reward discrimination task.

Behavioural Brain Research 368, 111903. https://doi.org/10.1016/j.bbr.2019.111903

Gruene, T.M., Flick, K., Stefano, A., Shea, S.D., Shansky, R.M., 2015. Sexually divergent

expression of active and passive conditioned fear responses in rats. eLife 4, e11352.

https://doi.org/10.7554/eLife.11352.001

Haaker, J., Maren, S., Andreatta, M., Merz, C.J., Richter, J., Richter, S.H., Meir Drexler, S.,

Lange, M.D., Jüngling, K., Nees, F., Seidenbecher, T., Fullana, M.A., Wotjak, C.T.,

Lonsdorf, T.B., 2019. Making translation work: Harmonizing cross-species

methodology in the behavioural neuroscience of Pavlovian fear conditioning.

Neuroscience and Biobehavioral Reviews 107, 329–345.

https://doi.org/10.1016/j.neubiorev.2019.09.020

Keiser, A.A., Turnbull, L.M., Darian, M.A., Feldman, D.E., Song, I., Tronson, N.C., 2017. Sex

Differences in Context Fear Generalization and Recruitment of Hippocampus and

Amygdala during Retrieval. Neuropsychopharmacol 42, 397–407.

https://doi.org/10.1038/npp.2016.174

Laxmi, T.R., Stork, O., Pape, H.C., 2003. Generalisation of conditioned fear and its

behavioural expression in mice. Behavioural Brain Research 145, 89–98.

https://doi.org/10.1016/S0166-4328(03)00101-3

Lenth, R.V., Bolker, B., Buerkner, P., Giné-Vázquez, I., Herve, M., Jung, M., Love, J., Miguez,

F., Piaskowski, J., Riebl, H., Singmann, H., 2024. emmeans: Estimated Marginal

Means, aka Least-Squares Means.

López-Moraga, A., Luyten, L., Beckers, T., 2025. Generalization and extinction of platform-

mediated avoidance in male and female rats. Sci Rep 15, 9730.

https://doi.org/10.1038/s41598-025-94265-x

[dataset] Lopez-Moraga, A., Luyten, L., Beckers, T. High threat intensity increases

conditioned fear generalization and makes extinction of a generalization stimulus less

effective. Open Science Framework. 2023. https://doi.org/10.17605/OSF.IO/9YD83

Lüdecke, D., Ben-Shachar, M.S., Patil, I., Waggoner, P., Makowski, D., 2021. performance:

An R Package for Assessment, Comparison and Testing of Statistical Models. Journal

of Open Source Software 6, 3139. https://doi.org/10.21105/joss.03139

Luyten, L., Schroyens, N., Luyck, K., Fanselow, M.S., Beckers, T., 2016. No effect of glucose

administration in a novel contextual fear generalization protocol in rats. Transl

Psychiatry 6, e903–e903. https://doi.org/10.1038/tp.2016.183

Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., Bethge, M., 2018.

DeepLabCut: markerless pose estimation of user-defined body parts with deep

learning. Nat Neurosci 21, 1281–1289. https://doi.org/10.1038/s41593-018-0209-y

Mitchell, J.R., Trettel, S.G., Li, A.J., Wasielewski, S., Huckleberry, K.A., Fanikos, M., Golden,

E., Laine, M.A., Shansky, R.M., 2022. Darting across space and time: parametric

modulators of sex-biased conditioned fear responses. Learn Mem 29, 171–180.

https://doi.org/10.1101/lm.053587.122

Nath, T., Mathis, A., Chen, A.C., Patel, A., Bethge, M., Mathis, M.W., 2019. Using

DeepLabCut for 3D markerless pose estimation across species and behaviors. Nat

Protoc 14, 2152–2176. https://doi.org/10.1038/s41596-019-0176-0

Noble, L.J., Meruva, V.B., Hays, S.A., Rennaker, R.L., Kilgard, M.P., McIntyre, C.K., 2019.

Vagus nerve stimulation promotes generalization of conditioned fear extinction and

reduces anxiety in rats. Brain Stimulation 12, 9–18.

https://doi.org/10.1016/j.brs.2018.09.013

Pearce, J.M., 1987. A Model for Stimulus Generalization in Pavlovian Conditioning.

Psychological Review 94, 61–73. https://doi.org/10.1037/0033-295X.94.1.61

Percie du Sert, N., Hurst, V., Ahluwalia, A., Alam, S., Avey, M.T., Baker, M., Browne, W.J.,

Clark, A., Cuthill, I.C., Dirnagl, U., Emerson, M., Garner, P., Holgate, S.T., Howells,

D.W., Karp, N.A., Lazic, S.E., Lidster, K., MacCallum, C.J., Macleod, M., Pearl, E.J.,

Petersen, O.H., Rawle, F., Reynolds, P., Rooney, K., Sena, E.S., Silberberg, S.D.,

Steckler, T., Würbel, H., 2020. The ARRIVE guidelines 2.0: Updated guidelines for

reporting animal research. PLoS Biol 18, e3000410.

https://doi.org/10.1371/journal.pbio.3000410

R Core Team, 2022. R: A language and environment for statistical computing (manual). R

Foundation for Statistical Computing, Vienna, Austria.

RStudio Team, 2021. RStudio: Integrated development environment for R (manual). RStudio,

PBC, Boston, MA.

Shanazz, K., Dixon-Melvin, R., Nalloor, R., Thumar, R., Vazdarjanova, A.I., 2022. Sex

Differences In Avoidance Extinction After Contextual Fear Conditioning: Anxioescapic

Behavior In Female Rats. Neuroscience 497, 146–156.

https://doi.org/10.1016/j.neuroscience.2022.06.031

Singmann, H., Bolker, B., Westfall, J., Aust, F., Ben-Shachar, M.S., Højsgaard, S., Fox, J.,

Lawrence, M.A., Mertens, U., Love, J., Lenth, R., Christensen, R.H.B., 2023. afex:

Analysis of Factorial Experiments.

Souza, R.R., Robertson, N.M., Mathew, E., Tabet, M.N., Bucksot, J.E., Pruitt, D.T., Rennaker,

R.L., Hays, S.A., McIntyre, C.K., Kilgard, M.P., 2020. Efficient parameters of vagus

nerve stimulation to enhance extinction learning in an extinction-resistant rat model of

PTSD. Progress in Neuro-Psychopharmacology and Biological Psychiatry 99,

109848. https://doi.org/10.1016/j.pnpbp.2019.109848

Struyf, D., Hermans, D., Vervliet, B., 2018. Maximizing the generalization of fear extinction:

Exposures to a peak generalization stimulus. Behaviour Research and Therapy 111,

1–8. https://doi.org/10.1016/j.brat.2018.09.005

Trott, J.M., Hoffman, A.N., Zhuravka, I., Fanselow, M.S., 2022. Conditional and unconditional

components of aversively motivated freezing, flight and darting in mice. eLife 11,

e75663. https://doi.org/10.7554/eLife.75663

Vervliet, B., Vansteenwegen, D., Baeyens, F., Hermans, D., Eelen, P., 2005. Return of fear in

a human differential conditioning paradigm caused by a stimulus change after

extinction. Behaviour Research and Therapy 43, 357–371.

https://doi.org/10.1016/j.brat.2004.02.005

Vervliet, B., Vansteenwegen, D., Eelen, P., 2004. Generalization of extinguished skin

conductance responding in human fear conditioning. Learning and Memory 11, 555–

558. https://doi.org/10.1101/lm.77404

Wickham, H., RStudio, 2022. tidyverse: Easily Install and Load the "Tidyverse."

Wong, A.H.K., Lovibond, P.F., 2020. Generalization of extinction of a generalization stimulus

in fear learning. Behaviour Research and Therapy 125, 103535.

https://doi.org/10.1016/j.brat.2019.103535

Xu, W., Südhof, T.C., 2013. A neural circuit for memory specificity and generalization.

Science 339, 1290–1295. https://doi.org/10.1126/science.1229534

Xuan, S.-M., Su, Y.-W., Liang, Y.-M., Gao, Z.-J., Liu, C.-Y., Fan, B.-F., Shi, Y.-W., Wang, X.-

G., Zhao, H., 2023. mGluR5 in amygdala modulates fear memory generalization.

Frontiers in Behavioral Neuroscience 17, 1072642.

Zhang, W.-H., Zhou, J., Pan, H.-Q., Wang, X.-Y., Liu, W.-Z., Zhang, J.-Y., Yin, X.-P., Pan, B.-

X., 2017. δ Subunit-containing GABAA receptor prevents overgeneralization of fear in

adult mice. Learn. Mem. 24, 381–384. https://doi.org/10.1101/lm.045856.117

Tables

 Table 1. Statistical test results of the manipulation check

Phase	Measured outcome	Statistical test	Result	Effect size
Habituation	Freezing	Non-parametric mixed ANOVA	Group shock: F(1, 62) = 1.58, p = 0.213 Stimuli: F(1, 62) = 0.23, p = 0.63 Gs*S: F(1, 62) = 0.11, p = 0.737	ω_p^2 < -0.01 [0,1] ω_p^2 < -0.01 [0,1] ω_p^2 < -0.01 [0,1]
Habituation	Discrimination index	Paired t-test	Stimuli: t(31) = 0.65, p = 0.52	d = 0.12 [-0.24, 0.47]
Acquisition	Freezing (trials)	Two-way ANOVA	Group shock: F(1, 60) = 1.58, p = 0.213 Sex: F(1, 60) = 10, **p = 0.002** Gs*S: F (1, 60) = 0.74, p = 0.393	ω_p^2 < -0.01 [0,1] ω_p^2 = 0.12 [0.02,1] ω_p^2 < -0.01 [0,1]
Acquisition test	Freezing	Mixed ANOVA	Group shock: F(1, 62) = 0.46, p = 0.502 Stimuli: F(1, 62) = 10.08, **p = 0.002** Gs*S: F(1, 62) = 5.22, **p = 0.026**	ω_p^2 < -0.01 [0,1] ω_p^2 = 0.03 [0,1] ω_p^2 = 0.02 [0,1]
Acquisition test	Freezing	Simple effects by group shock with Tukey correction	Moderate – Stimuli: t(124) = 2.63, **p = 0.01** Strong – Stimuli: t(124) = 0.43, p = 0.669	d = 0.66 [0.16, 1.16] d = 0.11 [-0.39, 0.6]
Acquisition test	Discrimination index	T-test	Group shock: t(62) = 1.47, p = 0.148	d = 0.37 [-0.13, 0.86]
Extinction training	Freezing (trials)	Non-parametric mixed ANOVA	Group: F(3, 60) = 8.54, **p < 0.001** Trial: F(17, 1020) = 16.06, **p < 0.001** G*T: F(51, 1020) = 1.32, p = 0.068	ω_p^2 = 0.24 [0.07,1] ω_p^2 = 0.13 [0.09,1] ω_p^2 < -0.01 [0,1]
Extinction training	Freezing (trials)	Pairwise comparisons with Tukey correction	MSCS – MSGS: t(60) = 2.58, p = 0.059 MSCS – SSCS: t(60) = -2.34, p = 0.1 MSCS – SSGS: t(60) = 1.05, p = 0.719 MSGS – SSCS: t(60) = -4.92, **p < 0.001** MSGS – SSGS: t(60) = -1.52, p = 0.43 SSCS – SSGS: t(60) = 3.39, **p = 0.006**	r = 0.18 [0.09, 0.27] r = -0.24 [-0.33, -0.15] r = 0.07 [-0.03, 0.16] r = -0.43 [-0.5, -0.35] r = -0.14 [-0.23, -0.05] r = 0.32 [0.23, 0.40]
Extinction training	Freezing (early vs late block)	Non-parametric mixed ANOVA	Sex: F(1, 56) = 4.66, **p = 0.035** Group: F(3, 56) = 8.43, **p = 0.001** Block: F(1, 56) = 93.47, **p < 0.001** S*G: F(3, 56) = 1.75, p = 0.168	ω_p^2 = 0.05 [0,1] ω_p^2 = 0.22 [0.06,1] ω_p^2 = 0.37 [0.21,1] ω_p^2 < 0.01 [0,1]

			S*B: F(1, 56) = 1.52, p = 0.223	ω_p^2 < 0.01 [0,1]
			G*B: F(3, 56) = 2.39, p = 0.078	ω_p^2 < -0.01 [0,1]
			S*G*B: F(3, 56) = 2.19, p = 0.099	ω_p^2 = 0.02 [0,1]
Extinction training	Freezing (early vs late block)	Pairwise comparisons with Tukey correction	MSCS – MSGS: t(56) = 2.7, **p = 0.044**	r = 0.27 [-0.01, 0.51]
			MSCS – SSCS: t(56) = -2.27, p = 0.12	r = -0.26 [-0.5, 0.02]
			MSCS – SSGS: t(56) = 0.76, p = 0.874	r = 0.05 [-0.23, 0.33]
			MSGS – SSCS: t(56) = -4.97, **p < 0.001**	r = -0.56 [-0.72, -0.33]
			MSGS – SSGS: t(56) = -1.95, p = 0.221	r = -0.22 [-0.47, 0.06]
			SSCS – SSGS: t(56) = 3.02, **p = 0.019**	r = 0.33 [0.06, 0.56]

724 **Table 2.** Statistical test results for the effects of generalized acquisition on generalization of extinction

Phase	Measured outcome	Statistical test	Result	Effect size
Extinction test	Freezing	Mixed ANOVA	Group: F(3, 56) = 8.45, **p < 0.001**	ω_p^2 = 0.27 [0.1,1]
			Sex: F(1, 56) = 1.39, p = 0.243	ω_p^2 < 0.01 [0,1]
			G*Se: F(3, 56) = 3.01, **p = 0.038**	ω_p^2 = 0.09 [0,1]
			Stimuli: F(1, 56) = 25.24, **p < 0.001**	ω_p^2 = 0.15 [0.03,1]
			G*St: F(3, 56) = 4.13, **p = 0.01**	ω_p^2 = 0.06 [0,1]
			Se*St: F(1, 56) = 0.02, p = 0.883	ω_p^2 < -0.01 [0,1]
			G*Se*St: F(3, 56) = 0.81, p = 0.493	ω_p^2 < -0.01 [0,1]
Extinction test	Freezing	Simple effects by stimuli with Tukey correction - CS	MSCS – MSGS: t(112) = -0.54, p = 0.949	d = -0.19 [-0.89, 0.51]
			MSCS – SSCS: t(112) = -3.20, **p = 0.01**	d = -1.13 [-1.85, -0.41]
			MSCS – SSGS: t(112) = -3.96, **p < 0.001**	d = -1.4 [-2.12, -0.67]
			MSGS – SSCS: t(112) = -2.65, **p = 0.044**	d = -0.94 [-1.65, -0.23]
			MSGS – SSGS: t(112) = -3.41, **p = 0.005**	d = -1.21 [-1.93, -0.49]
			SSCS – SSGS: t(112) = -0.76, p = 0.872	d = -0.27 [-0.97, 0.43]
Extinction test	Freezing	Simple effects by stimuli with Tukey correction - GS	MSCS – MSGS: t(112) = 2.73, **p = 0.036**	d = 0.97 [0.25, 1.68]
			MSCS – SSCS: t(112) = -1.38, p = 0.514	d = -0.49 [-1.19, 0.21]
			MSCS – SSGS: t(112) = 0.19, p = 0.997	d = 0.07 [-0.63, 0.77]
			MSGS – SSCS: t(112) = -4.11, **p < 0.001**	d = -1.45 [-2.18, -0.73]
			MSGS – SSGS: t(112) = -2.54, p = 0.06	d = -0.9 [-1.61, -0.19]
			SSCS – SSGS: t(112) = 1.58, p = 0.397	d = 0.56 [-0.15, 1.26]

Extinction test	Freezing	Simple effects by group with Tukey correction	MSCS – Sex: t(112) = 0.23, p = 0.822 MSGS – Sex: t(112) = 0.15, p = 0.882 SSCS – Sex: t(112) = -3.49, **p < 0.001** SSGS – Sex: t(112) = 0.53, p = 0.593	d = 0.08 [-0.62, 0.78] d = 0.05 [-0.65, 0.75] d = -1.23 [-1.95, -0.51] d = 0.19 [-0.51, 0.89]
Extinction test	Discrimination index	One-way ANOVA	Group: F(3, 60) = 10.89, **p < 0.001**	ω^2 = 0.32 [0.14, 1]
Extinction test	Discrimination index	Pairwise comparisons with Tukey correction	MSCS – MSGS: t(60) = -5.47, **p < 0.001** MSCS – SSCS: t(60) = -1.65, p = 0.358 MSCS – SSGS: t(60) = -3.24, **p = 0.01** MSGS – SSCS: t(60) = 3.82, **p = 0.002** MSGS – SSGS: t(60) = 2.23, p = 0.127 SSCS – SSGS: t(60) = -1.59, p = 0.39	d = -1.94 [-2.7, -1.17] d = -0.58 [-1.29, 0.12] d = -1.15 [-1.87, -0.42] d = 1.35 [0.6, 2.1] d = 0.79 [0.07, 1.51] d = -0.56 [-1.28, 0.15]

726 **Table 3.** Statistical tests for return of fear

Phase	Measured outcome	Statistical test	Result	Effect size
Reinstatement test	Freezing	Non-parametric mixed ANOVA	Group: F(3, 45) = 1.48, p = 0.231 Stimuli: F(1, 45) = 18.1, **p < 0.001** G*S: F(3, 45) = 1.44, p = 0.244	ω_p^2 = 0.03 [0,1] ω_p^2 = 0.07 [0.01,1] ω_p^2 = -0.02 [0,1]
Reinstatement test	Discrimination index	One-way ANOVA	Group: F(3, 45) = 1.62, p = 0.199	ω^2 = 0.04 [0, 1]
Spontaneous recovery test	Freezing	Mixed ANOVA	Group: F(3, 36) = 0.4, p = 0.757 Stimuli: F(1, 36) = 10.54, **p = 0.003** G*S: F(3, 36) = 1.99, p = 0.133	ω_p^2 = -0.05 [0,1] ω_p^2 = 0.06 [0,1] ω_p^2 = 0.02 [0,1]
Spontaneous recovery test	Discrimination index	One-way ANOVA	Group: F(3, 36) = 3.77, **p = 0.019**	ω^2 = 0.17 [0, 1]
Spontaneous recovery test	Discrimination index	Pairwise comparisons with Tukey correction	MSCS – MSGS: t(36) = -2.78, **p = 0.041** MSCS – SSCS: t(36) = 0.07, p = 0.999 MSCS – SSGS: t(36) = -1.2, p = 0.631 MSGS – SSCS: t(36) = 2.99, **p = 0.024** MSGS – SSGS: t(36) = 1.62, p = 0.378 SSCS – SSGS: t(36) = -1.33, p = 0.549	d = -1.28 [-2.26, -0.3] d = 0.03 [-0.88, 0.94] d = -0.55 [-1.49, 0.39] d = 1.31 [0.37, 2.25] d = 0.73 [-0.2, 1.65] d = -0.58 [-1.48, 0.31]

	Habituation	Acquisition	Acquisition Test	Extinction	Extinction Test	Reinstatement	Reinstatement Test	Spontaneous Recovery Test

MSCS

MSGS

SSCS

SSGS

♪ CS   ♪ GS

7 days

a Extinction training

MODERATE SHOCK → MSCS, MSGS

STRONG SHOCK → SSCS, SSGS

18x ♪ CS or ♪ GS

b MSCS, MSGS, SSCS, SSGS

c Female — * — Male