Small RNAs regulation and genomic harmony: insights into allopolyploid evolution in
marsh orchids (*Dactylorhiza*)

Mimmi C. Eriksson[1,2,3*], Matthew Thornton[1], Emiliano Trucchi[1,4], Thomas M. Wolfe[1,5],
Francisco Balao[1,6], Mikael Hedrén[7], Ovidiu Paun[1*]

[1]Botany and Biodiversity Research, University of Vienna, Rennweg 14, A-1030 Vienna,
Austria. [2]Vienna Graduate School of Population Genetics, Veterinärplatz 1, A-1210 Vienna,
Austria. [3]Department of Plant Biology, Swedish University of Agricultural Sciences, SE-750
07 Uppsala, Sweden. [4]Marche Polytechnic University, Ancona, Italy. [5]Institute of Forest
Entomology, Forest Pathology and Forest Protection (IFFF), BOKU, Peter-Jordan-Straße
82/I, 1190 Vienna, Austria. [6]Departamento de Biologia Vegetal y Ecologia, University of
Seville, Sevilla, Spain. [7]Department of Biology, Lund University, Sölvegatan 37, S-22362,
Lund, Sweden.

*Authors for correspondence: mimmi.c.eriksson@gmail.com; ovidiu.paun@univie.ac.at

ORCID
Mimmi C. Eriksson 0000-0002-9548-8226
Emiliano Trucchi 0000-0002-1270-5273
Thomas M. Wolfe 0000-0002-1509-0147
Francisco Balao 0000-0003-2104-3846
Mikael Hedrén 0000-0001-8194-4617
Ovidiu Paun 0000-0002-8295-4937

Total word count	5458	Acknowledgements:	169
Summary	196	Author contributions:	90
Introduction:	1094	No. figures:	5
Materials and Methods:	1069	No. tables:	0
Results:	1426	No Supporting information:	19 (Figs S1-S10, Tables S1-S9)
Discussion:	1869		

SUMMARY

▢   Hybridization and polyploidy are major drivers of plant diversification, often
accompanied by shifts in gene expression and genome composition. Small RNAs
(smRNAs) are thought to influence such genomic changes, particularly through their
interactions with transposable elements (TEs).
▢   We comparatively quantified smRNAs in established sibling allopolyploids
Dactylorhiza majalis and *D. traunsteineri* and their diploid progenitors to assess how
independent allopolyploidization events shaped smRNA landscapes.
▢   Despite independent origins, the allotetraploids exhibited substantial overlap in
smRNA composition, including transgressive accumulation of smRNA near genes
related to transcriptional regulation, cell division and stress response. Consistently,
TE-associated 24 nt smRNAs more closely resembled the paternal and larger
genome, while shorter smRNAs typically reflected the maternal and smaller genome.
Nevertheless, distinct patterns were also evident: the older *D. majalis* showed greater
accumulation of smRNAs near genes involved in transcriptional and translational
regulation, while the younger *D. traunsteineri* displayed stronger non-additive
patterns, suggesting ongoing resolution of post-polyploid meiotic and mitotic
instability.
▢   Our results reveal both convergence and divergence in smRNA landscapes among
independently formed allopolyploids. Our study highlights the central role of smRNAs
in resolving genomic conflict, with possible implications for functional divergence and
ecological innovation during polyploid evolution.

Key words
allopolyploidy, *Dactylorhiza*, gene regulation, marsh orchids, small RNAs, subgenome
dominance, transposable elements

Plain-language summary

We studied two sibling marsh orchid species that arose through hybridization and genome
doubling. By comparing their small RNA molecules and how these associate with genes, we
found both shared and species-specific patterns. These differences reflect how the orchids'
genomes have changed since formation, and may explain how repeated polyploidy events
contribute to genetic and ecological diversity in plants.

INTRODUCTION

Allopolyploidization—the combination of whole-genome duplication (WGD) and interspecific hybridization—can be a potent evolutionary force, generating genetic diversity, gene redundancy, and buffering capacity that may contribute to evolutionary innovation and ecological diversification. However, the initial merger of divergent genomes into a single nucleus can destabilize genome regulation, triggering "genomic shock" (McClintock, 1984), on the background of a severe genetic bottleneck.

Small noncoding RNAs (smRNAs) have emerged as key mediators during post-polyploidization genome stabilization (e.g., Ha *et al.*, 2009; Shen *et al.*, 2017; Cavé-Radet *et al.*, 2020). These short RNA molecules suppress transposable elements (TEs) and fine-tune gene expression, promoting both genome integrity and adaptive responses. Specifically, repeat-associated or heterochromatic small-interfering RNAs (ra-siRNAs/het-siRNAs; Shen *et al.*, 2017) help maintain chromatin stability, while microRNAs (miRNAs) can modulate gene expression with phenotypic implications (Ha *et al.*, 2009; Ghani *et al.*, 2014; Jiang *et al.*, 2021). In addition, plant smRNAs orchestrate abiotic and biotic stress responses (Kaur *et al.*, 2021; Tiwari & Rajam, 2022; Zhan & Meyers, 2024), potentially facilitating adaptation to novel, non-parental environments. These dual roles make smRNAs integral to allopolyploid success.

In general, smRNAs operate through three main mechanisms: (1) post-transcriptional gene silencing (PTGS), binding to RNAs to repress translation or promote cleavage and degradation; (2) transcriptional gene silencing (TGS), directing epigenetic silencing at homologous genomic loci; and (3) splicing regulation, interacting with pre-mRNAs to influence splice-site selection (Borges & Martienssen, 2015; Wang & Chekanova, 2016). In plants, smRNAs are classified by biogenesis into miRNAs (20–22 nt long, hairpin-derived), and small-interfering RNAs (siRNAs, from double stranded RNA) (Borges & Martienssen, 2015; Chen & Rechavi, 2021), with siRNAs further divided into: (*i*) ra-siRNAs (24 nt and primarily involved in TGS), (*ii*) secondary siRNAs (~21–22nt long, which can function in both TGS and PTGS pathways), and (*iii*) natural antisense transcript siRNA (nat-siRNA, 21–24 nt, acting both in *cis* and *trans* via PTGS) (Borges & Martienssen, 2015; Chen a& Rechavi, 2021).

Recent reviews (e.g., Ariel & Manavella, 2021; Tossolini *et al.*, 2025) highlight the pivotal link between TE mobilization and smRNA evolution. TEs not only contribute regulatory capacity through enhancer-derived transcripts (e.g., in wheat; TE-initiated enhancer-like elements, Xie *et al.*, 2023), but also spur the emergence of smRNAs and long non-coding RNAs that

diversify gene regulation and support genome plasticity (Hassan *et al.*, 2024; Tossolini *et al.*,
2025).

WGD and environmental stress are known to lead to TE activation, resulting in accumulation
of repeat-derived smRNAs, boosting both TGS and PTGS pathways (Ito *et al.*, 2011; Kenan-
Eichler *et al.*, 2011; Marí-Ordóñez *et al.*, 2013; Fultz *et al.*, 2015; Xiao & Wang, 2025). For
example, resynthesized *Arabidopsis* allotetraploids rapidly restore ra-siRNA populations
after WGD, while miRNAs show non-additive expression patterns and play key roles in
stabilizing hybrid genomes (Ha *et al.*, 2009). Similarly, early generations *Brassica napus*
allotetraploids exhibit a surge in TE-derived 21 nt siRNAs, especially from Gypsy elements,
helping to silence newly active TEs (Martinez Palacios *et al.*, 2019). Cavé-Radet *et al.* (2020,
2023) observed stronger miRNA responses in allopolyploid *Spartina* compared to homoploid
hybrids, suggesting a more pronounced regulatory impact of WGD than hybridization alone
(Cavé-Radet *et al.*, 2020). On the other hand, the divergence between progenitors
determines the extent of genomic shock in wheat allotetraploids, including enhanced smRNA
responses associated with TEs in hybrid crosses between parents with differing TE load,
compared to those with similar TE content (Jiao *et al.*, 2018).

Despite growing interest in the role of smRNAs in allopolyploid evolution, most research has
focused on synthetic or early-generation polyploids. To our knowledge, only Cavé-Radet *et*
al., (2020) have examined smRNAs in wild allopolyploids, albeit also in early stages
(*Spartina* F$_1$ hybrids and *S. anglica*, ~170 years old). Thus, the long-term dynamics,
regulatory roles and potential ecological relevance of smRNAs in more established
allopolyploid lineages remain poorly explored.

In this study, we focus on two ecologically-distinct, established, sibling orchid allotetraploids,
Dactylorhiza majalis s.str. (Rchb.) P.F.Hunt & Summerh. and *D. traunsteineri* (Saut. ex
Rchb.) Soó, and their parents, to address the long-term fate and potential roles of smRNAs
following allopolyploidization. The marsh orchids within *D. majalis s.l.* include several
independently formed sibling allotetraploids, each likely with multiple origins involving
overlapping stocks of parental populations (Brandrud *et al.*, 2020). The focal allotetraploids
here share a complex history of repeated polyploidization and/or hybridization and back-
crossing, but are today reasonably well separated in morphology and habitat preferences
(Fig. 1a; Wolfe *et al.*, 2023). They are estimated to have formed approximately 104k (*D.*
majalis) and 74k (*D. traunsteineri*) generations ago (Hawranek, 2021), from the diploid
progenitors *D. fuchsii* (Druce) Soó (maternal, smaller genome) and *D. incarnata* (L.) Soó
(paternal, larger genome) (Fig. 1a) (Aagard *et al.*, 2005; Brandrud *et al.*, 2020; Eriksson *et*
al., 2022). Transcriptomic comparisons of the diploid progenitors have revealed large-scale
gene expression differences, importantly, including differences in smRNAs associated with

TEs and PTGS pathways (Balao *et al.*, 2017). Further, Eriksson *et al.* (2022) reported

significant differences in both genome size and TE load between the two diploid progenitors.

Distinct ecological, spatial, morphological and physiological properties have been reported

for the two sibling allopolyploids (Paun *et al.*, 2010, 2011; Balao *et al.*, 2016; Wolfe *et al.*,

2023)*. D. majalis* is distributed across continental Europe, ranging from the Pyrenees to

Scandinavia, with a broader ecological niche (i.e., mesic to moist meadows) and less

geographically-driven genetic differentiation than *D. traunsteineri* which is found in disjunct

marshes and fens in the Alps, Scandinavia and British Isles. Wolfe and colleagues (2023)

reported physiological differences in photosynthesis and nutrient transport consistent with

the contrasting ecological conditions preferred by the two allotetraploids. These appear to

have helped the two allotetraploids adapt to different soil chemistry conditions, with *D.*

traunsteineri restricted to wet, nutrient-poor soils, in particular with extremely low levels of

available nitrate (Wolfe *et al.*, 2023). Furthermore, reciprocal transplant experiments have

shown that much of the gene expression divergence between the sibling allopolyploids is

environmentally plastic, and appeared to be associated to distinct, environmentally-driven,

biotic communities in their roots (Emelianova *et al.*, 2025). In stark contrast, analysis of the

genomes of several *Dactylorhiza* allopolyploids found relatively little difference between *D.*

majalis and *D. traunsteineri* both in genome size and TE activity, suggesting a strong control

of TEs following recurrent allopolyploidizations (Eriksson *et al.*, 2022).

In this study, we specifically investigate: (1) how smRNA abundance in the allopolyploids

reflects parental patterns—whether it follows additive, dominant, or transgressive trends—

and how consistent these patterns are across the sibling allopolyploids; (2) the extent and

similarity of smRNA association with TEs in the two sibling allotetraploids; and (3) how

variation in smRNA patterns may relate to distinct eco-physiological traits of *D. majalis* and

D. traunsteineri.

MATERIALS AND METHODS

Sampling

Adult specimens of each species of interest were transplanted from multiple localities across

Europe to a common garden in Vienna, Austria. *Dactylorhiza* includes perennial plants that

produce a new tuber yearly, which supports growth in the following year. Two years after

transplantation, in the morning of 14 May 2014, leaf tissue at a similar developmental stage

for all accessions was fixed in RNAlater (Sigma) and stored at -80°C until further processing.

Our final sampling included four accessions of each diploid (*D. incarnata* and *D. fuchsii*),

together with nine (*D. majalis*) and twelve (*D. traunsteineri*) individuals of each allotetraploid

(Supplementary Table S1). With few exceptions, each accession of each species originated

from different localities across Europe. The diploid smRNA data has been previously

investigated by Balao *et al.* (2017). Independent RNA isolations from the same tissue

fixations of *D. majalis* and *D. traunsteineri* were used for RNA-seq analyses by Wolfe *et al.*

(2023).

Library preparations and sequencing

Small RNA extraction, library preparation and sequencing for this experiment was previously

described for diploid investigations (Balao *et al.*, 2017). In short, isolation of smRNA was

performed with the mirVana miRNA Isolation Kit (Life Technologies). The smRNA-enriched

RNA extracts were further purified by gel size selection on 15% TBE-Urea pre-casted gels

(Life Technologies). The small RNA libraries were prepared with the NEBNext Multiplex

Small RNA Library Prep Set for Illumina (New England Biolabs). Illumina sequencing as 50

bp single-end reads was performed at the Vienna BioCenter Core Facilities (VBCF;

https://www.viennabiocenter.org/).

Read processing and quantifying

The libraries were demultiplexed using BamIndexDecoder v.1.03 (http://wtsi-

npg.github.io/illumina2bam/#BamIndexDecoder). Raw reads were adapter-trimmed, size-

selected for the 20-24 nt range, and then quality filtered using CLC Genomics Workbench

v.8.0 (QIAGEN). Filtered reads were mapped to the *Dactylorhiza incarnata* reference

genome v.1.0 (Wolfe *et al.*, 2023) with STAR v.2.7.3a (Dobin *et al.*, 2013), using --

alignIntronMax 1 and --outFilterMismatchNoverLmax 0.05. A single-pass alignment

approach was used, retaining multi-mapping reads below a threshold of 100 (--

outFilterMulitmapNmax 100).

An annotation-free approach was initially used to identify genomic regions (genic and non-

genic) that were potentially "targeted" by smRNAs, defined here as regions showing

sequence homology with smRNAs and therefore indicating an association between smRNAs

and the corresponding genomic sequence. Throughout this study, we use the term

"targeting" in a purely operational sense to describe genomic regions showing sequence

complementarity to smRNAs. This definition does not assume a direct causal or regulatory

interaction between smRNAs and the corresponding genomic loci. Rather, it denotes

potential sites of smRNA association inferred from sequence homology. While such

associations often coincide with transcriptionally or post-transcriptionally regulated regions

(e.g. in the case of heterochromatic siRNAs or miRNAs; Borges & Martienssen, 2015), they

may also reflect indirect or distal effects, or even passive sequence associations without

immediate regulatory function.

After sorting bam files by reference coordinates with Samtools v.1.10 (Danacek *et al.*, 2011),

smRNA read coverage was assessed for each individual in non-overlapping 100 bp windows

across the genome using bamCoverage from deepTools2 v.3.5.1 (Ramirez *et al.*, 2016).

Windows with a minimum coverage of ten reads in at least one sample were retained in a

bed file, and adjacent windows were merged using bedtools merge v.2.29.2 (Quinlan & Hall,

2010). The resulting bed file of retained regions (i.e., peaks of smRNA "targeting") was

converted to gff3 format using a custom python script. Counts were then obtained with

featureCounts from Subread v.2.0.0 (Liao *et al.*, 2014), accounting for multiple mappers

fractionally (-M --fraction), to include potential trans-associations. The full pipeline, scripts,

raw count file, and peak gff3 file are available at https://github.com/mc-er/dact-sRNAs.

202   Differential smRNA "targeting" analysis

Differently "targeted" (DT) regions were identified following a similar approach to that used

for detecting differentially expressed genes, using counts of smRNAs at peaks as input.

Analyses were performed in R v.4.1.1 and the Bioconductor package EdgeR v.3.34.0

(Robinson *et al.*, 2010; McCarthy *et al.*, 2012). The count matrix obtained from

featureCounts was filtered to retain genomic regions with a minimum smRNA mapping of

one count per million (CPM) in at least 25% of samples within a group and a minimum of two

samples overall. To account for differences in library size, data were normalised using the

weighted trimmed mean of M-values (TMM) method. A quasi-likelihood negative binomial

model was applied to identify DT regions across the genome, comparing either (i) the two

sibling allotetraploids or (ii) each allotetraploid to its parental diploid. P-values were corrected

for multiple testing using the Benjamini–Hochberg (BH) procedure, and only genomic

regions passing an FDR < 0.05 were kept for further analysis.

Analyses of genomic regions around TEs were performed separately for reads of 20–23 nt,

and for 24 nt reads (the latter representing potentially heterochromatic smRNAs, hetsiRNAs;

Borges & Martienssen, 2015). For analyses of genic regions, all read lengths (20–24 nt)

were combined into a single analysis per comparison.

219   Assessing genomic interactions

Differentially targeted peaks passing an FDR threshold of 0.05 were classified into three

categories of genomic interactions (Supplementary Table S2): (1) additivity—the polyploid

showed lower targeting relative to one diploid and higher targeting relative to the other, (2)

transgressive—the polyploid showed increased or decreased targeting relative to both

diploids, and (3) dominance—the polyploid showed no significant difference relative to one

diploid (the "dominant" parent) but differed significantly relative to the other diploid (illustrated

on the Y-axis of Fig. 4c-f). Deviation from the stochastically expected number of shared

peaks between the two tetraploids in each category was tested using a hypergeometric test

implemented in the Bioconductor package SuperExactTest (Wang *et al.*, 2015).

Functional interpretation of differentially targeted genes

To gain insights into potential roles for smRNA associations in shaping the divergent

ecological properties of the sibling allopolyploids, we performed gene ontology (GO)

enrichment analysis of DT genes. Because the molecular consequences of smRNA

association depend on the location of the targeted region, we analysed regions around

promoters separately from those within exons or introns. Genes associated with smRNAs

around promoters would be expected to be transcriptionally down-regulated or silenced,

whereas associations within gene bodies may indicate post-transcriptional effects or altered

splicing. GO enrichment analyses for genes potentially affected by DT peaks were

performed in R v.4.1.1 using the Bioconductor package topGO v.2.44.0 (Alexa &

Rahnenfuhre, 2021). Fisher's exact tests were run using the "weight01" algorithm to test for

over-representation of specific functional categories. P-values were adjusted for multiple

testing using the FDR method, with significance set at FDR < 0.05.

RESULTS

smRNAs and their genomic targets

Sequencing produced an average 13.5 M reads per accession (±5.9 M SD). After size

selection and quality filtering, an average of 5.8 M reads (±3.2 M SD) of 20–24 nt were

retained (Supplementary Table S3). On average, 83.1% (±4.2% SD) of these reads mapped

to the *D. incarnata* reference genome, with no apparent mapping bias across species

(Supplementary Table S3). Read length distributions matched the expected profile for

smRNAs (Borges & Martienssen, 2015), with modes at 21–22 nt and 24 nt in all species

(Fig. 2a).

D. fuchsii exhibited higher CPM of 20–22 nt smRNAs compared to *D. incarnata*, with a

significant difference observed only for 21 nt smRNAs (Fig 2a–c). The reverse trend was

found for the 23–24 nt smRNAs, which were more abundant in *D. incarnata* (Fig. 2a). The

two allotetraploids were more similar to each other than the diploids (Fig. 2a). For 21 nt

smRNAs, both allotetraploids displayed significantly higher CPM than the paternal species

(*D. incarnata*), but significantly lower CPM than the maternal species (*D. fuchsii*) (Fig. 2a). A

large proportion of 20–22 nt smRNAs mapped to exonic regions, while 24 nt smRNAs

predominantly mapped to TEs and intergenic regions (Fig. 2b–c, Supplementary Fig. S1).

Our annotation-free approach identified 2,454,439 putative smRNA peaks across the

reference genome. After filtering for CPM > 1 in at least two samples and 25% of samples

per group, between 102,222 and 370,440 peaks were retained depending on the

comparison. A multidimensional scaling (MDS) plot based on all 20–24 nt smRNAs across

212,434 peaks clearly separated the two diploid species, and distinguished diploids from

allotetraploids, with the latter overlapping centrally (Fig. 1b). Within *D. traunsteineri* a

geographical clustering was apparent (not shown).

smRNA patterns in ecologically divergent allopolyploids

We compared the two sibling allotetraploids using the combined 20–24 nt reads, retaining

152,503 peaks passing the CPM threshold (Supplementary Table S4). Among these, 2,106

DT peaks were identified between *D. traunsteineri* and *D. majalis*, of which approximately

half (1,016 peaks) were associated with genes. Gene-associated peaks were defined as

those overlapping a gene model (exons or introns) or within 1,000 bp upstream of a gene

model (hereafter referred to as 'promoters'; Supplementary Table S4).

Most genic DT peaks (762) were located within introns, and 82 overlapped multiple genic

features (intron, exon, and/or promoters) (Supplementary Table S4). Across all genic

categories, the older allotetraploid *D. majalis* displayed approximately three times as many

peaks with high smRNA targeting compared to *D. traunsteineri* (Supplementary Table S4,

Fig. 3a–c).

Comparison of DT peaks with previously identified differentially expressed (DE) genes from

the same experiment (Wolfe *et al.*, 2023) revealed that most genes with DT peaks did not

show differential expression between the two allopolyploids, consistent with potential post-

transcritional effects or alternative splicing rather than direct transcriptional regulation (Fig.

3f, Supplementary Table S5).

GO enrichment analyses of genes associated with DT peaks between the allotetraploids

revealed more enriched terms in *D. majalis* than *D. traunsteineri* (Fig. 5a, Supplementary

Table S6). Most enriched terms were for genes targeted within gene bodies and related to

transcription/translation regulation, transmembrane transport, response to biotic and abiotic

stress, DNA repair, and cell cycle control (Supplementary Table S6).

smRNA targeting and genomic interactions

We further compared 20-24 nt smRNA patterns of each allotetraploid to those of the parental

diploids. After filtering, 212,434 peaks were retained (Supplementary Table S4). Both

allotetraploids showed broadly similar targeting patterns relative to the diploids, although the

younger *D. traunsteineri* displayed greater divergence than *D. majalis* (Supplementary Table

S4, Supplementary Fig. S2). Allotetraploids showed larger differences from the maternal

parent (*D. fuchsii*) than from the paternal *D. incarnata* (Supplementary Table S4,

Supplementary Fig. S2).

DT peaks were classified into four interaction types: additive, transgressive, and dominance

towards either diploid. The majority of peaks showed no change relative to either diploid.

Among DT peaks, non-additive patterns predominated, with dominance most frequent,

particularly towards *D. incarnata* (Supplementary Table S7, Fig. 4a–f, Supplementary Fig.

S3–S4). Notably, in intergenic regions lacking annotated TEs, a higher proportion of smRNA

peaks showing dominance towards *D. fuchsii* than in other genomic contexts

(Supplementary Table S7, Fig. 4d, Supplementary Fig. S3). Among transgressive peaks,

both allotetraploids most often displayed higher smRNA targeting relative to both diploids

(Supplementary Table S7, Fig. 4a, Fig. 4c–f, Supplementary Fig. S3–S4). A notable general

pattern across interaction types was that a large proportion of DT peaks were shared

between the sibling allotetraploids (Supplementary Table S7, Fig. 4b–c, Supplementary Fig.

S3–S4), despite their independent origins (Brandrud *et al.*, 2020; Hawranek, 2021).

We also examined the distribution of interaction categories across genic regions (promoters,

exons, introns). In *D. majalis*, nearly 75% promoter-associated DT peaks showed dominance

towards *D. incarnata*, and almost 50% showed lower targeting than *D. fuchsii*

(Supplementary Fig. S5). In contrast, in *D. traunsteineri* dominance toward *D. fuchsii* was

more frequent across all genic regions (Supplementary Fig. S5).

Transgressively targeted genes in both allotetraploids were enriched for similar functional

categories, including stress response, transport, transcription/translation regulation, and cell

cycle control, though largely involving different genes and pathways in each species (Fig.

5b–c, Supplementary Tables S8–S9). In *D. traunsteineri*, transgressive peaks were

associated with enrichment for cell cycle regulation and stress response functions, and the

respective genes were targeted within exons or introns (Fig. 5b, Supplementary Tables S8–

S9). In *D. majalis*, transgressive peaks occurred mostly within introns, and enrichments were

associated with regulation of gene function (Fig. 5c, Supplementary Tables S8).

smRNAs associations with TEs

For TE-focused analyses, reads were divided into 24 nt smRNAs (likely het-siRNAs and

therefore expected to be more frequently associated with TEs; Fig2c, Supplementary Fig.

S1) and 20-23 nt smRNAs. After CPM filtering, for the comparative analysis of *D. majalis*

versus *D. traunsteineri* more 24 nt peaks (315,838) passed the threshold than 20–23 nt

peaks (102,222) (Supplementary Table S4). However, DT analysis uncovered the opposite

trend: 3,242 DT peaks were identified for the 20–23 nt smRNAs, compared to only 103 DT

peaks for the 24 nt smRNAs (Supplementary Table S4, Fig. 3d–e) suggesting a consistent

targeting of TEs by het-siRNAs across allopolyploids. Very few TE-associated peaks were

DT between the allotetraploids in both smRNA size classes simultaneously (Supplementary

Fig. S6). In both datasets, approximately half of the DT peaks overlapped annotated TEs

(Supplementary Table S4). Many of these were also located within gene introns, while a

smaller number were found in the vicinity of gene promoters (Supplementary Table S4).

Similar to the results for the genic regions, *D. majalis* exhibited more frequent higher smRNA

targeting of TEs than *D. traunsteineri* among DT peaks, except for 20–23 nt peaks

associated with TEs located around promoters, where *D. traunsteineri* showed higher

targeting (Supplementary Table S4). The majority of TE-associated DT peaks were located

within LTR retrotransposons, especially Ty3-*gypsy* elements (Supplementary Fig. S7–S8).

Among smRNAs targeting Ty3-*gypsy* elements, a slightly higher number targeted non-

chromoviruses than chromoviruses (Supplementary Fig. S8).

Comparisons of allotetraploids versus diploids showed similar trends. After filtering, 139,736

(20-23 nt) and 370,440 (24 nt) peaks were retained (Supplementary Table S4). As with genic

targets, DT peaks were classified into additive, transgressive, and dominance towards either

diploid. In both datasets, a significant number of DT peaks were found to be shared between

D. majalis and *D. traunsteineri* across interaction categories (p < 0.001, hypergeometric test;

Fig 3d-e, Supplementary Fig. S3, Fig. 4e–f, Supplementary Fig. S7, Supplementary Fig. S9–

10).

Dominance patterns again predominated, with more peaks showing dominance towards

either diploid than additive or transgressive patterns. For 20–23 nt smRNAs, slightly more

peaks showed dominance toward the diploid with lower targeting, whereas for 24 nt

smRNAs the opposite trend was observed (Fig. 4e–f, Supplementary Fig. S3,

Supplementary Fig. S9). The contrast between dominant types (higher vs. lower targeting)

was less pronounced for 20–23 nt but more distinct for 24 nt smRNAs (Fig. 4e–f,

Supplementary Fig. S3, Supplementary Fig. S9–10). Three general patterns observed for

smRNA-associated gene regions were also evident for TEs. First, for most additive peaks,

the smaller genome of *D. fuchsii* exhibited higher smRNA targeting than the larger *D.*

incarnata (Fig. 4e–f, Supplementary Fig. S3, Supplementary Fig. S9–10). Second, for

transgressive peaks, both allotetraploids generally showed elevated smRNA targeting than
either diploid (Fig. 4e–f, Supplementary Fig. S3, Supplementary Fig. S7, Supplementary Fig.
S9–10). Finally, *D. traunsteineri* displayed greater differences from the parental diploids than
D. majalis (Fig. 4e–f, Supplementary Fig. S3, Supplementary Fig. S7).

DISCUSSION

smRNAs are key regulators of genome integrity and gene expression in polyploids, but most
insights so far come from recent or resynthesized polyploids. In this study, we examine
natural polyploids that have evolved over extended evolutionary timescales to provide a
comprehensive overview of potential smRNA contributions to genomic and transcriptomic
diversification following repeated allopolyploidization in *Dactylorhiza majalis* s.l. We analysed
smRNA expression in two established sibling allotetraploids and their diploid progenitors, all
grown under common garden conditions, thus minimizing environmental sources of
variation. Our primary objectives were to comparatively characterize (*i*) patterns of genomic
interaction in the allopolyploids and (*ii*) smRNA associations to TE control and gene activity.

Inferring smRNA *"targeting"* from sequence homology offers an informative yet indirect view
of smRNA–genome interactions. Many plant smRNAs—particularly 24 nt siRNAs—act
locally to reinforce transcriptional silencing of homologous TEs or nearby genes (Matzke &
Mosher, 2014; Borges & Martienssen, 2015). Trans-acting and distal effects have
nonetheless been documented (Lewsey *et al.*, 2016; Liu & Chen, 2018), though the
prevalence and biological impact of such long-distance effects remain less well
characterized. Therefore, our analyses emphasize patterns of smRNA association rather
than assuming direct functional regulation. These patterns highlight potential regions of
epigenetic remodeling and provide a comparative framework for understanding smRNA
landscape evolution after allopolyploidization.

We acknowledge that the present design—limited to representative individuals of the two
extant diploid progenitors—cannot encompass the full spectrum of genetic and regulatory
polymorphism that may have existed at the time of allopolyploid formation. Given that both
parental lineages likely accumulated substantial variation over tens of thousands of
generations, some of the observed differences between the allopolyploids may partly reflect
ancestral or subsequent divergence in the diploid gene pools. Accordingly, we interpret our
results in terms of general and reproducible regulatory patterns rather than exact
reconstructions of the parental regulatory states.

Consistent dominance in the allopolyploids towards the paternal, larger genome

Allopolyploidization merges divergent genomes, giving rise to novel genetic and regulatory
interactions. Common outcomes include epistasis and non-additive genetic or epigenetic
effects, including smRNA-mediated gene regulation and TE silencing. Our results reveal a
consistent dominance of smRNA targeting aligned to the paternal, larger subgenome (*D.*
incarnata), especially for TE-associated 24 nt smRNAs. This pattern was observed across
both sibling allotetraploids (Fig. 4, Supplementary Fig. S5), and resulted in a comparatively
small number of DT peaks between them. Similar non-additive smRNA responses following
hybridization and/or whole-genome duplication have been documented in *Arabidopsis* (Ha *et*
al., 2009), *Brassica* (Fu *et al.*, 2016; Martinez Palacios *et al.*, 2019), and *Spartina* (Cavé-
Radet *et al.*, 2020), though it has remained unclear whether such patterns remain consistent
across longer evolutionary timescales.

Examining whether smRNA expression is additive, dominant, or transgressive allows us to
understand how regulatory interactions from the parental genomes combine in
allopolyploids. Additive patterns indicate intermediate regulation and minimal epistatic
interactions, dominance reflects the stronger influence of one parental subgenome, and
transgressive patterns reveal novel regulatory states beyond the parental range. These
distinctions are conceptually important because they highlight mechanisms of genome
stabilization, resolution of redundant gene copies, and the generation of potentially adaptive
phenotypes that can enable allopolyploids to occupy ecological niches outside the parental
range (Ha et al., 2009; Fu et al., 2016; Chen et al., 2020).

Given that *D. incarnata* has a larger genome and higher TE load than *D. fuchsii* (Eriksson *et*
al., 2022), our findings are consistent with prior reports showing that smRNA-mediated
silencing preferentially targets the TE-rich subgenome, as observed in wheat (Woodhouse *et*
al., 2014) and *Brassica rapa* (Cheng *et al.*, 2016). In contrast, gene expression dominance in
allopolyploids frequently shifts toward the subgenome with fewer TEs, a pattern that can
arise rapidly after allopolyploidization (for example in *Mimulus*; Edger *et al.*, 2017) and
persist across generations (Gaebelein *et al.*, 2019; Liu & Wang, 2023).

While our data indicate stronger dominance towards *D. incarnata*, some patterns may be
influenced by reference genome bias, as we only had access to a *D. incarnata* reference
genome. Accurate differentiation between homologous and homeologous targeting remains
challenging due to the short length of smRNAs and their tolerance for mismatches in plants.
Nonetheless, mapping success was comparable across samples (ca. 83%; Supplementary
Table S3), and intergenic smRNA dominance towards *D. fuchsii* was also observed,
suggesting that reference bias alone cannot account for our observations. Lineage-specific
TE degradation or divergence in *D. fuchsii* may have reduced smRNA detection from this

genome. Notably, *D. incarnata* and *D. fuchsii* diverged approximately 5.5 million years ago

(Brandrud *et al.*, 2020).

Divergent targeting of TEs by 24 nt and 20-23 nt smRNAs

As expected, 24 nt smRNAs—typically het-siRNAs—were strongly associated with TEs,

particularly LTR retrotransposons (Fig. 2c, Supplementary Fig. S1). Surprisingly, the number

of DT peaks for 24 nt smRNAs between the allotetraploids was low, suggesting that TE

silencing proceeds in a relatively consistent manner across these independently evolved

lineages (Fig.3e). In contrast, 20–23 nt smRNAs exhibited markedly greater divergence

between the sibling allopolyploids (Fig. 3d), consistent with a more dynamic, lineage-specific

regulatory role.

Previous studies have shown that TE-associated smRNA activity often peaks shortly after

hybridization or whole-genome duplication, followed by stabilization via DNA methylation or

other epigenetic mechanisms (Jiao *et al.*, 2018; Martinez Palacios *et al.*, 2019). In

Arabidopsis, Ha *et al.*, (2009) proposed that allopolyploids exploit smRNAs inherited from

the parents to buffer genomic shock and promote rapid stabilization. The comparatively low

divergence in 24 nt smRNA targeting observed here may reflect the advanced age of these

natural allotetraploids—approximately 74k (*D. traunsteineri*) and 104k generations (*D.

majalis*) (Hawranek, 2021). Nonetheless, for the few DT peaks detected, smRNA abundance

was higher in the older *D. majalis* (Fig. 3), possibly indicating continued regulatory

refinement associated with its broader ecological range or a more advanced stage of

genome diploidization.

Intriguingly, most smRNA differences associated with TE regulation in the ecologically

divergent allotetraploids involved 20–23 nt smRNAs, which likely include miRNAs and nat-

siRNAs. These classes are frequently implicated in responses to biotic and abiotic stress (Ito

et al., 2011; Marí-Ordóñez *et al.*, 2013; Creasey *et al.*, 2014; Lee *et al.*, 2020). Elevated

expression of miRNAs (typically 20–22 nt long) has also been observed in resynthesized

allopolyploids relative to diploid progenitors across several systems (Ha *et al.*, 2009; Kenan-

Eichler *et al.*, 2011; Ghani *et al.*, 2014), underscoring the potential role of shorter smRNAs in

stress adaptation and dynamic gene regulation in polyploids.

Despite *Ty1-copia* being the most abundant LTR-TE in *Dactylorhiza* genomes (Eriksson *et

al.*, 2022), we observed more 20–23 nt smRNAs mapping to, and possibly derived from,

Ty3-gypsy elements (Supplementary Figs. S6–S7), particularly non-chromovirus subgroups.

Notably, *Ty3-gypsy* elements displayed contrasting patterns between the diploid progenitors:

chromovirus subgroups were more prevalent in *D. fuchsii*, while non-chromoviruses were

more common in *D. incarnata* (Eriksson *et al.*, 2022). Elevated targeting of *Ty3-gypsy*

elements by shorter smRNAs in allopolyploids may reflect subgenome conflict or TE

activation following hybridization. A similar phenomenon occurs in *Gossypium*, where TEs

originally in the higher-TE A-subgenome transposed into the D-subgenome post-

allopolyploidization (Chen *et al.*, 2020), suggesting that transposition between subgenomes

may contribute to regulatory instability and diversification.

Genic smRNA targeting and possible regulatory implications

Polyploid formation introduces both meiotic/mitotic challenges and extensive genetic

redundancy. Accurate differentiation between homologous and homeologous chromosomes

is essential to ensure correct meiotic segregation. Preferential pairing of homologs has been

observed in early stages of polyploid formation in *Arabidopsis suecica* (Pecinka *et al.*, 2011)

and *Brassica napus* (Grandont *et al.*, 2014), although elevated homeolog mispairings has

also been reported (Xiong *et al.*, 2021). Simultaneously, allopolyploids often undergo genetic

diploidization, in which some redundant or misregulated gene copies may be silenced or

lose function over time (Lee & Chen, 2001; Kashkush *et al.*, 2002; Koh *et al.*, 2010). Some

duplicated copies can be retained and remain functional, for example if they confer adaptive

benefits, become subfunctionalized, or acquire new functions via neofunctionalization

(Adams *et al.*, 2003; Blanc & Wolfe, 2004; Duarte *et al.*, 2006; De Smet *et al.*, 2013).

In addition to silencing TEs, smRNAs—particularly 21–22 nt species—regulate gene

expression via PTGS or TGS (Borges & Martienssen, 2015). We observed widespread genic

smRNA targeting, especially in exons and 1 kb upstream of gene models (putative

promoters), with 21 nt smRNAs contributing the most. While such targeting may mediate

transcript degradation, alternative splicing, or cis-silencing (Guleria *et al.*, 2011; Moss, 2001),

some associations may also have no functional consequence. Nevertheless, these patterns

can reflect regulatory adjustments during early diploidization.

Substantial genic smRNA targeting was also detected in the diploids, especially *D. fuchsii*,

likely reflecting stress responses in the common garden experiment conducted outdoors in

Vienna, Austria—an area where only *D. incarnata* occurs naturally. Environmental mismatch

may have triggered miRNA- and nat-siRNA-mediated regulation in *D. fuchsii*, consistent with

previous reports of stress-induced smRNA upregulation (Zhang *et al.*, 2012; Mao *et al.*,

2021; Tiwari & Rajam, 2022).

Transgressive smRNA expression relative to the diploid progenitors may reflect how

allopolyploids respond to genome duplication while also acquiring novel ecological features,

potentially facilitating establishment in niches outside the parental range. We observed more

transgressive activity at peaks targeted by 20–23 nt smRNAs—compared to 24 nt—

especially when the allotetraploids showed higher smRNA levels relative to both diploid

parental species (Fig. 4e–f).

Genes transgressively targeted by smRNAs were enriched for functions related to mitosis,

meiosis, gene regulation, stress responses, and transport (Fig. 5). While these functional

categories were largely shared across the two allotetraploids, the specific genes and

pathways involved differed, suggesting species-specific regulatory routes. In the younger *D.*

traunsteineri, enrichment of smRNA targeting was observed for genes involved in cell cycle

control, possibly reflecting ongoing meiotic stabilization. By contrast, the older *D. majalis*

showed enrichments for genes involved in transcriptional and translational regulation,

suggesting a later phase of polyploid evolution focused on dosage control and fine-tuning of

gene expression.

Together, these observations support a dynamic evolution of smRNA-mediated regulation

following allopolyploidization: an early phase dominated by stabilization of chromosome

behavior during meiosis, together with refinement of gene redundancy and the emergence of

functional divergence. Similar dynamics have been proposed in other systems including

Arabidopsis, *Brassica*, and *Gossypium* (Ha *et al.*, 2009; Fu *et al.*, 2016; Chen *et al.*, 2020).

Conclusions and Future Directions

This study underscores the dual associations of smRNAs with genomic features in

allopolyploids—both with TEs and genic regions—reflecting their potential roles in genome

regulation. Despite their shared ancestry, *D. majalis* and *D. traunsteineri* exhibit clear

lineage-specific patterns of smRNA activity, particularly in genic regions, suggesting

divergent regulatory trajectories following allopolyploidization. Given their rapid action,

diversity, and regulatory plasticity, smRNAs likely play central roles in both early and long-

term genomic reorganization of allopolyploids. Our results support a model in which

smRNAs mediate both immediate stabilization and longer-term functional divergence after

genome merging. Future work should integrate smRNA expression profiles with homeolog-

specific reference genomes and DNA methylation landscapes. These approaches would

provide deeper insights into subgenome-specific regulation, the dynamics of smRNA–target

interactions, and the evolutionary consequences of smRNA-mediated gene and TE

regulation in natural allopolyploids.

ACKNOWLEDGEMENTS

This work was funded by the Austrian Science Fund (FWF) [grant DOI: 10.55776/Y661 awarded to OP and through the doctoral programme DOI: 10.55776/W1225, awarded to a faculty team including OP]. For open access purposes, the author has applied a CC BY public copyright license to any author accepted manuscript version arising from this submission. Faculty members, in particular Andreas Futschik, Robert Kofler and Christian Schlötterer, as well as the students and the scientific advisory board of the Vienna Graduate School of Population Genetics (www.popgen-vienna.at), Richard Bateman, and Ortrun Mittelsten Scheid are gratefully acknowledged for their valuable discussions and feedback throughout this work. We thank Juliane Baar, Daniela Paun and Maria Teresa Lorenzo for their support with laboratory procedures. Sequencing was performed at the Vienna BioCenter CoreFacilities (VBCF; https://www.viennabiocenter.org/). Computational resources were provided by the Life Science Compute Cluster (LiSC) of the University of Vienna. We also acknowledge Natural England and the Forestry Commission (UK), and regional county administrations in Austria, France, and Sweden for issuing the necessary collecting permits.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

Study conceived and designed by OP. Sampling and transplantation to the common garden conducted by OP, FB, MH, MCE and TMW. Read processing and mapping done by MT. Identification of putative smRNA peaks across the genome performed by MT and MCE. Data analyses of differentially targeted genes and TEs conducted by MCE. Analysis of overlapping differentially expressed and differentially targeted genes done by TMW. Interpretation of the results was undertaken by MCE. The manuscript was drafted by MCE with feedback from OP, and was revised and approved by all authors.

DATA AVAILABILITY

Newly reported raw Illumina smRNA-seq data are deposited on NCBI SRA (GenBank BioProject PRJNA317244, accessions SRR17191877–SRR17191897). Previously published Illumina RNA-seq and smRNA-seq are available from the same GenBank BioProject PRJNA317244. The pipeline, scripts and raw count file, together with the peak gff3 file are available at https://github.com/mc-er/dact-sRNAs.

REFERENCES

Adams KL, Cronn R, Percifield R, Wendel JF. 2003. Genes duplicated by polyploidy show unequal contributions to the transcriptome and organ-specific reciprocal silencing. *Proceedings of the National Academy of Sciences USA* **100**: 4649–4654. doi:10.1073/pnas.0630618100

Alexa A, Rahnenfuhrer J. 2021. topGO: Enrichment Analysis for Gene Ontology. R package version 2.44.0. [WWW document] URL https://bioconductor.org/packages/release/bioc/html/topGO.html. [accessed 15 May 2025].

Ariel FD, Manavella PA. 2021. When junk DNA turns functional: transposon-derived non-coding RNAs in plants. *Journal of Experimental Botany* **72**: 4132–4143. doi:10.1093/jxb/erab082

Balao F, Trucchi E, Wolfe TM, Hao BH, Lorenzo MT, Baar J, Paun O. 2017. Adaptive sequence evolution is driven by biotic stress in a pair of orchid species (*Dactylorhiza*) with distinct ecological optima. *Molecular Ecology* **26**: 3649–3662. doi:10.1111/mec.14123

Blanc G, Wolfe KH. 2004. Functional divergence of duplicated genes formed by polyploidy during *Arabidopsis* evolution. *Plant Cell* **16**: 1679–1691. doi:10.1105/tpc.021410

Borges F, Martienssen RA. 2015. The expanding world of small RNAs in plants. *Nature Reviews Molecular Cell Biology* **16**: 727–737. doi:10.1038/nrm4085

Brandrud MK, Baar J, Lorenzo MT, Athanasiadis A, Bateman RM, Chase MW, Hedrén M, Paun O. 2020. Phylogenomic relationships of diploids and the origins of allotetraploids in *Dactylorhiza*. *Systematic Biology* **69**: 91–109. doi:10.1093/sysbio/syz035

Carthew RW, Sontheimer EJ. 2009. Origins and mechanisms of miRNAs and siRNAs. *Cell* **136**: 642–665. doi:10.1016/j.cell.2009.01.035

Cavé-Radet A, Giraud D, Lima O, El Amrani A, Aïnouche ML, Salmon A. 2020. Evolution of small RNA expression following hybridization and allopolyploidization: insights from *Spartina* species (Poaceae, Chloridoideae). *Plant Molecular Biology* **102**: 55–72. doi:10.1007/s11103-019-00931-w

Cavé-Radet A, Salmon A, Tran Van Canh L, Ainouche M. 2023. Recent allopolyploidy alters *Spartina* microRNA expression in response to xenobiotic-induced stress. *Plant Molecular Biology* **111**: 309–328. doi:10.1007/s11103-022-01328-y

Chen X, Rechavi O. 2021. Plant and animal small RNA communications between cells and
organisms. *Nature Reviews Molecular Cell Biology* **22**: 750–766. doi:10.1038/s41580-
021-00425-y

Chen ZJ, Sreedasyam A, Ando A, Song Q, De Santiago LM, Hulse-Kemp AM, Schmutz
J. 2020. Genomic diversifications of five *Gossypium* allopolyploid species and their
impact on cotton improvement. *Nature Genetics* **52**: 525–533. doi:10.1038/s41588-020-
0614-5

Cheng F, Sun C, Wu J, Schnable J, Woodhouse MR, Liang J, Wang X. 2016. Epigenetic
regulation of subgenome dominance following whole genome triplication in *Brassica*
rapa. New Phytologist **211**: 288–299. doi:10.1111/nph.13884

Creasey KM, Zhai J, Borges F, Van Ex F, Regulski M, Meyers BC, Martienssen RA.
2014. miRNAs trigger widespread epigenetically activated siRNAs from transposons in
Arabidopsis. Nature 508: 411–415. doi:10.1038/nature13069

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, Depristo MA, Handsaker RE,
Lunter G, Marth GT, Sherry ST et al. 2011. The variant call format and VCFtools.
Bioinformatics **27**: 2156–2158. doi:10.1093/bioinformatics/btr330.

De Smet R, Adams KL, Vandepoele K, Van Montagu MCE, Maere S, Van de Peer Y.
2013. Convergent gene loss following gene and genome duplications creates single-copy
families in flowering plants. *Proceedings of the National Academy of Sciences, USA* **110**:
2898–2903. doi:10.1073/pnas.1300127110.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–
21. doi:10.1093/bioinformatics/bts635.

Duarte JM, Cui L, Wall PK, Zhang Q, Zhang X, Leebens-Mack J, DePamphilis CW.
2006. Expression pattern shifts following duplication indicative of subfunctionalization
and neofunctionalization in regulatory genes of *Arabidopsis. Molecular Biology and*
Evolution **23**: 469–478. doi:10.1093/molbev/msj051

Edger PP, Smith R, McKain MR, Cooley AM, Vallejo-Marin M, Yuan Y, Puzey JR. 2017.
Subgenome dominance in an interspecific hybrid, synthetic allopolyploid, and a 140-year-
old naturally established neo-allopolyploid monkeyflower. *Plant Cell* **29**: 2150–2167.
doi:10.1105/tpc.17.00010

Emelianova K, Hawranek A-S, Eriksson M, Wolfe TM, Paun O. 2025. Ecological
divergence of sibling allopolyploid marsh orchids is associated with species specific

plasticity and distinct fungal communities. *Plant Journal* **121**: e70001. doi:

10.1111/tpj.70001

Eriksson MC, Mandáková T, McCann J, Temsch EM, Chase MW, Hedrén M, Weiss-

Schneeweiss H, Paun O. 2022. Repeat dynamics across timescales: a perspective from

sibling allotetraploid marsh orchids (*Dactylorhiza majalis* s.l.). *Molecular Biology and*

Evolution **39**: msac167. doi:10.1093/molbev/msac167

Fedoroff NV. 2012. Transposable elements, epigenetics, and genome evolution. *Science*

338: 758–77. doi:10.1126/science.338.6108.758

Fu Y, Xiao M, Yu H, Mason AS, Yin J, Li J, Fu D. 2016. Small RNA changes in synthetic

Brassica napus. Planta **244**: 607–622. doi:10.1007/s00425-016-2529-z

Fultz D, Choudury SG, Slotkin RK. 2015. Silencing of active transposable elements in

plants. *Current Opinion in Plant Biology* **27**: 67–76. doi:10.1016/j.pbi.2015.05.027

Gaebelein R, Schiessl SV, Samans B, Batley J, Mason AS. 2019. Inherited allelic variants

and novel karyotype changes influence fertility and genome stability in *Brassica*

allohexaploids. *New Phytologist* **223**: 965-978. doi:10.1111/nph.15804.

Ghani MA, Li J, Rao L, Raza MA, Cao L, Yu N, Chen L. 2014. The role of small RNAs in

wide hybridisation and allopolyploidisation between *Brassica rapa* and *Brassica nigra*.

BMC Plant Biology **14**: 272. doi:10.1186/s12870-014-0272-9

Grandont L, Cuñado N, Coriton O, Huteau V, Eber F, Chèvre AM, Jenczewski E. 2014.

Homoeologous chromosome sorting and progression of meiotic recombination in

Brassica napus: ploidy does matter! *Plant Cell* **26**: 1448–1463.

doi:10.1105/tpc.114.122788

Guleria P, Mahajan M, Bhardwaj J, Yadav S. 2011. Plant small RNAs: biogenesis, mode of

action and their roles in abiotic stresses. *Genomics, Proteomics & Bioinformatics* **9**: 183–

199. doi:10.1016/S1672-0229(11)60022-9

Ha M, Lu J, Tian L, Ramachandran V, Kasschau KD, Chapman EJ, Chen ZJ. 2009.

Small RNAs serve as a genetic buffer against genomic shock in *Arabidopsis* interspecific

hybrids and allopolyploids. *Proceedings of the National Academy of Sciences USA* **106**:

17835–17840. doi:10.1073/pnas.0907003106

Hassan AH, Mokhtar MM, El Allali A. 2024. Transposable elements: multifunctional players

in the plant genome. *Frontiers in Plant Science* **14**: 1330127.

doi:10.3389/fpls.2023.1330127

Hawranek AS. 2021. *The impact of recurrent origins and gene flow on the genetic structure*

*of allopolyploid marsh orchids (*Dactylorhiza*, Orchidaceae).* MSc thesis, University of

Vienna, Vienna, Austria. doi:10.25365/thesis.70360

Ito H, Gaubert H, Bucher E, Mirouze M, Vaillant I, Paszkowski J. 2011. An siRNA

pathway prevents transgenerational retrotransposition in plants subjected to stress.

Nature 472: 115–120.

Jiang X, Song Q, Ye W, Chen ZJ. 2021. Concerted genomic and epigenomic changes

accompany stabilization of *Arabidopsis* allopolyploids. *Nature Ecology & Evolution* **5**:

1382–1393. doi:10.1038/s41559-021-01523-y.

Jiao W, Yuan J, Jiang S, Liu Y, Wang L, Liu M, Wang J, Zhang L, Liu Y, Xia C et al.

2018. Asymmetrical changes of gene expression, small RNAs and chromatin in two

resynthesized wheat allotetraploids. *Plant Journal* **93**: 828–842. doi:10.1111/tpj.13805.

Kashkush K, Feldman M, Levy AA. 2002. Gene loss, silencing and activation in a newly

synthesized wheat allotetraploid. *Genetics* **160**: 1651–1659.

doi:10.1093/genetics/160.4.1651.

Kaur R, Choudhury A, Chauhan S, Ghosh A, Tiwari R, Rajam MV. 2021. RNA

interference and crop protection against biotic stresses. *Physiology and Molecular*

Biology of Plants **27**: 2357–2373. doi:10.1007/s12298-021-01058-2.

Kenan-Eichler M, Leshkowitz D, Tal L, Noor E, Melamed-Bessudo C, Feldman M, Levy

AA. 2011. Wheat hybridization and polyploidization results in deregulation of small

RNAs. *Genetics* **188**: 263–272. doi:10.1534/genetics.111.128348.

Koh J, Soltis PS, Soltis DE. 2010. Homeolog loss and expression changes in natural

populations of the recently and repeatedly formed allotetraploid *Tragopogon mirus*

(Asteraceae). *BMC Genomics* **11**: 97. doi:10.1186/1471-2164-11-97.

Lee HS, Chen ZJ. 2001. Protein-coding genes are epigenetically regulated in *Arabidopsis*

polyploids. *Proceedings of the National Academy of Sciences, USA* **98**: 6753–6758.

doi:10.1073/pnas.121064898.

Lee SC, Ernst E, Berube B, Borges F, Parent J-S, Ledon P, Zuber H, Sperschneider J,

Boccara M, Lalonde S et al. 2020. *Arabidopsis* retrotransposon virus-like particles and

their regulation by epigenetically activated small RNA. *Genome Research* **30**: 576–588.

doi:10.1101/gr.259044.119.

Lewsey MG, Hardcastle TJ, Melnyk CW, Molnar A, Valli A, Ulrich MA, Nery JR,
Baulcombe DC, Ecker JR. 2016. Mobile small RNAs regulate genome-wide DNA
methylation. *Proceedings of the National Academy of Sciences, USA* **113**: E801-E810.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**: 923–930.
doi:10.1093/bioinformatics/btt656.

Liu C, Wang YG. 2023. Does one subgenome become dominant in the formation and
evolution of a polyploid? *Annals of Botany* **131**: 11–16. doi:10.1093/aob/mcac024.

Mao Y, Xu J, Wang Q, Li G, Tang X, Liu T, Zhang H, Wang W, Li T, Chen X et al. 2021. A
natural antisense transcript acts as a negative regulator for the maize drought stress
response gene ZmNAC48. *Journal of Experimental Botany* **72**: 2790–2806.
doi:10.1093/jxb/erab023.

Marí-Ordóñez A, Marchais A, Etcheverry M, Martin A, Colot V, Voinnet O. 2013.
Reconstructing de novo silencing of an active plant retrotransposon. *Nature Genetics* **45**:
1029–1033. doi:10.1038/ng.2703.

Martinez Palacios P, Jacquemot M-P, Tapie M, Rousselet A, Diop M, Remoué C,
Demangeat G, Berges H, Pichon M, Marande W et al. 2019. Assessing the response
of small RNA populations to allopolyploidy using resynthesized *Brassica napus*
allotetraploids. *Molecular Biology and Evolution* **36**: 709–726.
doi:10.1093/molbev/msz007.

Matzke MA, Mosher RA. 2014. RNA-directed DNA methylation: an epigenetic pathway of
increasing complexity. *Nature Reviews Genetics* **15**: 394-408. doi:10.1038/nrg3683.

McCarthy DJ, Chen Y, Smyth GK. 2012. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research* **40**:
4288–4297. doi:10.1093/nar/gks042.

McClintock B. 1984. The significance of responses of the genome to challenge. *Science*
226: 792–801. doi:10.1126/science.15739260.

Moss E. 2001. RNA interference: it's a small RNA world. *Current Biology* **11**: R772–R775.
doi:10.1016/S0960-9822(01)00411-1.

Pecinka A, Fang W, Rehmsmeier M, Levy AA, Mittelsten Scheid O. 2011.
Polyploidization increases meiotic recombination frequency in *Arabidopsis. BMC Biology*
9: 24. doi:10.1186/1741-7007-10-33.

Poretti M, Praz CR, Meile L, Kälin C, Schaefer LK, Schläfli M, Widrig V, Sanchez-Vallet
A, Wicker T, Parlange F et al. 2020. Domestication of high-copy transposons underlies
the wheat small RNA response to an obligate pathogen. *Molecular Biology and Evolution*
37: 839–848. doi:10.1093/molbev/msz272.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841–842. doi:10.1093/bioinformatics/btq033.

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F,
Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data
analysis. Nucleic Acids Research 44: W160–W165. doi:10.1093/nar/gkw257.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26: 139–
140. doi:10.1093/bioinformatics/btp616.

Shen Y, Sun S, Hua S, Shen E, Ye CY, Cai D, Timko MP, Zhu QH, Fan L, Xu L et al.
2017. Analysis of transcriptional and epigenetic changes in hybrid vigor of allopolyploid
Brassica napus uncovers key roles for small RNAs. Plant Journal 91: 874–893.
doi:10.1111/tpj.13605.

Tiwari R, Rajam MV. 2022. RNA- and miRNA-interference to enhance abiotic stress
tolerance in plants. Journal of Plant Biochemistry and Biotechnology 31: 1–10.
doi:10.1007/s13562-022-00770-9.

Tossolini I, Mencia R, Arce AL, Manavella PA. 2025. The genome awakens: transposon-
mediated gene regulation. Trends in Plant Science. doi:10.1016/j.tplants.2025.02.005.

Wang HLV, Chekanova JA. 2016. Small RNAs: essential regulators of gene expression
and defenses against environmental stresses in plants. Wiley Interdisciplinary Reviews:
RNA 7: 356–381. doi:10.1002/wrna.1340.

Wang M, Zhao Y, Zhang B. 2015. Efficient test and visualization of multi-set intersections.
Scientific Reports **5**: 16923. doi:10.1038/srep16923.

Wolfe TM, Balao F, Trucchi E, Bachmann G, Gu W, Baar J, Suda J, Paun O. 2023.
Recurrent allopolyploidizations diversify ecophysiological traits in marsh orchids
(*Dactylorhiza majalis* s.l.). *Molecular Ecology* **32**: 4777–4790. doi:10.1111/mec.17070.

Xiao Y, Wang J. 2025. Understanding the regulation activities of transposons in driving the
variation and evolution of polyploid plant genome. *Plants* **14**: 1160.
doi:10.3390/plants14081160.

Xie Y, Ying S, Li Z, Zhang Y, Zhu J, Zhang J, Li X, Liu H, Wang Y, Liu Y et al. 2023.
Transposable element-initiated enhancer-like elements generate the subgenome-biased
spike specificity of polyploid wheat. *Nature Communications* **14**: 7465.
doi:10.1038/s41467-023-42771-9.

Xiong Z, Gaeta RT, Edger PP, Cao Y, Zhao K, Zhang S, Wan T, Liu Y, Coate JE, Pires
JC. 2021. Chromosome inheritance and meiotic stability in allopolyploid *Brassica napus.*
G3: Genes, Genomes, Genetics **11**: jkaa011. doi:10.1093/g3journal/jkaa011.

Liu L, Chen X. 2018. Intercellular and systemic trafficking of RNAs in plants. *Nature Plants*
4: 869–878. doi:10.1038/s41477-018-0288-5.

Zhan J, Meyers BC. 2024. Plant small RNAs: their biogenesis, regulatory roles, and
functions. *Annual Review of Plant Biology* **75**: 21–51. doi:10.1146/annurev-arplant-
070122-035226.

Zhang X, Xia J, Lii YE, Barrera-Figueroa BE, Zhou X, Gao S, Lu L, Niu D, Chen Z, Liang
W et al. 2012. Genome-wide analysis of plant nat-siRNAs reveals insights into their
distribution, biogenesis and function. *Genome Biology* **13**: R20. doi:10.1186/gb-2012-13-
3-r20.

 FIGURE LEGENDS

Figure 1. Study system comprising two diploids (*Dactylorhiza fuchsii* and *D. incarnata*) and
their sibling allotetraploids (*D majalis* and *D. traunsteineri*). **(a)** Overview of the four species.
Coloured circles and their relative positions illustrate population densities. Cell icons indicate
difference in genome size between the diploids, while gender symbols show the direction of
hybridization leading to the formation of the allopolyploids. Additional symbols denote broad
ecological preferences. **(b)** Multidimensional scaling (MDS) plot based on small RNA count
data. Colours correspond to the species frames shown in (a); triangles indicate diploid
accessions, and circles, allotetraploid accessions.

Figure 2. Small RNA read length distributions across species. **(a)** Mean counts per million
(CPM) for each read length; error bars represent the standard error of the mean. Horizontal
lines indicate significant differences between species, with asterisks denoting significance
levels: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. **(b, c)** Proportions of reads by species and read
length mapped to genomic features: (b) exons and (c) transposable elements. fuc*, D. fuchsii*;
inc, *D. incarnata*; maj, *D. majalis*; tra, *D. traunsteineri*.

Figure 3. MA plots showing differences in small RNA (smRNA) "targeting" between
Dactylorhiza majalis and *D. traunsteineri*. The central dashed line marks logFC = 0, while
dotted lines indicate logFC = -1.5 and 1.5. Negative logFC values represent higher smRNA
targeting in *D. majalis* relative to *D. traunsteineri*, whereas positive logFC values indicate
higher targeting in *D. traunsteineri*. **(a–c)** Peaks differently targeted in genic regions by 20–
24 nt smRNAs: **(a)** promoter/1,000 bp upstream gene models, **(b)** exons and **(c)** introns. **(d)**
Transposable elements (TEs) targeted by 20–23 nt smRNAs. **(e)** TEs targeted by 24 nt
smRNAs. **(f)** MA plot of RNA-seq data from Wolfe *et al.* (2023), showing genes differentially
expressed (DE) and/or DT by smRNAs as indicated in the legend.

Figure 4. Patterns of genomic interactions in smRNA "targeting". **(a–b)** Comparison of logFC
values for *Dactylorhiza majalis* relative to either diploid parent (*D. fuchsii* on the Y-axis, *D.*
incarnata on the X-axis) for all peaks (20–24 nt smRNAs) in annotated genes. **(a)** Colored
symbols indicate additive and transgressive patterns; **(b)** shows dominance toward either
diploid, as indicated in the legend. **(c-f)** Number of differentially targeted (DT) peaks in each
type of genomic interaction, with peaks in *D. traunsteineri* shown in blue and *D. majalis* in
red. Striped bars denote DT peaks shared between the allotetraploids. The annotation
columns along the Y-axis represent the interaction type (colored bars, according to legend),
while dots connected by lines show the direction of the interaction. Letters above each
column indicate I - *D. incarnata*, P - polyploid, and F - *D. fuchsii.* **(c)** 20–24 nt smRNAs

associated with genic regions, **(d)** 20–24 nt smRNA peaks outside annotated genic and TE

regions, **(e)** 20–23 nt smRNAs in TEs, and **(f)** 24 nt smRNAs in TEs.

Figure 5. Summary of representative functional enrichments for **(a)** the allopolyploid

comparison, **(b)** transgressive smRNA "targeting" of genic regions in *Dactylorhiza*

traunsteineri, and **(c)** transgressive targeting in *D. majalis*. Bar length represents the

adjusted significance of each enriched GO term. Full results are provided in Supplementary

Tables S6, S8 and S9. Bar color indicates the z-score, computed from the logFC values of

the corresponding targeting, as indicated in the legends. Arrows on bars in panels (b) and (c)

mark GO terms shared between the two comparisons. Symbols next to the GO terms denote

the broad functional categories (see legend), with numbers indicating the count of enriched

genes within each category. Letters to the right of each plot specify the genetic region

targeted by smRNAs; p - 'promotor', e - exon, i - intron. Abbreviations used in GO terms: +

positive, - negative, DS double strand, pol polymerase, ts transcription.

SUPPLEMENTARY TABLES LEGENDS

Table S1. Sample information. Collector abbreviations are FB, Francisco Balao; MC, Mark Chase; MF, Michael Fay; MH, Mikael Hedren; and OP, Ovidiu Paun. Note: the smRNA-seq data has been collected within a common garden setup; here only the population of origin of the plants is indicated.

Table S2. Criteria for categories of genomic interactions. Abbreviations in the column names are as follows: 4x, allotetraploid; fuc - *D. fuchsii*; inc - *D. incarnata*. Abbreviations in the column Direction are as follows: F, *D. fuchsii*; I, *D. incarnata*; P, polyploid (i.e. *D. majalis* or *D. traunsteineri*).

Table S3. Mapping statistics for each sample. Accession IDs ending with 'r' indicate technical replicates.

Table S4. Number of smRNA targeting peaks in differential targeting (DT) analysis. Subscripts in the column names are as follow: A, all analyzed DT peaks; all, significant DT peaks across the whole genome; TE, significant DT peaks in annotated transposable elements; G, significant DT peaks in annotated genes; Gp, significant DT peaks 1k upstreams of annotated genes; Gi, significant DT peaks in introns of annotated genes; Ge, significant DT peaks in exons of annotated genes; Multiple regions, DT peaks spanning the border between two different genic features (exon, intron or promotor). Significant = adjusted p-value <=0.05. Abbreviations in the table are as follows: 4X, polyploid; 2x, diploid; maj, *D.majalis*; tra, *D.traunsteineri*; fuc - *D. fuchsii*; inc - *D. incarnata*.

Table S5. List of genes both differentially expressed between the allotetraploids in Wolfe *et al.*, (2023) and differentially targeted by smRNAs (this study). smRNA, small RNA values from this study; RNA, RNA-seq values from Wolfe et al. (2023); GI, Genomic Interaction; maj, *D. majalis*; tra, *D. traunsteineri*.

Table S6. Enrichment results for genes differentially targeted between the polyploids (D. majalis and D. traunsteineri). N_peaks, number of peaks in GO term; N_genes, number of genes in GO term; adj_pval, adjusted p-value from topGO; Z-score, normalized value based on number of genes and the average log fold change value.

Table S7. Number of peaks for each defined genomic interaction (additive, transgressive or dominant towards either diploid). Regions mean: all - genome wide, genes - around annotated genes, TEs - around annotated TEs, other regions - outside annotated TEs and genes. Abbreviations in the column names are as follows: maj - number of peaks in *D.*

majalis, tra - number of peaks in *D. traunsteineri*, sig. Significance level for hypergeometric
test done for shared. * < 0.05, ** < 0.01, *** < 0.001.

Table S8. Enrichment results for genes transgressivly differentially targeted between the
polyploid (D. majalis) and either of the diploids (D. fuchsii and D. incarnata). N_peaks,
number of peaks in GO term; N_genes, number of genes in GO term; adj_pval, adjusted p-
value from topGO; Z-score, normalized value based on number of genes and the average
log fold change value.

Table S9. Enrichment results for genes transgressivly differentially targeted between the
polyploid (D. traunsteineri) and either of the diploids (D. fuchsii and D. incarnata). N_peaks,
number of peaks in GO term; N_genes, number of genes in GO term; adj_pval, adjusted p-
value from topGO; Z-score, normalized value based on number of genes and the average
log fold change value.

SUPPLEMENTARY FIGURES LEGENDS

Figure S1. Distribution of normalised reads counts for different smRNA lengths over
genomic regions (promoter/1000 bp upstream, exons, introns and intergenic) in the four
tested species. fuc = *D. fuchsii*, inc = *D. incarnata*, maj = *D. majalis*, tra = *D. traunsteineri*.

Figure S2. Number of differentially targeted peaks in genes and TEs for each comparison
between a polyploid (4x) and a diploid (2x) for all three data sets, 20-24 nt 20-23 nt and 24 nt
smRNAs. The top panel contains the number of peaks where the polyploid is over regulated
compared to the diploid, whereas the bottom panel contains the number of peaks where the
diploid is over targeted. The coloured boxed below the figure represents the comparison
shown in the corresponding column.

Figure S3. Number of DT peaks in all datasets 20–24 nt, 20–23 nt and 24 nt smRNAs for
each genomic interaction (additive, transgressing or dominant to either diploid). The top row
contains DT peaks in genes, the middle row shows DT peaks in TEs and the bottom row
shows peaks found outside annotated genic and TE regions, e.g. intergenic. Colours
represent, red - *D. majalis*, blue - *D. traunsteineri* and striped pattern - number of DT peaks
showing the same genomic interaction in both *D. majalis* and *D. traunsteineri*. The two
annotation columns along the Y–axis represent the type (coloured bars according to legend)
of genomic interaction Fig. 4, while dots connected with lines show the direction. Letters
above each column of dots represent **F** - *D. fuchsii*, **P** - polyploid and **I** - *D. incarnata*.

Figure S4. Patterns of genomic interactions comparing logFC values for *D. majalis* (left) and
D. traunsteineri (right) towards either diploid, *D. fuchsii* on the Y-axis and *D. incarnata* on the
X-axis for all peaks (i.e., 20–24 nt smRNAs) found in annotated genes. (a-b) shows with
coloured symbols transgressive and additive patterns; whereas (c-d) is dominant to either
diploid, according to the legend.

Figure S5 Proportions of peaks within each genomic interaction (additive, transgressing or
dominant to either diploid) for each genomic region (1000 bp upstream/promoter, exon,
intron). Colours are according to legend. Letters in legend represent **F** - *D. fuchsii*, **p** -
polyploid and **I** - *D. incarnata* while symbols represent level of smRNA between the diploids
and polyploid.

Figure S6 MA plots showing differences between *D. majalis* and *D. traunsteineri* for TEs
types grouped together, the central dashed line marks 0 while the dotted lines mark logFC -
1.5 and 1.5. Negative logFC values represent over-targeting in *D. majalis* as compared to *D.*
traunsteineri, whereas positive logFC values indicate over-targeting in *D. traunsteineri*. (a)
and (b) show 20–23 nt smRNAs while (c) and (d) show 24 nt smRNAs. (a) and (c) show DT
peaks specific for each smRNA group, i.e. 20–23 nt or 24 nt, while (b) and (d) show DT
peaks for both 20–23 nt and 24 nt smRNAs.

Figure S7 MA plots showing differences between *D. majalis* and *D. traunsteineri* for TEs, the
central dashed line marks 0 while the dotted lines mark logFC -1.5 and 1.5. Negative logFC
values represent over-targeting in *D. majalis* as compared to *D. traunsteineri*, whereas
positive logFC values indicate over-targeting in *D. traunsteineri*. Colours correspond to
different types of TEs, indicated by column headers. Rows correspond to smRNAs, where
the 1st and 2nd row show 20–23 nt smRNAs and 3rd and 4th show 24 nt smRNAs. The 2nd
and 3rd row show peaks that are DT for both 20–23 nt and 24 nt smRNAs.

Figure S8 Count of DT peaks between *D. majalis* and *D. traunsteineri*, the coloured points of
the plots in the 1st row (i.e. 20–23 nt smRNAs) in Fig. S7. Here Ty3-gypsy has been divided
into the next annotation level.

Figure S9 Patterns of genomic interactions comparing logFC values for *D. majalis* (left) and
D. traunsteineri (right) towards either diploid, *D. fuchsii* on the Y-axis and *D. incarnata* on the
X-axis for 20-23 nt smRNAs peaks found in annotated TEs. Top row shows with coloured
symbols transgressive and additive patterns; whereas bottom row shows dominant to either
diploid.

Figure S10 Patterns of genomic interactions comparing logFC values for *D. majalis* (left)

and *D. traunsteineri* (right) towards either diploid, *D. fuchsii* on the Y-axis and *D. incarnata*

on the X-axis for 24 nt smRNAs peaks found in annotated TEs. Top row shows with coloured

symbols transgressive and additive patterns; whereas bottom row shows dominant to either

diploid.