Research Article

Downregulation of O-GlcNAcylation enhances etoposide-induced p53-mediated apoptosis in HepG2 human liver cancer cells

Jaehoon Lee, Gi-Bang Koo, Jihye Park, Byung-Cheol Han, Mijin Kwon, and Seung-Ho Lee[*]

R&D Headquarter Korea Ginseng Corporation, 65, Gwacheon-daero 7-gil, Gwacheon-si, 13840, Korea

* Correspondence

Seung-Ho Lee, R&D Headquarter Korea Ginseng Corporation, 65, Gwacheon-daero 7-gil, Gwacheon-si, 13840, Korea. Email: fptase@kgc.co.kr

Keywords: Apoptosis, etoposide, O-GlcNAc transferase inhibitor, p53, anticancer, chemotherapy

Abbreviations:

ANOVA	analysis of variance
ATCC	American Type Culture Collection)
BSA	bovine serum albumin
Cas-3/7	caspase-3/7
EMEM	Eagle's Minimum Essential Medium
OGA	O-GlcNAcase
O-GlcNAc	O-linked β-N-acetylglucosamine
O-GlcNAcylation	O-linked β-N-acetylglucosamine modification
OGT	O-GlcNAc transferase
PBS	phosphate-buffered saline
PBST	PBS containing 0.1% Tween 20
P/S	penicillin/streptomycin
SD	standard deviation
TBST	0.05% Tween 20 Tris-buffered saline
Topo II	topoisomerase II
UDP	uridine diphosphate

Running heading: Synergistic anticancer effects of etoposide and OSMI-1

Abstract

Etoposide, an anticancer drug that inhibits topoisomerase II, is commonly used in combination chemotherapy. However, the impact of O-GlcNAcylation regulation on etoposide's anticancer effects has rarely been investigated. This study evaluated the effect of etoposide on cellular O-GlcNAcylation and whether modulating this process enhances etoposide-induced apoptosis. O-GlcNAc expression was measured after 24 h of etoposide treatment, and the effect of O-GlcNAc transferase (OGT) inhibition by OSMI-1 on etoposide's anticancer activity in HepG2 human liver cancer cells was quantitatively analyzed. Additionally, molecular analyses were used to confirm that the observed effects were mediated by p53-induced apoptosis. Etoposide reduced O-GlcNAcylation in a dose-dependent manner without directly interacting with OGT. Cotreatment with 20 μM of OSMI-1 lowered the IC_{50} value for cell viability by approximately 1.64-fold to 60.68 μM and increased the EC_{50} value for cytotoxicity by around 4.07-fold to 43.41 μM. Furthermore, this synergistic effect was linked to the activation of the p53/caspase-3/PARP1 pathway. These findings suggest that downregulating O-GlcNAcylation may enhance the efficacy of etoposide-based chemotherapy and help overcome tumor resistance.

Introduction

O-linked β-N-acetylglucosamine modification (O-GlcNAcylation) is a well-known post-translational modification that occurs in response to endogenous or exogenous factors, primarily in the nucleus, cytoplasm, and mitochondria [1]. O-GlcNAcylation plays a critical role in regulating transcription, metabolism, cell signaling, and physiological activity [2-4]. Prior studies have linked precise O-GlcNAcylation regulation to immune [5, 6] and cardiac functions [7], metabolic diseases [8], autoimmune disorders [9], and neurodegenerative diseases, such as Parkinson's and Alzheimer's disease [10]. Interestingly, O-GlcNAcylation is regulated by two highly conserved enzyme pairs: O-GlcNAc transferase (OGT) and protein O-GlcNAcase (OGA) [11]. Therefore, OGT and OGA are targets for disease treatment studies.

OGT function and O-GlcNAcylation regulation are involved in tumor formation, survival, metastasis, and angiogenesis. OGT activity and O-GlcNAcylation are increased in most malignant tumors and correlate positively with tumor progression and recurrence [12]. For example, studies have shown that OGT inhibition reduces cell cycle progression in a c-MYC–dependent manner in prostate cancer cells [13] and decreases epithelial–mesenchymal transition, metastasis, and invasion *in vivo* [14, 15]. Therefore, modulating O-GlcNAcylation by inhibiting OGT activity may be a promising target for cancer treatment.

Etoposide, an anticancer drug, forms a ternary complex with topoisomerase II (Topo II) and DNA, inducing apoptosis by inhibiting DNA strand religation [16]. This mechanism makes etoposide effective against various cancers, particularly those with rapid cell division. p53, a key transcription factor in apoptosis, is mutated in over 50% of human tumors [17]. When p53 is dysfunctional, cell cycle regulation and apoptosis are impaired, leading to uncontrolled cell proliferation and tumor development [18]. Thus, p53 is a highly effective target for cancer therapy. Chemotherapeutic drugs, such as etoposide and cisplatin, increase p53 expression, which in turn regulates the expression of *CDKN1A*, *GADD45A*, *BAX*, and other genes to induce cancer cell death [19-21]. However, additional resistance or the tumor microenvironment can allow cancer cells to evade the p53-mediated apoptosis pathway. For example, in HepG2 liver cancer cells, hypoxia conditions silence etoposide-induced p53 expression [22-24], suggesting that etoposide alone may have limited effectiveness *in vivo*.

Previous studies have shown that OGT inhibition with OSMI-1, an OGT inhibitor, increases sensitivity to the anticancer drug doxorubicin in HepG2 human liver cancer cells and PC-3 prostate cancer cells [25, 26]. Additionally, downregulating OGT activity using siOGT has been found to enhance the apoptotic response to bortezomib in MCF-7 and T47D human breast cancer cells [27]. However, the effect of O-GlcNAcylation regulation on etoposide's anticancer activity remains unexplored.

In this study, we investigated how O-linked β-N-acetylglucosamine (O-GlcNAc) expression is regulated by etoposide treatment and demonstrated that further downregulation of O-GlcNAcylation enhances the anticancer effect of etoposide in multiple human cancer cells. We also confirmed that these effects occur through the p53-mediated apoptosis pathway. Our findings suggest that reducing O-GlcNAcylation could help overcome tumor resistance and improve therapeutic outcomes in etoposide-based chemotherapy.

Methods

General materials

Etoposide and Thiamet G were obtained from MedChemExpress (Monmouth Junction, NJ, USA). OSMI-1 and phalloidin-iFluor™ 555 conjugate were purchased from GlpBio (Montclair, CA, USA) and Cayman Chemical (Ann Arbor, MI, USA), respectively. Crystal violet solution, Tween 20, and bisBenzimide H 33342 trihydrochloride (Hoechst 33342) were acquired from Sigma-Aldrich (Saint Louis, MO, USA). Antibodies were sourced from the following companies: O-GlcNAc (monoclonal; #82332) for western blot analysis from Cell Signaling Technology (Danvers, MA, USA); O-GlcNAc (RL2; monoclonal; #MA1-072) antibodies for immunocytochemistry from Thermo Fisher Scientific (Waltham, MA, USA); OGT (ARC0790; monoclonal; #A3501), OGA (ARC65434; monoclonal; #A24124), PARP1 (ARC0075; monoclonal; #A19596), and cleaved-PARP1 (ARC0091; monoclonal; #A19612) antibodies from ABclonal Technology (Woburn, MA, USA); and β-actin (polyclonal; #ab8227) and p53 (PAb421; monoclonal; #NBP2-62555) antibodies from Abcam (Cambridge, United Kingdom) and Novus Biologicals (Centennial, CO, USA), respectively.

Cell culture

HepG2 and Hep3B human liver hepatocellular carcinoma, A549 human lung carcinoma, and Caco-2 human colorectal adenocarcinoma cell lines were obtained from the American Type Culture Collection (ATCC; Manassas, VA, USA). HepG2, Hep3B, Caco-2 cells were cultured in Eagle's Minimum Essential Medium (EMEM; ATCC, Chicago, IL, USA) containing 1.0 g/L D-glucose, 10% fetal bovine serum (FBS), 1% penicillin/streptomycin (P/S), 1.5 g/L sodium bicarbonate, 292.0 mg/L L-glutamine, and 110.0 mg/L sodium pyruvate. A549 cells were cultured in Kaighn's Modification of Ham's F-12 Medium (ATCC) containing 1.26 g/L D-glucose, 10% FBS, 1% P/S, 1.5 g/L sodium bicarbonate, 292.2 mg/L L-glutamine, and 220.0 mg/L sodium pyruvate. All cells were cultured at 37°C in an incubator with a 5% CO_2 humidified atmosphere.

Cell viability assay

HepG2 cells were seeded in 96-well plates at 1.0×10^4 cells per well and maintained for 24 h. The medium was replaced with different concentrations of drugs, and after 24 h, cell viability was assessed using the D-Plus™ CCK Cell Viability Assay Kit (Dongin LS, Seoul, South Korea).

Crystal violet staining

To evaluate the effects of etoposide and OSMI-1 on HepG2 growth, cells were seeded in EMEM in 48-well plates at 5.0×10^4 cells per well. After 24 h, the medium was replaced with the drugs. Following 24 h of treatment, the cells were stained with crystal violet for 20 min and then washed thoroughly with phosphate-buffered saline (PBS). ImageJ software (version 1.54g) was used to quantify crystal violet intensity.

Western blot analysis

HepG2 cells were harvested and lysed on ice for 5 min using Cell Culture Lysis 1X Reagent (Promega, Madison, WI, USA) containing a mixture of protease and phosphatase inhibitors (MedChemExpress, Monmouth Junction, NJ, USA). The insoluble debris was removed via centrifugation at $15,000 \times g$ for 20 min at

4°C, and the total protein concentration of the isolated supernatants was quantified using the Pierce^{TM} BCA

Protein Assay Kit (Thermo Fisher Scientific). Proteins (10–20 μg) were separated using the Bolt™ Bis-Tris

Plus Gel system (Thermo Fisher Scientific) and transferred to polyvinylidene difluoride membranes (Bio-Rad,

Hercules, CA, USA). The membranes were incubated in 0.05% Tween 20 Tris-buffered saline (TBST)

containing 3%–5% bovine serum albumin (BSA) for 1 h at room temperature, followed by incubation overnight

at 4°C in 5% BSA–TBST with diluted primary antibodies. The membranes were then washed three times with

TBST and incubated with horseradish peroxidase–conjugated secondary antibodies in 5% BSA–TBST for 1 h at

room temperature. Protein signals were developed on the membranes using ECL substrates (Thermo Fisher

Scientific) and analyzed using Fusion FX6.0 (Vilber, Collégien, France).

Immunocytochemistry

For immunocytochemistry, HepG2 cells were cultured with drugs at an appropriate density on gelatin-

coated coverslips (SPL Life Sciences, Pocheon-si, South Korea). Cells were fixed with 4% paraformaldehyde

for 20 min and permeabilized with 0.2% Triton X-100 for 5 min. After blocking with 10% normal donkey

serum (Abcam) for 1 h, the coverslips were incubated overnight with primary antibodies diluted in 10%

normal donkey serum. Subsequently, the coverslips were washed three times with PBS containing 0.1% Tween

20 (PBST) and incubated with Alexa Fluor 488–conjugated secondary antibodies for 30 min. After washing

with PBST, the coverslips were incubated with phalloidin-iFluor™ 555 conjugate and Hoechst 33342 for 10

min, followed by thorough washing. Finally, the coverslips were mounted on slides using ProLong™ Glass

Antifade Mountant (Thermo Fisher Scientific), and the slides were analyzed using LSM 900 with Airyscan 2

confocal microscopes (Carl Zeiss, Oberkochen, Germany).

OGT activity assay

OGT activity was assessed using the Uridine diphosphate (UDP)-Glo™ Glycosyltransferase Assay Kit

(Promega) according to the manufacturer's protocol. UDP-GlcNAc (Promega) was utilized as the UDP–sugar

substrate, while the OGT peptide substrate (KKKYPGGSTPVSSANMM) was sourced from AnaSpec (Fremont,

CA, USA). Recombinant human OGT protein was obtained from R&D Systems (Minneapolis, MN, USA). The

enzymatic reactions were performed in 1X OGT reaction buffer, consisting of 25 mM Tris (pH 7.5), 12.5 mM

MgCl, 0.06 mg/mL BSA, and 1 mM dithiothreitol.

Fluorescence-activated cell sorting

HepG2 cells were seeded at 1.0×10^6 cells per well in 6-well plates in the medium described above. After

24 h, the medium was replaced with fresh medium containing the desired concentrations of etoposide and

OSMI-1, after which it was maintained for 24 h. Cells were then harvested and stained using the Annexin V

Apoptosis Detection Kit with 7-AAD (STEMCELL Technologies, Vancouver, BC, Canada), following the

manufacturer's instructions, and analyzed using the FACSLyric™ System (BD Biosciences, Franklin Lakes, NJ,

USA).

Annexin V binding and membrane integrity assay

To assess the effect of drugs on apoptosis and necrosis in HepG2 cells, the RealTime-Glo™ Annexin V

Apoptosis and Necrosis Assay Kit (Promega) was used. Briefly, HepG2 cells were seeded in solid white 96-well

plates at 1.0×10^4 cells per well and maintained for 24 h. The medium was replaced with fresh medium

containing the drug and kit components according to the manufacturer's instructions. Luminescence and

fluorescence signals were measured using the GloMax® Discover Microplate Reader (Promega).

TUNEL assay

HepG2 cells cultured in 6-well plates were fixed with 4% formaldehyde for 30 min and permeabilized with

0.2% Triton X-100 and 0.5% BSA in PBS. The cells were then incubated with TUNEL reaction mixture

(Abbkine, Atlanta, GA, USA) for 2 h at 37°C, followed by phalloidin-iFluor™ 555 conjugate and 4',6-

diamidino-2-phenylindole (i.e., DAPI) for an additional 10 min at room temperature. After thorough washing

with PBS, the cells were analyzed using LSM 900 with Airyscan 2 confocal microscopes.

Viability (GF-AFC), cytotoxicity (bis-AAF-R110), and caspase-3/7 activity assay

To evaluate the effects of drugs on viability, cytotoxicity, and caspase-3/7 activity, HepG2 cells were

assessed using the ApoTox-Glo™ Triplex Assay Kit (Promega). Briefly, HepG2 cells were seeded in solid

white 96-well plates at 1.0×10^4 cells per well and maintained for 24 h. The medium was replaced with different

drug concentrations, and after 24 h, GF-AFC substrate and bis-AAF-R110 substrate were added according to the

manufacturer's protocol. After 2 h, fluorescence analysis (viability: 400/505 nm excitation/emission;

cytotoxicity: 485/520 nm excitation/emission) was conducted. A luminogenic caspase-3/7 substrate was then

added, and the plates were incubated for an additional 2 h at 37°C. Luminescence and fluorescence signals were

measured using the GloMax® Discover Microplate Reader (Promega).

Statistical analysis

Statistical analysis was performed using SPSS Statistics version 20.0 (IBM Corp., Armonk, NY, USA) and

GraphPad Prism 9 (GraphPad Software Inc., La Jolla, CA, USA). Data are presented as means ± standard

deviations (SDs). Analysis of variance (ANOVA) with the post-hoc Tukey HSD test was used for data analysis.

$P < 0.05$ was considered statistically significant.

Results

OGT inhibitor enhances sensitivity to etoposide in HepG2 cells

The effect of etoposide on the viability of HepG2 cells in the presence of the OGT inhibitor OSMI-1 was assessed using the CCK cell viability assay 24 h after treatment. Results showed that coadministration of etoposide with 20 μM OSMI-1 significantly reduced cell viability at all concentrations compared with etoposide alone (Fig. 1A). These findings were further validated through crystal violet staining (Fig. 1B and C). Viability analysis using GF-AFC showed that cotreatment with 20 μM OSMI-1 reduced the IC_{50} value of etoposide in HepG2 cells by approximately 1.64-fold, from 99.59 to 60.68 μM (Fig. 1D). Additionally, the membrane integrity assay, which serves as a cytotoxicity index, revealed that the EC_{50} value of etoposide decreased approximately 4.07-fold from 43.41 to 176.63 μM, following cotreatment with 20 μM OSMI-1 (Fig. 1E). The same trend was observed in other human cancer cell lines, including Hep3B, A549, and Caco-2, where cotreatment with etoposide and OSMI-1 resulted in a more significant reduction in cell viability (Fig. 1F). These results suggest that the anticancer effects of etoposide are influenced by O-GlcNAc regulation.

Etoposide decreases O-GlcNAcylation without directly interacting with OGT

To investigate the effect of etoposide on O-GlcNAc regulation, HepG2 cells were treated with etoposide for 24 h, and global O-GlcNAc expression was analyzed via western blot and immunocytochemistry analyses. Results revealed that etoposide downregulated O-GlcNAc expression in a concentration-dependent manner (Fig. 2A and B). When etoposide and OSMI-1 were administered either alone or in combination for 24 h, both treatments reduced O-GlcNAc expression, but cotreatment only reduced it to the same extent as OSMI-1 alone (Fig. 2C). To determine whether these O-GlcNAc modulatory effects were due to direct interaction with the OGT enzyme, an OGT inhibitor assay was conducted. The findings showed that etoposide did not significantly reduce OGT activity at concentrations below approximately 200 μM, whereas OSMI-1 had an IC_{50} value of 4.47 μM (Fig. 2D and E).

Anticancer effect of etoposide occurs through downregulation O-GlcNAc expression

To assess whether the anticancer effect of etoposide is regulated by global changes in O-GlcNAc expression, we evaluated HepG2 cell viability using the OGT inhibitor OSMI-1 and the OGA inhibitor Thiamet G. Results indicated that OSMI-1 significantly enhanced the anticancer effect of etoposide in a dose-dependent manner, whereas Thiamet G did not induce significant changes (Fig. 3A). These findings were confirmed through crystal violet staining (Fig. 3B). Although the etoposide-induced decrease in crystal violet intensity was partially reversed by Thiamet G treatment, this increase did not achieve statistical significance (Fig. 3C). Further investigation of O-GlcNAc expression via western blot and immunocytochemistry analyses after 24 h of treatment revealed that O-GlcNAc expression was more prominently reduced when cells were treated with OSMI-1 compared to etoposide alone, and Thiamet G restored the etoposide-induced reduction in O-GlcNAc expression (Fig. 3D and E). Additionally, etoposide alone or in combination with OSMI-1 reduced OGA expression in HepG2 cells, and Thiamet G restored the OGA expression lowered by etoposide. However, OGT expression remained unaffected. These results suggest that the potentiation of etoposide's anticancer effects by OSMI-1 occurs through O-GlcNAc downregulation.

OGT inhibitor significantly increases etoposide-induced apoptosis

To determine if the enhancement of etoposide's anticancer effect by OSMI-1 mediates the apoptosis pathway, fluorescence-activated cell sorting (FACS) analysis was performed. Annexin V was used to detect apoptotic cells, and the results revealed that cotreatment with 10 or 20 μM OSMI-1 and 20 μM etoposide for 24 h increased the proportion of cells in early apoptosis by 1.42- and 1.62-fold, respectively, compared with etoposide alone (Fig. 4A). However, 7-AAD staining did not significantly increase following 24 h of treatment. A luminescence assay was conducted to measure Annexin V binding to phosphatidylserine, yielding an EC_{50} value of 27.52 μM for the etoposide + 20 μM OSMI-1 cotreatment, which was 2.25-fold lower than the 61.94 μM value for etoposide treatment alone (Fig. 4B). Notably, etoposide and OSMI-1 cotreatment induced significant phosphatidylserine translocation from 4 h onward, with significant apoptosis occurring from 12 h onward, compared with etoposide treatment alone (Fig. 4C). Furthermore, TUNEL assay results confirmed that OSMI-1 treatment increased etoposide-induced apoptosis (Fig. 5A and B).

Synergistic anticancer effect of the OGT inhibitor is achieved through enhancement of the p53-mediated apoptotic pathway

To elucidate the mechanism underlying the synergistic anticancer effects of etoposide and the OGT inhibitor OSMI-1 p53 and cleaved-PARP1 expression levels were evaluated via western blot analysis following 24 h of treatment. Results showed that cotreatment with etoposide and 20 μM OSMI-1 significantly increased the expression of p53 and cleaved-PARP1 compared to etoposide treatment alone (Fig. 6A). Additionally, a caspase-3/7 activity assay, quantified via luminescence, revealed that etoposide alone enhanced caspase-3/7 activity in a dose-dependent manner, this activity was significantly increased further under the etoposide + 20 μM OSMI-1 cotreatment (Fig. 6B). These findings suggest that the enhanced downregulation of O-GlcNAcylation achieved by OSMI-1 potentiates activation of the p53-mediated apoptotic pathway, thereby augmenting the anticancer effects of etoposide.

Discussion

Etoposide, a plant-derived alkaloid, inhibits cell proliferation across various adenocarcinoma cell types by arresting cell division. Mammalian cells express two Topo II isoenzymes, namely Topo IIα and β, with Topo IIα often overexpressed in tumors, serving as a cell proliferation marker [28]. The primary cytotoxicity mechanism of etoposide involves targeting Topo IIα, which underpins its selective action on cancer cells [29]. However, tumors resistant to Topo II inhibitors and cell lines with decreased Topo II expression exhibit reduced sensitivity to etoposide, suggesting a direct correlation between resistance and reduced Topo IIα expression [30]. Consequently, combining Topo II inhibitors, including etoposide, with agents acting via alternative mechanisms of action has become a common strategy to counteract resistance.

Our findings demonstrate that OSMI-1 significantly enhanced the sensitivity of HepG2 cells to etoposide (Fig. 1D and E). Notably, the EC_{50} value for membrane integrity, a marker of cytotoxicity, decreased by more than 4-fold with OSMI-1 cotreatment. This result was validated by a marked reduction in cell viability under 10 μM etoposide treatment combined with OSMI-1 compared to etoposide alone (Fig. 1A). Although additional research is required to clarify the precise effect of reduced O-GlcNAcylation on etoposide's interaction with Topo II or subsequent molecular cascades, our study suggests that inhibiting O-GlcNAcylation may serve as an adjuvant strategy to enhance etoposide's anticancer efficacy. Previous studies have shown that reducing excessive O-GlcNAcylation can enhance the effectiveness of anticancer drugs [25, 26]. For example, the O-GlcNAcylation of the Topo IIα Ser1469 residue in breast cancer cells has been implicated in increased drug resistance to the chemotherapeutic drug adriamycin, and its inhibition has been shown to enhance therapeutic outcomes *in vitro* and *in vivo* [31]. This is consistent with findings that cancer cell lines and human tumors frequently exhibit elevated O-GlcNAcylation [32, 33]. In line with previous literature, our findings demonstrated that elevating O-GlcNAcylation via Thiamet G tended to decline the anticancer effect of etoposide (Fig. 3A and C). These observations suggest a potential causal relationship between heightened O-GlcNAcylation and tumorigenesis, although an analysis of 33 cancer types in the Cancer Genome Atlas database showed that mutations in O-GlcNAcylation cycling enzymes are rare compared with highly mutated genes, such as *p53* and *PIK3CA* [34], indicating that increased O-GlcNAcylation may be a phenotypic consequence of cancer progression.

Limited evidence exists on the effects of Topo II inhibitors on O-GlcNAcylation regulation. Doxorubicin, another Topo II inhibitor, has been reported to reduce O-GlcNAcylation in Cos-7 cells [35], but such effects have not been studied extensively for etoposide. Cotreatment with doxorubicin and OSMI-1 was previously shown to induce excessive apoptosis via the p53/PARP1 pathway in HepG2 cells [25], similar to our findings with etoposide in this study. However, studies using *Xenopus* egg extracts revealed that etoposide and doxorubicin exhibit differing Topo II inhibition mechanisms: etoposide traps Topo II behind replication forks in a Topo II–dependent manner, whereas doxorubicin stops the replication forks by inserting directly into the parental DNA in a Topo II–independent manner [36]. Therefore, the present study provides new insights into the interplay between Topo II inhibitors and O-GlcNAcylation regulation.

Our results confirmed that etoposide treatment reduces O-GlcNAcylation in a dose-dependent manner without directly interacting with OGT (Fig. 2A, C, and E). Although high etoposide concentrations (400 μM) appeared to reduce OGT activity by approximately 47.3%, this inhibitory effect was likely exaggerated owing to

etoposide solubility issues (Fig. 2E). These findings suggest that the apoptotic cascade initiated by etoposide's
inhibition of Topo II is associated with O-GlcNAcylation downregulation. This hypothesis is supported by the
dose-dependent increase in apoptosis observed with OSMI-1 cotreatment (Fig. 4A and 5B). Interestingly,
although etoposide treatment alone did not significantly alter OGT expression in HepG2 cells, it reduced OGA
levels, an effect further amplified by OSMI-1 cotreatment (Fig. 3D). This may reflect a compensatory
mechanism aimed at restoring reduced O-GlcNAcylation levels, as reported in a prior study [37].

Under cellular stress, whether arising internally or from external factors, p53 expression and stability are
elevated, triggering the activation of genes responsible for cell cycle arrest, DNA repair, apoptosis, and
autophagy [38]. Following DNA damage, the cell's fate depends on its severity, with damage either repaired or
cell death (e.g., apoptosis and necrosis) being initiated. In cases of minimal damage, the cell cycle is arrested,
enabling the activation of DNA repair mechanisms. PARP1 plays a crucial role in genome integrity by
recognizing DNA damage and facilitating the repair pathway [39]. However, when DNA damage becomes
irreparable, PARP1 is cleaved by caspase-3/7, a process regulated by the p53 pathway. This cleavage inactivates
PARP1, leading the cell toward apoptosis. [40]. Our findings demonstrated that cotreatment with OSMI-1
enhanced activation of the p53/PARP1 pathway to a greater extent than etoposide treatment alone (Fig. 6A).
This molecular insight aligns with the observed leftward shift in the phosphatidylserine–Annexin V binding
curve under etoposide + 20 μM OSMI-1 cotreatment conditions (Fig. 4B). Furthermore, the synergistic effect of
OSMI-1 was amplified over time compared with etoposide alone (Fig. 5C).

Etoposide has been extensively used in drug screening studies owing to its diverse cellular effects, which
include induction of autophagy, cell senescence, and apoptosis [41, 42]. These properties highlight the need for
further studies on the combination effects and mechanisms of etoposide and OSMI-1 across broader indications.
As OSMI-1 functions as an OGT inhibitor, downregulating global O-GlcNAcylation, future studies should focus
on identifying the specific O-GlcNAcylated residues in apoptosis-related proteins to clarify the molecular
mechanisms underlying this synergy. Although this study primarily used HepG2 human liver hepatocellular
carcinoma, the fundamental mechanism of etoposide's action is consistent across various cell types. This
suggests that similar effects can be anticipated in other cancers. Consistent with this hypothesis, we confirmed
the enhanced anticancer effect of the etoposide–OSMI-1 combination in Hep3B human liver hepatocellular
carcinoma, A549 human lung carcinoma, and Caco-2 human colorectal adenocarcinoma cell lines (Fig. 1F). It
should be confirmed whether these effects can be maintained *in vivo*, since our study did not verify its efficacy
in an animal model. Etoposide is a chemotherapeutic option used primarily for the treatment of lung or testicular
cancer; therefore, *in vivo* study using these other tumors could be considered based on existing studies.
Additionally, animal studies should be conducted to adjust the dosing ratio of etoposide and OSMI-1 to
maximize efficacy. Given that OGT is an important enzyme in systemic homeostasis, cotreatment of OSMI-1
may result in adverse effects. Therefore, future *in vivo* studies should be designed to simultaneously verify
safety and anticancer effects.

In conclusion, downregulation of global O-GlcNAcylation is integral to the apoptotic induction process
initiated by etoposide. Additional downregulation of O-GlcNAcylation via OGT inhibition significantly
amplifies etoposide's anticancer effects through activation of the p53-mediated pathway (Fig. 7). Although
further *in vivo* studies and molecular analyses are necessary to validate and elucidate this synergistic effect and
its mechanism, our results suggest that targeting O-GlcNAcylation presents a promising approach for

combination chemotherapy in cancer treatment.

Data accessibility

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Author contributions

JL and GK designed the experiments under the supervision of BH and SL. JL wrote the manuscript. JL, JP, and MK performed the experiments and analyzed the data. JL, GK, and BH performed literature research. JL designed the research template. All authors have read and agreed to the published version of the manuscript.

Acknowledgements

This study was funded by the R&D Headquarter of Korea Ginseng Corporation

Conflicts of interest

The authors are employees of Korea Ginseng Corporation, which funded this research. However, the company had no influence on the study design, data interpretation, or publication decision.

References

1. Kudlow, J. E. (2006) Post-translational modification by O-GlcNAc: another way to change protein
function, *J Cell Biochem.* **98**, 1062-75. doi:10.1002/jcb.20926

2. Hart, G. W. (2019) Nutrient regulation of signaling and transcription, *J Biol Chem.* **294**, 2211-2231.
doi:10.1074/jbc.AW119.003226

3. Yang, X. & Qian, K. (2017) Protein O-GlcNAcylation: emerging mechanisms and functions, *Nat*
Rev Mol Cell Biol. **18**, 452-465. doi:10.1038/nrm.2017.22

4. Wells, L., Whelan, S. A. & Hart, G. W. (2003) O-GlcNAc: a regulatory post-translational
modification, *Biochem Biophys Res Commun.* **302**, 435-41. doi:10.1016/s0006-291x(03)00175-x

5. Dong, H., Liu, Z. & Wen, H. (2022) Protein O-GlcNAcylation regulates innate immune cell function,
Front Immunol. **13**, 805018. doi:10.3389/fimmu.2022.805018

6. Lee, K. S., Lee, J., Lee, P., Jeon, B. C., Song, M. Y., Kwak, S., Lee, J., Kim, J. S., Kim, D. J., Kim,
J. H., Tesh, V. L., Lee, M. S. & Park, S. K. (2022) Inhibition of O-GlcNAcylation protects from Shiga
toxin-mediated cell injury and lethality in host, *EMBO Mol Med.* **14**, e14678.
doi:10.15252/emmm.202114678

7. Wang, H. F., Wang, Y. X., Zhou, Y. P., Wei, Y. P., Yan, Y., Zhang, Z. J. & Jing, Z. C. (2023)
Protein O-GlcNAcylation in cardiovascular diseases, *Acta Pharmacol Sin.* **44**, 8-18.
doi:10.1038/s41401-022-00934-2

8. Xie, Z., Xie, T., Liu, J., Zhang, Q. & Xiao, X. (2023) Emerging role of protein O-GlcNAcylation in
liver metabolism: Implications for diabetes and NAFLD, *Int J Mol Sci.* **24**. doi:10.3390/ijms24032142

9. Li, Y., Xie, M., Men, L. & Du, J. (2019) O-GlcNAcylation in immunity and inflammation: An intricate
system (Review), *Int J Mol Med.* **44**, 363-374. doi:10.3892/ijmm.2019.4238

10. Pratt, M. R. & Vocadlo, D. J. (2023) Understanding and exploiting the roles of O-GlcNAc in
neurodegenerative diseases, *J Biol Chem.* **299**, 105411. doi:10.1016/j.jbc.2023.105411

11. Vocadlo, D. J. (2012) O-GlcNAc processing enzymes: catalytic mechanisms, substrate
specificity, and enzyme regulation, *Curr Opin Chem Biol.* **16**, 488-97. doi:10.1016/j.cbpa.2012.10.021

12. Ferrer, C. M., Sodi, V. L. & Reginato, M. J. (2016) O-GlcNAcylation in cancer biology: Linking
metabolism and signaling, *J Mol Biol.* **428**, 3282-3294. doi:10.1016/j.jmb.2016.05.028

13. Itkonen, H. M., Minner, S., Guldvik, I. J., Sandmann, M. J., Tsourlakis, M. C., Berge, V.,
Svindland, A., Schlomm, T. & Mills, I. G. (2013) O-GlcNAc transferase integrates metabolic pathways
to regulate the stability of c-MYC in human prostate cancer cells, *Cancer Res.* **73**, 5277-87.
doi:10.1158/0008-5472.Can-13-0549

14. Gu, Y., Mi, W., Ge, Y., Liu, H., Fan, Q., Han, C., Yang, J., Han, F., Lu, X. & Yu, W. (2010)
GlcNAcylation plays an essential role in breast cancer metastasis, *Cancer Res.* **70**, 6344-51.
doi:10.1158/0008-5472.Can-09-1887

15. Lynch, T. P., Ferrer, C. M., Jackson, S. R., Shahriari, K. S., Vosseller, K. & Reginato, M. J.
(2012) Critical role of O-Linked β-N-acetylglucosamine transferase in prostate cancer invasion,
angiogenesis, and metastasis, *J Biol Chem.* **287**, 11070-81. doi:10.1074/jbc.M111.302547

16. Burden, D. A., Kingma, P. S., Froelich-Ammon, S. J., Bjornsti, M. A., Patchan, M. W., Thompson,
R. B. & Osheroff, N. (1996) Topoisomerase II.etoposide interactions direct the formation of drug-
induced enzyme-DNA cleavage complexes, *J Biol Chem.* **271**, 29238-44.
doi:10.1074/jbc.271.46.29238

17. Chen, X., Zhang, T., Su, W., Dou, Z., Zhao, D., Jin, X., Lei, H., Wang, J., Xie, X., Cheng, B., Li,
Q., Zhang, H. & Di, C. (2022) Mutant p53 in cancer: from molecular mechanism to therapeutic
modulation, *Cell Death & Disease.* **13**, 974. doi:10.1038/s41419-022-05408-1

18. Hafner, A., Bulyk, M. L., Jambhekar, A. & Lahav, G. (2019) The multiple mechanisms that
regulate p53 activity and cell fate, *Nat Rev Mol Cell Biol.* **20**, 199-210. doi:10.1038/s41580-019-0110-
x

19. Chresta, C. M., Masters, J. R. & Hickman, J. A. (1996) Hypersensitivity of human testicular
tumors to etoposide-induced apoptosis is associated with functional p53 and a high Bax:Bcl-2 ratio,
Cancer Res. **56**, 1834-41.

20. di Pietro, A., Koster, R., Boersma-van Eck, W., Dam, W. A., Mulder, N. H., Gietema, J. A., de
Vries, E. G. & de Jong, S. (2012) Pro- and anti-apoptotic effects of p53 in cisplatin-treated human
testicular cancer are cell context-dependent, *Cell Cycle.* **11**, 4552-62. doi:10.4161/cc.22803

21. Noh, H., Shoemaker, J. E. & Gunawan, R. (2018) Network perturbation analysis of gene
transcriptional profiles reveals protein targets and mechanism of action of drugs and influenza A viral
infection, *Nucleic Acids Research.* **46**, e34-e34. doi:10.1093/nar/gkx1314

22. Cosse, J. P., Sermeus, A., Vannuvel, K., Ninane, N., Raes, M. & Michiels, C. (2007) Differential
effects of hypoxia on etoposide-induced apoptosis according to the cancer cell lines, *Mol Cancer.* **6**,
61. doi:10.1186/1476-4598-6-61

23. Cosse, J. P., Ronvaux, M., Ninane, N., Raes, M. J. & Michiels, C. (2009) Hypoxia-induced
decrease in p53 protein level and increase in c-jun DNA binding activity results in cancer cell
resistance to etoposide, *Neoplasia.* **11**, 976-86. doi:10.1593/neo.09632

24. Sermeus, A., Cosse, J. P., Crespin, M., Mainfroid, V., de Longueville, F., Ninane, N., Raes, M.,
Remacle, J. & Michiels, C. (2008) Hypoxia induces protection against etoposide-induced apoptosis:
molecular profiling of changes in gene expression and transcription factor activity, *Mol Cancer.* **7**, 27.
doi:10.1186/1476-4598-7-27

25. Lee, S. J. & Kwon, O. S. (2020) O-GlcNAc transferase inhibitor synergistically enhances
doxorubicin-induced apoptosis in HepG2 cells, *Cancers (Basel).* **12**. doi:10.3390/cancers12113154

26. Makwana, V., Dukie, A. S. & Rudrawar, S. (2020) Investigating the impact of OGT inhibition on
doxorubicin- and docetaxel-induced cytotoxicity in PC-3 and WPMY-1 cells, *Int J Toxicol.* **39**, 586-593.
doi:10.1177/1091581820948433

27. Liu, Y., Wang, X., Zhu, T., Zhang, N., Wang, L., Huang, T., Cao, Y., Li, W. & Zhang, J. (2019)
Resistance to bortezomib in breast cancer cells that downregulate Bim through FOXA1 O-
GlcNAcylation, *J Cell Physiol.* **234**, 17527-17537. doi:10.1002/jcp.28376

28. Nitiss, J. L. (2009) DNA topoisomerase II and its growing repertoire of biological functions, *Nat*
Rev Cancer. **9**, 327-37. doi:10.1038/nrc2608

29.   Montecucco, A., Zanetta, F. & Biamonti, G. (2015) Molecular mechanisms of etoposide, *Excli j.*
14, 95-108. doi:10.17179/excli2015-561

30.   Ganapathi, R. N. & Ganapathi, M. K. (2013) Mechanisms regulating resistance to inhibitors of
topoisomerase II, *Front Pharmacol.* **4**, 89. doi:10.3389/fphar.2013.00089

31.   Liu, Y., Yu, K., Zhang, K., Niu, M., Chen, Q., Liu, Y., Wang, L., Zhang, N., Li, W., Zhong, X., Li,
G., Wu, S., Zhang, J. & Liu, Y. (2023) O-GlcNAcylation promotes topoisomerase IIα catalytic activity
in breast cancer chemoresistance, *EMBO Rep.* **24**, e56458. doi:10.15252/embr.202256458

32.   Kang, J. G., Park, S. Y., Ji, S., Jang, I., Park, S., Kim, H. S., Kim, S. M., Yook, J. I., Park, Y. I.,
Roth, J. & Cho, J. W. (2009) O-GlcNAc protein modification in cancer cells increases in response to
glucose deprivation through glycogen degradation, *J Biol Chem.* **284**, 34777-84.
doi:10.1074/jbc.M109.026351

33.   Phueaouan, T., Chaiyawat, P., Netsirisawan, P., Chokchaichamnankit, D., Punyarit, P.,
Srisomsap, C., Svasti, J. & Champattanachai, V. (2013) Aberrant O-GlcNAc-modified proteins
expressed in primary colorectal cancer, *Oncol Rep.* **30**, 2929-36. doi:10.3892/or.2013.2794

34.   Hanover, J. A., Chen, W. & Bond, M. R. (2018) O-GlcNAc in cancer: An Oncometabolism-fueled
vicious cycle, *Journal of Bioenergetics and Biomembranes.* **50**, 155-173. doi:10.1007/s10863-018-
9751-2

35.   Zachara, N. E., Molina, H., Wong, K. Y., Pandey, A. & Hart, G. W. (2011) The dynamic stress-
induced "O-GlcNAc-ome" highlights functions for O-GlcNAc in regulating DNA damage/repair and
other cellular pathways, *Amino Acids.* **40**, 793-808. doi:10.1007/s00726-010-0695-z

36.   Van Ravenstein, S. X., Mehta, K. P., Kavlashvili, T., Byl, J. A. W., Zhao, R., Osheroff, N., Cortez,
D. & Dewar, J. M. (2022) Topoisomerase II poisons inhibit vertebrate DNA replication through distinct
mechanisms, *Embo j.* **41**, e110632. doi:10.15252/embj.2022110632

37.   Weiss, M., Anderluh, M. & Gobec, M. (2021) Inhibition of O-GlcNAc transferase alters the
differentiation and maturation process of human monocyte derived dendritic cells, *Cells.* **10**, 3312.

38.   Brady, C. A. & Attardi, L. D. (2010) p53 at a glance, *J Cell Sci.* **123**, 2527-32.
doi:10.1242/jcs.064501

39.   Pascal, J. M. (2018) The comings and goings of PARP-1 in response to DNA damage, *DNA*
Repair (Amst). **71**, 177-182. doi:10.1016/j.dnarep.2018.08.022

40.   Shiokawa, D., Shika, Y., Araki, S., Sunaga, S., Mizuta, R., Kitamura, D. & Tanuma, S. (2007)
Stage-specific expression of DNasegamma during B-cell development and its role in B-cell receptor-
mediated apoptosis in WEHI-231 cells, *Cell Death Differ.* **14**, 992-1000. doi:10.1038/sj.cdd.4402086

41.   Saleh, T., Tyutyunyk-Massey, L., Murray, G. F., Alotaibi, M. R., Kawale, A. S., Elsayed, Z.,
Henderson, S. C., Yakovlev, V., Elmore, L. W., Toor, A., Harada, H., Reed, J., Landry, J. W. &
Gewirtz, D. A. (2019) Tumor cell escape from therapy-induced senescence, *Biochem Pharmacol.* **162**,
202-212. doi:10.1016/j.bcp.2018.12.013

42.   Katayama, M., Kawaguchi, T., Berger, M. S. & Pieper, R. O. (2007) DNA damaging agent-
induced autophagy produces a cytoprotective adenosine triphosphate surge in malignant glioma cells,
Cell Death Differ. **14**, 548-58. doi:10.1038/sj.cdd.4402030

Figure legends

Fig. 1. OSMI-1 enhances the sensitivity of HepG2 cells to etoposide.

(A-C) Effect of etoposide combined with OSMI-1 on HepG2 human liver cancer cell viability assessed through the CCK viability assay ($n = 5$) and crystal violet staining ($n = 5$) 24 h post-treatment. Cotreatment with 20 μM OSMI-1 significantly reduced cell viability at all etoposide concentrations compared with etoposide alone. *** $P < 0.001$ versus the control group; # $P < 0.05$, ## $P < 0.01$, and ### $P < 0.001$ versus the vehicle control in the etoposide treatment group at the same concentration (mean ± SD). (D) Viability analysis via GF-AFC revealed a 1.64-fold reduction in the IC_{50} value of etoposide, from 99.59 to 60.68 μM, with 20 μM OSMI-1 ($n = 3$). (E) Membrane integrity assays showed a 4.07-fold decrease in the EC_{50} value of etoposide, from 176.63 to 43.41 μM, when combined with 20 μM OSMI-1 ($n = 3$). (F) Cell viability assays in Hep3B human liver hepatocellular carcinoma, A549 human lung carcinoma, and Caco-2 human colorectal adenocarcinoma cell lines confirmed a greater reduction in viability with the etoposide and OSMI-1 cotreatment ($n = 4$). ** $P < 0.01$ and *** $P < 0.001$ versus the control group; # $P < 0.05$ and ## $P < 0.01$ versus etoposide treatment alone (mean ± SD). Data are means from at least three independent experiments. Significance statistical analysis was performed with one-way or two-way ANOVA with *post hoc* Tukey test. ns, not significant.

Fig. 2. Etoposide reduces O-GlcNAcylation in a dose-dependent manner without directly interacting with OGT.

(A) Etoposide treatment for 24 h resulted in dose-dependent reduction in O-GlcNAc expression in HepG2 cells. (B) Immunocytochemistry with O-GlcNAc antibody (RL2) (white scale bar: 10 μm) confirmed the findings in (A). (C) Etoposide and OSMI-1 cotreatment reduced O-GlcNAc expression levels comparable to OSMI-1 treatment alone. (D, E) OGT inhibition assays showed that etoposide did not significantly reduce OGT activity at concentrations below ~200 μM, whereas OSMI-1 exhibited an IC_{50} of 4.47 μM ($n = 3$; mean ± SD). Thus, etoposide decreases O-GlcNAc dose-dependently without directly interacting with OGT. Data are means from at least three independent experiments.

Fig. 3. Etoposide's anticancer effects are mediated by downregulation of O-GlcNAc expression.

(A) Cell viability assays demonstrated that OSMI-1 (an OGT inhibitor) significantly increased the anticancer effect of etoposide in a concentration-dependent manner, whereas Thiamet G (an OGA inhibitor) did not alter viability ($n = 4$; mean ± SD). (B, C) Crystal violet staining assay corroborated the findings in (A) ($n = 4$; mean ± SD). (D, E) Western blot and immunocytochemistry analyses at 24 h post-treatment revealed that etoposide and OSMI-1 cotreatment reduced O-GlcNAc expression levels more than etoposide treatment alone. Thiamet G restored the O-GlcNAc expression levels lowered by etoposide (white scale bar: 10 μm). ** $P < 0.01$ and *** $P < 0.001$ versus the control group; # $P < 0.05$ and ## $P < 0.01$ versus the etoposide treatment alone. Data are the means from at least three independent experiments. Significance statistical analysis was performed with one-way ANOVA with *post hoc* Tukey test. ns, not significant.

Fig. 4. OSMI-1 significantly enhances etoposide-induced apoptosis.

(A) FACS analysis using Annexin V and 7-AAD staining revealed that cotreatment with OSMI-1 and etoposide for 24 h significantly increased early apoptosis compared with etoposide treatment alone. (B) Quantitative

luminescence analysis showed a 2.25-fold reduction in the EC$_{50}$ of etoposide, from 61.94 to 27.52 μM, with 20 μM OSMI-1 ($n = 3$). (C) The apoptosis-inducing effect of the cotreatment was significantly greater than that of the control group from 4 h and that of etoposide treatment alone from 12 h post-treatment ($n = 4$). * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$ versus the control group at each time point; ### $P < 0.001$ versus the etoposide treatment alone at each time point (mean ± SD). Data are means from at least three independent experiments. Significance statistical analysis was performed with one-way ANOVA with *post hoc* Tukey test. ns, not significant.

Fig. 5. Etoposide and OSMI-1 cotreatment induces DNA fragmentation during apoptosis.

(A, B) TUNEL assays demonstrated dose-dependent DNA fragmentation in HepG2 cells treated with etoposide and OSMI-1 for 24 h. OSMI-1 induced double DNA breaks in a dose-dependent manner. Thus, reduced O-GlcNAcylation amplifies etoposide-mediated apoptosis induction ($n = 5$; white scale bar: 10 μm). ** $P < 0.01$ versus the etoposide treatment alone (mean ± SD). Significance statistical analysis was performed with one-way ANOVA with *post hoc* Tukey test. ns, not significant.

Fig. 6. OSMI-1 enhances etoposide's anticancer effect via activation of the p53/caspase-3/PARP1 pathway.

(A) Western blot analysis revealed increased expression of p53 and cleaved-PARP1 (apoptosis markers) with 24 h of 20 μM etoposide and 20 μM OSMI-1 cotreatment compared to etoposide alone. (B) Quantitative luminescence analysis showed that caspase-3/7 activity was also higher with etoposide and 20 μM OSMI-1 cotreatment ($n = 4$). *** $P < 0.001$ versus the etoposide treatment alone (mean ± SD). Data are means from at least three independent experiments. Significance statistical analysis was performed with two-way ANOVA with *post hoc* Tukey test. ns, not significant.

Fig. 7. Downregulation of O-GlcNAcylation augments etoposide-induced p53-mediated apoptosis.

Etoposide induces apoptosis by forming a ternary complex with topoisomerase II (Topo II) and DNA, inhibiting DNA strand religation and causing DNA damage. This activates the p53-mediated apoptosis pathway, increasing caspase-3/7 (Cas-3/7) activity. Activated caspase-3/7 cleaves PARP1, leading to apoptosis and thereby maintaining genome integrity. Although etoposide downregulates O-GlcNAcylation during this process, the exact mechanism remains unclear. Cotreatment with OSMI-1, an OGT inhibitor, further suppresses O-GlcNAcylation, enhancing etoposide's p53-mediated apoptosis pathway.

Graphical Abstract

Etoposide, a topoisomerase II inhibitor, reduces O-GlcNAcylation in HepG2 liver cancer cells. Further inhibition of O-GlcNAc transferase by OSMI-1 enhanced etoposide-induced apoptosis, lowering the IC50 for viability and increasing the EC50 for cytotoxicity. This synergy was mediated by the p53, caspase-3/7, PARP1 pathway, suggesting that targeting O-GlcNAcylation may improve etoposide efficacy.

Figure 1

Figure 2

Figure 3

Figure 4

(A)

(B) **(C)**

Figure 5

(A)

(B)

TUNEL Assay

Etoposide [µM]	—	20	20	20
OSMI-1 [µM]	—	—	10	20

Figure 6

(A)

(B)

Caspase-3/7 Activity

Figure 7