A novel multiplex RNAi therapy simultaneously targets *Hif1a* and *Hif2a* to defy retinal

degeneration in two models of AMD

Lynn J. A. Ebner[1,2], Cornelia Imsand[1], Duygu Karademir[1,3], Florian Peters[1,4], Eva Kiessling[1,5],

Antonia Fottner[1], Claudia Matter[1], Diego S. Fajardo[6], Luca Merolla[1], Gabriele M. Wögenstein[1],

Ioanna Tsioti[1], Larissa P. Govers[1], Frank Blaser[7], Isabelle Meneau[7], Sanford L. Boye[6],

Shannon E. Boye[8], Christian Grimm[1,+], Marijana Samardzija[1,*,+]

[1] Laboratory for Retinal Cell Biology, Department of Ophthalmology, University Hospital Zurich,
University of Zurich, Wagistrasse 14, Schlieren, 8952, Zurich, Switzerland.

[2] Present address: Roche Pharma Research and Early Development, Roche Innovative Center, F.
Hoffmann-La Roche Ltd, Grenzacherstrasse 124, Basel, 4070, Basel-Stadt, Switzerland
(lynn.ebner@roche.com)

[3] Present address: Institute of Molecular and Clinical Ophthalmology Basel, Switzerland

[4] Present address: Department of Clinical Research, University of Basel, University Hospital Basel,
Basel, Switzerland

[5] Present address: Lonza AG, Rottenstrasse 6, 3930 Visp, Switzerland

[6] Department of Pediatrics, Powell Gene Therapy Center, University of Florida; Gainesville, FL, USA.

[7] Department of Ophthalmology, University Hospital Zurich, Switzerland

[8] Department of Pediatrics, Division of Cellular and Molecular Therapy, University of Florida, Gainesville,
Florida, USA

[+] Equal contributions

[*] Corresponding author

lynn.ebner@roche.com
dallo.imsand@bluewin.ch
duygu.sigurdsson@iob.ch
florian.peters@usb.ch
eva.kiessling@lonza.com
antonia.fottner@usz.ch
Claudia.matter@usz.ch
diegof@ufl.edu
luca.merolla@uzh.ch
gabriele.woegenstein@uzh.ch
ioanna.tsioti@uzh.ch
larissa.govers@uzh.ch
frank.blaser@usz.ch
isabelle.meneau@usz.ch
sboye@ufl.edu
shaire@ufl.edu
cgrimm@opht.uzh.ch
marijana.samardzija@usz.uzh.ch

Abstract

Age-related tissue changes lead to reduced oxygen delivery to photoreceptors and the retinal pigment epithelium (RPE), and contribute to the pathology of age-related macular degeneration (AMD). The implication of hypoxia-inducible factors (HIFs) in this process makes them good candidates as therapeutic targets for AMD. We developed a multiplex dual-acting therapy utilizing the shRNAmir system, delivered by a single AAV, that reduces mRNA levels of *Hif1a* in photoreceptors and *Hif2a* in the RPE. This RNA interference-based strategy demonstrated a strong therapeutic effect, potently preserving photoreceptors and the RPE in two models of pseudo- and true hypoxia up to 61 weeks post-injection. The efficacy of our dual-acting virus proved superior to single-acting viruses targeting only *Hif1a* in photoreceptors or *Hif2a* in the RPE. By targeting a common, conserved disease pathway, this gene-agnostic RNAi therapy shows significant potential to protect tissues from chronic hypoxic insults in complex diseases such as AMD.

Keywords:

Hypoxia; HIF1; HIF2; retinal degeneration; age related macular degeneration; photoreceptor;

retinal pigment epithelium; gene therapy; cell specific RNA interference; AAV; multiplex RNAi

Background

Age-related macular degeneration (AMD) is a leading cause of vision loss in the elderly

worldwide, primarily affecting the photoreceptors and the retinal pigment epithelium (RPE) in

the macular region. These cells degenerate as the disease progresses, leading to reduced

visual acuity and eventually central vision loss. Current therapeutic strategies largely focus on

targeting secreted pro-angiogenic proteins involved in disease progression. However,

emerging RNA-based therapeutics hold considerable promise as alternatives, offering the

potential to provide more effective treatments by targeting intracellular pathways that are

otherwise difficult to drug [1].

Age-related tissue changes such as reduced choroidal thickness, vascular density and blood

flow, ghost vessel formation, thickening of Bruch's membrane and accumulation of drusen

deposits [2-5] lead to reduced tissue oxygenation and increased VEGFA secretion, an

established driver of neovascularization in the exudative (wet) form of AMD. Notably, these

age-related tissue changes may also be relevant to the non-exudative (dry) form of AMD, as

they have been correlated with drusen formation [2-5] and suggested to precede photoreceptor

or RPE degeneration [6].

Given the critical role of hypoxia-inducible factors (HIFs) in the cellular response to low oxygen

levels, chronic activation of the HIF pathway may contribute significantly to the development

of AMD. While chronic activation of HIF2, but not HIF1, in the RPE causes RPE atrophy and

retinal thinning [7], constitutively active HIF1, but not HIF2, in photoreceptor cells induces

photoreceptor degeneration in animal models of AMD [8,9]. Because cell loss and degeneration

can be prevented by the genetic inactivation of *Hif1a* in photoreceptors [8,9] and *Hif2a* in the

RPE [7], a targeted therapy to reduce the expression of the respective HIF isoform in the two

cell types may be an effective approach to manage disease pathology and preserve vision in

AMD patients.

The need to target two distinct genes in two different cell types presents a unique challenge in

developing an effective therapy. To meet this challenge, we have combined an RNA

interference (RNAi) strategy with the adeno-associated virus (AAV) delivery system that allows

targeted and long-lasting expression of therapeutic genes in retinal cells. Conventional

therapeutic RNA knockdown strategies using small interfering RNAs (siRNAs) are often limited

because they interfere with endogenous microRNA biogenesis, lead to an overload of transport

proteins, or cause other undesirable side effects [10-12]. To circumvent such problems, we have

generated a multiplex RNAi approach using the microRNA-based shRNAmir system, which

directs the therapeutic RNAs into the miRNA pathway already at the level of DROSHA in the

cell nucleus [13,14]. Sequential processing by DROSHA and DICER ensures the generation of

precisely cleaved siRNAs while minimizing interference with the natural microRNA processing

machinery [14]. This increases efficiency while reducing off-target effects and immune responses

[15,16]. The use of the miR-E instead of the conventional miR-30 backbone for shRNA expression

increases the knockdown efficiency of target genes and allows cell type specific expression

when placed in the 3'UTR of genes that are controlled by Pol-II promoters [17]. Using this system,

we developed a novel dual-acting RNA-based therapy that simultaneously targets *Hif1a* in

photoreceptors and *Hif2a* in the RPE, delivered by a single AAV. The strong and long-lasting

therapeutic potential of this system was demonstrated in two hypoxia-related animal models

of AMD.

Methods

Animals

Animal experimentation was approved by the Cantonal Veterinary Office of Zürich, Switzerland (license numbers: ZH019/2019 and ZH105/2022) and adhered to the ARVO statement for the use of Animals in Ophthalmic and Vision Research. Wild type (B6/J-Crl1*)* and transgenic mice were maintained as breeding colonies at the Laboratory Animal Service Center (LASC) of the University of Zürich with a 14/10 h light-dark cycle with an average light intensity of 60-150 lux at cage level. Mice had access to food and water *ad libitum*. *Vhl*$^{fl/fl}$ mice [18] were mated to *Vhl*$^{fl/fl}$ mice heterozygous for OpsinCre (LMOPC1, [19]) to generate *Vhl*$^{fl/fl}$*;OpsinCre* (*Rod*$^{\Delta Vhl}$) mice and *Vhl*$^{fl/fl}$ control littermates as described (Lange et al., 2011). *Vegfa*$^{fl/fl}$ animals were kindly provided by Christian Stockmann, Institute of Anatomy, University of Zürich [20]. Individual eyes of mice of both sexes were randomly assigned to treatment or control.

Human tissue samples

Human donor eyes without cornea and lens were received from the eyebank of the University Hospital Zürich in ice-cold PBS. The use of the tissue for research was approved by the ethics committee of Zürich, Switzerland (BASEC-Nr: PB_2017-00550 and 2020-01856) and adhered to the tenets of the Declaration of Helsinki. The summarized donor information is given in Table S1. The vitreous was carefully removed and four incisions were used to flatmount the tissue. Retinal and RPE samples were collected from the donor eyes with no clinical diagnosis of retinal disease. RNA was isolated from the samples using the RNeasy kit (Qiagen, Hilden, Germany). cDNA synthesis and real-time PCR were performed using human specific primer pairs (Table 1), following the same methods as described for the mouse samples (see below).

Cloning and generation of AAVs

shRNAmir DNA fragments containing *shHif1a*, *shHif2a* or *shCtrl* sequences were designed using the miR-E backbone following the recommendation by Fellmann *et al.* [17] (Fig. S1A-C) and produced by Bio-Synthesis (Bio-Synthesis Inc. TX, USA). The *shHif1a* and *shHif2a* RNAmiR fragments were cloned into the 3'UTRs of GFP and mCherry, respectively. To achieve cell-specificity, the G-protein-coupled kinase 1 (GRK1) promotor was used for expression of GFP (and thus *shHif1a*) in photoreceptors, and the vitelliform macular dystrophy 2 (VMD2, alias bestrophin 1) promoter for expression of mCherry (and thus *shHif2a*) in the RPE. This dual-acting viral construct is named shHif from here on (Fig. 2D). The shCtrl construct had a scrambled shRNA sequence cloned into the miR-E backbone in the 3'UTRs of both fluorescent genes (Fig. 2D). The complete plasmid sequences of plasmids *'shHif'* and *'shCtrl'* are given in supplemental data SD1. Plasmids for the single-acting AAVs contained either the GRK1-GFP-shHif1a or the VMD2-mCherry-shHif2a cassette (Fig. 2D). shHif, shHif1, shHif2 and shCtrl viruses were packed into an AAV8 with the three mutations Y447F, Y733F and T494V (AAV8-TM) [21]. AAV4::VMD2-Cre-P2A-GFP [22] was used to knockout *Vegfa* in the RPE of *Vegfa^{fl/fl}* mice and AAV4::VMD2-GFP served as control virus. Specifics of all AAVs are provided in Table 2. All AAVs had type 2 ITRs and were produced by the Viral Vector Facility of the University of Zürich.

Subretinal injections

Injections were performed between post-natal days (PND) 27-30 as described by Gasparini et al [23]. Briefly, pupils were dilated with 1% cyclogyl (Alcon Pharmaceuticals, Fribourg, Switzerland) and one drop of 5% neosynephrin (Ursapharm Schweiz GmbH, Hünenberg, Switzerland), and corneas were kept moist with viscotears (Bausch & Lomb Swiss AG, Zug, Switzerland). Mice were anesthetized with a subcutaneous injection of ketamine (85 mg/kg, Parke-Davis, Berlin, Germany) / xylazine (10 mg/kg, Bayer AG, Leverkusen, Germany) and placed on a heating pad set to 37°C during the procedure. The head was held by a stereotactic

adapter (Hugo Sachs Elektronik – Harvard Apparatus GmbH, March-Hugstetten, Germany)

and the temporal sclera punctured with a 30G needle just beneath the ora serrata. A 5 µL

Hamilton syringe (Hamilton Bonaduz AG, Bonaduz, Switzerland), placed in a micromanipulator

(H. Saur Laborbedarf, Reutligen, Germany), was inserted gently through the pre-punctured

site for transvitreal subretinal injections of 1 µL AAV solution approximately 15-25° nasally of

the optic nerve. All AAVs were applied at a concentration of $5x10^8$ vg/µL, except for

AAV4::VMD-Cre-P2A-eGFP ($1x10^9$ vg/µL) when used to generate $RPE^{\Delta Vegfa}$ mice for

characterization (Fig. 4). To visualize and control injections, the injection solution was

supplemented with 10% fluorescein (0.1 mg/mL, Akorn Inc., IL, USA). After injections,

anesthesia was reversed with atipazemole (2 mg/kg, Graeub, Bern, Switzerland) and mice

were placed on a heating pad until fully awake.

Funduscopy, optical coherence tomography, angiography, and volume scans

Anesthesia and pupil dilation were performed as described above. One drop of 2% methocel

(OmniVision AG, Neuhausen, Switzerland) was applied to keep the eyes moist. Fundus

images and optical coherence tomography (OCT) scans were acquired using the Micron IV

system (Phoenix Research Labs, Pleasanton, CA, USA) as described [24]. For fluorescein

angiography mice were injected intraperitoneally with 30 µL of 2% fluorescein (Akorn Inc., IL,

USA) and retinal vasculature was imaged with the Micron IV system. For *in vivo* quantification

of retinal thickness, volume scans were performed using a Spectralis OCT-2 device

(Heidelberg Engineering Inc., Heidelberg, Germany) with a 25-diopter lens. Animals were

placed on a custom-built holder and kept warm with an integrated water heating system. An

atraumatic focal contact lens was placed on the cornea moistened with a drop of viscotears

(Bausch & Lomb) to improve imaging quality. The optic scan area was aligned to the optic

nerve head and 25 consecutive high-resolution B-scans, each consisting of 1024 A-scans,

were acquired. Automatic real-time tracking was set to 50. The built-in semi-automated

segmentation tool (Spectralis software version 6.7.13.0, Eye Explorer software version

1.9.14.0) was utilized, and manual adjustments were done by blinded, experienced users (L.E.,

C.I., M.S.) to measure outer nuclear layer (ONL) thickness *in vivo*. To delineate fundus areas,

an Early Treatment Diabetic Retinopathy Study (ETDRS) circular grid with 9 subfields (1, 2.22,

3.45) was aligned to the optic nerve head (ON). ONL thickness was quantified in two ($RPE^{\Delta Vegfa}$

mice) or seven subfields ($Rod^{\Delta Vhl}$ mice). These subregions were marked with orange outlines

in the respective images. For quantification, the data were first averaged per subfield using

ETDRS grids (left), then the average across all subfields was calculated and plotted for each

eye (graphs, right).

Scotopic and photopic electroretinogram

Mice were dark-adapted overnight. Pupils were dilated and animals anesthetized as described

above. Electroretinograms (ERG) were recorded simultaneously from both eyes using the

Diagnosys Celeris rodent ERG device (Diagnosys, Cambridge, UK). Stimulus intensities

ranging from $8x10^{-6}$ to 3 cd*s/m^2 were used for scotopic measurements. After a 5-min exposure

to rod-suppressive background light (30 cd/m^2), photopic recordings were acquired using

single-flash intensities ranging from 1 to 200 cd*s/m^2 with the background light on. For each

stimulus intensity, five traces were averaged under scotopic and ten traces under photopic

conditions.

RNA isolation and real-time PCR

Mice were euthanized with CO_2 inhalation followed by decapitation. Retinas were isolated

through a slit in the cornea and snap-frozen in liquid nitrogen. In some instances, the remaining

eyecup (EC) containing the RPE was cleaned from excessive tissue including the cornea and

snap-frozen. RNA from tissues or cells was isolated using an RNA isolation kit (NucleoSpin

RNA, Macherey-Nagel GmbH & co.KG, Düren, Germany) according to the manufacturer's

instructions, including a DNase digestion step on the column. First-strand cDNA synthesis was

performed by M-MLV reverse transcriptase (Promega, Dübendorf, Switzerland) using 1 µg

total RNA and oligo-dT primers. Gene expression was analyzed via semi-quantitative real-time

PCR (QuantStudio 3, Thermo Fisher Scientific, MA, USA) using 10 ng of cDNA template and

PowerUp SYBR green Master Mix (Thermo Fisher Scientific). Primers pairs (Table 1) were

designed to span large intronic regions and avoid known single nucleotide polymorphisms in

the mouse sequence. Normalization was performed with β-Actin (*Actb)* as housekeeping gene,

and relative expression was calculated using the comparative threshold cycle method ($2^{-\Delta\Delta CT}$)

[25].

Morphology and immunofluorescence

Morphology and immunofluorescence analyses were performed as described [26]. For

morphology, dorsally marked eyes were enucleated from non-perfused (*Rod$^{\Delta Vhl}$*) or perfused

(*RPE$^{\Delta Vegfa}$*; 4% paraformaldehyde (PFA)) mice and subsequently fixed in 2.5% glutaraldehyde

at 4°C overnight. Eyes were postfixed in 1% osmium tetroxide and embedded in Epon812.

Nasal-temporal sections (0.5 µm) were cut through the optic nerve head, counterstained with

toluidine blue, and analyzed by light microscopy (AxioImager Z2, Carl Zeiss AG, Feldbach,

Switzerland). ONL thickness and photoreceptor segment (PS) length were measured every

200 µm (from the optic nerve head) along reconstructed panorama images using Adobe

Photoshop CS6 ruler tool (Adobe Systems, Inc.).

Eyes for immunofluorescence analysis were labelled dorsally and fixed in 4% PFA for 1 h.

After removal of the lens and post-fixation for an additional hour, eyes were cryoprotected in

30% sucrose (Sigma-Aldrich, St. Louis, MO, USA) overnight, embedded and frozen in freezing

medium (O.C.T., Leica Biosystems, Nussloch, Germany). Eyes were stored at -80°C until

sectioned into 12 µm thin nasal-temporal sections. Immunolabelling was performed overnight

at 4°C with primary antibodies (Table 3) in blocking solution containing 3% normal goat serum

(Sigma-Aldrich) and 0.3% Triton X-100 (Sigma-Aldrich) in phosphate buffer. On the next day,

sections were washed and incubated with the appropriate secondary antibodies in blocking

solution for 1 h. Counterstaining was done with 4',6-diamidino-2-phenylindole (DAPI) before

sections were mounted with MOWIOL (Sigma-Aldrich) antifade medium and imaged using a

fluorescence microscope (AxioImager Z2, Carl Zeiss AG). Pixel intensity measurements were

performed using ImageJ (Version 1.52a) by analyzing 250 µm long regions within transduced

and non-transduced areas. The automatic threshold function was used to identify signal and

background pixels. Pixels with an intensity of '0' were counted as background/gaps. The

cumulative length of these zero-intensity regions was calculated and presented as gap size

per 250 µm.

In-situ hybridization

In-situ hybridization was performed with the RNAscope 2.5 HD RED Detection Kit (Cat. No.

322350, BioTechne/ACD, Minneapolis, USA). Cryosections were prepared as described

above, baked at 60°C for 30 min, and post-fixed in 4% PFA for 15 min at 4°C. Slides were

dehydrated for 5 min each in 50%, 70% and 100% ethanol and dried for 10 min at 60°C.

Antigen retrieval was performed using hydrogen peroxide for 10 min, washed and followed by

overnight incubation with target retrieval buffer (BioTechne/ACD). Protease Plus treatment

was performed for 15 min at 40°C, followed by 2 x 30 s washings with H_2O, and incubation

with probes (Mm-Hif1a Cat. No. 313821, Mm-Epas1 Cat. No. 314371, Mm-Vegfa-O3 Cat. No.

521471; BioTechne/ACD) for 2 h at 40°C. Hybridized tissue sections were washed 2 x 5 min

with wash buffer and incubated at 40°C with each amplifier (1-6) for 15-30 min, with two 5-

minute washes conducted between each amplification step. The positive control probe _Mm-_

Polr2a (Cat. No. 312471, BioTechne/ACD) and the negative control probe diluent (Cat. No.

300041, BioTechne/ACD) were used to validate experimental conditions. Slides were stained

in 50% hematoxylin for 20 s, destained in H_2O, incubated in 0.02% ammonia water for 10 s,

and washed again. Finally, slides were dried for 30 min at 60°C and mounted with Ecomount

(Cat. No. 320409, BioTechne/ACD).

Retina- and RPE-flatmounts

After eyes were enucleated and incubated in 4% PFA for 10 min, corneas were opened and eyes incubated for another 20 min in PFA. The eyes were cut along the ora serrata to remove the cornea and placed on a nylon membrane (Boehringer Ingelheim, Ingelheim am Rhein, Germany). 5-6 radial incisions were made to flatten the eyecup and retina, and the lens was removed. Retina and eyecup were then separated and held flat between 2 membranes during an additional hour of fixation in 4% PFA. Flattened retinas or eyecups were removed from the membrane, washed with PBS, and blocked with 3% normal goat serum (Sigma-Aldrich, St. Louis, MO, USA) and 0.3% Triton X-100 (Sigma-Aldrich) in PBS for 1-2 h. Incubation with primary antibodies (Table 3) was overnight at 4°C. Flatmounts were washed 3 x 10 min with PBS and incubated with the appropriate secondary antibodies for 1 h at room temperature. Subsequently, retinas and eyecups were washed 3 x 10 min with PBS, mounted, and imaged using a fluorescence microscope AxioImager Z2 (Carl Zeiss AG) or Leica DMI6000B (Leica Microsystems, Wetzlar, Germany).

Choroidal flatmounts

After an initial 30 min fixation in 4% PFA, eyes for choroidal flatmounts were washed in PBS and incubated at 55°C in pre-warmed 10% H_2O_2 for 2 to 2.5 h to bleach RPE and choroidal pigments. The cornea was removed by cutting along the ora serrata, and the eyes were placed on a nylon membrane. Lens and retina were removed, and four radial incisions were made to flatten the choroidal eyecup. The flattened tissue, secured between 2 nylon membranes, underwent an additional fixation step in 4% PFA for 1 h. Staining for podocalyxin-like (PODXL) proceeded as above, with the modifications that incubation with the primary antibody lasted for 2 days, and 3% horse serum (Sigma-Aldrich) was used for blocking.

Lentiviral-mediated gene silencing in 661W cells and hypoxic exposure

661W [27] and HEK293T (ATCC CRL-3216) cells were cultured in DMEM, 10% heat-inactivated FBS (Gibco, Thermo Fisher Scientific) and 1% penicillin-streptomycin (Gibco, Thermo Fisher Scientific) at 37°C and 5% CO_2. Lentiviral-pseudotyped particles were used to generate stable cell lines expressing shRNA against *Hif1a* (TRCN0000232222) or *Hif2a* (TRCN0000082303, TRCN0000082307, TRCN0000436203, TRCN0000416474, TRCN0000428636, TRCN0000082306) or a non-targeted scrambled shRNA (shCtrl, SHC002) (Sigma-Aldrich). To prepare lentiviral particles, HEK293T cells were seeded in 75 cm^2 culture flasks and co-transfected with shRNAs in combination with the ViraPower lentiviral expression vector system and Lipofectamin 3000 (Invitrogen, Thermo Fisher Scientific) in serum-free medium. Medium was replaced the next day with medium containing FBS. The culture supernatants containing the lentivirus particles were collected 48-72 h post-transfection, cleared by centrifugation at 200 g for 5 min, and filtered through a 0.45 µm pore size filter (Merck & Cie, Schaffhausen, Switzerland). Filtrates were applied in a 1:1 mixture with medium containing 6 µg/mL polybrene (Sigma-Aldrich) to infect 661W cells. Infected cells were selected with 2 µg/mL puromycin for 2 weeks. Stable cell lines expressing *shHif1a*, *shHif2a,* and *shCtrl* were seeded at a density of 30,000 cells/mL in a 6-well plate and incubated overnight. Prior to hypoxic exposure, medium was exchanged, and cells were exposed to normoxia (21% O_2) or hypoxia (0.2% O_2 for 24 h). Subsequently, cells were harvested and prepared for Western blotting (see below) or real-time PCR (see above).

Western blotting

Cells were washed with PBS and lysed with 50 µL lysis buffer (1 M Tris, 150 mM NaCl, 1 mM Na_3VO_4, 1% Triton X-100 containing protease inhibitor (1 tablet per 10 ml solution; cOmplete ULTRA Tablets-Mini, F. Hoffmann-La Roche AG, Basel, Switzerland). Cells were scraped off the plate, homogenized by sonication for 10 min and centrifuged (3 min 1000 rcf) at room temperature. The protein concentration was determined in the supernatant with a BCA protein

assay kit (Pierce™, Thermo Scientific) according to manufacturer's instructions. DNA present

in the homogenate was digested at 37°C for 1 h with 125 units benzonase (Sigma-Aldrich, St.

Louis, MO, USA) before 70 µg of proteins were used for standard Western blotting. Primary

antibodies (Table 3) were diluted in 5% non-fat blocking milk (BioRad, Cressier, Switzerland)

in TBST and incubated over night at 4°C with mild agitation. HRP-conjugated secondary

antibodies were used for signal detection by the Western lightning chemiluminescence reagent

(PerkinElmer, Waltham, MA, USA) and exposure to X-ray films.

Statistical analysis

Statistical analysis was performed using GraphPad Prism (GraphPad Software, San Diego,

CA, USA). Significance was tested by an unpaired t-test when two independent groups were

compared. When more than 2 groups were compared, a one-way ANOVA followed by a

Tukey's (when all groups were compared to each other) or Dunnett's (when all groups were

compared to a specific group) multiple comparison test was applied. Data were considered to

be statistically different with $p \leq 0.05$. P-values are given directly in figures or in Table S2. All

tests and number of animals/samples are indicated in the figure legends.

Results

Age-related tissue changes impair oxygen delivery to the RPE and photoreceptors leading to a chronic molecular response to hypoxia that can be harmful to these delicate cells in the aging eye. RNA interference may be an ideal therapeutic strategy to attenuate this response and support long-term cell survival. To investigate the impact of ageing on the expression of known hypoxia-regulated genes [28-30] associated with AMD [29,31,32], we determined the expression of *VEGFA*, angiopoietin like 4 (*ANGPTL4*) and serpin family E member 1 (*SERPINE1*) in the retina and RPE of young (17 – 55 y, median: 35 y) and old (71 – 94 y, median: 82 y) human donor eyes (Table S1). G protein subunit alpha transducin 2 (*GNAT2*), a gene essential for phototransduction in cones [33], and retinoid isomerohydrolase RPE65 required for the visual cycle in the RPE [34] were included as controls. The observed increased expression of all three hypoxia-regulated genes, but not of the control genes, in the neuronal retina and the RPE of old donor eyes (Fig. 1) points to an elevated activity of HIF transcription factors with age. Reducing the expression of HIF transcription factors in the retina and RPE by RNA interference may attenuate their activity, reduce the expression of AMD-related HIF target genes and thus provide significant therapeutic benefits.

Efficient targeting of *Hif1a* and *Hif2a* using the shRNAmir system

Chronically active HIF1 is the main driver for degeneration of photoreceptors, while HIF2 is the pathologic isoform in RPE cells [7,8]. To prevent hypoxia-related degeneration, we developed an RNA interference strategy that simultaneously silences HIF1 in photoreceptors and HIF2 in the RPE by targeting the mRNAs for the respective HIF alpha subunits in the specific cell types. To reduce *Hif1a* expression, we used a published and previously validated shHif1a RNA sequence that targets both the mouse and human sequence [8] and refer to this sequence as shHif1. To identify an efficient shHif2a sequence, six different shRNAs (shHif2_1-6) were stably expressed in 661W cells and tested for their efficiency to reduce expression of *Hif2a*.

All six shRNAs were similarly efficient and reduced *Hif2a* mRNA levels in hypoxic cells (6 h,

0.2% O_2) to 2 - 9% when compared to the shCtrl sequence (Fig. S2). Since mouse shHif2_6

has only one mismatch to the human *HIF2A* RNA, all further experiments were performed

using this sequence, which will be referred to as shHif2.

Hif1a and *Hif2a* mRNA and protein levels were measured in normoxic and hypoxic (24 h, 0.2%

O_2) 661W cells stably expressing shHif1, shHif2 or shCtrl (Fig. 2B, C). *Hif1a* mRNA expression

was reduced by about 65% in both normoxic and hypoxic 661W-shHif1 cells when compared

to 661W-shCtrl cells while it remained at control levels in 661W-shHif2 cells (Fig. 2B). *Hif2a*

expression was reduced by roughly 80% in normoxic and hypoxic 661W-shHif2 but not 661W-

shHif1 cells. Interestingly, an 86% increase in *Hif2a* levels was observed in hypoxic 661W-

shHif1 cells, suggesting a compensatory induction of *Hif2a* upon *Hif1a* knockdown in hypoxia,

as reported by others [35]. The hypoxic state of the cells was confirmed by the significantly

increased expression level of solute carrier 2a1 (*Slc2a1,* alias *Glut1,* Fig. 2B), a gene primarily

regulated by HIF1. The diminished but still measurable *Slc2a1* expression levels in hypoxic

661W-shHif1 cells indicated a strongly but not fully blocked HIF1 activity, which is expected

after robust but not complete *Hif1a* gene silencing (Fig. 2B). Western blots confirmed increased

HIF1A and HIF2A protein levels in hypoxic cells and their specific reduction by the

corresponding shRNAs (Fig. 2C). Thus, the selected shRNAs efficiently downregulated *Hif1a*

and *Hif2a* at both the mRNA and protein levels.

To achieve a simultaneous and effective *in vivo* downregulation of *Hif1a* in photoreceptors and

Hif2a in the RPE, we placed the corresponding shRNAs (Fig. S1) into miR-E scaffolds [17] and

positioned them in the 3'UTRs of GFP (shHif1) and mCherry (shHif2). GRK1 [36] and VMD2 [37]

promoters were used to drive expression of GFP::miR-E-shHif1 in photoreceptors and

mCherry::miR-E-shHif2 in the RPE (Fig. 2D). Using a single vector to deliver the two shRNAs

to their specific cell type allowed for an efficient and precise intervention. A scrambled RNA

sequence replaced shHif1 and shHif2 in the shCtrl vector. To compare the potency of this

multiplex dual-acting AAV with single-acting AAVs, we generated two additional vectors to

deliver either shHif1 to the photoreceptors or shHif2 to the RPE (Fig. 2D).

Targeted expression in photoreceptors and RPE of wild type mice

The AAVs carrying shHif or shCtrl were injected into the subretinal space of wild type mice

(Fig. 3A) to evaluate transduction and expression profile of delivered constructs. Two weeks

after administration, the injection site was visible as a hyperreflective spot in the fundus (Fig.

3B,C; arrows). OCT scans taken at the injection site showed minor layer distortions likely

attributable to retinal injury caused by needle penetration. Beyond the injection site, however,

tissue integrity was preserved in shCtrl- or shHif-treated areas, as shown by OCT scans

through the optic nerve head, indicating that neither vector was toxic to retinal cells or the RPE

(Fig. 3B,C). Fluorescence funduscopy showed widespread GFP (as surrogate marker for

shHif1 expression) and mCherry (indicating shHif2 expression) positive areas surrounding the

injection site (Fig. 3B,C). Retinal cross-sections collected at 3 weeks post-injection (wpi)

showed GFP and mCherry signals localized to photoreceptors and the RPE, respectively (Fig.

3D,E). However, weak mCherry fluorescence was also detected in the neural retina close to

the injection site. The signal was much less robust than in the RPE and decreased to

undetectable levels with increasing distance from the injection site. Similarly, some GFP

fluorescence was observed in the RPE. While this may be attributed to phagocytosis of shed

GFP-positive outer segments by RPE cells, we cannot completely rule out expression of the

GRK1-driven gene in the RPE, as it has been observed for AAV8 in macaque before [38]. To

evaluate the transduction perimeter, GFP and mCherry fluorescence was examined on retinal

and RPE flatmounts at 3 wpi. The GFP-positive area covered approximately 40-50% of the

total retina and closely matched the mCherry-positive area in the corresponding RPE both in

size and position (Fig. 3F,G). Together with the reduced levels of *Hif1a* in photoreceptors and

Hif2a in the RPE in the transduced area (Fig. S3A,B), this demonstrates successful reduction

of the target mRNAs in the intended cell types. Potential inflammation was assessed by

immunofluorescence staining for allograft inflammatory factor 1 (IBA1, alias AIF1) on retina

and RPE flatmounts, and in retinal cross-sections at 3 wpi (Fig. 3F,G, Fig. S3C). Whereas

reactive (amoeboid) microglia and/or macrophages were detected in the retina close to the

injection site, ramified resting microglia populated the surrounding tissue. RPE flatmounts

presented few IBA1-positive cells only close to the injection site. The presence of microglia or

macrophages in this restricted area might have been caused by retinal detachment and scar

formation inflicted by the subretinal injections. Apart from the perturbations at the injection site,

the transduced areas showed normal staining patterns for cone arrestin (ARR3) and beta-

catenin (CTNB1) (Fig. S3D), and showed no signs of degeneration or functional impairment

up to 22 wpi (Fig. 3H,I; Fig. S3E). Taken together, these results demonstrate widespread and

selective transduction of photoreceptors and RPE cells by the AAV delivering the multiplex

RNA therapeutics with no apparent adverse effects beyond the injection site.

Mouse models for AMD

Our therapeutic strategy centers on targeting the alpha subunits of HIF1 and HIF2 transcription

factors, which are stabilized under reduced oxygen levels. Stabilization is initiated when the

activity of HIF-prolyl hydroxylases decreases due to reduced availability of O_2 as substrate.

Reduced hydroxylation of HIF1A and HIF2A prevents their recognition by the VHL protein

complex and subsequent degradation by proteasomes allowing them to accumulate and

function as transcription factors. This accumulation (and activation) of HIF1 and HIF2 is

mimicked in $Rod^{\Delta Vhl}$ mice, which lack VHL in rods. These mice represent a pseudohypoxia

model in which the hypoxic response is elicited in the presence of normal oxygen levels leading

to photoreceptor degeneration [8,9]. The stabilization of HIF1A by the chemical prolyl

hydroxylase inhibitor rodaxustat resulted in comparable cell death [39] supporting the validity of

the model. However, eyes of ageing individuals and those afflicted by AMD experience hypoxia

affecting both photoreceptors and the RPE. Thus, we established a second model by knocking

out *Vegfa* in RPE cells of adult mice ($RPE^{\Delta Vegfa}$; Fig. 4A). As reported by others, the lack of

VEGFA disrupts the homeostasis in the choriocapillaris leading to vasoconstriction in the choroid. This results in the development of true hypoxic conditions, including HIF activation, consequently causing RPE and photoreceptor degeneration [7,40,41]. To generate a local phenotype, we delivered Cre recombinase to the nasal RPE of 28-day-old *Vegfa*[fl/fl] mice [20] by the subretinal injection of an AAV4 expressing Cre-P2A-eGFP under control of the VMD2 promoter (Table 2). Inactivation of *Vegfa* expression in the transduced RPE was verified by *in situ* hybridization, and transduction was confirmed by staining for Cre in the consecutive section at 3 wpi. *Vegfa* expression was evident in the non-transduced area but absent in Cre-positive RPE cells in the transduced area (Fig. 4B). This mutually exclusive expression pattern indicated an effective suppression of *Vegfa* in transduced RPE cells. Choroidal flatmounts stained for podocalyxin (PODXL), an endothelial cell marker, demonstrated a strong reduction of choroidal vessel density at 3 wpi (Fig. 4C). To quantify the phenotype, we analyzed the staining pattern of PODXL in sections through control and transduced regions lacking *Vegfa* expression. Transduced areas showed notable gaps in the PODXL pattern and a significant increase in the gap size (Fig. 4C), suggesting a collapse of the choriocapillaris and rearrangement of endothelial cells in the absence of RPE-derived VEGFA.

At 3 wpi the transduced area was clearly visible as a brighter region in the fundus that correlated with the GFP-positive area and thus expression of Cre (Fig. 4D). Angiography indicated a predominantly regular retinal vasculature, and OCT showed normal retinal thickness, with the expected small perturbations around the injection site at this time (Fig. 4D, upper panel). At 39 wpi, however, the same eye presented various abnormalities including fundus discoloration, RPE hypertrophy, retinal thinning with an almost complete loss of the ONL and vascular leakage (Fig. 4D-F; Fig. S4). The weak GFP signal (Fig. 4D) suggested either a loss of transgene expression or the death of transduced cells at this time point. To quantify the effect *in vivo*, we employed OCT volume scans and measured the ONL thickness in two nasal subfields. Direct comparisons showed a significantly reduced ONL thickness in $RPE^{\Delta Vegfa}$ but not in non-injected (*Vegfa*[fl/fl]) or control-injected (*Vegfa*[fl/fl] *+PBS or +GFP)* mice

449 (Fig. 4E). Light microscopy and quantification of equatorial nasal-temporal sections supported

450 the *in vivo* findings and revealed significant thinning of the ONL and reduced length of PS that

451 was restricted to the transduced nasal retina of *RPE^{ΔVegfa}* mice (Fig. 4F,G). The phenotype was

452 not caused by subretinal injections or the activity of the VMD2 promotor *per se* since the

453 respective controls showed no signs of retinal degeneration (Fig. 4F,G).

Dual-acting shHif therapy restores tissue integrity in *Rod^{ΔVhl}* mice

Having validated the expression pattern of the shHif AAV vector in wt mice (Fig. 3), we

proceeded to investigate whether this vector can effectively interfere with the hypoxia pathway

in our two mouse models. We first used the *Rod^{ΔVhl}* mouse with its chronically activated

molecular response to hypoxia in rod photoreceptors due to the rod-specific deletion of *Vhl*

(Fig. 5A). The model is characterized by a HIF1-dependent, late onset progressive loss of

photoreceptors, occasional disturbances in the RPE, and a decline in visual function [8,9]. As

expected, untreated 26-week-old *Rod^{ΔVhl}* mice showed patches of photoreceptor cell loss in

the fundus, as well as RPE and photoreceptor irregularities in OCT images (Fig. 5B). Age-

matched *Rod^{ΔVhl}* mice injected with shCtrl vector displayed a comparable phenotype at 22 wpi,

with patches of degenerated areas in normal fundus that were prominently visible as dark

flecks in fundus fluorescence images capturing vector-derived GFP and mCherry signals. In

contrast, fundi and OCT images of shHif-treated mice displayed fewer irregularities and no

obvious signs of RPE disturbances (Fig. 5B). To quantify the extent of rescue *in vivo*, we

measured ONL thickness in the different groups using volumetric OCT scans. Although not

reaching the thickness of control *Vhl^{fl/fl}* mice (without Opsin-Cre), treatment of *Rod^{ΔVhl}* mice

with shHif resulted in a significantly thicker ONL compared to *Rod^{ΔVhl}* mice treated with shCtrl

(Fig. 5C). Light microscopy of retinal morphology at 22 wpi supported these findings and

showed that retinal areas that had received the shHif virus had less severe ONL thinning and

RPE disturbances than non- or shCtrl-transduced areas (Fig. 5D).

Gene expression analysis at 22 wpi showed reduced *Hif1a* but not *Hif2a* mRNA levels in the

retina, (Fig. 5E) and reduced *Hif2a* but not *Hif1a* mRNA expression in the eyecup tissue that

included the RPE (Fig. 5F). This suggests a high degree of cell-type specificity of our dual-

acting shHif gene silencing approach. The rather moderate reduction of the respective

expression levels of *Hif1a* and *Hif2a* by only 20-30% was not surprising because tissue

samples used for these experiments also contained cells from the transduced area that were

not targeted by our vector (e.g. 2nd and 3rd order neurons) and cells from non-transduced areas.

This may also be the reason why levels of *Casp1* (marker for degenerative processes) and

Pde6b (marker for rod integrity) in the retina and of *Rpe65* (marker for RPE integrity) in the

eyecup were only slightly reduced or increased, respectively (Fig. 5E). Together with the

significant rescue of retinal thickness our gene expression data collectively supports a

considerable potential of the shHif approach for alleviating the harmful effects of a chronic

hypoxic response in photoreceptors.

Dual-acting shHif therapy restores tissue integrity in *RPE$^{\Delta Vegfa}$* mice

We proceeded to assess the potential of the shHif approach to mitigate the degenerative

effects of true chronic hypoxia in the *RPE$^{\Delta Vegfa}$* model. Since this model also includes the

pathogenic activation of HIF2A in the RPE [7], we compared the efficacy of the dual-acting shHif

AAV with single-acting shHif1 (targeting photoreceptors) and shHif2a (targeting the RPE)

AAVs. Subretinal co-injections of AAV::VMD2-Cre-P2A-eGFP to inactivate *Vegfa* in the RPE,

along with either shCtrl, shHif, shHif1 or shHif2 were performed at 4 weeks of age followed by

analysis at 39 wpi. Funduscopy and OCT of *RPE$^{\Delta Vegfa}$* mice injected with shCtrl showed

discolored patches and affected tissue layers, indicating RPE disturbances and degeneration

at 39 wpi (Fig. 6A), a phenotype similar to untreated *RPE$^{\Delta Vegfa}$* mice (Fig 4D). In contrast, shHif-

treated animals displayed only minor disturbances around the injections site, but no obvious

retinal degeneration (Fig. 6A). The rescue effect by the treatment with the various viruses was

quantified *in vivo* using OCT volume scans. Treatment with the dual-acting shHif virus

preserved ONL thickness in *RPE$^{\Delta Vegfa}$* mice significantly better than treatment with the single-

acting or control AAVs. (Fig. 6B). The superior rescue effect by shHif was further verified by

light microscopy of retinal sections (Fig. 6C) and measurements of ONL thickness and PS

length (Fig. 6D) in a subset of mice. Eyes treated with shCtrl displayed a drastic thinning of the

ONL, loss of PS, and variable thickness of the RPE in transduced areas. In contrast, mice

treated with shHif had preserved PS with few disturbances, a thicker ONL, and only sporadic

regions with irregular RPE morphology. shHif1 and shHif2 viruses rescued the phenotype at

intermediate levels (Fig. 6C,D). Areas that were not transduced by the Cre-expressing virus

and thus with regular *Vegfa* expression in the RPE showed normal retinal morphologies for all

injected eyes. The superior rescue effect by the dual-acting shHif virus underscores the

importance of simultaneously targeting Hif1 in photoreceptors and Hif2 in the RPE to efficiently

mitigate the consequences of chronic hypoxia. The dual-targeting approach was not only

superior to targeting individual HIF transcription factors but was also long lasting and protected

tissue integrity at least up to 61 wpi (Fig. 7). While *RPE$^{\Delta Vegfa}$* mice treated with shCtrl exhibited

significant loss of GFP and mCherry fluorescence at 61 wpi when compared to 4 wpi, shHif

treated mice still showed strong fluorescence at this time point suggesting preservation of

photoreceptors and RPE (Fig. 7A,B). This was confirmed by light microscopy of retinal sections

and quantification of ONL thickness and PS length which showed strong preservation of retinal

morphology in shHif but not shCtrl treated mice (Fig 7C,D). The eyes injected with the GFP

control virus provided a baseline for non-degenerated retinas allowing to assess the overall

efficacy of the dual-acting shHif virus. Our results demonstrate the efficacy of our multiplex

shHif therapy in preserving retinal structures for more than one year despite the continued

presence of the toxic hypoxic stimulus. The strong protection in two different models of

hypoxia-induced retinal degeneration indicates the potential of the approach for a future

translation into a clinical setting.

Discussion

Age is the major non-genetic risk factor for the development of AMD. The aging ocular tissue undergoes several changes that result in reduced oxygen levels in the outer retina and increased expression of HIF target genes [2,3,8]. Some HIF target genes, such as *VEGFA* and *ANGPTL4*, are angiogenic factors that ultimately contribute to the development of neovascular AMD [29,32]. Others, such as *SERPINE1*, may facilitate disease development by remodeling the extracellular matrix [31] or through their connection to senescence and ageing [42]. Notably, the expression of all three aforementioned hypoxia-related and AMD-associated genes was increased in old donor eyes.

An integral part of the development and utilization of this therapy is that the nature of the retinal disease must be considered when targeting the hypoxia pathway as a gene-agnostic therapeutic approach, since a hypoxic environment triggers an intracellular response that adjusts major metabolic pathways, including glycolysis, oxidative phosphorylation, lipid metabolism, and others, to the new environmental conditions. This metabolic shift may reduce the susceptibility of photoreceptors to mutation-induced degeneration [43], but seems to contribute to cell death in otherwise normal photoreceptors and cells of the RPE. This divergent reaction of mutant and normal cells to an activated hypoxic response highlights the complex interplay between cell metabolism, aging, hypoxia, and other pathways involved in inherited retinal degeneration or AMD development [7,8,43-46]. In the case of AMD, reducing the HIF pathway may be favorable and benefit a wide range of patients affected by either the wet or dry form of the disease.

Here we established an efficient RNA interference strategy that simultaneously targets HIF1 in the photoreceptors and HIF2 in the RPE. This multiplex approach prevented retinal degeneration connected to a chronic hypoxic response in two mouse models of pseudo- or true hypoxia. Another key finding was the demonstration that this dual-acting therapy proved

to be more effective than single-target treatments. This finding highlights the potential and

superiority of multi-target therapies for complex multifactorial diseases such as AMD.

The use of shRNAs driven by Pol II promoters to downregulate target genes has several

advantages. (i) It allows targeting only those cell types that require treatment without affecting

other cells. (ii) It does not completely inactivate the target genes allowing their activity for

essential or housekeeping tasks, in contrast to knockout strategies. (iii) It ensures long-lasting

effects through continuous expression of the therapeutic RNAs as opposed to the transient

effects of short-lived synthetic siRNAs. (iv) It minimizes the possibility that a bolus injection of

siRNAs will overwhelm the endogenous RNAi pathway leading to toxic side effects [16,47].

Because our multiplex system is based on a single AAV vector, it has the additional advantage

that it does not rely on double transduction of individual cells, trans-splicing, or other

mechanisms that may reduce efficiency. The therapeutic approach is not only well tolerated,

but also potent and has protected tissue integrity for up to 61 weeks in two different models of

hypoxia-related retinal degeneration that share key aspects of AMD.

In contrast to our data for the chronic models presented here, the recent work of Babapoor-

Farrokhran and colleagues suggests a beneficial function of HIF1 for the survival of wild type

photoreceptors [48]. However, this conclusion was drawn after the induction of acute oxidative

stress by $NaIO_3$, which may less accurately model the effects of a long-lasting chronic hypoxic

response or environment. In addition, the chemicals used in their study can be toxic to

photoreceptors *per se* (digoxin; [49]) or exhibit side effects apart from HIF activation (DMOG;

[50,51]). Nevertheless, data from their acute model is interesting and may best be correlated with

the protective effects of hypoxic preconditioning against acute light-induced retinal

degeneration [52]. Since this indicates that acute and chronic hypoxic responses have distinct

consequences for retinal cells, we believe that our two genetic models for chronic hypoxia

represent the situation in the aged eye more accurately than drug-induced acute models.

While the $Rod^{\Delta Vhl}$ model has chronically active HIFs in normoxic rods causing HIF1-dependent

photoreceptor degeneration [8], the RPE-specific deletion of *Vegfa* in the $RPE^{\Delta Vegfa}$ model affects

the choroidal vasculature, causing true tissue hypoxia and leading to HIF2-dependent RPE

damage and to retinal degeneration [7]. Since the true tissue hypoxia in $RPE^{\Delta Vegfa}$ mice affects

multiple cell types and elicits broad effects including changes in metabolism [7], $RPE^{\Delta Vegfa}$ mice

may represent the complex, multifactorial environment in the aging eye with some accuracy.

To knockdown *Vegfa* in adult RPE cells, we delivered a VMD2-controlled Cre transgene to

Vegfa$^{fl/fl}$ mice. This approach is more specific, efficient and reproducible compared to methods

suffering from variable and patchy Cre expression in RPE of transgenic mice [41,53] or the lack

of cell type specificity after delivery of CjCas9 for indel generation [40].

The therapeutic vector was designed to express shHif1 in photoreceptors and shHif2 in the

RPE with GFP and mCherry as surrogate markers. Despite that the expression of GFP::shHif1

was controlled by the GRK1 promotor, GFP fluorescence was also visible in the RPE (Fig.

3D,E). This was likely due to phagocytosis of photoreceptor material by the RPE, or to some

expression of the gene in the RPE following delivery by AAV8, which has a known tropism for

the RPE, as observed in macaque studies [38]. This opens the possibility that some molecules

targeting *Hif1a* mRNA reached the RPE. Although phagocytosed material would likely be

confined to lysosomes, we cannot exclude some downregulation of *Hif1a* transcripts in the

RPE by such 'spills'. However, since the knockout of *Hif1a* in the mouse RPE does not cause

phenotypic changes [41] this possibility does not significantly influence the validity, specificity

and efficacy of our approach.

Although our dual-active RNA therapy rescued the phenotype in both mouse models, our

results suggest a higher efficiency in $RPE^{\Delta Vegfa}$ mice where degeneration is based on true

hypoxia due to pathological changes in the choroidal vasculature (Fig. 4C). The RNA-induced

silencing complex (RISC) that is formed after shRNA expression reduces stability or translation

of mRNA transcripts but does not affect preformed proteins. Thus, the efficacy of shRNA-

mediated downregulation of protein activity depends on the expression levels of the target

RNA. Considering reports suggesting that hypoxia can reduce levels and half-lives of *Hif1a*

and *Hif2a* transcripts [54,55], it appears possible that RNA interference is more effective in the

model of true hypoxia due to lower levels of target mRNAs. Clearly, unraveling the mechanisms

that contribute to the differences in treatment efficacies in the two models needs further

research and is important to further improve the RNA interference approach. A limitation of our

study lies in structural differences between the mouse and the human retina. Unlike humans,

mice do not possess a macula which prevents our models from presenting all aspects of AMD.

However, since the molecular response to hypoxia with the activation of HIF transcription

factors is quite universal, we nonetheless believe that our RNA therapy targeting this

conserved mechanism is translatable in the future to the human situation.

614   Conclusion

Since a reduction of oxygen delivery to the outer retina through the choroidal system affects

both RPE and photoreceptors, it is important that a therapeutic intervention targets the

degeneration-causing factors in both cell types simultaneously and specifically. We

approached these requirements through the dual-active multiplex RNAi therapy system using

a single AAV. Delivery of therapeutic genes by AAV is safe and has been evaluated in several

clinical trials [56,57]. The first FDA and EMA-approved gene therapy (Luxturna©) for Leber

congenital amaurosis type 2 not only supports the safety but also the efficacy of the delivery

of genetic material to target cells by subretinal AAV injections. The approach presented here

further expands the potential of AAVs as a therapeutic system by using a single AAV vector

with two specific Pol-II promotors to drive expression of shRNAs targeting two transcripts in

two distinct cell types. We believe that the system demonstrates potential for widespread

applicability to other multifactorial diseases that necessitate modifications of gene expression

across various cell types or tissues. Since RNAi-based therapeutics, including siRNA and

shRNA, have already been used in several clinical trials, e.g., to target cancer or genetic

disorders [58,59], the system developed here has the potential to be translated for the use in

patients, particularly for the treatment of retinal disorders associated with a chronic hypoxic

response such as AMD.

List of abbreviations

AAV: adeno-associated virus

AMD: age-related macular degeneration

ARR3:   cone arrestin

BM: Bruch's membrane

CC: choriocapillaris

CTNB1:   beta-catenin

Ctrl: control

DAPI: 4',6-diamidino-2-phenylindole

ERG: electroretinogram

Fl: flox

GCL: ganglion cell layer

GFP: green fluorescent protein

GRK1:   G-protein coupled kinase 1

HIF: hypoxia-inducible factor

IBA1:   allograft inflammatory factor 1

INL: inner nuclear layer

miR: microRNA

OCT: optical coherence tomography

ONL: outer nuclear layer

PFA: paraformaldehyde

PODXL: podocalyxin

PS: photoreceptor segment

VEGFA: vascular endothelial growth factor A

RPE: retinal pigment epithelium

siRNA: small interfering RNA

shRNA: small hairpin RNA

VHL: von Hippel Lindau

VMD2:   vitelliform macular dystrophy 2

WPI: weeks post injection

WT: wild type

3'UTR: 3' untranslated region

Declarations

Ethics approval and consent to participate

Analysis of human ocular tissue was approved by the Cantonal Ethic Commission (license numbers BASEC-Nr: PB_2017-00550 and 2020-01856) and adhered to the tenets of the Declaration of Helsinki. Animal experimentation was approved by the Cantonal Veterinary Office of Zürich, Switzerland (license numbers: ZH019/2019 and ZH105/2022) and adhered to the ARVO statement for the use of Animals in Ophthalmic and Vision Research.

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during this study are included in this article and its supplementary information files.

Competing interest

The authors declare that they have no competing interests.

Funding

This work was supported by Swiss National Science Foundation grants 31003A_173008 and
310030_200798, and a research grant of the University of Zürich to L.E. (FK-20-028).

Authors' contributions

L.E. M.S. and C.G. conceived the study. L.E. designed and performed experiments and the
analysis. I.M and F.B. collected human ocular tissue. M.S.; C.I.; D.K.; F.P; E.K.; A.F.; D.F.;
L.M.; G.W.; I.T., L.G., SL.B.; SE.B. contributed to data collection and performed experiments.
All authors contributed to project design, organization of the experiments, interpretation of
results. L.E., M.S. and C.G wrote the manuscript. All authors read and approved the final
manuscript.

Acknowledgements

The Viral Vector Facility (VVF) of the Neuroscience Center Zurich (ZNZ) produced all AAVs
and the Laboratory Animal Services Center (LASC) of the University of Zurich was responsible
for animal care.

Figure legends

Figure 1 | Gene expression in the human retina and RPE. Expression of *VEGFA,*
ANGPTL4, SERPINE1, GNAT2, and *RPE65* was determined by real-time PCR in the retina
(A) and RPE **(B)** of young (Y) and old (O) human donor eyes without diagnosed ocular
pathology. Shown are means (line) and individual data points on a log10 scale. Outliers were
identified and excluded using Grubbs method, and significance was determined by unpaired t-
tests. N = 15 (young) and 14 (old).

Figure 2 | *In vitro* validation of *Hif1a* and *Hif2a* shRNAs and design of the dual-acting AAV vector. (A) Graphical representation of tissue changes leading to hypoxia and degenerative processes during aging and AMD. **(B)** Gene expression analysis of 661W cells stably expressing shCtrl, shHif1 or shHif2 after normoxic or hypoxic (24 h, 0.2% O_2) exposure. Values were normalized to beta-actin (*Actb*) and expressed relatively to normoxic 661W-shCtrl (set to 1). Shown are individual values (n=3) and means ± SD. Significance was tested using a one-way ANOVA with Tukey's multiple comparison test. *: statistically significant; p-values of all comparisons are given in Table S2. **(C)** Western blot analysis for HIF1A, HIF2A and ACTB in 661W cells stably expressing shCtrl, shHif1 or shHif2 after normoxic and hypoxic exposure. The position of the band for HIF1A is indicated by an arrow. **(D)** Schematic representation of the DNA constructs packaged into AAV vectors. AAV-shHif: the construct consists of the GRK1 promotor driving photoreceptor-specific expression of shHif1a embedded in the miR-E scaffold in the 3'UTR of GFP, followed by the VMD2 promotor controlling RPE-specific expression of shHif2a embedded in the miR-E scaffold in the 3'UTR of mCherry. AAV-shCtrl: a scrambled non-targeting control shRNA sequence replaced shHif1a and shHif2a. AAV-shHif1: the GRK1 promotor drives photoreceptor-specific expression of shHif1a embedded in the miR-E scaffold in the 3'UTR of GFP. AAV-shHif2: the VMD2 promotor controls expression of shHif2a embedded in the miR-E scaffold in the 3'UTR of mCherry in the RPE. Sequences of shRNAs are provided in Fig. S1 and supplemental data SD1. CC: choriocapillaris, BM: Bruch`s membrane, RPE: retinal pigment epithelium, PS: photoreceptor segments, ONL: outer nuclear layer.

Figure 3 | Evaluation of the pattern and tolerability of shHif expression in wt mice after AAV-mediated delivery. (A) Illustration of transvitreal AAV-delivery to the subretinal space targeting photoreceptor cells (shown in green due to expression of GFP) and RPE cells (shown in red due to expression of mCherry). **(B, C)** Funduscopy, OCT scans and fundus fluorescence of wt mice injected with shCtrl (B) or shHif (C) at 2 wpi. Orange and red lines in the fundus

images indicate positions of the respective OCT scans. Fluorescence indicates transduced

areas in the retina (green, GFP) and RPE (red, mCherry). White arrows: injection site, *: optic

nerve head. **(D, E)** Retinal cross-sections of wt mice injected with shCtrl (D) or shHif (E) at 3

wpi. Green/white: GFP-positive photoreceptors / RPE. Red/white: mCherry-positive RPE cells.

Shown are merged images (left) and individual fluorescence channels (middle and right). **(F,**

G) Retina and their corresponding RPE flatmounts from wt mice injected with shCtrl (F) or

shHif (G) at 3 wpi. Shown are native GFP (green) and mCherry (red) fluorescence and

stainings for IBA1 (white). Boxed areas in the fluorescence images are shown in higher

magnification (middle). Arrow: injection site. **(H)** Representative retinal morphologies of wt

mice injected with shCtrl (left) or shHif (right) at 22 wpi. **(I)** Scotopic and photopic

electroretinograms (ERG) of wt mice injected with shCtrl (grey) or shHif (orange) at 22 wpi.

Shown are scotopic a- and b-wave and photopic b-wave amplitudes as a function of stimulus

intensity. Shaded areas indicate SD, n=4 per group. Scale bars: as indicated. BM: Bruch's

membrane, RPE: retinal pigment epithelium, PS: photoreceptor segments, ONL: outer nuclear

layer, INL: inner nuclear layer, GCL: ganglion cell layer, SRS: subretinal space.

Figure 4 | Assessment of the $RPE^{\Delta Vegfa}$ phenotype. (A) $RPE^{\Delta Vegfa}$ model schematics: *Vegfa*

deletion in the RPE leads to reduced choroidal vessel density, RPE alterations and

photoreceptor degeneration. **(B)** *Vegfa* (red) *in situ* hybridization and Cre (magenta)

immunofluorescence in areas that were or were not transduced with AAV::VMD2-Cre-P2A-

eGFP. **(C)** PODXL immunofluorescence in endothelial cells of choroidal flatmounts (top) and

cross-sections (lower panels) of indicated areas. Contrast-enhanced threshold images allowed

pixel intensity scan analysis (bottom) and quantification (right) in $RPE^{\Delta Vegfa}$ mice at 3 wpi. Data

from n=3 eyes, 18-19 measurements/area, analyzed by unpaired t-test. **(D)** Funduscopy, OCT

scans, fundus fluorescence and angiography of $RPE^{\Delta Vegfa}$ mice at 3 (top) and 39 (bottom) wpi.

Fundus images with orange/red lines indicate OCT scan positions. GFP-labelled panels show

green fluorescence confirming the presence of Cre at 3 wpi; background fluorescence

dominates at 39 wpi. Arrows: injection site, *: optic nerve head. **(E)** Representative color-coded

ONL thickness maps generated from OCT volume scans at 39 wpi. Right panel: boxplots of

ONL thickness in the two nasal subfields in mice. Shown are average values of individual eyes

(dots) and means of all eyes (line). Note: values for control injected mice (*Vegfa^{fl/fl}* + PBS and

Vegfa^{fl/fl} + GFP) were combined. n=4 (*Vegfa^{fl/fl}*); n=11 (*RPE^{ΔVegfa}*); n=6 (*Vegfa^{fl/fl}* + GFP /

Vegfa^{fl/fl} + PBS). One-way ANOVA with Tukey's multiple comparison test. P-values as shown.

(F) Representative retinal histology of *Vegfa^{fl/fl}* mice (left) subretinally injected with AAVs

expressing Cre (*RPE^{ΔVegfa}*) or GFP (*Vegfa^{fl/fl}* + GFP), or with PBS (*Vegfa^{fl/fl}* + PBS) at 39 wpi.

(G) ONL thickness and PS length quantified from histological sections 39 wpi. Shown are

spidergrams and average values in the transduced nasal area (boxplots, 400-2000 μm from

the optic nerve head (ON)) of indicated groups at 39 wpi. Shown are means ± SD and individual

data points. n=3 (*Vegfa^{fl/fl}*; *Vegfa^{fl/fl}* + GFP; *Vegfa^{fl/fl}* + PBS); n=2 (*RPE^{ΔVegfa}*). One-way ANOVA

with Tukey's multiple comparison test, p-values as indicated. CC: choriocapillaris, BM: Bruch's

membrane, RPE: retinal pigment epithelium, PS: photoreceptor segments, ONL: outer nuclear

layer, INL: inner nuclear layer, GCL: ganglion cell layer. DIC: differential interference contrast.

Figure 5 | Rescue of the *Rod^{ΔVhl}* phenotype by the dual-acting shHif RNA therapy. (A)

Schematic representation of *Rod^{ΔVhl}* mice and rescue of RPE (red) and photoreceptors (green)

through downregulation of *Hif2a* in the RPE and *Hif1a* in photoreceptors and *Hif2a* in the RPE

by shHif. **(B)** Funduscopy, OCT scans and fundus fluorescence assessed at 26 weeks of age

(22 wpi) in *Rod^{ΔVhl}* mice that were either non-injected (left) or treated with shCtrl (middle) or

shHif (right). Orange and red lines in the fundus images indicate positions of the respective

OCT scans. OCT scans of higher magnifications are included. Fundus fluorescence images

(last two rows) show the transduced areas in the retina (green, GFP) and RPE (red, mCherry).

*: optic nerve head. **(C)** Color-coded ONL thickness maps of *Vhl^{fl/fl}*, *Rod^{ΔVhl}* + shCtrl and *Rod^{ΔVhl}*

+ shHif mice generated from OCT volume scans at 22 wpi. Grids were aligned to the optic

nerve (ON). The ONL thickness was measured in seven subfields and indicated with numbers

(artificial units). Right panel: quantification of ONL thickness in the seven subfields (outlined

by the orange line in the grid). Shown are average values of the subfields in individual eyes

(dots) and means of all eyes (line) in box blots (n≥9). One-way ANOVA with Tukey's multiple

comparison test. P-values as shown. **(D)** Representative retinal morphologies of transduced

regions of *Vhl^{fl/fl}* control and *Rod^{ΔVhl}* mice injected with shCtrl or shHif. All mice were 26 weeks

of age (22 wpi) at time of analysis. **(E, F)** Relative gene expression levels in retinas (E) or

eyecups (F) of *Rod^{ΔVhl}* + shCtrl (grey) and *Rod^{ΔVhl}* + shHif (orange) mice at 22 wpi. Expression

was normalized to beta-actin (*Actb*) and shown relative to expression in *Rod^{ΔVhl}* + shCtrl mice

(set to 1). Shown are means ± SD of n = 11-12. Unpaired t-test. P-values as indicated. RPE:

retinal pigment epithelium, ONL: outer nuclear layer, INL: inner nuclear layer, GCL: ganglion

cell layer.

Figure 6 | Dual-acting multiplex shRNA therapy is more effective than single acting

shRNA vectors in rescuing pathology in *RPE^{ΔVegfa}* mice.

(A) Funduscopy, OCT scans and fundus fluorescence of *RPE^{ΔVegfa}* mice injected with shCtrl

(left) or shHif (right) at 39 wpi. Orange and red lines in the fundus images indicate positions of

the respective OCT scans that are shown in lower and higher magnifications. Fundus

fluorescence indicates transduced areas in the retina (green, GFP) and RPE (red, mCherry).

*: optic nerve head. **(B)** Color-coded ONL thickness maps generated from OCT volume scans

of *RPE^{ΔVegfa}* + shCtrl, *RPE^{ΔVegfa}* + shHif, *RPE^{ΔVegfa}* + shHif1 and *RPE^{ΔVegfa}* + shHif2 mice at 39

wpi. Grids were aligned to the optic nerve (ON). The ONL thickness was measured in the two

nasal subfields (outlined by the orange line in the grid) that were most affected in *RPE^{ΔVegfa}*

mice and indicated with numbers (artificial units). Right panel: quantification of ONL thickness

in the nasal two subfields. Shown are average values in individual eyes (dots) and means of

all eyes (line) in box blots. n=11 (shCtrl, shHif); n=9 (shHif1); n=8 (shHif2). One-way ANOVA

with Dunnett's multiple comparison test comparing all groups to shHif. p-values as shown. **(C)**

Representative retinal morphologies of transduced and non-transduced areas in *RPE^{ΔVegfa}*

mice treated with AAVs as indicated. Analysis was at 39 wpi. **(D)** Quantification of ONL

thickness and PS length of the four treatment groups. Shown are spidergrams and average

values of the transduced nasal area (400-2000 µm from the optic nerve head (ON)) of a subset

of mice that were already analyzed by volume scans (B) at 39 wpi. Grey: shCtrl; orange: shHif;

green shHif1; blue: shHif2. Shown are means ± SD and individual data points. One-way

ANOVA with Dunnett's multiple comparison test comparing all groups to shHif. P-values as

shown. shHif: n=4 (ONL) and n=3 (PS); shHif1: n=9 (ONL) and n=4 (PS); shHif2: n=8 (ONL)

and n=6 (PS); shCtrl: n=2 (ONL and PS). RPE: retinal pigment epithelium, PS: photoreceptor

segments, ONL: outer nuclear layer, INL: inner nuclear layer, GCL: ganglion cell layer, BM:

Bruch's membrane.

Figure 7 | Long-lasting rescue of the $RPE^{\Delta Vegfa}$ phenotype by the dual-acting shRNA

therapy

(A, B) Funduscopy and fundus fluorescence of $RPE^{\Delta Vegfa}$ mice injected with shCtrl (top) or

shHif (bottom) analyzed at 4 wpi (A) and 61 wpi (B). Fluorescence indicates transduced areas

in the retina (green, GFP) and RPE (red, mCherry). *: optic nerve head. **(C)** Representative

retinal morphologies of transduced and non-transduced areas of $RPE^{\Delta Vegfa}$ mice injected with

shCtrl or shHif, analyzed at 61 wpi. **(D)** Quantification of ONL thickness and PS length at 61

wpi, presented as spidergrams (left) and boxplots (right) presenting the averaged values of the

transduced nasal area (400-2000 µm from the optic nerve head (ON)). Treatment groups:

$Vegfa^{flox/flox}$ mice injected with GFP control virus (green), $RPE^{\Delta Vegfa}$ + shHif mice (orange) and

$RPE^{\Delta Vegfa}$ + shCtrl mice (grey) at 61 wpi. Note: ONL thickness and PS length in $Vegfa^{flox/flox}$ +

GFP mice are not shown in spidergrams to reduce complexity of the figure panels. Shown are

means ± SD and individual data points. One-way ANOVA with Tukey's multiple comparison

tests. P-values as shown. shHif: n=11 (ONL) and n= 9 (PS); shCtrl: n=10 (ONL) and n=6 (PS);

GFP: n=3 (ONL and PS).

References

1. Zhu, Y., Zhu, L., Wang, X. & Jin, H. RNA-based therapeutics: an overview and prospectus. *Cell Death Dis* **13**, 644 (2022).

2. Grunwald, J. E., Metelitsina, T. I., Dupont, J. C., Ying, G. S. & Maguire, M. G. Reduced foveolar choroidal blood flow in eyes with increasing AMD severity. *Invest Ophthalmol Vis Sci* **46**, 1033–1038 (2005).

3. Berenberg, T. L. *et al.* The association between drusen extent and foveolar choroidal blood flow in age-related macular degeneration. *Retina* **32**, 25–31 (2012).

4. Ramrattan, R. S. *et al.* Morphometric analysis of Bruch's membrane, the choriocapillaris, and the choroid in aging. *Invest Ophthalmol Vis Sci* **35**, 2857–2864 (1994).

5. Wakatsuki, Y., Shinojima, A., Kawamura, A. & Yuzawa, M. Correlation of Aging and Segmental Choroidal Thickness Measurement using Swept Source Optical Coherence Tomography in Healthy Eyes. *PLoS One* **10**, e0144156 (2015).

6. Mullins, R. F., Johnson, M. N., Faidley, E. A., Skeie, J. M. & Huang, J. Choriocapillaris vascular dropout related to density of drusen in human eyes with early age-related macular degeneration. *Invest Ophthalmol Vis Sci* **52**, 1606–1612 (2011).

7. Kurihara, T. *et al.* Hypoxia-induced metabolic stress in retinal pigment epithelial cells is sufficient to induce photoreceptor degeneration. *Elife* **5**, (2016).

8. Barben, M. *et al.* Hif1a inactivation rescues photoreceptor degeneration induced by a chronic hypoxia-like stress. *Cell Death Differ* **25**, 2071–2085 (2018).

9. Lange, C. *et al.* Retina-specific activation of a sustained hypoxia-like response leads to severe retinal degeneration and loss of vision. *Neurobiol Dis* **41**, 119–130 (2011).

10. Khan, A. A. *et al.* Transfection of small RNAs globally perturbs gene regulation by endogenous microRNAs. *Nat Biotechnol* **27**, 549–555 (2009).

11. Martin, J. N. *et al.* Lethal toxicity caused by expression of shRNA in the mouse striatum: implications for therapeutic design. *Gene Ther* **18**, 666–673 (2011).

12. Snøve, O. & Rossi, J. J. Toxicity in mice expressing short hairpin RNAs gives new insight into RNAi. *Genome Biol* **7**, 231 (2006).

13. Lee, Y. *et al.* The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**, 415–419 (2003).

14. Silva, J. M. *et al.* Second-generation shRNA libraries covering the mouse and human genomes. *Nat Genet* **37**, 1281–1288 (2005).

15.   Bauer, M. *et al.* Prevention of interferon-stimulated gene expression using microRNA-designed
hairpins. *Gene Ther* **16**, 142–147 (2009).

16.   Gu, S. *et al.* The loop position of shRNAs and pre-miRNAs is critical for the accuracy of dicer
processing in vivo. *Cell* **151**, 900–911 (2012).

17.   Fellmann, C. *et al.* An optimized microRNA backbone for effective single-copy RNAi. *Cell Rep* **5**,
1704–1713 (2013).

18.   Haase, V. H., Glickman, J. N., Socolovsky, M. & Jaenisch, R. Vascular tumors in livers with
targeted inactivation of the von Hippel-Lindau tumor suppressor. *Proc Natl Acad Sci U S A* **98**,
1583–1588 (2001).

19.   Le, Y. Z. *et al.* Mouse opsin promoter-directed Cre recombinase expression in transgenic mice.
Mol Vis **12**, 389–398 (2006).

20.   Gerber, H. P. *et al.* VEGF is required for growth and survival in neonatal mice. *Development* **126**,
1149–1159 (1999).

21.   Kay, C. N. *et al.* Targeting photoreceptors via intravitreal delivery using novel, capsid-mutated
AAV vectors. *PLoS One* **8**, e62097 (2013).

22.   Storti, F. *et al.* Impaired ABCA1/ABCG1-mediated lipid efflux in the mouse retinal pigment
epithelium (RPE) leads to retinal degeneration. *Elife* **8**, (2019).

23.   Gasparini, S. J. *et al.* Transplanted human cones incorporate into the retina and function in a
murine cone degeneration model. *J Clin Invest* **132**, (2022).

24.   Geiger, P., Barben, M., Grimm, C. & Samardzija, M. Blue light-induced retinal lesions, intraretinal
vascular leakage and edema formation in the all-cone mouse retina. *Cell Death Dis* **6**, e1985
(2015).

25.   Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402–408 (2001).

26.   Ebner, L. J. A. *et al.* Transcriptomic analysis of the mouse retina after acute and chronic
normobaric and hypobaric hypoxia. *Sci Rep* **11**, 16666 (2021).

27.   Tan, E. *et al.* Expression of cone-photoreceptor-specific antigens in a cell line derived from retinal
tumors in transgenic mice. *Invest Ophthalmol Vis Sci* **45**, 764–768 (2004).

28.   Meade, E. S., Ma, Y. Y. & Guller, S. Role of hypoxia-inducible transcription factors 1alpha and
2alpha in the regulation of plasminogen activator inhibitor-1 expression in a human trophoblast
cell line. *Placenta* **28**, 1012–1019 (2007).

29.   Qin, Y. *et al.* ANGPTL4 influences the therapeutic response of patients with neovascular age-
related macular degeneration by promoting choroidal neovascularization. *JCI Insight* **7**, (2022).

30.   Semenza, G. L. Targeting HIF-1 for cancer therapy. *Nat Rev Cancer* **3**, 721–732 (2003).

31.   Biasella, F. *et al.* Vitronectin and Its Interaction with PAI-1 Suggests a Functional Link to Vascular
912   Changes in AMD Pathobiology. *Cells* **11**, (2022).

913   32.   Witmer, A. N., Vrensen, G. F., Van Noorden, C. J. & Schlingemann, R. O. Vascular endothelial
914   growth factors and angiogenesis in eye disease. *Prog Retin Eye Res* **22**, 1–29 (2003).

915   33.   Ronning, K. E. *et al.* Loss of cone function without degeneration in a novel Gnat2 knock-out
916   mouse. *Exp Eye Res* **171**, 111–118 (2018).

917   34.   Redmond, T. M. *et al.* Rpe65 is necessary for production of 11-cis-vitamin A in the retinal visual
918   cycle. *Nat Genet* **20**, 344–351 (1998).

919   35.   Chhipa, R. R. *et al.* AMP kinase promotes glioblastoma bioenergetics and tumour growth. *Nat*
920   *Cell Biol* **20**, 823–835 (2018).

921   36.   Khani, S. C. *et al.* AAV-mediated expression targeting of rod and cone photoreceptors with a
922   human rhodopsin kinase promoter. *Invest Ophthalmol Vis Sci* **48**, 3954–3961 (2007).

923   37.   Esumi, N., Oshima, Y., Li, Y., Campochiaro, P. A. & Zack, D. J. Analysis of the VMD2 promoter
924   and implication of E-box binding factors in its regulation. *J Biol Chem* **279**, 19064–19073 (2004).

925   38.   Crosson, S. M. *et al.* Effects of Altering HSPG Binding and Capsid Hydrophilicity on Retinal
926   Transduction by AAV. *J Virol* **95**, (2021).

927   39.   Nsiah, N. Y., Morgan, A. B., Donkor, N. & Inman, D. M. Long-term HIF-1α stabilization reduces
928   respiration, promotes mitophagy, and results in retinal cell death. *Sci Rep* **13**, 20541 (2023).

929   40.   Kim, E. *et al.* In vivo genome editing with a small Cas9 orthologue derived from Campylobacter
930   jejuni. *Nat Commun* **8**, 14500 (2017).

931   41.   Kurihara, T., Westenskow, P. D., Bravo, S., Aguilar, E. & Friedlander, M. Targeted deletion of
932   Vegfa in adult mice induces vision loss. *J Clin Invest* **122**, 4213–4217 (2012).

933   42.   Eren, M., Boe, A. E., Klyachko, E. A. & Vaughan, D. E. Role of plasminogen activator inhibitor-1
934   in senescence and aging. *Semin Thromb Hemost* **40**, 645–651 (2014).

935   43.   Nolan, N. D. *et al.* CRISPR editing of anti-anemia drug target rescues independent preclinical
936   models of retinitis pigmentosa. *Cell Rep Med* **5**, 101459 (2024).

937   44.   Todorova, V. *et al.* Deficits in mitochondrial TCA cycle and OXPHOS precede rod photoreceptor
938   degeneration during chronic HIF activation. *Mol Neurodegener* **18**, 15 (2023).

939   45.   McHugh, K. J. *et al.* Computational modeling of retinal hypoxia and photoreceptor degeneration
940   in patients with age-related macular degeneration. *PLoS One* **14**, e0216215 (2019).

46.   Stefánsson, E., Geirsdóttir, A. & Sigurdsson, H. Metabolic physiology in age related macular
degeneration. *Prog Retin Eye Res* **30**, 72–80 (2011).

47.   Grimm, D. *et al.* Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA
pathways. *Nature* **441**, 537–541 (2006).

48.   Babapoor-Farrokhran, S. *et al.* Pathologic vs. protective roles of hypoxia-inducible factor 1 in RPE
and photoreceptors in wet vs. dry age-related macular degeneration. *Proc Natl Acad Sci U S A*
120, e2302845120 (2023).

49.   Landfried, B. *et al.* Digoxin-induced retinal degeneration depends on rhodopsin. *Cell Death Dis* **8**,
e2670 (2017).

50.   Chen, R., Ahmed, M. A. & Forsyth, N. R. Dimethyloxalylglycine (DMOG), a Hypoxia Mimetic
Agent, Does Not Replicate a Rat Pheochromocytoma (PC12) Cell Biological Response to
Reduced Oxygen Culture. *Biomolecules* **12**, (2022).

51.   Zhdanov, A. V., Okkelman, I. A., Collins, F. W., Melgar, S. & Papkovsky, D. B. A novel effect of
DMOG on cell metabolism: direct inhibition of mitochondrial function precedes HIF target gene
expression. *Biochim Biophys Acta* **1847**, 1254–1266 (2015).

52.   Grimm, C. *et al.* HIF-1-induced erythropoietin in the hypoxic retina protects against light-induced
retinal degeneration. *Nat Med* **8**, 718–724 (2002).

53.   Le, Y. Z. *et al.* Inducible expression of cre recombinase in the retinal pigmented epithelium. *Invest*
Ophthalmol Vis Sci **49**, 1248–1253 (2008).

54.   Bartoszewski, R. *et al.* Primary endothelial cell-specific regulation of hypoxia-inducible factor
(HIF)-1 and HIF-2 and their target gene expression profiles during hypoxia. *FASEB J* **33**, 7929–
7941 (2019).

55.   Chamboredon, S. *et al.* Hypoxia-inducible factor-1α mRNA: a new target for destabilization by
tristetraprolin in endothelial cells. *Mol Biol Cell* **22**, 3366–3378 (2011).

56.   Balakrishnan, B. & Jayandharan, G. R. Basic biology of adeno-associated virus (AAV) vectors
used in gene therapy. *Curr Gene Ther* **14**, 86–100 (2014).

57.   Mingozzi, F. & High, K. A. Immune responses to AAV vectors: overcoming barriers to successful
gene therapy. *Blood* **122**, 23–36 (2013).

58.   Burnett, J. C., Rossi, J. J. & Tiemann, K. Current progress of siRNA/shRNA therapeutics in clinical
trials. *Biotechnol J* **6**, 1130–1146 (2011).

59.   Whitehead, K. A., Langer, R. & Anderson, D. G. Knocking down barriers: advances in siRNA
delivery. *Nat Rev Drug Discov* **8**, 129–138 (2009).

Supplementary Figures

Figure S1 | Schematic representation of shRNAmir constructs with the miR-E backbone and shHif1a, shHif2a and shCtrl target sequences.

(A-C) Sequences of the shRNAmir constructs. Restriction enzymes used for cloning, the 5' complement, the basal stem, the target sequence, and the 3' flank are indicated.

Figure S2 | Evaluation of shHif2_1-6 in 661W cells

Hif2a mRNA levels in 661W cell lines stably expressing shCtrl or shHif2_1-6. Expression was normalized to beta-actin (*Actb*) and shown relative to expression of shCtrl (set to 1). Shown are means and individual values, n=2.

Figure S3 | Assessment of AAV-shHif injection in wt mice

In situ hybridization of retinal cross-sections from wt mice at 3 wpi detecting **(A)** *Hif1a* RNA in the shHif-transduced and the non-transduced retina or **(B)** *Hif2a* in the RPE transduced either with shHif or shCtrl AAVs. Shown are *in situ* fluorescence (red) and bright-field images (grey). **(C)** GFP fluorescence and immunostainings in retinal cross-sections of wt mice injected with PBS, shCtrl, or shHif at 3 wpi. Green: GFP; magenta/white: IBA1; grey: DAPI (nuclear stain). Shown are merged images (left) and single-channel images for IBA1 (right). RPE: retinal pigment epithelium. ONL: outer nuclear layer. INL: inner nuclear layer. GCL: ganglion cell layer. **(D)** Representative flatmount images of retina (top) and corresponding RPE (bottom) of wt mice injected with shCtrl (left) or shHif (right) at 3 wpi. GFP and mCherry fluorescence indicate the transduced area. Arrow: injection site. Antibodies for cone arrestin (ARR3, white) and beta-catenin (CTNB1, white) were used to stain cone photoreceptors and RPE cells, respectively. **(E)** Light microscopic images of panorama sections through retinas of wt mice injected with shCtrl (left) or shHif (right) at 22 wpi. Scale bar as indicated.

Figure S4 | Phenotype of $RPE^{\Delta Vegfa}$ mice.

Light microscopic panorama image of a retina section from an $RPE^{\Delta Vegfa}$ mouse (injected with

$1x10^9$ vg/µL AAV::VMD2-Cre-P2A-eGFP) at 39 wpi. The area transduced with the virus

resulting in an RPE-specific *Vegfa* knockout is marked. Insets: higher magnifications of boxed

areas. RPE: retinal pigment epithelium. PS: photoreceptor segments. ONL: outer nuclear

layer. INL: inner nuclear layer. GCL: ganglion cell layer. Scale bar as indicated.

1006

1007 **Supplemental data file 1 | AAV-plasmid sequences for shHif and shCtrl.**

1008 The shHif1, shHif2, and shCtrl sequences are highlighted in red, green, and blue, respectively.

Table 1 | Primer sequences for real-time PCR

Gene	Forward primer (5'-3')	Reverse Primer (5'-3')
<u>Mouse genes</u>		
Actb	CAACGGCTCCGGCATGTGC	CTCTTGCTCTGGGCCTCG
Hif1a	CAATGTCTCCTTTACCTTCATCGG	TGGATTCTTTGCCTCTGTGTCTTC
Hif2a	GGAGCTCAAAAGGTGTCAGG	CAGGTAAGGCTCGAACGATG
Slc2a1	AGTGTATCCTGTTGCCCTTCTG	GCCGACCCTCTTCTTTCATCTC
Casp1	GGCAGGAATTCTGGAGCTTCAA	GTCAGTCCTGGAAATGTGCC
Pde6b	TCTGGAGGATTGCCTGGTG	GGCGAATGAGCTGAAGTGG
Rpe65	AAAGCAGACAAGGAAGATCCAA	CCCAAAGACTCCACGAAGAA
<u>Human genes</u>		
ACTB	CCTGGCACCCAGCACAAT	GGGCCGGACTCGTCATAC
VEGFA	GTGGACATCTTCCAGGAGTACC	TGTTGTGCTGTAGGAAGCTCAT
ANGPTL4	AGAGGCAGAGTGGACTATTTG	TGGCGCCTCTGAATTACTG
SERPINE1	ACAAATCAGACGGCAGCAC	TAGGGCAGTTCCAGGATGTC
RPE65	TTGTGGATCTCTGCTGCTGGAA	GGAGCGTGACTAAATTCTTGCCT
GNAT2	CTGCTACTGCTGGGTGCT	TGGTGAATAGCCATCCTGGT

Table 2 | Treatment groups and AAVs

Treatment group	Mouse strain	Capsid	DNA construct
Single AAV injections			
WT + shCtrl	B6/J-Crl1	AAV8-TM	GRK1-GFP-shCtrl-VMD2-mCherry-shCtrl
WT + shHif	B6/J-Crl1	AAV8-TM	GRK1-GFP-shHif1-VMD2-mCherry-shHif2
Rod$^{\Delta Vhl}$ + shCtrl	*Vhl$^{fl/fl}$;Opsin-Cre*	AAV8-TM	GRK1-GFP-shCtrl-VMD2-mCherry-shCtrl
Rod$^{\Delta Vhl}$ + shHif	*Vhl$^{fl/fl}$;Opsin-Cre*	AAV8-TM	GRK1-GFP-shHif1-VMD2-mCherry-shHif2
Vegfa$^{fl/fl}$ + GFP	*Vegfa$^{fl/fl}$*	AAV4	VMD2-eGFP
RPE$^{\Delta Vegfa}$	*Vegfa$^{fl/fl}$*	AAV4	VMD2-Cre-P2A-eGFP
Double AAV injections (combined with AAV4/2::VMD2-Cre-P2A-GFP)			
RPE$^{\Delta Vegfa}$ + shCtrl	*Vegfa$^{fl/fl}$*	AAV8-TM	GRK1-GFP-shCtrl-VMD2-mCherry-shCtrl
RPE$^{\Delta Vegfa}$ + shHif	*Vegfa$^{fl/fl}$*	AAV8-TM	GRK1-GFP-shHif1-VMD2-mCherry-shHif2
RPE$^{\Delta Vegfa}$ + shHif1	*Vegfa$^{fl/fl}$*	AAV8-TM	GRK1-GFP-shHif1
RPE$^{\Delta Vegfa}$ + shHif2	*Vegfa$^{fl/fl}$*	AAV8-TM	VMD2-mCherry-shHif2
PBS and uninjected			
Vhl$^{fl/fl}$	*Vhl$^{fl/fl}$*	n.a.	n.a.
Vegfa$^{fl/fl}$	*Vegfa$^{fl/fl}$*	n.a.	n.a.
Vegfa$^{fl/fl}$ + PBS	*Vegfa$^{fl/fl}$*	n.a.	n.a. (PBS)

All AAVs have type 2 ITRs. AAV8-TM: AAV8-tripple mutation (Y447F + Y733F + T494V)

Table 3 | Primary antibodies used for immunofluorescence and Western blotting

Antibody	Cat. Number	Research Resource Identifier (RRID)	Vendor	Application	Dilution
IBA1	019-19741	AB_839504	Wako Pure Chemical Industries	Immunofluorescence	1:500
ARR3	AB15282	AB_1163387	Merck KGaA	Immunofluorescence	1:1000
PODXL	AF1556	AB_354858	R&D Systems	Immunofluorescence	1:200
CTNB1	610153	AB_397554	BD Transduction Laboratories™	Immunofluorescence	1:500
Cre	69050-3	AB_10806983	Merck	Immunofluorescence	1:300
HIF2A	AB199	AB_302739	Abcam	Western blotting	1:500
HIF1A	NB100-479	AB_10000633	Novus Biologicals	Western blotting	1:2000
ACTB	A5441	AB_476744	Sigma	Western blotting	1:10000

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A $Rod^{\Delta Vhl}$ $Rod^{\Delta Vhl}$ + shHif
BM
RPE
ONL
Vhl   *Hif2a*   *Hif1a*

B $Rod^{\Delta Vhl}$ 22 wpi
non-injected | + shCtrl | + shHif
Fundus | Fundus | Fundus
OCT | OCT | OCT
OCT | OCT | OCT
GFP | GFP | GFP
mCherry | mCherry | mCherry

C $Rod^{\Delta Vhl}$; *in vivo* quantification   22 wpi
$Vhl^{fl/fl}$ | + shCtrl | + shHif
ON 64 64 62 63 64 62 65 | ON 44 42 41 34 34 41 41 | ON 45 45 58 50 40 50 54

ONL thickness [units]
p<0.0001
p=0.0007
p=0.01
+ shCtrl
+ shHif
Ctrl ($Vhl^{fl/fl}$)

D $Rod^{\Delta Vhl}$; *morphology*   22 wpi
$Vhl^{fl/fl}$ | + shCtrl | + shHif
RPE | RPE | RPE
ONL | ONL | ONL
INL | INL | INL
GCL | GCL | GCL
50µm

E $Rod^{\Delta Vhl}$; retina   22 wpi
Hif1a p=0.018 | *Hif2a* p=0.854 | *Casp1* p=0.059 | *Pde6b* p=0.017
relative gene expression
+ shCtrl   + shHif

F $Rod^{\Delta Vhl}$; RPE / eyecup   22 wpi
Hif1a p=0.88 | *Hif2a* p=0.0372 | *Rpe65* p=0.528
relative gene expression
+ shCtrl   + shHif

Figure 6

A RPE$^{\Delta Vegfa}$ 39 wpi

+ shCtrl | + shHif

Fundus | OCT | OCT | GFP | mCherry | Fundus | OCT | OCT | GFP | mCherry

B RPE$^{\Delta Vegfa}$; *in vivo* quantification 39 wpi

+ shCtrl | + shHif | + shHif1 | + shHif2

24 24 ON | 50 56 ON | 35 38 ON | 38 43 ON

25 50 75 100 125 150

thickness [units]

p=<0.0001
p=0.0027
p=0.0450

○ + shCtrl
○ + shHif
○ + shHif1
○ + shHif2

C RPE$^{\Delta Vegfa}$; morphology 39 wpi

transduced area (nasal) | untransduced area (nasal)

+ shCtrl | + shHif | + shHif1 | + shHif2 | + shCtrl | + shHif | + shHif1 | + shHif2

RPE / ONL / INL / GCL   50μm | RPE / PS / ONL / INL / GCL   50μm

D RPE$^{\Delta Vegfa}$; *in vitro* quantification 39 wpi

ONL thickness | PS length

nasal (transduced) temporal (non-transduced) | nasal (transduced) temporal (non-transduced)

ONL thickness [μm] | PS length [μm]

Distance from ON [μm] | Distance from ON [μm]

p<0.0001
p=0.0091
p=0.099

p<0.0001
p=0.0372
p=0.0396

+ shCtrl
+ shHif
+ shHif1
+ shHif2

shCtrl shHif shHif1 shHif2

Figure 7